Dorso-Ventral and Night-Day Regulation of Extracellular K+ Dynamics in Mouse
Hippocampal Astrocytes

Nariman KIANI[1‡✉], Kabeer ABUBAKAR[1‡], Tin Luka PETANJEK[2], Monique ESCLAPEZ[1,3], Anton
IVANOV[1], and Christophe BERNARD[1✉]

[1] Aix Marseille Univ, INSERM, INS, Inst Neurosci Syst, Marseille, France

[2] University of Zagreb, Department of Physiology and Department of Neuroscience,
Croatian Institute for Brain Research, School of Medicine, Zagreb, Croatia

[3] Aix Marseille Univ, Inserm, INMED, Institut de Neurobiologie de la Méditerranée,
Marseille, France

‡ Equal contribution

✉ Corresponding authors

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author
upon reasonable request.

Orchid IDs

Nariman KIANI : https://orcid.org/0009-0003-8809-0521
Kabeer ABUBAKAR : https://orcid.org/0000-0001-7552-0383
Tin Luka PETANJEK : https://orcid.org/0000-0002-8950-8315
Monique ESCLAPEZ : https://orcid.org/0000-0002-8558-1363
Anton IVANOV : https://orcid.org/0000-0001-8305-1062
Christophe BERNARD: https://orcid.org/0000-0003-3014-1966

Acknowledgements

This work was supported by the European Union's Horizon 2020 Research and Innovation
Program under Grant Agreement Number 956325, the Fondation pour la Recherche
Médicale (FRM) FDT202404018111, the Croatian Science Foundation under project number
IP-2022-10-8493, and Petroleum Technology Development Fund
(PTDF/ED/OSS/PHD/KA/2015/22).

Correspondence: Nariman KIANI, Christophe BERNARD
Université Aix-Marseille, INS, Inserm, UMR1106, Faculté de Médecine de la Timone
27 boulevard jean moulin 13385 MARSEILLE, France
Nariman.KIANI@univ-amu.fr, Christophe.BERNARD@univ-amu.fr

Abstract

Astrocytes regulate extracellular potassium (K^+) through multiple mechanisms operating

across distinct spatiotemporal scales, yet whether this regulation exhibits regional and

circadian specificity remains unclear. Using real-time K^+ measurements combined with

electrophysiology and immunohistochemistry in astrocytes, we characterize K^+ buffering by

astrocytes across the dorsoventral hippocampal axis at three circadian times. We report

that hippocampal CA1 *stratum radiatum* astrocytes possess different functional K^+

buffering capacities depending on both anatomical location and time of day. At the early

light phase (ZT3), the ventral hippocampus (VH) exhibits faster K^+ accumulation and greater

peak amplitudes than the dorsal hippocampus (DH), despite similar neuronal network

activity. This regional divergence is driven by reduced Kir4.1 channel function in VH

astrocytes, which persists across all measured time points, and which is partially

compensated by enhanced Na^+/K^+-ATPase activity specifically at ZT3. Gap junction coupling

between astrocytes shows regional and time-dependent variation, with elevated coupling in

VH at ZT3 that subsequently normalizes. Kir4.1 protein expression exhibits circadian

dynamics, particularly in DH where expression declines from early light to late light and

recovers in early dark. These findings establish astrocytic K^+ buffering as a multiscale

phenomenon integrating regional heterogeneity with circadian regulation, with implications

for understanding regional and temporal differences in network excitability, particularly in

epilepsy.

Key words: Astrocytes, hippocampal regions, K+ homeostasis, Kir4.1 channels, circadian

rhythm

Introduction

Under physiological conditions, extracellular K^+ concentration ($[K^+]_o$) fluctuates between 2 and 5 mM, while increases above 10 mM are linked to pathological activities such as seizures and spreading depression (de Curtis et al., 2018; Enger et al., 2015; Erecińska & Silver, 1994). Maintaining $[K^+]_o$ within narrow physiological limits is therefore functionally important.

Astrocytes play an important role in regulating $[K^+]_o$ by taking up excess extracellular K^+, redistributing it through gap-junction–connected networks, and extruding it where $[K^+]_o$ is lower (Wallraff et al., 2006; Walz, 2000). This process relies on inward-rectifying Kir4.1 channels (Seifert et al., 2009), and the Na^+/K^+-ATPase pump. Impairment of either component can increase neuronal and network activity and excitability (Djukic et al., 2007; Romanos et al., 2020; Wallraff et al., 2006).

Neuronal activity and excitability are not uniform across the brain. Even within a single structure, distinct subregions exhibit marked differences in intrinsic excitability, synaptic properties, and responses to pharmacological or pathological challenges. For example, along the dorsoventral axis of the hippocampus, the ventral region (VH) displays higher intrinsic excitability, greater seizure susceptibility, distinct patterns of synaptic plasticity, and different metabolic properties compared to the dorsal region (DH) (Brancati et al., 2021; Dougherty et al., 2012; Papatheodoropoulos et al., 2005). Similarly, neuronal activity is not constant across time: circadian rhythms modulate neuronal excitability, synaptic transmission, and seizure threshold (B.-L. Chang et al., 2018; Debski et al., 2020; Quigg et

al., 2000), reflecting temporal regulation of molecular, metabolic, and physiological

processes (Brancati et al., 2021; Debski et al., 2020; McCauley et al., 2020).

This raises a fundamental question: do astrocytes exhibit similar spatial and temporal

heterogeneity in their K^+ buffering capacity? Do they "follow the pace" of neuronal activity,

adapting their homeostatic functions to match regional differences in excitability and

circadian variations in network state? If astrocytic K^+ buffering were spatially and temporally

invariant, regions of high neuronal excitability (such as the VH) and periods of elevated

seizure susceptibility (such as specific circadian phases) would be vulnerable to inadequate

K^+ homeostasis. Conversely, if astrocytes dynamically adjust their buffering capacity in

space and time, this could represent a sophisticated form of adaptive homeostasis

matched to local circuit demands.

The hippocampus provides an ideal system to address this question. Beyond the well-

documented differences in neuronal properties along the dorsoventral axis and across

circadian time, the hippocampus also exhibits regional and temporal heterogeneity in

astrocytic molecular markers, morphology, and calcium signaling (Dong et al., 2009; Jinno,

2011; Ryu et al., 2024; Thompson et al., 2008). However, whether these molecular and

morphological differences translate into functional differences in K^+ buffering—and how

they relate to the known regional and circadian variation in neuronal excitability—remains

unclear.

Here, we directly test whether astrocytic K^+ buffering capacity varies across the

dorsoventral hippocampal axis and across circadian time, using integrated functional,

electrophysiological, and molecular approaches. We use an extracellular K^+ sensor to

measure K^+ dynamics during standardized synaptic stimulation (Schaffer collateral

activation), monitoring multiple parameters of the K^+ transient (accumulation rate, peak

amplitude, clearance kinetics, and Na^+/K^+-ATPase-dependent undershoot) in dorsal and

ventral hippocampal slices at three Zeitgeber times: ZT3 (early light, baseline period), ZT8

(late light, period of heightened seizure susceptibility (W.-C. Chang et al., 2018; Debski et

al., 2020; Quigg et al., 2000), and ZT15 (early dark, active phase). We then employ whole-cell

patch-clamp electrophysiology to assess the cellular mechanisms underlying these

functional differences, examining gap-junction coupling, Kir4.1 channel function, and

astrocyte resting membrane potential. Finally, we use immunohistochemistry to quantify

regional and circadian variation in Kir4.1 protein expression. Our findings demonstrate that

astrocytic K^+ buffering capacity depends upon both anatomical location and circadian

phase, revealing a form of adaptive heterogeneity that parallels—but does not simply

mirror—the spatial and temporal organization of neuronal excitability.

Methods

Animals and Ethical Approval

All animal procedures were conducted in accordance with the European Directive

2010/63/EU and French regulations on animal experimentation. The study protocol was

approved by the relevant institutional and national ethics committees (APAFIS authorization

#44598-2023081812555435v4). A total of 50 male FVB mice (weight: 28–45 g; age: 2 months

or older) were used. Of these, 3 mice were dedicated to immunohistochemistry, while the

remaining animals were used for both patch-clamp experiments and K^+ dynamic recordings

in parallel. The single-sex design increased statistical power and ensured an adequate yield

of viable samples given the technically demanding nature of the experiments. The exclusion

of female mice is a limitation of the study, and future work will be required to assess the

generalizability of these findings across sexes.

Brain slice preparation

Mice anesthetized with sevoflurane, were decapitated, and the brain was swiftly removed

from the skull and submerged in ice-cold artificial cerebrospinal fluid (ACSF) (in mM): NaCl

126, KCl 3.50, NaH_2PO_4 1.25, $NaHCO_3$ 25, $CaCl_2$ 2, $MgSO_4$ 1.3, and glucose 10, with a pH

value maintained between 7.3 and 7.4. The ACSF was aerated with a 95% O_2/5% CO_2 gas

mixture. One hemisphere, left or right in alternation, was used to prepare dorsal slices via

coronal sections, whereas the other hemisphere was used to prepare ventral slices as

described previously (Dougherty et al., 2012). The brain was immersed in an ice-cold cutting

solution (in mM): K-gluconate 140, HEPES 10, Na-gluconate 15, EGTA 0.2, and NaCl 4, with

the pH adjusted to 7.2 with KOH. Using a tissue slicer (Leica VT 1200s, Leica Microsystem,

Germany), 300 μm-thick for patch-clamp recording and 350 μm- thick slices for field

recording were produced. Subsequently, to preserve the cells near the surface from

resulting damage caused by slicing, the 300 μm-thick slices were incubated for 15 to 20

minutes at room temperature (RT) in a choline chloride solution (in mM): Choline Chloride

110, KCl 2.5, NaH_2PO_4 1.25, $MgCl_2$ 10, $CaCl_2$ 0.5, $NaHCO_3$ 25, Glucose 10, Na-Pyruvate 5,

constantly bubbled with a 95% O_2/5% CO_2 gas mixture. To label the astrocytes, the slices

were incubated for 20 minutes in a ACSF solution containing 1 μM of sulforhodamine 101

(SR101, Merk), a fluorescent dye specifically absorbed by astrocytes and not neurons

(Nimmerjahn et al., 2004). The slices were allowed to recover in a dual-side perfusion

holding chamber with continuously circulating ACSF for at least one hour before

experimentation.

Synaptic stimulation and field potential recordings

In parallel, 350 µm-thick transferred to a dual perfusion recording chamber continuously

perfused (~6 ml/min) with ACSF containing 5 mM of glucose, warmed to 30°C. Schaffer

collateral (SC) was stimulated with a bipolar metal electrode connected to a DS2A-isolated

stimulator (Digitimer Ltd, UK). The electrode tip was placed in the *Stratum Radiatum* (*s.r.*)

between the CA3 and CA2 regions. Extracellular recording electrodes were pulled (Harvard

Apparatus Ltd, UK , O.D. = 1.2 and I.D. = 0.94 mm, Rp < 1 MΩ) using a horizontal puller (Sutter

Instruments, USA). The extracellular local field potential (LFP) was recorded with a glass

microelectrode filled with ACSF, placed in the *Stratum Oriens* (*s.o.*) towards the *Stratum*

Pyramidale (*s.p.*) of the CA1 area. The signal was amplified 1000 times with an EXT-02F

amplifier (NPI Electronic, Germany) operating with a low-pass 3kHz filter (DC mode). The

stimulation current intensity was adjusted to produce an LFP response of 60-70% of the

maximum amplitude. Each slice was stimulated with trains of stimulation pulses of different

frequencies ranging from 2.5Hz to 100Hz. The LFP responses were recorded simultaneously

with the K^+-dependent potential using Patch Master software. Each stimulation protocol

was repeated twice. The LFP responses to single pulse stimulation were monitored in

between train stimulations to check the slice stability. If the response decreased by more

than 20%, the slice was discarded. In this study, we used the response to the 10Hz30s trains

because this stimulation evoked a K+ signal with good SNR, allowing the measurement of

required parameters (peak amplitude, 25-75% rise time, decay time, and undershoot

amplitude) of the individual and not the averaged response.

Fabrication of Potassium Sensitive Microelectrode

K+-selective microelectrodes were prepared using the method described by Heinemann

and Arens (1992). In brief, electrodes were pulled from double-barrel theta glass (TG150-4,

Warner Instruments, Hamden, CT, USA). The reference barrel was filled with 154 mmol/L

NaCl solution. The silanized ion-sensitive barrel tip (5% trimethyl-1-chlorosilane in

dichloromethane) was filled with a potassium ionophore I cocktail A (60031 Fluka

distributed by Sigma-Aldrich, Lyon, France) and backfilled with 100 mmol/L KCl.

Measurements of K+-dependent potentials were performed using a high-impedance

differential DC amplifier equipped with negative capacitance feedback control, which

permitted the compensation of the microelectrode capacitance required for the recording

of the rapid changes of $[K^+]_o$. The electrodes were calibrated before each experiment. The

calibration curve, concentration vs potential, was fitted with an exponential function. The

obtained equation was used to convert $[K^+]_o$-dependent potential recordings into

concentration changes.

Electrophysiological recording of the astrocytes

For the electrophysiological recordings of astrocytes, 300 µm-thick slices were immersed in

a low-volume (2 ml) recording chamber and continuously perfused with ACSF at 32 - 34°C

and a perfusion rate of 5 mL/min. Sulforhodamine labeled astrocytes in the *stratum*

radiatum (*s.r.*) of the CA1 region of the hippocampus were identified as a fluorescent cells

with astrocytic morphology (excitation/emission wavelengths of 586/605 nm). Patch

pipettes were fabricated using the same horizontal puller from borosilicate glass tubing

(Harvard Apparatus Ltd, UK ,1.5 mm outer diameter (O.D.), 0.86 mm inner diameter (I.D.),

pipette resistance (Rp) = 5.78 ± 0.12 MΩ) and filled with an intrapipette solution (in mM): KCl

20, K-gluconate 115, HEPES 10, EGTA 1.1, MgATP 4, Na -phosphocreatine 10, Na_2GTP 0.4,

and 0.3-0.4% biocytin (Sigma, Germany). Signals were transmitted to a Multiclamp 700A

amplifier (Molecular Devices), digitized at a rate of 10 kHz using a DigiData 1550 interface

(Molecular Devices) connected to a personal computer, and analyzed using ClampFit

software (Molecular Devices). The resting membrane potential (RMP) of astrocytes was

measured using the patch-clamp technique, whole-cell configuration, in current clamp

mode.

Isolation of Kir4.1 currents

To isolate Kir4.1 currents and to keep constant the extracellular K^+, slices were incubated

for 36 ± 2 minutes in the baseline 5 mM glucose-containing ACSF supplemented with (in

µM): Meclofenamic acid (MFA, Sigma-Aldrich) 100 to block gap junctions, and Tetrodotoxin

(TTX, Abcam) 0.5, 6-Cyano-7-nitroquinoxaline-2,3-dione (CNQX, HelloBio) 10, and D-(-)-2-

Amino-5-phosphonopentanoic acid (D-AP5, HelloBio) 40 to stop neuronal firing and

minimize synaptic transmission. No effect of incubation time on Kir4.1 current was

detected. Upon establishment of the whole-cell patch clamp configuration, the patched

astrocyte was held at -70 mV in voltage clamp mode, and after 5 minutes of dialysis,

membrane currents were recorded at different holding potentials varying from -140mV to

50mV with 10mV steps. Then, to block Kir4.1 channels, 100 µM of $BaCl_2$ was added to the

baseline ACSF, and the currents elicited by the same voltage steps were recorded after 5

minutes. The current-voltage relationship of Ba^{2+} sensitive conductance was determined by

subtracting the currents recorded in Ba^{2+} from those recorded before its administration.

Access resistance was determined by biexponentially fitting the capacitive current resulting

from −70 to −60 mV voltage step. Cells were discarded if they met any of the following

criteria: an initial access resistance above 35 MΩ, a resting membrane potential (RMP) less

negative than -50 mV, or a final access resistance increase of more than 20%.

Coupling estimation

Astrocytes are extensively coupled via gap junctions formed by Cx43 and Cx30, supporting

intercellular redistribution of ions and metabolites, including potassium buffering.

Pharmacological inhibition of these gap junctions is expected to uncouple astrocytes and

thereby reducing their capacity for spatial potassium buffering. In principle, comparing K^+-

dependent currents evoked by Schaffer collateral stimulation before and after gap junction

blockade could provide an estimate of the coupling state of the recorded astrocyte.

However, available gap junction inhibitors lack specificity and profoundly affect neuronal

and glial function. Carbenoxolone alters both AMPA and $GABA_A$ receptor-mediated synaptic

transmission (Tovar et al., 2009). Meclofenamic acid is a potent activator of KCNQ2/Q3

(Kv7.2/7.3, M-type) channels (U. A. Anderson et al., 2009). and, at the 50–100 µM

concentrations commonly used to block gap junctions, hyperpolarizes CA1 neurons and

suppresses synaptic responses (Peretz et al., 2005). Connexin mimetic peptides such as

Gap26 and Gap27, while they are more specific at the channel level, also reduce basal

excitatory synaptic transmission, likely by inhibiting gliotransmitter release through Cx43

hemichannels (Dospinescu et al., 2025). Because all of these agents can alter overall

network activity, they inevitably affect not only K+ buffering but also activity-dependent K+

release. To avoid these confounding effects, we adopted an alternative approach to assess

astrocytic coupling, based on the spread of the gap junction–permeable intracellular tracer

biocytin following controlled whole-cell loading (Stephan et al., 2021).

In a separate set of experiments, astrocytes were patched in the absence of any blocker in

whole cell configuration and kept at -70 mV holding potential for 20 mins. Slices were then

fixed in 4% paraformaldehyde (AntigenFix, Diapath) overnight at 4°C. After three 30-minute

washes in 0.12 M phosphate buffer (PB), slices were cryoprotected in 20% sucrose

overnight, then deep-frozen on dry ice and stored at -80°C until the day of experimentation.

To reveal biocytin-containing astrocytes, slices were rinsed in PB twice, in 0.02 M potassium

phosphate buffered saline (KPBS) three times for 30 min, and then incubated with

streptavidin-conjugated with fluorophore Alexa 488 (Abcam) diluted (1/200) in KPBS

containing 0.3% Triton-X100, overnight at RT. After washing with KPBS for 3 x 30 min, slices

were mounted on Superfrost™ slides using Fluoromount-G.

To visualize the stained astrocytes coupling, slices were imaged using a Zeiss LSM 700

inverted confocal microscope with a 20x (0.5x zoom) objective. Z-stacks (15–35 optical

slices, length: 2048 pixels, width: 2048; 1 µm interval) were acquired in the CA1 *stratum*

radiatum using Zeiss ZEN software. Z-stacks were merged using Fiji software (NIH), and pixel

intensity was then normalized.

A blinded examiner then counted the number of stained astrocytes. Since only one astrocyte

per slice was patched, the stained astrocytes were mandatory, those connected with the

patched one directly or through other connected astrocytes. Therefore, we considered the

number of stained astrocytes as an estimation of the astrocytic coupling.

Immunohistochemistry and image analysis

Tissue preparation, Immunohistochemistry, and image analysis

Wild-type male FVB mice (P50–70) were anesthetized with an intraperitoneal injection of

ketamine and xylazine, followed by a lethal dose of pentobarbital administered at

designated Zeitgeber times (ZT3, ZT8, or ZT15). Mice were then transcardially perfused with

cold 0.12 M PB to remove the blood and then with AntigenFix to fix the tissue. Brains were

extracted, post-fixed for up to 4 hours in the same fixative, rinsed in PB, cryoprotected in

20% sucrose in PB overnight at 4°C, quickly frozen on dry ice and kept at -80°C. Hemisected

blocks of brain containing dorsal and ventral hippocampus were sectioned at 40 µm using

a cryostat. The Sections were collected sequentially in tubes containing an ethylene glycol-

based cryoprotective solution (Lu & Haber, 1992; Watson et al., 1986) and stored at –20°C

until histological processing.

Double labeling for GFAP and Kir4.1

Sections were processed for the simultaneous detection of GFAP to label astrocytes and

the Kir4.1 channel. They were rinsed for 30 min in PB and for 2x30 mins in KPBS. Sections

were then incubated for 1h at RT in 3% normal donkey serum (NDS) diluted in KPBS

containing 0.3% Triton X-100 and overnight at RT in a solution containing the polyclonal

guinea pig anti -GFAP (1:600, #AFP-001-GP, Alomone Labs) and polyclonal rabbit anti-Kir4.1

(1:500, #APC-035, Alomone Labs) diluted in KPBS containing 0.3% Triton X-100 and 1% NDS.

After 3, 30-minute rinses in KPBS, sections were incubated for 2 h in Alexa Fluor 488-

conjugated donkey anti-guinea pig IgG (1:100, #706-545-148, Jackson ImmunoResearch)

and Cy3-conjugated donkey anti-rabbit IgG (1:100, #711-167-003, Jackson

ImmunoResearch), diluted in KPBS and 3% NDS. Sections were then rinsed with 3x30 rinses

of KPBS and mounted on Superfrost™ slides using Fluoromount-G containing the nuclear

marker DAPI (Thermo Fisher).

Immunofluorescence was imaged on a Zeiss LSM 700 inverted confocal microscope using

a 63x oil-immersion objective. Z-stacks (12–19 optical slices at 1 μm intervals) of the CA1

stratum radiatum were acquired using Zeiss ZEN software. One field of view per section was

imaged. Z-projections were performed in Fiji (NIH), and GFAP signal was quantified by

measuring its integrated density. Thereafter, the GFAP was thresholded using triangle

method, creating a mask that defined the region of interest (ROI) for Kir4.1 analysis. Then,

Kir4.1 signal within the GFAP-defined ROI was quantified using the AND operation in the

Image Calculator, and the Kir4.1/GFAP integrated density ratio was calculated (Bataveljic et

al., 2024).

Statistical analysis

Statistical analyses were performed using R software (version 4.5.1). K^+ transient

parameters were all normalized to that of DH at ZT3, and subsequent analyses were

performed on these normalized data. To provide a comprehensive analysis of the data, we

used two different approaches: permutation-based hypothesis testing to identify significant

main effects and Bayesian multilevel modeling to estimate the magnitude and uncertainty

of those effects.

Permutation-Based Hypothesis Testing

Analyses involving the effects of region (DH vs. VH) and time (ZT3, ZT8, and ZT15) were

performed using a resampling-based, Freedman-Lane permutation ANOVA from the

permuco package in R (Kherad-Pajouh & Renaud, 2015). This non-parametric approach was

selected to provide robust inference by reducing assumptions about data distribution—

such as normality and sphericity. The analysis was conducted with 10,000 permutations to

ensure stable p-value estimates. Effect sizes were calculated and reported as omega

squared (ω^2), which represents the proportion of variance in the tested parameter

accounted for by each effect in the population. The magnitude of regional differences was

also evaluated using an online estimation statistics tool, which reported the mean

difference and corresponding 95% confidence intervals (Ho et al., 2019).

Bayesian Multilevel Modeling

To further analyze the effects of region, time, and their interactions while accounting for the

hierarchical structure of our experimental design (e.g., multiple cells nested within

animals), we performed Bayesian multilevel modeling using the brms package (version

2.22.0) (Bürkner, 2017). This framework allowed us to move beyond binary significance

testing to provide a probabilistic estimation of parameter uncertainty. Models were

specified with fixed effects for region, timepoint, and their interaction.

We used the fitdistrplus package (version 1.2.4) to identify the optimal family distribution for

each variable based on the lowest Akaike Information Criterion (AIC), ensuring the model

accurately reflected the data's underlying structure (Delignette-Muller & Dutang, 2015).

Four independent Markov Chain Monte Carlo (MCMC) chains were run with 4,000 iterations

each (2,000 warm-up); convergence was confirmed via $\hat{R} < 1.01$ and sufficiently high

effective sample sizes. Parameter estimates are reported as posterior medians with 95%

credible intervals (CrI). Predictors were centered and scaled where appropriate to aid

convergence.

Results

Extracellular K+ dynamics exhibit regional and circadian specificity

We first characterized how extracellular ($[K^+]_o$) concentration changes in the hippocampus

during and after standardized neuronal network activity, examining whether these dynamics

differ between dorsal and ventral hippocampus across circadian time. We used a 10 Hz, 30-

second stimulation of Schaffer collaterals to evoke robust $[K^+]_o$ release from activated

neurons (Fig. 1A and B). This protocol was selected to ensure that the neuronal network

response reached true peak amplitude and steady state before stimulus termination,

enabling consistent comparison across experimental groups (see Methods).

Synaptic strength is comparable across space and time

To ensure valid comparisons of $[K^+]_o$ transient properties, we first verified that evoked

neuronal activity was equivalent between regions and time points. The magnitude of the

local field potential (LFP) integral served as a quantitative proxy for network response

intensity. Statistical analysis revealed no significant effects of hippocampal region (DH

versus VH), Zeitgeber time (ZT), or their interaction on evoked LFP strength (Fig. 1C). This

uniformity confirms the robustness of our stimulation paradigm and validates the

assumption that any observed differences in $[K^+]_o$ dynamics reflect astrocyte properties

rather than variations in neuronal input.

Ventral hippocampus accumulates $[K^+]_o$ more rapidly and to greater amplitudes,

particularly at early light phase

We assessed $[K^+]_o$ accumulation by analyzing the peak amplitude of $[K^+]_o$ transients.

Permutation analysis of variance (ANOVA) shows a significant main effect of the regions only

($F(1, 98) = 12.05$, *permutation p* = .0010, $\omega^2 = 0.0950$), where the pool of VH samples shows

65% [95%CI 26%, 101%] larger $[K^+]_o$ peak amplitude than DH ones. Bayesian regression

confirmed that this difference is predominantly driven by ZT3. At this time point, $[K^+]_o$ in the

VH was 105% higher than in the DH [95% CrI: 40.5% , 200.4%]. In contrast, at ZT15, the DH

exhibited a peak amplitude that was 71.6% greater than at ZT3 [95% CrI: 17.4% , 150.9%],

suggesting that the transition from ZT3 (light cycle) to ZT15 (dark cycle) promotes greater

$[K^+]_o$ accumulation in the DH (Fig. 1D).

Analysis of the slope of the 25–75% rise phase of $[K^+]_o$ transients revealed that VH reached

peak $[K^+]_o$ 305.5% faster than the DH [95% CI: 124.8%, 646.3%] at ZT3. In the DH, the

transition from ZT3 to ZT8 produced no clear change [27.1%, 95% CrI: −26.7%, 120.3%],

whereas the transition to ZT15 resulted in a substantial increase in accumulation rate

[122.6%, 95% CrI: 28.4%, 293.5%]. Importantly, the regional advantage of the VH diminished

at later time points. Relative to its ZT3, the VH effect was attenuated by 41.7% at ZT8 [95%

CrI: 19.0%, 93.1%] and by 33.3% at ZT15 [95% CrI: 14.9%, 72.7%], confirming a significant,

negative interaction between region and time. (Fig. 1E).

Collectively, these results demonstrate that comparable neuronal network activity

produces different extracellular $[K^+]_o$ dynamics between dorsal and ventral hippocampus,

suggesting regional differences in astrocytic K^+ buffering capacity. This regional divergence

is most pronounced at the early light phase.

$[K^+]_o$ clearance kinetics show similar rates despite regional differences in

accumulation

To distinguish whether faster $[K^+]_o$ accumulation in VH reflects impaired $[K^+]_o$ removal or

differential regulation at the rising phase of activity, we analyzed $[K^+]_o$ decay kinetics by

fitting monoexponential decay (τ_{decay}) to the falling phase of the transient. Permutation

ANOVA revealed a significant main effect of region ($F(1, 98) = 3.93$, permutation $p = 0.0496$,

$\omega^2 = 0.0278$), with VH showing an overall 14.5% faster decay than DH [95% CI: -0.20%,

28.1%]. However, Bayesian regression analysis incorporating temporal variation did not

identify statistically robust regional differences across time points. When analyzed within

specific circadian phases, differences were modest and credible intervals crossed zero: at

ZT3, DH exhibited 22.3% slower decay than VH [95% CrI: -5.9%, 50.2%]; at ZT8 and ZT15,

differences were 15.2% [95% CrI: -14%, 41.5%] and 17.0% [95% CrI: -11%, 42.1%],

respectively (Fig. 1F).

This paradoxical finding – accelerated $[K^+]_o$ accumulation in VH despite similar or faster

clearance rates – indicates that differential $[K^+]_o$ dynamics are driven by regional difference

at the rising phase of the transient rather than the decay phase, suggesting that distinct

buffering mechanisms may be preferentially engaged during accumulation versus recovery.

Ventral hippocampus displays enhanced putative Na$^+$/K$^+$-ATPase activity specifically

at early light phase

The similar decay kinetics between regions despite higher $[K^+]_o$ accumulation in VH

prompted us to examine the recovery phase. After prolonged stimulation (10 Hz, 30 s), $[K^+]_o$

return to baseline is followed by an undershoot (Fig. 1B, right panel, black arrows), a

transient decrease of K^+ concentration below baseline level that reflects Na$^+$/K$^+$-ATPase

activity (Chever et al., 2010; D'Ambrosio et al., 2002). Analysis of undershoot amplitude

revealed significant regional effects and complex temporal dynamics: permutation ANOVA

revealed a significant effect of region ($F(1, 105) = 26.65$, permutation $p = 0.0001$, $\omega^2 = 0.180$)

as well as significant interaction of time with region ($F(2, 105) = 3.074$, permutation $p =$

0.0482, $\omega^2 = 0.0290$). Bayesian regression analysis revealed that at ZT3, VH showed a 239%

increase in undershoot amplitude compared to DH [95% CrI: 54%, 624%]. While VH

maintained elevated undershoot amplitudes at ZT8 and ZT15, the magnitude of this regional

difference was attenuated at these later timepoints [ZT8: -49%, 95% CrI: -82% , +42%; ZT15:

-48%, 95% CrI: -81% , +48%] (Fig. 1G).

These findings suggest that VH astrocytes display enhanced Na^+/K^+-ATPase activity,

particularly pronounced at the early light phase. Despite this elevated pump activity, overall

$[K^+]_o$ clearance rates remain unchanged, suggesting that the enhanced pump activity

primarily reflects the greater $[K^+]_o$ accumulation in VH (which provides greater driving force

for pump activation) rather than fundamentally altering the rate constant of $[K^+]_o$ removal.

This raises the question: what cellular mechanisms limit $[K^+]_o$ buffering capacity in the

ventral hippocampus?

Cellular basis of regional $[K^+]_o$ buffering differences: a role for reduced Kir4.1 function

in ventral astrocytes

Astrocytic Gap Junction Coupling Shows Regional and Temporal Variation

Potassium redistribution through astrocytic networks depends on gap junction-mediated

coupling, which expands the effective volume available for $[K^+]_o$ uptake. We examined

whether astrocytic coupling—measured by quantifying the number of biocytin-labeled cells

connected to a patched astrocyte—exhibits regional and circadian variation (Fig. 2A).

Across all time points, VH slices showed a trend toward more coupled astrocytes than DH

[mean difference: 13 cells, 95% CI: -4, 30], but this difference did not reach statistical

significance in permutation ANOVA ($F(2, 63) = 2.75$, permutation $p = 0.0686$, $\omega^2 = 0.0483$).

However, Bayesian regression analysis revealed significant regional and temporal effects.

Specifically, at ZT3, VH slices exhibited 39 more coupled cells than DH [95% CrI: +15 , +69],

corresponding to a 36.3% increase [95% CrI: 8.3% , 70.0%]. This regional difference was

absent at ZT8. At ZT15, DH coupling increased by 27% compared to its ZT3 value [95% CrI:

1% , 60%], while VH coupling declined by 16.2% [95% CI: 6.8% , 52.3%] (Fig. 2A and B).

Enhanced gap junction coupling in VH at ZT3 would be expected to facilitate $[K^+]_o$

redistribution and thus improve buffering capacity. The fact that VH nevertheless shows

greater $[K^+]_o$ accumulation at this timepoint suggests that reduced K^+ conductance

mechanisms (rather than impaired redistribution) are the primary determinant of regional

differences.

Ventral hippocampal astrocytes display reduced Kir4.1-mediated conductance across

all circadian phases

Kir4.1 channels mediate the primary K^+-dependent astrocytic conductance and are central

to spatial buffering, particularly during the rising phase of $[K^+]_o$ transients when astrocytic

depolarization drives K^+ influx through these channels (Larsen et al., 2014; Seifert et al.,

2009). To directly assess Kir4.1 function, we performed whole-cell patch-clamp recordings

(Fig. 3A) in the presence of gap junction blocker (MFA) and neuronal transmission blockers

(TTX, CNQX, D-AP5), isolating Kir4.1-specific currents via subsequent Ba^{2+} application

(Wallraff et al., 2006). Indeed, application of MFA was demonstrated to be effective in

uncoupling astrocytes, as shown by the number of dye-coupled cells (Fig. 3B-C).

Resting membrane potential did not vary significantly between regions or across time (Fig.

3D). Application of Ba^{2+} caused an average depolarization of 5.5 ± 0.64 mV consistent with

blockade of inward-rectifying currents, confirming effective channel blockade. Permutation

ANOVA revealed a significant main effect of region on Ba^{2+}-induced depolarization (F(1, 43)

= 8.26, permutation p = 0.0051, ω^2 = 0.124), with DH astrocytes depolarizing 3.2 mV more

than VH astrocytes following Ba^{2+} application [95% CI: 0.90, 5.70], indicating higher

functional Kir4.1 expression in DH (Fig. 3E).

To more directly quantify Kir4.1-mediated conductance, we analyzed Ba^{2+}-sensitive current

at -130 mV holding potential, a voltage optimal for Kir4.1 channel activity (Seifert et al.,

2009). Permutation ANOVA revealed a significant effect of region (F(1, 43) = 13.30,

permutation p = 0.0006, ω^2 = 0.209). Bayesian regression confirmed a significant main effect

of region, with no effect of time or region-time interaction. On average, Ba^{2+}-sensitive

conductance represents 29% of total transmembrane current in DH astrocytes compared

to 17% in VH, with a mean difference of 12% [95% CI: 5.8% , 17.6%] (Fig. 3F-G). This

reduction in functional Kir4.1 conductance in VH relative to DH persists across all examined

circadian phases, suggesting that regional differences in Kir4.1 function are stable

properties of these astrocytes rather than acutely regulated.

This reduced Kir4.1-mediated conductance in VH astrocytes provides a mechanistic

explanation for the observed rapid and pronounced K^+ accumulation in this region: with

lower K^+ conductance, astrocytes can passively take up less K^+ in response to

depolarization, resulting in higher $[K^+]_o$ despite the presence of enhanced gap junction

coupling that could support redistribution.

We noticed that in some astrocytes the IV relationship of Ba^{2+}-sensitive current was linear

without rectification. We checked whether such astrocytes are more frequent in one of the

explored regions. Current rectification was assessed by dividing I-V curve slopes between –

140 to –90 mV by that of 0 to +50 mV voltage ranges. Cells with slope ratios >1 were classified

as rectifying. Fisher's exact test revealed no statistically significant association between

region and rectification status (p = 0.104), though the estimated odds ratio of 0.28 [95% CI:

0.04, 1.31] suggested a non-significant trend toward greater rectification in DH astrocytes

(Fig. 3H). Our findings indicate that Kir4.1-mediated currents are regionally regulated, with

higher functional expression in the DH compared to VH but are not modulated by circadian

time under these experimental conditions. This difference partly explains why, at the same

rate of potassium release, its accumulation is faster and greater in VH than in DH.

Kir4.1 protein expression exhibits circadian dynamics distinct between dorsal and

ventral hippocampus

To investigate whether regional differences in Kir4.1 current reflect corresponding

differences in protein abundance, and to assess circadian regulation of Kir4.1 expression,

we performed double immunolabeling for GFAP (astrocyte marker) and Kir4.1 on

hippocampal sections across the regions (Fig. 4A-B) as well as time points.

Analysis of GFAP-positive area fraction revealed a main effect of region (F(1, 62) = 13.48,

permutation p = 0.0009, ω^2 = 0.157), although Bayesian analysis detected no statistically

robust effects (Fig. 4C). To account for regional differences in astrocytic Kir4.1 expression,

Kir4.1 integrated density was normalized to GFAP-positive area (astrocyte-specific

quantification). Values were further normalized within each immunolabeling batch to DH

samples to control batch variability.

Permutation ANOVA revealed a significant region × time interaction ($F_{(2, 62)} = 4.14$,

permutation $p = 0.0119$, $\omega^2 = 0.092$). Bayesian regression analysis revealed nuanced

circadian regulation: at ZT3 (early light), DH showed significantly higher Kir4.1 levels than VH

[39.3% increase, 95% CrI: 14.8%, 56.8%]. The transition from ZT3 to ZT8 (late light) was

marked by significant decline in DH Kir4.1 expression [34.3% decrease, 95% CrI: 10.4%,

51.8%], whereas VH showed the opposite pattern with a 24.4% increase [95% CrI: 22.1%,

197.4%], resulting in substantial convergence of expression levels. No reliable regional

differences were detected at ZT15 (early dark) (Fig. 4D).

Importantly, these circadian changes in Kir4.1 protein abundance do not translate into

circadian modulation of Kir4.1-mediated currents (which showed only regional, not

temporal, variation in functional assays). This dissociation between protein expression

dynamics and current magnitude suggests complex post-translational regulation of Kir4.1

channel function.

Integration and mechanistic synthesis

Our results reveal a multiscale organizational principle for astrocytic K^+ buffering in

hippocampus: regional and circadian factors converge to determine buffering capacity

through complementary mechanisms operating at different phases of $[K^+]_o$ transients (Fig.

5).

At the early light phase (ZT3), the ventral hippocampus exhibits a state of heightened

extracellular K^+ accumulation characterized by: 1) reduced Kir4.1-mediated K^+

conductance (41% lower than DH) that persists across all times, 2) enhanced gap junction

coupling (35% more coupled astrocytes) that could support redistribution, 3) elevated

Na^+/K^+-ATPase-dependent undershoot (239% greater than DH) indicating compensatory

pump activation. These properties produce rapid and pronounced $[K^+]_o$ accumulation

despite the presence of potentially advantageous gap junction coupling. The putative

enhanced Na^+/K^+-ATPase activity likely reflects activation by the greater local $[K^+]_o$

accumulation and astrocytic depolarization, coupled with potentially higher pump

expression. Thus, at ZT3, ventral astrocytes are limited in their capacity to prevent $[K^+]_o$

accumulation during the rising phase.

As the circadian cycle progresses through the light phase (ZT3 to ZT8 to ZT15): 1) Kir4.1

protein expression increases in VH and decreases in DH, partially equalizing regional

differences by ZT15, 2) gap junction coupling normalizes between regions, 3) Na^+/K^+-ATPase

compensatory activity diminishes.

By the early dark phase (ZT15), the circadian pattern reverses, with DH $[K^+]_o$ accumulation

increasing relative to ZT3, suggesting that dorsal hippocampal astrocytes are subject to

time-dependent suppression of buffering capacity during the light phase, with restoration

during the dark phase. The nature of this circadian suppression remains to be determined

but may reflect circadian regulation of Kir4.1 trafficking or channel modulation by clock-

controlled signaling pathways.

Discussion

This study addresses the question of spatial and temporal heterogeneity in astrocytic

function. Using integrated functional, electrophysiological, and molecular approaches, we

demonstrate that astrocytic K^+ buffering in the hippocampus is jointly regulated by regional

identity and circadian phase, with important implications for understanding network

excitability, seizure susceptibility, and the temporal organization of brain states.

Astrocytes show regional specialization but do not simply mirror neuronal excitability

patterns

Our findings reveal that ventral hippocampal astrocytes possess fundamentally different

$[K^+]_o$ buffering properties compared to their dorsal counterparts. At the early light phase

(ZT3), when regional differences are most pronounced, VH accumulates $[K^+]_o$ more rapidly

and to higher peak amplitudes than DH despite equivalent neuronal network activity. This

regional divergence reflects intrinsic differences in astrocyte physiology rather than

variations in neuronal input, establishing that astrocytic heterogeneity is a primary

organizational principle of hippocampal circuits.

However, astrocytic regional specialization does not simply track neuronal excitability in a

compensatory manner. The VH is known to exhibit higher intrinsic neuronal excitability and

greater seizure susceptibility than the DH (Debski et al., 2020; Dougherty et al., 2012;

Papatheodoropoulos et al., 2005). If astrocytes were optimized to compensate for regional

differences in neuronal activity, one would predict enhanced $[K^+]_o$ buffering capacity in VH

to counterbalance its elevated excitability. Instead, we observe the opposite: VH astrocytes

display reduced Kir4.1-mediated conductance (12% lower than DH), resulting in greater

$[K^+]_o$ accumulation during neuronal activity. This counterintuitive arrangement suggests that

astrocytic properties are not configured to suppress regional differences in excitability but

rather may amplify or enable them, supporting the functionally distinct roles mediated by

dorsal and ventral hippocampus in cognition and behavior.

The paradox of enhanced accumulation with preserved clearance: differential

mechanisms at rising vs. falling phases

A central finding of this study is the paradoxical observation that VH exhibits both faster $[K^+]_o$

accumulation and similar (or slightly faster) clearance rates compared to DH. This apparent

contradiction is resolved by recognizing that distinct buffering mechanisms dominate during

different phases of the K^+ transient, with Kir4.1 channels playing a central role during

accumulation and Na^+/K^+-ATPase predominating during recovery.

In this study, Na^+/K^+ ATPase activity was quantified using the amplitude of the post-stimulus

K^+ undershoot rather than by pharmacological pump inhibition because it does not

selectively reduce $[K^+]_o$ clearance. Indeed, the pump inhibition depolarizes neurons, alters

firing patterns, and modifies synaptic transmission (T. R. Anderson et al., 2010), thereby

changing K+ release during Schaffer collateral stimulation. Consequently, $[K^+]_o$ transients

recorded after pump blockade reflect a mixture of impaired clearance and altered K^+

release, which cannot be disentangled. In contrast, the $[K^+]_o$ undershoot that follows a

defined activation epoch arises from activity-dependent stimulation of the Na^+/K^+ ATPase

and reflects the excess pump-mediated uptake of K^+ once neuronal firing has ceased. For

these reasons, undershoot analysis offers a more specific and interpretable functional

measure of Na^+/K^+ ATPase activity in our experimental condition.

During the rising phase of neuronal activity, when K^+ is actively released from depolarizing

neurons, $[K^+]_o$ elevation depolarizes nearby astrocytes, driving passive K^+ influx through

Kir4.1 channels down the electrochemical gradient (Larsen & MacAulay, 2014). This Kir4.1-

mediated uptake represents the fastest way to limit $[K^+]_o$ accumulation. Our

electrophysiological data demonstrate that VH astrocytes possess significantly reduced

Ba^{2+}-sensitive (Kir4.1-mediated) conductance compared to DH astrocytes, a decrease that

persists across all circadian phases examined. This reduced conductance directly explains

why VH accumulates $[K^+]_o$ more rapidly: with fewer functional Kir4.1 channels, astrocytes

cannot buffer K^+ as efficiently during the rising phase, resulting in higher extracellular

concentrations.

In contrast, during the decay phase after neuronal activity ceases, $[K^+]_o$ clearance is

dominated by active Na^+/K^+-ATPase pumping rather than passive Kir4.1-mediated influx

(D'Ambrosio et al., 2002; MacAulay, 2020). Our observation that VH exhibits a dramatically

enhanced $[K^+]_o$ undershoot (239% larger than DH at ZT3)—a signature of robust Na^+/K^+-

ATPase activity—indicates that pump-mediated clearance is elevated in VH. Importantly,

this enhanced pump activity likely reflects a compensatory response to the greater $[K^+]_o$

accumulation itself: higher $[K^+]_o$ and greater astrocytic depolarization both stimulate

Na^+/K^+-ATPase activity (Wang et al., 2012). Thus, the similar decay kinetics between regions

emerge not from equivalent intrinsic clearance capacity but from activity-dependent

upregulation of pumping in VH that compensates for its reduced Kir4.1 function.

This mechanistic model—reduced Kir4.1-mediated uptake during accumulation,

compensated by enhanced pump-mediated clearance during recovery—reconciles the

paradoxical K^+ dynamics and highlights the phase-specific contributions of different

buffering mechanisms.

Gap junction coupling: enhanced connectivity cannot overcome reduced conductance

An unexpected finding was that VH astrocytes exhibit enhanced gap junction coupling at ZT3

(35% more coupled cells than DH), precisely when and where K^+ accumulation is greatest.

At first glance, this enhanced coupling appears maladaptive: expanded astrocytic networks

should increase the effective volume available for K^+ redistribution and thus improve

buffering capacity (Wallraff et al., 2006). Yet VH accumulates more $[K^+]_o$.

This paradox highlights an important distinction between network connectivity (assessed by

gap junction coupling) and cellular K^+ conductance (determined primarily by Kir4.1

expression). Enhanced coupling expands the astrocytic syncytium, but K^+ must first enter

astrocytes through Kir4.1 channels before it can be redistributed through the network. With

Kir4.1 conductance reduced by 12% in VH astrocytes, the membrane represents a

bottleneck that limits K^+ uptake regardless of how extensive the downstream network may

be. Thus, gap junction coupling appears to represent a partial compensatory adaptation—

expanding the potential redistribution capacity—but this adaptation cannot fully overcome

the primary deficit in membrane conductance.

This interpretation suggests that Kir4.1 channel function, rather than gap junction coupling,

is the rate-limiting step for K^+ buffering during active neuronal stimulation in VH.

Circadian regulation superimposed on stable regional identity

Our temporal analysis reveals that circadian phase modulates—but does not abolish—the

stable regional differences in K^+ buffering. Two key parameters exhibit no circadian

variation: (1) the time constant of K^+ decay, which remains similar between regions and

constant across time, and (2) Ba^{2+}-sensitive Kir4.1 current, which shows consistent regional

differences (lower in VH) at all timepoints despite circadian changes in Kir4.1 protein

expression.

The stability of Kir4.1 current despite dynamic protein expression points to complex post-

translational regulation. At ZT3, DH exhibits 41% higher Kir4.1 protein than VH; by ZT8, this

regional difference disappears due to reduced expression in DH and increased expression

in VH. Yet functional Kir4.1 currents remain regionally differentiated throughout the day. This

dissociation suggests that Kir4.1 channel activity is regulated not only by protein abundance

but also by trafficking to the membrane, phosphorylation state, interaction with scaffolding

proteins, or lipid microenvironment—mechanisms that may themselves be under circadian

control and differ between regions.

The most pronounced circadian effect is observed in the convergence of multiple

parameters at ZT3 versus later timepoints. At early light phase, the regional divergence

between DH and VH is maximal: K^+ peak amplitude differs 2.1-fold, gap junction coupling

differs by 35%, and Kir4.1 protein expression differs by 41%. By ZT8 and ZT15, these

differences attenuate or reverse. This temporal pattern suggests that early light phase—

corresponding to the rest period in nocturnal rodents—may be a time when regional

specialization is most important, possibly reflecting distinct roles for dorsal and ventral

hippocampus in sleep-dependent memory consolidation, sharp-wave ripple generation, or

metabolic recovery (Pronier et al., 2023).

Functional implications: regional K$^+$ dynamics shape circuit computations

The distinct K$^+$ buffering properties of DH and VH astrocytes have important implications for

how these regions process information and contribute to hippocampal function.

Extracellular K$^+$ accumulation exerts multiple effects on neuronal excitability, synaptic

transmission, and network oscillations, making astrocytic control of $[K^+]_o$ a powerful

regulator of circuit dynamics.

Ventral hippocampus: permissive K$^+$ accumulation enables synchronized network events

but imposes low-pass filtering. The rapid and pronounced $[K^+]_o$ accumulation in VH has both

facilitatory and inhibitory consequences. Elevated $[K^+]_o$ depolarizes neurons toward action

potential threshold, enhancing NMDA receptor activation by relieving Mg^{2+} block and

increasing the spatial range of glutamate spillover (Shih et al., 2013). These effects promote

network synchronization and may facilitate the generation of hippocampal sharp-wave

ripples (SWRs)—brief, high-frequency oscillations essential for memory consolidation.

Indeed, VH slices more readily generate spontaneous SWRs than DH slices (Schlingloff et

al., 2014), and CA3 pyramidal neurons in VH possess stronger recurrent connectivity and

higher intrinsic excitability (Sun et al., 2020), properties that synergize with permissive K$^+$

accumulation to support synchronized discharges.

However, sustained K$^+$ elevation during prolonged activity also inactivates voltage-gated

Na$^+$ channels and reduces the driving force for K$^+$ efflux through voltage-gated K$^+$ channels,

leading to synaptic depression and reduced responsiveness to repetitive stimulation (Meeks

& Mennerick, 2004). This creates a functional low-pass filter, where VH responds robustly to

initial or low-frequency inputs but saturates during sustained high-frequency activity. Such

filtering may prevent runaway excitation and protect against seizure propagation, while

simultaneously biasing VH toward processing sparse, salient events rather than sustained

information streams.

Dorsal hippocampus: tight K^+ control preserves high-fidelity transmission and supports

frequency-dependent facilitation. In contrast to VH, the strong Kir4.1-mediated buffering in

DH maintains lower $[K^+]_o$ during neuronal activity, preserving neuronal excitability and

enabling synaptic facilitation at frequencies between 10-50 Hz—precisely the range that

encodes behaviorally relevant information (Papaleonidopoulos et al., 2017). Lower K^+

accumulation prevents synaptic depression, supporting sustained information transfer and

communication with neocortical targets (Moreno et al., 2016). This configuration suits DH

for high-pass filtering: reliable transmission of sustained, structured activity patterns

required for spatial navigation, episodic memory encoding, and cognitive map formation

(Fanselow & Dong, 2010).

The metabolic consequences of these regional differences are consistent with recent

findings that VH relies more heavily on aerobic glycolysis—generating cytosolic ATP rapidly

but inefficiently—compared to the oxidative phosphorylation favored in DH (Brancati et al.,

2021). The elevated Na^+/K^+-ATPase activity required to clear excess K^+ in VH imposes

substantial energetic demands that may be preferentially met by glycolytic metabolism.

Pathological implications: regional K$^+$ dysregulation and epilepsy

The reduced K$^+$ buffering capacity of VH astrocytes provides a mechanistic framework for understanding the well-established vulnerability of VH to seizures. Evidence from human temporal lobe epilepsy (TLE) and animal models identifies VH as the primary seizure initiation zone (Babb et al., 1984; Bernasconi et al., 2003; Buckmaster et al., 2022; Toyoda et al., 2013). Our findings suggest that intrinsic differences in astrocytic K$^+$ buffering—present even in healthy tissue—predispose VH to hyperexcitability.

During pathological conditions such as epilepsy, when neuronal activity is abnormally intense and synchronized, the limited Kir4.1 capacity in VH would result in excessive $[K^+]_o$ accumulation, reaching levels (>10 mM) known to be associated with or even trigger seizures (de Curtis et al., 2018; Enger et al., 2015). Furthermore, epileptic tissue exhibits downregulation of Kir4.1 expression and function (Bataveljic et al., 2024; Romanos et al., 2020), exacerbating the baseline deficit in VH and creating a positive feedback loop, i.e. reduced buffering results in greater K$^+$ accumulation, which enhances excitability and thus more intense neuronal firing, favoring further K$^+$ release.

The circadian modulation of K$^+$ buffering we observe also provides a potential mechanism for the circadian rhythmicity of seizure occurrence documented in epilepsy patients and animal models (B.-L. Chang et al., 2018; Proix et al., 2021; Quigg et al., 2000). Seizure susceptibility peaks during specific circadian phases, and our data suggest that times of maximal regional divergence in K$^+$ buffering (e.g., ZT3) may correspond to windows of heightened vulnerability in VH. Conversely, circadian phases when regional differences

attenuate may represent periods of reduced seizure risk. However, given the extensive

reorganization of circuits in epilepsy, it will be important to assess all the above-described

properties in experimental epilepsy.

Dissociation between Kir4.1 protein expression and functional current: post-

translational complexity

One of the most intriguing findings is the dissociation between Kir4.1 protein expression

dynamics (which vary significantly across circadian time) and functional Kir4.1-mediated

currents (which show stable regional differences but no temporal variation). This

dissociation implies that channel function is regulated at multiple levels beyond simple

protein abundance.

Several post-translational mechanisms could account for this complexity. Kir4.1 channels

must traffic to the plasma membrane to be functional, and this trafficking is regulated by

interactions with scaffolding proteins such as dystrophin-associated protein complex and

α-syntrophin (reviewed in (Olsen & Sontheimer, 2008)). Circadian changes in trafficking

machinery or cytoskeletal organization could alter the proportion of synthesized channels

that reach the membrane. Additionally, Kir4.1 activity is modulated by phosphorylation, pH,

and membrane lipid composition (Hibino et al., 2010)—all of which exhibit circadian

oscillations (McCauley et al., 2020; Ryu et al., 2024). Finally, Kir4.1 channels interact with

other membrane proteins in macromolecular complexes, and changes in these interactions

could alter channel open probability or conductance without changing total protein levels.

The persistence of regional differences in functional current despite convergence of protein

expression also suggests that DH and VH astrocytes may differ in their complement of

regulatory proteins or post-translational modifications. Identifying these region-specific

regulatory mechanisms represents an important direction for future research and could

reveal novel targets for modulating astrocytic K^+ buffering in disease states.

Comparative context: astrocytic heterogeneity as a general principle

Our findings contribute to a growing recognition that astrocytic heterogeneity is a

fundamental organizational principle across the brain. Astrocytes differ between brain

regions in molecular marker expression, morphology, calcium signaling dynamics,

glutamate uptake capacity, and contribution to synaptic function (reviewed in (Khakh &

Deneen, 2019)). Within the hippocampus, dorsoventral differences in astrocytic gene

expression, morphology, and calcium dynamics have been documented (Dong et al., 2009;

Jinno, 2011; Ryu et al., 2024; Thompson et al., 2008), but the functional consequences of

this heterogeneity for circuit physiology have remained unclear.

Our demonstration that astrocytic K^+ buffering—a core homeostatic function—exhibits

pronounced regional and temporal variation establishes that astrocytic heterogeneity is not

merely a descriptive phenomenon but has direct, measurable impacts on network

excitability and information processing. This principle likely extends beyond the

hippocampus to other brain structures where astrocytic properties may be tuned to local

circuit demands.

Importantly, our findings challenge the traditional view of astrocytes as passive, uniform

support cells that maintain brain homeostasis in a spatially and temporally invariant

manner. Instead, astrocytes emerge as active, heterogeneous participants in circuit

function whose properties are dynamically regulated in space and time. This has

implications for understanding both normal brain function and the pathophysiology of

neurological disorders.

Reconciling slice physiology with in vivo function: caveats and considerations

Several important caveats limit the interpretation of our findings. First, our measurements

were performed in acute brain slices, which lack the intact vascular system, interstitial fluid

flow, and long-range connectivity present in vivo. K^+ clearance mechanisms that rely on

vascular uptake or long-distance redistribution through white matter tracts may be

compromised in slices, potentially altering the relative contributions of different buffering

mechanisms. Evidence suggests that gap junction-mediated spatial buffering contributes

more to K^+ clearance in vivo than in slices, raising the possibility that the enhanced gap

junction coupling we observe in VH may be more functionally significant in intact brain than

our slice experiments suggest (Breithausen et al., 2020; Cooper et al., 2025).

Second, patch-clamp approach in astrocytes samples a small radius of somata (Zhou et al.,

2021), which may not fully represent the properties of fine perisynaptic processes where K^+

buffering most directly impacts synaptic function. Recent advances in voltage imaging

reveal that distal astrocytic processes experience larger depolarizations in response to

neuronal   activity   than   somata   do   (Armbruster et al., 2022),   and   spatial

compartmentalization of ion channels may create local K^+ buffering microdomains that our

somatic recordings cannot detect. Additionally, we used GFAP immunolabeling to define

astrocytic regions for Kir4.1 quantification, but GFAP labels only ~15% of astrocyte volume

(Bushong et al., 2002), biasing our analysis toward the most GFAP-rich cellular

compartments and underestimating Kir4.1 in fine processes.

Third, our sampling of three circadian timepoints (ZT3, ZT8, ZT15) provides valuable initial

insight into temporal regulation but may miss peak effects occurring at other phases. Higher

temporal resolution studies examining more timepoints throughout the 24-hour cycle would

provide a more complete picture of circadian dynamics.

Despite these limitations, our findings reveal fundamental principles of astrocytic K^+

regulation that likely persist—and may be amplified—in vivo. The regional differences we

document are substantial (2-fold differences in K^+ peak amplitude, 41% difference in Kir4.1

current) and consistent across multiple experimental approaches, suggesting they reflect

robust biological properties rather than experimental artifacts.

Future directions and therapeutic implications

Our findings open several important avenues for future investigation. First, determining the

molecular mechanisms underlying circadian regulation of Kir4.1 function—including

identification of post-translational modifications, trafficking pathways, and regulatory

proteins that differ between DH and VH—could reveal novel targets for modulating

astrocytic K^+ buffering. Second, extending these studies to in vivo preparations using

genetically encoded K^+ sensors and astrocyte-specific manipulations would clarify how the

slice-based findings translate to intact circuits and behavior. Third, investigating whether

similar regional and temporal heterogeneity exists in human hippocampus and whether

these properties are altered in epilepsy could open the way to novel therapeutic avenues.

From a therapeutic perspective, our findings suggest that strategies to enhance astrocytic

K^+ buffering might need to be both regionally and temporally targeted. Global upregulation

of Kir4.1 expression throughout the brain could have unintended consequences in regions

where current expression levels are functionally appropriate. Similarly, chronotherapeutic

approaches that time interventions to periods of maximum vulnerability—when regional

differences in buffering capacity are most pronounced—may prove more effective than

continuous treatment.

Gene therapy approaches to selectively enhance Kir4.1 expression in VH astrocytes

represent a particularly promising avenue. Adeno-associated viral vectors with astrocyte-

specific promoters (Tyurikova et al., 2025) could deliver Kir4.1 cDNA to VH, potentially

normalizing its K^+ buffering capacity to DH-like levels and reducing seizure susceptibility.

Alternatively, small molecules that enhance Kir4.1 channel open probability or trafficking to

the membrane could achieve similar therapeutic effects through pharmacological means.

Conclusion

This study establishes that astrocytic K^+ buffering in the hippocampus is neither spatially

uniform nor temporally static but instead exhibits regional and circadian heterogeneity.

Ventral hippocampal astrocytes possess intrinsically reduced Kir4.1-mediated K^+

conductance, resulting in rapid K^+ accumulation during neuronal activity that is partially

compensated by putative enhanced Na^+/K^+-ATPase activity and expanded gap junction

coupling. These regional specializations do not simply mirror neuronal excitability patterns

but may actively shape them, creating distinct computational regimes in dorsal versus

ventral hippocampus. Superimposed on this stable regional identity is a circadian layer of

regulation that modulates—but does not abolish—regional differences, with maximal

divergence during early light phase.

Our findings support the view that astrocytes as active, heterogeneous participants in circuit

function whose homeostatic properties are dynamically tuned in space and time. This has

implications extending beyond the hippocampus to other brain regions and beyond K^+

buffering to other astrocytic functions. By revealing how regional and temporal factors

converge to determine astrocytic K^+ regulation, this work provides a mechanistic framework

for understanding hippocampal circuit dynamics in health and offers potentially new

therapeutic strategies for epilepsy based on targeted modulation of astrocytic function at

specific times and locations.

Figure Captions

Figure 1. Network level extracellular K+ recording

A) Example of anatomical structures of dorsal (DH: red) and ventral (VH: blue) hippocampal slices.

B) Schematic arrangement of stimulation of Schaffer Collateral (SC), recording of local field potential (LFP) from the *stratum oriens* (*s.o.*), and K+ recording $[K^+]_o$ from *stratum radiatum* (*s.r.*). Example trace of $[K^+]_o$ in DH (red) and VH (blue) in response to 10Hz30s stimulation.

C-**G**) Normalized LFP integral, 25-75% rise time, peak amplitude, tau decay of monoexponentially fitted, and undershoot amplitude across DH and VH in three ZT3, ZT8, and ZT15

Analyses, based on a lognormal distribution model, are reported as posterior median estimates with 95% credible intervals.

* $P < 0.05$, **$P < 0.01$. Statistical significance was calculated using two-sided unpaired Mann–Whitney U test.

[DH_ZT3: n =16 (N = 8) , DH_ZT8: n =16 (N= 8), DH_ZT5: n =19 (N = 10), VH_ZT3: n =16 (N=8), VH_ZT8: n =18 (N= 9), VH_ZT5: n =19 (N=10)]

Figure 2. Gap junction coupling across hippocampal regions and time of day

A) Inverted example image of the coupled astrocytes in a hippocampal slice, as obtained after biocytin diffusion from a single cell, held in whole-cell voltage clamp for 20 mins. Scale bar is equal to 50 μm.

821 **B**) Summary of the number of biocytin-coupled cell across DH and VH in different time

822 points (ZT3, 8, 15). Analyses, based on a lognormal distribution model, are reported as

823 posterior median estimates with 95% credible intervals. *p < 0.05 Mann–Whitney U test;

824 *s.p., Stratum pyramidale*; *s.r., stratum radiatum*.

825 [DH_ZT3: n =11 (N = 6) , DH_ZT8: n =15 (N= 6), DH_ZT5: n =11 (N = 6), VH_ZT3: n =12 (N=7),

826 VH_ZT8: n =11 (N= 7), VH_ZT5: n =9 (N=6)]

827 **Figure 3. Regional differences across regions and time**

828 **A**) Wide-field image of stained astrocytes with SR101.

829 **B**) Left: example image of biocytin stained astrocytes in absence of MFA versus incubated in

830 MFA (right). *S.r., stratum radiatum; s.l.m., stratum lacunosum moleculare*. Scale bar is

831 equal to 20 μm.

832 **C**) statistical summary of the coupled cell number in both groups. Addition of MFA results in

833 an average reduction of -86 [95%CI -95, -72], *p* < .00001.

834 **D**) Initial resting membrane potential (RMP; in mV) of the patched astrocytes across DH and

835 VH in ZT3, ZT8, and ZT15.

836 **E**) Ba^{2+}-induced depolarization across the regions and time. No significant effect of region

837 and time was observed, except in an isolated incident of ZT8.

838 **F**) Ba^{2+}-sensitive current obtained by subtracting membrane currents elicited in whole-cell

839 configuration (from -140 mV to +50 mV, 10 mV increment; holding potential = −70 mV) from

840 the baseline: DH (red) and VH (blue). The rectification of measurements are based on the

barium-sensitive currents. The IV curve slope of -140 to -100mV interval (green) was divided

to the slope of 10mV to 50mV (magenta), and rectification profile of the tested cell was

determined given the ratios.

G)) Kir4.1 current across the regions and time of day. Significant main effect of region was

observed without the influence of time.

Analyses are reported as posterior median estimates with 95% credible intervals.

H) Proportion of rectifying astrocytes (blue) versus non-rectifying (orange) across DH and

VH.

[DH_ZT3: n = 9 (N = 7) , DH_ZT8: n = 5 (N= 3), DH_ZT5: n =8 (N = 6), VH_ZT3: n =9 (N=6),

VH_ZT8: n =8 (N= 5), VH_ZT5: n =10 (N=5)]

* $P < 0.05$, ***$P < 0.001$. Statistical significance was calculated using two-sided unpaired

Mann–Whitney U test

Figure 4. Regional differences in Kir4.1 expression along the dorsoventral axis depends

on time

A) Representative examples of confocal images of immunofluorescent labeling of Kir4.1

(red) and astrocyte marker glial fibrillary acidic protein (GFAP, green) , and DAPI (blue) for

DH (Left) and VH (Right), at 20x magnification (scale bar = 100 µm).

B) Higher magnification of confocal images of Kir4.1 (red) and astrocyte marker glial fibrillary

acidic protein (GFAP, green), and the merging of the two. Images were taken from the s.r.

C) Fraction of GFAP positive area with Kir4.1 protein is significantly smaller in DH -1.72%

[95%CI -2.6%, -0.88%], and this effect is driven by ZT8 -2.21% [95%CI -3.12%, -1.11%].

D) Quantification of Kir4.1 integrated density normalized to GFAP. The relationship was only

significant in ZT3, where VH showed -46% (less) than DH [95%CI -74%, -19%].

Analyses, based on a student t-distribution model, are reported as posterior median

estimates with 95% credible intervals.

* *P* < 0.05. Statistical significance was calculated using two-sided unpaired Mann–Whitney

U test

[DH_ZT3: n = 11 (N = 1) , DH_ZT8: n = 16 (N= 1), DH_ZT5: n =9 (N = 1), VH_ZT3: n =12 (N=1),

VH_ZT8: n =15 (N= 1), VH_ZT5: n =11 (N=1)]

Figure 5. Graphical summary of the results

References

Anderson, T. R., Huguenard, J. R., & Prince, D. A. (2010). Differential effects of Na+–K+ ATPase blockade on cortical layer V neurons. *The Journal of Physiology*, *588*(22), 4401–4414. https://doi.org/10.1113/jphysiol.2010.191858

Anderson, U. A., Carson, C., & McCloskey, K. D. (2009). KCNQ currents and their contribution to resting membrane potential and the excitability of interstitial cells of Cajal from the guinea pig bladder. *The Journal of Urology*, *182*(1), 330–336. https://doi.org/10.1016/j.juro.2009.02.108

Armbruster, M., Naskar, S., Garcia, J. P., Sommer, M., Kim, E., Adam, Y., Haydon, P. G., Boyden, E. S., Cohen, A. E., & Dulla, C. G. (2022). Neuronal activity drives pathway-specific depolarization of peripheral astrocyte processes. *Nature Neuroscience*, *25*(5), Article 5. https://doi.org/10.1038/s41593-022-01049-x

Babb, T. L., Brown, W. J., Pretorius, J., Davenport, C., Lieb, J. P., & Crandall, P. H. (1984). Temporal Lobe Volumetric Cell Densities in Temporal Lobe Epilepsy. *Epilepsia*, *25*(6), 729–740. https://doi.org/10.1111/j.1528-1157.1984.tb03484.x

Bataveljic, D., Pivonkova, H., de Concini, V., Hébert, B., Ezan, P., Briault, S., Bemelmans, A.-P., Pichon, J., Menuet, A., & Rouach, N. (2024). Astroglial Kir4.1 potassium channel deficit drives neuronal hyperexcitability and behavioral defects in Fragile X syndrome mouse model. *Nature Communications*, *15*, 3583. https://doi.org/10.1038/s41467-024-47681-y

Bernasconi, N., Bernasconi, A., Caramanos, Z., Antel, S. B., Andermann, F., & Arnold, D. L. (2003). Mesial temporal damage in temporal lobe epilepsy: A volumetric MRI study

of the hippocampus, amygdala and parahippocampal region. *Brain*, *126*(2), 462–469. https://doi.org/10.1093/brain/awg034

Brancati, G. E., Rawas, C., Ghestem, A., Bernard, C., & Ivanov, A. I. (2021). Spatio-temporal heterogeneity in hippocampal metabolism in control and epilepsy conditions. *Proceedings of the National Academy of Sciences*, *118*(11), e2013972118. https://doi.org/10.1073/pnas.2013972118

Breithausen, B., Kautzmann, S., Boehlen, A., Steinhäuser, C., & Henneberger, C. (2020). Limited contribution of astroglial gap junction coupling to buffering of extracellular K$^+$ in CA1 stratum radiatum. *Glia*, *68*(5), 918–931. https://doi.org/10.1002/glia.23751

Buckmaster, P. S., Reyes, B., Kahn, T., & Wyeth, M. (2022). Ventral Hippocampal Formation Is the Primary Epileptogenic Zone in a Rat Model of Temporal Lobe Epilepsy. *Journal of Neuroscience*, *42*(39), 7482–7495. https://doi.org/10.1523/JNEUROSCI.0429-22.2022

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, *80*, 1–28. https://doi.org/10.18637/jss.v080.i01

Bushong, E. A., Martone, M. E., Jones, Y. Z., & Ellisman, M. H. (2002). Protoplasmic Astrocytes in CA1 Stratum Radiatum Occupy Separate Anatomical Domains. *The Journal of Neuroscience*, *22*(1), 183–192. https://doi.org/10.1523/JNEUROSCI.22-01-00183.2002

Chang, B.-L., Leite, M., Snowball, A., Lieb, A., Chabrol, E., Walker, M. C., Kullmann, D. M., Schorge, S., & Wykes, R. C. (2018). Semiology, clustering, periodicity and natural

history of seizures in an experimental occipital cortical epilepsy model. *Disease*

Models & Mechanisms, *11*(12), dmm036194. https://doi.org/10.1242/dmm.036194

Chang, W.-C., Kudlacek, J., Hlinka, J., Chvojka, J., Hadrava, M., Kumpost, V., Powell, A. D.,

Janca, R., Maturana, M. I., Karoly, P. J., Freestone, D. R., Cook, M. J., Palus, M.,

Otahal, J., Jefferys, J. G. R., & Jiruska, P. (2018). Loss of neuronal network resilience

precedes seizures and determines the ictogenic nature of interictal synaptic

perturbations. *Nature Neuroscience*, *21*(12), 1742–1752.

https://doi.org/10.1038/s41593-018-0278-y

Chever, O., Djukic, B., McCarthy, K. D., & Amzica, F. (2010). Implication of Kir4.1 Channel

in Excess Potassium Clearance: An In Vivo Study on Anesthetized Glial-Conditional

Kir4.1 Knock-Out Mice. *Journal of Neuroscience*, *30*(47), 15769–15777.

https://doi.org/10.1523/JNEUROSCI.2078-10.2010

Cooper, M. L., Selles, M. C., Cammer, M., Gildea, H. K., Sall, J., Chiurri, K. E., Saab, A. S.,

Liddelow, S. A., & Chao, M. V. (2025). Astrocytes connect specific brain regions

through plastic gap junctional networks. *bioRxiv: The Preprint Server for Biology*,

2025.07.18.665573. https://doi.org/10.1101/2025.07.18.665573

D'Ambrosio, R., Gordon, D. S., & Winn, H. R. (2002). Differential role of KIR channel and

Na(+)/K(+)-pump in the regulation of extracellular K(+) in rat hippocampus. *Journal*

of Neurophysiology, *87*(1), 87–102. https://doi.org/10.1152/jn.00240.2001

de Curtis, M., Uva, L., Gnatkovsky, V., & Librizzi, L. (2018). Potassium dynamics and

seizures: Why is potassium ictogenic? *Epilepsy Research*, *143*, 50–59.

https://doi.org/10.1016/j.eplepsyres.2018.04.005

Debski, K. J., Ceglia, N., Ghestem, A., Ivanov, A. I., Brancati, G. E., Bröer, S., Bot, A. M.,

Müller, J. A., Schoch, S., Becker, A., Löscher, W., Guye, M., Sassone-Corsi, P.,

Lukasiuk, K., Baldi, P., & Bernard, C. (2020). The circadian dynamics of the

hippocampal transcriptome and proteome is altered in experimental temporal lobe

epilepsy. *Science Advances*, *6*(41), eaat5979.

https://doi.org/10.1126/sciadv.aat5979

Delignette-Muller, M. L., & Dutang, C. (2015). fitdistrplus: An R Package for Fitting

Distributions. *Journal of Statistical Software*, *64*, 1–34.

https://doi.org/10.18637/jss.v064.i04

Djukic, B., Casper, K. B., Philpot, B. D., Chin, L.-S., & McCarthy, K. D. (2007). Conditional

Knock-Out of K_{ir} 4.1 Leads to Glial Membrane Depolarization, Inhibition of

Potassium and Glutamate Uptake, and Enhanced Short-Term Synaptic

Potentiation. *The Journal of Neuroscience*, *27*(42), 11354–11365.

https://doi.org/10.1523/JNEUROSCI.0723-07.2007

Dong, H.-W., Swanson, L. W., Chen, L., Fanselow, M. S., & Toga, A. W. (2009). Genomic–

anatomic evidence for distinct functional domains in hippocampal field CA1.

Proceedings of the National Academy of Sciences, *106*(28), 11794–11799.

https://doi.org/10.1073/pnas.0812608106

Dospinescu, V.-M., Mascarenhas, A., Butler, J., Nijjar, S., Taborda, K. de O., Connors, S.,

Huang, L., & Dale, N. (2025). CO2-dependent opening of Connexin 43

hemichannels. *eLife*, *14*. https://doi.org/10.7554/eLife.105989.2

Dougherty, K. A., Islam, T., & Johnston, D. (2012). Intrinsic excitability of CA1 pyramidal

neurones from the rat dorsal and ventral hippocampus. *The Journal of Physiology*,

590(Pt 22), 5707–5722. https://doi.org/10.1113/jphysiol.2012.242693

Enger, R., Tang, W., Vindedal, G. F., Jensen, V., Johannes Helm, P., Sprengel, R., Looger, L.

L., & Nagelhus, E. A. (2015). Dynamics of Ionic Shifts in Cortical Spreading

Depression. *Cerebral Cortex (New York, NY)*, *25*(11), 4469–4476.

https://doi.org/10.1093/cercor/bhv054

Erecińska, M., & Silver, I. A. (1994). Ions and energy in mammalian brain. *Progress in*

Neurobiology, *43*(1), 37–71. https://doi.org/10.1016/0301-0082(94)90015-9

Fanselow, M. S., & Dong, H.-W. (2010). Are the Dorsal and Ventral Hippocampus

Functionally Distinct Structures? *Neuron*, *65*(1), 7–19.

https://doi.org/10.1016/j.neuron.2009.11.031

Heinemann, U., & Arens, J. (1992). Production and calibration of ion-sensitive

microelectrodes. *Practical Electrophysiological Methods*, 206–212.

Hibino, H., Inanobe, A., Furutani, K., Murakami, S., Findlay, I., & Kurachi, Y. (2010).

Inwardly rectifying potassium channels: Their structure, function, and physiological

roles. *Physiological Reviews*, *90*(1), 291–366.

https://doi.org/10.1152/physrev.00021.2009

Ho, J., Tumkaya, T., Aryal, S., Choi, H., & Claridge-Chang, A. (2019). Moving beyond P

values: Data analysis with estimation graphics. *Nature Methods*, *16*(7), 565–566.

https://doi.org/10.1038/s41592-019-0470-3

Jinno, S. (2011). Regional and laminar differences in antigen profiles and spatial

distributions of astrocytes in the mouse hippocampus, with reference to aging.

Neuroscience, *180*, 41–52. https://doi.org/10.1016/j.neuroscience.2011.02.013

Khakh, B. S., & Deneen, B. (2019). The Emerging Nature of Astrocyte Diversity. *Annual*

Review of Neuroscience, *42*, 187–207. https://doi.org/10.1146/annurev-neuro-

070918-050443

Kherad-Pajouh, S., & Renaud, O. (2015). A general permutation approach for analyzing

repeated measures ANOVA and mixed-model designs. *Statistical Papers*, *56*(4),

947–967. https://doi.org/10.1007/s00362-014-0617-3

Larsen, B. R., Assentoft, M., Cotrina, M. L., Hua, S. Z., Nedergaard, M., Kaila, K., Voipio, J.,

& MacAulay, N. (2014). Contributions of the Na^+/K^+-ATPase, NKCC1, and Kir4.1 to

hippocampal K^+ clearance and volume responses: Mechanisms of K^+ Clearance in

the Brain. *Glia*, *62*(4), 608–622. https://doi.org/10.1002/glia.22629

Larsen, B. R., & MacAulay, N. (2014). Kir4.1-mediated spatial buffering of K+: Experimental

challenges in determination of its temporal and quantitative contribution to K+

clearance in the brain. *Channels*, *8*(6), 544–550.

https://doi.org/10.4161/19336950.2014.970448

Lu, W., & Haber, S. N. (1992). In situ hybridization histochemistry: A new method for

processing material stored for several years. *Brain Research*, *578*(1), 155–160.

https://doi.org/10.1016/0006-8993(92)90243-3

MacAulay, N. (2020). Molecular mechanisms of K+ clearance and extracellular space

shrinkage—Glia cells as the stars. *Glia*, *68*(11), 2192–2211.

https://doi.org/10.1002/glia.23824

McCauley, J. P., Petroccione, M. A., D'Brant, L. Y., Todd, G. C., Affinnih, N., Wisnoski, J. J.,

Zahid, S., Shree, S., Sousa, A. A., De Guzman, R. M., Migliore, R., Brazhe, A.,

Leapman, R. D., Khmaladze, A., Semyanov, A., Zuloaga, D. G., Migliore, M., &

Scimemi, A. (2020). Circadian Modulation of Neurons and Astrocytes Controls

Synaptic Plasticity in Hippocampal Area CA1. *Cell Reports*, *33*(2), 108255.

https://doi.org/10.1016/j.celrep.2020.108255

Meeks, J. P., & Mennerick, S. (2004). Selective Effects of Potassium Elevations on

Glutamate Signaling and Action Potential Conduction in Hippocampus. *The Journal*

of Neuroscience, *24*(1), 197–206. https://doi.org/10.1523/JNEUROSCI.4845-

03.2004

Moreno, A., Morris, R. G. M., & Canals, S. (2016). Frequency-Dependent Gating of

Hippocampal-Neocortical Interactions. *Cerebral Cortex (New York, N.Y.: 1991)*,

26(5), 2105–2114. https://doi.org/10.1093/cercor/bhv033

Nimmerjahn, A., Kirchhoff, F., Kerr, J. N. D., & Helmchen, F. (2004). Sulforhodamine 101 as

a specific marker of astroglia in the neocortex in vivo. *Nature Methods*, *1*(1), 31–37.

https://doi.org/10.1038/nmeth706

Olsen, M. L., & Sontheimer, H. (2008). Functional implications for Kir4.1 channels in glial

biology: From K+ buffering to cell differentiation. *Journal of Neurochemistry*, *107*(3),

589–601. https://doi.org/10.1111/j.1471-4159.2008.05615.x

Papaleonidopoulos, V., Trompoukis, G., Koutsoumpa, A., & Papatheodoropoulos, C.

(2017). A gradient of frequency-dependent synaptic properties along the

longitudinal hippocampal axis. *BMC Neuroscience*, *18*, 79.

https://doi.org/10.1186/s12868-017-0398-4

Papatheodoropoulos, C., Moschovos, C., & Kostopoulos, G. (2005). Greater contribution

of N-methyl-d-aspartic acid receptors in ventral compared to dorsal hippocampal

slices in the expression and long-term maintenance of epileptiform activity.

Neuroscience, *135*(3), 765–779.

https://doi.org/10.1016/j.neuroscience.2005.06.024

Peretz, A., Degani, N., Nachman, R., Uziyel, Y., Gibor, G., Shabat, D., & Attali, B. (2005).

Meclofenamic Acid and Diclofenac, Novel Templates of KCNQ2/Q3 Potassium

Channel Openers, Depress Cortical Neuron Activity and Exhibit Anticonvulsant

Properties. *Molecular Pharmacology*, *67*(4), 1053–1066.

https://doi.org/10.1124/mol.104.007112

Proix, T., Truccolo, W., Leguia, M. G., Tcheng, T. K., King-Stephens, D., Rao, V. R., & Baud,

M. O. (2021). Forecasting seizure risk in adults with focal epilepsy: A development

and validation study. *The Lancet Neurology*, *20*(2), 127–135.

https://doi.org/10.1016/S1474-4422(20)30396-3

Pronier, É., Morici, J. F., & Girardeau, G. (2023). The role of the hippocampus in the

consolidation of emotional memories during sleep. *Trends in Neurosciences*,

46(11), 912–925. https://doi.org/10.1016/j.tins.2023.08.003

Quigg, M., Clayburn, H., Straume, M., Menaker, M., & Bertram III, E. H. (2000). Effects of

Circadian Regulation and Rest—Activity State on Spontaneous Seizures in a Rat

Model of Limbic Epilepsy. *Epilepsia*, *41*(5), 502–509. https://doi.org/10.1111/j.1528-

1157.2000.tb00202.x

Romanos, J., Benke, D., Pietrobon, D., Zeilhofer, H. U., & Santello, M. (2020). Astrocyte

dysfunction increases cortical dendritic excitability and promotes cranial pain in

familial migraine. *Science Advances*, *6*(23), eaaz1584.

https://doi.org/10.1126/sciadv.aaz1584

Ryu, J. E., Shim, K.-W., Roh, H. W., Park, M., Lee, J.-H., & Kim, E. Y. (2024). Circadian

regulation of endoplasmic reticulum calcium response in mouse cultured

astrocytes. *eLife*, *13*. https://doi.org/10.7554/eLife.96357.1

Schlingloff, D., Káli, S., Freund, T. F., Hájos, N., & Gulyás, A. I. (2014). Mechanisms of Sharp

Wave Initiation and Ripple Generation. *The Journal of Neuroscience*, *34*(34), 11385–

11398. https://doi.org/10.1523/JNEUROSCI.0867-14.2014

Seifert, G., Huttmann, K., Binder, D. K., Hartmann, C., Wyczynski, A., Neusch, C., &

Steinhauser, C. (2009). Analysis of Astroglial K+ Channel Expression in the

Developing Hippocampus Reveals a Predominant Role of the Kir4.1 Subunit. *Journal*

of Neuroscience, *29*(23), 7474–7488. https://doi.org/10.1523/JNEUROSCI.3790-

08.2009

Shih, P.-Y., Savtchenko, L. P., Kamasawa, N., Dembitskaya, Y., McHugh, T. J., Rusakov, D.

A., Shigemoto, R., & Semyanov, A. (2013). Retrograde Synaptic Signaling Mediated

1063   by K+ Efflux through Postsynaptic NMDA Receptors. *Cell Reports*, *5*(4), 941–951.

1064   https://doi.org/10.1016/j.celrep.2013.10.026

1065 Stephan, J., Eitelmann, S., & Zhou, M. (2021). Approaches to Study Gap Junctional

1066   Coupling. *Frontiers in Cellular Neuroscience*, *15*, 640406.

1067   https://doi.org/10.3389/fncel.2021.640406

1068 Sun, Q., Jiang, Y.-Q., & Lu, M. C. (2020). Topographic heterogeneity of intrinsic excitability

1069   in mouse hippocampal CA3 pyramidal neurons. *Journal of Neurophysiology*, *124*(4),

1070   1270–1284. https://doi.org/10.1152/jn.00147.2020

1071 Thompson, C. L., Pathak, S. D., Jeromin, A., Ng, L. L., MacPherson, C. R., Mortrud, M. T.,

1072   Cusick, A., Riley, Z. L., Sunkin, S. M., Bernard, A., Puchalski, R. B., Gage, F. H.,

1073   Jones, A. R., Bajic, V. B., Hawrylycz, M. J., & Lein, E. S. (2008). Genomic anatomy of

1074   the hippocampus. *Neuron*, *60*(6), 1010–1021.

1075   https://doi.org/10.1016/j.neuron.2008.12.008

1076 Tovar, K. R., Maher, B. J., & Westbrook, G. L. (2009). Direct Actions of Carbenoxolone on

1077   Synaptic Transmission and Neuronal Membrane Properties. *Journal of*

1078   *Neurophysiology*, *102*(2), 974–978. https://doi.org/10.1152/jn.00060.2009

1079 Toyoda, I., Bower, M. R., Leyva, F., & Buckmaster, P. S. (2013). Early Activation of Ventral

1080   Hippocampus and Subiculum during Spontaneous Seizures in a Rat Model of

1081   Temporal Lobe Epilepsy. *The Journal of Neuroscience*, *33*(27), 11100–11115.

1082   https://doi.org/10.1523/JNEUROSCI.0472-13.2013

1083 Tyurikova, O., Kopach, O., Zheng, K., Rathore, D., Codadu, N., Wu, S.-Y., Shen, Y.,

1084   Campbell, R. E., Wykes, R. C., Volynski, K., Savtchenko, L. P., & Rusakov, D. A.

1085   (2025). Astrocyte Kir4.1 expression level territorially controls excitatory

1086   transmission in the brain. *Cell Reports*, *44*(2), 115299.

1087   https://doi.org/10.1016/j.celrep.2025.115299

1088 Wallraff, A., Köhling, R., Heinemann, U., Theis, M., Willecke, K., & Steinhäuser, C. (2006).

1089   The Impact of Astrocytic Gap Junctional Coupling on Potassium Buffering in the

1090   Hippocampus. *The Journal of Neuroscience*, *26*(20), 5438–5447.

1091   https://doi.org/10.1523/JNEUROSCI.0037-06.2006

1092 Walz, W. (2000). Role of astrocytes in the clearance of excess extracellular potassium.

1093   *Neurochemistry International*, *36*(4), 291–300. https://doi.org/10.1016/S0197-

1094   0186(99)00137-0

1095 Wang, F., Smith, N. A., Xu, Q., Fujita, T., Baba, A., Matsuda, T., Takano, T., Bekar, L., &

1096   Nedergaard, M. (2012). Astrocytes Modulate Neural Network Activity by Ca2+-

1097   Dependent Uptake of Extracellular K+. *Science Signaling*, *5*(218), ra26.

1098   https://doi.org/10.1126/scisignal.2002334

1099 Watson, R. E., Wiegand, S. J., Clough, R. W., & Hoffman, G. E. (1986). Use of cryoprotectant

1100   to maintain long-term peptide immunoreactivity and tissue morphology. *Peptides*,

1101   *7*(1), 155–159. https://doi.org/10.1016/0196-9781(86)90076-8

1102 Zhou, M., Du, Y., Aten, S., & Terman, D. (2021). On the electrical passivity of astrocyte

1103   potassium conductance. *Journal of Neurophysiology*, *126*(4), 1403–1419.

1104   https://doi.org/10.1152/jn.00330.2021

1105

 Figure 1

1108   Figure 2

1109

1110

1111   Figure 3

1112   Figure 4

A

B

C

 D

1113

Time (ZT) | Dorsal astrocytes | Evoked [K+]$_O$ | Ventral astrocytes

Kir4.1 Channnels   Gap Junctions   Potassium Ion