Article

Somatostatin-expressing neurons regulate sleep deprivation and recovery

Kenta Kobayashi [1] and **Y-h. Taguchi** [2]*

1   Department of Physics, Graduate School of Science and Engineering, Chuo University, 1-13-27 Kasuga, Bunkyo-ku, Tokyo, 112-8551, Japan; vv8vv8kenta@gmail.com
2   Department of Physics, Chuo University, 1-13-27 Kasuga, Bunkyo-ku, Tokyo, 112-8551, Japan;tag@granular.com
*   Correspondence: tag@granular.com; Tel.: +81-3-3817-1791 (F.L.)

Abstract

We re-analyzed gene expression profiles from the male mouse cortex under conditions of sleep deprivation using tensor decomposition-based unsupervised feature extraction. The analysis revealed that the selected genes were enriched in various brain regions, as identified through clustering in the Allen Brain Atlas. Somatostatin (Sst)-expressing neuronal clusters were among the most highly enriched. These finding suggests that Sst-expressing neurons may play a key role in sleep deprivation.

Keywords: Somatostatin; Sleep deprivation; Tensor Decomposition; Sleep debt;

1. Introduction

Sleep deprivation is believed to increase sleep needs, which is reduced by subsequent recovery sleep. Popescu et al. [1] re-analyzed gene expression profiles and observed that various pathways behaved as expected; that is, gene expression was altered during sleep deprivation and returned to baseline following recovery sleep.

Although they hypothesized that recovery sleep alleviates the sleep pressure induced by sleep deprivation, they proved that this hypothesis-driven approach is limited in its ability to fully exploit large datasets, as it may overlook unexpected findings. In contrast, a data-driven approach can determine what has happened in the dataset, that is, the process between sleep deprivation and sleep recovery.

In this study, we aimed to re-analyze the dataset from Popescu et al. using a fully data-driven approach, tensor decomposition (TD)-based unsupervised feature extraction (FE) [2]. We observed two patterns of dependence: home cage (control) vs. sleep deprivation vs. recovery sleep. These patterns were expressed as singular value vectors obtained using TD. Genes whose expression correlated with these two patterns were identified and observed that they were enriched in various biological processes, neuronal categories (based on Allen Brain Atlas), miRNA target genes, and histone modifications. It is strongly associated with the somatostatin (Sst) cell cluster in the Allen Brain Atlas, which is related to sleep regulation.

2. Materials and Methods

2.1. Gene expression profiles

The gene expression profiles analyzed in this study were downloaded from Gene Expression Omnibus (GEO) using the GEO ID GSE237419 [1]. The file GSE237419_gse_HCSDRS

_WT_SleepIntegration_salmon.txt was retrieved and loaded into R using the read.csv command.

2.2. Tensor

Gene expression profiles are formatted as a tensor $x_{ijkm} \in \mathbb{R}^{N \times 3 \times 5 \times 5}$ which represents the expression of ith gene of jth condition ($j = 1$: home cage (HC), $j = 2$: sleep deprivation (SD), and $j = 3$: recovery sleep (RS)) of kth replicate at mth hour. ($m = 1$: three hours, $m = 2$: five hours, $m = 3$: six hours, $m = 4$: seven hours, and $m = 5$: 12 h), where $N = 54,308$ denotes the total number of genes. As not all combinations of i, j, k, and m were measured, the missing values were filled with zeros (Table 1).

Table 1. Correspondence between indices, j, k, and m, of the tensor, x_{ijkm}, and experimental conditions. When the number of replicates is not five but three, only x_{ijkm} with $2 \le k \le 4$ have values. x_{ijkm}s whose $(j, k.m)$s are not listed are filled with zero. HC: Home cage (control), SD: Sleep deprivation, and RS: Recovery sleep. The numbers adjusted to two alphabets, HC, SD, and RS, represent hours. For more details, see [1]

m	$j = 1$	$j = 2$	$j = 3$
1	HC3 ($2 \le k \le 4$)	SD3 ($2 \le k \le 4$)	—
2	HC5 ($1 \le k \le 5$)	SD5 ($1 \le k \le 5$)	RS2 ($1 \le k \le 5$)
3	HC6 ($1 \le k \le 5$)	SD6 ($2 \le k \le 4$)	RS6 ($2 \le k \le 4$)
4	HC7 ($1 \le k \le 5$)	—	—
5	HC12 ($1 \le k \le 5$)	—	—

2.3. Tensor Decomposition

Before applying TD to the obtained tensor, x_{ijkm} and x_{ijkm} are standardized as follows:

$$\sum_{i=1}^{N} x_{ijkm} = 0 \tag{1}$$

$$\sum_{i=1}^{N} x_{ijkm}^2 = N \tag{2}$$

After applying HOSVD [2] to x_{ijkm}, we get

$$x_{ijkm} = \sum_{\ell_1=1}^{N} \sum_{\ell_2=1}^{3} \sum_{\ell_3=1}^{5} \sum_{\ell_4=1}^{5} G(\ell_1 \ell_2 \ell_3 \ell_4) u_{\ell_1 i} u_{\ell_2 j} u_{\ell_3 k} u_{\ell_4 m} \tag{3}$$

where $G \in \mathbb{R}^{N \times 3 \times 3 \times 5}$ denotes the core tensor representing the contribution of the $u_{\ell_1 i} u_{\ell_2 j} u_{\ell_3 k} u_{\ell_4 m}$ to x_{ijkm}, $u_{\ell_1 i} \in \mathbb{R}^{N \times N}$, $u_{\ell_2 j} \in \mathbb{R}^{3 \times 3}$, $u_{\ell_3 k} \in \mathbb{R}^{5 \times 5}$, and $u_{\ell_4 m} \in \mathbb{R}^{5 \times 5}$ are the singular value matrices, and orthogonal matrices.

2.4. Gene selection

Although the procedure was described in detail in a recent book [2], it is briefly explained below. To select $u_{\ell_1 i}$ for gene selection, we need to know which $u_{\ell_1 j}$, $u_{\ell_3 k}$, and $u_{\ell_4 m}$ are of interest, as follows:

- Since the gene expression is ought to be distinct between controls and others, u_{1j} should be distinct from u_{2j} and/or u_{2j}.
- Since the gene expression should not be distinct between replicates, $u_{\ell_3 k}$ should be constant as well as possible.
- Since we do not have any pre-judges dependence upon time, we do not specially mind the dependence of $u_{\ell_4 m}$ upon m.

After identifying ℓ_2, ℓ_3, and ℓ_4 of interest, we investigate $G(\ell_1\ell_2\ell_3\ell_4)$ by fixing ℓ_2. ℓ_3 and ℓ_4 to the selected ones, and select ℓ_1 whose associated G has the largest absolute value.

*P*values are assigned to *i*s by assuming the null hypothesis that $u_{\ell_1 i}$ is Gaussian as

$$P_i = P_{\chi^2}\left[> \left(\frac{u_{\ell_1 i}}{\sigma_{\ell_1}} \right)^2 \right] \tag{4}$$

where $P_{\chi^2}[> x]$ is the cumulative Gaussian distribution, where the argument is larger than x and σ_{ℓ_1} is the standard deviation of $u_{\ell_1 i}$, which is optimized (see below). Finally, genes associated with *P*-values adjusted by the BH criterion [2] to less than 0.01 are selected.

The motivation for optimizing σ_{ℓ_1} is as follows: If we compute σ_{ℓ_1} using all $u_{\ell_1 i}$ including those attributed to *i*s to be selected as outliers, σ_{ℓ_1} is likely to be overestimated and P_i might be overestimated. Using the overestimated P_i, we identify fewer *i*s associated with significant P_is. Thus, it is important to estimate σ_{ℓ_1} excluding i to be selected as outliers.

Suppose that $h_s(1 - P_i)$ is the histogram of *s*th bin ($1 \le s \le S$) of $1 - P_i$ computed by substituting the arbitrary value of σ_{ℓ_1} into eq. (4), excluding i values associated with *P*values adjusted by BH criterion of less than 0.01. We compute the standard deviation of h_s, σ_{h_2} as follows:

$$\langle h_s \rangle = \frac{1}{S} \sum_{s=1}^{S} h_s \tag{5}$$

$$\sigma_{h_s} = \sqrt{\frac{1}{S} \sum_{s=1}^{S} (h_s - \langle h_s \rangle)^2} \tag{6}$$

2.5. AlphaGenome

In this subsection, we summarize how we retrieved features from the AplhaGenome.

2.5.1. Conversion to genomic regions

As AlphaGenome does not accept the list of gene names/IDs, the Ensembl gene IDs of selected genes were converted to genomic regions using BSgenome.Mmusculus.UCSC.mm39 [3].

2.5.2. Upload to AlphaGenome and retrieving features

The genomic regions obtained were uploaded to AlphaGenome using the Python API [4]. DNase-Seq prediction by AlphaGenome of uploaded regions was performed using the Python API.

2.5.3. The selection of top ranked genes

The brain regions considered were "Brain," "left cerebral cortex," "right cerebral cortex," "layer of hippocampus," and "frontal cortex." Comparisons were performed within four pairwise comparisons, i.e., "Brain" vs "left cerebral cortex", "Brain" vs "right cerebral cortex," "Brain" vs "layer of hippocampus," and "Brain" vs "frontal cortex." Logarithmic ratios were computed (10^{-6} was added to the denominator to avoid division by zero), and the top 100 genes with the larger/smaller logarithmic ratios were selected. The 100 top ranked genes were uploaded to Enrichr.

3. Results

3.1. Gene selection

First, we investigated $u_{\ell_2 j}$ to determine the type of the dependence among the HC, SD, and RS conditions (Fig. 1). Contrary to the expectation, no simple "altered in SD and

Figure 1. From left to right: u_{1j}, u_{2j}, and u_{3j}. $j = 1$: HC, control, $j = 2$:SD, and $j = 3$:RS. Red broken lines are baselines.

recovery in RS" pattern is observed but "altered in SD and not recovery in RS" (u_{2j}) and "altered in SD and overshoot in RS" (u_{3j}) patterns are observed. However, u_{1k} (Fig. 2, left

Figure 2. Left: u_{1k} and right u_{1m}. Red broken lines are baselines.

panel) and u_{1m} (Fig. 2, right panel) do not express dependence on k and m . Thus, we sought the largest $G(\ell_1, 2, 1, 1)$ (Fig. 3, left panel) and $G(\ell_1, 3, 1, 1)$ (Fig. 3, right panel) values. We decided to use $\ell_1 = 2, 3$ for the gene selection.

Figure 3. Left: $G(\ell_1, 2, 1, 1)$ and right $G(\ell_1, 3, 1, 1)$. Horizontal axis is $1 \le \ell_1 \le 10$ from left to right.

Next, we attempt to optimize σ_{ℓ_1} for $\ell_1 = 2, 3$ (see Figs. 4). As expected, we successfully

Figure 4. Left column: σ_h as a function of $sigma_{\ell_1}$. Vertical red arrows suggest the selected σ_{ℓ_1}. Right column: The histogram of $1 - P_i$ computed by eq. (4) using the selected σ_{ℓ_1}. Upper: $\ell_1 = 2$. Lower: $\ell_1 = 3$.

optimize σ_{ℓ_1} to minimize σ_h. P_i is computed using Eqs. (4) and adjusted using the BH criterion [2]. As a result, 2,182 for $\ell_1 = 2$ and 2,792 for $\ell_1 = 3$ ensemble genes associated with the corrected P-values less than 0.01. The ensemble genes were converted to 2,081 (for $\ell_1 = 2$) and 2,089 (for $\ell_1 = 3$) gene symbols. Figure 5 shows a Venn diagram of the gene symbols selected by $\ell_1 = 2$ and $\ell_1 = 3$. Despite significant overlap, these two remain distinct.

3.2. Metascape

These gene symbols were uploaded to Metascape [5] to biologically evaluate the selected genes. Table 2 lists the top 20 clusters. These include terms such as "central nervous system neuron axonogenesis," which is consistent with the fact that the samples were obtained from brain tissue, and terms like "autophagy" and "metabolism," which were highlighted in the analysis by Popescu et al. [5]. Furthermore, endoplasmic reticulum stress is known to be induced by sleep loss [6] ("GO:0034976, response to endoplasmic reticulum stress"), sleep loss can alter the structures of various organelles, including mitochondria, nucleus, and Golgi apparatus [7] ("GO:0048193, Golgi vesicle transport"). Additionally, locomotory behavior was identified, which aligns with previous studies

Table 2. Top 20 clusters and their representative enriched terms (one per cluster) identified using Metascape, based on 2,081 (for $\ell_1 = 2$) and 2,089 (for $\ell_1 = 3$) gene symbols uploaded. "Count" refers to the number of user-provided genes associated with a given ontology term. "%" indicates the percentage of all of the user-provided genes that are found in the given ontology term (only input genes with at least one ontology term annotation are included in the calculation). "Log10(P)" is the p-value in log base 10. "Log10(q)" is the multi-test adjusted p-value in log base 10. PC2: $\ell_1 = 2$ and PC3: $\ell_1 = 3$.

PC2	PC3	GO/REAC	Category	Description	Count	%	Log10(P)	Log10(q)
	○	R-MMU-72766	REAC	Translation	65	4.16	-21.75	-17.49
○	○	R-MMU-8953854	REAC	Metabolism of RNA	112	7.17	-21.43	-17.47
○	○	GO:0016072	GO BP	rRNA metabolic process	59	3.77	-15.85	-12.54
○		GO:0061024	GO BP	membrane organization	119	7.61	-15.08	-11.82
	○	GO:0006351	GO BP	DNA-templated transcription	95	6.08	-14.75	-11.53
	○	R-MMU-199991	REAC	Membrane Trafficking	92	5.89	-13.88	-10.80
○	○	GO:0030705	GO BP	cytoskeleton-dependent intracellular transport	50	3.20	-13.72	-10.68
○	○	GO:0006886	GO BP	intracellular protein transport	92	5.89	-13.58	-10.63
○	○	GO:0006914	GO BP	autophagy	68	4.35	-13.57	-10.63
○		GO:0048858	GO BP	cell projection morphogenesis	95	6.08	-13.12	-10.24
○	○	GO:0022411	GO BP	cellular component disassembly	58	3.71	-12.49	-9.67
○	○	R-MMU-6791226	REAC	Major pathway of rRNA processing...	43	2.75	-11.76	-9.04
○		GO:0070848	GO BP	response to growth factor	91	5.82	-10.63	-7.95
○	○	GO:0007626	GO BP	locomotory behavior	55	3.52	-10.60	-7.93
○	○	GO:0048193	GO BP	Golgi vesicle transport	53	3.39	-9.98	-7.34
○		GO:1903829	GO BP	positive regulation of protein localization	84	5.37	-9.62	-7.02
	○	GO:0034976	GO BP	response to endoplasmic reticulum stress	48	3.07	-9.60	-7.01
	○	R-MMU-983799	REAC	Mitochondrial protein degradation	24	1.54	-9.29	-6.76
	○	GO:0021955	GO BP	central nervous system neuron axonogenesis	18	1.15	-9.20	-6.68

Figure 5. Venn diagram between gene symbols selected by $\ell_1 = 2$ and $\ell_1 = 3$, respectively. PC2: $\ell_1 = 2$ and PC3: $\ell_1 = 3$.

discussing motor control mechanisms during sleep [8] ("GO:0007626, locomotory behavior "). A full enrichment analysis, shown in Figure S1, reveals further overlap with Popescu et al.'s findings, including "Wnt signaling pathway," "Notch signaling pathway," and "Mapk signaling pathway." These results indicate that the genes identified are biologically functional.

To further validate the biological relevance of the identified genes, we examined the protein–protein interaction network and MCODE components generated from the gene lists (Figure 6) using Metascape. These results appear biologically plausible, as they include "MTOR signalling" and "Pathways in cancer," which were are reported in the study by Popescu et al. [5] and multiple Rho GTPases related terms, which were a focus of our previous work [9] were detected.

3.3. Enrichr

We have uploaded the selected genes to Enrichr [10] and observed that many enriched cell clusters exist in "Allen Brain Atlas 10x scRNA 2021" category (Figs 7 and 8). We observed that the cell cluster labeled as "Sst," referring to "MGE-derived GABAergic subclass typically expressing Sst," is highly enriched in $\ell_1 = 2$ and $\ell_1 = 3$.

Gabaergic Sst may play a critical role in SD and RS. All brain functions depend on a delicate balance between excitatory and inhibitory signals [11]. In the mature central

Figure 6. Protein–protein interaction network and MCODE components identified in the gene lists. For annotation of cluster, see Tables 3 and 4.

Table 3. Cluster annotation associated with Fig. 6

Color	MCODE	GO	Description	Log10(P)
Red	MCODE_1	mmu05214	Glioma - Mus musculus (house mouse)	-7.8
Red	MCODE_1	WP1763	Mechanisms associated with pluripotency	-7.7
Red	MCODE_1	mmu05200	Pathways in cancer - Mus musculus (house mouse)	-7.2
Blue	MCODE_2	GO:0051123	RNA polymerase II preinitiation complex assembly	-11.7
Blue	MCODE_2	GO:0070897	transaction preinitiation complex assembly	-11.2
Blue	MCODE_2	GO:0060261	positive regulation of transcription initiation by RNA polymerase II	-10.8
Green	MCODE_3	GO:0032543	mitochondrial translation	-22.4
Green	MCODE_3	GO:0140053	mitochondrial gene expression	-20.8
Green	MCODE_3	R-MMU-5419276	Mitochondrial translation termination	-20.7
Purple	MCODE_4	GO:0042254	ribosome biogenesis	-19.5
Purple	MCODE_4	GO:0022613	ribonucleoprotein complex biogenesis	-17.6
Purple	MCODE_4	R-MMU-8868773	rRNA processing in the nucleus and cytosol	-17.5
Orange	MCODE_5	R-MMU-165159	MTOR signalling	-8.7
Orange	MCODE_5	GO:1903432	regulation of TORC1 signaling	-7.0
Orange	MCODE_5	R-MMU-166208	mTORC1-mediated signalling	-6.9
Yellow	MCODE_6	mmu03083	Polycomb repressive complex - Mus musculus (house mouse)	-9.9
Yellow	MCODE_6	R-MMU-8953750	Transcriptional Regulation by E2F6	-7.2
Yellow	MCODE_6	R-MMU-212436	Generic Transcription Pathway	-3.7
Brown	MCODE_7	R-MMU-983169	Class I MHC mediated antigen processing & presentation	-5.7
Brown	MCODE_7	mmu04512	ECM-receptor interaction - Mus musculus (house mouse)	-5.6
Brown	MCODE_7	mmu04120	Ubiquitin mediated proteolysis - Mus musculus (house mouse)	-4.9

Table 4. Cluster annotation associated with Fig. 6

Color	MCODE	GO	Description	Log10(P)
Teal	MCODE_10	GO:0036503	ERAD pathway	-8.7
Teal	MCODE_10	GO:0034976	response to endoplasmic reticulum stress	-7.1
Teal	MCODE_10	GO:0010498	proteasomal protein catabolic process	-5.8
Lavender	MCODE_12	R-MMU-9013424	RHOV GTPase cycle	-8.5
Lavender	MCODE_12	R-MMU-9012999	RHO GTPase cycle	-5.2
Lavender	MCODE_12	R-MMU-194315	Signaling by Rho GTPases	-4.7
Coral	MCODE_13	R-MMU-9006936	Signaling by TGFB family members	-6.9
Coral	MCODE_13	mmu04350	TGF-beta signaling pathway - Mus musculus (house mouse)	-6.9
Steel Blue	MCODE_14	R-MMU-6794361	Neurexins and neuroligins	-8.6
Steel Blue	MCODE_14	R-MMU-6794362	Protein-protein interactions at synapses	-7.6
Steel Blue	MCODE_14	GO:0034329	cell junction assembly	-5.5

nervous system, gamma-aminobutyric acid (GABA)is the main inhibitory neurotransmitter, and plays a vital role in preventing the excessive excitation of neural networks and regulating information transmission [12]. Although GABAergic interneurons account for only approximately 10–20% of all neurons, their diversity and unique connectivity significantly enhance cortical computations and information processing [13]. Dysfunctions in these interneurons are associated with several neuropsychiatric disorders, including epilepsy, schizophrenia, and anxiety.

Among GABAergic interneurons, Sst-expressing neurons form a major class, comprising approximately 30% of cortical interneurons [14]. These neurons co-release the GABA and Sst, potentially enhancing GABAergic inhibitory effects [15]. Anatomically and electrophysiologically, these cells are heterogeneous, including Martinotti cells and low-threshold spiking cells [16]. Sst neurons are widely distributed in brain regions such as the neocortex and hippocampus, which are important for higher cognitive functions and memory [17].

Research on Sst neurons revealed two seemingly contradictory findings. On the one hand, these neurons play an essential role in generating electroencephalogram (EEG) slow-wave oscillations, characteristic of slow-wave sleep, which contributes to deep and restorative sleep [18]. On the other hand, they are crucial for generating brain wave (EEG), a process that is known as "slow-wave sleep". In this context, Sst neurons function as "sleep builders." At the same time, however, they have been shown to be deeply involved in processes that actively interfere with sleep, such as stress-induced insomnia and cognitive dysfunction due to sleep deprivation [19]. In the following section, we aim to answer this central paradox: How can the same types of neurons promoting and

Figure 7. Volcano plot of cell clusters in "Allen Brain Atlas 10x scRNA 2021" category. Red ones are associated with adjusted P-values less than 0.05. PC2: $\ell_1 = 2$ and PC3: $\ell_1 = 3$.

disrupt sleep? The answer lies in the "context-dependence function" of Sst neurons, which is not determined by their neurotransmitters, but by the specific neural circuits they are embedded in and the dynamic state of the brain. This report aims to systematically explain the complex relationship between Sst neurons and sleep regulation by exploring this context dependence.

Non-rapid eye movement (NREM) sleep, particularly in its deepest stage, slow-wave sleep, is essential for brain recovery, memory consolidation, and metabolite elimination [20]. The EEG of this sleep stage is characterized by low-frequency, high-amplitude "slow waves" below 1 Hz [18]. These slow waves are caused by a periodic cycle of "ON states" (periods of depolarization and firing) and "OFF states" (periods of hyperpolarization and silence), in which a large number of cortical neurons are synchronously active and cease to be active altogether [18].

Recent studies have shown that cortical Sst neurons play a central role in triggering synchronous OFF states. Using optogenetics, brief activation of cortical Sst neurons in the ON state induces a long OFF state that persists beyond the stimulation time, thereby silencing a broad population of cortical neurons [18]. This effect is attributed to the ability of Sst neurons, particularly Martinotti cells, to project extensive axons onto cortical layer 1, where the distal dendrites of pyramidal cells are densely located, causing strong and sustained GABA$_B$ receptor-mediated inhibition [14].

Figure 8. The bar chart shows the top 10 enriched terms in the chosen library, along with their corresponding p-values. Colored bars correspond to terms with significant p-values (< 0.05). An asterisk (*) next to a p-value indicates the term also has a significant adjusted p-value (< 0.05). Upper: $\ell_1 = 2$ and lower: $\ell_1 = 3$.

Furthermore, sustained activation of Sst neurons using chemogenetics significantly increases slow wave activity (SWA), steepens the slope of individual slow waves, and lengthens the total duration of NREM sleep. Conversely, the inhibition of these neurons significantly reduces SWA [18]. This function is specific to Sst neurons. For example, the activation of another major interneuron, parvalbumin-expressing neurons, increases NREM sleep duration but significantly decreases SWA, causing an overall inhibition rather than a structured oscillation in cortical activity [21]. These findings suggest that Sst neurons do not merely facilitate sleep but act as "builders" of slow waves that determine restorative sleep quality.

During sleep, Sst neurons play a role in disrupt sleep and impair cognitive function under certain circumstances. This duality indicates that the function of Sst neurons is strongly dependent on the brain region and state.

One example is its role in the hippocampus during sleep deprivation. Sleep deprivation has been shown to selectively activates Sst interneurons in the hippocampus1. This

activation acts as a "gate" that inhibits activity in the dentate gyrus, which is essential for memory fixation and interferes with synaptic plasticity and information transmission. Experimentally, simply mimicking the activation of Sst neurons inhibits memory fixation, even in animals that are normally sleeping. Conversely, inhibiting these neurons during post-learning sleep promotes memory fixation [19]. This indicates that Sst neurons are part of the direct mechanism by which sleep deprivation causes cognitive impairment.

Another important example is the circuit in the central nucleus of the amygdala (CeA) that links stress and sleep disturbances, where Sst neurons (CeA^{SST+}) in the CeA are activated by acute stress [22]. Optogenetic activation of these neurons induces a rapid transition from NREM sleep to wakefulness and prolongs the time required to fall asleep (onset latency) [22]. Critically, the inhibition of the activity of these neurons ameliorates stress-induced insomnia but has no effect on normal physiological sleep [22]. This shows that CeA^{SST+} neurons are not part of the normal sleep–wake cycle but are the basis of pathological sleep disturbances caused by specific stress conditions.

These findings show that the function of Sst neurons changes dramatically from "sleep promoting" to "sleep inhibiting" depending on the brain region (cortex, hippocampus, amygdala) and brain state (normal sleep, sleep deprivation, stress). This is not a contradiction but evidence of sophisticated context-dependent circuit regulation, showing that different systems (homeostasis, stress response, and memory) mobilize the powerful network regulators of Sst neurons for different and sometimes conflicting purposes.

The effects of Sst neurons on sleep are tightly controlled by neural circuits between specific brain regions. Of particular importance are the top-down control from the prefrontal cortex to the hypothalamus and structural changes caused by sleep deprivation in hippocampal and neocortical circuits.

The prefrontal cortex (PFC) is responsible for higher executive functions and is highly sensitive to sleep deprivation. Recent studies have revealed the existence of a top-down control mechanism for sleep homeostasis, in which the PFC monitors the internal state of sleep debt and directs appropriate physiological and behavioral responses [23].

Activity-dependent gene labelling techniques using c-Fos have identified a specific subpopulation of PFC Sst-GABAergic neurons that are particularly excitatory and activated in response to sleep deprivation [24]. These activated PFC Sst neurons have long-range projections to two distinct nuclei in the hypothalamus, each of which has been shown to be responsible for a distinct function by optogenetic manipulation.

Projection to the lateral preoptic area (LPO): Selective stimulation of the axon terminals of these PFC Sst neurons with the LPO elicited sleep preparation behaviors, such as nesting behavior. This behavior is accompanied by physiological changes, such as theta wave enhancement and increased body temperature, but does not induce sleep itself [23]. The LPO is part of the preoptic area, a long-established sleep facilitation center [25] that provides a new mechanism by which the cortex communicates with the sleep center and initiates preparatory behaviors before sleep. This provides a new mechanism by which the cortex communicates with the sleep center to initiate preparatory behaviors before sleep.

Projection to the lateral hypothalamus (LH): In contrast, stimulation of the axon terminals of the same group of neurons in the LH does not induce nesting behavior but instead induces robust NREM sleep (recovery sleep) with strong delta waves, accompanied by a decrease in body temperature [23].

This circuit shows the existence of a sophisticated top-down control, in which the PFC senses sleep debt and commands the hypothalamus via parallel Sst neuronal pathways to "first prepare a safe place to sleep (via LPO) and then initiate restorative sleep (via LH)," a two-stage coordinated homeostatic response. This challenges the classical view that sleep is exclusively driven by subcortical, bottom-up mechanisms.

The effects of sleep deprivation go beyond functionality and leave physical traces on Sst interneuron circuits. Using state-of-the-art imaging techniques, studies have reported that sleep deprivation for only a few hours can induce region-specific structural changes in the dendritic spines (postsynaptic structures) of Sst interneurons in the hippocampus and neocortex [26]. The changes were not uniform; for example, there was a dramatic increase in the density and size of spines in the CA3 region of the hippocampus and a slight decrease in the visual cortex.

As dendritic spines are key sites of excitatory input, this rapid and widespread structural change indicates that sleep deprivation fundamentally alters excitatory input to Sst interneurons. This disrupts the excitatory (E-I) balance across critical neural networks through changes in the inhibitory output from these neurons [26]. This finding provides a physical mechanism at the cellular level underlying the cognitive decline and changes in network activity observed after sleep deprivation, a process by which the brain's inhibitory circuits are dynamically and physically sculpted by sleep, and whose disruption is the core mechanism of sleep deprivation-induced dysfunction. This suggests that it is a core mechanism of sleep deprivation-induced dysfunction.

Table 5. Region-Specific Roles of Sst Neurons in Sleep–Wake Regulation

Brain Region	Context / Trigger	Function of Sst Neurons	Effect on Sleep / Wakefulness	Refs
Neocortex	Normal Sleep	Generates "OFF state" of slow waves	SWA ↑, NREM sleep time ↑	[18]
Hippocampus	Sleep Deprivation	Inhibitory gate for memory consolidation	Inhibits sleep function (memory)	[19]
Central Amygdala (CeA)	Acute Stress	Mediates stress-induced arousal	Sleep latency ↑, Arousal ↑	[22]
Prefrontal Cortex (PFC) Projection	Sleep Deprivation (Homeostatic Response)	Coordinates recovery sleep	Induces preparatory behavior and recovery NREM sleep	[23]

In conclusion, it is reasonable to detect Sst expressive neurons that regulate SD and RS (Table 5).

4. Discussion

To determine whether the detection of Sst enrichment was accidental and might not be supported when using other samples, we uploaded the two sets of selected genes to AlphaGenome [27] and tried to identify if Sst enrichment can be observed in AlphaGenome. AlphaGenome is a generative AI server that can predict various omics features (gene expression, DNA methylation, etc.) only from DNA sequences (for more details on how to obtain predicted features from AlphaGenome, see the Methods section).

Table 6 summarizes if Sst is in the top-ranked 10 cell types and/or associated with adjusted P-values less than 0.05 in "Allen Brain Atlas 10x scRNA 2021" (The full list of corresponding Figures S2 to S21 and Tables S1 to S21 is in supplementary files 5 and 6). Sst is almost always ranked in the top 10 cell types, and in 16 out of 80 cases. Sst is associated with adjusted Pvalues less than 0.05. All eight pairwise comparisons were associated with at least one case with adjusted Pvalues less than 0.05. As DNase-Seq identified the transcriptionally active sites, which are expected to correspond with actively expressed genes, the Sst enrichment observed in DNase-Seq predicted by AlphaGenome suggests that this enrichment is not accidental and supported by other samples as well.

Table 6. Confirmation of Sst enrichment in "Allen Brain Atlas 10x scRNA 2021" category using DNase-Seq predicted by AlphaGenome. ○: Sst is in top ranked 10 cell types and associated with adjusted P-values less than 0.05. △: Sst is in top ranked 10 cell types but not associated with adjusted P-values less than 0.05. ×: Sst is neither in top ranked 10 cell types nor associated with adjusted P-values less than 0.05. $\ell_1 = 2$ only and $\ell_1 = 3$ only mean that common genes between $\ell_1 = 2$ and $\ell_1 = 3$ are excluded from $\ell_1 = 2$ and $\ell_1 = 3$, respectively.

Pairwise comparisons			$\ell_1 = 2$	$\ell_1 = 3$	$\ell_1 = 2$ and $\ell_1 = 3$	$\ell_1 = 2$ only	$\ell_1 = 3$ only
Left cerebral cortex	>		△	△	○	△	△
	<		○	△	○	○	△
Right cerebral cortex	>		○	△	○	△	△
	<	Brain	○	○	△	○	×
Layer of hippocampus	>		○	△	○	△	△
	<		○	△	○	○	×
Frontal cortex	>		△	△	○	△	△
	<		○	△	△	△	×

5. Conclusions

In this study, we analyzed time-dependent gene expression in the prefrontal cortex of mice during SD and RS. Compared with control, we identified two distinct expression patterns using TD, 'altered during SD but not recovered during RS' and 'altered during SD with an overshoot during RS,' which differ from the commonly expected pattern "altered during SD and recovered during RS." We extracted two sets of genes associated with these patterns and performed enrichment analysis, which revealed enrichment in Sst-expressing cell types. To confirm that the Sst enrichment was not due to chance, we uploaded both gene sets to AlphaGenome and retrieved predicted DNase-Seq. The top-ranked genes distinguishing various brain subregions from the whole brain were enriched in Sst cell types. This suggests that Sst enrichment is not accidental, even when other samples are used.

Author Contributions: Y.-h.T. designed the study and edited the manuscript. Y-h. T and K.K. performed the analyses and wrote the manuscript. All authors read and agreed to the published version of the manuscript.

Funding: This work was supported by KAKENHI [grant numbers 24K15168] to YHT.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data used in this study in GSE237419.

Acknowledgments: We thank Mr. Hiroyuki Kanaya for critically reviewing the manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Popescu, A.; Ottaway, C.; Ford, K.; Medina, E.; Patterson, T.W.; Ingiosi, A.; Hicks, S.C.; Singletary, K.; Peixoto, L. Transcriptional dynamics of sleep deprivation and subsequent recovery sleep in the male mouse cortex. *Physiological Genomics* **2025**, *57*, 431–445, [https://doi.org/10.1152/physiolgenomics.00128.2024]. PMID: 40315180, https://doi.org/10.1152/physiolgenomics.00128.2024.
2. Taguchi, Y.h. *Unsupervised Feature Extraction Applied to Bioinformatics: A PCA Based and TD Based Approach*, 2 ed.; Unsupervised and Semi-Supervised Learning, Springer International Publishing: Switzland, 2024.
3. Team, T.B.D. *BSgenome.Mmusculus.UCSC.mm39: Full genome sequences for Mus musculus (UCSC genome mm39, based on GRCm39)*, 2021. R package version 1.4.3.
4. DeepMind, G. AlphaGenome, 2025. Accessed: 2025-09-19.

5. Zhou, Y.; Zhou, B.; Pache, L.; Chang, M.; Khodabakhshi, A.H.; Tanaseichuk, O.; Benner, C.; Chanda, S.K. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nature Communications* **2019**, *10*, 1523. https://doi.org/10.1038/s41467-019-09234-6.

6. Naidoo, N. Roles of Endoplasmic Reticulum and Energetic Stress in Disturbed Sleep. *NeuroMolecular Medicine* **2012**, *14*, 213–219. https://doi.org/10.1007/s12017-012-8179-9.

7. Wang, L.; Aton, S.J. Perspective – ultrastructural analyses reflect the effects of sleep and sleep loss on neuronal cell biology. *Sleep* **2022**, *45*, zsac047, [https://academic.oup.com/sleep/article-pdf/45/5/zsac047/43517156/zsac047.pdf]. https://doi.org/10.1093/sleep/zsac047.

8. Nishimaru, H.; Matsumoto, J.; Setogawa, T.; Nishijo, H. Neuronal structures controlling locomotor behavior during active and inactive motor states. *Neuroscience Research* **2023**, *189*, 83–93. https://doi.org/https://doi.org/10.1016/j.neures.2022.12.011.

9. Taguchi, Y.H.; Turki, T. L1 GABAergic neurones in the frontal cortex and L6 glutamatergic neurones in the prefrontal cortex use RHO GTPase to differentiate between unconsciousness states. *bioRxiv* **2025**, [https://www.biorxiv.org/content/early/2025/06/26/2025.06.]. https://doi.org/10.1101/2025.06.21.660840.

10. Xie, Z.; Bailey, A.; Kuleshov, M.V.; Clarke, D.J.B.; Evangelista, J.E.; Jenkins, S.L.; Lachmann, A.; Wojciechowicz, M.L.; Kropiwnicki, E.; Jagodnik, K.M.; et al. Gene Set Knowledge Discovery with Enrichr. *Current Protocols* **2021**, *1*, e90, [https://currentprotocols.onlinelibrary.wiley.com/doi/pdf/10.1002/cpz1.90]. https://doi.org/https://doi.org/10.1002/cpz1.90.

11. Wu, C.; Sun, D. GABA receptors in brain development, function, and injury. *Metabolic Brain Disease* **2015**, *30*, 367–379. https://doi.org/10.1007/s11011-014-9560-1.

12. Marín, O. Development of GABAergic Interneurons in the Human Cerebral Cortex. *European Journal of Neuroscience* **2025**, *61*, e70136, [https://onlinelibrary.wiley.com/doi/pdf/10.1111/ejn.70136]. e70136 EJN-2024-12-32323.R1, https://doi.org/https://doi.org/10.1111/ejn.70136.

13. Tremblay, R.; Lee, S.; Rudy, B. GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits. *Neuron* **2016**, *91*, 260–292. doi: 10.1016/j.neuron.2016.06.033, https://doi.org/10.1016/j.neuron.2016.06.033.

14. Urban-Ciecko, J.; Barth, A.L. Somatostatin-expressing neurons in cortical networks. *Nature Reviews Neuroscience* **2016**, *17*, 401–409. https://doi.org/10.1038/nrn.2016.53.

15. Liguz-Lecznar, M.; Dobrzanski, G.; Kossut, M. Somatostatin and Somatostatin-Containing Interneurons—From Plasticity to Pathology. *Biomolecules* **2022**, *12*. https://doi.org/10.3390/biom12020312.

16. Fee, C.; Prevot, T.D.; Misquitta, K.; Knutson, D.E.; Li, G.; Mondal, P.; Cook, J.M.; Banasr, M.; Sibille, E. Behavioral Deficits Induced by Somatostatin-Positive GABA Neuron Silencing Are Rescued by Alpha 5 GABA-A Receptor Potentiation. *International Journal of Neuropsychopharmacology* **2021**, *24*, 505–518, [https://academic.oup.com/ijnp/article-pdf/24/6/505/38956206/pyab002.pdf]. https://doi.org/10.1093/ijnp/pyab002.

17. Scheyltjens, I.; Arckens, L. The Current Status of Somatostatin-Interneurons in Inhibitory Control of Brain Function and Plasticity. *Neural Plasticity* **2016**, *2016*, 8723623, [https://onlinelibrary.wiley.com/doi/pdf/10.1155/2016/8723623]. https://doi.org/https://doi.org/10.1155/2016/8723623.

18. Funk, C.M.; Peelman, K.; Bellesi, M.; Marshall, W.; Cirelli, C.; Tononi, G. Role of Somatostatin-Positive Cortical Interneurons in the Generation of Sleep Slow Waves. *Journal of Neuroscience* **2017**, *37*, 9132–9148, [https://www.jneurosci.org/content/37/38/9132.full.pdf]. https://doi.org/10.1523/JNEUROSCI.1303-17.2017.

19. Delorme, J.; Wang, L.; Kuhn, F.R.; Kodoth, V.; Ma, J.; Martinez, J.D.; Raven, F.; Toth, B.A.; Balendran, V.; Medina, A.V.; et al. Sleep loss drives acetylcholine- and somatostatin interneuron–mediated gating of hippocampal activity to inhibit memory consolidation. *Proceedings of the National Academy of Sciences* **2021**, *118*, e2019318118, [https://www.pnas.org/doi/pdf/10.1073/pnas.2019318118]. https://doi.org/10.1073/pnas.2019318118.

20. Wilson, D.A.; Fleming, G.; Williams, C.R.O.; Teixeira, C.M.; Smiley, J.F.; Saito, M. Somatostatin neuron contributions to cortical slow wave dysfunction in adult mice exposed to developmental ethanol. *Frontiers in Neuroscience* **2023**, *Volume 17 - 2023*. https://doi.org/10.3389/fnins.2023.1127711.

21. Zielinski, M.R.; Atochin, D.N.; McNally, J.M.; McKenna, J.T.; Huang, P.L.; Strecker, R.E.; Gerashchenko, D. Somatostatin+/nNOS+ neurons are involved in delta electroencephalogram activity and cortical-dependent recognition memory. *Sleep* **2019**, *42*, zsz143, [https://academic.oup.com/sleep/article-pdf/42/10/zsz143/30122359/zsz143.pdf]. https://doi.org/10.1093/sleep/zsz143.

22. Yao, W.; Huang, S.X.; Zhang, L.; Li, Z.S.; Huang, D.Y.; Huang, K.Q.; Huang, Z.X.; Nian, L.W.; Li, J.L.; Chen, L.; et al. Central amygdala somatostatin neurons modulate stress-induced sleep-onset insomnia. *Communications Biology* **2025**, *8*, 381. https://doi.org/10.1038/s42003-025-07679-8.

23. Tossell, K.; Yu, X.; Giannos, P.; Soto, B.A.; Nollet, M.; Yustos, R.; Miracca, G.; Vicente, M.; Miao, A.; Hsieh, B.; et al. Somatostatin neurons in prefrontal cortex initiate sleep-preparatory behavior and sleep via the preoptic and lateral hypothalamus. *Nature Neuroscience* **2023**, *26*, 1805–1819. https://doi.org/10.1038/s41593-023-01430-4.

24. Tossell, K.; Yu, X.; Soto, B.A.; Vicente, M.; Miracca, G.; Giannos, P.; Miao, A.; Hsieh, B.; Ma, Y.; Yustos, R.; et al. Sleep deprivation triggers somatostatin neurons in prefrontal cortex to initiate nesting and sleep via the preoptic and lateral hypothalamus. *bioRxiv* **2020**, [https://www.biorxiv.org/content/early/2020/07/02/2020.07.01.179671.full.pdf]. https://doi.org/10.1101/2020.07.01.179671.

25. Saito, Y.; Tsujino, N.; Hasegawa, E.; Akashi, K.; Abe, M.; Mieda, M.; Sakimura, K.; Sakurai, T. GABAergic neurons in the preoptic area send direct inhibitory projections to orexin neurons. *Frontiers in Neural Circuits* **2013**, *Volume 7 - 2013*. https://doi.org/10.3389/fncir.2013.00192.

26. Raven, F.; Medina, A.V.; Schmidt, K.; He, A.; Vankampen, A.A.; Balendran, V.; Aton, S.J. Brief sleep disruption alters synaptic structures among hippocampal and neocortical somatostatin-expressing interneurons. *bioRxiv* **2024**, [https://www.biorxiv.org/content/early/2024/11/08/2024.07.22.604591.full.pdf]. https://doi.org/10.1101/2024.07.22.604591.

27. Avsec, Ž.; Latysheva, N.; Cheng, J.; Novati, G.; Taylor, K.R.; Ward, T.; Bycroft, C.; Nicolaisen, L.; Arvaniti, E.; Pan, J.; et al. AlphaGenome: advancing regulatory variant effect prediction with a unified DNA sequence model. *bioRxiv* **2025**, [https://www.biorxiv.org/content/early/2025/07/11/2025.06.25.661532.full.pdf]. https://doi.org/10.1101/2025.06.25.661532.