Jam2 Signaling Functions Downstream of Hand2 To Initiate The Formation Of Organ-Specific Vascular Progenitors In Zebrafish

Martyna Griciunaite*, Julius Martinkus*, Sanjeeva Metikala*, Ricardo DeMoya,

Suman Gurung, Diandra Rufin Florat, Saulius Sumanas[1]

Department of Pathology and Cell Biology

University of South Florida, Tampa, FL, USA

* Equal contribution

[1]Corresponding author. Address: USF Heart Institute, MDD 638, 560 Channelside Dr, Tampa, FL 33602. E-mail: ssumanas @usf.edu

ABSTRACT

The mechanisms regulating the formation of organ-specific vasculature are still poorly understood. We have previously identified a population of late-forming endothelial progenitor cells in zebrafish embryos (termed secondary vascular field, SVF), which emerge from the lateral plate mesoderm after 24 hpf stage, after blood circulation has been initiated. Here we investigate the functional role of SVF cells and the molecular mechanisms that govern the emergence of these SVF cells and their contribution to vasculature. We identified that the bHLH transcription factor Hand2 and Junctional Adhesion Molecule Jam2b are expressed in the SVF-forming region and are required for the emergence of SVF cells in zebrafish embryos. Time-lapse imaging and *jam2b:Cre* based lineage tracing showed that SVF cells serve as the major source of the intestinal vasculature, including the supraintestinal artery (SIA) and subintestinal vein (SIV), which are subsequently remodeled to provide the blood flow to many internal organs. To analyze the functional role of *jam2b* and a related *jam2a* gene in vascular development, we generated double maternal-zygotic *jam2a; jam2b* mutants, which display a greatly reduced number of SVF cells and show defects in the intestinal vasculature development. Further analysis showed that *hand2* functions in the SVF-forming region upstream of *jam2b* and is required to induce expression of transcription factor *etv2/etsrp,* a known master regulator of vasculogenesis. In summary, our results identify new roles for Jam2 signaling and Hand2 function in the emergence of organ-specific vascular progenitors. The presence of similar progenitors in mammalian embryos suggests that this mechanism is evolutionarily conserved.

INTRODUCTION

It has been well established that organ-specific blood vessels exhibit a large diversity of specialized endothelial cells. However, their developmental origin has been debatable. Previous studies have suggested that the progenitors of internal organ vasculature in mammalian embryos form by differentiation *de novo,* or vasculogenesis.[1-4] Different studies have demonstrated that the endocardium, and endoderm or mesothelium may contribute to the organ vasculature in mammals and avian embryos.[5-7] In contrast, several studies have argued that the posterior cardinal vein (PCV) serves as the major source of organ-specific vascular endothelial cells in zebrafish embryos.[8-10] According to this model, individual cells from the PCV migrate at 30-48 hours post fertilization (hpf) and coalesce into the intestinal vessels, which include the supraintestinal artery (SIA) and the subintestinal vein (SIV). These vessels subsequently undergo remodeling and ultimately provide blood supply for the internal organs, including the pancreas and the liver.[10] However, this model in zebrafish contrasts with the *de novo* vasculogenesis model, suggested in mammalian embryos. In addition, it is unclear if the PCV is the only or even the main source of internal organ-specific vascular progenitors.

In zebrafish, similar to mammalian embryos, the earliest vascular progenitors originate from the lateral plate mesoderm (LPM) during late gastrulation and somitogenesis, and coalesce into the major axial vessels, the dorsal aorta (DA) and the PCV.[11-14] This process is thought to be largely complete by the 24 hpf stage, when blood circulation is first initiated. We have recently demonstrated that between 24 and 48 hpf a separate group of late-forming vascular progenitors emerges from the LPM along the yolk extension at the site termed the secondary vascular field (SVF).[15] SVF cells can uniquely incorporate into the PCV after blood circulation has already

been established, and show high expression of vascular progenitor markers, including *etv2/etsrp,* *tal1/scl, npas4l,* while they are negative for the expression of more differentiated endothelial markers such as *kdrl* or *fli1a.* Previous time-lapse imaging using *etv2* reporter lines demonstrated that SVF cells can contribute to the intestinal vasculature, including the SIA and SIV.[15] However, the ultimate fate of SVF cells has not been clear due to the absence of genetic tools to perform the lineage tracing. Furthermore, mechanisms regulating the emergence of SVF cells and their contribution to vasculature remain largely unknown.

To identify the molecular signature of SVF cells, we have previously performed transcriptomic analysis of the zebrafish trunk region at 30 hpf.[16] This analysis identified some SVF marker genes, which have not been previously associated with vascular development, including *junctional adhesion molecule 2b (jam2b).*[15] Vertebrate JAM homologs have been implicated in diverse processes including leukocyte-endothelial interaction and transendothelial migration, maintenance of hematopoietic stem cells (HSCs), vascular development, and others.[17,18] A related zebrafish paralog *jam2a* has been previously implicated in HSC formation.[19] However, the role of *jam2b* in vascular development has not been described.

Here we analyzed the role of zebrafish *jam2b* in the formation of intestinal vasculature and the emergence of SVF cells. We demonstrate that *jam2b* is expressed along the lateral plate mesoderm in the region that includes SVF cells. By using genetic lineage tracing and time-lapse imaging we demonstrate that *jam2b*-positive cells contribute to multiple tissues, including the mesothelium, pericardium, fin buds and venous and intestinal vasculature. To study the role of Jam2 homologs in vascular development, we have generated genetic mutants in *jam2b* and a related homolog *jam2a*. Double *jam2a; jam2b* maternal-zygotic mutants show a reduced number of SVF cells, which correlates with the defects in the formation of the intestinal and venous

vasculature, and display reduced regenerative potential after endothelial cell ablation. We further demonstrate that the transcription factor *hand2* functions upstream of *jam2b* and is required for the formation of SVF cells. Altogether, our results identify a new role for Jam2 signaling and Hand2 function in the emergence of organ-specific vascular progenitors.

RESULTS

***Jam2b* expression analysis.** During the previous transcriptomic analysis, we have identified *jam2b* as one of the markers for SVF cells, which are late-forming vascular progenitors that emerge along the yolk extension at 24-48 hpf.[15] As we and others have previously demonstrated, *jam2b* expression is localized to the ventrolateral region of the LPM along the anterior and trunk regions during 20 somite – 48 hpf stages.[15,20,21] It is bilaterally expressed along the yolk extension during 24-48 hpf, where SVF cells originate (Suppl. Fig. S1). We have previously demonstrated that *jam2b* and the vascular progenitor marker *etv2/etsrp* expression in the SVF cells partially overlaps along the medial portion of *jam2b* expression domain within the mid-trunk region.[15]

Generation and characterization of *jam2b$^{Gt(2A-Gal4)}$* knock-in reporter line. To analyze *jam2b+* cells in live embryos, we used CRISPR / Cas9-mediated non-homologous recombination approach [22] to insert Gal4 transcriptional activator into the genomic locus of *jam2b* (Suppl. Fig. S2A). Similar to the endogenous *jam2b* mRNA expression, *jam2b$^{Gt(2A-Gal4)ci55}$;UAS:GFP* (further abbreviated as *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP*) expression was observed bilaterally in the lateral plate mesoderm along the yolk and yolk extension at 24 hpf (Suppl. Fig. S2B; Fig. 1A,B). At 3 and 5 dpf *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP* expression was observed in the cell

layer surrounding the yolk and yolk extension, which presumptively corresponds to the mesothelium (Fig. 1C-F). GFP expression was also observed in a subset of muscle cells within the skeletal and craniofacial muscle, and in the cells at the periphery of the eye (Fig. 1C-F). GFP expression in the trunk portion of the LPM partially overlapped with *etv2* expression in SVF cells at 30 hpf (Fig. 1G-I). Interestingly, GFP-positive cells were observed in the vasculature between 30 hpf and 3 dpf (Fig. 1J-R). The majority of GFP+ cells were located in the PCV, while a few cells were found within the ISV and SIA, and in less frequent cases contribution to the DA and parachordal lymphangioblasts (PLs) was observed (Suppl. Fig. S3). In addition, some GFP cells were located in the thoracic duct at 4 dpf, suggesting that SVF cells contribute to lymphatic vasculature (Fig. 1S-U). Due to very bright GFP expression over the yolk, we could not reliably analyze GFP cell presence in the SIV vessel, which is also positioned over the yolk and yolk extension. Overall, these findings suggest that *jam2b+* cells contribute to the vasculature, including the PCV, ISVs, intestinal vasculature and lymphatics. These findings are consistent with our previously reported contribution of SVF cells to the PCV and intestinal vasculature [15].

Jam2b+ cells are derived from the lateral plate mesoderm and incorporate into functional vasculature. To visualize directly the contribution of *jam2b+* cells to vasculature, we performed time-lapse imaging of *jam2b^Gt(2A-Gal4)*;*UAS:GFP*; *Tg(kdrl:mCherry)* embryos starting at 24 hpf. This imaging showed that a subset of GFP-positive cells migrated dorsally away from the LPM region, integrated into the PCV, proliferated and initiated *kdrl:mCherry* expression as they differentiated into vascular endothelial cells (Fig. 2A-F and Movie 1). This data shows that mesenchymal cells can contribute to and incorporate into the functional vasculature, even after

blood circulation has initiated, in agreement with our previous observations using $etv2^{Gt(2A-Venus)}$ reporter line.[15]

In an attempt to identify the origin of SVF cells, we analyzed $jam2b^{Gt(2A-Gal4)}$;UAS:GFP expression at the 18-somite stage. Similar to *jam2b* mRNA expression, GFP-positive cells were located bilaterally in the lateral plate mesoderm, mostly lateral and ventral to the emerging *etv2+* vascular progenitors (Fig. 2G-I). There was no apparent overlap between $jam2b^{Gt(2A-Gal4)}$;UAS:GFP and *etv2* expression at this stage. This domain largely overlapped with the expression domain of *prrx1a* (Fig. 2J-L), which labels the lateral portion of the LPM.[23] To analyze the behavior of these cells, we performed time-lapse imaging of $jam2b^{Gt(2A-Gal4)}$;UAS:GFP embryos starting at the 18-somite stage. GFP-positive cells remained in the same LPM region from the 18-somite stage to 24-30 hpf stages (Movie 2). As the embryo continued to undergo convergent extension, the same LPM region became bilaterally located along the yolk extension. Thus, *jam2b*-expressing cells observed at 24-30 hpf along the yolk extension are derived from the ventrolateral portion of the LPM.

Genetic lineage tracing of *jam2b*+ cells. To perform genetic lineage tracing of *jam2b*-expressing cells, we generated *UAS:Cre* and *UAS:CreERT2* zebrafish transgenic lines using Tol2 transgenesis. Subsequently they were crossed to $jam2b^{Gt(2A-Gal4)}$; *actb2:loxP-BFP-loxP-dsRed; fli1:GFP* adults, which ubiquitously express floxed BFP under *actb2* promoter, in combination with the $jam2b^{Gt(2A-Gal4)}$ knock-in driver and vascular endothelial *fli1:GFP* reporter (Fig. 3A). This approach permanently labels *jam2b*-derived cells, which switch their color from BFP to dsRed due to Cre-LoxP recombination. The embryos were then analyzed at 3 dpf by confocal microscopy, revealing switched dsRed-positive cells in the presumptive mesothelial layer surrounding the yolk and yolk extension, a subset of skeletal muscle cells and vasculature (Fig.

3B,E). In addition, both epicardial and myocardial layers within the heart were labeled by dsRed fluorescence (Fig. 3C,D). An embryo negative for $jam2b^{Gt(2A-Gal4)}$ expression was included for autofluorescence control (Fig. 3H). The majority of labeled vascular endothelial cells were positioned in the SIA and PCV, while a few cells were positioned in the DA, PL, and ISVs (Fig. 3F,G,I-K). Most ISVs, which showed dsRed expression, were venous, and very little contribution to arterial ISVs was observed (Fig. 3F,G,K). Due to intense dsRed fluorescence in the mesothelial layer along the yolk and yolk extension we were unable to reliably determine the contribution of *jam2b+* cells to the SIV. To determine when *jam2b+* cells contribute to vasculature, *UAS:CreERT2* line was crossed to $jam2b^{Gt(2A-Gal4)}$; *actb2:loxP-BFP-loxP-dsRed; fli1:GFP* adults and 4-hydroxitamoxifen (4-OHT) was added at different time points to induce Cre-LoxP recombination. 4-OHT addition at 7 hpf mostly in labeled SIA with a smaller fraction of cells present in the SIV, PCV and venous ISVs (Fig. 3L). Only cells in the SIA were labeled when 4-OHT was added at 22 hpf. Addition of 4-OHT at 7 hpf and subsequent washing out at 22 hpf resulted in the labeling of SIA and a small number of cells within the PCV (Fig. 3L). These results argue that *jam2b+* cells contribute to vasculature both before and after 22 hpf. However, contribution after 22 hpf was largely limited to the intestinal vasculature, which supports our findings that SVF cells show major contribution to the SIA and SIV.

Intestinal vasculature forms largely by new vasculogenesis. Previous studies have argued that the intestinal vasculature, including the SIA and SIV, are derived from the PCV [8-10]. In contrast, our current data suggest that *jam2b*-positive SVF cells make a major contribution to the intestinal vasculature. However, the number of intestinal vascular cells labeled by $jam2b^{Gt(2A-Gal4)}$; *UAS:Cre* or *UAS:CreERT2* recombination was rather small. This could be due to only a small contribution of SVF cells to the intestinal vasculature, or alternatively, due to low

recombination efficiency observed in *jam2b*[Gt(2A-Gal4)]*; UAS:Cre* and *UAS:CreERT2* lines. To clarify the contribution of new vasculogenesis towards the intestinal vasculature, we photoconverted all vascular endothelial cells in *etv2:Kaede* transgenic line at 24 hpf stage, which expresses green to red photoconvertible Kaede in all vascular endothelial cells,[24] and then analyzed the percentage of red and green labeled cells at 4 dpf in the intestinal vasculature over the yolk extension region, where SVF cells are observed. Notably, 78.5% of cells in the SIA and 80.5% cells in the SIV showed only green expression of newly synthesized Kaede, and had no red fluorescence of photoconverted protein, arguing that they originated after photoconversion at 24 hpf stage (Fig. 4). These data suggest that most intestinal progenitors in the mid and posterior trunk regions form by new vasculogenesis from SVF cells.

Transcriptomic analysis of SVF cells. We have previously performed single-cell RNA sequencing (scRNA-seq) of the entire trunk region of zebrafish embryos in order to identify the transcriptional profile of SVF cells.[15,16] While this analysis identified a set of SVF marker genes that included *etv2, tal1, lmo2* and *jam2b,* only a limited number of marker genes were identified with this approach, because SVF cells comprise only a small portion of cells in the entire trunk. Therefore, we used *jam2b* and *etv2* reporter lines to perform additional transcriptomic analysis in order to characterize the profiles of SVF cells and their transition to vascular endothelial cells in greater detail. mCherry and Venus-positive cells were FACS-sorted from *jam2b*[Gt(2A-Gal4)]*;UAS:mCherry-NTR; etv2*[Gt(2A-Venus)] embryos at 36 hpf and subjected to scRNA-seq using the Chromium platform (10x Genomics). Bioinformatic analysis using Seurat 4.0 identified a total of 37 cell clusters, including the SVF cell cluster, five venous, three arterial clusters, and a capillary endothelial cluster (Fig. 5A-C, Suppl. Table 1). The SVF cell cluster (cluster #15) was identified based on top marker genes, which included *npas4l, tal1, lmo2,* and *etv2* (Fig. 5D,E,

Suppl. Table 1). Additional SVF top marker genes included *gfra3, sox7, si:dkey-52l18.4, egfl7* and *tmem88a* (Suppl. Table 1). The key SVF marker genes, including *etv2, tal1, lmo2, sox7, si:dkey-52l18.4* and *egfl7* were also identified in our earlier study,[15] arguing that the current approach has successfully identified the transcriptional signature of SVF cells.

To identify transitional events during SVF cell differentiation into vascular endothelium, we performed cell lineage trajectory analysis using Monocle 3.[25] Cell lineage trajectory showed that SVF cells can directly transition into both venous and arterial clusters (Fig. 5F). Such differentiation trajectory agrees with experimental observations of SVF cells contributing both to venous (PCV, SIV) and arterial vasculature (SIA). Unfortunately, it is currently not possible to distinguish SIA or SIV-specific cells on the lineage trajectory from other types of arterial or venous cells due to the absence of specific markers for these cells.

***Jam2b* mutants undergo normal vascular development.** To analyze the functional role of Jam2b in vascular development, we generated *jam2b* mutants using CRISPR/Cas9 mediated genome editing. Two different guide RNAs were used to delete the *jam2b* promoter region, and a stable mutant line was established (Suppl. Fig. S4A). As indicated by ISH analysis, *jam2b-/-* embryos showed no detectable *jam2b* expression in the LPM region, suggesting that the mutant may be a null or close to a null (Suppl. Fig. S4C,D). However, both zygotic and maternal zygotic *jam2b-/-* embryos appeared morphologically normal and did not show any apparent vascular defects based on *kdrl:GFP* expression analysis (data not shown). *Jam2b* homozygous mutants were viable as adults and did not exhibit any apparent defects. *Etv2* expression in SVF cells was also not affected in maternal-zygotic *jam2b* mutant embryos (Suppl. Fig. S5A-C). We have previously demonstrated that SVF cells contribute to vascular recovery after endothelial cell ablation in *etv2*[Gt(2A-Gal4)]*; UAS:mCherry-NTR* embryos.[15] We then tested if vascular recovery was

affected in *jam2b* deficient embryos. Metronidazole (MTZ) was added at approximately 10 hpf stage and then washed out at 45 hpf stage in zygotic *jam2b-/-* and sibling control *jam2b+/-* embryos in *etv2*^{Gt(2A-Gal4)}*; UAS:mCherry-NTR; kdrl:GFP* background. This resulted in nearly complete ablation of vascular endothelial cells. As we previously demonstrated, such treatment results in a complete ablation of all vascular endothelial cells except for SVF cells, which emerge later than other endothelial cells.[15] Afterwards, the recovery was analyzed at 72 hpf, when the partial regeneration of vascular cords was apparent, which as our previous study suggested, were derived from SVF cells.[15] However, no difference in vascular recovery was observed between *jam2b-/-* and *jam2b+/-* embryos (Suppl. Fig. S5D,E).

jam2a; jam2b mutants display defects in SVF cell formation and intestinal and venous vasculature. We then explored the possibility that *jam2b* may function redundantly with a related homolog *jam2a*. While *jam2a* is broadly expressed in the somitic muscle,[20,21,26] our scRNA-seq analysis indicated that it also showed significant expression in the SVF cell population, although it was not enriched among marker genes.[15] We used CRISPR / Cas9 mutagenesis to remove the entire coding sequence of *jam2a* (Suppl. Fig. S4B,E,F). To expedite the generation of double mutant embryos, *jam2a* mutant generation was performed in *jam2b-/-* mutant background. Stable *jam2a;jam2b* mutant line was generated and used for further experiments. Zygotic *jam2a-/-* / maternal *jam2b-/-* mutant embryos, obtained by the in-cross *of jam2a+/-;jam2b-/-* adults did not show any apparent defects in vascular development or SVF cell formation (data not shown). Subsequently, maternal-zygotic (MZ) mutant embryos were obtained and analyzed for SVF cell formation and vascular defects. Interestingly, MZ *jam2a-/-;jam2b-/-* embryos showed significant reduction in the SVF cell number when compared to control double heterozygous embryos in the same background. The number of *etv2, tal1* and

npas4l-positive SVF cells was greatly reduced in MZ *jam2a;jam2b* mutant embryos, while no other morphological defects were apparent (Fig. 6). Overall vascular patterning appeared largely unaffected in MZ *jam2a;jam2b* mutant embryos based on *kdrl:GFP* fluorescence analysis at 3 dpf (Fig. 7A,B). However, MZ *jam2a;jam2b* mutant embryos showed greater incidence of incompletely extended ISVs (Fig. 7A-C). Venous ISVs, derived from the PCV, made 97% (33 out of 34) of such truncated ISVs, while only 3% were arterial. This suggested that the number of venous cells could be reduced, possibly due to reduced contribution of SVF cells to the PCV. Indeed, the number of ECs in the floor of the PCV was slightly yet significantly reduced in MZ *jam2a; jam2b* mutant embryos (Fig. 7D-F). However, the most pronounced defects were observed in the formation of the intestinal vasculature, in particular the SIA, which was underdeveloped and showed gaps or missing segments in MZ *jam2a;jam2b* mutants (Fig. 7G-I). In addition, a significant fraction of MZ *jam2a;jam2b* mutants showed gaps in the SIV (Fig. 7G-I).

Because our data suggested that *jam2b+* cells may contribute to lymphatics, we analyzed the formation of the thoracic duct using *prox1:RFP, fli1:GFP* reporter embryos. Approximately 37% of MZ *jam2a;jam2b* mutants had gaps in the thoracic duct compared to 5% of control *jam2a+/-;jam2b+/-* embryos (Fig. 7J-L). These results argue that *jam2a* and *jam2b* function is required for the formation of the intestinal vasculature and lymphatics.

We then tested if vascular recovery after endothelial cell ablation was affected in MZ *jam2a; jam2b* mutant embryos in *etv2*$^{Gt(2A-Gal4)}$*; UAS:mCherry-NTR* background. In order to ablate all vascular endothelial cells, MTZ was added at 6 hpf and then washed out at approximately 45 hpf. Subsequently, embryos were analyzed at 72 hpf by confocal imaging. A substantial recovery was observed in the control *jam2a; jam2b* double heterozygous embryos,

while it was greatly reduced in MZ *jam2a, jam2b* mutant embryos (Fig. 7M-O). These results argue that the loss of SVF cells in MZ *jam2a; jam2b* mutants leads to defects in venous, intestinal and lymphatic vessel development, and is also associated with the reduced vascular recovery after EC ablation.

Hand2 **functions upstream of** *jam2b* **in SVF cell formation.** Recent work has identified that a bHLH transcription factor *hand2,* previously implicated in cardiac development, is also expressed in the lateral plate mesoderm along the yolk extension and labels mesothelial and pericardial progenitors, similar to *jam2b.*[27,28]. Intriguingly, *hand2-2A-Cre* based lineage tracing showed contribution to the venous and intestinal vasculature,[29] similar to our results obtained using *jam2b*[Gt(2A-Gal4)]*; UAS:Cre*. To determine if both genes are co-expressed in the same cells, we performed fluorescent hybridization chain reaction (HCR) analysis at 24 hpf. Indeed, expression of *hand2* and *jam2b* largely overlapped in the LPM region along the yolk extension (Fig. 8A-F). To test if *hand2* may function upstream of *jam2b,* we analyzed *jam2b* expression in the previously generated *hand2* mutants by in situ hybridization. Indeed, *jam2b* expression was absent in *hand2* mutants from most of the LPM, including the SVF-forming region along the yolk extension (Fig. 8G-J). Only the anterior domain of *jam2b* expression adjacent to the yolk was not affected in *hand2* mutants (Fig. 8G-J, white arrowheads). We then analyzed if *hand2* function is required for *etv2* expression in SVF cells. Indeed, *hand2* mutants showed complete absence of SVF cells at 30 hpf (Fig. 8K,L). In addition, *hand2* mutants showed reduced *etv2* expression in the PCV, which is consistent with its previously demonstrated requirement in the formation of venous endothelium.[30] These results argue that *hand2* functions upstream of *jam2b* in the specification of SVF cells within the trunk region of the LPM.

DISCUSSION

Our previous work has demonstrated that new *etv2+* progenitors emerge from the trunk region of the LPM along the yolk extension and contribute to the intestinal and axial vasculature.[15] Here we implicate *jam2a* and *jam2b* function in the emergence of SVF cells and their contribution to vasculature. Our results show that *jam2b* is expressed along the broad region of LPM, including the site where SVF cells emerge. Lineage tracing studies revealed that *jam2b*-expressing cells show major contribution to the intestinal vasculature, and a smaller contribution to the PCV and venous ISVs (Fig. 9). MZ j*am2a;jam2b* mutants show a reduced number of SVF cells and display defects in the intestinal and venous vessel development. We further show that *hand2* transcription factor functions upstream of *jam2b* in the specification of SVF cells within the trunk region of the LPM.

Previous studies have argued that internal organ-specific vessels, including the intestinal vessels, are derived from the PCV.[8-10] According to the recent model, individual cells migrate and assemble into the two vessels, SIA and SIV, which can further branch and give rise to interconnecting vessels, ICV.[8-10] Our recent study challenged this model and showed that SVF cells can contribute to intestinal vasculature.[15] However, it has not been clear if SVF cells have only a minor contribution to the intestinal vasculature. Our current results suggest that SVF cells may be the main source of the intestinal vasculature, while the PCV only has a minor contribution. *Etv2:Kaede* photoconversion showed that most cells in the intestinal vasculature emerge after the 24 hpf stage and, therefore, they were not present in the PCV at the time of the photoconversion. Although *jam2b*[Gt(2A-Gal4);] *UAS:Cre* based lineage tracing resulted in only a fraction of the PCV and intestinal vessels labeled with switched dsRed expression, this could be due to the relative inefficiency or mosaic expression of Cre. Based on our earlier time-lapse lineage tracing experiments, it appears that some SVF cells migrate directly to the SIA and SIV, while others integrate into the PCV first, and only afterwards exit the PCV and migrate to the intestinal vasculature.[15] Thus, our model can be reconciled with the previous observations if some SVF cells first integrate into the PCV and subsequently contribute to the intestinal vasculature. It is also possible that the earlier studies focused largely on the anterior portion of the intestinal vasculature, which forms the plexus-like structure and is largely derived from the

PCV. Our study focused on the mid-trunk region along the yolk extension, where SVF cells are positioned, which is posterior to the intestinal plexus, analyzed in the previous studies.

jam2b expression within the LPM is observed during somitogenesis stages. It is evident that this region gives rise to diverse cell types, including mesothelial, pectoral fin, pericardium and muscle. Similar contribution has also been demonstrated for hand2-positive LPM cells,[28] which largely overlap with jam2b expression. Only a small fraction of jam2b-positive cells contributes to the vascular endothelium. The cells fated to become endothelium are located at the edge of the medial portion of the jam2b expression domain in the trunk region, closest to the midline (Fig. 9). It is tempting to speculate that all jam2b cells are multipotent initially, yet they experience different signals based on their relative position. The cells closest to the midline in the trunk region become induced to differentiate into vascular endothelium. The inductive signal is currently unknown and will require further investigation. Interestingly, vegfaa is expressed in the region where intestinal vessels form.[8].= Furthermore, vegfaa mutants show reduced number of SVF cells, which correlates with the defects in the intestinal vessel formation,[8,15] suggesting that vegfaa could be a candidate signal.

During embryonic vasculogenesis arterial and venous progenitors originate from the LPM during somitogenesis stages.[11-14] While the arterial progenitors originate close to the midline, venous progenitors emerge from the more lateral portion of the LPM.[24] As we show here, early venous progenitors emerge adjacent to jam2b-positive LPM cells, which are positioned further laterally from the venous progenitors. While jam2b and etv2 expression overlaps in later forming SVF cells, it does not appear to overlap in early endothelial progenitors, which was also corroborated in our scRNA-seq data.[15,23] We also did not observe jam2b^{Gt(2A-Gal4)}; UAS:GFP expression in the PCV prior to 24 hpf stage, suggesting that jam2b may not be involved in early vasculogenesis. It appears that jam2b+ LPM cells maintain vasculogenic potential until later stages past 24 hpf.

The mechanism of how jam2a; jam2b mutants regulate SVF cell specification is unclear. A previous study has demonstrated that antibodies against murine Jam-B (Jam2) inhibited angiogenesis in vitro and interfered with the VEGFR2 signaling pathway.[31] Our previous data indicate that VEGF signaling is required to upregulate etv2 expression during early vasculogenesis, and it is also required for SVF cell emergence.[15,32] It is possible that Jam2

signaling is required to promote VEGF signaling, which will have to be investigated in future studies.

Previous studies have demonstrated that *hand2* function is required for the PCV development. *npas4l* and *etv2* expression were reduced in the venous progenitors, arguing that *hand2* functions upstream of these transcription factors.[30,33] We show here that SVF cells are also absent in *hand2* mutants. It is possible that *hand2* directly regulates *npas4l* and /or *etv2* expression. Alternatively, *hand2* may function through *jam2b* and indirectly be involved in SVF specification through VEGF or a different pathway.

In summary, this study demonstrates that *jam2b*-positive LPM serves as a major source of internal organ-specific vascular progenitors and identifies the functional requirement for *jam2a* and *jam2b* in promoting *etv2* expression and vasculogenesis. Because analogous late-forming vascular progenitors have been also identified in the mammalian embryos,[15] it is likely that the mechanisms regulating their emergence are also evolutionarily conserved. These results will be important to understand the molecular mechanisms involved in the formation of different vascular beds and will promote the development of new approaches aimed at vascular regeneration.

MATERIALS AND METHODS

Zebrafish lines. The following previously established lines were used in the study: wild-type AB (Zebrafish International Resource Center), *Tg(kdrl:mCherry)^{ci5},*[34] *Tg(fli1a:GFP)^{y1},*[35] *hand2^{s6},*[27] *etv2$^{Gt(2A-Gal4)ci32}$;UAS:mCherry-NTR,*[36] *etv2$^{Gt(2A-Venus)}$,*[15] *Tg(actb2:loxP-BFP-loxP-dsRed)^{sd27},*[37] *Tg(kdrl:GFP)^{s843},*[14] *TgBAC(prox1a:KalT4-UAS:uncTagRFP)^{nim5}* [38] (abbreviated as *prox1:RFP*), *Tg(lyve1b:dsRed)^{nz101}.*[39]

Generation of jam2b$^{Gt(2A-Gal4)}$ knock-in line. To generate the *jam2b$^{Gt(2A-Gal4)ci55}$;UAS:GFP* knock-in line (further abbreviated as *jam2b$^{Gt(2A-Gal4)}$*; UAS:GFP), a gBlock Gene Fragment was made by gene synthesis (IDT) which included *jam2b* sgRNA site, the P2A sequence, *Gal4* DNA binding domain and poly-A tail. This fragment was then A-tailed using dATP and Taq DNA Polymerase (NEB, cat. no. M0273) followed by TA cloning into TOPO dual vector using manufacturer's instructions (ThermoFisher, cat. no. 450640). A mixture of 300 pg *Cas9* mRNA and 50 pg *jam2b-sgRNA-P2A-Gal4* plasmid DNA was injected into

Tg(*UAS:GFP*)[40] embryos at 1-cell stage. Embryos showing any specific *GFP* expression were raised through adulthood and screened to identify a founder which produced progeny with a specific expression pattern.

   Generation of jam2a and jam2b mutant lines. To generate *jam2b*[sfl2] line, two synthetic gRNAs (Synthego) targeting *jam2b* promoter region were co-injected with Cas9 protein (NEB, EnGen Spy Cas9 NLS, Cat No. M0646) into zebrafish embryos at 1-cell stage. The following sgRNA sequences were used: GCAATTCTGGAAGAAGTCAA and AATAAGCAGGAGAAGAGCAG. Successful deletion was confirmed using the following primers outside of the deletion site: *jam2b_P3.4.5_F:* GCATCACTCAAAGGCAAATG and *jam2b_P1.2_R:* ACAGTTCCCACCATTTCAGC. A single founder with 1,071 bp deletion, which encompasses *jam2b* proximal promoter region and a portion of exon 1, was identified and used to establish the mutant line. The genomic sequence flanking the deleted region is shown here:

TATAGACCGGGCAGATTAGCCATTG………TCTTCTCCTGCTTATTTACAGTAAG.

   To generate *jam2a*[sfl3] line, two synthetic gRNAs (Synthego) were designed to delete *jam2a* coding sequence with the following sequences: TCTGTGTGCCTCTAGGTGAG and CAGAGCACAAACCCACGCAA. Both gRNAs were co-injected together with Cas9 protein into one-cell stage zebrafish embryos, homozygous for *jam2b*[sfl2] mutation. Successful deletion was confirmed using the following primers that flank the deletion site: *jam2a-sg1_2_F:* CAAACACCTGTGACCTTACAAAA and *jam2a-sg1_2_R:* TATCAATGGCAACACAGTGG. A founder was identified with the deletion of 15,771 bp, which includes the entire *jam2a* coding sequence, and used to establish the stable line for further studies. The genomic sequence flanking the deleted region is shown here:

CCTACAGGTGGTTTGTCAAGCCCAC……….AAAGGTAGGGAGAACATGCAAACTC

   Generation of UAS:Cre and UAS:CreERT2 lines. To generate *UAS:Cre*[sfl4] and *UAS:CreERT2*[sfl5] lines, the gene fragment containing 5xUAS sequence, *hbae3* 5'and 3'UTR sequences, zebrafish Kozak sequence,[41] NLS-Cre, and NLS-CreERT2 codon optimized for zebrafish expression using iCodon approach,[42] SV40 polyA site, and attL4 and attL3 sequences specific to LR recombinase, was synthesized by gene synthesis and subcloned into pUC57 vector (Genscript). LR recombination into the Gateway Destination vector containing Tol2 transposase sites and lens specific alpha crystallin-dsRed sequence[43] was performed using LR recombinase

(ThermoFisher). Zebrafish embryos were co-injected with 25-40pg of each construct together with 75-150 pg *tol2* mRNA. Founders were screened for the presence of lens fluorescence in their progeny. Recombination activity was tested by mating to *jam2b[Gt(2A-Gal4)]; Tg(actb2:BFP-loxP-dsRed-loxP)* driver / reporter line. One of each lines, *UAS:Cre* and *UAS:CreERT2,* which showed good recombination activity, was selected and used for all subsequent experiments.

Confocal microscopy imaging and image processing. For confocal imaging, embryos were anesthetized in 0.016% Tricaine solution, mounted in 0.6% low melting point agarose and imaged using Nikon Eclipse or Nicon AX confocal microscope. Series of z-slices were acquired using NIS Elements software. Nikon AI Denoise function was used to reduce the image noise. Maximum intensity projection was generated using Fiji or Nikon Elements software. Images were adjusted using Levels function in Adobe Photoshop to enhance the brightness and contrast. In all cases, the same adjustments were performed in control and experimental embryo samples.

Hybridization chain reaction (HCR). Whole mount fluorescent *in situ* hybridization was performed using the hybridization chain reaction version 3.0 as previously described.[44] Fluorescently labeled RNA probes for *etv2*, *prrx1a, hand2* and *jam2b* were purchased from Molecular Instruments, Inc.

Whole mount *in situ* hybridization (ISH) was performed as previously described.[45] DIG-labeled antisense RNA probes for *jam2b,*[15] *npas4l,*[15,46] *etv2/etsrp,*[47] *scl/tal1*[48] were synthesized as previously described. Following ISH, embryos were whole mounted in 0.6% low melting point agarose and imaged under light microscopy at 10X or 20X with Nikon Eclipse compound microscope. A series of z-slices were acquired using NIS Elements software to produce extended focus projected images.

4-hydroxitamoxifen (OHT) treatments. To induce tissue-specific recombination at a specific time point, *jam2b[Gt(2A-Gal4)];UAS:CreERT2* line was crossed with the *Tg(actb2-loxP-BFP-loxP-dsRed; fli1:GFP)* reporter line. Embryos were treated with 10 μM 4-OHT (Sigma-Aldrich) solution starting at 7 or 22 hpf stage. Alternatively, to analyze the timing requirement, embryos were treated at 7 hpf stage and subsequently washed at 22 hpf by rinsing a few times in embryo water. Confocal z-stacks were acquired at 3 dpf and analyzed with NIS Elements Software. Recombination was quantified by counting dsRed-positive cells within vasculature.

Endothelial cell (EC) count. To assess EC number in *jam2a;jam2b* double homozygous mutants and control heterozygous embryos, confocal images were analyzed using NIS Elements

Imaging Software. An area consisting of 5 intersegmental vessels in the trunk region was selected, and ECs located at the bottom of the posterior cardinal vein (PCV) were manually counted.

Subintestinal vessel and thoracic duct (TD) measurements. To evaluate SIA, SIV and TD in *jam2a; jam2b* double homozygous mutants and control heterozygous embryos, confocal images were analyzed using NIS Elements 5.41.02 Imaging Software. SIA, SIV and TD were categorized into normal and abnormal groups according to morphological characteristics. Normal vessels were considered uniform and without any gaps throughout the whole embryo; everything else was considered abnormal.

Chemical cell ablation. To assess vascular tissue ablation and regeneration, *jam2a;jam2b* homozygous mutant zebrafish in *kdrl:GFP* background were crossed with *jam2a;jam2b* homozygous mutants carrying the *etv2$^{Gt(2A-Gal4)}$; UAS:NTR-mCherry* transgene. As a control group, heterozygous embryos were obtained by crossing *jam2a; jam2b* homozygous mutants carrying the *etv2$^{Gt(2A-Gal4)}$; UAS:NTR-mCherry* transgene with wild-type *kdrl:GFP*. Embryos were collected and treated with 10 mM metronidazole (MTZ) dissolved in 0.1 % DMSO at 6 or 10 hpf stages. At 2 dpf, embryos were dechorionated, thoroughly washed, and transferred to fresh embryo water to allow for recovery. Confocal imaging was performed at 3 dpf to assess vascular regeneration. NIS-Elements Imaging Software (version 5.41.02) was used to quantify tissue ablation and recovery. The total length of regenerated vessels was calculated by measuring and adding the lengths of all recovered vessels based on mCherry and GFP expression.

Statistical analysis. Data representing graphs were generated and statistical analyses (t-test and Fisher's exact test) were performed using GraphPad Prism software.

Single-cell RNA-seq analysis. Transgenic *etv2$^{Gt(2A-Venus)}$* fish were crossed to *jam2b$^{Gt(2A-Gal4)}$;UAS:NTR-mCherry* fish. Double-positive embryos expressing both *Venus* and *mCherry* fluorescent proteins were selected at 36 hpf. Embryos were dechorionated and immediately transferred into a 1.5 ml Eppendorf tube for the generation of single-cell suspension. Whole embryos were then dissociated into a single-cell suspension by cold protease tissue dissociation protocol [49]. Cells expressing *Venus* only, *mCherry* only, and both (*Venus* and *mCherry*) were sorted using a BD/FACSAria II flow cytometer cell sorter. After sorting, cells expressing both

Venus and *mCherry* were equally divided and added into the *Venus* only and *mCherry* only populations manually. Suspensions of ~10,000 single Venus and mCherry-positive cells were loaded separately onto the 10XGenomics Single Cell 3' (V3 chemistry) chip. 12 cDNA amplification cycles were used to generate cDNA. Samples were sequenced on an Illumina NovaSeq 6000 instrument (Illumina, San Diego, CA) running an S2 flow cell with parameters as follows: Read1, 28 cycles; Index Read1: 8 cycles; Read 2: 91 cycles at the CCHMC DNA Sequencing core. The fastq files obtained from the sequencing core were then mapped to the Danio rerio genome (version zv11) to generate single-cell feature counts using Cell Ranger version 3.0.2. Counts were generated using fastq files from each of the populations individually. Utilizing R packages: Seurat,[50] ggplot2,[51] dplyr,[52] reshape2,[53] and tidyverse[54] the data was filtered for high quality cells with 200-3500 features and less than 5% mitochondrial gene content. The datasets were then integrated and the figure plots were made using custom written R code.[55-57] To find a stable cluster resolution we tested resolutions between 0-3 in 0.1 increments and decided on 1.6, revealing the SVF cells in cluster 15. We then selected clusters for trajectory analysis using the Monocle3 package. The UMAP, feature plots, and violin plots were made using Seurat, and the lineage plot was made using Monocle3.[25] Cluster annotation was done manually based on the top marker gene expression. In some cases, AI ChatGPT 5 assistance was used to help with the cell type prediction based on 40-60 top marker gene expression, which was subsequently verified manually based on published gene expression patterns. Single-cell RNA-seq dataset has been submitted to NCBI GEO database under accession number: GSE179926.

Acknowledgements. This research was supported by NIH R01 HL153005 to S.S., a supplemental postdoctoral diversity award R01 HL163161-S1 to R.D., AHA 19POST34400016 award to S.M. and AHA 19POST34450021 award to S.G. We thank Olivia Nester and Kanwal Jussa for their assistance with zebrafish experiments.

REFERENCES

1. Gouysse, G., Couvelard, A., Frachon, S., Bouvier, R., Nejjari, M., Dauge, M.C., Feldmann, G., Henin, D., and Scoazec, J.Y. (2002). Relationship between vascular development and vascular differentiation during liver organogenesis in humans. J Hepatol *37*, 730-740. 10.1016/s0168-8278(02)00282-9.

2. Kingsbury, J.W., Alexanderson, M., and Kornstein, E.S. (1956). The development of the liver in the chick. Anat Rec *124*, 165-187. 10.1002/ar.1091240204.

3. Matsumoto, K., Yoshitomi, H., Rossant, J., and Zaret, K.S. (2001). Liver organogenesis promoted by endothelial cells prior to vascular function. Science *294*, 559-563. 10.1126/science.1063889.

4. Sherer, G.K. (1991). The Development of the Vascular System (Karger).

5. Goldman, O., Han, S., Hamou, W., Jodon de Villeroche, V., Uzan, G., Lickert, H., and Gouon-Evans, V. (2014). Endoderm generates endothelial cells during liver development. Stem Cell Reports *3*, 556-565. 10.1016/j.stemcr.2014.08.009.

6. Perez-Pomares, J.M., Carmona, R., Gonzalez-Iriarte, M., Macias, D., Guadix, J.A., and Munoz-Chapuli, R. (2004). Contribution of mesothelium-derived cells to liver sinusoids in avian embryos. Dev Dyn *229*, 465-474. 10.1002/dvdy.10455.

7. Zhang, H., Pu, W., Tian, X., Huang, X., He, L., Liu, Q., Li, Y., Zhang, L., He, L., Liu, K., et al. (2016). Genetic lineage tracing identifies endocardial origin of liver vasculature. Nat Genet *48*, 537-543. 10.1038/ng.3536.

8. Koenig, A.L., Baltrunaite, K., Bower, N.I., Rossi, A., Stainier, D.Y., Hogan, B.M., and Sumanas, S. (2016). Vegfa signaling promotes zebrafish intestinal vasculature development through endothelial cell migration from the posterior cardinal vein. Dev Biol *411*, 115-127. 10.1016/j.ydbio.2016.01.002.

9. Goi, M., and Childs, S.J. (2016). Patterning mechanisms of the sub-intestinal venous plexus in zebrafish. Dev Biol *409*, 114-128. 10.1016/j.ydbio.2015.10.017.

10. Hen, G., Nicenboim, J., Mayseless, O., Asaf, L., Shin, M., Busolin, G., Hofi, R., Almog, G., Tiso, N., Lawson, N.D., and Yaniv, K. (2015). Venous-derived angioblasts generate organ-specific vessels during zebrafish embryonic development. Development *142*, 4266-4278. 10.1242/dev.129247.

11. Eriksson, J., and Lofberg, J. (2000). Development of the hypochord and dorsal aorta in the zebrafish embryo (Danio rerio). J Morphol *244*, 167-176. 10.1002/(SICI)1097-4687(200006)244:3<167::AID-JMOR2>3.0.CO;2-J.

12. Fouquet, B., Weinstein, B.M., Serluca, F.C., and Fishman, M.C. (1997). Vessel patterning in the embryo of the zebrafish: guidance by notochord. Dev Biol *183*, 37-48. 10.1006/dbio.1996.8495.

13. Childs, S., Chen, J.N., Garrity, D.M., and Fishman, M.C. (2002). Patterning of angiogenesis in the zebrafish embryo. Development *129*, 973-982. 10.1242/dev.129.4.973.

14. Jin, S.W., Beis, D., Mitchell, T., Chen, J.N., and Stainier, D.Y. (2005). Cellular and molecular analyses of vascular tube and lumen formation in zebrafish. Development *132*, 5199-5209. 10.1242/dev.02087.

15. Metikala, S., Warkala, M., Casie Chetty, S., Chestnut, B., Rufin Florat, D., Plender, E., Nester, O., Koenig, A.L., Astrof, S., and Sumanas, S. (2022). Integration of vascular progenitors into functional blood vessels represents a distinct mechanism of vascular growth. Dev Cell *57*, 767-782 e766. 10.1016/j.devcel.2022.02.015.

16. Metikala, S., Casie Chetty, S., and Sumanas, S. (2021). Single-cell transcriptome analysis of the zebrafish embryonic trunk. PLoS One *16*, e0254024. 10.1371/journal.pone.0254024.

17.	Ebnet, K. (2017). Junctional Adhesion Molecules (JAMs): Cell Adhesion Receptors With Pleiotropic Functions in Cell Physiology and Development. Physiol Rev *97*, 1529-1554. 10.1152/physrev.00004.2017.

18.	Kobayashi, I., Kobayashi-Sun, J., Hirakawa, Y., Ouchi, M., Yasuda, K., Kamei, H., Fukuhara, S., and Yamaguchi, M. (2020). Dual role of Jam3b in early hematopoietic and vascular development. Development *147*. 10.1242/dev.181040.

19.	Kobayashi, I., Kobayashi-Sun, J., Kim, A.D., Pouget, C., Fujita, N., Suda, T., and Traver, D. (2014). Jam1a-Jam2a interactions regulate haematopoietic stem cell fate through Notch signalling. Nature *512*, 319-323. 10.1038/nature13623.

20.	Powell, G.T., and Wright, G.J. (2012). Genomic organisation, embryonic expression and biochemical interactions of the zebrafish junctional adhesion molecule family of receptors. PLoS One *7*, e40810. 10.1371/journal.pone.0040810.

21.	Thisse, B., and Thisse, C. (2004). Fast Release Clones: A High Throughput Expression Analysis.

22.	Auer, T.O., Duroure, K., De Cian, A., Concordet, J.P., and Del Bene, F. (2014). Highly efficient CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent DNA repair. Genome Res *24*, 142-153. 10.1101/gr.161638.113.

23.	Chestnut, B., Casie Chetty, S., Koenig, A.L., and Sumanas, S. (2020). Single-cell transcriptomic analysis identifies the conversion of zebrafish Etv2-deficient vascular progenitors into skeletal muscle. Nat Commun *11*, 2796. 10.1038/s41467-020-16515-y.

24.	Kohli, V., Schumacher, J.A., Desai, S.P., Rehn, K., and Sumanas, S. (2013). Arterial and venous progenitors of the major axial vessels originate at distinct locations. Dev Cell *25*, 196-206. 10.1016/j.devcel.2013.03.017.

25.	Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos, S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional landscape of mammalian organogenesis. Nature *566*, 496-502. 10.1038/s41586-019-0969-x.

26.	Powell, G.T., and Wright, G.J. (2011). Jamb and jamc are essential for vertebrate myocyte fusion. PLoS Biol *9*, e1001216. 10.1371/journal.pbio.1001216.

27.	Yelon, D., Ticho, B., Halpern, M.E., Ruvinsky, I., Ho, R.K., Silver, L.M., and Stainier, D.Y. (2000). The bHLH transcription factor hand2 plays parallel roles in zebrafish heart and pectoral fin development. Development *127*, 2573-2582. 10.1242/dev.127.12.2573.

28.	Prummel, K.D., Crowell, H.L., Nieuwenhuize, S., Brombacher, E.C., Daetwyler, S., Soneson, C., Kresoja-Rakic, J., Kocere, A., Ronner, M., Ernst, A., et al. (2022). Hand2 delineates mesothelium progenitors and is reactivated in mesothelioma. Nat Commun *13*, 1677. 10.1038/s41467-022-29311-7.

29.	Ming, Z., Liu, F., Moran, H.R., Lalonde, R.L., Adams, M., Restrepo, N.K., Joshi, P., Ekker, S.C., Clark, K.J., Friedberg, I., et al. (2025). Lineage labeling with zebrafish hand2 Cre and CreERT2 recombinase CRISPR knock-ins. Dev Dyn. 10.1002/dvdy.70022.

30.	Perens, E.A., Garavito-Aguilar, Z.V., Guio-Vega, G.P., Pena, K.T., Schindler, Y.L., and Yelon, D. (2016). Hand2 inhibits kidney specification while promoting vein formation within the posterior mesoderm. Elife *5*. 10.7554/eLife.19941.

31.	Meguenani, M., Miljkovic-Licina, M., Fagiani, E., Ropraz, P., Hammel, P., Aurrand-Lions, M., Adams, R.H., Christofori, G., Imhof, B.A., and Garrido-Urbani, S. (2015). Junctional adhesion molecule B interferes with angiogenic VEGF/VEGFR2 signaling. FASEB J *29*, 3411-3425. 10.1096/fj.15-270223.

32.	Casie Chetty, S., Rost, M.S., Enriquez, J.R., Schumacher, J.A., Baltrunaite, K., Rossi, A., Stainier, D.Y., and Sumanas, S. (2017). Vegf signaling promotes vascular endothelial differentiation by modulating etv2 expression. Dev Biol *424*, 147-161. 10.1016/j.ydbio.2017.03.005.

33. Perens, E.A., and Yelon, D. (2025). Drivers of vessel progenitor fate define intermediate mesoderm dimensions by inhibiting kidney progenitor specification. Dev Biol *517*, 126-139. 10.1016/j.ydbio.2024.09.008.

34. Proulx, K., Lu, A., and Sumanas, S. (2010). Cranial vasculature in zebrafish forms by angioblast cluster-derived angiogenesis. Dev Biol *348*, 34-46. 10.1016/j.ydbio.2010.08.036.

35. Lawson, N.D., and Weinstein, B.M. (2002). In vivo imaging of embryonic vascular development using transgenic zebrafish. Dev Biol *248*, 307-318. 10.1006/dbio.2002.0711.

36. Chestnut, B., and Sumanas, S. (2020). Zebrafish etv2 knock-in line labels vascular endothelial and blood progenitor cells. Dev Dyn *249*, 245-261. 10.1002/dvdy.130.

37. Bertrand, J.Y., Chi, N.C., Santoso, B., Teng, S., Stainier, D.Y., and Traver, D. (2010). Haematopoietic stem cells derive directly from aortic endothelium during development. Nature *464*, 108-111. 10.1038/nature08738.

38. Dunworth, W.P., Cardona-Costa, J., Bozkulak, E.C., Kim, J.D., Meadows, S., Fischer, J.C., Wang, Y., Cleaver, O., Qyang, Y., Ober, E.A., and Jin, S.W. (2014). Bone morphogenetic protein 2 signaling negatively modulates lymphatic development in vertebrate embryos. Circ Res *114*, 56-66. 10.1161/CIRCRESAHA.114.302452.

39. Okuda, K.S., Astin, J.W., Misa, J.P., Flores, M.V., Crosier, K.E., and Crosier, P.S. (2012). lyve1 expression reveals novel lymphatic vessels and new mechanisms for lymphatic vessel development in zebrafish. Development *139*, 2381-2391. 10.1242/dev.077701.

40. Asakawa, K., Suster, M.L., Mizusawa, K., Nagayoshi, S., Kotani, T., Urasaki, A., Kishimoto, Y., Hibi, M., and Kawakami, K. (2008). Genetic dissection of neural circuits by Tol2 transposon-mediated Gal4 gene and enhancer trapping in zebrafish. Proc Natl Acad Sci U S A *105*, 1255-1260. 10.1073/pnas.0704963105.

41. Grzegorski, S.J., Chiari, E.F., Robbins, A., Kish, P.E., and Kahana, A. (2014). Natural variability of Kozak sequences correlates with function in a zebrafish model. PLoS One *9*, e108475. 10.1371/journal.pone.0108475.

42. Diez, M., Medina-Munoz, S.G., Castellano, L.A., da Silva Pescador, G., Wu, Q., and Bazzini, A.A. (2022). iCodon customizes gene expression based on the codon composition. Sci Rep *12*, 12126. 10.1038/s41598-022-15526-7.

43. Villefranc, J.A., Amigo, J., and Lawson, N.D. (2007). Gateway compatible vectors for analysis of gene function in the zebrafish. Dev Dyn *236*, 3077-3087. 10.1002/dvdy.21354.

44. Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. Development *145*. 10.1242/dev.165753.

45. Jowett, T. (1999). Analysis of protein and gene expression. Methods Cell Biol *59*, 63-85. 10.1016/s0091-679x(08)61821-x.

46. Reischauer, S., Stone, O.A., Villasenor, A., Chi, N., Jin, S.W., Martin, M., Lee, M.T., Fukuda, N., Marass, M., Witty, A., et al. (2016). Cloche is a bHLH-PAS transcription factor that drives haemato-vascular specification. Nature *535*, 294-298. 10.1038/nature18614.

47. Sumanas, S., Jorniak, T., and Lin, S. (2005). Identification of novel vascular endothelial-specific genes by the microarray analysis of the zebrafish cloche mutants. Blood *106*, 534-541. 10.1182/blood-2004-12-4653.

48. Liao, E.C., Paw, B.H., Oates, A.C., Pratt, S.J., Postlethwait, J.H., and Zon, L.I. (1998). SCL/Tal-1 transcription factor acts downstream of cloche to specify hematopoietic and vascular progenitors in zebrafish. Genes Dev *12*, 621-626. 10.1101/gad.12.5.621.

49. Potter, A.S., and Steven Potter, S. (2019). Dissociation of Tissues for Single-Cell Analysis. Methods Mol Biol *1926*, 55-62. 10.1007/978-1-4939-9021-4_5.

50. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*, 3573-3587 e3529. 10.1016/j.cell.2021.04.048.

51. Wickham, H. (2016). ggplot2 : Elegant Graphics for Data Analysis. Use R!,. 2nd ed. Springer International Publishing : Imprint: Springer,.

52. Wickham H, A.M., Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A, Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D, Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H (2019). Welcome to the tidyverse. Journal of Open Source Software. 10.21105/joss.01686.

53. Wickham, H. (2007). Reshaping Data with the reshape Package. Journal of Statistical Software *21(12)*, 1-20.

54. Wickham H, F.R., Henry L, Müller K, Vaughan D (2023). dplyr: A Grammar of Data Manipulation.

55. Hao, Y., Stuart, T., Kowalski, M.H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., and Satija, R. (2024). Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol *42*, 293-304. 10.1038/s41587-023-01767-y.

56. R Core Team (2025). R: A language and environment for statistical computing.

57. Rstudio Team (2020). RStudio: Integrated Development for R (RStudio, Inc.).

FIGURE LEGENDS

Figure 1. Characterization of *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP* expression at 1-5 dpf. (A-F) *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP* shows strong GFP expression within the lateral plate mesoderm (LPM) along the yolk extension at 26 hpf (white arrows, A,B), which becomes presumptive mesothelium at 3-5 dpf (white arrows, C-F). GFP expression is also apparent in a subset of skeletal muscle cells (white arrowheads, A,C,E), craniofacial muscle (yellow arrows, E,F) and in the cells surrounding the eye (yellow arrowheads, E,F). (G-I) Co-expression of *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP* and *etv2* (red) in the lateral plate mesoderm of the trunk region at 30 hpf. Note that GFP and *etv2* expression overlaps in the SVF cells (arrowheads), located in the most dorsal region of GFP expression domain. (J-R) Expression of *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP* in the LPM along the yolk extension and in muscle cells from 30 to 66 hpf. A few GFP-positive cells are present in the PCV and overlap with vascular endothelial marker *kdrl:mCherry* expression (white arrowheads and insets). (S-U) Analysis of *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP* expression in *lyve1b:dsRed* reporter background at 4 dpf, which labels PCV and the thoracic duct. Note GFP-positive cells in the TD (arrowheads). Bright GFP expression in the skeletal muscle fibers is also apparent. PCV, posterior cardinal vein; DA, dorsal aorta; TD, thoracic duct. Lateral views of the trunk region, anterior is to the left. Scale bars: 100 μm.

Figure 2. Jam2b-positive cells are derived from the lateral plate mesoderm and contribute to the vasculature. (A-F) Time-lapse images of *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP ; Tg(kdrl:mCherry)* embryo starting at 24 hpf show two GFP-positive cells (arrowheads), which migrate dorsally, proliferate, integrate into the PCV and initiate *kdrl:mCherry* expression (see Movie 1). Time is in minutes after 24 hpf. PCV, posterior cardinal vein; DA, dorsal aorta. Scale bar: 50 μm.

(G-I), HCR against *etv2* in a *jam2b^(Gt(2A-Gal4))*;*UAS:GFP* embryo at the 18-somite stage showing

jam2b expression (green) lateral and ventral to *etv2* positive vascular progenitors (red). (J-L)

HCR against *prrx1a* in a *jam2b^(Gt(2A-Gal4))*;*UAS:GFP* embryo at the 18-somite stage showing

colocalization of *prrx1a* mRNA and GFP fluorescence. Dorsolateral view of the trunk region is

shown, anterior is to the left.

Figure 3. Genetic lineage tracing analysis of *jam2b*+ cell contribution to vasculature. (A)

Schematic diagram of recombination in *jam2b^(Gt(2A-Gal4);)* *UAS:Cre; actb2-loxP-BFP-loxP-dsRed*

embryos. Ubiquitous BFP expression switches to dsRed in Cre-positive cells. (B-J) Lineage

tracing analysis in embryos at 3 dpf obtained by crossing *jam2b^(Gt(2A-Gal4))* and *UAS:Cre;*

actb2:loxP-BFP-loxP-dsRed; fli1:GFP parents. Red cells, indicating the genetic switch, are

observed in the mesothelial layer surrounding the yolk (arrow, B,E), skeletal muscle (yellow

arrowheads, B), and the epicardial (arrows, C,D) and myocardial (arrowheads, C,D) heart layers.

Switched cells in the vasculature are observed in the venous ISVs (white arrowheads, B,E,F,G),

the PCV (blue arrowheads, F,G) and SIA (arrow, I,J). A control embryo negative for *jam2b^(Gt(2A-*

Gal4)) is shown in (H) to indicate autofluorescence. (F,G) show larger magnification views of

(B,E). (K) Quantification of endothelial cell labeling in different vascular beds in *jam2b^(Gt(2A-Gal4))*;

UAS:Cre; actb2-loxP-BFP-loxP-dsRed embryos at 3 dpf. (L) Quantification of endothelial cell

labeling in different vascular beds in *jam2b^(Gt(2A-Gal4))*; *UAS:CreERT2; actb2-loxP-BFP-loxP-*

dsRed embryos at 3 dpf. 4-OHT was added starting at 7 hpf or 22 hpf stages. In a third group of

embryos, 4-OHT was added at 7 hpf and washed out at 22 hpf stages. Note that cell contribution

to SIV could not be analyzed reliably due to very strong red fluorescence in the mesothelial layer

surrounding the yolk.

Figure 4. Intestinal vasculature in the middle and posterior trunk region forms largely by new vasculogenesis. (A,B). *etv2:Kaede* embryos were photoconverted from green to red at 24 hpf and analyzed at 4 dpf. Merged and red only channels are shown. (C-E) Higher magnification images of the boxed area in (A,B). Merged, red and green channels are shown. Note that SIA and SIV are comprised mostly from green only cells (white arrowheads), indicating that they have originated after the photoconversion. (F) Quantification of green only (new) and green+red (pre-existing) cells in the SIA and SIV at 4 dpf. The analysis focused on the middle and posterior trunk region over the yolk extension area.

Figure 5. scRNA-seq analysis of cells from *jam2b*$^{Gt(2A-Gal4)}$*;UAS:mCherry-NTR; etv2*$^{Gt(2A-Venus)}$ embryos at 36 hpf. (A) UMAP plot showing 37 cell clusters identified by Seurat analysis. LPM, lateral plate mesoderm; SVF, secondary vascular field; RBC, red blood cells. (B) Dot plot of selected marker gene expression level in each cell cluster. (C) 2D UMAP plot showing the distribution of Venus and mCherr*y* positive cells. Note that Venus-positive cells mostly correspond to different vascular endothelial lineages, while mCherry-positive cells label different subsets of LPM. (D) Feature plots showing the expression of SVF cell markers, enriched in the cluster 15. (E) Violin plots show SVF cell marker expression in different clusters. (F) Pseudotime trajectory of vascular endothelial and SVF clusters made using Monocle 3. Note that SVF cells (cluster 15) can transition into both venous and arterial cell types.

Figure 6. SVF cell number is reduced in maternal-zygotic (MZ) *jam2a; jam2b* mutant embryos. (A-C) In situ hybridization analysis (ISH) for *etv2* expression in SVF cells (arrows, A,B) in MZ *jam2a-/-; jam2b-/-* embryos and control *jam2a+/-; jam2b+/-* embryos in *kdrl:GFP* background, obtained by crossing double homozygous *jam2a-/-; jam2b-/-; kdrl:GFP* adults to wild-type *kdrl:GFP*. (D-I) ISH analysis for *tal1* and *npas4l* expression in SVF cells (arrows) in

MZ *jam2a-/-; jam2b-/-* embryos and control embryos in *fli1:GFP* background. Controls were obtained by mating *jam2a+/-; jam2b-/-* embryos to wild-type *fli1:GFP,* resulting in a mixture of *jam2a+/+; jam2b+/-* and *jam2a+/-; jam2b+/-* genotypes. ***p<0.001, ****p<0.0001, two-tailed t-test, error bars show ±s.d. Data are combined from four (C) or two (F,I) independent replicate experiments. Het labels in (C,F,I) refer to double heterozygous in (C) and a mixture of *jam2a+/+; jam2b+/-* and *jam2a+/-; jam2b+/-* genotypes in (F,I), while Mut label refers to *jam2a-/-; jam2b-/-* MZ mutant embryos.

Figure 7. Analysis of vascular defects in MZ *jam2a; jam2b* mutant embryos. (A-C) Intersegmental vessel extension analysis in MZ *jam2a-/-;jam2b-/-* and *jam2a+/-;jam2b+/-* embryos in *kdrl:GFP* background at 3 dpf. Note that some ISVs are truncated in MZ *jam2a; jam2b* mutant embryos (arrow, B, an inset shows higher magnification view of the boxed trunk region). (D-F) Endothelial cell number in the PCV is reduced in MZ *jam2a; jam2b* mutant embryos compared to *jam2a+/-;jam2b+/-* controls at 3 dpf. Endothelial cells were counted in the floor of the PCV (arrowheads) in the selected area of the trunk region based on *fli1:GFP* expression. (G-I) SIA and SIV are fragmented and show increased incidence of gaps in MZ *jam2a;jam2b* mutant embryos compared to heterozygous controls at 3 dpf. Analysis was performed in the background of *etv2*[Gt(2A-Gal4)]; *UAS:mCherry-NTR* transgene, which shows expression in all vasculature. Note the underdeveloped SIA (arrowheads) and gaps in SIV (arrow) in *jam2a;jam2b* mutants. Higher magnification views are shown in the insets. Diagrams in (I) show percentage of embryos with gaps in SIA (left) and SIV (right). (J-L) Thoracic duct analysis in *prox1:RFP; fli1:GFP* transgenic embryos at 6 dpf. Note the gaps in the thoracic duct (arrows) observed in MZ *jam2a;jam2b* mutant embryos. (L) Shows the percentage of embryos with normal or fragmented thoracic duct. (M-O) Vascular recovery analysis after endothelial cell

ablation. *jam2a;jam2b* mutant and control heterozygous embryos in *etv2* $^{Gt(2A-Gal4)}$*; UAS:NTR-mCherry* background were treated with MTZ starting at 6 hpf to ablate ECs and subsequently washed out at 45 hpf. A partial vascular recovery is observed in the control embryos, while it is greatly reduced in MZ *jam2a;jam2b* mutants. (O) shows length measurements of the recovered vasculature. (C,F,O) graphs show two tailed t-test, while (I,L) show Fisher's exact test. *p<0.05, **p<0.01, ****p<0.0001. DA, dorsal aorta; PCV, posterior cardinal vein; SIA, supraintestinal artery; SIV, subintestinal vein.

Figure 8. *hand2* is required for *jam2b* expression in the lateral plate mesoderm and SVF cells. (A-F) HCR analysis for *hand2* (red) and *jam2b* (magenta) expression in *kdrl:GFP* embryos at 24 hpf. Anterior (A-C) and posterior (D-F) regions of the trunk are shown. Note the overlapping expression in the LPM along the yolk and yolk extension (arrows), which does not overlap with *kdrl:GFP* expression. (G-J) Whole-mount in situ hybridization analysis for *jam2b* expression in *hand2* mutant and wt control embryos at 26 hpf. Lateral (G,H) and dorsal (I,J) views. Note that *jam2b* expression in *hand2* mutants is absent from the posterior domain of the LPM along the yolk extension (black arrowheads), while the anterior portion of the LPM (white arrowheads) is only mildly affected. (K,L) Whole-mount in situ hybridization for *etv2* expression in *hand2* mutant and control wild-type embryos at 30 hpf. Note that *etv2* expression in SVF cells (arrows) is absent in *hand2* mutants. In addition, its expression in the PCV (arrowheads) is greatly reduced or absent. ISV, intersegmental vessel; DA, dorsal aorta; PCV, posterior cardinal vein. Scale bars: 100 μm.

Figure 9. A diagram illustrating the emergence of SVF cells from *jam2b*-positive lateral plate mesoderm. SVF cells contribute to the intestinal vasculature, including the supraintestinal artery (SIA), subintestinal vein (SIV), as well as the posterior cardinal vein (PCV).

Figure 1. Characterization of *jam2b^{Gt(2A-Gal4)}*;UAS:GFP* expression at 1-5 dpf. (A-F) *jam2b^{Gt(2A-Gal4)}*;UAS:GFP* shows strong GFP expression within the lateral plate mesoderm (LPM) along the yolk extension at 26 hpf (white arrows, A,B), which becomes presumptive mesothelium at 3-5 dpf (white arrows, C-F). GFP expression is also apparent in a subset of skeletal muscle cells (white arrowheads, A,C,E), craniofacial muscle (yellow arrows, E,F) and in the cells surrounding the eye (yellow arrowheads, E,F). (G-I) Co-expression of *jam2b^{Gt(2A-Gal4)}*;UAS:GFP* and *etv2* (red) in the lateral plate mesoderm of the trunk region at 30 hpf. Note that GFP and *etv2* expression overlaps in the SVF cells (arrowheads), located in the most dorsal region of GFP expression domain. (J-R) Expression of *jam2b^{Gt(2A-Gal4)}*;UAS:GFP* in the LPM along the yolk extension and in muscle cells from 30 to 66 hpf. A few GFP-positive cells are present in the PCV and overlap with vascular endothelial marker *kdrl:mCherry* expression (white arrowheads and insets). (S-U) Analysis of *jam2b^{Gt(2A-Gal4)}*;UAS:GFP* expression in *lyve1b:dsRed* reporter background at 4 dpf, which labels PCV and the thoracic duct. Note GFP-positive cells in the TD (arrowheads). Bright GFP expression in the skeletal muscle fibers is also apparent. PCV, posterior cardinal vein; DA, dorsal aorta; TD, thoracic duct. Lateral views of the trunk region, anterior is to the left. Scale bars: 100 μm.

Figure 2. *Jam2b*-positive cells are derived from the lateral plate mesoderm and contribute to the vasculature. (A-F) Time-lapse images of *jam2b^Gt(2A-Gal4)^;UAS:GFP* ; *Tg(kdrl:mCherry)* embryo starting at 24 hpf show two GFP-positive cells (arrowheads), which migrate dorsally, proliferate, integrate into the PCV and initiate *kdrl:mCherry* expression (see Movie 1). Time is in minutes after 24 hpf. PCV, posterior cardinal vein; DA, dorsal aorta. Scale bar: 50 μm.

(G-I) HCR against *etv2* in a *jam2b^Gt(2A-Gal4)^;UAS:GFP* embryo at the 18-somite stage showing *jam2b* expression (green) lateral and ventral to etv2 positive vascular progenitors (red).

(J-L) HCR against *prrx1a* in a *jam2b^Gt(2A-Gal4)^;UAS:GFP* embryo at the 18-somite stage showing colocalization of *prrx1a* mRNA and GFP fluorescence. Dorsolateral view of the trunk region is shown, anterior is to the left.

Figure 3. Genetic lineage tracing analysis of *jam2b*+ cell contribution to vasculature.
(A) Schematic diagram of recombination in *jam2b^Gt(2A-Gal4)^; UAS:Cre; actb2-loxP-BFP-loxP-dsRed* embryos. Ubiquitous BFP expression switches to dsRed in Cre-positive cells. (B-J) Lineage tracing analysis in embryos at 3 dpf obtained by crossing *jam2b^Gt(2A-Gal4)^* and *UAS:Cre; actb2:loxP-BFP-loxP-dsRed; fli1:GFP* parents. Red cells, indicating the genetic switch, are observed in the mesothelial layer surrounding the yolk (arrow, B,E), skeletal muscle (yellow arrowheads, B), and the epicardial (arrows, C,D) and myocardial (arrowheads, C,D) heart layers. Switched cells in the vasculature are observed in the venous ISVs (white arrowheads, B,E,F,G), the PCV (blue arrowheads, F,G) and SIA (arrow, I,J). A control embryo negative for *jam2b^Gt(2A-Gal4)^* is shown in (H) to indicate autofluorescence. (F,G) show larger magnification views of (B,E). (K) Quantification of endothelial cell labeling in different vascular beds in *jam2b^Gt(2A-Gal4)^; UAS:Cre; actb2-loxP-BFP-loxP-dsRed* embryos at 3 dpf. (L) Quantification of endothelial cell labeling in different vascular beds in *jam2b^Gt(2A-Gal4)^; UAS:CreERT2; actb2-loxP-BFP-loxP-dsRed* embryos at 3 dpf. 4-OHT was added starting at 7 hpf or 22 hpf stages. In a third group of embryos, 4-OHT was added at 7 hpf and washed out at 22 hpf stages. Note that cell contribution to SIV could not be analyzed reliably due to very strong red fluorescence in the mesothelial layer surrounding the yolk.

Figure 4. Intestinal vasculature in the middle and posterior trunk region forms largely by new vasculogenesis.

(A,B). *etv2:Kaede* embryos were photoconverted from green to red at 24 hpf and analyzed at 4 dpf. Merged and red only channels are shown. (C-E) Higher magnification images of the boxed area in (A,B). Merged, red and green channels are shown. Note that SIA and SIV are comprised mostly from green only cells (white arrowheads), indicating that they have originated after the photoconversion. (F) Quantification of green only (new) and green+red (pre-existing) cells in the SIA and SIV at 4 dpf. The analysis focused on the middle and posterior trunk region over the yolk extension area.

Figure 5. scRNA-seq analysis of cells from *jam2b^Gt(2A-Gal4)*; *UAS:mCherry-NTR*; *etv2^Gt(2A-Venus)* embryos at 36 hpf.
(A) UMAP plot showing 37 cell clusters identified by Seurat analysis. LPM, lateral plate mesoderm; SVF, secondary vascular field; RBC, red blood cells. (B) Dot plot of selected marker gene expression level in each cell cluster. (C) 2D UMAP plot showing the distribution of Venus and mCherry positive cells. Note that Venus-positive cells mostly correspond to different vascular endothelial lineages, while mCherry-positive cells label different subsets of LPM. (D) Feature plots showing the expression of SVF cell markers, enriched in the cluster 15. (E) Violin plots show SVF cell marker expression in different clusters. (F) Pseudotime trajectory of vascular endothelial and SVF clusters made using Monocle 3. Note that SVF cells (cluster 15) can transition into both venous and arterial cell types.

Figure 6. SVF cell number is reduced in *jam2a; jam2b* mutant embryos.
(A-C) In situ hybridization analysis (ISH) for *etv2* expression in SVF cells (arrows, A,B) in maternal-zygotic (MZ) *jam2a-/-; jam2b-/-* embryos and control *jam2a+/-; jam2b+/-* embryos in *kdrl:GFP* background, obtained by crossing double homozygous *jam2a-/-; jam2b-/-; kdrl:GFP* adults to wild-type *kdrl:GFP*. (D-I) ISH analysis for *tal1* and *npas4l* expression in SVF cells (arrows) in MZ jam2a-/-; jam2b-/- embryos and control embryos in *fli1:GFP* background. Controls were obtained by mating *jam2a+/-; jam2b-/-* embryos to wild-type *fli1:GFP*, resulting in a mixture of *jam2a+/+; jam2b+/-* and *jam2a+/-; jam2b+/-* genotypes. ***p<0.001, ****p<0.0001, two-tailed t-test, error bars show ±s.d. Data are combined from four (C) or two (F,I) independent replicate experiments. Het labels in (C,F,I) refer to double heterozygous in (C) and a mixture of *jam2a+/+; jam2b+/-* and *jam2a+/-; jam2b+/-* genotypes in (F,I), while Mut label refers to *jam2a-/-; jam2b-/-* MZ mutant embryos.

Figure 7. Analysis of vascular defects in MZ *jam2a; jam2b* mutant embryos.
(A-C) Intersegmental vessel extension analysis in MZ *jam2a-/-;jam2b-/-* and *jam2a+/-;jam2b+/-* embryos in *kdrl:GFP* background at 3 dpf. Note that some ISVs are truncated in *jam2a;jam2b* mutant embryos (arrow, B, an inset shows higher magnification view of the boxed trunk region). (D-F) Endothelial cell number in the PCV is reduced in MZ *jam2a; jam2b* mutant embryos compared to *jam2a+/-; jam2b+/-* controls at 3 dpf. Endothelial cells were counted in the floor of the PCV (arrowheads) in the selected area of the trunk region based on *fli1:GFP* expression. (G-I) SIA and SIV are fragmented and show increased incidence of gaps in MZ *jam2a;jam2b* mutant embryos compared to heterozygous controls at 3 dpf. Analysis was performed in the background of *etv2^{Gt(2A-Gal4)}; UAS:mCherry-NTR* transgene, which shows expression in all vasculature. Note the underdeveloped SIA (arrowheads) and gaps in SIV (arrow) in MZ *jam2a;jam2b* mutants. Higher magnification views are shown in the insets. Diagrams in (I) show percentage of embryos with gaps in SIA (left) and SIV (right). (J-L) Thoracic duct analysis in *prox1:RFP; fli1:GFP* transgenic embryos at 6 dpf. Note the gaps in the thoracic duct (arrows) observed in MZ *jam2a;jam2b* mutant embryos. (L) Shows the percentage of embryos with normal or fragmented thoracic duct. (M-O) Vascular recovery analysis after endothelial cell ablation. MZ *jam2a;jam2b* mutant and control heterozygous embryos in *etv2^{Gt(2A-Gal4)}; UAS:NTR-mCherry* background were treated with MTZ starting at 6 hpf to ablate ECs and subsequently washed out at 45 hpf. A partial vascular recovery is observed in the control embryos, while it is greatly reduced in MZ *jam2a;jam2b* mutants. (O) shows length measurements of the recovered vasculature. (C,F,O) graphs show two tailed t-test, while (I,L) show Fisher's exact test. *p<0.05, **p<0.01, ****p<0.0001. DA, dorsal aorta; PCV, posterior cardinal vein; SIA, supraintestinal artery; SIV, subintestinal vein.

Figure 8. *hand2* is required for *jam2b* expression in the lateral plate mesoderm and SVF cells.
(A-F) HCR analysis for *hand2* (red) and *jam2b* (magenta) expression in *kdrl:GFP* embryos at 24 hpf. Anterior (A-C) and posterior (D-F) regions of the trunk are shown. Note the overlapping expression in the LPM along the yolk and yolk extension (arrows), which does not overlap with *kdrl:GFP* expression. (G-J) Whole-mount in situ hybridization analysis for *jam2b* expression in *hand2* mutant and wt control embryos at 26 hpf. Lateral (G,H) and dorsal (I,J) views. Note that *jam2b* expression in *hand2* mutants is absent from the posterior domain of the LPM along the yolk extension (black arrowheads), while the anterior portion of the LPM (white arrowheads) is only mildly affected. (K,L) Whole-mount in situ hybridization for *etv2* expression in *hand2* mutant and control wild-type embryos at 30 hpf. Note that *etv2* expression in SVF cells (arrows) is absent in *hand2* mutants. In addition, its expression in the PCV (arrowheads) is greatly reduced or absent. ISV, intersegmental vessel; DA, dorsal aorta; PCV, posterior cardinal vein. Scale bars: 100 µm.

Figure 9. A diagram illustrating the emergence of SVF cells from *jam2b*-positive lateral plate mesoderm. SVF cells contribute to the intestinal vasculature, including the supraintestinal artery (SIA), subintestinal vein (SIV), as well as the posterior cardinal vein (PCV).

Supplementary Figure S1. In situ hybridization analysis of *jam2b* expression between the 20-somite and 48 hpf stages. Note strong *jam2b* expression in the lateral plate mesoderm along the yolk and yolk extension (arrows). (A,B,C,E) lateral view, (D) ventral view.

A

CCA**AGCTCAGATCCTGTTACGGT**G*GGAAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAGGCTGGAGACGTGGAGGAGAACCCTGGACCT*GTCGACACC**ATGAAGCTACTG.....**

B

Supplementary Figure S2. Generation of *jam2b^Gt(2A-Gal4)^;UAS:GFP* line using CRISPR-Cas9 mediated non-homologous recombination.
(A) A diagram of *sgRNA-2A-Gal4* construct and its insertion into the *jam2b* genomic locus. The targeting construct contained *jam2b* sgRNA site, followed by in-frame fusion to the viral peptide P2A, Gal4 DNA-binding domain and the SV40 polyA sequence. *jam2b* sgRNA targets the second exon of *jam2b* genomic sequence. (B) Comparison of GFP fluorescence in *jam2b^Gt(2A-Gal4)^;UAS:GFP* embryo and *jam2b* mRNA expression analyzed by in situ hybridization at 24 hpf. Note the bilateral *jam2b* expression and GFP fluorescence observed in the lateral mesoderm along the yolk and yolk extension. Scale bars: 100 μm.

Supplementary Figure S3. Quantification of *jam2b^Gt(2A-Gal4)*;*UAS:GFP* cell contribution to different vascular beds. All analysis at 3 dpf, except for TD, which is at 4 dpf. n=7 embryos except for TD, for which n=17. PCV, posterior cardinal vein, DA, dorsal aorta, ISV, intersegmental vessel, PL, parachordal lymphangioblast, SIA, supraintestinal artery, TD, thoracic duct.

Supplementary Figure S4. Generation of *jam2b* and *jam2a* mutants using CRISPR-Cas9 genome editing. (A) A schematic diagram illustrating generation of *jam2b* mutants using two sgRNAs designed to delete the promoter region. (B) A schematic diagram illustrating generation of *jam2a* mutants using two sgRNAs designed to delete the entire *jam2a* coding sequence. (C,D) In situ hybridization (ISH) analysis for *jam2b* expression in wild-type *jam2b+/+* and *jam2b-/-* mutant embryos at 30 hpf. Note that *jam2b* expression along the lateral plate mesoderm is absent in *jam2b* mutants (arrows). Some residual staining is due to background and some pigment cells along the horizontal myoseptum. (E,F) ISH analysis for *jam2a* expression in wild-type *jam2a+/+* and *jam2a-/-* mutants at 26 hpf. The strongest *jam2a* expression is observed in the posterior somites, which is absent in *jam2a* mutants (arrows).

Supplementary Figure S5. *jam2b* **mutant embryos do not show any apparent defects in SVF cell formation or vascular recovery after endothelial cell ablation.** (A-C) Quantification of SVF cells (white arrowheads) by in situ hybridization analysis for *etv2* expression at 30 hpf. Experimental and control embryos were obtained from the incross of *jam2b-/-* or wild-type (*jam2b+/+*) sibling parents in *kdrl:GFP* background, respectively. (D,E) Quantification of vascular recovery after endothelial cell ablation. *jam2b+/-* and *jam2b-/-* adults in *etv2*^*Gt(2A-Gal4)*^*; UAS:mCherry-NTR; Tg(kdrl:GFP)* background were crossed with each other. Embryos were treated with 10mM MTZ from 10 hpf to 45 hpf, which was then washed and embryos were allowed to undergo vascular recovery until 72 hpf, when they were imaged and genotyped. Control embryos were treated with 0.1% DMSO. The length of new vascular cords (white lines) at the position of the PCV and SIV was measured and compared in *jam2b+/-* and *jam2b-/-* mutant embryos. Note that the bright *kdrl:GFP* expression along the yolk extension corresponds to the pronephros and was not affected in these experiments. Lateral view is shown, anterior is to the left. Mean±sd is shown; ns, not significant. Scale bars: 100 μm.

MOVIE LEGENDS

Movie 1. Migration of *jam2b*-expressing cells from the lateral plate mesoderm into the PCV: A time-lapse video of a *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP;Tg(kdrl:mCherry)* embryo shows GFP positive cells migrating dorsally, integrating into the PCV, proliferating and initiating *kdrl:mCherry* expression (white arrowheads) (see Fig. 2A-F). Imaging was started at 24 hpf. PCV, posterior cardinal vein; DA, dorsal aorta. Scale bar: 20 μm.

Movie 2. *jam2b*- expressing cells are derived from the lateral plate mesoderm: A time-lapse video of a *jam2b$^{Gt(2A-Gal4)}$;UAS:GFP;Tg(kdrl:mCherry)* embryo showing from the 18-somite stage to 24 hpf. GFP positive cells can be seen in the ventrolateral LPM region at the 18-somite stage and remain in the same region until 24 hpf as the embryo undergoes convergent extension. PCV, posterior cardinal vein; DA, dorsal aorta. Scale bar: 50 μm.