DupyliCate: mining, classifying, and characterizing gene duplications

Shakunthala Natarajan[1] & Boas Pucker[1*]

[1] Plant Biotechnology and Bioinformatics, Institute for Cellular and Molecular Botany, University of Bonn, Kirschallee 1, 53115, Bonn, Germany

* Corresponding author: pucker@uni-bonn.de

Abstract

Paralogs, copies of a gene, form an important basis for novelty during evolution. Analysis of such gene duplications is important to understand the emergence of novel traits during evolution. DupyliCate is a Python tool that has been developed for this purpose. With the ability to process multiple datasets concurrently, flexible features, and parameters to set species-specific thresholds, DupyliCate offers a high-throughput method for gene copy identification and analysis. The different available parameters and modes are explored in detail based on the *Arabidopsis thaliana* datasets. Proof of concept for the tool is presented by characterizing well known duplications in different plants, and its broad applicability is demonstrated by running it on diverse datasets including complex plant genome sequences with high heterozygosity. Further, two case studies involving the evolution of *flavonol synthase* (*FLS*) genes in Brassicales, and the evolution of flavonol synthesis regulating myeloblastosis (MYB) transcription factors- *MYB12* and *MYB111* across a large number of plant species, are presented as exemplar use cases. The tool's applicability beyond plants is demonstrated on *Escherichia coli, Saccharomyces cerevisiae, and Caenorhabditis elegans* datasets. DupyliCate is available at: https://github.com/ShakNat/DupyliCate.

Key words: Gene duplications, comparative genomics, expression analysis, high-throughput, Ka/Ks, evolution

Introduction

Gene duplication products also called paralogs are homologous genes originating from a common ancestor gene. In contrast to orthologs, they arise through duplication events and not speciation events[1]. They are an important source of evolutionary novelty in all organisms and are major drivers for the diversification of biosynthetic pathways seen in plants and other organisms. Gene duplications can broadly be divided into small-scale duplications and large-scale duplications. The small-scale duplication group encompasses tandem duplications, proximal duplications, transposon-derived duplications, and retroduplications. Whereas, the large-scale duplication group includes whole genome duplications, and segmental duplications[2]. Identification of gene

38   duplications is fraught with difficulties like functional redundancy of gene duplicates, and extensive
39   small-scale duplication events. It is also compounded by gene loss, rearrangement, mutation
40   events across evolutionary time scales and presence of multidomain protein structures[2,3]. Lastly,
41   gene duplicates can be formed by a number of mechanisms, and have different ages and
42   evolutionary fates, further increasing the difficulty of detection and analysis[1].
43
Nevertheless, there have been huge research efforts in this direction resulting in a plethora of
tools and methodologies to identify gene duplications. GenomeHistory[4], was among the first tools
to provide a basic framework for gene duplication detection with a combination of local and global
alignment approaches followed by pairwise computation of Ka/Ks values to evaluate selection
pressures. But this software's web page no longer exists and is hence not accessible. Following
this, several tools like OrthoMCL[5], OrthoFinder[6], MCScanX[7] were developed to identify orthologs
and as a result also distinguish them from paralogs using tree-based approaches and synteny
approaches, respectively. All these tools focus more on the ortholog identification aspects. Next,
DuplicationDetector[8] was a Perl script developed to identify putative recent segmental duplicates
in homozygous organisms from next-generation sequencing data. It focuses on identifying large
scale duplications like segmental duplications, at the DNA level. Later, with a detailed study on
gene duplications and evolution in plants, DupGen_finder[9], a Perl based script was developed to
identify gene duplicate pairs, and comprehensively classify gene duplicates into tandem
duplicates, proximal duplicates, transposed duplicates and whole genome duplicates. The study
also provided separate scripts for Ka/Ks analysis by integrating KaKs_Calculator 2.0[10], a
dedicated tool for Ka/Ks calculations. Ka/Ks is defined as the ratio of nonsynonymous mutation
rate to that of synonymous mutation rate. Ka/Ks ratios are useful for analysing selection pressures
and sequence divergence between genes. Thus, the Ka/Ks computation was integrated with the
tool, to study the sequence divergence of the identified gene duplicates [9,10]. The work further
involved expression divergence analysis of the identified gene duplicates, but this analysis was
not a part of the DupGen_finder main script, and is not expandable to wide-ranging user data.
Zhang and co-workers later developed HSDFinder[11], a BLAST-based web tool to annotate,
categorize and visualize highly similar duplicates across species. More recently, an updated
version of this tool, called HSDSnake[12] was released as a user-friendly SnakeMake pipeline, with
added features to classify duplicate pairs by integrating the DupGen_finder script and compute
Ka/Ks ratios. WGDI[13] was another Python-based command-line tool that was developed to
identify and characterize whole genome duplication related events in species, thus focusing on
large scale duplications. Another Python-based tool integrating machine learning framework to
detect orthologs, annotate them, identify gene losses and duplications across alignable genomes
is TOGA (Tool to infer Orthologs from Genome Alignments)[14]. TOGA leverages alignments
between intergenic regions and introns to identify orthologs, and perform other auxiliary analyses.
Following this, doubletrouble[15] was released as an R package to classify and identify gene
duplications. It works on a framework similar to DupGen_finder, identifies duplicate pairs,
classifies them, calculates Ka/Ks ratios, but also goes one step ahead in helping identify
retrotransposon derived duplicates and DNA transposon derived duplicates. Finally, SegmantX[16]
is another recent Python script that uses a novel approach of local alignment followed by a
chaining algorithm to identify segmental duplications at the DNA level and also provides a docker
container for ease of use.

82
From the varying tools discussed above, the following aspects are evident - first, most of the gene
duplication identification tools identify duplicates in pairs. But in a biological context there can be
arrays of duplicates, and identifying them all and maintaining the link between them is crucial for
further evolutionary understanding. Next, most tools have been adapted for use on standardized
data from databases like Ensembl and NCBI, limiting the use cases on widely varying
bioinformatics data formats. Given the heterogeneity of General Feature Format (GFF) files from
different sources, and databases, it is necessary to come up with tools that can work across
different GFF file flavours. Furthermore, many tools focus on either identifying large scale
duplications and characterizing relevant evolutionary events or classifying duplications. But it is
also important to integrate expression analysis, to understand the functional divergence of
identified duplicates and also study the nature of the identified duplicates by adding details on
genomic order conservation with synteny, in cases of comparative genomic studies between two
species. Finally, from a biological perspective, different species have different duplication
landscapes in their genomes. However, existing methods work based on defined thresholds
necessitating the use of species-specific thresholds[2].
98
In this context, DupyliCate is presented here as a solution for gene duplication array identification,
and broad classification of the duplicates. It provides options for downstream expression analysis
of the identified duplicates to understand the expression divergence, and gene-level Ka/Ks
computations implemented independently as a part of the script, to infer selection pressures. It
has been deployed here on datasets from NCBI, Phytozome, and varying additional sources, to
demonstrate its ability to process widely varying input file formats. It also leverages BUSCO-
based metrics to determine species-specific thresholds for gene duplicates identification, and
offers the users a wide range of parameters that can be fine-tuned according to the objectives at
hand.

Results

Benchmarking results

DupyliCate was compared against the tools doubletrouble and DupGen_finder. The
benchmarking results of DupyliCate runs using manual threshold and BUSCO-based threshold
for singleton-duplicate segregation were different from one another as well from the other tools
compared (Table 1). The run with manual thresholds (a) gave numbers comparable to the results
output by both doubletrouble and DupGen_finder. The number of gene duplicates classified by
DupyliCate (a) as tandems and proximals is comparable to that output by doubletrouble, while
DupGen_finder gives a lower number of tandem and proximal duplicates. When comparing the
total number of duplicates, DupyliCate and DupGen_finder have close estimates while
doubletrouble gives a lower number of total duplicates. Singleton genes show the maximum
difference across the three tools. While DupyliCate and DupGen_finder output ~4000 and ~20
singletons respectively, doubletrouble does not have a singleton output. Further, when looking at
the classifications assigned to all the input genes, both doubletrouble and DupGen_finder have
remnant unclassified or unaccounted genes while DupyliCate classifies all the input genes.

The BUSCO-based threshold run used a self-normalized bit score cutoff of 0.696 based on the
BUSCO-analysis and results in approximately two-thirds of the genes being classified as
singletons.
126
127   **Table 1:** Consolidated results of benchmarking analysis on the *A. thaliana* Col-0 dataset. 'Misc.'
128   gene duplicates in the table indicate all gene duplicates that are not tandem and proximal
129   duplicates.
130

Tool	Tandem	Proximal	*Misc.	Singleton	Non-recurrant tandem and proximal duplicates	Non-recurrant total gene duplicates	Non-recurrant total duplicates and singletons	Total no. of coding genes input for classification	Total no. of unclassified genes
DupyliCate (a)	3584	1530	20092	3829	4844	23615	27444	27444	0
DupyliCate (b) (BUSCO-based threshold)	1756	705	5387	19909	2374	7535	27444	27444	0
doubletrouble	3045	3581	22224	-	4970	22869	22869	27444	4575
DupGen_finder	1316	2978	20921	19	4294	23340	23359	27444	4085

131
132   When compared to the other tools, DupyliCate additionally produces a duplication landscape plot
133   from the distribution of the self-normalized bit scores in the genome of a sample analyzed for
134   gene duplications (described in workflow sub-section 'Step 5: Threshold determination for
135   singleton-duplicate segregation and self-alignment'). The plot of *A. thaliana* Columbia-0 (Col-0) is
136   shown in Fig. 1. More details on the plot interpretation are presented in the discussion section.

Figure 1: Duplication landscape plot of *A. thaliana* Col-0 output by DupyliCate - shows the distribution of self-normalized bit scores from the sequence alignments. The plot can help get an idea of the genome-wide duplications in the sample being analyzed and can also help decide optimal normalized bit score thresholds in case a user would like to set them manually without using the auto option. Left and right skewed distributions in the plot could suggest genomes with extremely low gene duplications and genomes with extensive gene duplications or polyploidization, respectively. KDE stands for the Kernel Density Estimate. It is a smoothed, continuous probability density curve estimated from the discrete histogram data. The KDE reveals the underlying distribution shape independent of the histogram bin width, enabling clearer detection of skewness and multiple peaks.

167 In terms of runtime performance, doubletrouble and DupyliCate without BUSCO had comparable
168 runtimes while DupGen_finder and DupyliCate with BUSCO had comparable runtimes (Fig. 2).

169
170 **Figure 2:** Runtime comparison across DupyliCate, doubletrouble, and DupGen_finder

Runtime analyses

172 The different parameters tunable in DupyliCate were explored in different runs of the tool using
173 *A. thaliana* Col-0 dataset on an Ubuntu 24.04 LTS machine with 30 cores. For runs involving
174 comparative analysis with a reference, *A. thaliana* Niederzenz-1 (Nd-1) was used as the sample
175 and *A. thaliana* Col-0 was used as the reference. All the runs were performed using the default
176 BUSCO-based thresholding method. The runtimes for the different runs are shown in Table 2 .
177
178 **Table 2:** Runtime compilation when using different modes and parameters of DupyliCate on *A.*
179 *thaliana* datasets
180

Key parameter changed	Runtime (mins)

Overlap mode	6.234
Strict mode	6.422
Phylogeny-based ortholog assignment in the presence of a reference	8.918
Local alignment-based ortholog assignment in the presence of a reference	8.0346
BLAST aligner	14.368
MMSeqs2 aligner	6.433
Expression analysis of all gene duplicates	71.82
Expression analysis of specific genes	28.317
Ka/Ks computation of all gene duplicates	25.5

The specific expression analysis of the *Rapid Alkalinization Factor* (*RALF*), *Xeroderma*
pigmentosum (*XPB*), and *β-L arabinofuranosidases* (*BLA*) duplicates resulted in the gene
expression plots shown in Fig. 3. The *RALF* genes expression plot (Fig. 3(a)) captures the
pairwise expression of some genes in the *RALF* array[17]. The expression plot pattern for the *XPB*
duplicates potentially point towards a subfunctionalization fate shown by the strong correlation in
the gene expression of the gene duplicates with both the copies being functional (Fig. 3(b)).
Subfunctionalization is usually denoted by expression in different parts of the plant or under
different stress conditions. One such example of subfunctionalization is the spatially different gene
expression patterns of the sub-group 7 myeloblastosis (*MYB*) transcription factor (TF) gene
copies - *MYB11*, *MYB12* and *MYB111*, in *Arabidopsis thaliana* seedlings. Each of these TFs,
MYB11, *MYB12* and *MYB111,* are expressed in specific domains of the seedling, roots and
cotyledons, respectively[18]. But since the spatial context of the gene expression data is not known
here, based on the correlation analysis in DupyliCate (Supplementary Table S4) and experimental
evidence from Masuda and co-workers[19], the subfunctionalization fate of *XPB* duplicates is
explained here. Whereas, the expression plot of the *BLA* duplicates indicates a potential
neofunctionalization fate, where only one of the duplicate copies is highly expressed while the
other shows very low expression (Fig. 3(c))[20]. Neofunctionalization is defined as the fate where
one of the copies diverges and develops a new function. While the low expression of one of the
gene copies can be attributed to pseudogenization as well, it is important again to note that overall
expression is considered here and spatial gene expression information is not included here. But
the experimental evidence from the work of Tao and co-workers shows that the copy *AT5G12960*
is pollen-specific while the other copy *AT5G12950* is ubiquitously expressed in all other tissues[20].
This supports the hypothesis that *AT5G12950* was the original copy while *AT5G12960* diverged
after duplication and acquired new tissue-specific functions not found in the original copy leading
to their fate being called neofunctionalization. The correlation analysis of these duplicates
computed with DupyliCate also indicate expression divergence (Supplementary Table S4) and
the low overall expression of *AT5G12960* supports its pollen-specific function. Further, the gene
expression plots can be obtained in the compact mode as shown in Fig. 3, as well as matrix mode
wherein the genes in a duplicate array are arranged in the form of a matrix and the respective

expression plots are displayed at the intersection of the correct gene pair in the matrix
representation (Supplementary Figure 11). Moving on, the runtime trajectory analyses were
performed using the Brassicales plant datasets used for the *FLS* case study. The runtime
trajectory for duplicates mining (reference-free) (Fig. 4(a)), and duplicates mining combined with
orthology assignments for comparative gene copy number analysis (with phylogeny) showed a
linear time increase trend (Fig. 4(b)). The runtime for the large-scale study involving plant datasets
from Phytozome with *A. thaliana* as the reference took 34.8 hours to complete. This demonstrates
the suitability of DupyliCate for high-throughput use cases involving gene duplicates mining as
well as comparative genomic studies needing orthology assignments with respect to a reference.
220
221
222

224 **Figure 3:** Pairwise gene expression plots of *A. thaliana* Col-0 tandem genes encoding (a) Rapid
225 Alkalinization Factors (b) Xeroderma pigmentosum, and (c) β-L arabinofuranosidases.
226 'n' represents the number of samples used to create the gene expression plot. The x and y axes
227 show the gene expression in log(1+TPM). TPM stands for Transcripts Per Million.

228
229 **Figure 4:** Runtime trajectory analysis for (a) Gene duplicates mining across twelve Brassicales
230 genome sequences (b) Gene duplicates mining, and comparative genomics involving ortholog
231 assignments across twelve Brassicales genome sequences with respect to the reference *A.*
232 *thaliana.*

Proof of Concept analyses

Proof of Concept (POC) analyses of DupyliCate were performed using well-known gene duplications reported in different plants. DupyliCate correctly identified tandem duplication of the *SEC10* gene in *A. thaliana* Nd-1 with respect to *A. thaliana* Col-0[21,22]. Next, tandem duplication of a gene in *Gardenia jasminoides* with respect to *Coffea canephora*[23] was correctly mined as a tandem duplicate array of four genes. A proximal duplication of genes was detected in the nematode tolerance locus of the nematode tolerant sugarbeet line StrubeU2Bv whereas the susceptible line KWS2320 had just a single copy of the gene[24]. DupyliCate confirmed the dispersed duplication of the sterol Δ^{24}-isomerase (24ISO) gene in *W. somnifera* responsible for withanolides biosynthesis[25]. These genes were also found to show up as copies along with the closely related sterol side chain reductase (SSR) genes in a proximal-dispersed gene duplicate group in the duplicate relationships file.

GeMoMa activating flag in the tool produced annotations for *V. amurensis* and *V. rotundifolia* with *V. vinifera* as the reference, and helped identify the gene duplicates for species with missing annotation files. *V. amurensis* had the highest number of gene duplicates (Table 3).
Among the analyzed rice and related genomes, *Echinochloa colona* and *Echinochloa oryzicola*, *which* are rice weeds, had the highest number of gene duplicates. All the other rice varieties and their relatives had a comparative number of duplicates (Table 3). The copy number table (Supplementary Table S5) gave a visual presence-absence matrix of the orthologs of the different stress genes in *O. sativa* indica across the analyzed rice relatives.

DupyliCate also identified gene duplicates and singletons in the non-plant species - *E. coli, S. cerevisiae, and C. elegans*. As expected, *E. coli* and *S. cerevisiae* had a low number of gene duplicates whereas *C. elegans* showed a relatively high number of gene duplicates (Table 3).

Table 3: Consolidated results of gene duplicates identified and classified in *Vitis* species, *Oryza* species, rice relatives, and non-plant samples

Samples	Tandem duplicates	Proximal duplicates	Dispersed duplicates	Singletons	Low confidence duplicate groups	Moderate confidence duplicate groups	High confidence duplicate groups
V. vinifera	4856	2095	8751	10832	98	26	5867
V. rotundifolia	5281	2958	10829	10898	58	23	6849
V. amurensis	5422	3741	17364	10463	83	24	9632
L. perrieri	1336	1006	5883	21241	47	7	3587
O. barthii	1518	1470	6623	25389	43	9	4254
O. glaberrima	1631	922	9641	21406	47	11	5065
O. nivara	1282	1269	7253	26941	47	9	4326

O. rufipogon	1252	1342	7018	27885	56	9	4237
O. sativa japonica cv.	1788	1686	7300	25582	42	9	4676
Z. latifolia	1033	724	17702	24747	169	8	8317
E. oryzicola	5428	2933	53451	9743	921	275	24672
E. colona	5922	2798	77576	10854	1347	362	28510
E. coli	29	22	570	3673	0	0	267
S. cerevisiae	101	42	971	4936	12	1	468
C. elegans	2277	978	3609	13727	5	41	2806

FLS evolution in Brassicales

Among the Brassicales plants analyzed in the *FLS* (flavonol synthase) case study*, I. tinctoria, E.*
vesicaria, S. alba and *B. napus* showed a high number of gene duplications. Tables consolidating
the gene duplicates mined and the orthologs and copy numbers of the *Arabidopsis thaliana FLS*
(*AtFLS*) genes in the other Brassicales plants are included in Supplementary Table S3 and
Supplementary Table S6, respectively. The presence-absence variation in the copy number table
showed the presence of *FLS1* - the functional *FLS* in all the Brassicales plants analyzed. But all
of the Brassicales plants were missing at least one or more of the other *FLS* genes. The
phylogenetic tree used to validate the ortholog assignments is shown in Fig. 5. Most of the
ortholog assignments in DupyliCate and the clade grouping in the phylogenetic tree match, except
for a few sequences (coloured with red strips in the tree (Fig. 5). The *C. papaya* gene identified
as *FLS1* (evm.TU.supercontig_37.77) does not exactly group in the *AtFLS1* clade but is a sister
to this clade. No other *FLS* genes were identified in *C. papaya.* The *B. oleracea* gene that groups
in the *AtFLS2* clade (Bol019362.v1.0) shows up as an *FLS1* ortholog in DupyliCate. On a similar
note, a group of genes marked by the red strips in the *FLS4* clade in the tree are assigned as
orthologs of *FLS1* or *FLS3* in DupyliCate results. A consistent occurrence of this pattern among
these genes that form a separate subclade with respect to *AtFLS4* is interesting. The phylogenetic
tree also shows a high number of sequences in the *FLS3* and *FLS4* clades, indicating a potential
expansion of these genes in Brassicales.
281

Figure 5: Phylogenetic analysis of *FLS* genes across twelve Brassicales plant genomes. The red striped sequences have different ortholog assignments in DupyliCate and the phylogenetic tree. The collapsed clade contains flavanone 3-hydroxylase (F3H) sequences as outgroups

MYB evolution across a broad phylogenetic range of plant species

A large-scale analysis of plant datasets from Phytozome facilitated the understanding of *MYB* (Myeloblastosis) transcription factors - *AtMYB12,* and *AtMYB111* evolution across a wide range of plant species. A summary of the number of gene duplicates mined across the 153 plant species along with the copy number table representing the *A. thaliana MYB12* and *MYB111* orthologs across the analyzed datasets were obtained (Supplementary Table S7). Among the analyzed plants, orthologs for *MYB12* and *MYB111* were not found in algae, mosses, liverworts, and ferns. Interestingly, the *MYB12,* and *MYB111* orthologs were found in lycophytes (club mosses). In the other plants, orthologs were found for at least one or both of *AtMYB12* and *AtMYB111*. The phylogenetic tree used to validate the ortholog assignments is shown in Fig. 6. The phylogenetic analysis also reveals an interesting evolutionary pattern of *MYB12* and *MYB111*. As seen in Fig. 6, *AtMYB12* and *AtMYB111* fall in separate clades that consist of sequences from *A. thaliana* and other dicot plants (pink and purple clades in Fig. 6). Monocots, a few other dicots, and some basal angiosperms on the other hand do not show separate lineages of these flavonol biosynthesis-regulating MYBs.

302
303
304 **Figure 6:** Phylogenetic analysis of *MYB12* and *MYB111* genes across a broad phylogenetic
305 range of plants. All the sequences not covered by the yellow strips belong to dicots. The
306 collapsed clade contains *MYB75* sequences as outgroups.
307
308 Further, the tree revealed that there are *MYB111* and *MYB12* orthologs captured correctly, as
309 well as false positive sequences (uncoloured sequences bounded by the outgroup MYB75 clade
310 and the *A. thaliana* flavonol MYB clade coloured in blue). The ortholog assignments in DupyliCate
311 and the clade grouping in the tree were compared (Fig. 7).

312
313 **Figure 7:** Pie chart representing the validation of DupyliCate's ortholog assignments with
314 respect to clade grouping in the phylogenetic analysis
315

316 The validation analysis shows that nearly 50% of the ortholog assignments in DupyliCate matched
317 exactly with the phylogenetic tree. More details on the validation analysis can be found in
318 Supplementary Methods.
319

Parameter tuning results

321 The parameter tuning analyses revealed that --score, --self_simcut, and --seq_aligner are the top
322 sensitive parameter flags that influenced the final results the most (Supplementary Table S10). A
323 parameter sensitivity ranking table was created based on these tuning trials and a total of five
324 parameter flags were found to be the most sensitive influencers of the final results
325 (Supplementary Table S11).
326

The BUSCO vs manual threshold comparative analyses were performed and the results were
consolidated in Supplementary Table S12. While the total number of gene duplicates were higher
in the manual threshold method for *A. thaliana*, *D. purpurea,* and *D. dumetorum*, for *V. cruziana*,
the BUSCO threshold-based method gave more gene duplicates. But the overall proportion of
gene duplicates and singletons in the BUSCO-based threshold approach corresponded well with
the respective duplication landscape plots of the different plant species (Supplementary Results
1).
334
335

Discussion

The benchmarking analyses reveal that DupyliCate performs on par or better than doubletrouble[15], and DupGen_finder[9] when using a manual threshold specification. This can be seen from the remnant number of unclassified genes in the other two tools (Table 1). doubletrouble does not specify or produce a singleton output file, hence it can be explained that it inherently considers the genes not assigned to the duplicate groups as singletons. DupGen_finder on the other hand, outputs only those genes that do not have self-local alignment hits as singletons. This ignores genes that hit up only against themselves in the local self-alignment. In the manual threshold specification to DupyliCate, genes that do not have self hits, genes that have only themselves as self hits, and genes whose second best hit within the genome fail to pass the threshold are classified as singleton genes, resulting in a few thousand singleton genes. But when using a BUSCO-based threshold for obtaining gene duplicates, genes that do not have self hits, genes that have only themselves as self hits, and genes whose second best hit within the genome fail to pass the BUSCO-derived threshold are classified as singleton genes. This causes the results to significantly differ giving around 27.5 % of gene duplicates in *A. thaliana*. However, the relatively lower number of gene duplicates in *A. thaliana* is supported by a low BUSCO duplication percentage (Supplementary Table S2). A potential biological explanation could be that only a certain proportion of genes were retained as duplicates after the polyploidization events reported in *A. thaliana*[26]. These estimates are also close to the estimate of 28% gene duplicates in *A. thaliana* previously reported[27] after the most recent alpha polyploidy event (Supplementary Figure 1)[26,28]. This puts forth the biological implication of the BUSCO-based thresholding method that distinguishes genes that escaped ancient duplication events and those that were retained in gene families or underwent more recent duplications. It thus allows for species-specific gene duplicates identification and classification. But it is also important to note that there is no clear boundary between singleton genes and duplicated genes. Both are rather extremes on a gradient. Therefore, the identification is imperfect, influenced by the cutoffs and other selection criteria determined by the users. Further the different number of singleton gene outputs in the different existing tools could as well be attributed to the fluidity of the singleton-duplicate segregation criteria. However, the BUSCO-based approach developed in this study offers one possible way of segregating singleton and duplicated genes with a reliable biological support. Additionally, the duplication landscape plot of *A. thaliana* output by DupyliCate (Fig. 1) also offers a visual inference on the genome-wide duplication status. Left and right skewed distributions in the plot could suggest genomes with extremely low gene duplications (as seen for *E. coli, S. cerevisiae* (Supplementary Results 1)) and genomes with extensive gene duplications or polyploidization (Supplementary Results 1), respectively. The approximately central, bell-shaped curve for *A. thaliana* (Fig. 1) corresponds well with the polyploidization-gene duplications loss events described for *A. thaliana*[27,29]. Looking at the performance metrics of the three tools (Fig. 2), DupGen_finder had the highest runtime, while doubletrouble had the lowest runtime. However, since an outgroup was specified in DupGen_finder, the inclusion of a forward alignment step leads to an increase in runtime. Thus doubletrouble and DupGen_finder can be considered to be on par with respect to runtimes. Since DupyliCate incorporates FASTA cleaning, GFF validation, and duplicate group clustering steps, the runtime is slightly longer, especially when BUSCO analysis is included. Also steps like isoform removal and execution of local alignment,

379 which need to be performed separately by the user prior to running doubletrouble and
380 DupGen_finder, are integrated in DupyliCate's run. Although this increases the runtime, it could
381 lead to possibly reduced hands-on time when using DupyliCate. Moreover, it is expected that the
382 availability of DupyliCate via docker and conda would further simplify the installation and use of
383 the tool, and expand its usability for users with different levels of bioinformatics expertise.
384
The runtime analyses involving different parameter and mode selection with the *Arabidopsis*
datasets give an idea of the runtimes for the different functionalities. As seen from Table 2, a
complete expression analysis, and Ka/Ks analysis of gene duplicates take longer. Hence, when
such analyses are intended, it is recommended to use as many cores as available to complete
the run in the shortest possible time. The expression analysis plots output by the use of the
specific duplicate analysis flag in the absence of a reference (Fig. 3) help analyze the functional
fates of the duplicated genes and determine expression divergence among them. The matrix
pattern of the plot helps visualize the gene expression trends among all the genes in the same
duplicate group at once. The entire RALF tandem duplicates array reported by Cao and Shi[17] -
AT2G19020, AT2G19030, AT2G19040, AT2G19045, was identified in the RALF tandem group
by DupyliCate along with the genes *AT2G19050* and *AT2G19060.* However, *AT2G19020,*
AT2G19040, and *AT2G19045* were found to be perceived (putative) pseudogenes based on the
expression analysis. Thus Fig. 3(c) shows the pairwise expression plots of the other genes in the
identified tandem duplicates array. But it is important to note that including information on the
spatial aspects of gene expression in the different tissues and under different conditions is
necessary to confidently ascertain the subfunctionalization, neofunctionalization, or
pseudogenization fates of the gene duplicates. From Fig. 4(b) it is evident that dataset size and
inclusion of phylogeny in the presence of a reference, are factors that influence the runtime. Large
datasets and inclusion of phylogeny for ortholog assignment in the presence of a reference lead
to an increase in the runtime. But incorporating phylogeny during ortholog assignments will
produce better ortholog assignments, as local alignment-based approaches alone are not enough
to identify orthologs correctly[6]. Hence phylogeny is recommended when a reference is used. In
case the user just wants to have a first hand assessment of gene copy numbers with comparative
analysis, runtime can be prioritized over accuracy by deactivating the phylogeny analysis.
The *SEC10* gene is incorrectly collapsed in the Col-0 genome sequence, but is properly
represented in the *A. thaliana* Nd-1 assembly and annotation[21,22,30]. This was confirmed through
the DupyliCate analysis where the *SEC10* gene duplication was identified as a tandem duplication
in the Nd-1 dataset compared to TAIR (Col-0). The first dedicated gene in the crocine biosynthesis
pathway of *Gardenia jasminoides* is a part of a four-gene tandem array that shares just one
ortholog with the *Coffea canephora* genome. This tandem duplication-led expansion of the gene
cluster is responsible for the initial critical steps in crocine biosynthesis that is not found in coffee[23].
The entire tandem cluster was correctly identified by DupyliCate demonstrating the usefulness of
representing gene duplicates in cluster groups instead of gene pairs. In the analysis of nematode
resistance locus between tolerant and susceptible sugarbeet lines[24], DupyliCate identified that
the crucial genes from the locus belong to a proximal duplicate cluster. This shows that proximal
duplication was responsible for the nematode tolerance phenotype seen in the tolerant line
StrubeU2Bv while exemplifying the application of DupyliCate to answer biological questions.
Finally, the 24ISO gene responsible for the first steps of withanolide biosynthesis in *W. somnifera*

was identified to be present in a dispersed duplicate group. It was also found to appear in the
duplicates relationships results along with the closely related SSR genes[25] providing a PoC for
the usefulness of the duplicates relationships file in the overlap mode and the mixed duplicates
file in the strict mode. These different PoC analyses show the ability of DupyliCate to correctly
identify gene duplicates and classify them into the different duplication classes. The gene
duplicates identification in the *Vitis* species lacking annotation was facilitated through the
GeMoMa flag of DupyliCate (Table 4). This option could be useful in cases where the samples of
interest lack structural annotation or a good quality structural annotation, but have a closely
related species with a good structural annotation that can serve as the hint for gene prediction
with GeMoMa. It is also helpful to note that the hint for GeMoMa annotation and the reference for
comparative analysis with DupyliCate could be the same or different. This means, the GeMoMa
flag can be used in the presence as well as in the absence of a reference for DupyliCate analysis,
but requires the specification of a hint for creating the annotation for the sample. In either case,
the species whose annotation is to be used by GeMoMa for the new annotation must be specified
in the GeMoMa config TXT file along with the names of the samples to be annotated. More details
are provided in the GitHub repository. In the rice genome sequence analysis, the rice weeds
showed a high number of gene duplicates (Table 4). The BUSCO analyses
(Supplementary_Table_S2) showed that these rice weeds have a high percentage of duplicated
BUSCO genes (>80%), and the pseudo-ploidy numbers (apparent indicators of ploidy of the
sample based on BUSCO gene frequencies) of *E. colona* and *E. oryzicola* were ~3 and ~2,
respectively. Their duplication landscape plots (Supplementary_Results_1) were also distinctively
right skewed suggesting a recent polyploidization event. The other rice varieties had a
comparable number of gene duplicates and showed a very similar trend of duplication landscape
plots except *Oryza glaberrima.* The plot of *O. glaberrima* was almost flat except for a marked high
in the number of genes with a self-normalized bit score of 1 indicating that the duplicates in the
genome are recent or have not diverged greatly (Supplementary Results 1). DupyliCate's
applicability to non-plant samples was shown by mining duplicate genes in *E. coli, S. cerevisiae*,
and *C. elegans*. The high number of gene duplicates identified in *C. elegans* corroborates the
widespread occurrence of gene duplications in *C. elegans*, when compared to other non-plant
eukaryotes like yeast and fruit fly[31].
453
Among the analyzed Brassicales plants for the *FLS* case study, *I. tinctoria, E. vesicaria*, and *S.*
alba had a pseudo-ploidy number of ~2 indicating their diploidy, while *B. napus* with a pseudo-
ploidy of ~3 was classified as triploid (Supplementary Table S2). This explains their high gene
duplication numbers. Coming to the ortholog assignments in the copy number analysis, *C. papaya*
FLS1 (evm.TU.supercontig_37.77) groups as a sister of the *AtFLS1* clade and is the only *FLS*
ortholog identified. It also has a good BUSCO completeness (>80%) (Supplementary Table S2)
indicating a biological phenomenon. *C. papaya* belongs to Caricaceae, one of the most basal
families in Brassicales[32]. This along with the phylogenetic tree (Fig. 5) suggests a possible
hypothesis on *FLS* evolution in the Brassicales - the true *FLS* had appeared in the early
Brassicales plants followed by an expansion of the *FLS* gene family in Brassicaceae after the
divergence of Caricaceae and Brassicaceae. The *B. oleracea* gene showing up in the *AtFLS2*
clade (Bol019362.v1.0) could probably be an ancient *FLS2* duplicate that cannot be placed
correctly only based on local alignment methods. Similarly, almost all the sequences falling in the

later subclade of *AtFLS4* (coloured by the red strips in *AtFLS4* clade in Fig. 5), have *AtFLS4*
appear as only the fourth or the fifth best forward hits of these genes in the forward local
alignment. These genes could also be ancient duplications of *FLS4* and hence difficult to assign
orthologs based on candidates from similarity based methods. Such difficult cases of ortholog
detection can be resolved with higher phylogenetic resolution by including more sequences that
give better placement and context to the query gene within its gene family supported by better
taxon sampling and global alignment cues[33]. Further, from the phylogenetic analysis, there is an
evident expansion of the *FLS3* and *FLS4* lineages happening in Brassicaceae, which is quite
intriguing. This is also observed in the *FLS* phylogenetic tree presented by Schilbert et al.[34].
Expression analyses performed in DupyliCate, found a majority of genes in the *FLS3/5* clades to
be pseudogenes. In the *FLS4* clade most of the *I. tinctoria* genes turned out to be pseudogenes.
A potential explanation for the *FLS3/4* expansion could come from the expression analysis. Since
these are non-functional *FLS* genes, they could be under relaxed selection pressure. Expansion
might be tolerated because additional copies do not impose strong fitness costs, allowing them
to accumulate mutations and gradually become pseudogenes.
The transcriptional regulation of flavonol biosynthesis in plants is achieved by the action of
MYB11, MYB12, and *MYB111* transcription factors (TFs)[35]. Schilbert and Glover analyzed the
evolution of flavonol regulators in the family Brassicaceae and reported the duplication of *MYB111*
and *MYB12* TFs in the family[35]. The MYB analysis presented here serves as a case study of these
flavonol regulators' evolution across a large phylogenetic range of plants. Flavonoid biosynthesis
is well known and established in land plants. Flavonols, a class of flavonoids that play a major
role in protecting against abiotic stress factors like UV exposure, are also widespread in land
plants[36–38]. These factors provide a basis for the reason behind the absence of flavonol regulating
MYB12 and *MYB111* orthologs in non-land plants and early land plants like mosses. The *MYB12,*
and *MYB111* orthologs mined in club mosses were examined and were found to belong to very
low confidence ortholog groups, which could mean that these species could also lack the *MYB12*
and *MYB111* orthologs. An alternative explanation could also be that algae, mosses, and other
basal plants might have other flavonol biosynthesis regulating MYBs. Looking at the other plants
in the analyses, all assignments that had very high confidence orthologous group confidence
scores (OGCS) in DupyliCate were found to match exactly with the phylogeny results. While
groups with moderate and low OGCS were found to have mismatches with the tree or have false
positive sequences. This demonstrates the usefulness of the OGCS scoring system developed
that can help refine and obtain more reliable results. Further, nearly half of the examined plant
species' ortholog assignments are exact matches with the tree with nearly one third having
matches with the presence of false positive sequences (Fig. 7). One possible reason for the
ortholog misassignments observed could be the widely varying evolutionary distance differences
between the reference *A. thaliana* and the analyzed sample species. This is also supported in the
phylogenetic tree where most of the sequences near the outgroup clade belong to either
gymnosperms, basal angiosperms, monocots, or polyploid plants like cotton. The phylogenetic
tree (Fig. 6) also provides some biological insights. It indicates the duplication of *MYB12* and
MYB111 in some dicots (purple and pink clades) that could have been derived from an ancestral
flavonol regulating MYB which is still observed in gymnosperms, basal angiosperms, monocots,
and some dicots. In conclusion, the MYB case study demonstrates the usability of the tool for

large scale analysis involving gene duplicates mining, gene copy number analysis in nominal
time, and its ability to provide biological insights.

Further, the parameter tuning analyses helped analyse the effect of the different settings
controlled by the user. As seen in Supplementary Table S10, --score and --self_simcut
parameters influenced the results significantly. As the values of both these parameters were
gradually increased, the number of identified gene duplicates lowered, while a decrease in these
cutoffs increased the gene duplicate numbers identified due to the relaxed thresholds. Next, the
type of sequence alignment tool used, influenced the results with an albeit low difference between
the different runs. Since the duplicate and singleton segregation relies on the self-normalized bit
score cutoff, differences in the bit scores put out by the different aligners can lead to slightly
different results. Nevertheless, considering the low runtime and small difference between the
gene duplicate numbers reported by the three aligners, the default DIAMOND aligner was found
to work well. The next most important parameter is the --proximity flag that determines the
maximum number of intervening genes between duplicates for them to be considered proximal
duplications. As expected, when the proximity value was increased, the number of proximal
duplicates increased and the number of dispersed duplicates decreased. The parameter --evalue
is the third most sensitive parameter and it slightly influences the total number of singletons and
duplicates identified. Low --evalue cutoffs led to slightly more gene duplicates being identified
while higher cutoffs lowered the number of gene duplicates identified (Supplementary Table S11).
The other parameter flags in the table --fwd_simcut, --scoreratio, --hits, --occupancy, --
synteny_score, --flank, and --side do not impact the overall gene duplicates singletons
segregation and classification. This is because all these parameter flags are relevant for forward
local alignment in case a reference is provided, and hence play a role in finding orthologs for the
sample gene duplicates in the reference organism. There is a detailed discussion on adjusting
these parameters for ortholog detection in the presence of a reference organism along with some
suggestions for parameter tuning of DupyliCate, in the Supplementary Discussion.

Moving on to the BUSCO vs manual threshold run results, there is an increase in gene duplicates
for manual mode with *A. thaliana*, *D. purpurea*, and *D. dumetorum* and a decrease in gene
duplicates in the manual mode with *V. cruziana*. This can be explained by looking at the manually
set normalized bit score cutoff used and the BUSCO-based normalized bit score cutoff used. The
BUSCO-derived self-normalized bit score threshold for *A. thaliana*, *D. purpurea*, *D. dumetorum*,
and *V. cruziana* were 0.696, 0.726, 0.822, and 0.430, respectively. Since the first three species
have a higher threshold when compared to the manual threshold of 0.5, the number of gene
duplicates were lower in the BUSCO mode and higher in the manual mode due to the lowering of
the cutoff. However, the scenario for *V. cruziana* is inverted since its species-specific threshold is
lower than 0.5, explaining the decrease in the gene duplicate numbers in the manual mode due
to an increase in the cutoff. The duplication landscape plots of these four species (Supplementary
Results 1) reveal that *A. thaliana* has an almost bell-shaped curve, while the other three species
show a right skewed curve, with *V. cruziana* having sufficient number of genes with low
normalized bit scores as well. The shape of *A. thaliana*'s plot combined with duplication loss
events reported in the species conform well with the low number of gene duplicates identified, as
discussed before. The lower normalized bit score threshold for *V. cruziana*, combined with the

almost similar proportions of gene duplicates and singletons, explain the right skewed shape of the curve with a significantly populated tail representing a good number of singletons too. The duplication plots for the other two plant species - *D. purpurea*, and *D. dumetorum* correspond well with the high proportion of gene duplicates identified. Thereby, these results show the applicability of DupyliCate to a wide taxonomic range of species with the parameters contributing to species-specific results that are reproducible.

In conclusion, DupyliCate is a high-throughput Python script developed for gene duplicates identification, classification, and integrated expression analysis. It has a number of adjustable parameters, species-specific singleton duplicates segregation, flexibility to handle a variety of GFF files, synteny analysis of small-scale duplicates (in the presence of a reference) and optional phylogeny-based ortholog detection for comparative gene copy number analysis. DupyliCate can thus be a valuable addition to the bioinformatics toolkit to support research in the domains of gene duplications research and comparative genomics for both plant and non-plant species.

Materials and methods

Datasets

Datasets of *Arabidopsis thaliana* Col-0 and Nd-1[22] were used to demonstrate the available modes, different aligner options, possible Ka/Ks analysis and expression analysis functionalities of DupyliCate. The two accessions were also used to demonstrate the PoC for *SEC10* duplication in the *A. thaliana* Col-0 dataset. *Gardenia jasminoides* and *Coffea canephora* datasets[23] were used to demonstrate the PoC for tandem duplication of the first gene in crocine biosynthesis in *G. jasminoides* with respect to *C. canephora*. Datasets of the *Beta vulgaris* lines StrubeU2Bv and KWSONT2320Bv[24] were used as PoC for proximal duplication of the nematode tolerance gene in the tolerant line StrubeU2Bv with respect to KWS2320. Datasets of *Withania somnifera* and *Physalis pruinosa*[25] were included to show PoC for dispersed duplication of the 24-Isomerase gene in *W. somnifera* with respect to *P. pruinosa*. DupyliCate was used on three *Vitis* datasets (*V. vinifera, V. rotundifolia,* and *V. amurensis*) to demonstrate applicability on complex genome sequences and also to demonstrate the use case of GeMoMa feature of the tool using the datasets of *V. rotundifolia* and *V. amurensis* that lacked structural annotation. A collection of 10 genome sequences of rice (*Oryza sativa*) and close relatives from the RiceRelativesGD database[39] were leveraged to demonstrate the performance of DupyliCate on monocot plants. Gene duplicates were mined in *O. sativa indica* variety and its relatives. The stress related genes in *O. sativa indica* sp. were used to create a copy number table to analyse the number of the corresponding orthologs in the other rice relatives. *Escherichia coli*[40]*, Saccharomyces cerevisiae*[41], and *Caenorhabditis elegans*[42] datasets were used to show PoC for deployment on non-plant species.

Application case studies

Arabidopsis thaliana is reported to have six copies of the flavonol synthase (*FLS1-6*) gene encoding a key enzyme in flavonol biosynthesis[43]. Of the copies, only *AtFLS1* is known to produce a catalytically functional protein whereas the functions of the other gene copies are yet to be uncovered fully. Comparative analyses of the *FLS* gene family in the Brassicales order combined with expression analysis can help better understand the evolutionary dynamics of this gene. A total of 12 datasets of the Brassicales family (detailed in Supplementary Table S1), for which expression data were available, were chosen to do an application case study on flavonol synthase evolution in the Brassicales using *Arabidopsis thaliana FLS1-6* as reference genes.

Next, a large collection of 153 plant datasets was retrieved from Phytozome (details specified in Supplementary Table S1). These datasets were used to undertake a large-scale gene duplicates mining study across plant datasets to demonstrate the high-throughput utility of the tool. Further, these datasets were also used to study the evolution of the key flavonol regulator MYBs (Myeloblastosis genes) - *MYB12* and *MYB111* across the plant kingdom with *A. thaliana MYB12* and *MYB111* as reference genes, to demonstrate the biological relevance of DupyliCate's results. The forward normalized bit score and forward similarity cutoffs were lowered to 0.2 and 30%, respectively, since the run involved samples quite distant from the reference *A. thaliana* to avoid missing potential orthologs in the samples.

The results of DupyliCate from the two application case studies were also validated by building a phylogenetic tree. For the FLS tree, flavanone 3-hydroxylase (F3H) sequences were used as outgroups. For the MYB tree the ortholog results from DupyliCate were visualized in a phylogenetic tree with anthocyanin regulating *MYB75* sequences as outgroups. Other closely related *A. thaliana* MYBs *AtMYB5, AtMYB11, AtMYB21, AtMYB24, AtMYB57, AtMYB99, AtMYB82*, and *AtMYB123* were also included in the tree. MAFFTv7.5.2.6[44] was used with the G-INS-i accuracy-oriented method for alignment of the amino acid sequences. pxclsq from PHYX[45] was used to remove alignment columns with occupancy below 10%. Phylogenetic trees were inferred using IQ-TREE3[46] with the parameters -m MFP -wsr and --alrt 1000 -B 1000[47]. The best-fit evolutionary model inferred with ModelFinder[48] for the FLS case study was Q.PLANT+R5 and for the MYB case study the best fitting model was JTT+R8. The phylogenetic trees were visualized using iTOL v7.2.2[49].

All the datasets were analyzed with DupyliCate v1.0 in the overlap mode with BUSCO-specific threshold option activated. In case, a reference was used, phylogeny was used to determine the ortholog relationships with the reference genes.

Workflow

A pipeline for gene duplicates identification, classification, and further downstream analyses was implemented in Python3 (Fig. 8). The overall workflow involves validation of input files followed by self-alignment of the sample(s) (sample(s) - species in which gene duplicates are to be identified). Species-specific thresholds are then used to segregate singleton genes from

duplicates and duplicates are categorized as tandem, proximal, dispersed, and mixed (a class
where genes can belong to more than one duplication class; this helps retain the relationship
across the different duplicate groups). In the presence of a reference organism (reference - the
species with respect to which the sample's genes are assigned orthologs and synteny analysis is
performed), forward alignment is performed. The classified gene duplicates are assigned to
orthologs in the reference and synteny analysis is performed for small-scale duplicate groups of
tandem duplicates and proximal duplicates. Depending on the mode chosen, a fourth category of
mixed duplicates is returned. DupyliCate also facilitates optional gene expression analysis of gene
duplicates. Further, it provides the user with the option to calculate Ka/Ks values of the duplicates.
The DupyliCate script was tested with its dependencies and tools whose version numbers are
specified in the workflow steps detailed below.

645
Figure 8: Schematic representation of DupyliCate's workflow. The steps of the pipeline are
numbered and the output files are alphanumerically labelled. The blue boxes indicate user
provided input files and green boxes indicate intermediate output files produced by the pipeline
that serve as inputs for further steps. Pipeline steps are colored yellow. Output files are shown
by the cyan coloured boxes. External tools are represented by the ovals. Dotted outlines around

the shapes indicate that they are not a part of the mandatory inputs, and dependencies or
routine outputs of the pipeline.

Step 1: Optional structural annotation: DupyliCate requires the structural annotation file(s)
(GFF3) along with one of the following (1) assembly sequence, or (2) coding sequences, or (3)
polypeptide sequences in FASTA format. In case structural annotation is not available for a
sample in which duplicates need to be analyzed, there is an option to provide the annotation of a
related species as a hint for the gene prediction process with GeMoMa v1.9[50].

Step 2: Input file(s) check and validation: The input files are first checked and validated before
the actual run starts. If the check fails, the script exits and the errors are recorded and displayed
sample-wise. In case of GFF errors, there is a helper script (AGAT_wrapper.py) provided along
with the main script that uses AGAT v1.5.1[51] to process and correct the GFF files.

Step 3: GFF-FASTA files standardization, error correction, and polypeptide file generation:
GFF files have widely varying attribute fields denoting the information about the different genomic
features. It is also necessary for the headers in the protein FASTA file to match correctly with the
text denoted by these different attribute fields. This is a major limitation hampering the deployment
of a number of existing tools on different GFF files. To address this issue, this work provides a
two-step solution. First, it offers another helper script (FASTA_fix.py) for the users to process the
FASTA file headers according to the attribute field text in the respective GFF files. Next, the main
DupyliCate script offers an option to the user to indicate the specific GFF attributes to be used by
the script when processing the GFF file of a particular species, through a simple config TXT file.
This helps to circumvent GFF-FASTA mismatch errors, and expand the usability of DupyliCate
on a number of GFF file flavours. The corrected and validated files then enter the main analysis
and are processed to generate the FASTA files of polypeptide sequences without considering
alternate transcripts.

Step 4: Quality check of input polypeptide files: Since the output depends on the quality of
the input files and is also influenced by the ploidy of the samples being analyzed, a BUSCO[52] -
based QC step is included. This provides detailed QC reports containing the BUSCO
completeness, duplication, and BUSCO-derived pseudo-ploidy number that indicates the ploidy
of the input sample(s).

Step 5: Threshold determination for singleton-duplicate segregation and self-alignment:
Before moving on to the duplication analysis, it is important to segregate singletons and duplicates
correctly. This in turn depends on the duplication landscape within each analyzed species. A new
method for such species-specific threshold determination based on BUSCO v6.0.0 is
implemented here. For this, two cutoffs - one based on self-normalized bit score (bit score of the
hit of a polypeptide sequence with another polypeptide sequence within the genome (second best
hit) relative to the bit score against itself) and self-similarity are offered. The complete BUSCO
single copy genes are retrieved for each sample. If enough BUSCO single copy genes are
present, the normalized bit score of the 95th percentile of the BUSCO single copy genes is picked
as the threshold for duplicate and singleton identification. By default, this BUSCO-based auto

threshold method is chosen. If BUSCO is not available or if the sample has a very low number of
BUSCO single copy genes, then the fallback is to go for a self-similarity threshold instead of
normalized bit score threshold. There is also an option to manually set the normalized bit score
threshold and self-similarity threshold, allowing for fine grained analysis. The detailed steps of
threshold determination can be found in Supplementary Figure 2.

Next, self-alignment of the sequences of a sample is performed. Options are provided to choose
between DIAMOND v2.1.14[53], BLAST v2.17.0[54], and MMseqs2[55] for the alignment step. Following
self-alignment, duplicates and singletons are segregated using the previously determined
threshold. Independent of the thresholding method, a plot folder called
Duplication_landscape_plots is produced. It contains the normalized self-bit score distribution
plots of samples. The overall pattern of the duplication landscape plot can be used to infer the
genome-wide duplication status of the species.

Step 6: Ortholog determination: If sequences of a reference are provided, forward alignment of
the sequences of each sample is performed against the sequences of this reference. Here again,
two thresholds based on similarity value of the hits in the forward alignment and forward
normalized bit score (bit score of a forward alignment hit of a polypeptide sequence relative to the
bit score of the hit of the polypeptide sequence against itself in self-alignment) are applied to
retain good quality forward hits for the further downstream steps. Next follows a comprehensive
ortholog detection step in the presence of a reference. The top n candidates (default value of
n=10) from local alignment algorithms like DIAMOND are subjected to a global alignment using
MAFFT v7.5.2.6[44], synteny analysis (the synteny analysis is performed by code developed as a
part of DupyliCate and hence does not rely on external dependencies), and gene tree inference
using FastTree2[56]. Phylogeny derived evolutionary distance assessments using DendroPy[57] are
utilized to obtain orthologs of genes in the samples among the reference genes. The detailed
steps in ortholog detection are shown in Supplementary Figure 3. Since a multi-evidence
approach is used for ortholog search, a confidence scoring scheme was developed to ascertain
the confidence of each of the assigned orthologs based on the number of evidence points needed
to determine orthology. The ortholog confidence scoring scheme is detailed in Supplementary
Table S8.

Step 7: Gene duplicates grouping and classification: Next, the duplicates clustering and
grouping step is performed per sample to give rise to the different gene duplicate groups shown
in Fig. 9 below. There are two main modes of DupyliCate. Gene duplicates are classified as
tandems, proximal, and dispersed duplicates in both modes. Differences arise between the
modes in the following aspects -

'Overlap': Some genes, repeat across groups and this mode also produces an output file called
'Duplicates_relationships' that connects the different genes repeating across the duplicate groups

'Strict': Gene duplicates do not repeat across groups and this mode produces a fourth class of
duplicates type called 'Mixed_duplicates' where the gene duplicates from different duplicate
groups are merged together to retain their individual classification and also their relationship
across the groups

Figure 9: Different gene duplicate classifications utilized by DupyliCate. The red and black lines in the dispersed duplicates, mixed duplicates and gene duplicates relationships representations indicate different contigs/ scaffolds/ chromosomes.

The classification and clustering approach is detailed in Supplementary Figure 4. As a clustering approach is used to obtain gene duplicate groups/arrays, for all the duplicate groups in the output, there is an internal confidence scoring scheme used to classify the gene duplicate group as low, moderate and high confidence group (Supplementary Figure 5) that can help in interpreting the results accordingly. Along with the different duplicate group files, the singletons in a sample species are also provided per species.

In case the user has a set of desired genes in the reference whose copy number variation (CNV) needs to be assessed in the sample, a specific analysis flag option (--specific_genes) is available. Use of this flag produces two files. One is a comprehensive file providing information about the duplication and singleton status of the orthologs of the desired reference genes in the sample. Second is a copy number TSV file that provides the copy numbers of the specified reference genes across the different samples along with the orthologous group confidence score (OGCS) (OGCS scoring explained in Supplementary Table S9) to convey the confidence of the identified CNVs. This step also outputs a copy number XLSX file that has cells with no orthologs in the samples automatically colour coded in red, to allow for easy visual interpretation of presence-absence variation of the specified reference genes across the species analyzed.

Step 8: Synteny analysis: In the presence of a reference, synteny analysis is also carried out for small-scale gene duplicate groups of tandem and proximal duplicates, to add a confidence layer of genomic positional context with respect to the reference. If a reference is involved, detailed gene duplicate group nature details including gene group expansion and *de novo* duplication are provided for small-scale gene duplicate groups of tandem and proximal duplicates in the output. More information on the synteny analysis and gene duplicate group nature attributes have been described in Supplementary Figure 6 and Supplementary Figure 7, respectively. In the reference-free mode, ortholog detection, synteny analysis, and the gene group nature analysis steps are skipped.

Step 9: Ka/Ks computation: Optional downstream evolutionary analyses like Ka/Ks computation is performed for all gene duplicates on an individual gene basis using either Nei Gojobori (NG)[58] or Modified Yang Nielsen (MYN)[59] methods. These methods have been reimplemented from Ka_Ks_Calculator[10,60]. There is also in-built code facilitating amino-acid to codon-alignments and trimming of the multiple sequence alignments, thus removing the need for external tools in these steps. While an exact implementation of the NG method is available, the MYN method implemented is an approximation of the MYN method implemented in Ka_Ks_Calculator (Supplementary Figure 8 and Supplementary Results 1). Supplementary Figure 9 details the computational steps involved in the Ka/Ks analysis.

Step 10: Gene duplicates expression analysis and statistical analysis: If expression data is available for the sample(s), optional expression analysis of gene duplicates can also be performed by DupyliCate. This step gives out comprehensive information about the correlation among genes in duplicate groups, generates pairwise gene expression plots in a matrix figure and also helps determine divergent expression among duplicates as shown in Supplementary Figure 10. This step also outputs a list of perceived (putative) pseudogenes that have a low average expression value across samples. Since spatial information of gene expression in the different tissues is not involved in this analysis, genes with low average expression value across the samples are designated as perceived pseudogenes. The method used in this expression analysis was adopted and modified from the method implemented by Qiao and co-workers[9].

Benchmarking

Arabidopsis thaliana Col-0 and Nd-1 accessions were used to benchmark DupyliCate v1.0, DupGen_finder v1.0.0[9], and doubletrouble v1.8.0[15]. DupyliCate was used in the overlap mode with *A. thaliana* Col-0 as the sample. DupyliCate additionally offers species-specific cutoff for segregating singletons and duplicates using BUSCO. Hence two runs of the tool were performed with *Arabidopsis thaliana* Col-0 dataset using BUSCO-based cutoff method and user-specified cut-off method. For the user-specified cutoff method results to be comparable with the other tools, a cutoff of 0 was specified for both normalized bit score and self similarity parameter flags. Since the standard classification scheme of doubletrouble was the closest to the classification scheme of DupyliCate, it was adopted for the doubletrouble run with *A. thaliana* Col-0 as the target. Transposed duplicates (TDs) are not classified in the standard scheme, hence, the doubletrouble run involved the use of only the target species *A. thaliana* Col-0. For DupGen_finder, *A. thaliana* Nd-1 was the outgroup and *A. thaliana* Col-0 as the target, as DupGen_finder does not offer

793 classification schemes without TD classification, necessitating the use of an outgroup species.
794 BLAST was used as the aligner for all three tools with a resource allocation of 15 threads during
795 the benchmarking runs to avoid differences due to different sensitivity parameters in DIAMOND.
796
DupyliCate mainly assigns duplicates to tandems, proximals, and dispersed groups. Therefore,
all the other duplicate group classifications apart from tandem and proximal duplicates in
DupGen_finder and doubletrouble were processed with custom Python scripts to obtain a single
miscellaneous group ('Misc.' in Table. 1) that is comparable to the dispersed duplicates group in
DupyliCate. The benchmarking was therefore done mainly with respect to the number of genes
classified as tandem, proximal, and dispersed duplicates by the three tools. Also, the runtime of
the three tools was compared. Five replicate runs were performed on the *A. thaliana* dataset using
each of the three tools to estimate the runtime variability given the use of the same computational
resources. Since local alignment is a step included within a run of DupyliCate, the local alignment
times were also included in the runtime for the other two tools to standardize the comparison. The
additional codes, and commands used for running doubletrouble, running DupGen_finder, and
performing the benchmarking analysis can be found in the GitHub repository of DupyliCate
(https://github.com/ShakNat/DupyliCate).

Runtime analyses

The runtime of DupyliCate depends on the selected mode and parameter settings. The following
approach was adopted to offer a comprehensive runtime analysis when using the different modes
and parameter settings. The runtime for each run of the tool using *A. thaliana* datasets was
recorded for the following run variations: 1) overlap and strict mode; 2) phylogeny and non-
phylogeny-based ortholog detection in the presence of a reference; 3) expression analysis of all
identified gene duplicates; 4) expression analysis of specified gene duplicates (XPB tandem
duplication[19], BLA[61], and RALF tandem duplicates array[17] reported in *A. thaliana* Col-0 were
chosen for demonstrating the specific expression analysis feature) 5) Ka/Ks analysis of all
identified gene duplicates; and 6) run with different aligners - BLAST/ DIAMOND/ MMseqs2.
Next, to analyze the tool runtime increase with an increasing number of input datasets, a
comprehensive runtime analysis was conducted with the Brassicales plants used for the FLS case
study. The different datasets were randomly grouped into groups of sizes 2, 4, 6, 8, 10, and 12,
respectively. Runtimes were recorded using each group as an input for DupyliCate. To account
for the variation in the file sizes of the datasets in each group, there were five random sets of the
same group length generated, and the runtime was recorded for every such replicate run.
Finally, to get an idea of the performance of DupyliCate in large-scale studies, the runtime was
also analyzed for the tool run on the 153 plant datasets used for the MYB case study that was
done in the overlap mode using DIAMOND as the aligner, *A. thaliana* as the reference, and
integrated phylogeny for ortholog detection.
830

Parameter tuning analysis

DupyliCate offers a number of qualitative and quantitative parameter flags that can be adjusted
and tuned according to the specific needs of the user. Among these, most of the qualitative

parameters take in the input file paths, and intermediate dependency tool paths and do not
contribute to variability in the final results. Hence, twelve parameters that could significantly
impact the results in a concrete quantitative manner were selected for tuning trials that were
performed with the *A. thaliana* Col-0 dataset as the sample and *A. thaliana* Nd-1 dataset as the
reference. One factor at a time (OFAT) approach was employed for these trials wherein only one
parameter was changed at a time, and the other parameters remained constantly at default. For
almost all the parameters, tuning was performed with three values - the default value, a value
25% lower than the default and a value 25% higher than the default. In case of parameters that
accepted only integer values, the low and high values were rounded off to the next nearest integer.
For the --score parameter alone, a slightly different approach was adopted. The default for this
parameter flag is 'auto' which activates the BUSCO-based threshold method, and calculates a
species-specific normalized bit score threshold for singleton-duplicate segregation. The other
option for this --score flag is manual where the user needs to enter specific values. Since the auto
mode does not depend on user input at all and is entirely species-dependent, the self-similarity
threshold is switched off or is automatically set to 0 (--self_simcut is 0). But in the manual mode
the --score flag must be provided by the user and the default value of 50% for the --self_simcut
parameter flag is used. But this --self_simcut value can also be adjusted by the user in the manual
mode. Thus the --self_simcut flag jointly influences the singleton-duplicates segregation along
with the --score flag. Therefore the user needs to know the contributions of self-normalized bit
score cutoff and self-similarity cutoff individually and taken together to select suitable values when
setting them manually. Hence, a detailed combinatorial analysis was performed with these two
parameters, where one flag was set to 0 and the other was set to the low, default, and high values,
to demonstrate their individual effects. Another set of analyses that combined the low, default and
high values from both these parameters were also performed to show their combined effects.
858
Further, after the tuning trials, a set of BUSCO-based threshold vs. manually set threshold runs
were performed using DupyliCate on the plant species *Digitalis purpurea*[62], *Victoria cruziana*[63],
and *Dioscorea dumetorum*[64] along with *Arabidopsis thaliana* Col-0. The manual threshold runs
were performed with --score of 0.5 (0.5 was chosen since it is a median-normalized bit score
cutoff that is bounded by 0 and 1) and --self_simcut of 50% (default value of the --self_simcut
flag). The BUSCO vs manual runs were performed to give a comparative analysis of the results
between the two approaches that can be used for adjusting the parameters of a DupyliCate run.
The above mentioned plant species were specifically chosen for this analysis in order to show the
applicability of the method and the reproducibility of results over a wide variety of plant species.

Data Availability

869 DupyliCate, the helper scripts, and test data sets can be accessed at:
870 https://github.com/ShakNat/DupyliCate. Underlying data sets are publicly available at the NCBI,
871 Phytozome, and other databases as specified in Supplementary Table S1. Details of the
872 supplementary data can be found in Supplementary Information.

AI Statement

AI tools (ChatGPT and Claude) were used for optimizing, restructuring, and debugging the DupyliCate script. Claude was also used to check the language and grammar of the manuscript text.

Reference

1. Lallemand, T., Leduc, M., Landès, C., Rizzon, C. & Lerat, E. An Overview of Duplicated Gene Detection Methods: Why the Duplication Mechanism Has to Be Accounted for in Their Choice. *Genes* **11**, 1046 (2020).

2. Zhang, X. & Smith, D. R. An overview of online resources for intra-species detection of gene duplications. *Front. Genet.* **13**, (2022).

3. Durand, D. & Hoberman, R. Diagnosing duplications – can it be done? *Trends Genet.* **22**, 156–164 (2006).

4. Conant, G. C. & Wagner, A. GenomeHistory: a software tool and its application to fully sequenced genomes. *Nucleic Acids Res.* **30**, 3378–3386 (2002).

5. Li, L., Stoeckert, C. J. & Roos, D. S. OrthoMCL: Identification of Ortholog Groups for Eukaryotic Genomes. *Genome Res.* **13**, 2178–2189 (2003).

6. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* **20**, 238 (2019).

7. Wang, Y. *et al.* MCScanX: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Res.* **40**, e49 (2012).

8. Djedatin, G., Monat, C., Engelen, S. & Sabot, F. *DuplicationDetector*, a light weight tool for duplication detection using NGS data. *Curr. Plant Biol.* **9–10**, 23–28 (2017).

9. Qiao, X. *et al.* Gene duplication and evolution in recurring polyploidization–diploidization cycles in plants. *Genome Biol.* **20**, 38 (2019).

10. Wang, D., Zhang, Y., Zhang, Z., Zhu, J. & Yu, J. KaKs_Calculator 2.0: A Toolkit

Incorporating Gamma-Series Methods and Sliding Window Strategies. *Genomics*

Proteomics Bioinformatics **8**, 77–80 (2010).

11. Zhang, X., Hu, Y. & Smith, D. R. HSDFinder: A BLAST-Based Strategy for Identifying

Highly Similar Duplicated Genes in Eukaryotic Genomes. *Front. Bioinforma.* **1**, (2021).

12. Zhang, X., Hu, Y., Smith, D. R., Cheng, Z. & Archibald, J. M. HSDSnake: a user-friendly

SnakeMake pipeline for analysis of duplicate genes in eukaryotic genomes. *Bioinformatics*

41, btaf325 (2025).

13. Sun, P. *et al.* WGDI: A user-friendly toolkit for evolutionary analyses of whole-genome

duplications and ancestral karyotypes. *Mol. Plant* **15**, 1841–1851 (2022).

14. Kirilenko, B. M. *et al.* Integrating gene annotation with orthology inference at scale.

Science **380**, eabn3107 (2023).

15. Almeida-Silva, F. & Van de Peer, Y. doubletrouble: an R/Bioconductor package for the

identification, classification, and analysis of gene and genome duplications. *Bioinformatics*

btaf043 (2025) doi:10.1093/bioinformatics/btaf043.

16. Hanke, D. M. & Dagan, T. SegMantX: a novel tool for detecting DNA duplications

uncovers prevalent duplications in plasmids. Preprint at

https://doi.org/10.1101/2025.03.14.643293 (2025).

17. Cao, J. & Shi, F. Evolution of the RALF Gene Family in Plants: Gene Duplication and

Selection Patterns. *Evol. Bioinforma. Online* **8**, 271–92 (2012).

18. Stracke, R. *et al.* Differential regulation of closely related R2R3-MYB transcription factors

controls flavonol accumulation in different parts of the Arabidopsis thaliana seedling. *Plant*

J. **50**, 660–677 (2007).

19. Masuda, H. P. *et al.* Evidence for sub-functionalization of tandemly duplicated *XPB*

nucleotide excision repair genes in *Arabidopsis thaliana*. *Gene* **754**, 144818 (2020).

20. Tao, F., Sollapura, V., Robert, L. S. & Fan, C. Neofunctionalization of tandem duplicate

genes encoding putative β-L-arabinofuranosidases in Arabidopsis. *Plant Physiol.* **192**,

2855–2870 (2023).

21. Pucker, B. *et al.* A De Novo Genome Sequence Assembly of the Arabidopsis thaliana

Accession Niederzenz-1 Displays Presence/Absence Variation and Strong Synteny. *PLOS*

ONE **11**, e0164321 (2016).

22. Pucker, B. *et al.* A chromosome-level sequence assembly reveals the structure of the

Arabidopsis thaliana Nd-1 genome and its gene set. *PLOS ONE* **14**, e0216233 (2019).

23. Xu, Z. *et al.* Tandem gene duplications drive divergent evolution of caffeine and crocin

biosynthetic pathways in plants. *BMC Biol.* **18**, 63 (2020).

24. Sielemann, K. *et al.* Genomic characterization of a nematode tolerance locus in sugar

beet. *BMC Genomics* **24**, 748 (2023).

25. Hakim, S. E. *et al.* Phylogenomics and metabolic engineering reveal a conserved gene

cluster in Solanaceae plants for withanolide biosynthesis. *Nat. Commun.* **16**, 6367 (2025).

26. Barker, M. S., Vogel, H. & Schranz, M. E. Paleopolyploidy in the Brassicales: Analyses

of the Cleome Transcriptome Elucidate the History of Genome Duplications in Arabidopsis

and Other Brassicales. *Genome Biol. Evol.* **1**, 391–399 (2009).

27. Charon, C., Bruggeman, Q., Thareau, V. & Henry, Y. Gene duplication within the Green

Lineage: the case of TEL genes. *J. Exp. Bot.* **63**, 5061–5077 (2012).

28. Gao, B. *et al.* Evolution by duplication: paleopolyploidy events in plants reconstructed by

deciphering the evolutionary history of VOZ transcription factors. *BMC Plant Biol.* **18**, 256

(2018).

29. Blanc, G. & Wolfe, K. H. Functional Divergence of Duplicated Genes Formed by

Polyploidy during Arabidopsis Evolution. *Plant Cell* **16**, 1679–1691 (2004).

30. Vukašinović, N. *et al.* Dissecting a Hidden Gene Duplication: The Arabidopsis thaliana

SEC10 Locus. *PLOS ONE* **9**, e94077 (2014).

31. Woollard, A. Gene duplications and genetic redundancy in C. elegans. in *WormBook:*

The Online Review of C. elegans Biology [Internet] (WormBook, 2005).

32. Delaux, P.-M., Séjalon-Delmas, N., Bécard, G. & Ané, J.-M. Evolution of the plant–

microbe symbiotic 'toolkit'. *Trends Plant Sci.* **18**, 298–304 (2013).

33. Emms, D. M. & Kelly, S. SHOOT: phylogenetic gene search and ortholog inference.

Genome Biol. **23**, 85 (2022).

34. Schilbert, H. M. *et al.* Characterization of the Brassica napus Flavonol Synthase Gene

Family Reveals Bifunctional Flavonol Synthases. *Front. Plant Sci.* **12**, (2021).

35. Schilbert, H. M. & Glover, B. J. Analysis of flavonol regulator evolution in the

Brassicaceae reveals MYB12, MYB111 and MYB21 duplications and MYB11 and MYB24

gene loss. *BMC Genomics* **23**, 604 (2022).

36. Davies, K. M. *et al.* The Evolution of Flavonoid Biosynthesis: A Bryophyte Perspective.

Front. Plant Sci. **11**, (2020).

37. Davies, K. M. *et al.* The evolution of flavonoid biosynthesis. *Philos. Trans. R. Soc. B Biol.*

Sci. **379**, 20230361 (2024).

38. Yonekura-Sakakibara, K., Higashi, Y. & Nakabayashi, R. The Origin and Evolution of

Plant Flavonoid Metabolism. *Front. Plant Sci.* **10**, 943 (2019).

39. Mao, L. *et al.* RiceRelativesGD: a genomic database of rice relatives for rice research.

Database **2019**, baz110 (2019).

40. Karberg, K. A., Olsen, G. J. & Davis, J. J. Similarity of genes horizontally acquired by

Escherichia coli and Salmonella enterica is evidence of a supraspecies pangenome. *Proc.*

Natl. Acad. Sci. **108**, 20154–20159 (2011).

41. Cliften, P. F., Fulton, R. S., Wilson, R. K. & Johnston, M. After the Duplication: Gene

Loss and Adaptation in Saccharomyces Genomes. *Genetics* **172**, 863–872 (2006).

42. Ma, F., Lau, C. Y. & Zheng, C. Young duplicate genes show developmental stage- and

cell type-specific expression and function in Caenorhabditis elegans. *Cell Genomics* **4**,

100467 (2024).

43. Owens, D. K. *et al.* Functional Analysis of a Predicted Flavonol Synthase Gene Family in

Arabidopsis. *Plant Physiol.* **147**, 1046–1061 (2008).

44. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

45. Brown, J. W., Walker, J. F. & Smith, S. A. Phyx: phylogenetic tools for unix.

Bioinformatics **33**, 1886–1888 (2017).

46. Wong, T. K. F. *et al.* IQ-TREE 3: Phylogenomic Inference Software using Complex

Evolutionary Models. https://ecoevorxiv.org/repository/view/8916/ (2025).

47. Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:

Improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

48. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,

587–589 (2017).

49. Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to the

phylogenetic tree display and annotation tool. *Nucleic Acids Res.* **52**, W78–W82 (2024).

50. Keilwagen, J., Hartung, F. & Grau, J. GeMoMa: Homology-Based Gene Prediction

Utilizing Intron Position Conservation and RNA-seq Data. in *Gene Prediction: Methods and*

Protocols (ed. Kollmar, M.) 161–177 (Springer, New York, NY, 2019). doi:10.1007/978-1-

4939-9173-0_9.

51. Dainat, J. *et al.* NBISweden/AGAT: AGAT v1.5.1. Zenodo

https://doi.org/10.5281/zenodo.16317950 (2025).

52. Tegenfeldt, F. *et al.* OrthoDB and BUSCO update: annotation of orthologs with wider

sampling of genomes. *Nucleic Acids Res.* **53**, D516–D522 (2025).

53. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale

using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

54. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment

search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

55. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for

the analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

56. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately Maximum-

Likelihood Trees for Large Alignments. *PLOS ONE* **5**, e9490 (2010).

57. Sukumaran, J. & Holder, M. T. DendroPy: a Python library for phylogenetic computing.

Bioinformatics **26**, 1569–1571 (2010).

58. Nei, M. & Gojobori, T. Simple methods for estimating the numbers of synonymous and

nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* **3**, 418–426 (1986).

59. Wang, D.-P., Wan, H.-L., Zhang, S. & Yu, J. γ-MYN: a new algorithm for estimating Ka

and Ks with consideration of variable substitution rates. *Biol. Direct* **4**, 20 (2009).

60. Zhang, Z. KaKs_Calculator 3.0: Calculating Selective Pressure on Coding and Non-

Coding Sequences. *Genomics Proteomics Bioinformatics* **20**, 536–540 (2022).

61. Ugalde, J. M. & Straube, H. New genes on the block: Neofunctionalization of tandem

duplicate genes with putative new functions in Arabidopsis. *Plant Physiol.* **192**, 2574–2576

(2023).

62. Horz, J. M., Wolff, K., Friedhoff, R. & Pucker, B. Genome sequence of the ornamental

plant Digitalis purpurea reveals the molecular basis of flower color and morphology

variation. 2024.02.14.580303 Preprint at https://doi.org/10.1101/2024.02.14.580303 (2025).

63. Nowak, M. S. *et al.* Genome sequence and RNA-seq analysis reveal genetic basis of

flower coloration in the giant water lily Victoria cruziana. 2024.06.15.599162 Preprint at

https://doi.org/10.1101/2024.06.15.599162 (2024).

64. Siadjeu, C., Pucker, B., Viehöver, P., Albach, D. C. & Weisshaar, B. High Contiguity de

novo Genome Sequence Assembly of Trifoliate Yam (Dioscorea dumetorum) Using Long

Read Sequencing. *Genes* **11**, 274 (2020).

1027

Acknowledgements

The authors thank the Plant Biotechnology and Bioinformatics group (University of Bonn) for valuable inputs and inspiring discussions. SN was supported by the doctoral scholarship of the German Academic Exchange Service (DAAD). This work was supported by the de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI) and ELIXIR-DE (Forschungszentrum Jülich and W-de.NBI-001, W-de.NBI-004, W-de.NBI-008, W-de.NBI-010, W-de.NBI-013, W-de.NBI-014, W-de.NBI-016, W-de.NBI-022). Open access funding was enabled by Project DEAL.

Funding

Open access funding was enabled by Project DEAL. There was no specific funding for this research.

Author information

Authors and Affiliations

Plant Biotechnology and Bioinformatics, Institute for Cellular and Molecular Botany, University of Bonn, Kirschallee 1, 53115, Bonn, Germany

Shakunthala Natarajan & Boas Pucker

Contributions

S.N. and B.P. designed the study. S.N. wrote the software and conducted the bioinformatic analyses. S.N. and B.P. interpreted the results and wrote the manuscript. All authors read the final version of the manuscript and agreed to its submission.

Corresponding author

Correspondence to pucker@uni-bonn.de

Ethics declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.