1

Conservation and divergence of sex-biased gene expression across

50 million years of *Drosophila* evolution

7   Amanda Glaser-Schmitt* and John Parsch*

12   Division of Evolutionary Biology, Faculty of Biology, Ludwig-Maximilians-Universität

13   München, Munich, Germany

16   **Running title:** Sex-biased gene expression in *Drosophila*

18   *Corresponding authors:

19   Email: glaser@bio.lmu.de (AGS); parsch@bio.lmu.de (JP)

Abstract

In sexually dimorphic species, sex-biased gene expression plays an important role in driving morphological, physiological, and behavioral differences between males and females. Here, we examined patterns of sex-biased gene expression within and among 6 *Drosophila* species with divergence times ranging from 2–50 million years. We detected contrasting patterns of sex bias conservation and turnover between heads and bodies, with more extensive sex-biased expression and greater conservation of sex-biased expression in the body, but more species-specific turnover of sex-biased expression in the head, where conserved, unbiased expression was common. Interestingly, in both body parts lineage-specific gains of sex-biased expression occurred most often via concordant expression changes in both sexes that differed only in their magnitude, with this pattern being particularly strong in the head, suggesting that the majority of lineage-specific sex bias gains do not represent a resolution of sexual antagonism, but instead reflect regulatory changes shared between the sexes. We detected an enrichment of putatively positively selected expression changes among sex-biased genes in both body parts. Taken together, our findings suggest sex-biased changes may often be facilitated by selection acting on the sex with lower expression, especially for male expression of female-biased genes in the body. We also detected differences in the proportion of sex-biased genes located on the X chromosome depending on sex bias and body part. Overall, our results provide novel insights into the dynamics of sex-biased gene expression, as well as the molecular mechanisms and selective forces underlying its turnover, across short and long evolutionary timescales.

Introduction

Gene expression variation underlies much of the phenotypic variation that we observe among species, populations, and individuals of the same species (King and Wilson 1975; Wray et al. 2003; Buchberger et al. 2019). In species with sexual dimorphism, this gene expression variation is particularly important as the vast majority or, in some cases, all of the genome is shared by both sexes (Ellegren and Parsch 2007). Therefore, sex-biased (SB) gene expression, i.e. the differential expression of a shared genome between males and females, plays an important role in driving morphological, physiological, and behavioral differences between the sexes (reviewed in Parsch and Ellegren 2013; Grath and Parsch 2016; Tosto et al. 2023). Indeed, quantifying and understanding patterns of sex-biased gene expression over both short and longer evolutionary timescales can help us better understand sexual

conflict and the evolutionary forces that shape and maintain phenotypic variation, especially between the sexes, which in turn can shed light on biodiversity and the evolution of complex phenotypic traits (reviewed in Tosto et al. 2023). For example, sex-specific gene expression and regulation have been implicated in the susceptibility and presentation of several human diseases (e.g. Shi et al. 2016; Khodursky et al. 2022; Fass et al. 2024), as well as variation in human height (Naqvi et al. 2019), all of which represent complex, highly polygenic phenotypes.

Studies have shown that sex-biased gene expression is extensive (Meiklejohn et al. 2003; Yang et al. 2006; Oliva et al. 2020), can be tissue-, developmental stage-, and cell type-specific (Yang et al. 2006; Ma et al. 2018; Naqvi et al. 2019; Oliva et al. 2020; Khodursky et al. 2022; Rodríguez-Montes et al. 2023), and evolves rapidly (Assis et al. 2012; Whittle and Johannesson, 2013; Harrison et al. 2015; Yang et al. 2016; Xie et al. 2025). The shared genome has long been thought to impose genetic constraints on the sexes (Lande 1980; Griffen et al. 2013), and SB gene expression is thought to be an indication of this ongoing sexual conflict (i.e. sexual antagonism) or its resolution (Connallon and Knowles 2005; Innocenti and Morrow 2010). Indeed, previous studies of SB gene expression in diverse taxa have found that sexual conflict is often resolved via SB gene expression and/or sex-specific regulation (e.g. Cheng and Kirkpatrick 2016; Dean et al. 2016; Wright et al. 2018).

The model insect, *Drosophila melanogaster*, was utilized in some of the earliest studies of SB gene expression (Jin et al. 2001; Arbeitman et al. 2002; Ranz et al. 2003; Parisi et al. 2003) and has continued to be an important model system for studying its evolution at both the interspecific and intraspecific levels (e.g. Meiklejohn et al. 2003; Ranz et al 2004; Gibson et al. 2004; Catalán et al. 2012; Llopart et al. 2012; Meiklejohn et al. 2014; Huylmans and Parsch 2014; Khodursky et al. 2020). Despite this progress, transcriptomic studies that systematically compare SB gene expression among species while also assessing the full range of within-species expression diversity are currently lacking. Previous studies in the genus *Drosophila* were restricted in their phylogenetic scale (e.g. Ranz et al. 2003; Grath and Parsch 2012; Müller et al. 2012; Khodursky et al. 2020) or had limited samples for determining within-species expression variation (e.g. Zhang et al. 2007; Pal et al. 2023).

The characterization of expression variation both within and between species is important for inferring the type of selection that influences gene expression, as the two levels of variation are thought to represent, respectively, the short-term and long-term outcomes of the same underlying evolutionary process (Kimura 1983; Hudson et al. 1987;

McDonald and Kreitman 1991). To address this issue, we measured gene expression in two

body parts (head and body) of adult males and females of 6 *Drosophila* species spanning

divergence times of up to 50 million years (Tamura et al. 2004). Within each species, we

included 5–8 strains derived from diverse geographic locations. These data allowed us to

detect the conservation and turnover of SB gene expression across the genus in samples that

include or exclude the gonad, which is the tissue with the highest proportion of SB genes

(Parisi et al. 2004). Furthermore, we were able to compare levels of expression

polymorphism and divergence to identify genes showing evidence for adaptive regulatory

evolution. We found that there is a much greater extent and conservation of SB expression

in the body than in the head. Most genes in the head showed conserved unbiased (UB)

expression, with reversals in the sex bias of a gene, i.e. from male-biased (MB) to female-

biased (FB) or vice versa, occurring only very rarely. Interestingly, in both body parts,

lineage-specific gains of SB expression typically involved concordant expression changes in

both sexes that differed only in their magnitude, with this pattern being particularly strong in

the head. We detected an enrichment of putatively positively selected gene expression

changes among SB genes in both body parts, and found evidence suggesting that these

changes may often be facilitated by selection acting on expression in the sex with lower

expression, especially for male expression of FB genes in the body. Together, these results

suggest that, intriguingly, the most common route to a gain in sex bias may be through

shared regulatory mechanisms between the sexes, rather than sex-specific regulation.

Overall, our results shed light on the long-term evolutionary dynamics of sex-biased gene

expression in the *Drosophila* genus.

Results

In order to study the evolution of sex-biased gene expression over a range of

timescales, we performed RNA-seq of male and female heads and bodies of 6 *Drosophila*

species with divergence times ranging between 2–50 million years (Fig 1), including *D.*

melanogaster (*Dmel*), *D. simulans* (*Dsim*), *D. suzukii* (*Dsuz*), *D. ananassae* (*Dana*), *D.*

subobscura (*Dsub*), and *D. immigrans* (*Dimm*). To get a comprehensive view of expression

variation within each species, we sampled 5–8 strains from a broad geographic distribution,

encompassing strains from multiple populations across three continents, including strains

from the putative ancestral range for *Dmel*, *Dsim*, *Dana*, and *Dsuz* (Supplementary Table S1).

We analyzed expression in 9,016 1:1 orthologs present in all species (Thiébaut et al. 2024),

7,570–8,161 of which were expressed in the head and 8,936–9,006 of which were expressed

in the body of each species. When we examined gene expression variation across all samples

in all species and sexes, the strains clustered strongly by body part (Fig 2A). In the head, the

sexes showed only slight separation (Fig 2B), while in the body, males and females were

clearly differentiated (Fig 2A, C). Within each sex and body part, strains of each species

clustered together, although these clusters often overlapped among species. *Dimm*, which is

the most distantly related to the other species (Fig 1), also showed the greatest interspecific

differentiation (Fig 2B, C). Within each species, patterns were similar, with body parts and

sexes clustering by sample type but differentiation between the sexes much stronger in the

body than the head (Supplementary Fig S1).

131

Fig 1: Phylogeny and sex-biased expression turnover in the examined *Drosophila* species in

the A) head and B) body. Branch lengths are shown only roughly to scale. Numbers at each

tip or node indicate sex bias gains (+) and losses (-) for female- (red) and male-biased (blue)

genes. The subgenera are shown in the light blue boxes, the species groups in the gray

boxes, and the subgroups in the light green boxes (*subobscura*: *sub*; *ananassae*: *ana*). *Gains

and losses could not be polarized for *Dimm* due to the lack of an outgroup; therefore, the

number of genes with different sex bias relative to the other species are shown.

and C) the body. The legend at the bottom right indicates the color associated with each

species and the shape indicates the sex (male: M; female: F) and body part.

Patterns and magnitude of sex-biased expression differ among body parts and species

Within each species, SB gene expression was extensive in the body, with 71.9–81.6% of genes showing sex bias versus 2.0–21.7% of genes in the head (Table 1). In comparison to the other species, sex bias in *Dsub* heads was very low (2.0% versus 10.1–21.7%; Table 1), especially when considering MB genes (32 versus 212–814 genes; Table 1). We detected more genes as SB in the *Dmel* head in comparison to the other species, but this excess was most likely driven by the sampling of additional strains in this species, as down-sampling resulted in the detection of similar numbers of SB genes as in the other species (Supplementary Table S2). Due to the large amount of sex bias in the body, the majority of genes that could be examined in both body parts (64.2–73.9%) were SB in the body but not the head (Fig 3C, Supplementary Fig S2). The next largest category of overlap was genes UB in both the body and head (14.1–25.2%), with genes showing the same sex bias in both body parts being the third largest category (0.9–13.0%). Interestingly, genes that switched sex bias between the head and body were more common than genes SB in the head but not the body

159   (0.7–5.6% versus 0.4–3.1%, respectively). Thus, genes SB in the head were 3–6 times more

160   likely to also be SB in the body, and this sex bias was often, but not always, in the same

161   direction (Fig 3C, Supplementary Fig S2).

163   **Table 1:** Sex bias in the head and body of six *Drosophila* species

Species	Head				Body			
	Genes	MB	FB	UB	Genes	MB	FB	UB
Dmel	7899	814 (10.3)	898 (11.4)	6187 (78.3)	9006	4229 (47.0)	3121 (34.7)	1656 (18.4)
Dsim	7828	370 (4.7)	520 (6.6)	6940 (88.7)	9002	3998 (44.4)	3166 (35.2)	1838 (20.4)
Dsuz	7591	304 (4.0)	562 (7.4)	6725 (88.6)	8936	3715 (41.6)	3035 (34.0)	2186 (24.6)
Dana	7570	490 (6.5)	548 (7.2)	6532 (86.3)	8955	3727 (41.6)	3007 (33.6)	2221 (24.8)
Dsub	8161	32 (0.4)	135 (1.7)	7994 (98.0)	8981	3727 (41.5)	3004 (33.4)	2250 (25.1)
Dimm	7715	212 (2.7)	565 (7.3)	6938 (89.9)	8938	3527 (39.5)	2896 (32.4)	2515 (28.1)

The number of analyzed genes, MB, FB, and UB genes in the head and body of each species

are shown. Percentages of analyzed genes rounded to the nearest decimal are shown in

parentheses.

Fig 3: Sex bias within and among species. A,B) Pairwise overlap in sex bias categories among

species in the A) head and B) body. Shown are the proportion of overlapping genes that had

conserved SB (Cons SB; white) or conserved UB expression (Cons UB; dark blue), were SB in

only one of the examined species (SB 1 species; light blue), or switched sex bias (Switch;

gray) between *Dmel* (*m*), *Dsim* (*si*), *Dsuz* (*su*), *Dana* (*a*), *Dsub* (*sb*), and *Dimm* (*im*). C)

Proportion of overlapping genes that had conserved SB or conserved UB expression, were SB

in only the body (Gain SB B; light blue) or head (Gain SB H; medium blue), or switched sex

bias between the head and body are shown. D) Spearman's ρ correlations between τ and

magnitude of sex bias in the head and body for MB, FB, and UB genes. Significance was

assessed with a Mann-Whitney *U* test. ***$P < 0.005$. Non-significant comparisons not

shown.

The magnitude of sex bias was significantly higher in the body than the head for all

species for both MB and FB genes (*t*-test; BH corrected $P < 1.5 \times 10^{-15}$ for all; Supplementary

Fig S3). Within the head, the degree of sex bias was higher for FB than for MB genes, while in

the body, the opposite pattern was detected, which was significant for all comparisons in all

species (*t*-test; BH corrected $P < 4.5 \times 10^{-7}$ for all; Supplementary Fig S3), except *Dana* and

Dsub heads (BH corrected $P = 1$). We also detected significant differences in the degree of

sex bias within the same body part and sex bias category (MB or FB) among species,

especially in the head, for many but not all comparisons (Supplementary Fig S3). Despite

these differences in the magnitude of sex bias, overall expression divergence (as measured

by 1 - ρ) between sexes was lowest in the head within species (0.8–1.3%; Supplementary

Table 3), while divergence in the body was fairly similar to divergence between body parts

regardless of sex (27.7–36.5% versus 25.2–43.6%%; Supplementary Tables 3–4). Divergence

between species within the same body part and sex was higher than in the head but lower

than in the body between sexes within species (1.8–12.0%; Supplementary Tables 3–4).

Thus, we detected differences in the magnitude of expression bias based on the type of sex

bias, the body part, and the species, but overall expression levels showed the strongest

correlation (between sexes in the head) and weakest correlation (between the examined

body parts in either sex) within species.

Tissue specificity and expression level are associated with differences in sex bias

According to FlyAtlas2 data (Krause et al. 2022; Supplementary Table S5), genes we

identified as MB in the head or body were often most highly expressed in testes or a somatic

tissue, while FB genes were most highly expressed in the ovary or a somatic tissue. To better

understand the relationship between expression level, breadth, and sex bias in our dataset,

we first examined the overall expression level of SB and UB genes in the head and body of

each species. We detected significant effects of both species and sex bias category on the

expression level in both body parts (ANOVA; $P < 1.5 \times 10^{-6}$ for all; Supplementary Fig S4). In

both the head and body, FB genes were the most highly expressed, followed by MB genes,

and then UB genes, except in *Dsub* heads, where MB genes were the most lowly expressed

(Supplementary Fig S4). Thus, differences in sex bias were associated with differences in

overall gene expression levels.

Next, we used the *Dmel* FlyAtlas2 adult tissue expression data (Krause et al. 2022) to

calculate the tissue specificity index, τ, which ranges from 0 to 1 with values closer to 1

indicating higher tissue specificity, for each gene. When we compared tissue specificity

among SB and UB genes in each species and body part, we detected a significant effect of

sex bias category in both body parts as well as an effect of species in the head (ANOVA; $P <$

1.6×10^{-10} for all; Supplementary Fig S5). Tissue specificity was highest for MB genes in both

the head and body for all species (Supplementary Fig S5). In the head, τ was mostly similar

between FB and UB genes, while in the body, τ was lower in FB than in UB genes

(Supplementary Fig S5). Tissue specificity was significantly positively correlated with the

magnitude of sex bias in both body parts in all sex bias categories and all examined species

(Spearman's ρ; $P < 0.003$ for all) except MB genes in the head of *Dsub* ($P = 0.2483$). Over all

species, this correlation between the degree of sex bias and τ was highest for MB genes in

both body parts (Fig 3D), which was significantly higher than UB genes in both body parts

and FB genes in the body (Mann-Whitney *U* test; $P < 0.005$ for all), while the correlation for

FB genes was only significantly higher than UB genes in the head ($P < 0.005$; Fig 3). Thus,

tissue specificity was positively associated with sex bias, with the strongest correlation in MB

genes, which were also the most tissue specific. This correlation was particularly strong in

the body, which might be driven at least in part by the presence of the testes, which is the

tissue with the highest proportion of MB genes (Parisi et al. 2004).

Patterns of sex bias turnover among species differ between the heads and body

In order to better understand how sex bias is conserved among species, we identified

genes that were SB or UB in all 6 species. In the head, we identified 4,363 (53.5–57.6%)

genes with conserved UB expression in all species, but only 1 (< 0.02%) and 35 (< 0.5%)

genes as having conserved MB or FB expression, respectively. On the other hand, we

identified only 119 (1.3%) genes with UB expression in all species in the body, but 1,638

(18.2–18.3%) and 1,949 (21.6–21.8%) genes with conserved FB and MB expression, respectively. In order to understand how turnover in sex bias occurs among species, we identified lineage-specific gains or losses of FB or MB that occurred on only one branch of the phylogeny (Fig 1). In the head, we detected many more sex bias gains than losses for all species except *Dsub* (Fig 1A; 1,629 total lineage-specific gains versus 135 losses, 103 of which were in *Dsub*); while, in the body we detected more sex bias losses than gains (1,227 losses versus 663 gains; Fig 1B), although the pattern was not as extreme as in the head. Across the entire phylogeny, only 171 genes switched sex bias on a specific lineage, all of which occurred in the body and the majority of which (64.3%) switched from FB to MB (Supplementary Table S6).

To better understand how sex bias changed among species, we made pairwise comparisons of sex bias in all examined genes between species for each body part (Fig 3A–B, Supplementary Figs S6–S7). In the head, genes that were SB in one species were usually UB in the other species (10.2–22.7% of genes), but if a gene was SB in both species, it almost always shared the same sex bias between species (1.0–7.9% of genes), with only 0–9 genes switching their sex bias (i.e. from MB to FB or vice versa) between species (0–0.12% of genes; Fig 3A,B, Supplementary Fig S6). In the body, SB genes were often shared between species, with the majority showing sex bias towards the same sex (55.0–70.2% of genes), but there were also many SB genes that were UB in the other species (18.2–29.9% of genes; Supplementary Figs S7, S8) as well as a small proportion (1.4–4.5%) that switched sex bias between species. However, we should note that because the body is a composite of multiple tissues, it is possible that some of these switches represent opposing changes in different tissues rather than a switch in the same tissue. Thus, we detected a relatively large amount of sex bias turnover in both body parts among species, which occurred in a mostly species-specific manner in the head (Figs 1–2). Moreover, when turnover occurred in the head, a gene's expression switched between being UB and biased towards a single sex, but never between the two sexes.

Sex bias is most often gained or lost via correlated expression changes in both sexes, especially in the head

In order to better understand how expression changes during the gain or loss of sex bias, we examined how expression in a focal species changed relative to the inferred ancestral expression within each sex (see Materials and Methods) for species-specific gains

or losses of SB expression. In the head, the majority of expression changes occurred in the same direction (either up-regulation or down-regulation) in both sexes (75.6% of all sex bias gains and losses; Fig 4), and this pattern was even stronger for genes with large expression changes (log$_2$ fold-change $\geq$ 1), 95.2% of which had concordant expression changes between the sexes (versus 74.3% of genes with small expression changes; Supplementary Fig S8). Although concordant expression changes were also more common than opposing changes in the body (54.3% versus 45.7%), the direction of expression changes in males versus females was more evenly distributed compared to the head (P < 2.2 x 10^{-16}; χ^2 test; Fig 4). Genes with large expression changes (log$_2$ fold-change $\geq$ 2) were more similar to patterns in the head, with 66.7% of genes showing concordant expression changes between sexes (versus 53.3% of genes with small expression changes; Supplementary Fig S8). In keeping with the observation that expression changes were often concordant between sexes, expression changes associated with SB gains were significantly positively correlated in the head and body in all species (Spearman's ρ; P < 0.05 for all) except for MB gains in the *Dsub* head and FB gains in the *Dmel* and *Dsim* body (Supplementary Figs S9–S10). This correlation in gene expression changes between the sexes was significantly higher in the head than in the body (paired Mann-Whitney U test; P = 0.0003). However, this pattern was less apparent for SB losses, for which this correlation was sometimes negative and only significant in half of the comparisons (Supplementary Figs S9–S10), although in the head part of this observation might be at least partially explained by the low number of SB losses in each species. Indeed, correlations in gene expression changes between the sexes were significantly higher for SB gains than for losses in both the head and the body (P < 0.05 for all; Fig 4).

Fig 4: Expression changes underlying species-specific gains and losses of SB gene expression.

As an example, scatterplots depicting the change between *Dmel* and ancestral expression in

males versus females for genes with FB gains in the A) head and D) body are shown. All other

scatterplots can be found in Supplementary Figs S9–S10. A,B,D,E) Genes with concordant

(con) changes in expression between the sexes are located in quadrants ii and iii; while

genes with opposing (opp) expression changes between the sexes are located in quadrants i

and iv. B, E) Total genes in each quadrant for the entire phylogeny in the B) head and E) body

are shown. In parentheses, the proportion of total gene expression changes are shown. C, F)

Spearman's ρ correlations between male and female expression changes within each species

for SB gains and losses in the C) head and F) body. Significance was assessed with a Mann-

Whitney *U* test. ***$P < 0.005$, **$P < 0.01$, *$P < 0.05$. Non-significant comparisons not shown.

Next, we examined how the size of expression changes and ancestral sex bias

affected the distribution of expression changes during SB loss or gain. For the majority of

large gene expression changes (log$_2$ fold-change ≥ 1 in the head or ≥ 2 in the body; 56.7–

75.2% of genes), gain or loss of sex bias occurred via a down-regulation in both sexes with

only 10–20% of genes up-regulated (Supplementary Fig S8). In contrast, up- or down-

regulation in both sexes was more evenly distributed for genes with small expression

changes (32.3–36.5% and 21.0–37.8% of genes for head and body, respectively;

Supplementary Fig S8). Due to the lack of SB reversals and the low number of SB losses for

most species in the head, it is difficult to determine if the ancestral sex bias of a gene

affected the distribution of expression changes, although MB and FB losses tended to

overlap with no clear separation based on the ancestral sex bias (Supplementary Fig S9). We

observed a similar pattern for MB and FB gains in the body, with no clear clustering based on

whether sex bias was gained from an UB or SB gene (Supplementary Fig S10). On the other

hand, FB and MB losses in the body were clearly separated (Supplementary Fig S10), with

genes that lost female bias tending to show down-regulation in females and/or up-

regulation in males and the opposite for MB losses. Thus, a gain or loss of sex bias most

often involved an expression change in the same direction (up-regulation or down-

regulation) in both sexes, but of greater magnitude in one sex than the other, with larger

expression changes tending to be associated with down-regulated expression in both sexes.

Overall, reversal-of-expression changes were less common but were almost as frequent as

concordant expression changes in the body.

Contrasting distributions of X versus autosomal SB genes in the head and body

Previous studies found that, in whole flies and gonads, FB genes were

overrepresented on the X chromosome, while MB genes were underrepresented (Parisi et

al. 2003; Ranz et al. 2003; Sturgill et al. 2007). In contrast, in the head and brain, genes with

SB expression, and especially MB expression, were enriched on the X chromosome (Catalán

et al. 2012; Huylmans and Parsch 2015; Khodursky et al. 2020). We detected similar patterns

across the species included in our study, with the proportion of X-linked genes depending

upon the body part and, in some cases, the species in which SB gene expression was

detected (Fig 5; Supplementary SX). In the head, SB genes were generally over-represented

on the X chromosome, but whether this enrichment occurred for MB or FB genes depended

upon the examined species, and in some cases (e.g. FB genes in *Dsuz* or MB genes in *Dana*)

there was a slight underrepresentation of SB genes on the X (Fig 5A). Interestingly, in *Dsub*

MB genes were highly enriched on the X chromosome but highly underrepresented on the

autosomes (Fig 5A). In the body, FB genes were enriched, while MB genes were

underrepresented on the X chromosome for all species (Fig 5B). Reciprocally, MB genes were

enriched while FB genes were underrepresented on the autosomes for all species except

Dana (Fig 5B). Thus, SB genes displayed unique chromosomal distributions depending on the

examined body part.

Fig 5: Chromosomal distribution of SB genes in the A) head and B) body. Shown are the

ratios of observed (obs) to expected (exp) number of genes in each SB category on the X

versus the autosomes based on sex bias category and the number of analyzed on the X

versus the autosomes. Significances was assessed with a χ^2 test. ***$P < 0.005$, **$P < 0.01$, *P

< 0.05, marginally non-significant: .$P < 0.1$. Non-significant comparisons not shown.

Signs of positive selection at the gene expression level are pervasive in SB genes

A previous study in the brain of three *melanogaster* subgroup species found that SB

genes, and MB genes in particular, are enriched for signatures of directional selection

(Khodursky et al. 2020). In order to test for signs of putative positive selection at the gene

expression level, we calculated Δx (see Materials and Methods), which acts similarly to a

McDonald-Kreitman (MK) test (McDonald and Kreitman 1991), in both sexes for all species

relative to the outgroup species *Dimm*. Briefly, in cases where -1 < Δx > 1, divergence of

expression in the focal species from *Dimm* exceeds expression polymorphism within that

species, indicating that positive selection may have occurred. We detected putative signs of

positive selection in 6.3–76.5% of genes depending upon the sex, species, and sex bias

category (Supplementary Table S7; Supplementary Fig S11), with the proportion of

putatively positively selected genes higher for both FB and MB than for UB genes for all

comparisons. Indeed, we detected significantly more FB and/or MB genes as putatively

positively selected than expected in both sexes in the head for all species (Fig 6A; χ^2 test; $P <$

0.05), except in *Dana* females and *Dsub* males, which were marginally non-significant ($P <$

0.09). Putatively positively selected MB or FB genes were similarly enriched to a lesser

extent in one or both sexes in all species in the body, although we detected fewer FB genes

as putatively positively selected than expected in *Dsuz* females (Fig 6B). This enrichment of

putatively positively selected MB and/or FB genes occurred on both the X and the

autosomes, although in some cases we also detected fewer SB genes than expected both on

the X and the autosomes, especially in the body (Supplementary Fig S12). Interestingly, SB

genes were not always enriched for putative positive selection in the sex in which they were

detected as SB (Fig 6A, B). Indeed, SB genes with signs of positive selection were enriched in

the opposite sex (i.e. the sex with lower expression) in at least one sex in all species in both

the head and body, although this enrichment was marginally non-significant in females in

the *Dsub* body (Fig 6A, B). When we compared Δx between males and females, we detected

significant differences in the magnitude of Δx for the majority of sex bias categories and

species in both head and body (80% for both; Fig 6C, D), with the absolute value of Δx most

often shifted higher in males, especially in the body (75 and 92% of significant shifts in the

head and body, respectively), and the mean absolute value in the significantly higher sex

usually close to or above 1 (88% of significant shifts; Fig 6C, D; Supplementary Fig S11).

Despite this shift in magnitude, Δx was significantly positively correlated between the sexes

in all sex bias categories and species (Spearman's ρ; $P <2.2 \times 10^{-16}$ for all; Supplementary

Table S8), except *Dsub* MB genes in the head ($P = 0.153$), and these positive correlations

were significantly stronger in the body than in the head (mean of 0.827 vs 0.585 in the body

and head, respectively; paired Mann-Whitney U test; $P = 5.37 \times 10^{-3}$). Thus, we detected an

enrichment of putatively positively selected genes among SB genes in both the head and

body, and this enrichment often occurred in the sex with lower expression, mainly involving

the male expression of FB genes. Moreover, the overall magnitude of Δx was significantly

higher in males in comparison to females, especially in the body.

Fig 6: Signs of putative positive selection at the gene expression level. Shown are the ratios of observed (obs) to expected (exp) number of genes under putative positive selection (-1 < Δx > 1) in each SB category and sex in the A) head and B) body. C, D) Shown is the magnitude of Δx values for FB, MB and UB genes in females and males in the C) head and D) body. The dashed gray line represents the cut-off above which a gene shows signs of putative positive selection (absolute value of Δx > 1). For better visualization, absolute Δx values above 5 are not shown. A, B) Significance was assessed with a χ² test. C, D) Significant differences between sexes were assessed with a *t*-test. ***$P < 0.005$, **$P < 0.01$, *$P < 0.05$, marginally non-significant: .$P < 0.1$. Non-significant comparisons not shown.

Discussion

We characterized SB gene expression in the head and body of 6 *Drosophila* species,

encompassing a broad geographic distribution within species and 2–50 million years of

evolution (Fig 1; Supplementary Table S1). Overall, we detected contrasting patterns of SB

gene expression and turnover between the head and body (Table 1; Figs 1–3; Supplementary

Figs S3, S6–S8), as well as differences in expression level, tissue-specificity, and magnitude of

sex bias depending on the species, sex bias category, body part and/or sex (Supplementary

Figs S3–S5). Our findings that *i*) sex bias was less extensive (Table 1), *ii*) conserved sex bias

was rare while conserved UB expression was extensive (Fig 3A), and *iii*) the least amount of

expression divergence occurred in the head (Supplementary Tables 3–4) suggest that gene

expression is more constrained in the head than in the body. Indeed, sex bias in the head

was largely gained in a species-specific manner (Fig 1). Unlike in the body, where sex bias

sometimes switched between species, these gains almost exclusively occurred in genes that

were either UB or biased towards the same sex in the head of other species (99.88–100% of

examined genes; Fig 3A,B, Supplementary Figs S6–S7), suggesting that there is greater

constraint against sex bias reversals in the head throughout the *Drosophila* genus.

Moreover, SB genes in the head were usually also SB in the body irrespective of sex (Fig 3;

Supplementary Fig S2), suggesting that sex bias is more likely to arise in the head when the

underlying regulatory mechanisms for sex bias already exist for that gene. Taken together,

these findings suggest that the gain of sex bias is much more constrained in the head than in

the body. This pattern is likely driven, at least in part, by expression in the brain, where

expression is known to be more constrained than in other tissues (Khaitovich et al. 2005,

2006). In keeping with our findings, Khodursky et al. (2020) also found that expression was

constrained in the brain with SB expression being largely species-specific; however, they also

detected a much larger proportion of switching in sex bias in their dataset. This discrepancy

between studies is likely due to their more limited sampling within species, which included

either a limited number of isofemale lines (N = 2) or lines derived from a single population (N

= 6). On the other hand, our dataset included more lines (N = 5–8) per species, which were

derived from multiple, geographically diverse populations. Thus, our samples represented a

broader overview of within species variation and likely "averaged out" potential population-

or line-specific sex bias. Consistent with this interpretation, a recent study found that

somatic SB genes are more variable than UB genes among individuals within sub-species of

mice (Xie et al. 2025). This study also detected a larger proportion of sex bias reversals

among subspecies, but this discrepancy with our study might be due to their shorter

evolutionary timescale (Xi et al. 2025).

Interestingly, lineage-specific gains or losses of sex bias most often occurred via

concordant expression changes in both sexes, but with a greater change in one sex than the

other (75.6% in the head and 54.3% in the body; Fig 4, Supplementary Figs S9–S10), and this

pattern was even stronger for sex bias gains (Fig 4C,F) as well as genes with large expression

changes (95.2% in the head and 66.7% in the body; Supplementary Fig S8), the majority of

which occurred via a down-regulation of expression in both sexes. This concordance

suggests that these lineage-specific sex bias gains are not a way to resolve sexual

antagonism, as in such a case we would expect up-regulated expression in one sex and

down-regulated expression in the other (quadrants i and iv in Fig 4A,B,D,E). Such changes

represented the minority of SB gains in our data (24.4% in the head and 45.7% in the body),

especially in the head and among genes with large expression changes (4.8% in the head and

33.3% in the body). Thus, in general, the presence of SB expression is likely to be a poor

proxy for ongoing or resolved sexual antagonism in *Drosophila*. In keeping with our findings,

a recent study in *Dmel* found that sexually antagonistic effects on starvation resistance could

not be explained by SB gene expression (Chen et al. 2025). Moreover, another study of 5

mammalian species found that only 27% of lineage-specific sex bias changes could be

associated with turnover in SB transcription factor motifs (Naqvi et al. 2019), suggesting that

in such cases the gain of sex-specific regulation may be driving the gain or loss of sex bias.

Thus, while opposing sex-specific regulatory changes contribute to the gain or loss of

lineage-specific SB gene expression, our results suggest that the majority of lineage-specific

SB gains occur via concerted changes in expression with a larger magnitude of change in one

sex than the other, suggesting that the underlying regulatory architecture is largely shared

between the sexes. Indeed, a previous study in humans found that sex differences in the

genetic regulation of gene expression are not very common (Oliva et al. 2020), while another

study of the brain of three *melanogaster* group species found that male and female

expression were positively correlated relative to ancestral expression (Khodursky et al.

2020). We similarly detected a positive correlation in lineage-specific SB gains and losses, but

the strength of this correlation was stronger for SB gains than losses (Fig 4), underscoring

that many species-specific SB gains are likely the result of differences in regulation that is

shared between the sexes.

We also detected contrasting patterns in the chromosomal distribution of SB genes

between the head and body (Fig 5). In the body, we detected a consistent enrichment of FB

and a paucity of MB genes on the X chromosome (Fig 5B), which is consistent with the

"feminization" and concurrent "de-masculinization" of the X that has been reported in

previous studies (e.g. Parisi et al. 2003; Ranz et al. 2003; Sturgill et al. 2007). On the other

hand, in the head, we often detected an enrichment of SB genes (MB, FB, or both) on the X

chromosome (Fig 5A). Previous work in *melanogaster* group species found that genes with

SB expression, and particularly MB expression, in the brain were enriched on the X

chromosome (Catalán et al. 2012; Khodursky et al. 2020) and that this signal from the brain

may contribute to a detectable enrichment of X-linked SB genes in studies of whole heads

(Huylmans and Parsch 2015). Thus, expression in the brain likely contributed to the

enrichment of X-linked SB genes that we detected in the head (Fig 5A).

SB genes, especially MB genes, are often more tissue-specific than UB genes (Meisel

2011; Dean et al. 2016; Khodursky et al. 2020). In keeping with previous findings, we found

that tissue specificity was highest in MB genes (Supplementary Fig S5). Moreover, tissue

specificity was positively associated with the magnitude of sex bias, and this pattern was

strongest in MB genes, especially in the body (Fig 3D). Together, these findings suggest that

increased tissue specificity may facilitate larger changes in SB expression, especially for MB

genes, which is in keeping with the idea that increased tissue specificity may be a driving

force in the evolution of SB gene expression, and MB gene expression in particular (Miesel

2011; Dean et al. 2016). However, we should note that in this study we focused on

expression in body parts, which are a composite of multiple tissues. While the head is

composed of relatively few tissues, the body is composed of many, including reproductive

tissues; therefore, it is possible we may have missed some smaller or tissue-specific changes

in SB gene expression.

When we examined Δx as a measure of putative positive selection at the gene

expression level (where -1 < Δx > 1 is indicative of positive selection), we found that *i*) the

absolute value of Δx was most often shifted higher in males compared to females, with the

magnitude of male Δx usually close to or above 1 (Fig 6; Supplementary Table S8), *ii*) SB

genes were often enriched for signs of positive selection in the opposite sex (i.e. the more

lowly expressed sex), especially FB genes for male expression (Fig 6), and *iii*) Δx tended to

shift in the same direction in both sexes (Supplementary Table S8). At the same time,

changes in gene expression relative to ancestral expression during sex bias gains tended to

be positively correlated between the sexes (Supplementary Figs S9–S10), as did gene

expression between the sexes within body parts (Supplementary Tables S3–S4). Taken

together, these findings suggest that lineage-specific sex bias changes may often be

facilitated by selection acting on expression in the sex with lower expression (i.e. selection

may be occurring for lower expression, often resulting in a net gain of sex bias in the

opposite sex), especially for male expression of FB genes in the body. Indeed, we found that

lineage-specific sex bias changes were most often concordant between the sexes, especially

in the head, for SB gains, and for genes with large expression changes (Fig 4; Supplementary

Fig 8), the majority of which occurred via a down-regulation of expression in both sexes.

Thus, our results suggest that most lineage-specific sex bias gains may be the result of

selection acting on the asymmetrical down- (or up-) regulation of expression in both sexes

via shared regulatory architecture, and this mechanism may be particularly important when

expression is highly constrained, such as in the head or brain, or when there are large overall

expression changes. This finding is rather counterintuitive, as SB gene expression is known to

evolve rapidly (Assis et al. 2012; Whittle and Johannesson 2013; Harrison et al. 2015; Yang et

al. 2016; Tosto et al. 2023; Xie et al. 2025), and theory predicts that shared genetic

architecture should slow down the evolution of sexual dimorphism (Lande 1980) given the

constraints that the shared genome imposes on gene expression (Griffin et al. 2013;

Khodursky et al. 2020). However, a recent study using mouse and human data detected

overlapping variances of SB gene expression between the sexes in somatic tissues,

suggesting that patterns and evolutionary dynamics of SB genes are likely more nuanced

than previously thought. Therefore, in cases where sex bias is beneficial, but constraint on

gene expression is relatively high, working within the constraint of a shared genome, rather

than around it, may provide an alternative, more common route to the establishment of SB

gene expression.

Materials and Methods

***Drosophila* strains and RNA-seq**

All *Drosophila* species were maintained at 21°C with a 14 h light:10 h dark cycle on

standard cornmeal–yeast–molasses medium. For all species, strains were established as

isofemale lines and maintained by transferring 20–50 flies per generation for at least 10

generations. For all of the *Dmel* and some of the *Dsim* strains, additional inbreeding was

done by performing single sib-pair crosses for 5–10 generations. Expression was examined in

5–8 strains of 6 *Drosophila* species (Fig 1). To provide a broad geographical overview of

expression within each species, the *Drosophila* strains were sampled from locations across

Europe, Asia, and Africa and were either collected by ourselves or kindly provided by others

(see Supplementary Table S1 for details on sampling locations and provenance). Briefly, we

included strains from at least two populations per species and included strains from the

inferred ancestral region for *Dmel*, *Dsim*, *Dana*, and *Dsuz*. For each strain, groups of 30–120

heads and 8–80 bodies of 3–6-day-old flies were dissected in cold 1X PBS and frozen in RNA-

DNA shield (Zymo research; Freiburg, Germany) at -70°C until RNA extraction. RNA was

extracted using a Qiagen RNeasy Mini or Plus Universal Kit (Qiagen; Hilden, Germany). Poly-

A selection, fragmentation, reverse transcription, library construction, and high- throughput

RNA-seq was performed by Novogene (Martinsried, Germany) using the Illumina NextSeq

550 platform (Illumina; San Diego, CA) with 150-bp paired-end reads. We sequenced one

biological replicate per strain, sex, and body part, resulting in 35 libraries for each sample

type (140 libraries in total). Library size ranged from 22,874,617 to 34,733,318 million

paired-end reads, 43.29–78.74% of which could be mapped (Supplementary Table S9; see

description of read mapping below).

Read mapping and estimation of gene expression levels

In order to prevent potential bias due to lower quality annotations in some species

(for example, no full transcript information was available for *Dimm*; whereas, in *Dmel* UTRs

are better annotated than in the other, less studied species), we mapped RNA-seq reads to

CDS sequences (excluding UTRs) for each species. Reference CDS sequences were

downloaded (April 25, 2024) from the following sources: NCBI (*Dsub* genome assembly

UCBerk_Dsub_1.0, *Dana* genome assembly ASM1763931v2, D*suz* Genome assembly

LBDM_Dsuz_2.1.pri, and *D simulans* genome assembly Prin_Dsim_3.1), FlyBase (version

6.57; Öztürk-Çolak et al. 2024; *Dmel*), and Li et al. (2022; *Dimm*). Links to the downloaded

references are provided in Supplementary Table S10.

RNA-seq reads were mapped to the respective species reference using NextGenMap

(Sedlazeck et al. 2013) in paired-end mode. Read pairs mapping to different genes were

discarded, while read pairs matching more than one isoform of a gene were randomly

assigned to one of the isoforms of that gene. For downstream analyses, we analyzed the

sum of read counts across all of a gene's isoforms (across all annotated exons), i.e. on the

individual gene-level. In order to make direct comparisons of gene expression among species

and sample types, we used our gene counts to calculate levels of gene expression as transcripts per million (TPM; Wagner et al. 2013) for all genes in each library. For TPM calculations, CDS lengths were extracted from the *Dmel* reference files or calculated from the reference files for all other species using the faidx function in *samtools* (Danecek et al. 2021). In cases of multiple isoforms, the longest CDS length was used.

Identification of 1:1 orthologs

In order to keep the datasets for each species comparable, we restricted our analyses to 1:1 orthologs present in all species. To identify 1:1 orthologs, we downloaded (April 29, 2024) pairwise lists of orthologs between *Dmel* and all other species from Li et al. (2022; for *Dimm*) or DrosOMA (Thiébaut et al. 2024; for all other species; Supplementary Table S10), as well as the full pairwise orthologs list from DrosOMA (Thiébaut et al. 2024; Supplementary Table S10). First, all non-1:1 orthologs were removed from each *Dmel* ortholog list and the FlyBase batch download and/or ID validator tools (Öztürk-Çolak et al. 2024) were used to obtain FBgn numbers for *Dmel* IDs from each list. Non-gene or non-coding matches were removed and in cases for which submitted IDs yielded multiple matches, only the FBgn number that exactly corresponded the original ID was retained. For *Dmel* IDs with multiple matches but no exact match or duplicate FBgn numbers due to ID changes/gene re-naming, we used a combination of FlyBase (Öztürk-Çolak et al. 2024), DrosOMA (Thiébaut et al. 2024), and UniProt (Ahmad et al. 2025) to validate the IDs. IDs that did not match between our reference files and our ortholog lists were validated in the same manner. The orthologous *Dmel* sequences corresponding to IDs from the *Dimm* list (Li et al. 2022) that could not be validated using this method were blasted (blastn; Altschul et al. 1990) to identify the current *Dmel* gene, and their isoform was verified in DrosOMA. All IDs that could not be verified were removed from our 1:1 ortholog list as were IDs that were no longer classified as genes by NCBI, not included in our reference files, and IDs that had been incorporated into a gene model already represented in our list (i.e. duplicate IDs). Next, we used the full pairwise orthologs list from DrosOMA (Thiébaut et al. 2024; Supplementary Table S10) to remove genes that were not 1:1 orthologs between the non-*melanogaster* species. Orthologs that could not be confirmed as belonging to the same, known OMA group were also removed as were orthologs belonging to the same OMA group (i.e. OMA groups that appeared more than once in our list). After these filtering steps, we obtained a list of 9,016 1:1 orthologs for use in our expression analyses.

Expression analyses

We identified sex-biased genes (Supplementary Data S1) using gene counts in the 9,016 1:1 orthologs and a negative binomial test as implemented in DESeq2 (Love et al. 2014) in R (R core team 2022) with a 5% false discovery rate (FDR). Head and body samples for each species were analyzed separately using a one-factor model design with two levels corresponding to each sex. For PCAs within each species, we implemented a 2-factor (sex and body part) model design with two levels for each factor (male versus female and head versus body). For the PCA including all samples, we similarly implemented a 3-factor design (sex, body part, and species), with the additional third factor consisting of 6 levels corresponding to the 6 examined species. Gene expression divergence between 2 sample types was calculated as $1 -$ Spearman's ρ between the mean TPM of the 2. Significant differences in divergence or magnitude of sex bias between samples were assessed with a *t*-test with a Benjamini-Hochberg correction (Benjamini and Hochberg 1995) for the number of comparisons made per sample. We tested for an effect of sex bias category and species on overall expression levels using the mean of the $\log_2$ TPM of each gene in each species and body part and a type II ANOVA.

To identify lineage-specific gains or losses of sex bias, we employed a conservative approach in which a terminal species (or all species of a terminal clade) was only considered to have a change in sex bias if all species on that lineage shared the same sex bias, and this sex bias differed from that of all other outgroup species, which were also all required to share the same sex bias. Due to the lack of an outgroup, we could not polarize sex bias changes as gains or losses on the lineage leading to *Dimm*; therefore, we categorized the gain or loss of expression relative to the other analyzed species. In order to compare how expression changed in a particular species relative to the inferred ancestral expression, we calculated the ancestral expression for each gene, sex and body part for all internal nodes of our phylogenetic tree using the mean TPM values in each species and a Brownian motion model fitted by maximum likelihood as implemented in the ace function in the R package *ape* (Paradis and Schliep 2019). Changes in expression within each sex, species, and body part in comparison to the inferred ancestral expression were calculated as $\log_2(\text{TPM}_{\text{Focal}}/\text{TPM}_{\text{Anc}})$, where $\text{TPM}_{\text{Focal}}$ was the mean TPM in the focal species for a given body part and sex and TPM_{Anc} was the inferred ancestral TPM at the nearest internal node. We tested for differences in the distribution of the direction (up- or down-regulation) of

expression changes in males versus females using a χ^2 test for the number of genes in

quadrants i–iv (see Fig 4). We further categorized genes as having a large or small expression

change based on a $\log_2$ fold-change threshold of ≥ 2 in the body or ≥ 1 in the head, where

the extent (Table 1) and magnitude (Supplementary Fig S3) of sex bias was lower. To test for

significant correlations between male and female expression changes during changes in sex

bias, we calculated Spearman's ρ between expression changes in males versus female for all

genes that gained or lost sex bias within each species. We then tested for differences in

these correlations between MB or FB gains and SB losses using a Mann-Whitney U test or a

paired Mann-Whitney U test to test for differences between body parts.

Chromosomal location and calculation of τ and Δx

Chromosomal locations of all protein-coding genes for *Dsim*, *Dsuz*, *Dana*, and *Dsub*

were downloaded (July 11, 2025) from NCBI. For *Dsuz* the CBGP_Dsuzu_IsoJpt1.0 assembly,

which is chromosome-level, was used to determine the chromosomal location of analyzed

genes. *Dmel* chromosomal locations were extracted from the reference CDS file. Because a

chromosome-level assembly was not available for *Dimm*, this species was excluded from all

chromosomal location analyses. For all comparisons between X and autosomal genes, we

compared genes that could be assigned to the X chromosome (also known as Muller

element "A") to all genes that could be assigned to any autosome within each species.

Separately for the X and the autosomes, we tested for differences in the number of observed

genes in each sex bias category versus the expected number of genes in each category based

on the overall proportion of genes in each category for that species using a χ^2 test.

In order to examine tissue specificity, we downloaded the FlyAtlas2 FPKM data

(Krause et al 2022) for all adult *Dmel* tissues, excluding fat body, spermatheca, and heart,

which were sequenced using a different chemistry from the other tissues, from

https://motif.mvls.gla.ac.uk/downloads/FlyAtlas2_gene_data_2025.xlsx (April 24, 2025). We

then calculated the tissue-specificity index τ (Yanai et al 2005) for each gene (Supplementary

Data S1) and identified the tissue with the highest expression. We tested for an effect of sex

bias category and species on τ in each body part using a type II ANOVA. In order to test for

correlations between sex bias and tissue specificity, we calculated Spearman's ρ between τ

and the magnitude of sex bias for each sex bias category in each body part and species. We

tested for differences in these correlations among sex bias categories using a Mann-Whitney

U test.

In order to identify putative signs of positive selection at the gene expression level,

for each species and body part we calculated Δx (Supplementary Data S1) similar to

Moghadam et al. (2012) within each sex as

$$x = \frac{d}{fr}, (1)$$

673   where, *d* is the divergence in gene expression between a focal species and *Dimm*, *r* is the

674   range of expression levels for the focal species, and *f* is a modifier correcting for differences

675   in sample size among species. To calculate Δx, we used the $\log_2$ TPM as a measure of

676   expression for each gene in each fly strain and calculated the divergence (*d*) in expression

677   between the focal species expression (X_{Focal}) and *Dimm* expression (X_{Imm}) as

$$d = \frac{mean(X_{Focal}) - mean(X_{Imm})}{mean(X_{Focal})}, (2)$$

679   the expression range (*r*) as

$$r = \frac{max(X_{Focal}) - min(X_{Imm})}{mean(X_{Focal})}, (3)$$

681   and the correction factor (*f*) as

$$f = \sqrt{\frac{mN - 1}{mN - m}}, (4)$$

where *m* is the number of strains for the focal species and *N* is the most common number of

strains (for our data, *N* = 5). For all genes, and for the X chromosome and autosomes

separately, we tested for differences in the number of observed putatively positively

selected genes in each sex bias category and the number of putatively selected genes

expected based on the proportion of genes in each sex bias category for each sex, species

and body part using a χ^2 test. For each body part, species, and sex bias category, we tested

for differences in the magnitude of Δx between males and females using a *t*-test. We also

tested for correlations in Δx between the sexes for each body part, species, and sex bias

category using Spearman's ρ. We tested for differences in the strengths of these correlations

between the body and the head using a paired Mann-Whitney *U* test

Data Availability

The RNA-seq data that support the findings of this study are publicly available from the

European Nucleotide Archive (ENA) at EMBL-EBI under Project accession number

XXXXXXXXXXX. All other relevant data are within the manuscript and its Supplementary

Material files.

Acknowledgements

We thank Hilde Lainer for excellent technical assistance in the laboratory. We are grateful to members of the European *Drosophila* Populations Genomics Consortium (DrosEU) and the following people for providing the fly lines used in this study: Jessica Abbott, Nicolas Gompel, Josefa González, Sonja Grath, Katja Hoedjes, Mihailo Jelić, Maaria Kankare, Viola Nolte, Darren Obbard, Marta Pascual, John Pool, Christian Schlötterer, Mads Fristrup Schou, Marina Stamenković, Fabian Staubach, Cristina Vieira, Jorge Vieira, and Ricardo Wilches. This study was supported by the Deutsche Forschungsgemeinschaft priority program "The genomic basis of evolutionary innovations" (SPP2349; Project No. 503272152 awarded to JP).

References

1. Ahmad S, Jose da Costa Gonzales L, Bowler-Barnett EH, Rice DL, Kim M, Wijerathne S, et al. The UniProt website API: facilitating programmatic access to protein knowledge. *Nucleic Acids Res.* 2025;53(W1):W547-W53.

2. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215(3):403-10.

3. Arbeitman MN, Furlong EE, Imam F, Johnson E, Null BH, Baker BS, et al. Gene expression during the life cycle of *Drosophila melanogaster*. Science. 2002;297(5590):2270-5.

4. Assis R, Zhou Q, Bachtrog D. Sex-biased transcriptome evolution in *Drosophila*. Genome Biol Evol. 2012;4(11):1189-200.

5. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*. 1995;57(1):289-300.

6. Buchberger E, Reis M, Lu TH, Posnien N. Cloudy with a Chance of Insights: Context Dependent Gene Regulation and Implications for Evolutionary Studies. *Genes* (Basel). 2019;10(7).

7. Catalán A, Hutter S, Parsch J. Population and sex differences in *Drosophila melanogaster* brain gene expression. *BMC Genomics*. 2012;13:654.

8. Chen J, Liu C, Li W, Clark AG, Wang Z, Lu J. Pervasive sex-dependent effects in the genetic architecture of starvation resistance in. *Sci Adv*. 2025;11(39):eadt3560.

9. Cheng C, Kirkpatrick M. Sex-Specific Selection and Sex-Biased Gene Expression in Humans and Flies. *PLoS Genet*. 2016;12(9):e1006170.

10. Connallon T, Knowles LL. Intergenomic conflict revealed by patterns of sex-biased gene expression. *Trends Genet*. 2005;21(9):495-9.

11. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
SAMtools and BCFtools. Gigascience. 2021;10(2).

12. Dean R, Mank JE. Tissue Specificity and Sex-Specific Regulatory Variation Permit the
Evolution of Sex-Biased Gene Expression. *Am Nat.* 2016;188(3):E74-84.

13. Ellegren H, Parsch J. The evolution of sex-biased genes and sex-biased gene expression.
Nat Rev Genet. 2007;8(9):689-98.

14. Fass SB, Mulvey B, Chase R, Yang W, Selmanovic D, Chaturvedi SM, et al. Relationship
between sex biases in gene expression and sex biases in autism and Alzheimer's disease.
Biology of Sex Differences. 2024;15(1):47.

15. Gibson G, Riley-Berger R, Harshman L, Kopp A, Vacha S, Nuzhdin S, et al. Extensive sex-
specific nonadditivity of gene expression in *Drosophila melanogaster*. *Genetics*.
2004;167(4):1791-9.

16. Grath S, Parsch J. Rate of amino acid substitution is influenced by the degree and
conservation of male-biased transcription over 50 myr of *Drosophila* evolution. *Genome
Biol Evol*. 2012;4(3):346-59.

17. Grath S, Parsch J. Sex-Biased Gene Expression. *Annu Rev Genet*. 2016;50:29-44.

18. Griffin RM, Dean R, Grace JL, Rydén P, Friberg U. The shared genome is a pervasive
constraint on the evolution of sex-biased gene expression. *Mol Biol Evol*. 2013;30(9):2168-
76.

19. Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, et al. Sexual
selection drives evolution and rapid turnover of male gene expression. *Proc Natl Acad Sci U
S A*. 2015;112(14):4393-8.

20. Hudson RR, Kreitman M, Aguadé M. A test of neutral molecular evolution based on
nucleotide data. *Genetics*. 1987;116(1):153-9.

21. Huylmans AK, Parsch J. Population- and sex-biased gene expression in the excretion
organs of *Drosophila melanogaster*. *G3* (Bethesda). 2014;4(12):2307-15.

22. Huylmans AK, Parsch J. Variation in the X:Autosome Distribution of Male-Biased Genes
among *Drosophila melanogaster* Tissues and Its Relationship with Dosage Compensation.
Genome Biol Evol. 2015;7(7):1960-71.

23. Innocenti P, Morrow EH. The sexually antagonistic genes of *Drosophila melanogaster*.
PLoS Biol. 2010;8(3):e1000335.

24. Jin W, Riley RM, Wolfinger RD, White KP, Passador-Gurgel G, Gibson G. The
contributions of sex, genotype and age to transcriptional variance in *Drosophila
melanogaster*. Nat Genet. 2001;29(4):389-95.

25. Khaitovich P, Enard W, Lachmann M, Pääbo S. Evolution of primate gene expression.
Nat Rev Genet. 2006;7(9):693-702.

26. Khaitovich P, Hellmann I, Enard W, Nowick K, Leinweber M, Franz H, et al. Parallel
patterns of evolution in the genomes and transcriptomes of humans and chimpanzees.
Science. 2005;309(5742):1850-4.

27. Khodursky S, Svetec N, Durkin SM, Zhao L. The evolution of sex-biased gene expression
in the *Drosophila* brain. *Genome Res*. 2020;30(6):874-84.

28. Kimura M. The neutral theory of molecular evolution. Cambridge: Cambridge University
Press; 1983.

29. King MC, Wilson AC. Evolution at two levels in humans and chimpanzees. *Science*.
1975;188(4184):107-16.

30. Krause SA, Overend G, Dow JAT, Leader DP. FlyAtlas 2 in 2022: enhancements to the
Drosophila melanogaster expression atlas. *Nucleic Acids Research*. 2022;50(D1):D1010-D5.

31. Lande R. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.
Evolution. 1980;34(2):292-305.

32. Li F, Rane RV, Luria V, Xiong Z, Chen J, Li Z, et al. Phylogenomic analyses of the genus
Drosophila reveals genomic signals of climate adaptation. *Mol Ecol Resour*.
2022;22(4):1559-81.

33. Llopart A. The rapid evolution of X-linked male-biased gene expression and the large-X
effect in *Drosophila yakuba*, *D. santomea*, and their hybrids. *Mol Biol Evol*.
2012;29(12):3873-86.

34. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):550.

35. Ma WJ, Veltsos P, Toups MA, Rodrigues N, Sermier R, Jeffries DL, et al. Tissue Specificity
and Dynamics of Sex-Biased Gene Expression in a Common Frog Population with
Differentiated, Yet Homomorphic, Sex Chromosomes. *Genes* (Basel). 2018;9(6).

36. McDonald JH, Kreitman M. Adaptive protein evolution at the Adh locus in *Drosophila*.
Nature. 1991;351(6328):652-4.

37. Meiklejohn CD, Coolon JD, Hartl DL, Wittkopp PJ. The roles of *cis*- and *trans*-regulation
in the evolution of regulatory incompatibilities and sexually dimorphic gene expression.
Genome Res. 2014;24(1):84-95.

38. Meiklejohn CD, Parsch J, Ranz JM, Hartl DL. Rapid evolution of male-biased gene
expression in *Drosophila*. *Proc Natl Acad Sci U S A*. 2003;100(17):9894-9.

39. Meisel RP. Towards a more nuanced understanding of the relationship between sex-
biased gene expression and rates of protein-coding sequence evolution. *Mol Biol Evol*.
2011;28(6):1893-900.

40. Moghadam HK, Pointer MA, Wright AE, Berlin S, Mank JE. W chromosome expression
responds to female-specific selection. *Proc Natl Acad Sci U S A*. 2012;109(21):8207-11.

41. Müller L, Grath S, von Heckel K, Parsch J. Inter- and intraspecific variation in *Drosophila*
genes with sex-biased expression. *Int J Evol Biol*. 2012;2012:963976.

42. Naqvi S, Godfrey AK, Hughes JF, Goodheart ML, Mitchell RN, Page DC. Conservation,
acquisition, and functional impact of sex-biased gene expression in mammals. *Science*.
2019;365(6450).

43. Oliva M, Muñoz-Aguirre M, Kim-Hellmuth S, Wucher V, Gewirtz ADH, Cotter DJ, et al.
The impact of sex on gene expression across human tissues. *Science*. 2020;369(6509).

44. Öztürk-Çolak A, Marygold SJ, Antonazzo G, Attrill H, Goutte-Gattat D, Jenkins VK, et al.
FlyBase: updates to the *Drosophila* genes and genomes database. *Genetics*. 2024;227(1).

45. Pal S, Oliver B, Przytycka TM. Stochastic Modeling of Gene Expression Evolution
Uncovers Tissue- and Sex-Specific Properties of Expression Evolution in the. *J Comput Biol.*
2023;30(1):21-40.

46. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*. 2019;35(3):526-8.

47. Parisi M, Nuttall R, Edwards P, Minor J, Naiman D, Lü J, et al. A survey of ovary-, testis-,
and soma-biased gene expression in *Drosophila melanogaster* adults. *Genome Biol.*
2004;5(6):R40.

48. Parsch J, Ellegren H. The evolutionary causes and consequences of sex-biased gene
expression. *Nat Rev Genet*. 2013;14(2):83-7.

49. R Core Team. R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing 2022.

50. Ranz JM, Castillo-Davis CI, Meiklejohn CD, Hartl DL. Sex-dependent gene expression and
evolution of the *Drosophila* transcriptome. *Science*. 2003;300(5626):1742-5.

51. Ranz JM, Namgyal K, Gibson G, Hartl DL. Anomalies in the expression profile of
interspecific hybrids of *Drosophila melanogaster and Drosophila simulans*. *Genome Res*.
2004;14(3):373-9.

52. Rodríguez-Montes L, Ovchinnikova S, Yuan X, Studer T, Sarropoulos I, Anders S, et al.
Sex-biased gene expression across mammalian organ development and evolution. *Science*.
2023;382(6670):eadf1046.

53. Sedlazeck FJ, Rescheneder P, von Haeseler A. NextGenMap: fast and accurate read
mapping in highly polymorphic genomes. *Bioinformatics*. 2013;29(21):2790-1.

54. Shi L, Zhang Z, Su B. Sex Biased Gene Expression Profiling of Human Brains at Major
Developmental Stages. *Sci Rep*. 2016;6:21181.

55. Sturgill D, Zhang Y, Parisi M, Oliver B. Demasculinization of X chromosomes in the
Drosophila genus. *Nature*. 2007;450(7167):238-41.

56. Tamura K, Subramanian S, Kumar S. Temporal patterns of fruit fly (*Drosophila*)
evolution revealed by mutation clocks. *Mol Biol Evol*. 2004;21(1):36-44.

57. Thiébaut A, Altenhoff AM, Campli G, Glover N, Dessimoz C, Waterhouse RM. DrosOMA:
the *Drosophila* Orthologous Matrix browser. *F1000Res*. 2024;12:936.

58. Tosto NM, Beasley ER, Wong BBM, Mank JE, Flanagan SP. The roles of sexual selection
and sexual conflict in shaping patterns of genome and transcriptome variation. *Nat Ecol*
Evol. 2023;7(7):981-93.

59. Wagner GP, Kin K, Lynch VJ. A model based criterion for gene expression calls using
RNA-seq data. *Theory in Biosciences*. 2013;132(3):159-64.

60. Whittle CA, Johannesson H. Evolutionary dynamics of sex-biased genes in a
hermaphrodite fungus. *Mol Biol Evol*. 2013;30(11):2435-46.

61. Wray GA, Hahn MW, Abouheif E, Balhoff JP, Pizer M, Rockman MV, et al. The Evolution
of Transcriptional Regulation in Eukaryotes. *Molecular Biology and Evolution*.
2003;20(9):1377-419.

62. Wright AE, Fumagalli M, Cooney CR, Bloch NI, Vieira FG, Buechel SD, et al. Male-biased
gene expression resolves sexual conflict through the evolution of sex-specific genetic
architecture. *Evol Lett*. 2018;2(2):52-61.

63. Xie C, Künzel S, Tautz D. Fast evolutionary turnover and overlapping variances of sex-
biased gene expression patterns defy a simple binary sex classification of somatic tissues.
Elife. 2025;13.

64. Yanai I, Benjamin H, Shmoish M, Chalifa-Caspi V, Shklar M, Ophir R, et al. Genome-wide
midrange transcription profiles reveal expression level relationships in human tissue
specification. *Bioinformatics*. 2005;21(5):650-9.

65. Yang L, Zhang Z, He S. Both Male-Biased and Female-Biased Genes Evolve Faster in Fish
Genomes. *Genome Biol Evol*. 2016;8(11):3433-45.

66. Yang X, Schadt EE, Wang S, Wang H, Arnold AP, Ingram-Drake L, et al. Tissue-specific
expression and regulation of sexually dimorphic genes in mice. *Genome Res*.
2006;16(8):995-1004.

67. Zhang Y, Sturgill D, Parisi M, Kumar S, Oliver B. Constraint and turnover in sex-biased
gene expression in the genus *Drosophila*. *Nature*. 2007;450(7167):233-7.
871

Supplementary Material Captions

Supplementary Table S1: Overview of *Drosophila* lines assayed in this study

Supplementary Table S2: Sex biased genes in *D. melanogaster* in a random sub-sampling of

5 lines

Supplementary Table S3: Expression divergence within species

Supplementary Table S4: Expression divergence among species

Supplementary Table S5: Number of sex-biased genes expressed highest in testis, ovary, or a

somatic tissue

Supplementary Table S6: Lineage-specific gains and losses of sex biased gene expression

Supplementary Table S7: Sex-biased genes on the X versus autosomes

Supplementary Table S8: Genes showing signs of putative positive selection at the

expression level

Supplementary Table S9: Library size and mapping efficiency

Supplementary Table S10: Download location of files used for analysis

Supplementary Fig S1: Principal component analysis of gene expression profiles in A) *Dmel*,

B) *Dsim*, C) *Dsuz*, D) *Dana*, E) *Dsub*, and F) *Dimm*. Triangles and diamonds indicate female

and male body, respectively, while circles and squares indicate female and male head,

respectively.

Supplementary Fig S2: Overlap in SB genes between head (H) and body (B). Shown are upset

plots demonstrating the overlap in UB, FB, or MB genes in A) *Dmel* (mel), B) *Dsim* (sim), C)

Dsuz (suz), D) *Dana* (ana), E) *Dsub* (sub), and F) *Dimm* (imm). Horizontal bars represent the

total number of genes in a body part and sex bias category combination. Vertical bars

represent the number of genes in an intersection class. Connected, filled circles underneath

a vertical bar indicate that a body part and sex bias category combination is included in an

intersection class. Blue represents genes UB in one body part and SB in the other, dark gray

represents gene in the same bias category, and light blue indicates genes that switch sex bias

between head and body.

Supplementary Fig S3: Magnitude of sex bias in the A) head and B) body of *D. melanogaster*

(mel), *Dsim* (sim), *Dsuz* (suz), *Dana* (ana), *Dsub* (sub), and *Dimm* (im/imm). Significant

differences between species for MB and FB genes were assessed with a *t*-test. Shown are the

BH-corrected *P*-values in the C) head and D) body. Non-significant *P*-values are shown in

bold.

Supplementary Fig S4: Overall expression levels in the A) head and B) body for male (M) and

female (F) biased and unbiased (U) genes in *D. melanogaster* (mel), *D. simulans* (sim), *D.*

suzukii (suz), *D. ananassae* (ana), *D. subobscura* (sub), and *D. immigrans* (imm). Significance

was assessed for each body part with a type II ANOVA with sex bias (SB) category and

species as factors.

Supplementary Fig S5: Tissue specificity τ in the A) head and B) body for MB (M), FB (F), and

UB (U) genes in *Dmel*, *Dsim*, *Dsuz*, *Dana*, *Dsub*, and *Dimm*. Significance was assessed for each

body part with a type II ANOVA with sex bias (SB) category and species as factors.

Supplementary Fig S6: Overlap in SB genes within the head (H) between species. Shown are

upset plots demonstrating the overlap in UB, FB, or MB genes in pairwise comparisons

between A–E) *Dmel* (mel), A, F–I) *Dsim* (sim), B,F,J–L) *Dsuz* (suz), C,G,J,M,N) *Dana* (ana),

D,H,K,M,O) *Dsub* (sub), and E,I,L,N,O) *Dimm* (imm). Horizontal bars represent the total

number of genes in a species and sex bias category combination. Vertical bars represent the

number of genes in an intersection class. Connected, filled circles underneath a vertical bar

indicate that a species and sex bias category combination is included in an intersection class.

Blue represents genes UB in one species and SB in the other, dark gray represents genes in

the same sex bias category, and light blue indicates genes that switch sex bias between

species.

Supplementary Fig S7: Overlap in sex biased genes within the body (B) between species.

Shown are upset plots demonstrating the overlap in UB, FB, or MB genes in pairwise

comparisons between A–E) *Dmel* (mel), A, F–I) *Dsim* (sim), B,F,J–L) *D. suz* (suz), C,G,J,M,N)

Dana (ana), D,H,K,M,O) *Dsub* (sub), and E,I,L,N,O) *Dimm* (imm). Horizontal bars represent

the total number of genes in a species and sex bias category combination. Vertical bars

represent the number of genes in an intersection class. Connected, filled circles underneath

a vertical bar indicate that a species and sex bias category combination is included in an

intersection class. Blue represents genes UB in one species and SB in the other, dark gray

represents genes in the same sex bias category, and light blue indicates genes that switch

sex bias between species.

Supplementary Fig S8: All SB gains and losses for genes with A,B) large and C,) small

expression changes between the focal and ancestral species in the A,C) head and B,D) body.

The cut-off for a large expression change was a $\log_2$ fold-change of magnitude 1 or higher in

the head and 2 or higher in the body. Genes with concordant (con) changes in expression

between the sexes are located in quadrants ii and iii; while genes with opposing (opp)

expression changes between the sexes are located in quadrants i and iv.

Supplementary Fig S9: Expression changes during sex bias turnover in the head. Shown are

scatterplots of expression in A–C) *Dmel*, D–F) *Dsim*, G–I) *Dsuz*, J–L) *Dana*, M–O) *Dsub*, and P–

R) *Dimm* in each sex relative to ancestral (anc) expression for A,D,G,J,M,P) FB or B,E,H,K,N,O)

MB gains or C,F,I, L,O,R) losses. Spearman's ρ and the associated *P* value for each category

are shown.

Supplementary Fig S10: Expression changes during sex bias turnover in the body. Shown are

scatterplots of expression in A–C) *Dmel*, D–F) *Dsim*, G–I) *Dsuz*, J–L) *Dana*, M–O) *Dsub*, and P–

R) *Dimm* in each sex relative to ancestral (anc) expression for A,D,G,J,M,P) FB or B,E,H,K,N,O)

MB gains or C,F,I, L,O,R) losses. Spearman's ρ and the associated *P* value for each category

are shown.

Supplementary Fig S11: Δx in the A) head and B) body of 5 *Drosophila* species. Shown are

Δx values for FB, MB and UB genes in females (F) and males (M) of *Dmel*, *Dsim*, *Dsuz*, *Dana*,

and *Dsub*. The dashed gray line represents the cut-off above or below which a gene shows

signs of putative positive selection. For better visualization, Δx values above 5 or below -5

are not shown.

Supplementary Fig S12: Chromosomal distribution of signs of putative positive selection at

the gene expression level. Shown are the ratios of observed (obs) to expected (exp) number

of genes under putative positive selection (-1 < Δx > 1) in each sex bias category and sex on

the X chromosome and autosomes (auto) in the A) head and B) body. Significance was

assessed with a $χ^2$ test. ***P < 0.005, **P < 0.01, *P < 0.05, marginally non-significant: .*P* <

0.1. Non-significant comparisons not shown.

Supplementary Data S1: Data underlying analyses.