**Parallel elevational replacement of hosts and parasites in a highly susceptible songbird**

**genus**

Daniele L. F. Wiley[1,2], Jessie L. Williamson[1,3], Silas E. Fischer[4], Selina M. Bauernfeind[1],

Henry M. Streby[4], Kathy Granillo[5], Christopher C. Witt[1,2], and Lisa N. Barrow[1,2]

[1]Museum of Southwestern Biology, University of New Mexico, 302 Yale Blvd NE,

Albuquerque, NM 87131, USA

[2]Department of Biology, University of New Mexico, 302 Yale Blvd NE, Albuquerque, NM

87131, USA

[3]Department of Zoology and Physiology, University of Wyoming, 1000 E. University Ave,

Laramie, WY 82071, USA

[4]Department of Environmental Sciences, University of Toledo, Wolfe Hall Room 1235, 3050

West Towerview Blvd., Toledo, OH 43606-3390, USA

[5]United States Fish and Wildlife Service, Socorro, NM 87801, USA; retired

*Corresponding author: Daniele L. F. Wiley, dwiley7@unm.edu

Address: Department of Biology, University of New Mexico, Castetter Hall 1480, MSC03-

2020, 219 Yale Blvd. NE, Albuquerque, NM 87131, USA

ABSTRACT

Elevational replacement distribution patterns underpin montane diversity and reflect

the interaction of both biotic and abiotic pressures, but the degree to which parasites exhibit

elevational zonation remains unclear. Investigating infection patterns in related host species

across elevational gradients can reveal whether parasites and hosts show concordant patterns

of elevational turnover, potentially due to shared historical and ecological factors. Here, we

assessed patterns of elevational replacement in haemosporidian parasite assemblages that

infect three congeneric songbird species: Bell's Vireo (*Vireo bellii),* Gray Vireo *(V. vicinior),*

and Plumbeous Vireo (*V. plumbeus*), each of which breeds across distinct elevations and

habitats in the southwestern United States. We screened a total of 248 individuals using

cytochrome *b* PCR and microscopy. We identified 19 haemosporidian haplotypes, including

eight novel lineages. We found that each of the three vireo species exhibited high

haemosporidian prevalence (55.0–86.2%), with nearly all infections from the genus

*Haemoproteus* (subgenus *Parahaemoproteus*). Haemosporidian assemblages varied across

elevations; each sampled range of elevations harbored abundant, yet host-specific lineages

with different environmental associations. Bell's and Plumbeous Vireos, but not Gray Vireos,

hosted several phylogenetically distinct, putative generalist lineages, likely reflecting

spillover from more diverse local breeding bird communities. Repeated infections in

individuals across breeding seasons, together with moderate parasitemia ($\bar{x} \approx 1\%$) suggest

that these focal vireo species harbor chronic infections during their respective breeding

seasons. These results demonstrate that elevational replacement patterns in avian hosts may

be mirrored by their haemosporidian parasites, particularly among host-specialized lineages.

Keywords: Avian malaria, diversity, elevational range, haemosporidian, parasite,

*Parahaemoproteus*, Vireonidae, *Vireo*, turnover

## 1. INTRODUCTION

Mountains harbor exceptional biodiversity, in part because elevational gradients tend to be finely partitioned by related species with ranges that are elevationally parapatric (Cadena and Loiselle, 2007; von Humboldt and Bonpland, 2009; Freeman et al., 2016). Yet, it remains unclear whether parasites of elevational replacement host species mirror this range stratification. Generally, host-parasite interactions can strongly influence biogeographic patterns, with reciprocal effects on fitness (Combes, 1997, 2001; Hasik and Siepielski, 2022; Poulin, 2011), population dynamics (Albon et al., 2002), and community structure (Minchella and Scott, 1991, McNew et al., 2021). Strong host-specificity, combined with shared environmental constraints, should result in parasite turnover that parallels the turnover of hosts along environmental gradients. Conversely, aspects of parasite life-history—such as insulation from climatic factors (e.g., endoparasites), dependency on secondary hosts and vectors (e.g., complex life cycle), and degree of host-specificity (e.g., host range and degree of host switching)—should be expected to cause discordant patterns from those observed in hosts (Gage et al. 2008; Krasnov and Poulin, 2010; Poulin and Krasnov, 2010; Ellis et al., 2015).

Regional surveys across elevational gradients reflect these complex relationships. In various mountain systems, parasite turnover has been found to exceed host turnover (Galen and Witt, 2014; Barrow et al., 2021), and host turnover has been found to predict parasite turnover (Ellis et al., 2015; Williamson et al., 2019; McNew et al., 2021). Gradients of precipitation and temperature, as well as landscape features, have also been found to shape parasite communities (Illera et al., 2017; McNew et al., 2021). Investigating the parasite communities of closely related host taxa that are elevational replacements provides a natural experiment for disentangling the relative roles of host associations, parasite traits, and environmental gradients in shaping parasite biogeography, while controlling for host

phylogeny, life history, and parasite susceptibility (Medeiros et al., 2013; Barrow et al.,

2019).

Among blood parasites, haemosporidians (order Haemosporida; Danilewsky, 1885) are a

cosmopolitan group of protozoans with complex life cycles dependent upon vertebrate taxa,

including birds, mammals, and reptiles (Ricklefs and Fallon, 2002; Duval et al., 2007;

Bensch et al., 2009; Galen et al., 2018; Boysen et al., 2022), and invertebrate vectors in the

order Diptera (Linnaeus, 1758). A long history of co-evolution with avian hosts (Ricklefs and

Fallon, 2002; Ricklefs et al., 2004; Lauron et al., 2015; Galen et al., 2018) has given rise to a

remarkable diversity of avian haemosporidian lineages (Valkiūnas, 2005; Hellgren et al.,

2007; Perkins, 2014), with many still yet to be described (Perkins, 2014; Borner et al., 2016;

Galen et al., 2018). This diversity is reflected both phylogenetically and phenotypically, with

variable degrees of pathogenicity (reviewed in Bennett et al., 1993) and host specificity (Ellis

and Bensch, 2018; Fecchio et al., 2019) recorded among and within genera.

Given the high diversity and complexity of haemosporidian parasites, it has remained

challenging to characterize the distribution patterns of haemosporidian lineages across large

geographic scales (Clark, 2018). However, within regions and genera, spatial trends appear—

reflecting the influence of climate directly, through physiological interactions during

transmission and development (Ikemoto 2008; Mordecai et al., 2013), and indirectly, by

influencing host and vector abundance and susceptibility (Filion et al., 2020). As a result, the

fluctuating climate of temperate and montane regions produce seasonal, annual, and habitat-

level waves of infection (Schrader et al., 2003; Bensch et al., 2007; Kim and Tsuda, 2010;

Pérez-Rodríguez et al., 2013; Lutz et al., 2015; Reinoso-Pérez et al., 2024), whereas areas

with higher, less seasonal temperatures, such as tropical regions and lowlands, can have

persistent haemosporidian pressure (Zamora-Vilchis et al., 2012).

Infection intensity, commonly measured as parasitemia (defined as the 'estimated

percentage of haemosporidians circulating in host blood') provides insight into host condition

and stage of haemosporidian infection (i.e., acute or chronic; Valkiūnas, 2005). Like

prevalence (here defined as 'proneness to infection'), parasitemia is also influenced by

environmental factors, with peaks in temperate regions during the breeding season, when

warmer, wetter conditions favor vector activity (Valkiūnas, 2005; Huang et al., 2020;

Reinoso-Pérez et al., 2024) and when chronically infected hosts experience tissue-to-blood

recrudescence (Atkinson and van Riper, 1991). Therefore, montane regions provide a

valuable system for studying avian haemosporidian infection dynamics because of the sharp

local variation in both climate and habitat over relatively small geographic distance (e.g.,

Illera et al., 2017; Rodríguez-Hernández et al., 2021; Williamson et al., 2019).

The southwestern U.S. presents a particularly compelling region for studying

haemosporidian biogeography and dynamics due to its climatic heterogeneity, wide range of

habitat types, and steep elevational gradients in sky island mountains. Although arid overall,

the region exhibits strong variation in precipitation timing and intensity, shaped by both

temperate seasonality and the North American monsoon system. These climatic patterns

interact with topographic variation (reviewed in Coblentz and Riitters, 2004) to create a

mosaic of ecologically distinct elevational zones ranging from lowland riparian corridors

with high relative humidity and permanent water sources, to mid-elevation desert scrublands

and savannas reliant on seasonal rainfall, to cooler montane forests with shorter growing

seasons and higher annual precipitation (Sheppard et al., 2002). While community-level

surveys suggest moderate overall prevalence of haemosporidian infection among avian

communities in the region (mean: ~36%; Barrow et al., 2021), and while environmental

conditions—particularly related to elevation, have been shown to strongly influence spatial

patterns of infection (Williamson et al., 2019)—previous studies have primarily described

environmental and ecological contributions across mid-high elevation habitats and sky

islands. Therefore, further work is needed to expand our understanding of whether observed

environmental and elevational trends persist across broader gradients and habitat types,

especially among particularly susceptible hosts.

Among the many avian species previously surveyed in the desert Southwest, the

Plumbeous Vireo (*Vireo plumbeus*) has consistently stood out for its high haemosporidian

prevalence across sites and years (Marroquin-Flores et al., 2017; Barrow et al., 2021).

Migratory populations of this insectivorous songbird breed in mid- to high-elevation

coniferous woodlands (from ~1,200–3,000 m) across the western U.S. and Mexico (Barlow,

1977; Curson and Goguen, 1998) and spend the nonbreeding season along the Pacific Slope

and in the lowlands of central Mexico (Sibley and Monroe, 1990). Two closely related

migratory vireos, Bell's Vireo (*Vireo bellii*) and Gray Vireo (*Vireo vicinior*), have relatively

similar breeding distributions, yet inhabit distinct habitats and elevations. Gray Vireos breed

in arid mid-elevation juniper (*Juniperus* spp.) savannas and chaparral, ~400–1,900 m

(Hubbard, 1970; Barlow, 1977) and spend the nonbreeding season primarily in lowland

desert and coastal areas of Sonora and Baja California (Barlow et al., 1999; Fischer et al.,

2025). Bell's Vireos breed in lowland riparian habitats, generally below ~1,500 m, with

comparatively high access to water and vegetative cover (Brown, 1993) and spend the

nonbreeding season along the Pacific coast of Mexico, from Baja California and Sonora

south to El Salvador. Given the similarities in ecology and life history among vireos, local

differences in breeding elevational range make these three focal species an ideal comparative

system through which to examine haemosporidian infection patterns across elevational

gradients.

In this study, we tested whether haemosporidian biogeography, community assemblage,

and infection dynamics vary predictably among three species of vireos that occupy distinct

elevations and habitats at sites in New Mexico and Utah, USA. Using a combination of

molecular screening and microscopy, we (1) compared parasite prevalence and infection

intensity among host species, including the first parasite survey for regionally vulnerable

Gray Vireos; (2) characterized haemosporidian haplotypes and host specificity; and (3) tested

the extent to which community composition was linked to elevational zones and/or their

associated host species and habitats. We predicted finding extensive parasite sharing between

elevational replacement vireo species, considering the proximity and interdigitated nature of

their ranges. Based on previously published haemosporidian surveys in vireos, we anticipated

finding high infection prevalence across all three host species; however, we also expected

prevalence and parasitemia to vary among species and environments. Lastly, because the

Southwest is arid, we predicted higher prevalence and parasite diversity in more mesic

habitats suitable for dipterid vectors, such as in the low-elevation riparian corridors used by

Bell's Vireos and high-elevation woodlands occupied by Plumbeous Vireos.

2. MATERIALS AND METHODS

*2.1 Sample collection*

We sampled 248 wild-caught individuals of three vireo species (Bell's Vireo: $n = 20$,

Gray Vireo: $n = 170$, and Plumbeous Vireo: $n = 58$; Fig. 1A; Table S1). Sampled individuals

included adults and juveniles (Table S1). All individuals were sampled during the summer

months (May–August), with the majority sampled prior to monsoon season (~early-July–

September) from 1995–2019 across 28 unique sites in New Mexico and a single site in Utah.

During 2017–2019, targeted fieldwork was conducted to sample Gray Vireos: most

individuals were captured in mist nets, with the exception of hatch-year birds sampled at

nests. Blood samples were taken from the brachial vein and stored in lysis buffer, and

individuals were released at capture sites for ongoing study of Gray Vireo annual cycle

ecology (Fischer, 2020; Fischer et al., 2022; Fischer et al., 2025). In 2018 only, we prepared

and air-dried blood smears in the field ($n = 53$). We also recaptured and collected a second

whole blood sample from fourteen Gray Vireos (approximately one year after their initial

sample collection in 2017), yielding two total whole blood samples and one smear for this

subset of individuals (Table S2). Additional Gray Vireo tissues for four individuals were

deposited at the Museum of Southwestern Biology (hereafter, MSB) at the University of New

Mexico following protocols approved by the UNM Institutional Animal Care and Use

Committee. Samples for Bell's and Plumbeous Vireos were requested as tissue loans from

the MSB. Detailed sampling protocol for Bell's and Plumbeous Vireos are described in

(Marroquin-Flores et al., 2017; Barrow et al., 2021; Gyllenhaal, 2024).

All sampled birds either had specific coordinates recorded at the time of capture or

were caught within 100 m of an existing sampling location. For the latter, we assigned

geographically proximate coordinates based on the nearest documented sampling site or,

when birds were captured between two closely spaced sites, by taking the midpoint between

them (Table S3).

### 2.2 Parasite screening and genetic data collection

We extracted DNA from pectoral muscle ($n = 82$) and whole blood ($n = 166$) using

the QIAGEN DNeasy Blood and Tissue Kit following manufacturer's recommendations.

Birds were screened for *Haemoproteus* and *Plasmodium* parasites using three nested

polymerase chain reaction (PCR) protocols to amplify a 478-base pair (bp) fragment of the

haemosporidian mitochondrial cyt *b* gene (Hellgren et al., 2004; Waldenström et al., 2004).

We amplified parasite DNA with the outer primer pairs HaemNFI/HaemNR3 and

HaemNF/HaemNR2, followed by the nested primer pair HaemF/HaemR2. We prepared outer

PCR reactions in 25 µL volumes, containing 1.25 U of AmpliTaq Gold DNA Polymerase

(Applied Biosystems, Foster City, CA, USA), 1× PCR Buffer II, 2.5 mM MgCl2, 0.2 mM

dNTPs, 0.5 μM of each primer, and 20 ng of template DNA. Thermal cycling conditions

were modified from Galen and Witt (2014), with an initial denaturation at 95 °C for 8 mins,

followed by 20 cycles of 94 °C for 30 seconds, 50 °C for 30 secs, 72 °C for 45 secs, and a

final extension at 72 °C for 10 mins.

For nested PCR, we used 1 μL of the outer PCR product as the template, with reaction

conditions identical to the outer PCR, except for an increase to 35 cycles. Each reaction set

included negative and positive controls to monitor for contamination and confirm

amplification success, respectively. We visualized PCR products on 2% agarose gels stained

with SYBR Safe Gel Stain (Invitrogen, Carlsbad, CA, USA) to verify the presence of

amplicons of the expected size. Successfully amplified products were purified using ExoSap-

IT (Affymetrix, Inc., Santa Clara, CA, USA) and submitted for Sanger sequencing at

Psomagen (Rockville, MD, USA). Gray Vireo blood samples from recaptured individuals

(i.e., samples from the same individuals from 2017 and 2018) underwent PCR amplification,

but only a single sample was sequenced (Table S2).

*2.3 Determining prevalence and haplotype diversity*

We determined positive infections upon successful amplification of haemosporidian

cyt *b* sequences and calculated pathogen prevalence within each host species by summing the

number of infected individuals for that species, divided by the total number of samples

screened for that species. Next, we calculated 95% binomial confidence intervals using the

'exact' method available in the *binom* package in R (version 4.3.2; R Core Team, 2023;

Sundar, 2006).

Haemosporidian cyt *b* forward and reverse reads were trimmed to remove primers,

resulting in the target fragment size of 478 bp and were then assembled using the default

alignment algorithm in Geneious (version 2025.03; Biomatters Ltd; Kearse et al., 2012; Table

S4). To identify haplotypes, we compared cleaned sequences to published records stored in

the public databases GenBank (National Center for Biotechnology Information, US National

Library of Medicine) and MalAvi (Bensch et al., 2009) by using the Basic Local Alignment

Search Tool (BLAST). Additionally, to ensure the accuracy of both established and novel

haplotypes, we downloaded all haemosporidian cyt *b* haplotype sequence files from MalAvi

and compared the number of differences, if any, to our sequences via a distance matrix

calculated in Geneious. Haplotypes that differed by one or more base pairs (~0.2% sequence

divergence) from published sequences on GenBank or MalAvi were considered novel and

named following MalAvi conventions. We compared coinfections with existing haplotypes in

our dataset to determine individual haplotypes when possible.

*2.4 Phylogenetic analysis*

We estimated phylogenetic relationships among haemosporidian haplotypes using a

maximum-likelihood (ML) framework in RAxML, v8.2.10 (Stamatakis, 2014). Using the

GTR+G model of nucleotide substitution, we conducted a rapid bootstrap analysis with 1000

replicates, after which we searched for the best-scoring ML tree. We rooted the tree with

*Leucocytozoon* (COLBF21, GenBank Accession MK947795) based on the current

phylogenetic hypothesis for avian haemosporidians (Borner et al., 2016; Galen et al., 2018).

Nodes with <50% bootstrap support were collapsed in TreeGraph2 (Stöver and Müller,

2010). We determined the relationships among the three vireo study species based on the

posterior distribution of likely trees available on birdtree.org with the 'Hackett All Species'

option (Hackett et al., 2008; Jetz et al., 2012). We downloaded 100 phylogeny subsets and

selected the first tree because the relationships were consistent among the 100 trees.

### 2.5 Estimating parasite species richness and sampling completeness

We used the *iNEXT* R package (version 3.0.1; Chao et al., 2014; Hsieh et al., 2024) to estimate parasite lineage diversity and evaluate sampling completeness for host species with adequate data (i.e., Gray and Plumbeous Vireos). In separate analyses for each host species, we used infection counts of each parasite haplotype to generate rarefaction and extrapolation curves for parasite species richness ($q = 0$), with the endpoint set at 400 individuals (i.e., infections). We applied the 'iNEXT()' function to compute species richness estimates, including observed richness, extrapolated richness, and associated confidence intervals using standard function parameters. Additionally, we used the output to estimate sampling completeness (Chao et al., 2014) and defined this value as the minimum number of individual birds required to detect 95% of the total estimated parasite haplotypes infecting each host species. To visualize the rarefaction curves, we used the 'ggiNEXT()' function and *ggplot2* package for style modifications (Wickham, 2016).

### 2.6 Microscopy and parasitemia calculations

We obtained blood smears from the MSB for Plumbeous Vireos, when available ($n = 28$). Blood smears were made and air dried in the field for Gray Vireos ($n = 53$). We did not obtain blood smears for Bell's Vireos. Each slide was fixed with absolute methanol and stained for 50 mins with Giemsa solution (pH 7.0; Sigma-Aldrich, St. Louis, MO, USA). We then examined each slide to confirm infection status using light microscopy on an Olympus BX 53 Microscope following the protocol described in Valkiūnas (2005), where 10,000 erythrocytes were scanned at 1000x magnification with an oil immersion lens to identify and count *Parahaemoproteus* and *Plasmodium* infections (Valkiūnas, 2005). We took digital photographs of representative haemosporidian specimens, which were archived at the MSB and are searchable via the Arctos database (https://arctos.database.museum/). If parasites

were found free-floating outside of erythrocytes, either due to the parasite's developmental

stage or by erythrocyte lysis due to factors during the preparation of the blood smear, they

were separately tallied and not included in the parasitemia calculation (Table S1). We

calculated parasitemia, or the estimated percentage of erythrocytes infected with

haemosporidian parasites, as the number of infected erythrocytes out of 10,000 screened. To

account for lack of normality in the data, we used a bootstrap method with 1,000 replications

to calculate parasitemia 95% confidence intervals via the R package *boot* (Kushary, 2000).

***2.7 Characterizing environmental variation***

To obtain elevation in meters (m) from coordinates that did not contain original

elevation data, we used the 'elevation()' function from the R package *rgbif* (Chamberlain and

Boettiger, 2017; Chamberlain et al., 2024). We identified and removed elevation outliers

prior to analysis, as described in (Williamson and Witt, 2021).

We characterized climatic variation using 19 bioclimatic variables at 30 sec (~1 km²)

resolution from the WorldClim 2.1 database (Fick and Hijmans, 2017). We used principal

components analysis (PCA) for temperature (Bio1–11) and precipitation (Bio12–19)

variables to create a composite measure of temperature and precipitation across the gradient.

The first two axes for temperature (hereafter, tempPC1 and tempPC2) explained a combined

84.2% of the variation and the first two axes for precipitation (hereafter, precipPC1 and

precipPC2) explained 95.1% of the variation (Fig. S1; Table S5-6). Variable loadings

indicated that tempPC1 primarily represented year-round temperatures, with higher values

relating to lower average temperatures, while tempPC2 represented greater seasonality and

less temperature stability, with higher values relating to more extreme temperature

fluctuations throughout the year and day-night cycle. Similarly, precipPC1 described the

overall precipitation amount, with higher values indicating wetter conditions, whereas

precipPC2 represented precipitation seasonality, with higher values representing more stable

and less seasonal precipitation (Table S5-6).

*2.8 Assessing infection dynamics among host species and infection types*

We tested whether haemosporidian infection status differed between categorical

variables such as host species and tissue type (i.e., pectoral muscle or whole blood samples),

using Pearson's chi-squared tests via the *stats* package. To examine the relative magnitude of

effect for each comparison, we calculated Cramér's V effect sizes were using the 'CramerV()'

function in the *rcompanion* package (Mangiafico, 2024). Next, to assess differences in

parasitemia, we first normalized the data using a natural log $\ln(1+x)$ transformation and

removed two influential outliers with parasitemia >4% (Gray Vireo IDs: 272132427 and

272132642). We then evaluated residual variance with the 'var.test()' function in the *stats*

package and compared log-transformed parasitemia between Gray and Plumbeous Vireos and

infection types (i.e., single vs. coinfected) using two-tailed Student's *t*-tests.

*2.9 Intraspecific infection prevalence and parasitemia across environments*

We employed a two-pronged approach to assess environmental associations with

infection prevalence: First, we used univariate, non-parametric tests to characterize and

compare intraspecific infection patterns across abiotic variables relevant to parasite and

vector ecology, including elevation, overall temperature (tempPC1) and precipitation

(precipPC1), and the seasonality of temperature (tempPC2) and precipitation (precipPC2).

Then, we built intraspecific multivariate linear models and applied nested and exhaustive

model selection to evaluate the strength and direction of environmental effects. We adopted

this approach to account for the structure of our data (i.e., non-overlapping elevational and

environmental bands distinctly correlated to species identity) and limitations of our dataset

(i.e., limited sample sizes, degree of environmental variation, and infection class imbalances).

We first assessed data normality both visually, using the 'qqplot()' function from the

*car* package (Fox and Weisberg, 2019), and then statistically, using the Shapiro-Wilk test

using the 'shapiro.test()' function from the *stats* package. Given the non-normal distribution

of environmental and geographic data in our dataset, we applied a rank-sum approach via

Mann-Whitney U-tests to compare the distributions of environmental variables between

infected and uninfected birds. Wilcoxon effect sizes of each comparison were then calculated

via the 'wilcox_effsize()' function in the *rstatix* package (Kassambara, 2023).

We then constructed species-specific generalized linear models using glm() from the

*stats* package. Prevalence model sets used a binomial distribution (hereafter: "BEVI_prev"

for Bell's Vireo, "GRVI_prev" for Gray Vireo, and "PLVI_prev" for Plumbeous Vireo sets),

while parasitemia model sets used a Gaussian distribution ("GRVI_para", "PLVI_para"). We

evaluated linearity between the logit (log(probabilities/1-probabilities)) and predicted values

in prevalence models visually via *ggplot2* package and assessed residual diagnostics in

parasitemia models and with 'autoplot()' within the *ggfortify* package (Tang et al., 2016).

Multicollinearity between variables was assessed by calculating variance inflation factors

(VIF) via the *car* package. As expected, there was high collinearity with aspects of

geographic and topographic variation, i.e. latitude and elevation, with other environmental

variables. We retained elevation in all model sets because it captured variation in habitat and

vector dynamics important to infection that were not explained by temperature and

precipitation alone (Ishtiaq and Barve, 2018); however, we removed latitude from all linear

models to reduce overfitting.

For model sets with adequate sampling (i.e., GRVI_prev, PLVI_prev, GRVI_para,

PLVI_para), we evaluated all possible additive combinations of five predictor variables:

elevation, tempPC1, tempPC2, precipPC1, precipPC2. We used the 'dredge' function in the

*MuMIn* package (Bartoń, 2024) for exhaustive model selection. For host datasets with < 30

data points (BEVI_prev and PLVI_para) we restricted candidate models to simple,

biologically relevant structures to avoid overfitting. Specifically, we included univariate

models nested within simple additive models containing at most two predictor variables, as

well as null and global models for comparison. We then compared model performance and

evaluated the trade-off between model fit and complexity using Akaike's Information

Criterion corrected for small sample sizes (AICc; Table 1).

To examine the effect of each predictor in top-ranked models, we standardized and

calculated model-averaged regression coefficients using the *MuMIN* package (Table 2). For

top-ranked models where the null model was not favored, we assessed goodness of fit via

Hosmer-Lemeshow tests as part of the *ResourceSelection* package (Lele et al., 2023) and

checked for overdispersion by dividing residual deviance by the degrees of freedom.

Predictive accuracy was also assessed using 10-fold cross-validation via the *caret* package

(Kuhn, 2008).

3. RESULTS

*3.1 Haemosporidian prevalence, abundance, and diversity*

We uncovered an unusually high prevalence of haemosporidian infections in all three

vireo species, with 175 infected birds of 248 screened (70.6%; 95% CI [64.5 – 76.2%]). The

highest documented prevalence was found in Plumbeous Vireos (84.5% infected; 95% CI

[72.6 – 92.7%]), followed by Gray Vireos (67.7%; 95% CI [60.0 – 74.7%]), and lastly, Bell's

Vireos (55%; 95% CI [31.5 – 76.9%]; Fig. 1B). Prevalence was significantly lower in Bell's

than in Plumbeous Vireos ($\chi^2(1) = 5.72$, $p = 0.017$, $\varphi = 0.31$), indicating a moderate effect

size. While most birds were infected by single haemosporidian haplotype infections (88.1%;

95% CI [82.3 – 92.5%]), we found 21 individuals that were coinfected with two or more haplotypes (12.0%; 95% CI [7.54 – 17.7%]), and one Gray Vireo that was infected by three haplotypes (VIRVIC01, VIRVIC02, VIRVIC03). Of the 14 Gray Vireos recaptured about a year after initial sampling, 12 were infected with haemosporidians in both years, while two that were uninfected in 2017 tested positive in 2018.

Nearly all infections were caused by parasites in the genus *Haemoproteus*, specifically the subgenus *Parahaemoproteus* (99.5%), which comprised 18 of 19 haplotypes identified. *Plasmodium* was rare in our data, with a single infection (lineage VIRPLU13) recorded from a Plumbeous Vireo (MSB:Bird:60726; Table S4). Of the 19 total parasite haplotypes identified, 8 were novel (Fig. 1D; Table S4).

### 3.2 Estimates of parasite species richness and sampling completeness

We identified a total of three haemosporidian haplotypes from Bell's Vireos, four from Gray Vireos, and 14 from Plumbeous Vireos (Fig. 1D). These differences were partly, but not entirely, attributable to differences in sample size. Gray Vireo *Parahaemoproteus* infections were substantially less diverse than those of Plumbeous Vireos despite more intensive screening efforts. Extrapolation from rarefaction curves showed considerably lower estimated parasite richness for Gray Vireos ($4.99 \pm 0.53$ SE; 95% CI [4.00–6.03]; Table S7) and estimated that much smaller sample sizes would be needed to reach 95% sampling completeness (14 infections; Fig. 2; Table S8). This result contrasted our results for Plumbeous Vireos, where parasite richness was estimated to be ~10x higher than Gray Vireos ($53.8 \pm 23.3$ SE; 95% CI [14.0–99.6]) and require much greater sampling effort (~26x more than estimated for Gray Vireos) to reach an estimated 95% completeness (estimated 364 infections). Bell's Vireos also exhibited low lineage diversity ($3.92 \pm 0.784$ SE; 95% CI [3.00-5.45] and an estimated 95% sample completeness at only 33 infections, consistent with

the limited diversity recovered (Table S7-S8). However, we caution overinterpreting these

results due to the small Bell's Vireo sample size (*n* = 12 infected) in our study and

recommend additional sampling to generate more precise estimates.

Of the 19 total haplotypes identified, we found evidence of potential host specificity

with at least one species-specific haplotype infecting each of the three vireo species (Table

S4). Of note, Gray Vireos were infected by novel haemosporidian haplotypes only,

(VIRVIC01-VIRVIC04). This finding contrasted with Plumbeous Vireos, which showed

novel haplotypes (i.e. VIRPLU10-13) and haplotypes previously found only in Plumbeous

Vireos (i.e. VIRPLU03, VIRPLU07-09), as well as lineages known to infect other non-vireo

species (i.e. SETAUD14, TROAED12, VIGIL07, VIRPLU01, and VIRPLU04). Sampling

was comparatively lower for Bell's Vireos; no novel lineages were detected. We did detect,

however, a recently described lineage (VIRBEL01) which was previously documented in

Bell's Vireos sampled from Arizona and Texas (Fecchio et al., 2023), as well as other more

generalist lineages (i.e. TROAED12 and VIGIL07; Table S4).

***3.3 Intensity of infection (parasitemia) across hosts and haplotypes***

We screened 102 individuals via microscopy (Gray *n* = 71, Plumbeous *n* = 31). A

total of 81 samples showed evidence of infection and had parasitemia *calculated*

*(Parahaemoproteus:* Gray *n* = 53 and Plumbeous *n* = 27; *Plasmodium*: Plumbeous *n* = 1).

Gray Vireos typically had slightly higher infection loads (1.00%; 95% CI [0.782 – 1.27%])

compared to Plumbeous Vireos (0.786%; 95% CI [0.584 – 1.01%]), partly due to two outlier

Gray Vireo samples with >4% parasitemia (ID 272132427 and 27213264; Fig 1C). After

removing these outliers, we found that mean parasitemia did not differ significantly between

host species (t(77) = 0.656, p = 0.514). When we compared single-haplotypes and

coinfections, we also found no significant difference in parasitemia between species (t(73) = -

0.728, p = 0.469), though we note the sampling pool of coinfected birds was small compared

to single infections ($n = 7$ and $n = 68$, respectively).

Our comparison of infections using PCR and microscopy was largely consistent, with

few detection errors identified between the two methods. We detected four positively infected

birds via microscopy that did not successfully amplify by PCR (Table S1). These samples

had a range of parasitemia spanning 0.70% to 2.68%. Additionally, we failed to detect

haemosporidian infections via microscopy for two birds that were later successfully amplified

using PCR; one individual was the single coinfected Gray Vireo with three

*Parahaemoproteous* haplotypes that were identified by sequencing (Table S1).

*3.4 Environmental patterns of infection status and parasitemia*

*Parahaemoproteus* prevalence showed idiosyncratic patterns among host species and

their associated habitats (Fig 3; Table S9). For example, we found that infected Bell's Vireos

tended to be found at lower latitudes ($z = -2.09$, $p = 0.040$), with warmer year-round

temperatures (tempPC1; $z = -2.25$, $p = 0.027$ Fig. 3; Fig. S2; Table S9). Similarly, probability

of infection increased as overall precipitation (precipPC1) and temperature seasonality

(tempPC2) increased (Table 1; Table 2). In contrast, we found higher proportions of infected

Gray Vireos at lower elevations ($z = -2.07$, $p = 0.038$), Fig. 1C; Table S9) and at sites with

more stable precipitation (precipPC2; $z = 1.72$, $p = 0.057$), Fig 3; Table S9). However, this

result was dampened when we excluded the eleven infected Gray Vireos from the Abajo

Mountain site in Utah, which were outliers in many of the environmental PCA dimensions.

The top model, however, showed that infection was negatively associated with elevation and

positively associated with temperature (tempPC1; Table 1; Table 2) even when excluding the

Utah site. Lastly, Plumbeous Vireo infections tended to be higher in areas with more

seasonally stable precipitation (precipPC2; $z = 2.04$, $p = 0.042$), Fig. 3). Despite the top-

ranked infection probability model showing no significant linear trends, precipitation

(precipPC1 and precipPC2) was present in the majority of top candidate models (Table 1).

We did not find any association between parasitemia and environmental variables in

Plumbeous Vireos, with the null model having the strongest support and no clear predictor

variable significantly reported across top models (Table S10). Gray Vireos, however, showed

a significant pattern of increased parasitemia as overall precipitation (precipPC1) increased

(Table S10), but this model had low explanatory power ($R^2 = 0.05$).

4. DISCUSSION

Vireo species breeding in the southwestern U.S. are especially susceptible to

haemosporidian infection, particularly by parasites in the *Parahaemoproteus* subgenus. In

this study, we uncovered that the elevational limits of haemosporidian parasites closely

parallel those of their avian hosts. We observed striking differences in host specificity and

parasite richness across host species and their associated habitats, revealing that the high-

elevation Plumbeous Vireos harbored the most diverse and phylogenetically distinct parasite

lineages, especially compared to another well-sampled, lower elevation host species, the

Gray Vireo. Although parasite prevalence differed significantly between Bell's Vireos (55%

infected) and Plumbeous Vireos (86.2% infected), which breed under different levels of

precipitation and temperature, we found variable links between infection status and climatic

factors in our dataset. Specifically, we found that infection probability was strongly

influenced by aspects of temperature for Bell's Vireos, elevation for Gray Vireos, and

precipitation for Plumbeous Vireos, likely reflecting subtle differences in habitat suitability

for arthropod vectors. Lastly, we found low to moderate parasitemia values in both Gray and

Plumbeous Vireos, consistent with chronic infections. We examine these results in relation to

previous haemosporidian research in the southwest U.S. and in vireos, both in their breeding

and nonbreeding grounds, and discuss remaining questions in the sections below.

*4.1 Elevational replacement and haemosporidian diversity patterns*

Avian haemosporidians exhibit variable patterns of turnover across elevations (Galen

and Witt, 2014; Illera et al., 2015; McNew et al., 2021), which is often independent of

geographic distance between hosts (Williamson et al., 2019; Barrow et al., 2021). We

demonstrated a strong case of parallel elevational replacement in hosts and parasites, with

high turnover among *Parahaemoproteus* assemblages infecting vireo species breeding at

different elevations. This elevational replacement pattern could be the result of a few

different scenarios. For example, shared historical and ecological forces, such as habitat

specialization in both hosts and vectors (Ellis et al., 2020), may shape the elevational range

limits of both vireos and their *Parahaemoproteus* parasites. Alternatively, species-specific

factors that co-evolved independently in response to specialized parasite lineage pressures,

such as immune defenses (e.g., Armour et al., 2025), could also be at play. Whether parasite

elevational replacement is host-driven or independently mediated by the environmental

gradient, global patterns of spatial diversity in parasites should be tightly coupled with those

of their hosts.

Of the 30 cyt *b Parahaemoproteus* and 12 *Plasmodium* haplotypes currently reported

to infect vireo species (MalAvi; downloaded 1/1/2025), our study adds seven novel

*Parahaemoproteus* and one novel *Plasmodium* haplotype. The near absence of *Plasmodium*

infections, even at lower elevations, contrasts findings from other temperate region studies

(e.g. in Alaska: Oakgrove et al., 2014; California: Walther et al., 2016; Oklahoma: Wyckoff

et al., 2024) but is consistent with previous screening at high elevations in New Mexico

(Marroquin-Flores et al., 2017; Williamson et al., 2019; Barrow et al., 2021). While the

cooler temperatures experienced at higher elevations have been proposed as a potential

limiting factor for haemosporidian infection (e.g. LaPointe et al., 2012; Barrow et al., 2021),

the low prevalence of *Plasmodium* at lower elevations in this region may also reflect limited

suitable habitat for mosquito vectors prior to the onset of monsoon rains. Broader and

temporally replicated surveys of potential vectors and vector abundance in this region are

needed to fully explain this apparent absence.

We identified an interesting pattern among the *Parahaemoproteus* lineages found in

our focal species: Plumbeous Vireos harbored a broad diversity of parasite haplotypes

relative to the other two sampled species, while Gray Vireos exhibited a depauperate and

likely specialized haemosporidian assemblage (Fig. 1D). This disparity suggests that

Plumbeous Vireos may support more permissive or ecologically flexible parasite

associations, while Gray Vireos may be undergoing, or may have already undergone, a

transition toward fixed symbiosis with a narrower set of haemosporidian lineages.  Although

the degree of cyt *b* divergence among lineages is generally low (<5% in most haplotypes

based on post-hoc Jukes–Cantor distance assessments), a few lineages (e.g., SETAUD14,

VIRPLU07) diverge more substantially (>5%) from others such as VIRVIC01–03 and

TROAED12 (Hellgren et al., 2007; Table S11). The contrasting pattern of haemosporidian

diversity between vireo species could alternatively be caused by differences in avian host

richness, and the low diversity found in Gray Vireos could reflect the depauperate breeding

bird communities of the arid and structurally simple juniper savanna habitats.

*4.2 Conserved high susceptibility to haemosporidians among vireo species*

Vireos stand out among southwestern songbirds for their consistently high

haemosporidian infection rates, providing a valuable opportunity to examine parasite

dynamics in hyper-susceptible hosts. In the three species screened here, we found that 70.6%

of 248 individuals screened had at least one haemosporidian infection, which is greater than

double the prevalence reported in broader community-wide surveys conducted in the

southwestern U.S. over a similar period. For example, Marroquin-Flores et al. (2017)

screened 186 birds of 49 species sampled in New Mexico from June-July 2016, including 13

individual vireos of two species: $n = 4$ Warbling Vireos (*Vireo gilvus*) and $n = 9$ Plumbeous

Vireos. The average proportion of infection was 36.6% across all studied species, but 92% of

vireos tested positive. A subsequent large-scale study by Barrow et al. (2021) reported similar

patterns, with an overall haemosporidian prevalence of 36.1% among 776 total individuals of

61 species. Of these, vireos again exhibited significantly higher infection proportions, with

Plumbeous and Warbling Vireos demonstrating the highest prevalence in the study at 87%

and 78%, respectively (Barrow et al., 2021).

This observation of high susceptibility in vireos is not limited to birds sampled in the

southwestern U.S., as other studies across breeding sites have also reported higher relative

prevalences in the *Vireo* genus. Granthon and Williams (2017) reported that 100% of the 62

PCR-screened Red-eyed Vireos (*Vireo olivaceus*) from Pennsylvania were infected with

*Plasmodium*, *Haemoproteus*, or *Leucocytozoon* species (Granthon and Williams, 2017).

Rodriquez et al. (2021) found *Haemoproteus* prevalence of 59% in 22 individual Warbling

Vireos sampled from Colorado breeding sites (Rodriguez et al., 2021). Shared susceptibility

within closely related species has been extensively documented (e.g. González et al., 2014;

Lutz et al., 2015; Barrow et al., 2019; Pigeault et al., 2022) but attempts to identify the

specific factors that contribute to high prevalence in vireos are still lacking. Of the conserved

host traits recorded to have consistent interaction with haemosporidian prevalence in other

avian clades, thus far, shared life history traits such as open cup nest type and insectivorous

diets (as reported and reviewed in Braga et al., 2011; Rodriguez et al., 2021) may increase

exposure risk, at least partially explaining the high prevalence in this group. However, a

combination of traits, including behavior (e.g., time spent incubating and/or brooding at the

nest), features of nest microhabitat (e.g., height of nest from ground), and aspects of

immunity (e.g., haemosporidian tolerance) may also contribute to the elevated susceptibility

(González et al., 2014; Lutz et al., 2015; Muriel, 2020; Rodriguez et al., 2021; Pigeault et al.,

2022).

Interestingly, of the 14 Gray Vireos that we recaptured and screened for parasites

across two sampling years (2017 and 2018), 12 were positive via PCR in both years, and the

remaining two birds became infected by the second year (Table S2). Although more data is

needed to determine whether these represent chronic infections or reinfections, and pinpoint

where transmission occurs, these results indicate Gray Vireos likely experience sustained

parasite pressure during the breeding season. Continued monitoring of haemosporidian

infection across vireo populations, coupled with investigations into the mechanisms

underlying their susceptibility, could yield valuable insights into the spatial and temporal

dynamics of infection risk in this clade.

*4.3 Environmental determinants of haemosporidian infection vary by host species*

Previous studies have found strong links between haemosporidian prevalence and

topographic, geographic, and climatic variables (Pérez-Rodríguez et al., 2013; Rodriguez et

al., 2021), but the direction and magnitude of these effects vary by region and

haemosporidian clade (Williamson et al., 2019). In the arid southwest, we expected

haemosporidian prevalence to correspond with water availability, which is important for

insect vector (Valkiūnas, 2004; Santiago-Alarcon et al., 2012), such as those associated with

potential water sources (i.e. riparian habitats) and areas with higher, more seasonally stable

precipitation (i.e. montane woodlands). Somewhat contrary to expectations given predictions

about the relative aridity of habitat during the pre-monsoon sampling period, we found that

interspecific differences in prevalence increased linearly with elevation and precipitation:

Bell's Vireos showed the lowest prevalence and Gray Vireos exhibited unexpectedly high

infection rates. We were unable to determine whether observed infection patterns were driven

by species-specific traits or abiotic environmental differences, and increased sampling of

Bell's Vireos may help clarify or reduce these differences. Within-species analyses revealed

no consistent predictors of infection across all taxa (Fig. S2-6); instead, different

environmental variables were associated with infection in each species (Fig. 3; Table S9-

S10). Expanded screening of haemosporidians across low- to mid-elevation habitats, among

co-distributed species, at variable distance from water sources, and across precipitation

gradients would further elucidate cross-elevational trends and enhance predictions of

infection probability in avian communities of the southwestern U.S.

### 4.4 Moderate and consistent infection loads across environmental gradients

Gray and Plumbeous Vireos exhibited similar average infection loads (~1% in Gray

Vireos, 0.786% in Plumbeous, range: 0.03–4.5%) with no significant differences between

species (Fig. 1B). These values slightly exceed the regional cross-species average (<1%;

Marroquin-Flores et al., 2017) and fall within the range of moderate infection intensities

reported in experimental studies of other passerines (e.g. Asghar et al., 2012). We did note,

however, two Gray Vireos with elevated parasitemia (4.5 and 4.49%) were sampled at

distinct sites in the Sevilleta National Wildlife Refuge. The elevated parasitemia in these

individuals likely represented acute infection acquired shortly before capture and prior to

immune response, or recrudescence of chronic infections exacerbated by physiological

stressors such as nutritional deficits or hormonal fluctuations (Arriero et al., 2018; Names et

al., 2021). The absence of high parasitemia in Bell's and Plumbeous Vireos could be an

artifact of limited sampling, as birds with severe infections often exhibit reduced mobility

and capture probability (Mukhin et al., 2016), suggesting that heavily infected individuals in

these species may be underrepresented.

Additionally, we found no relationship between parasitemia and any tested

environmental variables for Plumbeous Vireos. In Gray Vireos we saw a significant linear

trend in parasitemia with increased overall precipitation (precipPC1; Table S10), but this

model had weak explanatory power ($R^2 = 0.05$). These results are consistent with previous

findings which suggest that haemosporidian parasitemia patterns are shaped by complex

interactions between host and parasite biology not easily captured in models with abiotic

factors alone (Loiseau et al., 2013).

*4.5 Remaining questions*

First, expanding screening to include *Leucocytozoon* parasites of vireos would

provide a useful contrast with *Parahaemoproteus*; this could be especially important at high

elevations, since *Leucocytozoon* is more common at cooler, high-elevation sites and has been

detected in Plumbeous Vireos in this region (Galen and Witt, 2014; Marroquin-Flores et al.,

2017; Barrow et al., 2019; Barrow et al., 2021). Second, incorporating a multi-locus nuclear

DNA data (as described in Barrow et al., 2018), together with morphological data, could

provide greater resolution for distinguishing closely related *Parahaemoproteus* haplotypes,

offering clearer insights into their evolutionary history and host specificity. Finally, the host

species-specific environmental patterns that we captured might reflect dynamics of arthropod

vectors, which still remain largely uncharacterized in the southwestern United States.

*4.6 Conclusions*

In this study, we found that parasite elevational distributions mirror the elevational

replacement pattern of host species among vireos in southwestern North America.This

pattern may have resulted from host specialization, climate variation independently acting on

hosts and parasites, climate variation acting indirectly through habitat or broader host and

vector communities, or a combination of those forces. We found that vireos were consistently

infected at high rates and largely harbor chronic, low-to-moderate intensity infections

dominated by *Parahaemoproteus*, suggesting the potential for exclusive, reciprocal selective

pressures that facilitate host-parasite coadaptation. The contrast in parasite richness between

Plumbeous Vireos and Gray Vireos was striking: Gray Vireos harbored a limited number of

haemosporidian lineages, all of which were host specific, while Plumbeous Vireos were

infected with a diverse set of lineages. These differences imply species-specific variability in

resistance, tolerance, or exposure to generalist haemosporidian lineages. In sum, our work

adds to a growing body of evidence about clade-specific avian host susceptibility and host-

parasite turnover across elevational gradients, highlighting vireos as an ideal focal group for

future work on host-parasite dynamics across space and time.

ACKNOWLEDGEMENTS

The authors sincerely thank Bethany Abrahamson, Michael Andersen, Matthew Baumann,

Alison Boyer, Serina Brady, Sara Brant, Mariel Campbell, Andrea Chavez, Mina Carnicom,

Lida Crooks, Rosario Marroquin-Flores, Chauncey Gadek, Spencer Galen, Ethan Gyllenhaal,

Mike Hartshorne, Andy Johnson, Tae Kunisawa, Michael Lelevier, Joseph Manthey, Xena

Mapel, Taylor Martinez, Jenna McCullough, Moses Michelsohn, Kathleen Ramsay, George

Rosenberg, C. Jonathan Schmitt, Donna Schmitt, Brenda Villanueva, and Toby Weisenhaus

for their support and contributions.

REFERENCES

Albon, S.D., Stien, A., Irvine, R.J., Langvatn, R., Ropstad, E., Halvorsen, O., 2002. The role of parasites in the dynamics of a reindeer population. Proc. R. Soc. Lond. B Biol. Sci. 269, 1625–1632. https://doi.org/10.1098/rspb.2002.2064.

Arriero, E., Pérez-Tris, J., Ramírez, A., Remacha, C., 2018. Trade-off between tolerance and resistance to infections: an experimental approach with malaria parasites in a passerine bird. Oecologia 188, 1001–1010. https://doi.org/10.1007/s00442-018-4290-4.

Armour, C., Hone A.L., Dunn J.C., 2025. Do specialist haemoparasites induce tolerance in their hosts? Parasitology. 152, 374–380. https://doi.org/10.1017/ S0031182025000393.

Asghar, M., Westerdahl, H., Zehtindjiev, P., Ilieva, M., Hasselquist, D., Bensch, S., 2012. Primary peak and chronic malaria infection levels are correlated in experimentally infected great reed warblers. Parasitology 139, 1246–1252. https://doi.org/10.1017/S0031182012000510.

Atkinson, C.T., van Riper, C., 1991. Pathogenicity and epizootiology of avian haematozoa: *Plasmodium, Leucocytozoon, and Haemoproteus*, in: Loye, J.E., Zuk, M. (Eds.), Bird-Parasite Interactions: Ecology, Evolution, and Behavior. Oxford Academic Press, pp. 19–48.

Barlow, J.C., 1977. Effects of habitat attrition on vireo distribution and population density in the northern Chihuahuan Desert, in: Wauer, R.H., Riskind, D.H. (Eds.), Transactions of the Symposium on the Biological Resources of the Chihuahuan Desert Region, United States and Mexico. U.S. Department of the Interior National Park Service Trans. Proc., pp. 591–596.

Barlow, J.C., Leckie, S.N., Baril, C.T., 1999. Gray Vireo (*Vireo vicinior*), in: Poole, A., Gill, F. (Eds.), The birds of North America. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, D.C.

Barrow, L.N., Bauernfeind, S.M., Cruz, P.A., Williamson, J.L., Wiley, D.L.F., Ford, J.E., Baumann, M.J., Brady, S.S., Chavez, A.N., Gadek, C.R., Galen, S.C., Johnson, A.B., Mapel, X.M., Marroquin-Flores, R.A., Martinez, T.E., McCullough, J.M., McLaughlin, J.E.,

663   Witt, C.C., 2021. Detecting turnover among complex communities using null models: a case

664     study with sky-island haemosporidian parasites. Oecologia 195, 435–451.

665     https://doi.org/10.1007/s00442-021-04854-6.

666   Barrow, L.N., Lemmon, A.R., Lemmon, E.M., 2018. Targeted sampling and target capture:

667     assessing phylogeographic concordance with genome-wide data. Syst. Biol. 67, 979–996.

668     https://doi.org/10.1093/sysbio/syy021

669   Barrow, L.N., McNew, S.M., Mitchell, N., Galen, S.C., Lutz, H.L., Skeen, H., Valqui, T.,

670     Weckstein, J.D., Witt, C.C., 2019. Deeply conserved susceptibility in a multi-host, multi-

671     parasite system. Ecol. Lett. 22, 987–998. https://doi.org/10.1111/ele.13263.

672   Bartoń, K. 2024. MuMIN: Multi-model inference. R package version 1.48.4. https://CRAN.R-

673     project.org/package=MuMIn.

674   Bennett, G.F., Peirce, M.A., Ashford, R.W., 1993. Avian Haematozoa: mortality and pathogenicity.

675     J. Nat. Hist. 27, 993–1001. https://doi.org/10.1080/00222939300770621.

676   Bensch, S., Hellgren, O., Pérez-Tris, J., 2009. MalAvi: a public database of malaria parasites and

677     related haemosporidians in avian hosts based on mitochondrial cytochrome *b* lineages. Mol.

678     Ecol. Resour. 9, 1353–1358. https://doi.org/10.1111/j.1755-0998.2009.02692.x.

679   Bensch, S., Waldenström, J., Jonzén, N., Westerdahl, H., Hansson, B., Sejberg, D., Hasselquist, D.,

680     2007. Temporal dynamics and diversity of avian malaria parasites in a single host species. J.

681     Anim. Ecol. 76, 112–122. https://doi.org/10.1111/j.1365-2656.2006.01176.x.

682   Billerman, S. M., B. K. Keeney, G. M. Kirwan, F. Medrano, N. D. Sly, and M. G. Smith, Ed. 2025.

683     Birds of the World. Cornell Laboratory of Ornithology, Ithaca, NY, USA.

684     https://birdsoftheworld.org/bow/home.

685   Borner, J., Pick, C., Thiede, J., Kolawole, O.M., Kingsley, M.T., Schulze, J., Cottontail, V.M.,

686     Wellinghausen, N., Schmidt-Chanasit, J., Bruchhaus, I., Burmester, T., 2016. Phylogeny of

687     haemosporidian blood parasites revealed by a multi-gene approach. Mol. Phylogenet. Evol.

688     94, 221–231. https://doi.org/10.1016/j.ympev.2015.09.003.

Boysen, K.E., Perkins, S.L., Hunjan, S., Oliver, P., Gardner, M.G., Balasubramaniam, S., Melville, J., 2022. Diversity and phylogenetic relationships of haemosporidian and hemogregarine parasites in Australian lizards. Mol. Phylogenet. Evol. 167, 107358. https://doi.org/10.1016/j.ympev.2021.107358.

Braga, É.M., Silveira, P., Belo, N.O., Valkiūnas, G., 2011. Recent advances in the study of avian malaria: an overview with an emphasis on the distribution of Plasmodium spp in Brazil. Mem. Inst. Oswaldo Cruz 106, 3–11. https://doi.org/10.1590/S0074-02762011000900002.

Brown, B.T., 1993. Bell's Vireo (*Vireo bellii*), in: Poole, A., Gill, F. (Eds.), The birds of North America. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, D.C.

Cadena C.D., Loiselle B.A., 2007. Limits to elevational distributions in two species of emberizine finches: disentangling the role of interspecific competition, autoecology, and geographic variation in the environment. Eco. 30, 491–504

Chamberlain, S., Barve, V., Mcglinn, D., Oldoni, D., Desmet, P., Geffert, L., Ram, K., 2024. rgbif: Interface to the global biodiversity information facility API.

Chamberlain, S.A., Boettiger, C., 2017. R Python, and Ruby clients for GBIF species occurrence data. https://doi.org/10.7287/peerj.preprints.3304v1.

Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K., Ellison, A.M., 2014. Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies. Ecol. Monogr. 84, 45–67. https://doi.org/10.1890/13-0133.1.

Cicero, C., Koo, M.S., Braker, E., Abbott, J., Bloom, D., Campbell, M., Cook, J.A., Demboski, J.R., Doll, A.C., Frederick, L.M., Linn, A.J., Mayfield-Meyer, T.J., McDonald, D.L., Nachman, M.W., Olson, L.E., Roberts, D., Sikes, D.S., Witt, C.C., Wommack, E.A., 2024. Arctos: Community-driven innovations for managing natural and cultural history collections. PLOS ONE 19, e0296478. https://doi.org/10.1371/journal.pone.0296478.

Clark N.J., 2018. Phylogenetic uniqueness, not latitude, explains the diversity of avian blood

parasite communities worldwide. Glob Ecol Biogeogr 27, 744–755.

https://doi.org/10.1111/geb.12741.

Coblentz, D.D., Riitters, K.H., 2004. Topographic controls on the regional-scale biodiversity of the

south-western USA. J. Biogeogr. 31, 1125–1138. https://doi.org/10.1111/j.1365-

2699.2004.00981.x.

Combes, C., 2001. Parasitism: the ecology and evolution of intimate interactions. University of

Chicago Press, Chicago and London.

Combes, C., 1997. Fitness of parasites: Pathology and selection. Int. J. Parasitol. 27, 1–10.

https://doi.org/10.1016/S0020-7519(96)00168-3.

Curson, D.R., Goguen, C.B., 1998. Plumbeous Vireo (*Vireo plumbeus*), in: Poole, A., Gill, F.

(Eds.), The birds of North America, Inc., Philadelphia, PA.

Duval, L., Robert, V., Csorba, G., Hassanin, A., Randrianarivelojosia, M., Walston, J., Nhim, T.,

Goodman, S.M., Ariey, F., 2007. Multiple host-switching of Haemosporidia parasites in

bats. Malar. J. 6, 157. https://doi.org/10.1186/1475-2875-6-157.

Ellis, V.A., Bensch, S., 2018. Host specificity of avian haemosporidian parasites is unrelated among

sister lineages but shows phylogenetic signal across larger clades. Int. J. Parasitol. 48, 897–

902. https://doi.org/10.1016/j.ijpara.2018.05.005.

Ellis, V. A., Collins, M. D., Medeiros, M. C. I., Sari, E. H. R., Coffey, E. D., Dickerson, R. C.,

Lugarini, C., Stratford, J. A., Henry, D. R., Merrill, L., Matthews, A. E., Hanson, A. A.,

Roberts, J. R., Joyce, M., Kunkel, M. R., Ricklefs, R. E., 2015. Local host specialization,

host-switching, and dispersal shape the regional distributions of avian haemosporidian

parasites. Proc. Natl. Acad. Sci. 112(36), 11294–11299.

https://doi.org/10.1073/pnas.1515309112.

Ellis, V. A., Huang, X., Westerdahl, H., Jönsson, J., Hasselquist, D., Neto, J. M., Nilsson, J.,

Nilsson, J., Hegemann, A., Hellgren, O., Bensch, S., 2020. Explaining prevalence, diversity

and host specificity in a community of avian haemosporidian parasites. Oikos 129(9), 1314–

1329. https://doi.org/10.1111/oik.07280.

Fecchio, A., Bell, J.A., Williams, E.J., Dispoto, J.H., Weckstein, J.D., De Angeli Dutra, D., 2023.

Co-infection with *Leucocytozoon* and other haemosporidian parasites increases with latitude

and altitude in New World bird communities. Microb. Ecol. 86, 2838–2846.

https://doi.org/10.1007/s00248-023-02283-x.

Fecchio, A., Wells, K., Bell, J.A., Tkach, V.V., Lutz, H.L., Weckstein, J.D., Clegg, S.M., Clark,

N.J., 2019. Climate variation influences host specificity in avian malaria parasites. Ecol.

Lett. 22, 547–557. https://doi.org/10.1111/ele.13215.

Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. Int. J. Climatol. 37, 4302–4315. https://doi.org/10.1002/joc.5086.

Filion, A., Eriksson, A., Jorge, F., Niebuhr, C.N., Poulin, R., 2020. Large-scale disease patterns

explained by climatic seasonality and host traits. Oecologia 194, 723–733.

https://doi.org/10.1007/s00442-020-04782-x.

Fischer, S.E., 2020. Post-fledging and migration ecology of gray vireos (*Vireo vicinior*) and using

ArtScience to explore gender and identity. Master's thesis, The University of Toledo,

Toledo, OH, USA. https://doi.org/10.13140/RG.2.2.31260.41603.

Fischer, S.E., Granillo, K., Streby, H.M., 2022. Post-fledging survival, movements, and habitat

associations of gray vireos in New Mexico. Avian Conserv. Ecol. 17, art13.

https://doi.org/10.5751/ACE-02053-170113.

Fischer, S.E., Kramer, G.R., Baumann, M.J., Fahy, E.F., Granillo, K., Streby, H.M., 2025.

Migratory connectivity and potential nonbreeding sexual segregation in Gray Vireos (*Vireo*

*vicinior*). Wilson J. Ornithol. https://doi.org/10.1080/15594491.2025.2497654.

Fox, J., Weisberg, S., 2019. An R companion to applied regression, Third. ed. Sage, Thousand

Oaks, CA.

Freeman, B. G., Class Freeman, A. M., Hochachka, W. M., 2016. Asymmetric interspecific

aggression in New Guinean songbirds that replace one another along an elevational gradient.

Ibis 158(4), 726–737. https://doi.org/10.1111/ibi.12384.

Gage K.L., Burkot T.R., Eisen R.J., Hayes, E.B.,. 2008. Climate and vectorborne diseases. Am. J.

Prev. Med. 35, 436–450. https://doi.org/10.1016/j.amepre.2008.08.030.

Galen, S.C., Borner, J., Martinsen, E.S., Schaer, J., Austin, C.C., West, C.J., Perkins, S.L., 2018.

The polyphyly of *Plasmodium*: comprehensive phylogenetic analyses of the malaria

parasites (order Haemosporida) reveal widespread taxonomic conflict. R. Soc. Open Sci. 5,

171780. https://doi.org/10.1098/rsos.171780.

Galen, S.C., Witt, C.C., 2014. Diverse avian malaria and other haemosporidian parasites in Andean

house wrens: evidence for regional co-diversification by host-switching. J. Avian Biol. 45,

374–386. https://doi.org/10.1111/jav.00375.

González, A.D., Matta, N.E., Ellis, V.A., Miller, E.T., Ricklefs, R.E., Gutiérrez, H.R., 2014. Mixed

species flock, nest height, and elevation partially explain avian haemoparasite Prevalence in

Colombia. PLOS ONE 9, e100695. https://doi.org/10.1371/journal.pone.0100695.

Granthon, C., Williams, D.A., 2017. Avian malaria, body condition, and blood parameters in four

species of songbirds. Wilson J. Ornithol. 129, 492–508. https://doi.org/10.1676/16-060.1

Gyllenhaal, E.F., 2024. Crossing the ocean: the importance of gene flow in island evolution.

University of New Mexico, Albuquerque, NM, USA.

Hackett, S.J., Kimball, R.T., Reddy, S., Bowie, R.C.K., Braun, E.L., Braun, M.J., Chojnowski, J.L.,

Cox, W.A., Han, K.-L., Harshman, J., Huddleston, C.J., Marks, B.D., Miglia, K.J., Moore,

W.S., Sheldon, F.H., Steadman, D.W., Witt, C.C., Yuri, T., 2008. A phylogenomic study of

birds reveals their evolutionary history. Science 320, 1763–1767.

https://doi.org/10.1126/science.1157704.

Hasik, A.Z., Siepielski, A.M., 2022. Parasitism shapes selection by drastically reducing host fitness
and increasing host fitness variation. Biol. Lett. 18, 20220323.
https://doi.org/10.1098/rsbl.2022.0323.

Hellgren, O., Waldenström, J., Bensch, S., 2004. A new PCR assay for simultaneous studies of
*Leucocytozoon*, *Plasmodium*, and *Haemoproteus* from avian blood. J. Parasitol. 90, 797–
802.

Hellgren, O., Waldenström, J., Peréz-Tris, J., Szöll, E., Si, Ö., Hasselquist, D., Krizanauskiene, A.,
Ottosson, U., Bensch, S., 2007. Detecting shifts of transmission areas in avian blood
parasites — a phylogenetic approach. Mol. Ecol. 16, 1281–1290.
https://doi.org/10.1111/j.1365-294X.2007.03227.x.

Hsieh, T.C., Ma, K.H., Chao, A., 2024. iNEXT: iNterpolation and EXTrapolation for species
diversity.

Huang, X., Jönsson, J., Bensch, S., 2020. Persistence of avian haemosporidians in the wild: a case
study to illustrate seasonal infection patterns in relation to host life stages. Int. J. Parasitol.
50, 611–619. https://doi.org/10.1016/j.ijpara.2020.05.006.

Hubbard, J.P., 1970. Check-list of the birds of New Mexico. N. M. Ornithol. Soc. 3, 108.

Illera, J.C., López, G., García-Padilla, L., Moreno, Á., 2017. Factors governing the prevalence and
richness of avian haemosporidian communities within and between temperate mountains.
PLOS ONE 12, e0184587. https://doi.org/10.1371/journal.pone.0184587.

Ikemoto T., 2008. Tropical malaria does not mean hot environments. J. Med. Entomol. 45(6), 963–
969.

Ishtiaq, F., Barve, S., 2018. Do avian blood parasites influence hypoxia physiology in a high
elevation environment? BMC Ecol. 18. https://doi.org/10.1186/s12898-018-0171-2.

Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K., Mooers, A.O., 2012. The global diversity of birds
in space and time. Nature 491, 444–498. https://doi.org/10.1038/nature11631.

Kassambara, A., 2023. rstatix: Pipe-friendly framework for basic statistical tests. R package version

0.7.2. https://CRAN.R-project.org/package=rstatix.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper,

A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012.

Geneious basic: An integrated and extendable desktop software platform for the

organization and analysis of sequence data. Bioinformatics 28, 1647–1649.

https://doi.org/10.1093/bioinformatics/bts199.

Kim, K.S., Tsuda, Y., 2010. Seasonal changes in the feeding pattern of *Culex pipiens pallens*

govern the transmission dynamics of multiple lineages of avian malaria parasites in

Japanese wild bird community: vector study of avian malaria transmission. Mol. Ecol. 19,

5545–5554. https://doi.org/10.1111/j.1365-294X.2010.04897.x.

Krasnov, B.R. and Poulin, R., 2010. Ecological properties of a parasite: species-specific stability

and geographical variation. The biogeography of host–parasite interactions. Oxford

University Press, Oxford, pp. 99-113.

Kuhn, M., 2008. Building Predictive Models in *R* Using the caret Package. J. Stat. Softw. 28.

https://doi.org/10.18637/jss.v028.i05.

Kushary, D., 2000. Bootstrap methods and their application. Technometrics 42, 216–217.

https://doi.org/10.1080/00401706.2000.10486018.

LaPointe, D.A., Atkinson, C.T., Samuel, M.D., 2012. Ecology and conservation biology of avian

malaria. Ann. N. Y. Acad. Sci. 1249, 211–226. https://doi.org/10.1111/j.1749-

6632.2011.06431.x.

Lauron, E.J., Loiseau, C., Bowie, R.C.K., Spicer, G.S., Smith, T.B., Melo, M., Sehgal, R.N.M.,

2015. Coevolutionary patterns and diversification of avian malaria parasites in African

sunbirds (Family Nectariniidae). Parasitology 142, 635–647.

https://doi.org/10.1017/S0031182014001681.

Lele, S.R., Keim, J.L., Solymos, P., 2023. ResourceSelection: resource selection (probability)

functions for use-availability data. https://CRAN.R-project.org/package=ResourceSelection.

Loiseau, C., Harrigan, R.J., Bichet, C., Julliard, R., Garnier, S., Lendvai, Á.Z., Chastel, O., Sorci,

G., 2013. Predictions of avian *Plasmodium* expansion under climate change. Sci. Rep. 3,

1126. https://doi.org/10.1038/srep01126.

Lutz, H.L., Hochachka, W.M., Engel, J.I., Bell, J.A., Tkach, V.V., Bates, J.M., Hackett, S.J.,

Weckstein, J.D., 2015. Parasite prevalence corresponds to host life history in a diverse

assemblage of Afrotropical birds and haemosporidian parasites. PLOS ONE 10, e0121254.

https://doi.org/10.1371/journal.pone.0121254.

Mangiafico, S.S., 2024. rcompanion: Functions to Support Extension Education Program

Evaluation. R package version 2.4.36. https://CRAN.R-project.org/package=rcompanion.

Marroquin-Flores, R.A., Williamson, J.L., Chavez, A.N., Bauernfeind, S.M., Baumann, M.J.,

Gadek, C.R., Johnson, A.B., McCullough, J.M., Witt, C.C., Barrow, L.N., 2017. Diversity,

abundance, and host relationships in the avian malaria community of New Mexico pine

forests. PeerJ 5, e3700–e3700. https://doi.org/10.7717/peerj.3700.

McNew, S.M., Barrow, L.N., Williamson, J.L., Galen, S.C., Skeen, H.R., DuBay, S.G., Gaffney,

A.M., Johnson, A.B., Bautista, E., Ordoñez, P., Schmitt, C.J., Smiley, A., Valqui, T., Bates,

J.M., Hackett, S.J., Witt, C.C., 2021. Contrasting drivers of diversity in hosts and parasites

across the tropical Andes. Proc. Natl. Acad. Sci. 118, e2010714118.

https://doi.org/10.1073/pnas.2010714118.

Medeiros, M.C.I., Hamer, G.L., Ricklefs, R.E., 2013. Host compatibility rather than vector–host-

encounter rate determines the host range of avian *Plasmodium* parasites. Proc. R. Soc. B

Biol. Sci. 280, 20122947. https://doi.org/10.1098/rspb.2012.2947.

Minchella, D.J., Scott, M.E., 1991. Parasitism: A cryptic determinant of animal community

structure. Trends Ecol. Evol. 6, 250–254. https://doi.org/10.1016/0169-5347(91)90071-5.

Mordecai, E. A., Paaijmans, K. P., Johnson, L. R., Balzer, C., Ben-Horin, T., De Moor, E., McNally, A., Pawar, S., Ryan, S. J., Smith, T. C., & Lafferty, K. D., 2013. Optimal temperature for malaria transmission is dramatically lower than previously predicted. Ecol. Lett. 16(1), 22–30. https://doi.org/10.1111/ele.12015.

Mukhin, A., Palinauskas, V., Platonova, E., Kobylkov, D., Vakoliuk, I., Valkiūnas, G., 2016. The strategy to survive primary malaria infection: an experimental study on behavioural changes in parasitized birds. PLOS ONE 11, e0159216. https://doi.org/10.1371/journal.pone.0159216.

Muriel, J., 2020. Ecophysiological assessment of blood haemosporidian infections in birds. Ecosistemas 29. https://doi.org/10.7818/ecos.1979.

Names, G.R., Schultz, E.M., Krause, J.S., Hahn, T.P., Wingfield, J.C., Heal, M., Cornelius, J.M., Klasing, K.C., Hunt, K.E., 2021. Stress in paradise: effects of elevated corticosterone on immunity and avian malaria resilience in a Hawaiian passerine. J. Exp. Biol. 224. https://doi.org/10.1242/jeb.242951.

Oakgrove, K.S., Harrigan, R.J., Loiseau, C., Guers, S., Seppi, B., Sehgal, R.N.M., 2014. Distribution, diversity and drivers of blood-borne parasite co-infections in Alaskan bird populations. Int. J. Parasitol. 44, 717–727. https://doi.org/10.1016/j.ijpara.2014.04.011.

Pérez-Rodríguez, A., Fernández-González, S., De La Hera, I., Pérez-Tris, J., 2013. Finding the appropriate variables to model the distribution of vector-borne parasites with different environmental preferences: climate is not enough. Glob. Change Biol. 19, 3245–3253. https://doi.org/10.1111/gcb.12226.

Perkins, S.L., 2014. Malaria's many mates: past, present, and future of the systematics of the order Haemosporida. J. Parasitol. 100, 11–25. https://doi.org/10.1645/13-362.1.

Pigeault, R., Chevalier, M., Cozzarolo, C.S., Baur, M., Arlettaz, M., Cibois, A., Keiser, A., Guisan, A., Christe, P., Glaizot, O., 2022. Determinants of haemosporidian single- and co-infection

risks in western palearctic birds. Int. J. Parasitol. 52, 617–627.

https://doi.org/10.1016/j.ijpara.2022.05.002.

Poulin, R., 2011. Evolutionary ecology of parasites. Princeton University Press, Princeton, New

Jersey.

Poulin, R., Krasnov, B. R., 2010. Similarity and variability of parasite assemblages across

geographical space. The biogeography of host–parasite interactions. Oxford University

Press, Oxford, pp. 115-127.

R Core Team, 2023. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reinoso-Pérez, M.T., Dhondt, K.V., Dulcet, H., Katzenstein, N., Sydenstricker, A.V., Dhondt,

A.A., 2024. Seasonal variation in detection of Haemosporidia in a bird community: a

comparison of nested PCR and microscopy. J. Wildl. Dis. 60. https://doi.org/10.7589/JWD-

D-23-00023.

Ricklefs, R.E., Fallon, S.M., 2002. Diversification and host switching in avian malaria parasites.

Proc. R. Soc. Lond. B Biol. Sci. 269, 885–892. https://doi.org/DOI 10.1098/rspb.2001.1940.

Ricklefs, R.E., Fallon, S.Y.M., Bermingham, E., 2004. Evolutionary relationships, cospeciation,

and host switching in avian malaria parasites. Syst. Biol. 53.

Rodriguez, M.D., Doherty, P.F., Piaggio, A.J., Huyvaert, K.P., 2021. Sex and nest type influence

avian blood parasite prevalence in a high-elevation bird community. Parasit. Vectors 14,

145. https://doi.org/10.1186/s13071-021-04612-w.

Rodríguez-Hernández, K., Álvarez-Mendizábal, P., Chapa-Vargas, L., Escobar, F., González-

García, F., Santiago-Alarcon, D., 2021. Haemosporidian prevalence, parasitaemia and

aggregation in relation to avian assemblage life history traits at different elevations. Int. J.

Parasitol. 51, 365–378. https://doi.org/10.1016/j.ijpara.2020.10.006.

Santiago-Alarcon, D., Palinauskas, V., Schaefer, H.M., 2012. Diptera vectors of avian

Haemosporidian parasites: untangling parasite life cycles and their taxonomy. Biol. Rev. 87,

928–964. https://doi.org/10.1111/j.1469-185X.2012.00234.x.

Schrader, M.S., Walters, E.L., James, F.C., Greiner, E.C., 2003. Seasonal prevalence of a

haemotozoan parasite of red-bellied woodpeckers (*Melanerpes carolinus*) and its association

with host condition and overwinter survival. The Auk 120, 130.

https://doi.org/10.1642/0004-8038.

Sheppard, P., Comrie, A., Packin, G., Angersbach, K., Hughes, M., 2002. The climate of the US

Southwest. Clim. Res. 21, 219–238. https://doi.org/10.3354/cr021219.

Sibley, C.G., Monroe, J., 1990. Distribution and taxonomy of birds of the world. Yale University

Press, New Haven, CT.

Slowinski, S.P., Geissler, A.J., Gerlach, N., Heidinger, B.J., Ketterson, E.D., 2022. The probability

of being infected with haemosporidian parasites increases with host age but is not affected

by experimental testosterone elevation in a wild songbird. J. Avian Biol. 2022, jav.02819.

https://doi.org/10.1111/jav.02819.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics 30, 1312–1313. https://doi.org/10.1093/bioinformatics/btu033.

Stöver, B.C., Müller, K.F., 2010. TreeGraph 2: combining and visualizing evidence from different

phylogenetic analyses. BMC Bioinformatics 11, 7. https://doi.org/10.1186/1471-2105-11-7.

Sundar D., 2006. binom: Binomial confidence intervals for several parameterizations. R package

version 1.1-1.1.  https://CRAN.R-project.org/package=binom.

Tang, Y., Horikoshi, M., Li, W., 2016. ggfortify: Data visualization tools for statistical analysis

results. https://CRAN.R-project.org/package=ggfortify.

Valkiūnas, G., 2005. Avian malaria parasites and other Haemosporidia, 1st ed. CRC Press, Boca

Raton.

von Humboldt A., Bonpland A., 2009. Essay on the geography of plants—with a physical tableau

of the equinoctial regions (1807). In: Jackson ST (ed) Essay on the geography of plants.

University of Chicago Press, Chicago, pp 61–155.

Waldenström, A.J., Bensch, S., Hasselquist, D., Östman, Ö., Pryor, G.S., Greiner, E.C., Hall, B.,

2004. A new nested polymerase chain reaction method very efficient in detecting

Plasmodium and Haemoproteus infections from avian blood. J. Parasitol. 90, 191–194.

https://doi.org/10.1645/GE-3221RN.

Walther, E.L., Carlson, J.S., Cornel, A., Morris, B.K., Sehgal, R.N.M., 2016. First molecular study

of prevalence and diversity of avian haemosporidia in a Central California songbird

community. J. Ornithol. 157, 549–564. https://doi.org/10.1007/s10336-015-1301-7.

Wickham, H., 2016. ggplot2, Use R! Springer International Publishing, Cham.

https://doi.org/10.1007/978-3-319-24277-4.

Williamson, J.L., Witt, C.C., 2021. Elevational niche-shift migration: Why the degree of elevational

change matters for the ecology, evolution, and physiology of migratory birds. Ornithology

138, ukaa087. https://doi.org/10.1093/ornithology/ukaa087.

Williamson, J.L., Wolf, C.J., Barrow, L.N., Baumann, M.J., Galen, S.C., Schmitt, C.J., Schmitt,

D.C., Winter, A.S., Witt, C.C., 2019. Ecology, not distance, explains community

composition in parasites of sky-island Audubon's Warblers. Int. J. Parasitol. 49, 437–448.

https://doi.org/10.1016/j.ijpara.2018.11.012.

Wyckoff, S.T., Judkins, T.C., Nemeth, N.M., Ruder, M.G., Martin, J.A., Kunkel, M.R., Garrett,

K.B., Adcock, K.G., Mead, D.G., Yabsley, M.J., 2024. Surveillance for selected pathogens

and parasites of northern bobwhite (*Colinus virginianus*) from western Oklahoma, USA.

2018–20. J. Wildl. Dis. 60. https://doi.org/10.7589/jwd-d-23-00102.

Zamora-Vilchis, I., Williams, S. E., & Johnson, C. N., 2012. Environmental temperature affects

prevalence of blood parasites of birds on an elevation gradient: implications for disease in a

warming climate. PLOS ONE 7(6), e39208. https://doi.org/10.1371/journal.pone.0039208

DATA STATEMENT

Specimen and parasite data are available in Supplementary Tables S1–S11 and are searchable via

Arctos (arctos.database.museum; Cicero et al., 2024). Parasite sequences are archived in GenBank

(accession nos. PV295634-PV295830). Vouchered specimens and frozen tissue samples are

archived at the Museum of Southwestern Biology Division of Birds and Division of Genomic

Resources, respectively, where they are available for further study (https://msb.unm.edu/). All R

code and datasets are available on FigShare (https://doi.org/10.6084/m9.figshare.28585943).

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest to declare.

ETHICS STATEMENT

Research activities were conducted under Institutional Animal Care and Use Protocol 16-200406-

MC and appropriate state and federal scientific collecting permits (New Mexico Department of

Game and Fish Authorization Number 3217; US Fish and Wildlife Permit Numbers MB094297-0

and MB0942970-1).

FUNDING STATEMENT

This work was supported by the National Science Foundation (Graduate Research Fellowship NSF-

DGE-2439853 to DLFW; Postdoctoral Research Fellowship in Biology NSF-DBI-1611710 to

LNB; and NSF-DEB-1146491 to CCW), the Bureau of Land Management Rio Puerco Field Office

(via the Colorado Plateau Cooperative Eco-systems Studies Unit agreement), New Mexico

Department of Game and Fish Share with Wildlife Grant to HMS/SEF, and a Sevilleta LTER

Graduate Summer Research Fellowship and New Mexico Ornithological Society Ryan Beaulieu

Research Grant to SEF.

TABLES

| Top five infection probability fixed effects additive models | | | | | |
|---|---|---|---|---|---|
| Bell's Vireo (*n* = 20) | | | | | |
| *Rank* | *Predictors* | *K* | *AICc* | *ΔAICc* | *Weight* |
| *1* | tempPC2 + precipPC1 | 3 | 22.6 | - | 0.324 |
| *2* | tempPC1 | 2 | 23.6 | *0.967* | 0.200 |
| *3* | precipPC1 | 2 | 24.7 | *0.348* | 0.113 |
| *4* | tempPC1 + precipPC1 | 3 | 25.9 | *0.199* | 0.064 |
| *5* | Elevation + tempPC1 | 3 | 25.9 | *0.197* | 0.064 |
| Gray Vireo (*n* = 170) | | | | | |
| *Rank* | *Predictors* | *K* | *AICc* | *ΔAICc* | *Weight* |
| *1* | Elevation + tempPC1 | 3 | 204 | - | 0.419 |
| *2* | Elevation + precipPC1 + tempPC1 | 4 | 206 | *1.72* | 0.177 |
| *3* | Elevation + precipPC2 + tempPC1 | 4 | 206 | *1.73* | 0.177 |
| *4* | Elevation + tempPC1 + tempPC2 | 4 | 206 | *2.07* | 0.149 |
| *5* | Elevation + tempPC1 + tempPC2, precipPC1 + precipPC2 | 5 | 207 | *3.35* | 0.078 |
| Plumbeous Vireo (*n* = 58) | | | | | |
| *Rank* | *Predictors* | *K* | *AICc* | *ΔAICc* | *Weight* |
| *1* | Null model | 1 | 52.1 | - | 0.301 |
| *2* | precipPC2 | 2 | 52.4 | *0.409* | 0.246 |
| *3* | precipPC1 | 2 | 53.3 | *1.17* | 0.168 |
| *4* | tempPC2 | 2 | 53.4 | *1.29* | 0.158 |
| *5* | PrecipPC1 + precipPC2 | 2 | 53.9 | *1.73* | 0.127 |

991 **Table 1. Top five models assessing environmental predictors of haemosporidian infection for**

992 **each species.** Sample sizes for datasets and directionality of best model outputs are listed at the top.

993 Models were ranked by AICc, with ΔAICc values relative to the top model and corresponding

994 Akaike weights (Weight) indicating model support.

|  | Bell's Vireo top models | | | | Gray Vireo top models | | | | Plumbeous Vireo top models | | | |
|  | Model-averaged estimate and 95% confidence limits | | | | Model-averaged estimate and 95% confidence limits | | | | Model-averaged estimate and 95% confidence limits | | | |
| Parameter | β | SE | Lower CL | Upper CL | β | SE | Lower CL | Upper CL | β | SE | Lower CL | Upper CL |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Intercept | 0.845 | 0.876 | -0.989 | 2.68 | 1.14 | 0.357 | 0.433 | 1.84 | 1.74 | 0.377 | 0.983 | 2.49 |
| Elevation | 0.062 | 0.371 | -0.703 | 0.827 | -1.72 | 0.763 | -3.23 | -0.214 | - | - | - | - |
| tempPC1 | -0.938 | 1.44 | -3.85 | 1.98 | 1.08 | 0.546 | 0.006 | 2.16 | - | - | - | - |
| tempPC2 | 1.80 | 2.65 | -3.54 | 7.14 | -0.048 | 0.717 | -1.46 | 1.37 | 0.051 | 0.179 | -0.305 | 0.406 |
| precipPC1 | 2.83 | 3.47 | -4.15 | 9.80 | -0.163 | 0.588 | -1.32 | 0.996 | 0.099 | 0.24 | -0.378 | 0.577 |
| precipPC2 | NA | NA | NA | NA | -0.278 | 0.901 | -2.05 | 1.50 | 0.172 | 0.309 | -0.439 | 0.783 |

**Table 2. Standardized model-averaged regression coefficients used to estimate effects of**

**predictors and precision of effects across species-specific candidate model sets that assess**

**environmental predictors of haemosporidian infection.** Regression coefficients (β), standard

error (SE), and 95% confidence limits (CL) are given. Dashes indicate that the parameter was not

present in a final model set, and NA values indicate that the parameter was not tested due to lack of

variation present in model set.

1003

1005

**Figure 1)** A) Sampling distribution for each host species (Bell's Vireo: *Vireo bellii,* Gray Vireo: *V.*

*vicinior,* and Plumbeous Vireo: *V. plumbeus*) in the southwestern USA; B) Left: summary of

infection prevalence, with significant differences between groups determined using multiple

Pearson chi-squared tests and Bonferroni-corrected p-values indicated by letters. Error bars

represent 95% confidence intervals for positive infection proportions. Right: percent parasitemia for

Gray and Plumbeous vireos. C) Proportion of individuals infected versus uninfected across the

1012    sampled elevational gradient. Asterisk denotes significant p-values (< 0.05) based on Mann-

1013    Whitney U-tests; D) Phylogeny of 19 haemosporidian cyt *b* haplotypes and their respective

1014    abundances per species. Stars denote novel haplotypes and bootstrap values >50 are shown at

1015    nodes. Bird images courtesy Birds of the World (Billerman et al., 2025).

1016

1017

**Figure 2)** Estimate of haemosporidian lineage diversity in Bell's (*Vireo belli*), Gray (*V. vicinior*)

and Plumbeous Vireos (*V. plumbeus*) based on iNEXT rarefaction and extrapolation. Bell's Vireos

curves, shown in yellow, were created using 12 avian haemosporidian infections and three

haplotypes; Gray Vireo curves, shown in gray, were created using 123 avian haemosporidian

infections and four haplotypes; Plumbeous Vireo curves, shown in blue, used 62 haemosporidian

infections and 14 haplotypes. Points indicate the reference sample, solid line the rarefaction

estimate, and dotted line the extrapolation. The analysis suggested that sampling approximately 33

total infections in Bell's, 14 total infections for Gray, and 364 total infections for Plumbeous Vireos

would capture >95% of the haemosporidian lineage diversity in this community. The total

1027     haplotype richness, rounded to nearest whole number, is estimated to be four for Bell's Vireos (95%

1028     CI [3-6]), five for Gray Vireos (95% CI [4 – 6]) and 54 for Plumbeous Vireos (95% CI [14 – 101]).

1029

1030

1031     **Figure 3)** Intraspecific distribution of infected and uninfected vireos across associated

1032     environmental variables. Higher tempPC1 values relate to cooler overall temperatures and higher

1033     precipPC2 relates to sites with more seasonally stable precipitation. Significantly different

1034     distributions, determined by Mann-Whitney U-tests are denoted by an asterisk ($p < 0.05$).

1035