Parallel elevational replacement of hosts and parasites in a highly susceptible songbird genus

[1]Daniele L. F. Wiley[1,2], Jessie L. Williamson[1,3], Silas E. Fischer[4], Selina M. Bauernfeind[1], Henry

M. Streby[4], Kathy Granillo[5], Christopher C. Witt[1,2], and Lisa N. Barrow[1,2]

[1]Museum of Southwestern Biology, University of New Mexico, 302 Yale Blvd NE, Albuquerque,

NM 87131, USA

[2]Department of Biology, University of New Mexico, 302 Yale Blvd NE, Albuquerque, NM 87131,

USA

[3]Department of Zoology and Physiology, University of Wyoming, 1000 E. University Ave, Laramie,

WY 82071, USA

[4]Department of Environmental Sciences, University of Toledo, Wolfe Hall Room 1235, 3050 West

Towerview Blvd., Toledo, OH 43606-3390, USA

[5]United States Fish and Wildlife Service, Socorro, NM 87801, USA; retired

*Corresponding author: Daniele L. F. Wiley, dwiley7@unm.edu

Address: Department of Biology, University of New Mexico, Castetter Hall 1480, MSC03-2020,

219 Yale Blvd. NE, Albuquerque, NM 87131, USA

---

[1]Authors' Contributions: DLFW, JLW, CCW, and LNB conceptualized and developed methodology for this project. JLW, SEF, HMS, SMB, KG, CCW, and LNB performed fieldwork and subsequent data curation. DLFW, JLW, and SMB performed molecular analysis and generated sequence data. JLW, SEF, HMS, KG, CCW, and LNB secured funding and provided supervision. DLFW and JLW performed formal analyses and created visualizations. DLFW and JLW wrote the original draft of the manuscript, and DLFW wrote the final version with input from all co-authors.

Abstract

Elevational replacement distribution patterns underpin montane diversity and reflect the interaction of both biotic and abiotic pressures, but the degree to which parasites exhibit elevational zonation remains unclear. Investigating infection patterns in related host species across elevational gradients can reveal whether parasites and hosts show concordant patterns of elevational turnover, potentially due to shared historical and ecological factors. Here, we assessed patterns of elevational replacement in haemosporidian parasite assemblages that infect three congeneric songbird species: Bell's vireo (*Vireo bellii),* gray vireo *(V. vicinior),* and plumbeous vireo (*V. plumbeus*), each of which breeds across distinct elevations and habitats in the southwestern United States. We screened a total of 248 individuals using cytochrome *b* PCR and microscopy. We identified 19 haemosporidian haplotypes, including eight novel lineages. We found that each of the three vireo species exhibited high haemosporidian prevalence (55.0–86.2%), with nearly all infections from the genus *Haemoproteus* (subgenus *Parahaemoproteus*). Haemosporidian assemblages varied across elevations; each sampled range of elevations harbored abundant, yet host-specific lineages with different environmental associations. Bell's and plumbeous vireos, but not gray vireos, hosted several phylogenetically distinct, putative generalist lineages, likely reflecting spillover from more diverse local breeding bird communities. Repeated infections in individuals across breeding seasons, together with moderate parasitemia ($\bar{x} \approx 1\%$) suggest that these focal vireo species harbor chronic infections during their respective breeding seasons. These results demonstrate that elevational replacement patterns in avian hosts may be mirrored by their haemosporidian parasites, particularly among host-specialized lineages.

Keywords: Avian malaria, diversity, elevational range, haemosporidian, turnover

Introduction

Mountains harbor exceptional biodiversity, in part because elevational gradients tend to be finely partitioned by related species with ranges that are elevationally parapatric (Cadena and Loiselle, 2007; von Humboldt and Bonpland, 2009; Freeman et al., 2016). Yet, it remains unclear whether parasites of elevational replacement host species mirror this range stratification. Generally, host-parasite interactions can strongly influence biogeographic patterns, with reciprocal effects on fitness (Combes, 1997, 2001; Poulin, 2011; Hasik and Siepielski, 2022), population dynamics (Albon et al., 2002), and community structure (Minchella and Scott, 1991, McNew et al., 2021). Strong host-specificity, combined with shared environmental constraints, should result in parasite turnover that parallels the turnover of hosts along environmental gradients. Conversely, aspects of parasite life-history—such as insulation from climatic factors (e.g., endoparasites), dependency on secondary hosts and vectors (e.g., complex life cycle), and degree of host-specificity (e.g., host range and degree of host switching)—should be expected to cause discordant patterns from those observed in hosts (Gage et al. 2008; Krasnov and Poulin, 2010; Ellis et al., 2015).

Regional surveys across elevational gradients reflect these complex relationships. In various mountain systems, parasite turnover has been found to exceed host turnover (Galen and Witt, 2014; Barrow et al., 2021), and host turnover has been found to predict parasite turnover (Ellis et al., 2015; Williamson et al., 2019). Gradients of precipitation and temperature, as well as landscape features, have also been found to shape parasite communities (Illera et al., 2017; McNew et al., 2021). Investigating the parasite communities of closely related host taxa that are elevational replacements provides a natural experiment for disentangling the relative roles of host associations, parasite traits, and environmental gradients in shaping parasite biogeography, while controlling for host phylogeny, life history, and parasite susceptibility (Medeiros et al., 2013; Barrow et al., 2019).

Among blood parasites, haemosporidians (order Haemosporida; Danilewsky, 1885) are a cosmopolitan group of protozoans with complex life cycles dependent upon vertebrate taxa, including birds, mammals, and reptiles (Ricklefs and Fallon, 2002; Duval et al., 2007; Boysen et al.,

2022), and invertebrate vectors in the order Diptera (Linnaeus, 1758). A long history of co-

evolution with avian hosts (Ricklefs et al., 2004; Lauron et al., 2015; Galen et al., 2018) has given

rise to a remarkable diversity of avian haemosporidian lineages (Valkiūnas, 2005; Hellgren et al.,

2007; Perkins, 2014), with many still yet to be described (Borner et al., 2016). This diversity is

reflected both phylogenetically and phenotypically, with variable degrees of pathogenicity

(reviewed in Bennett et al., 1993) and host specificity (Ellis and Bensch, 2018) recorded among and

within genera.

Given the high diversity and complexity of haemosporidian parasites, it has remained

challenging to characterize the distribution patterns of haemosporidian lineages across large

geographic scales (Clark, 2018). However, within regions and genera, spatial trends appear—

reflecting the influence of climate directly, through physiological interactions during transmission

and development (Ikemoto 2008; Mordecai et al., 2013), and indirectly, by influencing host and

vector abundance and susceptibility (Filion et al., 2020). As a result, the fluctuating climate of

temperate and montane regions produce seasonal, annual, and habitat-level waves of infection

(Bensch et al., 2007; Pérez☐Rodríguez et al., 2013; Lutz et al., 2015; Reinoso-Pérez et al., 2024),

whereas areas with higher, less seasonal temperatures, such as tropical regions and lowlands, can

have persistent haemosporidian pressure (Zamora-Vilchis et al., 2012).

Infection intensity, commonly measured as parasitemia (defined as the 'estimated percentage of

haemosporidians circulating in host blood') provides insight into host condition and stage of

haemosporidian infection (i.e., acute or chronic; Valkiūnas, 2005). Like prevalence (here defined as

'proneness to infection'), parasitemia is also influenced by environmental factors, with peaks in

temperate regions during the breeding season, when warmer, wetter conditions favor vector activity

(Valkiūnas, 2005; Reinoso-Pérez et al., 2024) and when chronically infected hosts experience

tissue-to-blood recrudescence (Atkinson and van Riper, 1991). Therefore, montane regions provide

a valuable system for studying avian haemosporidian infection dynamics because of the sharp local

variation in both climate and habitat over relatively small geographic distance (e.g., Illera et al.,

2017; Rodríguez-Hernández et al., 2021; Williamson et al., 2019).

The southwestern U.S. presents a particularly compelling region for studying haemosporidian

biogeography and dynamics due to its climatic heterogeneity, wide range of habitat types, and steep

elevational gradients in sky island mountains. Although arid overall, the region exhibits strong

variation in precipitation timing and intensity, shaped by both temperate seasonality and the North

American monsoon system. These climatic patterns interact with topographic variation (reviewed in

Coblentz and Riitters, 2004) to create a mosaic of ecologically distinct elevational zones ranging

from lowland riparian corridors with high relative humidity and permanent water sources, to mid-

elevation desert scrublands and savannas reliant on seasonal rainfall, to cooler montane forests with

shorter growing seasons and higher annual precipitation (Sheppard et al., 2002). While community-

level surveys suggest moderate overall prevalence of haemosporidian infection among avian

communities in the region (mean: ~36%; Barrow et al., 2021), and while environmental conditions–

–particularly related to elevation, have been shown to strongly influence spatial patterns of infection

(Williamson et al., 2019)—previous studies have primarily described environmental and ecological

contributions across mid-high elevation habitats and sky islands. Therefore, further work is needed

to expand our understanding of whether observed environmental and elevational trends persist

across broader gradients and habitat types, especially among particularly susceptible hosts.

Among the many avian species previously surveyed in the desert Southwest, the plumbeous

vireo (*Vireo plumbeus*) has consistently stood out for its high haemosporidian prevalence across

sites and years (Marroquin-Flores et al., 2017; Barrow et al., 2021). Migratory populations of this

insectivorous songbird breed in mid- to high-elevation coniferous woodlands (from ~1,200–3,000

m) across the western U.S. and Mexico (Barlow, 1977; Curson and Goguen, 1998) and spend the

nonbreeding season along the Pacific Slope and in the lowlands of central Mexico (Sibley and

Monroe, 1990). Two closely related migratory vireos, Bell's vireo (*Vireo bellii*) and gray vireo

(*Vireo vicinior*), have relatively similar breeding distributions, yet inhabit distinct habitats and

elevations. gray vireos breed in arid mid-elevation juniper (*Juniperus* spp.) savannas and chaparral,

~400–1,900 m (Hubbard, 1970; Barlow, 1977) and spend the nonbreeding season primarily in

lowland desert and coastal areas of Sonora and Baja California (Barlow et al., 1999; Fischer et al.,

2025). Bell's vireos breed in lowland riparian habitats, generally below ~1,500 m, with

comparatively high access to water and vegetative cover (Brown, 1993) and spend the nonbreeding

season along the Pacific coast of Mexico, from Baja California and Sonora south to El Salvador.

Given the similarities in ecology and life history among vireos, local differences in breeding

elevational range make these three focal species an ideal comparative system through which to

examine haemosporidian infection patterns across elevational gradients.

In this study, we tested whether haemosporidian biogeography, community assemblage, and

infection dynamics vary predictably among three species of vireos that occupy distinct elevations

and habitats at sites in New Mexico and Utah, USA. Using a combination of molecular screening

and microscopy, we (1) compared parasite prevalence and infection intensity among host species,

including the first parasite survey for regionally vulnerable gray vireos; (2) characterized

haemosporidian haplotypes and host specificity; and (3) tested the extent to which community

composition was linked to elevational zones and/or their associated host species and habitats. We

predicted finding extensive parasite sharing between elevational replacement vireo species,

considering the proximity and interdigitated nature of their ranges. Based on previously published

haemosporidian surveys in vireos, we anticipated finding high infection prevalence across all three

host species; however, we also expected prevalence and parasitemia to vary among species and

environments. Lastly, because the Southwest is arid, we predicted higher prevalence and parasite

diversity in more mesic habitats suitable for dipterid vectors, such as in the low-elevation riparian

corridors used by Bell's vireos and high-elevation woodlands occupied by plumbeous vireos.

Materials and methods

Sample collection

We sampled 248 wild-caught individuals of three vireo species (Bell's vireo: $n = 20$, gray

vireo: $n = 170$, and plumbeous vireo: $n = 58$; Fig. 1A; Table S1). Sampled individuals included

adults and juveniles (Table S1). All individuals were sampled during the summer months (May–

August), with the majority sampled prior to monsoon season (~early-July–September) from 1995–

2019 across 28 unique sites in New Mexico and a single site in Utah. During 2017–2019, targeted

fieldwork was conducted to sample gray vireos: most individuals were captured in mist nets, with

the exception of hatch-year birds sampled at nests. Blood samples were taken from the brachial vein

and stored in lysis buffer, and individuals were released at capture sites for ongoing study of gray

vireo annual cycle ecology (Fischer, 2020; Fischer et al., 2022; Fischer et al., 2025). In 2018 only,

we prepared and air-dried blood smears in the field ($n = 53$). We also recaptured and collected a

second whole blood sample from fourteen gray vireos (approximately one year after their initial

sample collection in 2017), yielding two total whole blood samples and one smear for this subset of

individuals (Table S2). Additional gray vireo tissues for four individuals were deposited at the

Museum of Southwestern Biology (hereafter, MSB) at the University of New Mexico following

protocols approved by the UNM Institutional Animal Care and Use Committee. Samples for Bell's

and plumbeous vireos were requested as tissue loans from the MSB. Detailed sampling protocol for

Bell's and plumbeous vireos are described in (Marroquin-Flores et al., 2017; Barrow et al., 2021;

Gyllenhaal, 2024).

All sampled birds either had specific coordinates recorded at the time of capture or were

caught within 100 m of an existing sampling location. For the latter, we assigned geographically

proximate coordinates based on the nearest documented sampling site or, when birds were captured

between two closely spaced sites, by taking the midpoint between them (Table S3).

Parasite screening and genetic data collection

We extracted DNA from pectoral muscle ($n = 82$) and whole blood ($n = 166$) using the

QIAGEN DNeasy Blood and Tissue Kit following manufacturer's recommendations. Birds were

screened for *Haemoproteus* and *Plasmodium* parasites using three nested polymerase chain reaction

(PCR) protocols to amplify a 478-base pair (bp) fragment of the haemosporidian mitochondrial cyt

b gene (Hellgren et al., 2004; Waldenström et al., 2004). We amplified parasite DNA with the outer

primer pairs HaemNFI/HaemNR3 and HaemNF/HaemNR2, followed by the nested primer pair

HaemF/HaemR2. We prepared outer PCR reactions in 25 µL volumes, containing 1.25 U of

AmpliTaq Gold DNA Polymerase (Applied Biosystems, Foster City, CA, USA), 1× PCR Buffer II,

2.5 mM MgCl2, 0.2 mM dNTPs, 0.5 µM of each primer, and 20 ng of template DNA. Thermal

cycling conditions were modified from Galen and Witt (2014), with an initial denaturation at 95 °C

for 8 mins, followed by 20 cycles of 94 °C for 30 seconds, 50 °C for 30 secs, 72 °C for 45 secs, and

a final extension at 72 °C for 10 mins.

For nested PCR, we used 1 µL of the outer PCR product as the template, with reaction

conditions identical to the outer PCR, except for an increase to 35 cycles. Each reaction set included

negative and positive controls to monitor for contamination and confirm amplification success,

respectively. We visualized PCR products on 2% agarose gels stained with SYBR Safe Gel Stain

(Invitrogen, Carlsbad, CA, USA) to verify the presence of amplicons of the expected size.

Successfully amplified products were purified using ExoSap-IT (Affymetrix, Inc., Santa Clara, CA,

USA) and submitted for Sanger sequencing at Psomagen (Rockville, MD, USA). gray vireo blood

samples from recaptured individuals (i.e., samples from the same individuals from 2017 and 2018)

underwent PCR amplification, but only a single sample was sequenced (Table S2).

Determining prevalence and haplotype diversity

We determined positive infections upon successful amplification of haemosporidian cyt *b*

sequences and calculated pathogen prevalence within each host species and calculated 95%

binomial confidence intervals using the 'exact' method available in the *binom* package in R

(version 4.3.2; R Core Team, 2023; Sundar, 2006).

Haemosporidian cyt *b* forward and reverse reads were trimmed to remove primers, resulting

in the target fragment size of 478 bp and were then assembled using the default alignment algorithm

in Geneious (version 2025.03; Biomatters Ltd; Kearse et al., 2012; Table S4). To identify

haplotypes, we compared cleaned sequences to published records stored in the public databases

GenBank (National Center for Biotechnology Information, US National Library of Medicine) and

MalAvi (Bensch et al., 2009) by using the Basic Local Alignment Search Tool (BLAST).

Additionally, to ensure the accuracy of both established and novel haplotypes, we downloaded all

haemosporidian cyt *b* haplotype sequence files from MalAvi and compared the number of

differences, if any, to our sequences via a distance matrix calculated in Geneious. Haplotypes that

differed by one or more base pairs (~0.2% sequence divergence) from published sequences on

GenBank or MalAvi were considered novel and named following MalAvi conventions.

Phylogenetic analysis

We estimated phylogenetic relationships among haemosporidian haplotypes using a

maximum-likelihood (ML) framework in RAxML, v8.2.10 (Stamatakis, 2014). Using the GTR+G

model of nucleotide substitution, we conducted a rapid bootstrap analysis with 1000 replicates, after

which we searched for the best-scoring ML tree. We rooted the tree with *Leucocytozoon* (COLBF21,

GenBank Accession MK947795) based on the current phylogenetic hypothesis for avian

haemosporidians (Borner et al., 2016; Galen et al., 2018). Nodes with <50% bootstrap support were

collapsed in TreeGraph2 (Stöver and Müller, 2010). We determined the relationships among the

three vireo study species based on the posterior distribution of likely trees available on birdtree.org

with the 'Hackett All Species' option (Hackett et al., 2008; Jetz et al., 2012). We downloaded 100

phylogeny subsets and selected the first tree because the relationships were consistent among the

100 trees.

Estimating parasite species richness and sampling completeness

We used the *iNEXT* R package (version 3.0.1; Chao et al., 2014; Hsieh et al., 2024) to

estimate parasite lineage diversity and evaluate sampling completeness for host species with

adequate data (i.e., gray and plumbeous vireos). In separate analyses for each host species, we used

infection counts of each parasite haplotype to generate rarefaction and extrapolation curves for

parasite species richness (q = 0), with the endpoint set at 400 individuals (i.e., infections). We

applied the 'iNEXT()' function to compute species richness estimates, including observed richness,

extrapolated richness, and associated confidence intervals using standard function parameters.

Additionally, we used the output to estimate sampling completeness (Chao et al., 2014) and defined

this value as the minimum number of individual birds required to detect 95% of the total estimated

parasite haplotypes infecting each host species. To visualize the rarefaction curves, we used the

'ggiNEXT()' function and *ggplot2* package for style modifications (Wickham, 2016).

Microscopy and parasitemia calculations

We obtained available blood smears from the MSB for plumbeous vireos ($n = 28$) and made

and air dried blood smears in the field for gray vireos ($n = 53$). Each slide was fixed with absolute

methanol and stained for 50 mins with Giemsa solution (pH 7.0; Sigma-Aldrich, St. Louis, MO,

USA). We then examined each slide to confirm infection status using light microscopy on an

Olympus BX 53 Microscope following the protocol described in Valkiūnas (2005), where 10,000

erythrocytes were scanned at 1000x magnification with an oil immersion lens to identify and count

Parahaemoproteus and *Plasmodium* infections (Valkiūnas, 2005). We took digital photographs of

representative haemosporidian specimens, which were archived at the MSB and are searchable via

the Arctos database (https://arctos.database.museum/). We calculated parasitemia, or the estimated

percentage of erythrocytes infected with haemosporidian parasites, as the number of infected

erythrocytes out of 10,000 screened. To account for lack of normality in the data, we used a

bootstrap method with 1,000 replications to calculate parasitemia 95% confidence intervals via the

R package *boot* (Kushary, 2000).

Characterizing environmental variation

We characterized elevation (m) and climatic variation using 19 bioclimatic variables at 30

sec (~1 km²) resolution from the WorldClim 2.1 database (Fick and Hijmans, 2017). We used

principal components analysis (PCA) for temperature (Bio1–11) and precipitation (Bio12–19)

variables to create a composite measure of temperature and precipitation across the gradient. The

first two axes for temperature (hereafter, tempPC1 and tempPC2) explained a combined 84.2% of

the variation and the first two axes for precipitation (hereafter, precipPC1 and precipPC2) explained

95.1% of the variation (Fig. S1; Table S5-6). Variable loadings indicated that tempPC1 primarily

represented year-round temperatures, with higher values relating to lower average temperatures,

while tempPC2 represented greater seasonality and less temperature stability, with higher values

relating to more extreme temperature fluctuations throughout the year and day-night cycle.

Similarly, precipPC1 described the overall precipitation amount, with higher values indicating

wetter conditions, whereas precipPC2 represented precipitation seasonality, with higher values

representing more stable and less seasonal precipitation (Table S5-6).

Assessing infection dynamics among host species and infection types

233   We tested whether haemosporidian infection status differed between categorical variables

such as host species and tissue type (i.e., pectoral muscle or whole blood samples), using Pearson's

chi-squared tests via the *stats* package. To examine the relative magnitude of effect for each

comparison, we calculated Cramér's V effect sizes were using the 'CramerV()' function in the

rcompanion package (Mangiafico, 2024). Next, to assess differences in parasitemia, we first

normalized the data using a natural log $\ln(1+x)$ transformation and removed two influential outliers

with parasitemia >4% (gray vireo IDs: 272132427 and 272132642). We then evaluated residual

variance with the 'var.test()' function in the *stats* package and compared log-transformed

parasitemia between gray and plumbeous vireos and infection types (i.e., single vs. coinfected)

using two-tailed Student's *t*-tests.

Intraspecific infection prevalence and parasitemia across environments

244   We employed a two-pronged approach to assess environmental associations with infection

prevalence: First, we used univariate, non-parametric tests to characterize and compare intraspecific

infection patterns across abiotic variables relevant to parasite and vector ecology, including

elevation, overall temperature (tempPC1) and precipitation (precipPC1), and the seasonality of

temperature (tempPC2) and precipitation (precipPC2). Then, we built intraspecific multivariate

linear models and applied nested and exhaustive model selection to evaluate the strength and

direction of environmental effects. We adopted this approach to account for the structure of our data

(i.e., non-overlapping elevational and environmental bands distinctly correlated to species identity)

and limitations of our dataset (i.e., limited sample sizes, degree of environmental variation, and

infection class imbalances).

We first assessed data normality both visually and then statistically, using the Shapiro-Wilk

test using the 'shapiro.test()' function from the *stats* package. Given the non-normal distribution of

environmental and geographic data in our dataset, we applied a rank-sum approach via Mann-

Whitney U-tests to compare the distributions of environmental variables between infected and

uninfected birds. Wilcoxon effect sizes of each comparison were then calculated via the

'wilcox_effsize()' function in the *rstatix* package (Kassambara, 2023).

We then constructed species-specific generalized linear models using glm() from the *stats*

package. Prevalence model sets used a binomial distribution (hereafter: "BEVI_prev" for Bell's

vireo, "GRVI_prev" for gray vireo, and "PLVI_prev" for plumbeous vireo sets), while parasitemia

model sets used a Gaussian distribution ("GRVI_para", "PLVI_para"). We evaluated linearity

between the logit and predicted values in prevalence models and assessed residual diagnostics in

parasitemia models and with 'autoplot()' within the *ggfortify* package (Tang et al., 2016).

Multicollinearity between variables was assessed by calculating variance inflation factors (VIF) via

the *car* package. As expected, there was high collinearity with aspects of geographic and

topographic variation, i.e. latitude and elevation, with other environmental variables. We retained

elevation in all model sets because it captured variation in habitat and vector dynamics important to

infection that were not explained by temperature and precipitation alone (Ishtiaq and Barve, 2018);

however, we removed latitude from all linear models to reduce overfitting.

For model sets with adequate sampling (i.e., GRVI_prev, PLVI_prev, GRVI_para,

PLVI_para), we evaluated all possible additive combinations of five predictor variables: elevation,

tempPC1, tempPC2, precipPC1, precipPC2. We used the 'dredge' function in the *MuMIn* package

(Bartoń, 2024) for exhaustive model selection. For host datasets with < 30 data points (BEVI_prev

and PLVI_para) we restricted candidate models to simple, biologically relevant structures to avoid

overfitting. Specifically, we included univariate models nested within simple additive models

containing at most two predictor variables, as well as null and global models for comparison. We

then compared model performance and evaluated the trade-off between model fit and complexity

using Akaike's Information Criterion corrected for small sample sizes (AICc; Table 1).

To examine the effect of each predictor in top-ranked models, we standardized and

calculated model-averaged regression coefficients using the *MuMIN* package (Table 2). For top-

ranked models where the null model was not favored, we assessed goodness of fit via Hosmer-

Lemeshow tests as part of the *ResourceSelection* package (Lele et al., 2023) and checked for

overdispersion by dividing residual deviance by the degrees of freedom. Predictive accuracy was

also assessed using 10-fold cross-validation via the *caret* package (Kuhn, 2008).

Results

Haemosporidian prevalence, abundance, and diversity

We uncovered an unusually high prevalence of haemosporidian infections in all three vireo

species, with 175 infected birds of 248 screened (70.6%; 95% CI [64.5 – 76.2%]). The highest

documented prevalence was found in plumbeous vireos (84.5% infected; 95% CI [72.6 – 92.7%]),

followed by gray vireos (67.7%; 95% CI [60.0 – 74.7%]), and lastly, Bell's vireos (55%; 95% CI

[31.5 – 76.9%]; Fig. 1B). Prevalence was significantly lower in Bell's than in plumbeous vireos

($\chi^2(1) = 5.72$, p = 0.017, $\varphi = 0.31$), indicating a moderate effect size. While most birds were

infected by single haemosporidian haplotype infections (88.1%; 95% CI [82.3 – 92.5%]), we found

21 individuals that were coinfected with two or more haplotypes (12.0%; 95% CI [7.54 – 17.7%]),

and one gray vireo that was infected by three haplotypes (VIRVIC01, VIRVIC02, VIRVIC03). Of

the 14 gray vireos recaptured about a year after initial sampling, 12 were infected with

haemosporidians in both years, while two that were uninfected in 2017 tested positive in 2018.

Nearly all infections were caused by parasites in the genus *Haemoproteus*, specifically the

subgenus *Parahaemoproteus* (99.5%), which comprised 18 of 19 haplotypes identified.

Plasmodium was rare in our data, with a single infection (lineage VIRPLU13) recorded from a

plumbeous vireo (MSB:Bird:60726; Table S4). Of the 19 total parasite haplotypes identified, 8

were novel (Fig. 1D; Table S4).

Estimates of parasite species richness and sampling completeness

We identified a total of three haemosporidian haplotypes from Bell's vireos, four from gray

vireos, and 14 from plumbeous vireos (Fig. 1D). These differences were partly, but not entirely,

attributable to differences in sample size. gray vireo *Parahaemoproteus* infections were

substantially less diverse than those of plumbeous vireos despite more intensive screening efforts.

Extrapolation from rarefaction curves showed considerably lower estimated parasite richness for

gray vireos (4.99 ± 0.53 SE; 95% CI [4.00–6.03]; Table S7) and estimated that much smaller

sample sizes would be needed to reach 95% sampling completeness (14 infections; Fig. 2; Table

S8). This result contrasted our results for plumbeous vireos, where parasite richness was estimated

to be ~10x higher than gray vireos (53.8 ± 23.3 SE; 95% CI [14.0–99.6]) and require much greater

sampling effort (~26x more than estimated for gray vireos) to reach an estimated 95% completeness

(estimated 364 infections). Bell's vireos also exhibited low lineage diversity (3.92 ± 0.784 SE; 95%

CI [3.00-5.45] and an estimated 95% sample completeness at only 33 infections, consistent with the

limited diversity recovered (Table S7-S8). However, we caution overinterpreting these results due

to the small Bell's vireo sample size (*n* = 12 infected) in our study and recommend additional

sampling to generate more precise estimates.

Of the 19 total haplotypes identified, we found evidence of potential host specificity with at

least one species-specific haplotype infecting each of the three vireo species (Table S4). Of note,

gray vireos were infected by novel haemosporidian haplotypes only, (VIRVIC01-VIRVIC04). This

finding contrasted with plumbeous vireos, which showed novel haplotypes (i.e. VIRPLU10-13) and

haplotypes previously found only in plumbeous vireos (i.e. VIRPLU03, VIRPLU07-09), as well as

lineages known to infect other non-vireo species (i.e. SETAUD14, TROAED12, VIGIL07,

VIRPLU01, and VIRPLU04). Sampling was comparatively lower for Bell's vireos; no novel

lineages were detected. We did detect, however, a recently described lineage (VIRBEL01) which

was previously documented in Bell's vireos sampled from Arizona and Texas (Fecchio et al., 2023),

as well as other more generalist lineages (i.e. TROAED12 and VIGIL07; Table S4).

Intensity of infection (parasitemia) across hosts and haplotypes

We screened 102 individuals via microscopy (gray n = 71, plumbeous n = 31). A total of 81

samples showed evidence of infection and had parasitemia *calculated (Parahaemoproteus:* gray n =

53 and plumbeous n = 27; *Plasmodium*: plumbeous n = 1). gray vireos typically had slightly higher

infection loads (1.00%; 95% CI [0.782 – 1.27%]) compared to plumbeous vireos (0.786%; 95% CI

[0.584 – 1.01%]), partly due to two outlier gray vireo samples with >4% parasitemia (ID

272132427 and 27213264; Fig 1C). After removing these outliers, we found that mean parasitemia

did not differ significantly between host species (t(77) = 0.656, p = 0.514). When we compared

single-haplotypes and coinfections, we also found no significant difference in parasitemia between

species (t(73) = -0.728, p = 0.469), though we note the sampling pool of coinfected birds was small

compared to single infections (n = 7 and n = 68, respectively).

Our comparison of infections using PCR and microscopy was largely consistent, with few

detection errors identified between the two methods. We detected four positively infected birds via

microscopy that did not successfully amplify by PCR (Table S1). These samples had a range of

parasitemia spanning 0.70% to 2.68%. Additionally, we failed to detect haemosporidian infections

via microscopy for two birds that were later successfully amplified using PCR; one individual was

the single coinfected gray vireo with three *Parahaemoproteous* haplotypes that were identified by

sequencing (Table S1).

Environmental patterns of infection status and parasitemia

Parahaemoproteus prevalence showed idiosyncratic patterns among host species and their

associated habitats (Fig 3; Table S9). For example, we found that infected Bell's vireos tended to be

found at lower latitudes (z = -2.09, p = 0.040), with warmer year-round temperatures (tempPC1; z =

-2.25, p = 0.027 Fig. 3; Fig. S2; Table S9). Similarly, probability of infection increased as overall

precipitation (precipPC1) and temperature seasonality (tempPC2) increased (Table 1; Table 2). In

contrast, we found higher proportions of infected gray vireos at lower elevations (z = -2.07, p =

0.038), Fig. 1C; Table S9) and at sites with more stable precipitation (precipPC2; z = 1.72, p =

0.057), Fig 3; Table S9). However, this result was dampened when we excluded the eleven infected

gray vireos from the Abajo Mountain site in Utah, which were outliers in many of the

environmental PCA dimensions. The top model, however, showed that infection was negatively

associated with elevation and positively associated with temperature (tempPC1; Table 1; Table 2)

even when excluding the Utah site. Lastly, plumbeous vireo infections tended to be higher in areas

with more seasonally stable precipitation (precipPC2; z = 2.04, p = 0.042), Fig. 3). Despite the top-

ranked infection probability model showing no significant linear trends, precipitation (precipPC1

and precipPC2) was present in the majority of top candidate models (Table 1).

We did not find any association between parasitemia and environmental variables in

plumbeous vireos, with the null model having the strongest support and no clear predictor variable

significantly reported across top models (Table S10). gray vireos, however, showed a significant

pattern of increased parasitemia as overall precipitation (precipPC1) increased (Table S10), but this

model had low explanatory power (R^2 = 0.05).

Discussion

Vireo species breeding in the southwestern U.S. are especially susceptible to

haemosporidian infection, particularly by parasites in the *Parahaemoproteus* subgenus. In this study,

we uncovered that the elevational limits of haemosporidian parasites closely parallel those of their

avian hosts. We observed striking differences in host specificity and parasite richness across host

species and their associated habitats, revealing that the high-elevation plumbeous vireos harbored

the most diverse and phylogenetically distinct parasite lineages, especially compared to another

well-sampled, lower elevation host species, the gray vireo. Although parasite prevalence differed

significantly between Bell's vireos (55% infected) and plumbeous vireos (86.2% infected), which

breed under different levels of precipitation and temperature, we found variable links between

infection status and climatic factors in our dataset. Specifically, we found that infection probability

was strongly influenced by aspects of temperature for Bell's vireos, elevation for gray vireos, and

precipitation for plumbeous vireos, likely reflecting subtle differences in habitat suitability for

arthropod vectors. Lastly, we found low to moderate parasitemia values in both gray and plumbeous

vireos, consistent with chronic infections. We examine these results in relation to previous

haemosporidian research in the southwest U.S. and in vireos, both in their breeding and

nonbreeding grounds, and discuss remaining questions in the sections below.

Elevational replacement and haemosporidian diversity patterns

388   Avian haemosporidians exhibit variable patterns of turnover across elevations (Galen and

Witt, 2014; Illera et al., 2015; McNew et al., 2021), which is often independent of geographic

distance between hosts (Williamson et al., 2019; Barrow et al., 2021). We demonstrated a strong

case of parallel elevational replacement in hosts and parasites, with high turnover among

Parahaemoproteus assemblages infecting vireo species breeding at different elevations. This

elevational replacement pattern could be the result of a few different scenarios. For example, shared

historical and ecological forces, such as habitat specialization in both hosts and vectors (Ellis et al.,

2020), may shape the elevational range limits of both vireos and their *Parahaemoproteus* parasites.

Alternatively, species-specific factors that co-evolved independently in response to specialized

parasite lineage pressures, such as immune defenses (e.g., Armour et al., 2025), could also be at

play. Whether parasite elevational replacement is host-driven or independently mediated by the

environmental gradient, global patterns of spatial diversity in parasites should be tightly coupled

with those of their hosts.

401   Of the 30 cyt *b Parahaemoproteus* and 12 *Plasmodium* haplotypes currently reported to

infect vireo species (MalAvi; downloaded 1/1/2025), our study adds seven novel

Parahaemoproteus and one novel *Plasmodium* haplotype. The near absence of *Plasmodium*

infections, even at lower elevations, contrasts findings from other temperate region studies (e.g. in

Alaska: Oakgrove et al., 2014; California: Walther et al., 2016; Oklahoma: Wyckoff et al., 2024)

but is consistent with previous screening at high elevations in New Mexico (Marroquin-Flores et al.,

2017; Williamson et al., 2019; Barrow et al., 2021). While the cooler temperatures experienced at

higher elevations have been proposed as a potential limiting factor for haemosporidian infection

(e.g. LaPointe et al., 2012; Barrow et al., 2021), the low prevalence of *Plasmodium* at lower

elevations in this region may also reflect limited suitable habitat for mosquito vectors prior to the

onset of monsoon rains. Broader and temporally replicated surveys of potential vectors and vector

abundance in this region are needed to fully explain this apparent absence.

We identified an interesting pattern among the *Parahaemoproteus* lineages found in our

focal species: plumbeous vireos harbored a broad diversity of parasite haplotypes relative to the

other two sampled species, while gray vireos exhibited a depauperate and likely specialized

haemosporidian assemblage (Fig. 1D). This disparity suggests that plumbeous vireos may support

more permissive or ecologically flexible parasite associations, while gray vireos may be undergoing,

or may have already undergone, a transition toward fixed symbiosis with a narrower set of

haemosporidian lineages. The contrasting pattern of haemosporidian diversity between vireo

species could alternatively be caused by differences in avian host richness, and the low diversity

found in gray vireos could reflect the depauperate breeding bird communities of the arid and

structurally simple juniper savanna habitats.

Conserved high susceptibility to haemosporidians among vireo species

Vireos stand out among southwestern songbirds for their consistently high haemosporidian

infection rates, providing a valuable opportunity to examine parasite dynamics in hyper-susceptible

hosts. In the three species screened here, we found that 70.6% of 248 individuals screened had at

least one haemosporidian infection, which is greater than double the prevalence reported in broader

community-wide surveys conducted in the southwestern U.S. over a similar period. For example,

Marroquin-Flores et al. (2017) screened 186 birds of 49 species sampled in New Mexico from June-

July 2016, including 13 individual vireos of two species: $n = 4$ warbling vireos (*Vireo gilvus*) and n

$= 9$ plumbeous vireos. The average proportion of infection was 36.6% across all studied species, but

92% of vireos tested positive. A subsequent large-scale study by Barrow et al. (2021) reported

similar patterns, with an overall haemosporidian prevalence of 36.1% among 776 total individuals

of 61 species. Of these, vireos again exhibited significantly higher infection proportions, with

plumbeous and warbling vireos demonstrating the highest prevalence in the study at 87% and 78%,

respectively (Barrow et al., 2021).

This observation of high susceptibility in vireos is not limited to birds sampled in the

southwestern U.S., as other studies across breeding sites have also reported higher relative

prevalences in the *Vireo* genus. Granthon and Williams (2017) reported that 100% of the 62 PCR-

screened red-eyed vireos (*Vireo olivaceus*) from Pennsylvania were infected with *Plasmodium*,

Haemoproteus, or *Leucocytozoon* species (Granthon and Williams, 2017). Rodriquez et al. (2021)

found *Haemoproteus* prevalence of 59% in 22 individual warbling vireos sampled from Colorado

breeding sites (Rodriguez et al., 2021). Shared susceptibility within closely related species has been

extensively documented (e.g. González et al., 2014; Lutz et al., 2015; Barrow et al., 2019; Pigeault

et al., 2022) but attempts to identify the specific factors that contribute to high prevalence in vireos

are still lacking. Of the conserved host traits recorded to have consistent interaction with

haemosporidian prevalence in other avian clades, thus far, shared life history traits such as open cup

nest type and insectivorous diets (as reported and reviewed in Braga et al., 2011; Rodriguez et al.,

2021) may increase exposure risk, at least partially explaining the high prevalence in this group.

However, a combination of traits, including behavior (e.g., time spent incubating and/or brooding at

the nest), features of nest microhabitat (e.g., height of nest from ground), and aspects of immunity

(e.g., haemosporidian tolerance) may also contribute to the elevated susceptibility (González et al.,

2014; Lutz et al., 2015; Muriel, 2020; Rodriguez et al., 2021; Pigeault et al., 2022).

Interestingly, of the 14 gray vireos that we recaptured and screened for parasites across two

sampling years (2017 and 2018), 12 were positive via PCR in both years, and the remaining two

birds became infected by the second year (Table S2). Although more data is needed to determine

whether these represent chronic infections or reinfections, and pinpoint where transmission occurs,

these results indicate gray vireos likely experience sustained parasite pressure during the breeding

season. Continued monitoring of haemosporidian infection across vireo populations, coupled with

investigations into the mechanisms underlying their susceptibility, could yield valuable insights into

the spatial and temporal dynamics of infection risk in this clade.

Environmental determinants of haemosporidian infection vary by host species

463   Previous studies have found strong links between haemosporidian prevalence and

topographic, geographic, and climatic variables (Pérez☐Rodríguez et al., 2013; Rodriguez et al.,

2021), but the direction and magnitude of these effects vary by region and haemosporidian clade

(Williamson et al., 2019). In the arid southwest, we expected haemosporidian prevalence to

correspond with water availability, which is important for insect vector (Valkiūnas, 2004;

Santiago☐Alarcon et al., 2012), such as those associated with potential water sources (i.e. riparian

habitats) and areas with higher, more seasonally stable precipitation (i.e. montane woodlands).

Somewhat contrary to expectations given predictions about the relative aridity of habitat during the

pre-monsoon sampling period, we found that interspecific differences in prevalence increased

linearly with elevation and precipitation: Bell's vireos showed the lowest prevalence and gray

vireos exhibited unexpectedly high infection rates. We were unable to determine whether observed

infection patterns were driven by species-specific traits or abiotic environmental differences, and

increased sampling of Bell's vireos may help clarify or reduce these differences. Within-species

analyses revealed no consistent predictors of infection across all taxa (Fig. S2-6); instead, different

environmental variables were associated with infection in each species (Fig. 3; Table S9-S10).

Expanded screening of haemosporidians across low- to mid-elevation habitats, among co-

distributed species, at variable distance from water sources, and across precipitation gradients

would further elucidate cross-elevational trends and enhance predictions of infection probability in

avian communities of the southwestern U.S.

Moderate and consistent infection loads across environmental gradients

Gray and plumbeous vireos exhibited similar average infection loads (~1% in gray vireos,

0.786% in plumbeous, range: 0.03–4.5%) with no significant differences between species (Fig. 1B).

These values slightly exceed the regional cross-species average (<1%; Marroquin-Flores et al.,

2017) and fall within the range of moderate infection intensities reported in experimental studies of

other passerines (e.g. Asghar et al., 2012). We did note, however, two gray vireos with elevated

parasitemia (4.5 and 4.49%) were sampled at distinct sites in the Sevilleta National Wildlife Refuge.

The elevated parasitemia in these individuals likely represented acute infection acquired shortly

before capture and prior to immune response, or recrudescence of chronic infections exacerbated by

physiological stressors such as nutritional deficits or hormonal fluctuations (Arriero et al., 2018;

Names et al., 2021). The absence of high parasitemia in Bell's and plumbeous vireos could be an

artifact of limited sampling, as birds with severe infections often exhibit reduced mobility and

capture probability (Mukhin et al., 2016), suggesting that heavily infected individuals in these

species may be underrepresented.

Additionally, we found no relationship between parasitemia and any tested environmental

variables for plumbeous vireos. In gray vireos we saw a significant linear trend in parasitemia with

increased overall precipitation (precipPC1; Table S10), but this model had weak explanatory power

($R^2 = 0.05$). These results are consistent with previous findings which suggest that haemosporidian

parasitemia patterns are shaped by complex interactions between host and parasite biology not

easily captured in models with abiotic factors alone (Loiseau et al., 2013).

Conclusions

We found that parasite elevational distributions mirror the elevational replacement pattern of

host species among vireos in southwestern North America. This pattern may have resulted from

host specialization, climate variation independently acting on hosts and parasites, climate variation

acting indirectly through habitat or broader host and vector communities, or a combination of those

forces. We found that vireos were consistently infected at high rates and largely harbor chronic,

low-to-moderate intensity infections dominated by *Parahaemoproteus*, suggesting the potential for

exclusive, reciprocal selective pressures that facilitate host-parasite coadaptation. The contrast in parasite richness between plumbeous vireos and gray vireos was striking: gray vireos harbored a limited number of haemosporidian lineages, all of which were host specific, while plumbeous vireos were infected with a diverse set of lineages. These differences imply species-specific variability in resistance, tolerance, or exposure to generalist haemosporidian lineages. In sum, our work adds to a growing body of evidence about clade-specific avian host susceptibility and host-parasite turnover across elevational gradients, highlighting vireos as an ideal focal group for future work on host-parasite dynamics across space and time.

Acknowledgements

The authors sincerely thank Bethany Abrahamson, Michael Andersen, Matthew Baumann, Alison Boyer, Serina Brady, Sara Brant, Mariel Campbell, Andrea Chavez, Mina Carnicom, Lida Crooks, Rosario Marroquin-Flores, Chauncey Gadek, Spencer Galen, Ethan Gyllenhaal, Mike Hartshorne, Andy Johnson, Michael Lelevier, Joseph Manthey, Xena Mapel, Taylor Martinez, Jenna McCullough, Moses Michelsohn, Kathleen Ramsay, George Rosenberg, C. Jonathan Schmitt, Donna Schmitt, Brenda Villanueva, and Toby Weisenhaus.

Declarations

Not applicable.

Funding

This work was supported by the National Science Foundation (Graduate Research Fellowship NSF-DGE-2439853 to DLFW; Postdoctoral Research Fellowship in Biology NSF-DBI-1611710 to LNB; and NSF-DEB-1146491 to CCW), the Bureau of Land Management Rio Puerco Field Office (via the Colorado Plateau Cooperative Eco-systems Studies Unit agreement), New Mexico Department of Game and Fish Share with Wildlife Grant to HMS/SEF, and a Sevilleta LTER Graduate Summer Research Fellowship and New Mexico Ornithological Society Ryan Beaulieu Research Grant to SEF.

Conflict of interest

535   The authors have no conflicts of interest to declare.

Ethics approval

537   Research activities were conducted under Institutional Animal Care and Use Protocol 16-200406-

538   MC and appropriate state and federal scientific collecting permits (New Mexico Department of

539   Game and Fish Authorization Number 3217; US Fish and Wildlife Permit Numbers MB094297-0

540   and MB0942970-1).

Availability of data and material

Specimen data are available in Supplementary Tables S1–S10 and are searchable via Arctos

(arctos.database.museum; Cicero et al., 2024). Parasite sequences are archived in GenBank

(accession nos. PV295634-PV295830). Vouchered specimens and frozen tissue samples are

archived at the Museum of Southwestern Biology Division of Birds and Division of Genomic

Resources, respectively (https://msb.unm.edu/). All R code and datasets are available on FigShare

(https://doi.org/10.6084/m9.figshare.28585943).

Authors' contributions

DLFW, JLW, CCW, and LNB conceptualized and developed methodology for this project. JLW,

SEF, HMS, SMB, KG, CCW, and LNB performed fieldwork and subsequent data curation. DLFW,

JLW, and SMB performed molecular analysis and generated sequence data. JLW, SEF, HMS, KG,

CCW, and LNB secured funding and provided supervision. DLFW and JLW performed formal

analyses and created visualizations. DLFW and JLW wrote the original draft of the manuscript, and

DLFW wrote the final version with input from all co-authors.

References

Albon SD, Stien A, Irvine RJ, Langvatn R, Ropstad E, Halvorsen O (2002) The role of parasites in

the dynamics of a reindeer population. Proc R Soc Lond B Biol Sci 269:1625–1632 doi:

10.1098/rspb.2002.2064

Arriero E, Pérez-Tris J, Ramírez A, Remacha C (2018) Trade-off between tolerance and resistance

to infections: an experimental approach with malaria parasites in a passerine bird. Oecologia

188:1001–1010 doi: 10.1007/s00442-018-4290-4

Armour C, Hone AL, Dunn JC (2025) Do specialist haemoparasites induce tolerance in their hosts?

Parasitol 152:374–380 doi: 10.1017/ S0031182025000393

Asghar M, Westerdahl H, Zehtindjiev P, Ilieva M, Hasselquist D, Bensch S (2012) Primary peak

and chronic malaria infection levels are correlated in experimentally infected great reed

warblers. Parasitol 139:1246–1252 doi: 10.1017/S0031182012000510

Atkinson CT, van Riper C (1991) Pathogenicity and epizootiology of avian haematozoa:

Plasmodium, Leucocytozoon, and Haemoproteus. In: Loye JE, Zuk M (eds) Bird-parasite

interactions: ecology, evolution, and behavior. Oxford Academic Press, Oxford, pp 19–48

Barlow JC, (1977) Effects of habitat attrition on vireo distribution and population density in the

northern Chihuahuan Desert. In: Wauer RH, Riskind DH (eds) Transactions of the

symposium on the biological resources of the Chihuahuan Desert region, United States and

Mexico. U.S. Department of the Interior National Park Service Trans Proc, pp 591–596

Barlow JC, Leckie SN, Baril CT (1999) Gray vireo (*Vireo vicinior*). In: Poole A, Gill F (eds) The

birds of North America. The Academy of Natural Sciences, Philadelphia, PA, and The

American Ornithologists' Union, Washington, D.C.

Barrow LN, Bauernfeind SM, Cruz PA, Williamson JL, Wiley DLF, Ford JE, Baumann MJ, Brady

SS, Chavez AN, Gadek CR, Galen SC, Johnson AB, Mapel XM, Marroquin-Flores RA,

Martinez TE, McCullough JM, McLaughlin JE, Witt CC (2021) Detecting turnover among

complex communities using null models: a case study with sky-island haemosporidian

parasites. Oecologia 195:435–451 doi: 10.1007/s00442-021-04854-6

Barrow LN, Lemmon AR, Lemmon EM (2018) Targeted sampling and target capture: assessing

phylogeographic concordance with genome-wide data. Syst Biol 67:979–996 doi:

10.1093/sysbio/syy021

Barrow LN, McNew SM, Mitchell N, Galen SC, Lutz HL, Skeen H, Valqui T, Weckstein JD, Witt

CC (2019) Deeply conserved susceptibility in a multi☐host, multi☐parasite system. Ecol

Lett 22:987–998 doi: 10.1111/ele.13263

Bartoń K (2024) MuMIN: Multi-model inference. R package version 1.48.4 https://CRAN.R-

project.org/package=MuMIn

Bennett GF, Peirce MA, Ashford RW (1993) Avian Haematozoa: mortality and pathogenicity. J

Nat Hist 27:993–1001 doi: 10.1080/00222939300770621

Bensch S, Hellgren O, Pérez☐Tris J (2009) MalAvi: a public database of malaria parasites and

related haemosporidians in avian hosts based on mitochondrial cytochrome *b* lineages. Mol

Ecol Resour 9:1353–1358 doi: 10.1111/j.1755-0998.2009.02692.x

Bensch S, Waldenström J, Jonzén N, Westerdahl H, Hansson B, Sejberg D, Hasselquist D (2007)

Temporal dynamics and diversity of avian malaria parasites in a single host species. J Anim

Ecol 76:112–122 doi: 10.1111/j.1365-2656.2006.01176.x

Billerman SM, Keeney BK, Kirwan GM, Medrano F, Sly ND, Smith MG (2025) Birds of the

World. Cornell Laboratory of Ornithology, Ithaca, New York

https://birdsoftheworld.org/bow/home

Borner J, Pick C, Thiede J, Kolawole OM, Kingsley MT, Schulze J, Cottontail VM, Wellinghausen

N, Schmidt-Chanasit J, Bruchhaus I, Burmester T (2016) Phylogeny of haemosporidian

blood parasites revealed by a multi-gene approach. Mol Phylogenet Evol 94:221–231 doi:

10.1016/j.ympev.2015.09.003

Boysen KE, Perkins SL, Hunjan S, Oliver P, Gardner MG, Balasubramaniam S, Melville J (2022)

Diversity and phylogenetic relationships of haemosporidian and hemogregarine parasites in

Australian lizards. Mol Phylogenet Evol 167:107358 doi: 10.1016/j.ympev.2021.107358

Braga ÉM, Silveira P, Belo NO, Valkiūnas G (2011) Recent advances in the study of avian malaria:

an overview with an emphasis on the distribution of *Plasmodium* spp in Brazil. Mem Inst

Oswaldo Cruz 106:3–11 doi: 10.1590/S0074-02762011000900002

Brown BT (1993) Bell's vireo (*Vireo bellii*). In: Poole A, Gill F (eds) The birds of North America.

The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists'

Union, Washington, D.C.

Cadena CD, Loiselle BA (2007) Limits to elevational distributions in two species of emberizine

finches: disentangling the role of interspecific competition, autoecology, and geographic

variation in the environment. Eco 30:491–504 doi: 10.1111/j.0906-7590.2007.05045.x

Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM (2014) Rarefaction

and extrapolation with Hill numbers: a framework for sampling and estimation in species

diversity studies. Ecol Monogr 84:45–67. doi: 10.1890/13-0133.1

Cicero C, Koo MS, Braker E, Abbott J, Bloom D, Campbell M, Cook JA, Demboski JR, Doll AC,

Frederick LM, Linn AJ, Mayfield-Meyer TJ, McDonald DL, Nachman MW, Olson LE,

Roberts D, Sikes DS, Witt CC, Wommack EA (2024) Arctos: Community-driven

innovations for managing natural and cultural history collections. PLOS ONE 19:e0296478

doi: 10.1371/journal.pone.0296478

Clark NJ (2018) Phylogenetic uniqueness, not latitude, explains the diversity of avian blood parasite

communities worldwide. Glob Ecol Biogeogr 27:744–755 doi: 10.1111/geb.12741

Coblentz DD, Riitters KH (2004) Topographic controls on the regional□scale biodiversity of the

southwestern USA. J Biogeogr 31:1125–1138 doi: 10.1111/j.1365-2699.2004.00981.x

Combes C (2001) Parasitism: the ecology and evolution of intimate interactions. University of

Chicago Press, Chicago and London

Combes C (1997) Fitness of parasites: Pathology and selection. Int J Parasitol 27:1–10. doi:

10.1016/S0020-7519(96)00168-3

Curson DR, Goguen CB (1998) Plumbeous vireo (*Vireo plumbeus*). In: Poole A, Gill, F (eds) The

birds of North America, Inc, Philadelphia, Pennsylvania

Duval L, Robert V, Csorba G, Hassanin A, Randrianarivelojosia M, Walston J, Nhim T, Goodman

SM, Ariey F (2007) Multiple host-switching of Haemosporidia parasites in bats. Malar J

6:157 doi: 10.1186/1475-2875-6-157

Ellis VA, Bensch S (2018) Host specificity of avian haemosporidian parasites is unrelated among

sister lineages but shows phylogenetic signal across larger clades. Int J Parasitol 48:897–902

doi: 10.1016/j.ijpara.2018.05.005

Ellis VA, Collins MD, Medeiros MCI, Sari EHR, Coffey ED, Dickerson RC, Lugarini C, Stratford

JA, Henry DR, Merrill L, Matthews AE, Hanson AA, Roberts JR, Joyce M, Kunkel MR,

Ricklefs RE (2015) Local host specialization, host-switching, and dispersal shape the

regional distributions of avian haemosporidian parasites. Proc Natl Acad Sci

112(36):11294–11299 doi: 10.1073/pnas.1515309112

Ellis VA, Huang X, Westerdahl H, Jönsson J, Hasselquist D, Neto JM, Nilsson J, Nilsson J,

Hegemann A, Hellgren O, Bensch S (2020) Explaining prevalence, diversity and host

specificity in a community of avian haemosporidian parasites. Oikos 129(9):1314–1329. doi:

10.1111/oik.07280.

Fecchio A, Bell JA, Williams EJ, Dispoto JH, Weckstein JD, De Angeli Dutra D (2023) Co-

infection with *Leucocytozoon* and other haemosporidian parasites increases with latitude and

altitude in New World bird communities. Microb Ecol 86:2838–2846 doi: 10.1007/s00248-

023-02283-x

Fecchio A, Wells K, Bell JA, Tkach VV, Lutz HL, Weckstein JD, Clegg SM, Clark NJ (2019)

Climate variation influences host specificity in avian malaria parasites. Ecol Lett 22:547–

557 doi: 10.1111/ele.13215

Fick SE, Hijmans RJ (2017) WorldClim 2: new 1 km spatial resolution climate surfaces for global

land areas. Int J Climatol 37:4302–4315 doi: 10.1002/joc.5086

Filion A, Eriksson A, Jorge F, Niebuhr CN, Poulin R (2020) Large-scale disease patterns explained

by climatic seasonality and host traits. Oecologia 194:723–733 doi: 10.1007/s00442-020-

04782-x

Fischer SE (2020) Post-fledging and migration ecology of gray vireos (*Vireo vicinior*) and using

ArtScience to explore gender and identity. Master's thesis, The University of Toledo,

Toledo, Ohio, USA doi: 10.13140/RG.2.2.31260.41603

Fischer SE, Granillo K, Streby HM (2022) Post-fledging survival, movements, and habitat

associations of gray vireos in New Mexico. Avian Conserv Ecol 17:art13 doi:

10.5751/ACE-02053-170113

Fischer SE, Kramer GR, Baumann MJ, Fahy EF, Granillo K, Streby HM (2025) Migratory

connectivity and potential nonbreeding sexual segregation in gray vireos (*Vireo vicinior*).

Wilson J Ornithol doi: 10.1080/15594491.2025.2497654

Freeman BG, Class Freeman AM, Hochachka WM (2016) Asymmetric interspecific aggression in

New Guinean songbirds that replace one another along an elevational gradient. Ibis

158(4):726–737 doi: 10.1111/ibi.12384

Gage KL, Burkot TR, Eisen RJ, Hayes, EB (2008) Climate and vectorborne diseases. Am J Prev

Med 35:436–450 doi: 10.1016/j.amepre.2008.08.030

Galen SC, Borner J, Martinsen ES, Schaer J, Austin CC, West CJ, Perkins SL (2018) The

polyphyly of *Plasmodium*: comprehensive phylogenetic analyses of the malaria parasites

(order Haemosporida) reveal widespread taxonomic conflict. R Soc Open Sci 5:171780. doi:

10.1098/rsos.171780

Galen SC, Witt CC (2014) Diverse avian malaria and other haemosporidian parasites in Andean

house wrens: evidence for regional co☐diversification by host☐switching. J Avian Biol

45:374–386 doi: 10.1111/jav.00375

González AD, Matta NE, Ellis VA, Miller ET, Ricklefs RE, Gutiérrez HR (2014) Mixed species

flock, nest height, and elevation partially explain avian haemoparasite prevalence in

Colombia. PLOS ONE 9:e100695 doi: 10.1371/journal.pone.0100695

Granthon C, Williams DA (2017) Avian malaria, body condition, and blood parameters in four

species of songbirds. Wilson J Ornithol 129:492–508 doi: 10.1676/16-060.1

Gyllenhaal EF (2024) Crossing the ocean: the importance of gene flow in island evolution.

University of New Mexico, Albuquerque, New Mexico, USA

Hackett SJ, Kimball RT, Reddy S, Bowie RCK, Braun EL, Braun MJ, Chojnowski JL, Cox WA,

Han KL, Harshman J, Huddleston CJ, Marks BD, Miglia KJ, Moore WS, Sheldon FH,

Steadman DW, Witt CC, Yuri T (2008) A phylogenomic study of birds reveals their

evolutionary history. Science 320:1763–1767 doi: 10.1126/science.1157704

Hasik AZ, Siepielski AM (2022) Parasitism shapes selection by drastically reducing host fitness

and increasing host fitness variation. Biol Lett 18:20220323 doi: 10.1098/rsbl.2022.0323

Hellgren O, Waldenström J, Bensch S (2004) A new PCR assay for simultaneous studies of

Leucocytozoon, *Plasmodium*, and *Haemoproteus* from avian blood. J Parasitol 90:797–802

doi: 10.1645/GE-184R1

Hellgren O, Waldenström J, Peréz☐Tris J, Szöll E, Si Ö, Hasselquist D, Krizanauskiene A,

Ottosson U, Bensch S (2007) Detecting shifts of transmission areas in avian blood parasites

– a phylogenetic approach. Mol Ecol 16:1281–1290 doi: 10.1111/j.1365-

294X.2007.03227.x

Hsieh T, Ma K, Chao A (2025). iNEXT: Interpolation and extrapolation for species diversity. R

package version 3.0.2 http://chao.stat.nthu.edu.tw/wordpress/software_download/

Hubbard JP (1970) Check-list of the birds of New Mexico. NM Ornithol Soc 3:108

Illera JC, López G, García-Padilla L, Moreno Á (2017) Factors governing the prevalence and

richness of avian haemosporidian communities within and between temperate mountains.

PLOS ONE 12:e0184587 doi: 10.1371/journal.pone.0184587

Ikemoto T (2008) Tropical malaria does not mean hot environments. J Med Entomol 45(6):963–969

doi: 10.1603/0022-2585(2008)45[963:tmdnmh]2.0.co;2

Ishtiaq F, Barve S (2018) Do avian blood parasites influence hypoxia physiology in a high elevation

environment? BMC Ecol 18 doi: 10.1186/s12898-018-0171-2

Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO (2012) The global diversity of birds in space

and time. Nature 491:444–498 doi: 10.1038/nature11631

Kassambara A (2023) rstatix: Pipe-friendly framework for basic statistical tests. R package version

0.7.2 https://CRAN.R-project.org/package=rstatix

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,

Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious

basic: An integrated and extendable desktop software platform for the organization and

analysis of sequence data. Bioinform 28:1647–1649 doi: 10.1093/bioinformatics/bts199

Krasnov BR and Poulin R (2010) Ecological properties of a parasite: species-specific stability and

geographical variation. The biogeography of host–parasite interactions. Oxford University

Press, Oxford, pp 99-113

Kuhn M (2008) Building predictive models in *R* using the caret package. J Stat Softw 28 doi:

10.18637/jss.v028.i05

Kushary D (2000) Bootstrap methods and their application. Technometrics 42:216–217 doi:

10.1080/00401706.2000.10486018

LaPointe DA, Atkinson CT, Samuel MD (2012) Ecology and conservation biology of avian malaria.

Ann NY Acad Sci 1249:211–226 doi: 10.1111/j.1749-6632.2011.06431.x

Lauron EJ, Loiseau C, Bowie RCK, Spicer GS, Smith TB, Melo M, Sehgal RNM (2015)

Coevolutionary patterns and diversification of avian malaria parasites in African sunbirds

(Family Nectariniidae). Parasitol 142:635–647 doi: 10.1017/S0031182014001681

Lele SR, Keim JL, Solymos P (2023) ResourceSelection: resource selection (probability) functions

for use-availability data. https://CRAN.R-project.org/package=ResourceSelection

Loiseau C, Harrigan RJ, Bichet C, Julliard R, Garnier S, Lendvai ÁZ, Chastel O, Sorci G (2013)

Predictions of avian *Plasmodium* expansion under climate change. Sci Rep 3:1126 doi:

10.1038/srep01126

Lutz HL, Hochachka WM, Engel JI, Bell JA, Tkach VV, Bates JM, Hackett SJ, Weckstein JD

(2015) Parasite prevalence corresponds to host life history in a diverse assemblage of

Afrotropical birds and haemosporidian parasites. PLOS ONE 10:e0121254 doi:

10.1371/journal.pone.0121254

Mangiafico SS (2024) rcompanion: Functions to support extension education program evaluation. R

package version 2.4.36 https://CRAN.R-project.org/package=rcompanion

Marroquin-Flores RA, Williamson JL, Chavez AN, Bauernfeind SM, Baumann MJ, Gadek CR,

Johnson AB, McCullough JM, Witt CC, Barrow LN (2017) Diversity, abundance, and host

relationships in the avian malaria community of New Mexico pine forests. PeerJ 5:e3700–

e3700 doi: 10.7717/peerj.3700

McNew SM, Barrow LN, Williamson JL, Galen SC, Skeen HR, DuBay SG, Gaffney AM, Johnson

AB, Bautista E, Ordoñez P, Schmitt CJ, Smiley A, Valqui T, Bates JM, Hackett SJ, Witt CC

(2021) Contrasting drivers of diversity in hosts and parasites across the tropical Andes. Proc

Natl Acad Sci 118:e2010714118 doi: 10.1073/pnas.2010714118

Medeiros MCI, Hamer GL, Ricklefs RE (2013) Host compatibility rather than vector–host-

encounter rate determines the host range of avian *Plasmodium* parasites. Proc R Soc B Biol

Sci 280:20122947 doi: 10.1098/rspb.2012.2947

Minchella DJ, Scott ME (1991) Parasitism: A cryptic determinant of animal community structure.

Trends Ecol Evol 6:250–254 doi: 10.1016/0169-5347(91)90071-5

Mordecai E A, Paaijmans KP, Johnson LR, Balzer C, Ben☐Horin T, De Moor E, McNally A,

Pawar S, Ryan SJ, Smith T C, Lafferty KD (2013) Optimal temperature for malaria

transmission is dramatically lower than previously predicted. Ecol Lett 16(1):22–30 doi:

10.1111/ele.12015

Mukhin A, Palinauskas V, Platonova E, Kobylkov D, Vakoliuk I, Valkiūnas G (2016) The strategy

to survive primary malaria infection: an experimental study on behavioural changes in

parasitized birds. PLOS ONE 11:e0159216 doi: 10.1371/journal.pone.0159216

Muriel J (2020) Ecophysiological assessment of blood haemosporidian infections in birds.

Ecosistemas 29 doi: 10.7818/ecos.1979

Names GR, Schultz EM, Krause JS, Hahn TP, Wingfield JC, Heal M, Cornelius JM, Klasing KC,

Hunt KE (2021) Stress in paradise: effects of elevated corticosterone on immunity and avian

malaria resilience in a Hawaiian passerine. J Exp Biol 224 doi: 10.1242/jeb.242951

Oakgrove KS, Harrigan RJ, Loiseau C, Guers S, Seppi B, Sehgal RNM (2014) Distribution,

diversity and drivers of blood-borne parasite co-infections in Alaskan bird populations. Int J

Parasitol 44:717–727 doi: 10.1016/j.ijpara.2014.04.011

Pérez￼Rodríguez A, Fernández￼González S, De La Hera I, Pérez￼Tris J (2013) Finding the

appropriate variables to model the distribution of vector￼borne parasites with different

environmental preferences: climate is not enough. Glob Change Biol 19:3245–3253 doi:

10.1111/gcb.12226

Perkins SL (2014) Malaria's many mates: past, present, and future of the systematics of the order

Haemosporida. J Parasitol 100:11–25 doi: 10.1645/13-362.1

Pigeault R, Chevalier M, Cozzarolo CS, Baur M, Arlettaz M, Cibois A, Keiser A, Guisan A, Christe

P, Glaizot O (2022) Determinants of haemosporidian single- and co-infection risks in

western palearctic birds. Int J Parasitol 52:617–627 doi: 10.1016/j.ijpara.2022.05.002

Poulin R (2011) Evolutionary ecology of parasites. Princeton University Press, Princeton, New

Jersey

R Core Team 2(023) R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria https://www.R-project.org/

Reinoso-Pérez MT, Dhondt KV, Dulcet H, Katzenstein N, Sydenstricker AV, Dhondt AA (2024)

Seasonal variation in detection of Haemosporidia in a bird community: a comparison of

nested PCR and microscopy. J Wildl Dis 60 doi: 10.7589/JWD-D-23-00023

Ricklefs RE, Fallon SM (2002) Diversification and host switching in avian malaria parasites. Proc

R Soc Lond B Biol Sci 269:885–892 doi: 10.1098/rspb.2001.1940

Ricklefs RE, Fallon SYM, Bermingham E (2004) Evolutionary relationships, cospeciation, and host

switching in avian malaria parasites. Syst Biol 53 doi: 10.1080/10635150490264987

Rodriguez MD, Doherty PF, Piaggio AJ, Huyvaert KP (2021) Sex and nest type influence avian

blood parasite prevalence in a high-elevation bird community. Parasit Vectors 14:145. doi:

10.1186/s13071-021-04612-w

Rodríguez-Hernández K, Álvarez-Mendizábal P, Chapa-Vargas L, Escobar F, González-García F,

Santiago-Alarcon D (2021) Haemosporidian prevalence, parasitaemia and aggregation in

relation to avian assemblage life history traits at different elevations. Int J Parasitol 51:365–

378 doi: 10.1016/j.ijpara.2020.10.006

Santiago☐Alarcon D, Palinauskas V, Schaefer HM (2012) Diptera vectors of avian

Haemosporidian parasites: untangling parasite life cycles and their taxonomy. Biol Rev

87:928–964 doi: 10.1111/j.1469-185X.2012.00234.x

Sheppard P, Comrie A, Packin G, Angersbach K, Hughes M (2002) The climate of the US

Southwest. Clim Res 21:219–238 doi: 10.3354/cr021219

Sibley CG, Monroe J (1990) Distribution and taxonomy of birds of the world. Yale University

Press, New Haven, Connecticut

Slowinski SP, Geissler AJ, Gerlach N, Heidinger BJ, Ketterson ED (2022) The probability of being

infected with haemosporidian parasites increases with host age but is not affected by

experimental testosterone elevation in a wild songbird. J Avian Biol 2022:jav.02819 doi:

10.1111/jav.02819

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinform 30:1312–1313 doi: 10.1093/bioinformatics/btu033

Stöver BC, Müller KF (2010) TreeGraph 2: combining and visualizing evidence from different

phylogenetic analyses. BMC Bioinform 11:7 doi: 10.1186/1471-2105-11-7

Sundar D (2006) binom: Binomial confidence intervals for several parameterizations. R package

version 1.1-1.1 https://CRAN.R-project.org/package=binom

Tang Y, Horikoshi M, Li W (2016) ggfortify: Data visualization tools for statistical analysis results.

https://CRAN.R-project.org/package=ggfortify

Valkiūnas G (2005) Avian malaria parasites and other Haemosporidia, 1st edn. CRC Press, Boca

Raton

von Humboldt A, Bonpland A (2009) Essay on the geography of plants – with a physical tableau of

the equinoctial regions (1807). In: Jackson ST (ed) Essay on the geography of plants.

University of Chicago Press, Chicago, pp 61–155

Waldenström AJ, Bensch S, Hasselquist D, Östman Ö, Pryor GS, Greiner EC, Hall B (2004) A new

nested polymerase chain reaction method very efficient in detecting *Plasmodium* and

Haemoproteus infections from avian blood. J Parasitol 90:191–194 doi: 10.1645/GE-

3221RN

Walther EL, Carlson JS, Cornel A, Morris BK, Sehgal RNM (2016) First molecular study of

prevalence and diversity of avian haemosporidia in a Central California songbird

community. J Ornithol 157:549–564 doi: 10.1007/s10336-015-1301-7

Wickham H (2016) ggplot2, Use R! Springer International Publishing, Cham doi: 10.1007/978-3-

319-24277-4

Williamson JL, Wolf CJ, Barrow LN, Baumann MJ, Galen SC, Schmitt CJ, Schmitt DC, Winter

AS, Witt CC (2019) Ecology, not distance, explains community composition in parasites of

sky-island Audubon's Warblers. Int J Parasitol 49:437–448 doi:

10.1016/j.ijpara.2018.11.012

Wyckoff ST, Judkins TC, Nemeth NM, Ruder MG, Martin JA, Kunkel MR, Garrett KB, Adcock

KG, Mead DG, Yabsley MJ (2024) Surveillance for selected pathogens and parasites of

northern bobwhite (*Colinus virginianus*) from western Oklahoma, USA. 2018–20. J Wildl

Dis 60 doi: 10.7589/jwd-d-23-00102

Zamora-Vilchis I, Williams SE, Johnson CN (2012) Environmental temperature affects prevalence

of blood parasites of birds on an elevation gradient: implications for disease in a warming

climate. PLOS ONE 7(6):e39208 doi: 10.1371/journal.pone.0039208

Tables
Table 1

Top five infection probability fixed effects additive models					
Bell's vireo ($n = 20$)					
Rank	*Predictors*	*K*	*AICc*	*ΔAICc*	*Weight*
1	tempPC2 + precipPC1	3	22.6	-	0.324
2	tempPC1	2	23.6	*0.967*	0.200
3	precipPC1	2	24.7	*0.348*	0.113
4	tempPC1 + precipPC1	3	25.9	*0.199*	0.064
5	Elevation + tempPC1	3	25.9	*0.197*	0.064
gray vireo ($n = 170$)					
Rank	*Predictors*	*K*	*AICc*	*ΔAICc*	*Weight*
1	Elevation + tempPC1	3	204	-	0.419
2	Elevation + precipPC1 + tempPC1	4	206	*1.72*	0.177
3	Elevation + precipPC2 + tempPC1	4	206	*1.73*	0.177
4	Elevation + tempPC1 + tempPC2	4	206	*2.07*	0.149
5	Elevation + tempPC1 + tempPC2, precipPC1 + precipPC2	5	207	*3.35*	0.078
plumbeous vireo ($n = 58$)					
Rank	*Predictors*	*K*	*AICc*	*ΔAICc*	*Weight*
1	Null model	1	52.1	-	0.301
2	precipPC2	2	52.4	*0.409*	0.246
3	precipPC1	2	53.3	*1.17*	0.168
4	tempPC2	2	53.4	*1.29*	0.158
5	PrecipPC1 + precipPC2	2	53.9	*1.73*	0.127

845 [2]

---

[2] **Table 1.** Top five models assessing environmental predictors of haemosporidian infection for each species. Sample sizes for datasets and directionality of best model outputs are listed at the top. Models were ranked by AICc, with ΔAICc values relative to the top model and corresponding Akaike weights (Weight) indicating model support.

Table 2

Param	Bell's vireo				gray vireo				plumbeous vireo			
	β	SE	LCL	UCL	β	SE	LCL	UCL	β	SE	LCL	UCL
Intercept	0.85	0.88	-0.99	2.7	1.1	0.36	0.43	1.8	1.7	0.40	0.98	2.5
Elevation	0.06	0.37	-0.70	0.82	-1.7	0.76	-3.2	-0.21	-	-	-	-
tempPC1	-0.94	1.5	-3.9	2.0	1.1	0.55	0.01	2.1	-	-	-	-
tempPC2	1.8	2.7	-3.5	7.1	-0.05	0.72	-1.5	1.4	0.05	0.18	-0.31	0.41
precipPC1	2.8	3.5	-4.2	9.8	-0.16	0.59	-1.3	1.0	0.10	0.24	-0.38	0.58
precipPC2	-	-	-	-	-0.28	0.90	-2.1	1.5	0.17	0.31	-0.44	0.78

846 [3]

---

[3] **Table 2**. Standardized model-averaged regression coefficients used to estimate effects of predictors and precision of effects across species-specific candidate model sets that assess environmental predictors of haemosporidian infection. Regression coefficients (β), standard error (SE), and 95% confidence limits (lower as LCL, upper as UCL) are given. Dashes indicate that the parameter was not present in a final model set, and NA values indicate that the parameter was not tested due to lack of variation present in model set.

Figure Legends

Figure 1) A) Sampling distribution for each host species (Bell's vireo: *Vireo bellii,* gray vireo: *V.*

vicinior, and plumbeous vireo: *V. plumbeus*) in the southwestern USA; B) Left: summary of

infection prevalence, with significant differences between groups indicated by letters. Error bars

represent 95% confidence intervals for positive infection proportions. Right: percent parasitemia for

gray and plumbeous vireos. C) Proportion of individuals infected versus uninfected across

elevational gradient. Asterisk denotes significant p-values (< 0.05); D) Phylogeny of

haemosporidian cyt *b* haplotypes and their respective abundances by host. Stars denote novel

haplotypes and bootstrap values >50 are shown at nodes. Bird images courtesy Birds of the World

(Billerman et al., 2025).☐ ☐

Figure 2) Estimate of haemosporidian lineage diversity in Bell's (*Vireo belli*), gray (*V. vicinior*)

and plumbeous vireos (*V. plumbeus*) based on iNEXT rarefaction and extrapolation. Bell's vireo

curves (yellow) were created using 12 avian haemosporidian infections and three haplotypes; gray

vireo curves (gray) used 123 avian haemosporidian infections and four haplotypes; plumbeous vireo

curves (blue) used 62 haemosporidian infections and 14 haplotypes. Points indicate the reference

sample, solid line the rarefaction estimate, and dotted line the extrapolation. The analysis suggested

that sampling approximately 33 infections in Bell's, 14 in grays, and 364 total infections for

plumbeous vireos would capture >95% of the haemosporidian lineage diversity in this community.

Total haplotype richness, rounded to nearest whole number, is estimated to be four for Bell's vireos

(95% CI [3-6]), five for gray vireos (95% CI [4 – 6]) and 54 for plumbeous vireos (95% CI [14 –

101]).☐

Figure 3) Intraspecific distribution of infected and uninfected vireos across associated

environmental variables. Higher tempPC1 values relate to cooler overall temperatures and higher

precipPC2 relates to sites with more seasonally stable precipitation. Significantly different

distributions are denoted by an asterisk (p < 0.05).☐