Multiple protein-protein interactions drive the assembly and budding of the Chikungunya virion

Miguel Angel Coronado-Ipiña[1,2,#], Shuyang Zhang[3,#], Siyu Li[3++], Pedro Gerardo Hernández-Sánchez[1], Denisse Cadena-Lopez[1,2,+], Vanessa Labrada-Martagón[5], Ajay Gopinathan[6], Suchetana Mukhopadya[7], Roya Zandi[3*], Mauricio Comas-Garcia[1,4,5,*]

[1]High-Resolution Microscopy Section, Research Center for Health Sciences and Biomedicine, Autonomous University of San Luis Potosi (UASLP), Mexico

[2]Life Sciences Graduate Program, Department of Sciences, UASLP, Mexico

[3]Department of Physics and Astronomy, University of California, Riverside, Riverside, CA 92521, USA.

[4]Molecular and Translational Medicine Section, Research Center for Health Sciences and Biomedicine, UASLP, Mexico

[5]Department of Sciences, UASLP, Mexico

[6]Department of Physics, University of California, Merced, CA 95343, USA.

[7]Department of Biology, Indiana University Bloomington

+ Current address: ThermoFisher Mexico

++ Current address: Department of Physics and Astronomy, California State Polytechnic University, Pomona, California 91768, USA.

*Correspondence: royaz@ucr.edu and mauricio.comas@uaslp.mx

#These two authors contribute equally

ABSTRACT

The assembly of enveloped viruses is a highly orchestrated process that depends on the coupling of multiple protein–protein interactions within a membrane environment. To gain mechanistic insight into this process, we use Chikungunya virus as a model system to study Alphavirus

assembly, focusing on the interplay between core–spike and spike–spike interactions. We begin with coarse-grained molecular dynamics simulations to systematically explore how the symmetry of the nucleocapsid core, together with the relative strengths of spike–core and spike–spike interactions, influences budding efficiency and the emergence of icosahedral particle symmetry. Building on these computational predictions, we perform site-directed mutagenesis on Chikungunya virus 181/25 and examine the consequences for particle assembly and budding in cultured cells, as well as the impact of these mutations during *in-cellulo* assembly. Our results reveal that canonical core–spike interactions, while necessary, are not sufficient for successful assembly. Instead, lateral interactions among glycoproteins emerge as critical determinants of efficient budding, particle stability, and the maintenance of icosahedral symmetry. Together, these findings provide an integrated computational and experimental framework for understanding the molecular principles governing Alphavirus assembly.

KEYWORDS: Enveloped virion budding, glycoprotein steric repulsions, Alphaviruses, Molecular Dynamics simulations, multiple protein interactions

The assembly of enveloped viruses is a complex process orchestrated by protein–protein and protein–membrane interactions to drive membrane deformation, budding, and packaging of the viral genome into a fully infectious particle[1-7]. For certain viruses, this process is known to rely primarily on matrix or capsid proteins that recruit host-cell machinery to mediate budding from the plasma membrane[6, 8-13]. In contrast, the mechanisms of assembly and budding in Alphaviruses remain less well characterized, with some conflicting models [2, 14-18].

Alphaviruses are arthropod-borne viruses infecting diverse animal species, classified as

New- or Old-World depending on whether they are endemic to the Americas or other regions [19].

Chikungunya virus (CHIKV), an Old-World Alphavirus, is the most prevalent Alphavirus

infecting humans and has caused recurrent epidemics of febrile illness and chronic polyarthritis

worldwide[20-25]. Alphaviruses have a spherical virion approximately 70 nm in diameter,

consisting of two concentric icosahedral protein shells separated by a lipid bilayer derived from

the host plasma membrane. The outer shell comprises 240 copies of E1–E2 heterodimers

arranged as 80 trimeric spikes, while the inner shell, nucleocapsid or core (NC), consists of 240

capsid proteins (CP) encapsidating the ~12 kb single-stranded positive-sense RNA genome[14].

These two layers are structurally coupled via interactions between the cytoplasmic tail of E2 and

a hydrophobic pocket in the capsid protein[14, 26].

Cryo-electron microscopy (cryo-EM), cryo-electron tomography (cryo-ET), biochemical

analysis, and molecular dynamics (MD) simulation studies of Alphaviruses have revealed four

major classes of interactions stabilizing the virion: (1) intra-dimer interactions[3, 4], (2) inter-dimer

E1–E2[2, 16, 27-29] , (3) inter-spike trimer–trimer interactions[30, 31], and (4) vertical contacts between

E2 and the capsid core[26, 30, 32-34]. These interactions coordinate virion formation—from initiating

budding and guiding icosahedral assembly to controlling particle yield, stability, and

infectivity[14, 35, 36] —though their precise roles and mechanisms remain unclear.

A central question in Alphavirus biology is how these two icosahedral layers assemble

coordinately during budding. Several models have been proposed: in the assembly-from-within

model, a preformed icosahedral core serves as a structural template for glycoprotein

assembly [15, 17]; in the assembly-from-without model, glycoproteins initiate budding and recruit

the capsid core *post hoc*[16]; and in the concerted assembly model, an initially irregular

cytoplasmic core gains order upon interacting with glycoproteins at the membrane[2, 18]. The latter

has gained support from recent cryo-ET studies, which show that cores lack icosahedral

symmetry in the cytoplasm but acquire structural order during budding[2, 18]. These studies did

not observe a pre-assembled lattice of spikes at the membrane or icosahedral cytoplasmic cores,

suggesting that E2–core binding and lateral glycoprotein interactions drive budding and shape

the geometry of the final virion. However, in the absence of direct, molecular-resolution

visualization of both the budding process and spike organization under budding-inhibited

conditions in infected cells, the mechanisms underlying icosahedral assembly and its inhibition

remain poorly defined.

To understand the factors contributing to Sindbis (Alphavirus) budding, Hagan and co-

workers developed a coarse-grained simulation model that incorporates spike–spike and spike–

core interactions[30]. They showed that while glycoprotein interactions alone can drive budding,

the resulting particles tend to be polydisperse and irregular. Even modest interactions with a rigid

core improve uniformity and reduce kinetic trapping. However, their focus was on the conditions

that permit or impede budding, rather than on achieving icosahedral symmetry in the final

product.

Here, we identify the molecular determinants that govern CHIKV budding and assembly

by pinpointing the specific interactions that coordinate these processes. One question that

naturally arises is whether capsid proteins (CPs) preassemble into icosahedral nucleocapsids

(NCs) that act as structural templates, or whether glycoprotein spikes instead drive the

coordinated formation of both the outer spike lattice and the inner NC shell. To address this, we

performed coarse-grained MD simulations to examine how spike–core and spike–spike

interactions, together with core symmetry, influence budding efficiency and the emergence of

icosahedral order. In particular, we tested whether immature, non-icosahedral NCs can serve as

imperfect scaffolds that promote the formation of icosahedral spike shells.

In our MD simulations, we find that when the core lacks icosahedral symmetry and is

constrained to a fixed geometry that prevents local dissociation and reassociation, the spikes fail

to assemble into a complete icosahedral shell. Nevertheless, the absence of an entire pentamer

does not completely abolish symmetry; spike assembly can still progress toward icosahedral

order, albeit with reduced efficiency. These results strongly support a concerted assembly

mechanism, in which tightly coordinated interactions across the membrane synchronize the

formation of both the inner nucleocapsid and the outer spike lattice.

Guided by these insights, we conducted targeted mutational experiments using a CHIKV

reporter virus to probe the functional roles of key glycoprotein interfaces. Specifically, we

introduced E2 Y400K/L402R to disrupt spike–core interactions, E2 N263Q to weaken

intertrimer contacts, and E1 K245N/R247T to perturb intradimer interactions. Our combined

computational and experimental findings demonstrate that E2–core interactions are essential for

initiating membrane curvature and nucleating particle formation, while strong lateral spike–spike

contacts are required for the completion and stabilization of fully formed virions. Thus, both

vertical (spike–core) and lateral (spike–spike) interactions act cooperatively to establish and

maintain icosahedral symmetry across the two structural layers of the virus. This cooperative

mechanism ensures proper virion architecture and infectivity and highlights potential antiviral

targets aimed at disrupting envelope-protein organization to block particle assembly or release.

RESULTS

Motivated by several experiments on CHIKV budding and the formation of icosahedral symmetry, carried out by Chiu's group [2, 18, 37] and others[27, 30, 38-40], we performed molecular dynamics (MD) simulations to investigate how specific spike–spike and spike–core interactions govern virus assembly. This approach isolates the effects of individual interactions in a controlled, simplified environment, and the resulting insights guided targeted mutational experiments using CHIKV 181/25 [41] to test their roles in particle formation in cultured cells— results presented in the sections following the simulations.

Simulations

We simulated the budding of a nucleocapsid through a membrane that, as described in the *Methods and Materials* section, exhibits fluid behavior[41] and contains inserted spike proteins (Figure 1A). Each spike comprises three domains: the ectodomain (ED), transmembrane domain (TMD), and cytoplasmic domain (CTD). The nucleocapsid is represented by a core with different geometries, illustrated in Figures 1B–1D, which can interact with the membrane spikes.

Figure 1. *Schematic representation of a coarse-grained spike protein and core.* A) The atomic structure of the CHIKV spike protein, a trimer of heterodimers (PDB ID: 8FCG). The shaded circles represent the components of the coarse-grained model, which consists of three spherical particles: ED (ecto domain,

blue), TMD (transmembrane domain, red), and CTD (cytoplasmic domain, yellow). The size asymmetry

between the domains allows the spike proteins to adopt a curved configuration, promoting membrane

bending. **B)** A simple spherical core. **C)** A spherical core decorated with 80 triangular units arranged with

icosahedral symmetry. **D)** A modified version of the core in which several triangles are missing, breaking

the icosahedral symmetry.

Spike–spike interactions alone were first tested to determine whether they could drive

budding in the absence of a core. Consistent with earlier work [29, 30, 42, 43], these interactions were

sufficient to induce budding but produced irregular, non-icosahedral structures. Introducing

a spherical core that interacts attractively with the spike cytoplasmic tails (see *Methods and*

Materials for potential details) improved local ordering but failed to yield global icosahedral

symmetry(see Supplementary Figure 3). Only when a preassembled icosahedral core was

incorporated (Figure 1C) did the system produce a symmetric, virus-like shell.

In this configuration, three interaction classes determined assembly behavior: (i) CTD–

core attraction (ε_{mc}), (ii)TMD–TMD lateral association (ε_{mm}), and (iii) excluded-volume

repulsion among spike ectodomains (ED–ED). By systematically varying ε_{mc} and ε_{mm} we

identified conditions that maximize budding efficiency and yield near-perfect icosahedral

structures. The resulting trends were compared to experiments using CHIKV 181/25 infection

and transfection of HEK-293T cells expressing a CMV-driven reporter gene. We note that

mutations at the intradimer interface were introduced experimentally and shown to affect

heterodimer interactions within a trimer; however, these were not modeled in our simulations,

which treat the trimer as the smallest structural unit in a coarse-grained framework.

Parameter Optimization for the Wild-Type System

Systematic exploration of the parameter space revealed an optimal regime corresponding to a

CTD–core interaction strength of $\varepsilon_{mc} = 74\,k_BT$, together with the default ED size and TMD–

TMD attraction (values listed in the figure caption). This condition, designated as the wild-type

(WT) baseline, yields an effective E2–core interaction of approximately $8\,k_BT$, consistent with

previous studies[30]. Under these conditions, budding proceeds efficiently, with about 80±5 spike

assemblies on the core to form a nearly icosahedral shell (Figure 2A). The inset in Figure 2B

highlights the final spike distribution (yellow spheres), and Figure 2C presents a cutaway view

showing their spatial arrangement around the core. Although the model lacks an explicit scission

mechanism, budding terminates naturally once the shell closes around the core.

Figure 2. *Snapshots of the budding process.* **A)** Snapshots of the budding process using an icosahedral

core at timesteps t=0,5,10, 80,250, and 700 (left to right, top to bottom). The system parameters include a

CTD–core interaction strength of 74 k_BT and a TMD-TMD interaction strength of 0.2 k_BT. The particle

diameters are set to 1.15 for ED, 0.8 for TMD, and 0.4 for CTD. **B)** Final state of the system after

budding. The inset highlights the location of CTD particles along with the core. **C)** Cross-sectional view

of the final state through the center of the assembled structure.

At weaker interaction strengths, however, this behavior changes markedly. When ε_{mc} falls

below a critical threshold, the core fails to bud from the membrane—consistent with previous

observations [17, 26, 28, 32, 33]. In this regime, the membrane remains unbudded (see Supplementary

Figure 4), reproducing the phenotype observed experimentally for the CT mutant described

below.

Modeling the Effects of Mutations by Tuning Attractive and Steric Interactions Between

Spike Protein Domains

Ribeiro-Filho *et al*. proposed that in the Mayaro virus, the glycosylation site equivalent to

CHIKV E2 N263 acts as a "*molecular handshake*" that stabilizes the viral particles[31]. This

glycosylation could influence trimer–trimer interactions through steric repulsions or hindrances,

and its removal may alter the local curvature or relative orientation between adjacent spikes,

thereby making budding more difficult.

To quantify the absence of this glycosylation site in E2, we conducted a series of

simulations with icosahedral cores and compared their relative success rates. To isolate the

effects of specific mutations, we introduced changes by modifying one or more of these

parameters, while holding the CTD–core interaction constant at $\varepsilon_{mc} = 74\,k_BT$, same as the WT

system interaction strength. For each condition, we performed eight independent simulation

trials. In the WT system ($\varepsilon_{mc} = 74\,k_BT$), 7 out of 8 simulations resulted in successful

assemblies, which we define as a 100% baseline success rate. A configuration was considered

successful if no more than five of the 80 spike proteins, expected in a perfect T = 4 structure,

were missing or mispositioned. This definition is consistent with recent Cryo-ET data showing

that Alphavirus budded virions can have defects at one pole of the particle[18].

As shown in Supplementary Table 1, reducing the ED size while keeping the CTD–core

fixed at 74 k_BT resulted in a drop in the success rate to approximately 57.1%. Eliminating the

TMD–TMD interactions while maintaining the WT ED size also reduced the success rate to

about 28.6%, highlighting the importance of attractive interactions between spike proteins. When

both the ED size was reduced and the TMD–TMD interaction was removed, the success rate

declined even further to 14.3%. These results suggest that decreasing ED size disrupts assembly,

likely by impairing the induction of membrane curvature, and that the absence of TMD–TMD

attraction further compromises proper assembly. Our findings indicate that both steric and

attractive interactions among spike proteins are essential for efficient and robust assembly. These

simulations reflect the behavior of the mutant ITS described in the experimental section.

We next investigated how changes in the CTD–core interaction strength affect assembly

outcomes in the mutated systems described above. Achieving the right balance is critical: an

overly strong attraction can lead to excessive binding of spike proteins to the core, while a too

weak interaction may fail to bring in enough spikes for successful assembly. To explore this, we

performed additional simulations in which CTD–core strength was varied while holding the

other parameters fixed at their mutated values. This allowed us to assess how changing the

interaction strength affects budding efficiency under each mutation scenario.

Supplementary Table 2 summarizes these findings. In the absence of TMD–TMD

interaction, increasing CTD–core strength from 74 to 80 k_BT improved the success rate to 85.7%

relative to WT. When only the ED size was reduced and the CTD–core was set to 75 k_BT, the

success rate reached 71.4%. However, when both the ED size was reduced and the TMD

interaction was removed, success fell again to 57.1%. Notably, while stronger CTD–core

attraction improves assembly efficiency, it also results in excessive spike adsorption in the WT

system, underscoring the importance of a finely tuned interplay among all interactions for

successful T=4 structure formation.

221   Budding of an Imperfect Icosahedral Core Through the Membrane

Recent experimental studies have shown that the CHIKV capsid does not exhibit perfect

icosahedral symmetry during budding[2]. To investigate this effect, we constructed a core missing

one pentamer, as shown in Figure 1D, introducing an imperfect geometry. As described later, we

tested this core under both wild-type conditions and in simulations incorporating mutations that

disrupt CTD–core and spike–spike interactions.

Compared with the perfect icosahedral core, the imperfect core yielded a lower success

rate of budding. Nevertheless, under optimized conditions—using ED = 1.15, a CTD–core

interaction strength of 77 k_BT, and a TMD-TMD = 0.1 k_BT —spikes near the site of the missing

pentamer remained correctly positioned (Figure 3), which shows the budding of an imperfect

core through the plasma membrane. Reducing the ED size or weakening the TMD–TMD

interactions disrupted the T = 4 symmetry by impairing inter-trimer contacts, further lowering

assembly efficiency. For the optimal values of the interaction energies, only 2 out of 12

simulations produced a successful configuration (~17%), and the success rate declined further

when both the ED size and the TMD–TMD interactions were reduced.

Figure 3. *Configurations of the membrane and an imperfect core with icosahedral symmetry.* (A) Initial configuration of the system with an imperfect core. (B) Final configuration after budding; the inset shows the CTD particles together with the core. System parameters: CTD–core interaction strength = 77 k_BT , TMD–TMD interaction strength = 0.1 k_BT. Particle diameters are 1.15 for ED, 0.8 for TMD, and 0.4 for CTD.

The relatively low success rate observed in our simulations, compared to experimental results[2], likely stems from the limitations of the current model, which has a fixed core size and immobile subunits. Experimental data suggest that the initial core is smaller than the final assembled structure[2], indicating that core subunits can locally dissociate and reassociate to achieve near-perfect icosahedral symmetry. In this study, our goal was to determine whether spike proteins can still assemble into an icosahedral shell in the presence of core defects, and to evaluate the roles of spike–spike and spike–core interactions in this process.

Capturing the dynamic remodeling of the core—from an imperfect to an icosahedral

symmetric structure—requires a model that explicitly allows internal rearrangement of capsid

subunits[44]. Simulating this process is computationally intensive, as it involves dissociation and

reassociation events that must overcome significant free energy barriers[45]. However, such local

restructuring is essential for correcting geometric imperfections during budding and achieving

stable icosahedral symmetry. In future work, we plan to develop models that incorporate core

flexibility and explicitly examine the role of the genome[46] in promoting protein rearrangement

and structural maturation.

While the theoretical model and simulations provide mechanistic insight into how

specific interactions contribute to CHIKV assembly, they necessarily simplify certain aspects of

the biological system, such as membrane heterogeneity and the dynamic behavior of viral

proteins. To assess the biological relevance of our findings and examine how these interactions

function in a cellular context, we next designed targeted mutational experiments to test the roles

of glycoprotein–glycoprotein and glycoprotein-core interactions in budding and infectivity.

The infectivity of CHIKV depends on multiple interactions between glycoproteins.

To complement and validate the simulation predictions, we constructed three CHIKV mutants.

To disrupt intertrimer interactions, we removed a glycosylation site in E2 via the mutation

N263Q (intertrimer surface mutant, or ITS), and to weaken spike–core interactions, we

introduced a double mutation in E2 Y400K/L402R (cytoplasmic tail mutant, or CT). To disrupt

the intradimer (E1/E2) surface, we introduced a glycosylation site in E1 at K245N/R247T

(intradimer surface mutant, or IDS). The ITS and CT mutants correspond to interaction changes

examined in our coarse-grained simulations. The effect of the IDS mutation was not studied in

the simulations but was included experimentally to further test the role of spike–core coupling.

Figure 4 shows a Cryo-EM reconstruction (PDB 3j2w) of the asymmetric subunit of

CHIKV, depicting the positions of amino acids targeted for mutation. To investigate the role of

the E1-E2 heterodimer interface, we generated an E1 double mutant K245N/R247T (CHIKV

IDS) (Figure 4A) that creates a glycosylation site at the interface between E1 and E2. This is

expected to introduce steric hindrance that weakens and/or destabilizes the E1/E2 heterodimer.

For other Alphaviruses, it has been shown that replacing the hydrophobic amino acids in E2 that

interact with the core with basic residues can inhibit budding[26, 47]. To this end, we generated the

E2 double mutant Y400K/L402R to block the interaction between the core and the glycoproteins

(Figure 4B) (CHIKV CT). Finally, we mutated N263 to a Q in E2 (CHIKV ITS) to remove the

glycosylation site at the outer part of the trimer of heterodimers (Figure 4C). The goal was to

decrease the interaction between trimers by removing the carbohydrates at this interface; we

hypothesize that these carbohydrates stabilize the trimers through steric repulsion. Figure 4D is a

schematic representation of the three interactions studied here, and Figure 4E depicts some of the

most likely effects of the mutations on these interactions.

Figure 4. *Structure of Chikungunya virus* A-C) Cryo-EM (PDB 3j2w) structure of the asymmetric unit of CHIKV virion with E1, E2, and C in salmon, gray, and teal. A) E1 K245 is at the interface between E1 and E2. B) The association of E2 with C is driven mainly by a hydrophobic pocket of E2 and Capsid. C) E2 N263 is glycosylated and might result in steric interactions between two E2 proteins at the trimer-trimer interface. D-E) Schematic representation of the interaction between glycoproteins and the core during assembly and budding. D) E1 and E2 dimerize in the cytoplasm, and once the glycoproteins are at the plasma membrane, they interact with the CHIKV cytoplasmic core, then the neighboring trimers interact with each other through high-molecular weight sugars at the E2 N263 site, E) In CHIKV IDS, a new glycosylation site was introduced in E1 amino acid 245; this mutation aims to decrease the E1/E2 heterodimer formation. CHIKV CT was designed to inhibit the interaction between E2 and the core by removing the hydrophobic pocket in E1 Y400/Y402. In CHIKV ITS, the N263 glycosylation site was removed to weaken the trimer-trimer interactions.

Determining the relative infectivity of mutants compared to WT using a reporter virus system.

To determine the effect of each mutation on virion assembly and budding, we transfected HEK-

293T cells with CMV-driven CHIKV plasmids based on the attenuated strain CHIKV 181/25

(hereafter referred to as WT). These plasmids contain a reporter gene between the C and E3

genes, which is post-translationally cleaved. Therefore, the amount of reporter gene is

proportional to the number of viral structural proteins. Then, HEK-293T cells were infected with

the virus rescued by transfection. At 24 hours post-infection (h.p.i.), the amount of virus

produced was determined by fluorescence microscopy as a relative infectivity to the WT. Each

infection was done in triplicate and with four biological replicas.

Figure 5A-D shows representative fluorescence images of infected cells (mKate2-

positive). Figure 5A shows that at 24 h.p.i. most cells are mKate2-positive cells (WT infected),

while panels B, C, and D show that the number of infected cells with CHIKV IDS, CT, and ITS

is extremely low compared to the WT.

Figure 5. *The infectivity of the viral particle depends on several protein-protein interactions.*

Representative fluorescent micrographs of HEK-283T cells infected with WT, IDS, CT, and ITS CHIKV

viruses. The genome of all these viruses has mKate2 as a reporter gene between the capsid and E3. A-D)

Cells infected with the WT, IDS, CT, and ITS viruses, respectively. E) Relative infectivity of each mutant

with respect to the WT virus. The medians represent values from four experiments (n=4). K-W analysis

with Dunn's post hoc showed differences in infectivity for the three mutants compared to the WT. The

mutants showed a decrease in the relative infectivity of approximately 50%, 90%, and 100%, respectively,

for ISD, CT, and ITS (* p<0.05). F) Viral infection kinetics. Infections with WT, IDS, and ITS reach a

maximum at 48 hours post-infection. The bars indicate the standard deviation.

The relative infectivity of these mutant CHIKV viruses is shown in Figure 5E. CHIKV

IDS and CT are about 90% and 100% less infectious than the WT, respectively, while CHIVK

ITS is about 50% less infectious than the WT. These results show that blocking the interaction

between the core and E2 (CT) reduces relative infectivity to below the detection limit. It has

been proposed that E2-core interaction is necessary for assembly [47]. However, the data in Figure

5E suggest that this interaction is insufficient; altering, at least, the interactions between

monomeric E1 and E2 and between trimers also decreases the infectivity.

These data suggest that the CHIKV mutants exhibit lower infectivity than the WT, likely

due to defects in assembly and/or particle release. However, it is also possible that these

mutations affect genome replication. To address this, infections with CHIKV WT, IDS, CT, and

ITS were monitored over 72 hours, with measurements taken every 12 hours. Figure 5F shows

the normalized average area of infected cells from three independent experiments. These results

indicate that the peak of normalized relative infectivity occurs at 48 h.p.i. for CHIKV WT, IDS,

and ITS. In contrast, CHIKV IDS and CT mutants show minimal, if any, replication. Therefore,

the differences in infectivity between the CHIKV WT, ITS, and IDS are not due to differences in

the kinetics of replication but are related to assembly/budding defects.

Budding of CHIKV particles depends on multiple interactions.

To determine which stage of the infectious cycle is affected by these mutations, we analyzed

transfected cells by thin-section transmission electron microscopy (TEM) (Supplementary Figure

5), focusing on how these mutations affect virion assembly and budding. CHIKV IDS and CT

exhibit 90% and 100% reductions in infectivity relative to WT, respectively, making it extremely

challenging to identify infected cells. Nonetheless, cells infected with Alphaviruses can often be

identified by the presence of cytopathic vesicle type I (CPV-I), a characteristic marker

observable by TEM[7]. Supplementary Figures 6A-D show the presence of CPV-I (arrows) in cells

infected with the four types of viruses. Once these vesicles were found, we searched the area

around the plasmatic membrane of that cell to characterize any possible assembly and or budding

process.

Figure 6 shows characteristic budding events that result in the doubled-layer icosahedral

viral particles. Figure 6A shows a CHIKV WT virion budding (red arrow) and an extracellular

virion. CHIKV IDS has a relative infectivity of 90% and most budding particles are "stuck" in

the initial budding stages, forming a group or collection of incomplete budding structures close

to each other. (Figure 6B). The presence of partial virions suggests that this mutation did not

completely block E1/E2 heterodimer formation. As expected, we could not find any budding

particles for CHIKV IDS (Figure 6C). Figure 6D shows one extracellular CHIKV ITS particle

and two still bound to the plasmatic membrane.

Figure 6. *Budding structures of WT and mutant viral particles.* Thin-section transmission electron

micrographs of HEK-293T cells infected with the WT, IDS, CT, and ITS, CHIKV viruses showing

budding events of the virions (A, B, and D). The WT, IDS, and ITS viruses bud by forming a neck, which

will eventually be pinched off, releasing the virus into the extracellular medium (arrows). C) CHIKV CT

does not bud out, and all the cores remain in the cytoplasm (red oval).

Virion morphology depends on the strength of the interactions between the glycoproteins.

To determine whether weakening the interaction between trimers by removing the "*molecular*

handshake" between them (CHIKV ITS) alters particle morphology, we measured the

hydrodynamic diameter of purified CHIKV WT and ITS particles using dynamic light scattering

(DLS). Figures 7A and 7B show the hydrodynamic diameter distribution for two independent

batches of CHIKV WT and ITS, respectively. The broken and solid lines indicate the

measurements for batches 1 and 2, respectively. The average hydrodynamic diameters for the

WT and ITS CHIKV virions are 67 nm +/- 3 nm and 113 +/- 10 nm, respectively. In addition,

these experiments indicate that CHIKV ITS virions are more heterogeneous than CHIKV WT

virions. This observation is consistent with the trends shown in Figure 7 and further supports our

proposed mechanism. Figures 7C and D show positive-stain TEM images of purified CHIKV

WT and ITS from batch 1, respectively.

Figure 7. *Removing the E2 N623 glycosylation site increases particle heterogeneity.* The hydrodynamic

diameter (nm) of purified virions was determined by using dynamic light scattering (DLS). Three

measurements of the hydrodynamic diameters of the WT virus (A) and ITS (B) from two independent

batches (broken lines are batch 1 and solid lines are batch 2. For the WT distribution of the hydrodynamic

radius is homogenous (67 nm +/- 3 nm), while the ITS is heterogeneous (113 +/- 10 nm). C-D) Negative-

staining TEM images of the purified samples for batch 1. The scale bar is 100 nm.

The data shown here suggest that altering the interaction between glycoproteins decreases

the ability of the particles to bud from the producer cells. Therefore, we analyzed the cellular

localization of 500 virions/cores for CHIKV WT, IDS, and ITS of transfected cells by thin-

section EM. These localizations were defined as cytoplasmic, budding, and extracellular

particles; the cytoplasmic particles are cores that do not interact with the plasma membrane, the

budding particles are those forming any neck or protrusion, and the extracellular particles are the

released virions. The distribution of virions in Supplementary Figure 7 shows that most CHIKV

WT particles are either at the budding stage or released (extracellular). Very few particles remain

as cytoplasmic cores. This distribution shows that when the E1/E2 heterodimer interface is

altered, most particles are found at the plasma membrane at some stage of budding. This suggests

that the mutant's inability to form proper lateral interactions almost completely blocks the

particle from pinching off the plasma membrane. As expected, from the WT, IDS, and ITS

particles, the WT are the most abundant in the extracellular space, with only about 1/3 of the

CHIKVK ITS with respect to the WT are found outside the cell, which is consistent with the

relative infectivity data and the MD simulations (*i.e.,* Supplementary Table 1).

Based on our data, we hypothesize that carbohydrates at the interface between two

trimers generate steric repulsion that contributes to the icosahedral ordering of the trimers. This

steric interaction was first proposed for another Old-World Alphavirus, Mayaro virus, and

described as a "*molecular handshake*" between the glycans of two different trimers [31]. Therefore,

removing the E2 glycosylation site at N263 is expected to decrease the stability of the viral

particles. To test this, we analyzed the stability of the particles to heating by taking the

supernatant from transfected cells, dividing it into aliquots, incubating it from 37 °C to 60 °C for

30 minutes, cooling it to 37 °C, and using it to infect HEK-293T cells. The relative infectivity

was determined by normalizing the areas of fluorescent cells for each virus type with the sample

incubated at 37 °C for that virus type (Figure 8). The normalized relative infectivity of CHIKV

WT at 40ºC and 45ºC remained constant, whereas a significant drop in relative infectivity

occurred around 50ºC. CHIKV ITS is more sensitive to changes in temperature compared to a

sample of CHIKV ITS incubated at 37ºC. These data were fit to an equation that assumes a

cooperative dissociation process. This model has two parameters: the temperature at which the

relative infectivity decreases by 50% (T_{RI50}) and a cooperative exponent (n). The values of T_{RI50}

and n for CHIKV WT and ITS were 50ºC and 99, and 44ºC and 16.5, respectively. These

parameters suggest that removing the glycan in E2 N263 significantly affects the particle

stability and that the dissociation process of the virion of the CHIKV ITS is less ordered and

might involve a weaker interaction than CHIKV WT.

Figure 8. *The thermal stability of the virions depends on trimer-trimer interactions.* The relative

infectivity of the WT and ITS viruses was calculated with respect to the area and fluorescence of infected

cells for each virus at 37ºC. The circles and bars are the mean and standard deviation of three independent

experiments. The relative infectivity of WT remains constant from 37ºC to 45ºC, reaching a 50% decrease

at 50ºC. In CHIKV ITS decreased at 40ºC, reaching a 50% decrease at 45ºC. The data was fitted with a

cooperative model that follows Hill's equation (see methods and materials). For the WT, the T_{RI50} and n

are 50ºC and 99, respectively, while for the ITS mutant, they are 44ºC and 16.5.

DISCUSSION

In this paper, we set out to determine how specific protein–protein interactions govern CHIKV

assembly and budding. We hypothesized that the assembly and budding depend not only on the

canonical E2-core interactions [3, 26, 35] but also on other protein-protein interactions. To achieve

this, we combined coarse-grained simulations with site-directed mutagenesis targeting key

interfaces within the CHIKV glycoprotein layer.

Our simulations revealed that efficient budding requires a finely tuned balance between

spike–core attraction and lateral glycoprotein interactions. Although this is not the first model to

describe Alphavirus budding, it addresses novel interactions between the ectopic domains of

trimers that have not been previously explored or compared with experimental data[30].

First, interactions between the CTD of the spikes and the underlying NC were essential

for initiating membrane deformation. When the strength of this interaction was reduced below a

critical threshold, budding was arrested—consistent with experimental results for CHIKV CT,

which disrupts the hydrophobic pocket in E2 that binds to a hydrophobic pocket on the

nucleocapsid core—leading to complete loss of budding and virion assembly. The effect of this

mutation has been demonstrated elsewhere[3, 26, 35] and prompted the proposal that E2-core

interactions primarily drive Alphavirus budding. However, the simulations and experiments

presented here show that this interaction is necessary but not sufficient for assembly and

budding.

Beyond initiating budding, interactions among glycoprotein trimers were found to be

critical for achieving proper spatial organization. Simulations showed that spike–spike

interactions alone could not guide self-assembly toward the correct geometry, nor could a smooth

spherical core. However, an icosahedral core with T = 4 symmetry was sufficient to support the

correct positioning of glycoproteins, even when imperfect. Notably, the absence of ED-ED

interactions significantly reduced budding efficiency, mirroring experimental results for CHIKV

ITS, in which deletion of a glycosylation site weakens trimer–trimer interactions, leading to

virions with diminished infectivity, increased particle heterogeneity, and decreased particle

stability. Despite a previous suggestion that spike-spike proteins between ectodomains through

glycans in E2 N263 confer stability to the particle[31], the effect of this interaction has not been

previously explored either through simulation or experiments.

Although our model does not explicitly resolve E1–E2 heterodimerization, experimental

analysis of the CHIKV IDS mutant—which introduces a glycosylation site in E1 at the

heterodimer interface—revealed severe assembly defects. These particles were largely non-

infectious, rarely completed budding, and accumulated at the plasma membrane, indicating that

stable E1–E2 heterodimer formation is essential for trimer assembly and for progression beyond

the early stages of envelopment and release.

High-resolution cryo-EM maps of CHIKV[2 3] □ □□□□[2] do not show electron density

corresponding to high-molecular-weight glycans at E2 N263; however, our experimental data

reveal a structural role for this site, consistent with the "molecular handshake" model proposed

for Mayaro virus[31]. Deletion of the E2 N263 glycan disrupts trimer–trimer contacts and decreases

the cooperativity of disassembly upon heating, underscoring its importance in stabilizing the

spike lattice.

To further probe structural determinants of assembly, we examined how the geometry of

the capsid core influences budding. Simulations performed with a defective core lacking one

pentamer revealed that budding can still occur under favorable conditions, but with markedly

reduced efficiency compared to a fully symmetric core. These results align with cryo-ET

observations showing that one pole of the CHIKV virion can initiate budding despite local

structural irregularities, emphasizing the need for geometric compatibility between the

nucleocapsid and the glycoprotein lattice.

Previous work demonstrated that glycoprotein spikes alone cannot reliably form

icosahedral structures[3], whereas some studies suggest that capsid proteins can assemble into T

= 4 shells before budding[1] [1]. However, the symmetry of such pre-budding cores remains

uncertain. Recent cryo-ET analyses[1] show that when the core deviates from perfect symmetry,

spike–core and lateral spike–spike interactions can still guide assembly toward icosahedral order.

These observations, together with evidence that the CHIKV core transitions from an irregular

cytoplasmic structure to a more symmetric state during budding[2], suggest that local

rearrangements of capsid subunits are required for forming stable icosahedral particles. Although

our current model assumes a rigid, preassembled core and cannot capture this dynamic behavior,

incorporating such flexibility in future simulations—alongside ongoing site-directed mutagenesis

experiments of the core—will be crucial for elucidating the coupling between nucleocapsid

remodeling and membrane budding.

CONCLUSIONS

Our integrated computational and experimental study demonstrates that CHIKV assembly and

budding are governed by a coordinated network of protein–protein interactions operating across

multiple spatial and temporal scales. More significantly, this work highlights the power of

simulation-guided mutagenesis as a rational strategy for dissecting and reprogramming virus

assembly. By selectively altering key interaction interfaces, we gain deep molecular insights

without the need for atomistic simulations or exhaustive mutational screens.

Disrupting E2–core attraction, E1–E2 heterodimerization, or trimer–trimer contacts

impair assembly through distinct but complementary mechanisms—preventing budding,

disrupting trimer formation, or destabilizing the glycoprotein lattice. These results counter the

view that E2–core interactions alone are sufficient for particle formation; instead, they are

necessary but not sufficient. Our simulations emphasize the essential roles of both vertical (core-

glycoprotein) and lateral (inter-spike) interactions, including E1–E2 contacts and steric

repulsions between adjacent trimers, in enabling budding and the emergence of T = 4 icosahedral

symmetry.

While this study focused on perturbing E1–E2 and intertrimer interfaces, additional

interactions—such as E1–E1′ and E2–E2′ contacts—are also likely to contribute to lattice

stability and proper assembly. Introducing a glycosylation site into E1, as in CHIKV IDS,

significantly reduced infectivity, further underscoring the delicate balance of interactions

required for efficient particle formation.

A more comprehensive understanding of glycoprotein interactions will be essential for

building a complete and predictive model of alphavirus assembly—one that incorporates

mechanistic diversity across viral lineages and explains variations in budding efficiency and

infectivity between Old- and New-World alphaviruses. Taken together, our findings support a

unified model in which vertical and lateral interactions act in concert to shape virion formation,

offering a strategic foundation for therapeutic interventions that target the assembly and release

of infectious virus particles.

METHODS AND MATERIALS

Coarse-grained simulations

To simulate CHIKV budding through the plasma membrane, we developed a model incorporating a fluid membrane, spike proteins, and the viral core, as described below.

Fluid membrane: We model the biomembrane as a triangular lattice[48], as illustrated in Supplementary Figure 1. The energy of this network includes contributions from both bond stretching and bending[49]. The stretching energy is represented by a harmonic potential summed over all bonds:

$$E_s = \sum_{bonds} \frac{1}{2} k_s (r - a)^2, \; \textbf{Eq. 1}$$

here, $a = 10 \; nm$ is the equilibrium length of the spring. The parameter $k_s = 25 \; k_B T * nm^{-2}$ is the stretching modulus. The bending energy is calculated by summing over all pairs of triangular subunits that share an edge, and is given by:

$$E_b = \sum_{pairs} k_b \left(1 - \cos(\theta - \theta_0)\right), \textbf{Eq. 2}$$

where $\theta_0 = \pi$ is the preferred dihedral angle, and $k_b = 20 \; k_B T$ is the bending modulus.

To ensure the incompressibility of our coarse-grained membrane, we have placed a spherical LJ potential particle, denoted by M (see Supplementary Figure 1), with a diameter *a* at each membrane vertex. Membrane fluidity is achieved by performing Monte Carlo (MC) simulations with the bond-flipping method, which permits dynamic changes in the membrane's connectivity[50, 51]. The procedure for bond flips is illustrated in Supplementary Figure 2. Each MC step consists of N_b=8781 attempts, where N_b is the total number of bonds. In each attempt, a

randomly selected bond connecting two adjacent triangles is removed and reattached between

two previously unconnected vertices, as shown in Supplementary Figure 2. These moves are

designed to preserve the total number of vertices and bonds, thereby maintaining the overall

topology of the membrane. By allowing lateral rearrangements of connectivity, they facilitate the

diffusion of topological defects across the membrane.

Spike proteins: Each coarse-grained spike represents a trimer of heterodimers (Figure 1). It

comprises three components: the ED (ectodomain, blue), the TMD (transmembrane domain,

red), and the CTD (C-terminal or cytosolic domain, yellow). ED protrudes outward from the

membrane into the lumenal/extracellular-facing side of the budding virus; TMD is hydrophobic

and embedded within the lipid bilayer; and CTD is located on the cytosolic side, where it

interacts with the viral core.

Core Particles: The capsid of CHIKV is illustrated in Figure 1. In our model, we use three types

of core particles. In all three types of cores, the sphere part has a diameter of 3.2a (see Eq. 1).

The first is a simple spherical core (Figure 1B) that interacts with the C-terminal domain of the

spike proteins (the yellow domain in Figure 1A). The second core (Figure 1C) is also spherical

and interacts specifically with the CTD of the spike proteins. Its surface exhibits icosahedral

symmetry and consists of 80 triangular faces. Note that only the spherical surface directly

interacts with the spike proteins, while the triangles interact with them exclusively via steric

repulsion. This type of interaction promotes the assembly of spikes into a T=4 shell around the

core—an organization that would not occur otherwise. Without such interactions, the proteins

would form a shell around a spherical core lacking any particular symmetry, which would be

inconsistent with the experimental data. The third core (Figure 1D) is similar in shape and size

but lacks icosahedral symmetry due to the absence of a pentamer.

Interactions: The attractive interactions between CTD-CTD and the core, as well as between

TMD proteins, are modeled using the Lennard-Jones potential,

$$E_{lj} = \sum_{<i,j>} 4\epsilon_{ij}\left(\left(\frac{\sigma}{r_{ij}}\right)^{12} - \left(\frac{\sigma}{r_{ij}}\right)^{6}\right) \quad \textbf{Eq. 3}$$

where $\sigma = 2^{-1/6}d$ with d the distance between any two particles in contact including, CTD-core

and TMD-TMD. ϵ_{ij} and r_{ij} denote the interaction strength and the distance between particles i

and j, respectively. The interactions are subject to a cutoff distance $r_{cut\_1} = 0.5a + d$. In

addition, the excluded volume interaction between all particles is presented with a shifted LJ

repulsion potential

$$E_{lj}^{rep} = \sum_{<i,j>} 4\epsilon\_2\left(\left(\frac{\sigma}{r_{ij}}\right)^{12} - \left(\frac{\sigma}{r_{ij}}\right)^{6}\right) + \epsilon\_2 \quad \textbf{Eq. 4}$$

with $\epsilon\_2 = 1$, $r_{cut\_2} = d$ and $\sigma = 2^{-1/6}d$.

Generation of cDNA constructs for mutant viruses.

The plasmid pACNR-CHIKV is a complete sequence clone of the genome of the Chikungunya

181/25 virus (WT) in a CMV-driven plasmid[41]. It allows for the generation of viral progeny by

transfecting HEK-293T cells and serves as a template for all mutations. This plasmid contains

the mKate2 reporter gene between CP and E3, which is released from the structural polyprotein.

The reporter gene is translated from the same subgenomic RNA as the structural proteins; hence,

there is a 1:1:1:1:1 ratio of CP:mKate2:E3:E2:E1 proteins in the cytoplasm. This protein allows

for the relative infectivity of the mutant viruses to be determined by fluorescence microscopy

compared to the WT virus.

The intradimer surface mutant (IDS) carries two mutations in E2 (Y400K/L402R), and

the intertrimer surface mutant (ITS) has N263 of E2 replaced by Q. The mutations were

introduced by site-directed mutagenesis using the protocol by Liu & Naismith[52]. All PCRs were

performed using HS Phusion II (ThermoFisher, Inc.) (see Supplementary Table 3). The PCR

products were digested with DpnI (ThermoFisher, Inc.) and analyzed by agarose/TBE gel

electrophoresis. *E. coli* DH5a bacteria (New England Biolabs) were transformed with the DpnI-

digested PCR products. The selected colonies were cultured in LB (Luria-Bertani) medium with

ampicillin, following standard protocols. The clones were purified with Promega's MiniPrep

Wizard kit, and the sequence identity was confirmed by Sanger sequencing at

LANBAMA/IPICyT.

Virus rescue through transfection.

The production of infectious and quantification of viral particles is similar to that done with other

viruses with reporter genes [53, 54]. HEK-293T cells (ATTC) were grown in DMEM medium

(Corning, Inc.) with glucose (1 g/L) and sodium pyruvate (110mg/L) with 10% fetal bovine

serum (Sigma-Aldrich, Inc.) at 37ºC and 5% CO_2. The transfections were done following the

protocol from Colunga-Saucedo et al.[41]. Briefly, transfections were performed in 12-well plates

using Lipofectamine 3000 (Invitrogen, Inc.) according to the vendor's protocol, with the

modification that 0.6 μg of plasmid was used to transfect 100,000 seeded cells. We always

transfected pACNR-CHIKV (positive control) and a mock transfection with lipofectamine,

without any plasmid, on the same plate as the mutants. At 24 hours post-transfection (h.p.t.), the

supernatant was removed, the cells were washed with PBS, and at 48 h.p.t., the supernatant was

collected, filtered using a 0.22 μm syringe filter, and stored at -80ºC in 0.1 mL aliquots. We

performed four independent transfections.

Assessing infection via a reporter assay.

The quantification of the viral particles produced by transfection was directly determined as the

relative infectivity of the mutants to the WT. This was achieved by infecting HEK293T cells in

24-well culture plates in triplicate with 30 μL of supernatant from transfected cells. Every culture

plate used for relative quantification assays contained a positive control (CHIKV 181/25) and the

supernatant of non-transfected cells (negative control), both of which were produced in the same

transfection as the mutants of interest. The infectivity assays were performed only with

supernatants from the same batch of transfected cells, thawed only once. The relative infectivity

was measured by quantifying the total area of cells expressing mKate2 at 24 hours post-infection

(h.p.i.) using an inverted fluorescence microscope LionHeart FX (Biotek, Inc.). 25 images were

taken per well (5x5 symmetrical grid automatically generated by the microscope) with a 4x

objective, using the Texas red filter (excitation 559 nm +/- 34 nm and emission 630 +/- 69 nm) to

visualize the cells expressing the reporter gene. The relative infectivity (R.I.) was determined by

measuring the area of fluorescent cells per well, averaging across wells, and setting the total

average area of fluorescent cells infected with the WT as 100%. Four independent experiments

(n=4) were performed, each in triplicate for each infection scenario.

Sample preparation for thin-section EM.

For thin-section Transmission Electron Microscopy (TEM), HEK293T cells were seeded in a 6-

well plate, transfected, and fixed with 2% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.2, for

2 hours at room temperature, following a previously published protocol[55]. The resin discs were

cut (70–80 nm thickness) using a diamond blade on a LEICA EM UC7 ultramicrotome, placed

on copper grids, and stained with 0.5% uranyl acetate and 0.5% lead citrate. The samples were

visualized on a JEOL JEM-2100 transmission electron microscope at 200 kV with a 4k Gatan

One View camera. The images were analyzed using ImageJ® (NIH) and Fiji (NIH) software.

Viral replication kinetics.

The replication kinetics were performed in three independent experiments, each in triplicate, by

infecting HEK293 T cells seeded in 24-well plates (50,000 cells per well) with 15 μL of

supernatant. The normalized area of the fluorescent cells was measured every 12 hours for three

days, taking 25 images per well, using the Texas red filter and a 20x objective. The average

fluorescent area of each experiment was normalized to WT, such that the mean fluorescence cell

area of the WT was a relative infection of 100%.

Thermal stability assays.

To determine the thermal stability of the viral particles, the media collected from the transfection

were diluted in OptiMem, incubated at 37°C, 40°C, 45°C, 50°C, 55°C, and 60°C for 15 minutes,

and then cooled to 37°C. HEK-293T cells were infected in 48-well plates in triplicate. The R.I.

of the WT at a temperature different than 37ºC was normalized to that of the WT at 37ºC, and the

infectivity of E2 N263Q was normalized with that of this mutant at 37ºC. The average R.I. was

plotted as a function of the incubation temperature. The data was fitted with QtiPlot 5.15.2 using

Hill's-like equation for competitive unbinding R.I. $= 100/(1+(T/T_{RI50})^n)$ where R.I., T, T_{RI50}, and

n are the R.I. at any temperature normalized the relative infectivity at 37ºC, the temperature at

which the virions were incubated, the temperature at which the infectivity decreases by 50%, and

a cooperative coefficient, respectively. Each experiment was done in triplicate with four

biological replicates.

Virus purification and characterization by Dynamic Light Scattering and Transmission

Electron Microscopy.

The production of CHIKV WT and ITS virions was performed in Vero E6 cells in two T75

flasks per virus for 3 days. The cells were grown in DMEM 10% FBS and antibiotics until they

reached 100% confluency (8 million cells per flask) before infection. Then, the media was

replaced with OptiMem, and the cells were infected with 1 ml of virus for each condition. The

medium was replaced after 2 hours, and the media were collected, filtered through a 0.22 μm

syringe filter, and stored at 4ºC every 24 hours for 3 days. The viral particles were purified with

a low-speed centrifugation protocol[56]. The following day, the supernatant was recovered, and the

pellet was resuspended with 200 μl of PBS.

677   The hydrodynamic diameter of virions was measured by Dynamic Light Scattering

678   (DLS) using a ZETA SIZE PRO Zetameter (Malvern, Inc). Each measurement was done in

679   triplicate with two biological replicas.

680   The purified samples were loaded onto 100-mesh copper grids coated with Formvar (Ted

681   Pella, Inc.) according to standard protocols, stained with uranyl acetate, and observed in a TEM

682   (JEM-2100, JEOL) at 200 kV.

Statistical analysis

Data analysis was performed using the Statistica 10® program. The Kolmogorov-Smirnov test was used to determine the normality of the data, and the statistical differences in the infectivity data were analyzed with the Shapiro-Wilk test. To determine whether significant differences exist ($p \leq 0.05$), the nonparametric Kruskal-Wallis test for multiple comparisons, followed by Dunn's post hoc test, was used.

ACKNOWLEDGMENT

M.A.C.I. (CVU: 1232434) acknowledges the support of SECIHTI through the Ph.D. fellowship 4044561 M.C-G. thanks SECIHTI for the grant CBF2023-2024-1125 and COPOCYT for the grant 2024-03-M07. RZ acknowledges support from NSF DMR-2131963 and MCB/PHY-2413062.

ABBREVIATIONS

CHIKV, Chikungunya virus

CMV, Cytomegalovirus

CP, Capsid protein

cryo-EM, Cryo-electron microscopy

cryo-ET, Cryo-electron tomography

CT, Cytoplasmic tail mutant

CTD, C-terminal or cytosolic domain

DLS, dynamic light scattering

ED, Ectodomain

h.p.i., Hours post-infection

IDS, Intradimer surfaced mutant

ITS, Intertrimer surface mutant

LJ, Lennard-Jones

MC, Monte Carlo

MD, Molecular dynamics

NC, Nucleocapsid or core

TMD, Transmembrane domain

TEM, Transmission Electron Microscopy

WT, Wild-type

REFERENCES

1.   Ruiz-Herrero, T.; Velasco, E.; Hagan, M. F., Mechanisms of budding of
nanoscale particles through lipid bilayers. *J Phys Chem B* **2012,** *116* (32), 9595-603.
2.   Chmielewski, D.; Schmid, M. F.; Simmons, G.; Jin, J.; Chiu, W., Chikungunya
virus assembly and budding visualized in situ using cryogenic electron tomography. *Nat*
Microbiol **2022,** *7* (8), 1270-1279.
3.   Byrd, E. A.; Kielian, M., The Alphavirus E2 Membrane-Proximal Domain Impacts
Capsid Interaction and Glycoprotein Lattice Formation. *J Virol* **2019,** *93* (4).
4.   Byrd, E. A.; Kielian, M., An Alphavirus E2 Membrane-Proximal Domain Promotes
Envelope Protein Lateral Interactions and Virus Budding. *mBio* **2017,** *8* (6).
5.   Jose, J.; Przybyla, L.; Edwards, T. J.; Perera, R.; Burgner, J. W., 2nd; Kuhn, R.
J., Interactions of the cytoplasmic domain of Sindbis virus E2 with nucleocapsid cores
promote alphavirus budding. *J Virol* **2012,** *86* (5), 2585-99.
6.   Hartlieb, B.; Weissenhorn, W., Filovirus assembly and budding. *Virology* **2006,**
344 (1), 64-70.
7.   Elmasri, Z.; Nasal, B. L.; Jose, J., Alphavirus-Induced Membrane
Rearrangements during Replication, Assembly, and Budding. *Pathogens* **2021,** *10* (8).
8.   Freed, E. O., HIV-1 assembly, release and maturation. *Nature Reviews*
Microbiology **2015,** *13* (8), 484-96.

9.   Lingappa, J. R.; Reed, J. C.; Tanaka, M.; Chutiraka, K.; Robinson, B. A., How HIV-1 Gag assembles in cells: Putting together pieces of the puzzle. *Virus Res* **2014,** *193*, 89-107.

10.   Sundquist, W. I.; Krausslich, H. G., HIV-1 assembly, budding, and maturation. *Cold Spring Harb Perspect Med* **2012,** *2* (7), a006924.

11.   Su, Z.; Wu, C.; Shi, L.; Luthra, P.; Pintilie, G. D.; Johnson, B.; Porter, J. R.; Ge, P.; Chen, M.; Liu, G.; Frederick, T. E.; Binning, J. M.; Bowman, G. R.; Zhou, Z. H.; Basler, C. F.; Gross, M. L.; Leung, D. W.; Chiu, W.; Amarasinghe, G. K., Electron Cryo-microscopy Structure of Ebola Virus Nucleoprotein Reveals a Mechanism for Nucleocapsid-like Assembly. *Cell* **2018,** *172* (5), 966-978 e12.

12.   Wan, W.; Kolesnikova, L.; Clarke, M.; Koehler, A.; Noda, T.; Becker, S.; Briggs, J. A. G., Structure and assembly of the Ebola virus nucleocapsid. *Nature* **2017,** *551* (7680), 394-397.

13.   Dong, S.; Yang, P.; Li, G.; Liu, B.; Wang, W.; Liu, X.; Xia, B.; Yang, C.; Lou, Z.; Guo, Y.; Rao, Z., Insight into the Ebola virus nucleocapsid assembly mechanism: crystal structure of Ebola virus nucleoprotein core domain at 1.8 A resolution. *Protein Cell* **2015,** *6* (5), 351-62.

14.   Button, J. M.; Qazi, S. A.; Wang, J. C.; Mukhopadhyay, S., Revisiting an old friend: new findings in alphavirus structure and assembly. *Curr Opin Virol* **2020,** *45*, 25-33.

15.   Garoff, H.; Simons, K., Location of the spike glycoproteins in the Semliki Forest virus membrane. *PNAS* **1974,** *71* (10), 3988-92.

16.   Forsell, K.; Xing, L.; Kozlovska, T.; Cheng, R. H.; Garoff, H., Membrane proteins organize a symmetrical virus. *EMBO J* **2000,** *19* (19), 5081-91.

17.   Lee, S.; Owen, K. E.; Choi, H. K.; Lee, H.; Lu, G.; Wengler, G.; Brown, D. T.; Rossmann, M. G.; Kuhn, R. J., Identification of a protein binding site on the surface of the alphavirus nucleocapsid and its implication in virus assembly. *Structure* **1996,** *4* (5), 531-41.

18.   Chmielewski, D.; Su, G. C.; Kaelber, J. T.; Pintilie, G. D.; Chen, M.; Jin, J.; Auguste, A. J.; Chiu, W., Cryogenic electron microscopy and tomography reveal imperfect icosahedral symmetry in alphaviruses. *PNAS Nexus* **2024,** *3* (3), pgae102.

19.   Weaver, S. C.; Winegar, R.; Manger, I. D.; Forrester, N. L., Alphaviruses: population genetics and determinants of emergence. *Antiviral Res* **2012,** *94* (3), 242-57.

20.   Robinson, M. C., An epidemic of a dengue-like fever in the southern province of Tanganyika. *Cent Afr J Med* **1956,** *2* (11), 394-6.

21.   Robinson, M. C., An epidemic of virus disease in Southern Province, Tanganyika Territory, in 1952-53. I. Clinical features. *Trans R Soc Trop Med Hyg* **1955,** *49* (1), 28-32.

22.   Frolov, I.; Frolova, E. I., Molecular Virology of Chikungunya Virus. *Curr Top Microbiol Immunol* **2022,** *435*, 1-31.

23.   Erasmus, J. H.; Auguste, A. J.; Kaelber, J. T.; Luo, H.; Rossi, S. L.; Fenton, K.; Leal, G.; Kim, D. Y.; Chiu, W.; Wang, T.; Frolov, I.; Nasar, F.; Weaver, S. C., A chikungunya fever vaccine utilizing an insect-specific virus platform. *Nat Med* **2017,** *23* (2), 192-199.

24.   Ganesan, V. K.; Duan, B.; Reid, S. P., Chikungunya Virus: Pathophysiology, Mechanism, and Modeling. *Viruses* **2017,** *9* (12).

25.   Weaver, S. C.; Lecuit, M., Chikungunya virus and the global spread of a
mosquito-borne disease. *N Engl J Med* **2015,** *372* (13), 1231-9.
26.   Owen, K. E.; Kuhn, R. J., Alphavirus budding is dependent on the interaction
between the nucleocapsid and hydrophobic amino acids on the cytoplasmic domain of
the E2 envelope glycoprotein. *Virology* **1997,** *230* (2), 187-96.
27.   Ren, S. C.; Qazi, S. A.; Towell, B.; Wang, J. C.; Mukhopadhyay, S., Mutations
at the Alphavirus E1'-E2 Interdimer Interface Have Host-Specific Phenotypes. *J Virol*
2022, *96* (5), e0214921.
28.   Suomalainen, M.; Liljestrom, P.; Garoff, H., Spike protein-nucleocapsid
interactions drive the budding of alphaviruses. *J Virol* **1992,** *66* (8), 4737-47.
29.   Lopez, S.; Yao, J. S.; Kuhn, R. J.; Strauss, E. G.; Strauss, J. H., Nucleocapsid-
glycoprotein interactions required for assembly of alphaviruses. *J Virol* **1994,** *68* (3),
1316-23.
30.   Lazaro, G. R.; Mukhopadhyay, S.; Hagan, M. F., Why Enveloped Viruses Need
Cores-The Contribution of a Nucleocapsid Core to Viral Budding. *Biophys J* **2018,** *114*
(3), 619-630.
31.   Ribeiro-Filho, H. V.; Coimbra, L. D.; Cassago, A.; Rocha, R. P. F.; Guerra, J.;
de Felicio, R.; Carnieli, C. M.; Leme, L.; Padilha, A. C.; Paes Leme, A. F.; Trivella, D.
B. B.; Portugal, R. V.; Lopes-de-Oliveira, P. S.; Marques, R. E., Cryo-EM structure of
the mature and infective Mayaro virus at 4.4 A resolution reveals features of
arthritogenic alphaviruses. *Nature communications* **2021,** *12* (1), 3038.
32.   Skoging, U.; Vihinen, M.; Nilsson, L.; Liljestrom, P., Aromatic interactions define
the binding of the alphavirus spike to its nucleocapsid. *Structure* **1996,** *4* (5), 519-29.
33.   Zhao, H.; Lindqvist, B.; Garoff, H.; von Bonsdorff, C. H.; Liljestrom, P., A
tyrosine-based motif in the cytoplasmic domain of the alphavirus envelope protein is
essential for budding. *EMBO J* **1994,** *13* (18), 4204-11.
34.   Forsell, K.; Griffiths, G.; Garoff, H., Preformed cytoplasmic nucleocapsids are not
necessary for alphavirus budding. *EMBO J* **1996,** *15* (23), 6495-505.
35.   Brown, R. S.; Wan, J. J.; Kielian, M., The Alphavirus Exit Pathway: What We
Know and What We Wish We Knew. *Viruses* **2018,** *10* (2).
36.   Mendes, A.; Kuhn, R. J., Alphavirus Nucleocapsid Packaging and Assembly.
Viruses **2018,** *10* (3).
37.   Kaelber, J. T.; Chmielewski, D.; Chiu, W.; Auguste, A. J., Alphavirus Particles
Can Assemble with an Alternate Triangulation Number. *Viruses* **2022,** *14* (12).
38.   Cheng, F.; Tsvetkova, I. B.; Khuong, Y. L.; Moore, A. W.; Arnold, R. J.;
Goicochea, N. L.; Dragnea, B.; Mukhopadhyay, S., The packaging of different cargo
into enveloped viral nanoparticles. *Mol Pharm* **2013,** *10* (1), 51-8.
39.   Cheng, F.; Mukhopadhyay, S., Generating enveloped virus-like particles with in
vitro assembled cores. *Virology* **2011,** *413* (2), 153-60.
40.   Mukhopadhyay, S.; Chipman, P. R.; Hong, E. M.; Kuhn, R. J.; Rossmann, M.
G., In vitro-assembled alphavirus core-like particles maintain a structure similar to that
of nucleocapsid cores in mature virus. *J Virol* **2002,** *76* (21), 11128-32.
41.   Colunga-Saucedo, M.; Rubio-Hernandez, E. I.; Coronado-Ipina, M. A.;
Rosales-Mendoza, S.; Castillo, C. G.; Comas-Garcia, M., Construction of a
Chikungunya Virus, Replicon, and Helper Plasmids for Transfection of Mammalian
Cells. *Viruses* **2022,** *15* (1).

42.   Ruiz-Guillen, M.; Gabev, E.; Quetglas, J. I.; Casales, E.; Ballesteros-Briones,
M. C.; Poutou, J.; Aranda, A.; Martisova, E.; Bezunartea, J.; Ondiviela, M.; Prieto, J.;
Hernandez-Alcoceba, R.; Abrescia, N. G.; Smerdou, C., Capsid-deficient alphaviruses
generate propagative infectious microvesicles at the plasma membrane. *Cell Mol Life*
Sci **2016,** *73* (20), 3897-916.
43.   Zhang, Y. N.; Deng, C. L.; Li, J. Q.; Li, N.; Zhang, Q. Y.; Ye, H. Q.; Yuan, Z.
M.; Zhang, B., Infectious Chikungunya Virus (CHIKV) with a Complete Capsid Deletion:
a New Approach for a CHIKV Vaccine. *J Virol* **2019,** *93* (15).
44.   Li, S.; Tresset, G.; Zandi, R., From disorder to icosahedral symmetry: How
conformation-switching subunits enable RNA virus assembly. *Sci Adv* **2025,** *11* (39),
eady7241.
45.   Timmermans, S.; Ramezani, A.; Montalvo, T.; Nguyen, M.; van der Schoot, P.;
van Hest, J. C. M.; Zandi, R., The Dynamics of Viruslike Capsid Assembly and
Disassembly. *J Am Chem Soc* **2022,** *144* (28), 12608-12612.
46.   Li, S.; Erdemci-Tandogan, G.; van der Schoot, P.; Zandi, R., The effect of RNA
stiffness on the self-assembly of virus particles. *J Phys Condens Matter* **2018,** *30* (4),
044002.
47.   Cheng, R. H.; Kuhn, R. J.; Olson, N. H.; Rossmann, M. G.; Choi, H. K.; Smith,
T. J.; Baker, T. S., Nucleocapsid and glycoprotein organization in an enveloped virus.
Cell **1995,** *80* (4), 621-30.
48.   Zhang, Y.; Anbir, S.; McTiernan, J.; Li, S.; Worcester, M.; Mishra, P.; Colvin,
M. E.; Gopinathan, A.; Mohideen, U.; Zandi, R.; Kuhlman, T. E., Synthesis, insertion,
and characterization of SARS-CoV-2 membrane protein within lipid bilayers. *Sci Adv*
2024, *10* (9), eadm7030.
49.   Panahandeh, S.; Li, S.; Dragnea, B.; Zandi, R., Virus Assembly Pathways
Inside a Host Cell. *ACS Nano* **2022,** *16* (1), 317-327.
50.   Panahandeh, S.; Li, S.; Zandi, R., The equilibrium structure of self-assembled
protein nano-cages. *Nanoscale* **2018,** *10* (48), 22802-22809.
51.   Zandi, R.; Dragnea, B.; Travesset, A.; Podgornik, R., On virus growth and form.
Physics Reports **2020,** *847*, 1-102.
52.   Liu, H.; Naismith, J. H., An efficient one-step site-directed deletion, insertion,
single and multiple-site plasmid mutagenesis protocol. *BMC Biotechnol* **2008,** *8*, 91.
53.   Pierson, T. C.; Diamond, M. S.; Ahmed, A. A.; Valentine, L. E.; Davis, C. W.;
Samuel, M. A.; Hanna, S. L.; Puffer, B. A.; Doms, R. W., An infectious West Nile virus
that expresses a GFP reporter gene. *Virology* **2005,** *334* (1), 28-40.
54.   Gadea, G.; Bos, S.; Krejbich-Trotot, P.; Clain, E.; Viranaicken, W.; El-
Kalamouni, C.; Mavingui, P.; Despres, P., A robust method for the rapid generation of
recombinant Zika virus expressing the GFP reporter gene. *Virology* **2016,** *497*, 157-162.
55.   Lara-Hernandez, I.; Munoz-Escalante, J. C.; Bernal-Silva, S.; Noyola, D. E.;
Wong-Chew, R. M.; Comas-Garcia, A.; Comas-Garcia, M., Ultrastructural and
Functional Characterization of Mitochondrial Dynamics Induced by Human Respiratory
Syncytial Virus Infection in HEp-2 Cells. *Viruses* **2023,** *15* (7).
56.   Rayaprolu, V.; Ramsey, J.; Wang, J. C.; Mukhopadhyay, S., Alphavirus
Purification Using Low-speed Spin Centrifugation. *Bio Protoc* **2018,** *8* (6), e2772.