Regenerative Grazing as a Climate Change Mitigation Strategy: A
Systematic Review

Authors:

Docker B. Clark, New York University

Shelby C. McClelland, New York University, Stony Brook University

Jasmine A. Dillon, Cornell University

Matthew N. Hayek*, New York University

*Corresponding Author - Matthew N. Hayek, (matthew.hayek@nyu.edu,

ORCID iD: https://orcid.org/0000-0001-9792-4362)

Abstract

Ruminant livestock production is a meaningful contributor to global greenhouse gas (GHG) emissions. Regenerative grazing has been proposed as a climate change mitigation strategy for offsetting livestock GHG emissions by potentially causing additional soil organic carbon (SOC) sequestration. We review the current state of the literature and collect reported and inferred SOC sequestration from regenerative grazing experiments, ranking their study designs by ascending evidentiary strength: observational, cross-sectional and longitudinal. Across 28 studies that compare regenerative to conventional grazing, the reported and inferred SOC sequestration rates in regenerative grazing treatments varied widely, with the highest values reported in lower-strength observational studies. Among higher-strength cross-sectional studies and longitudinal studies, median additional SOC sequestration rates in regenerative grazing treatments were not significantly different than zero, indicating no significant enhancement of SOC across all regenerative grazing interventions. Currently, higher-strength evidence does not support widespread claims of SOC sequestration from regenerative grazing. To clarify its potential for climate mitigation, a greater number of longitudinal studies are needed.

Keywords: regenerative agriculture, livestock, climate mitigation, SOC, grazing, carbon sequestration

1.   Introduction

Ruminant meat and dairy production account for 9% to 13% of global anthropogenic greenhouse gas (GHG) emissions (FAO, 2022; Halpern et al., 2022). One notable GHG mitigation strategy has sparked popular imagination: "regenerative grazing", which may result in soil organic carbon (SOC) sequestration. Regenerative grazing lacks an agreed-upon definition, but encompasses livestock management practices aimed at improving soil and ecosystem health, farm profit, and resiliency (Dillon and Machmuller, 2021). These practices often involve rotating herds across smaller subdivided pastures at high stocking densities to mimic natural grazing. Highly-visible public presentations have asserted that regenerative grazing meaningfully offsets cattle emissions, or even offset emissions from other economic sectors (Leu 2024).

Recent scientific assessments reach diverging conclusions about SOC sequestration from regenerative grazing (Eshel et al., 2025; Pelton et al., 2024). However, these studies derived estimates from only a few studies. Additionally, these reviews have not analyzed experimental designs, i.e. whether they were potentially prone to confounding variables.

Our systematic review assembles and evaluates evidence addressing the question: does regenerative grazing induce SOC sequestration? Specifically, what is the quantity and quality of evidence that SOC sequestration occurred and was additional to counterfactual conditions, e.g. under conventional grazing? We assemble SOC sequestration outcomes from all regeneratively-managed ruminant livestock grazing experiments to date, and evaluate SOC results based on a hierarchy of evidence based on their experimental designs.

2.   Methods

2.1. Grazing management definitions

To facilitate comparison between regenerative and conventional grazing, we define "conventional grazing" as methods that allow animals to access pastures for the entirety of the grazing season without providing rest. This is also commonly referred to as "continuous grazing," "traditional grazing," and "set-stocking". We define "regenerative grazing" as a suite

of methods that subdivide larger pastures into "paddocks" that are alternatively intensively

grazed by the entire herd or subdivisions of it, then rested intermittently with periods of no

grazing. This rotational approach is a frequent but not universal tenet of regenerative agriculture,

a term that lacks a consensus definition (Giller et al., 2021). Many terms in scientific and

agricultural parlance meet our functional definition, including: adaptive multi-paddock grazing,

mob grazing, management intensive grazing, and holistic planned management.

2.2. Literature review

We conducted a systematic review of the literature following PRISMA guidelines (Fig.

S1), for results quantifying SOC outcomes of regenerative grazing. To compile these articles, we

conducted series of database searches from 2002 extending to the present (accessed: Feb 5,

2025) using Web of Science and Google Scholar. We searched Web Of Science for all studies

containing a wide array of search terms related to regenerative grazing and its synonyms, plus

soil carbon outcomes (Box S1). This query returned 86 studies. We included studies that A)

compare regenerative grazing and/or conventional grazing management; B) report soil organic

matter (SOM%, SOC%, SOC stocks, or SOC flux; C) present original in-situ soil carbon

measurements; and D) report the duration of time pastures were under either grazing treatment.

A total of 23 of the 86 Web Of Science retrieved studies met our initial criteria.

Additional Google Scholar searches returned 1 unique study. Searching references from

literature reviews appearing in the initial search resulted in 4 more studies that met our criteria.

Lastly, references cited by the regenerative grazing SOC analyses that met our search criteria

were carefully examined for inclusion, but none met all of our criteria. In total, we found 28

unique studies meeting our inclusion criteria. Several studies (N = 11) reported data using a

management type falling outside of our definition of conventional grazing as a counterfactual for

regenerative grazing management. These data are accounted for in our auxiliary analysis but

were not suitable for our main analysis. In addition to SOC, we catalogued the studies' location,

species, animal stocking densities and rates, soil sampling depth(s), and land use history.

2.3. Categorization of study designs

We categorized the 28 studies that met our criteria into four experimental design

categories according to a hierarchy of evidentiary strength: (1) Time-Series Observational

studies, which reported SOC from regeneratively grazed sites, using either true time series or

chronosequence measurements, but without an observed analogous conventionally grazed site;

(2) Comparative Observational studies, that compared adjacent sites that differed in one or more

biotic or abiotic confounding variables (e.g., soil texture, climate, land use history) and reported

only a single SOC measurement; (3) Matched Quasi-Experimental (MQE) Cross-Sectional

studies, which directly compare adjacent regenerative and conventional sites that are otherwise

similar with respect to all reported biotic and abiotic properties the authors deemed likely to

affect SOC, but only report SOC at a single time point, and; (4) MQE Longitudinal studies

reporting at least two SOC measurements over a period of ≥1 year. MQE designs were ranked

higher than comparative observational studies because they attempt to control for confounding

variables including land use history and different starting baseline SOC quantities. No true

experimental studies, which would randomize grazing treatments among sites, were found in our

literature search.

2.4. SOC stocks

We converted all studies' SOC stock data into common units of metric tons of

carbon per hectare (MgC ha^{-1}) which some studies report directly (N = 16). In cases, soil carbon

concentration units were reported (%SOC or SOC g/kg) and we used the following equation to

convert to C stock:

105
$$Cstock\ (MgC/ha)\ =\ \sum_{i}^{n} SOC_i\left(\frac{g}{100g}\right) \times BD_i\left(\frac{g}{cm^3}\right) \times D_i(cm) \times 100$$

where BD = bulk density of a soil sample at depth i, D = the depth in centimeters of each

stratum of soil, SOC = soil organic carbon concentration at depth i, and 100 is the conversion

factor of gC cm^{-2} to MgC ha^{-1}. When soil C concentration was reported in %SOM, we first

converted SOM to SOC using a 0.58 conversion factor (McClelland et al., 2021; Lunt, 1931;

Broadbent, 1953; Howard, 1965). In cases where we were limited by a lack of BD measurements

by depth, we applied a single value to all depths: (1) Where only one BD was reported (N = 1),

we applied this to all depths; (2) Where none were reported (N = 3), an estimated value from

literature (USDA, 2019) was applied to all depth strata based on soil textures reported within

their respective studies.

2.5. SOC fluxes

Annual fluxes from these studies (MgC ha^{-1} y^{-1}) were either reported directly, or we

calculated them according to the common method within other studies in their evidentiary

category. All Time-Series Observational studies reported their own SOC flux rates which we

incorporated directly. However, for the comparative observational studies that did not quantify

flux, we took the difference in total reported SOC stock between treatment and control sites

divided by the duration of the treatment.

For cross-sectional studies, differences in C stock between treatment and control sites

were divided by the number of years since the site was given contrasting grazing treatments. If

treatment durations differed, we calculated the mean of the durations. Some studies did not

explicitly organize adjacent regenerative and conventional pasture sites into pairs. We paired

these sites according to similar animal stocking densities/rates. If no stocking information was

available, SOC was averaged across treatment sites and compared to the average across control

sites.

For MQE Longitudinal studies, fluxes were calculated by taking a difference of

differences: the difference in SOC stock from the first (baseline) and last (most recent) sampling

year, then the difference between paired regenerative and conventional sites, dividing the result

by number of years since first and last measurements.

When fluxes were reported directly we collected them from their parent study regardless

of how the authors calculation methods differed from those we outline above.

2.6. Statistical analysis

All statistical analysis was conducted using R version 4.3.2 (R Core Team, 2023).

Data manipulation and visualizations were facilitated by use of the "tidyverse", "_matrixStats",

and "_ggrepel" packages (Wickham et al. 2019; Bengtsson, 2025; Slowikowski 2024). Means

and medians for each category were calculated with 95% confidence intervals using

bootstrapping by randomly resampling SOC fluxes over N = 1001 synthetic vectors with

replacement (Efron, 1979).

3. Results

We compiled 28 regenerative grazing studies in which SOC fluxes were either reported or could be inferred using methods common in the literature. Combined, these studies represent 49 pasture sites (one pair of co-located continuous vs. regenerative case-control pastures = one pasture site). In most studies, pastures were grazed by cattle (N = 16), while others were grazed by sheep (N = 5), or had mixed cattle with sheep and/or goats (N = 5). The Americas and Australia were disproportionately represented: 93% of studies, i.e. all but two. In descending order: 46% of studies or 57% of sites were in North America; 32% of studies or 27% of sites were in Australia; 14% of studies and 10% of sites were in South America. Two remaining studies, with a combined three sites, concerned sheep grazing in China and Spain.

Time-series observational studies show the greatest maximum values for annual SOC sequestration. Median and mean SOC sequestration rates were 3.59 (2.29 - 7.10 95% CI) MgC ha^{-1} y^{-1} and 4.33 (2.29 – 7.10) MgC ha^{-1} y^{-1}, respectively, both of which were significantly greater than zero (Table 1). Comparative observational studies' median SOC sequestration was 0.33 MgC ha^{-1} y^{-1} and was not significantly greater than zero; the mean was significantly greater than zero at 2.42 (0.02 − 4.93) MgC ha^{-1} y^{-1}, but overlaps zero when studies with insufficient soil depth data are excluded (-0.43 − 6.61) MgC ha^{-1} y^{-1}.

The largest category by study count was MQE Cross-Sectional (N = 16). Median SOC sequestration was 0.09 MgC ha^{-1} y^{-1} and was not significantly greater than zero (-0.05 – 0.42). This lack of statistical significance remained when SOC data from studies with sparse depth measurements were excluded (Table 1). In both cases, however, mean SOC sequestration in the MQE cross-sectional study category was significantly greater than zero (Table 1) due to a non-symmetrical distribution of SOC outcomes in this category.

MQE longitudinal studies (N = 5) showed the smallest range of SOC fluxes, and median sequestration of -0.03 MgC ha^{-1} y^{-1}, which was not significantly less than zero (-0.03 − 0.89), nor was the mean (Table 1). Three out of the five longitudinal studies measured surface soil only (10 cm).

The same tendencies in medians and means across study categories were observed when we excluded studies with sparse soil depth data (Table 1), when we treated sites as replicates

rather than studies (Fig S2), or when we normalized SOC numbers to a common depth (Figs, S3
& S4).

To evaluate farmed livestock species as a potentially confounding variable (Fig S5), we
calculated summary statistics for our dataset separating categories by grazed species (Table S1).
We evaluated the significance of SOC sequestration on cattle-grazed vs all other grazed ruminant
categories (Mixed Large and Small + Small Ruminants) using a Mann-Whitney U, which
returned a p-value of 0.0879, i.e. not significant at the 95% confidence level. We performed this
test on only studies within the MQE cross-sectional category as this provided the largest and
most even sample size of cattle-grazing (N = 9) and non-cattle ruminant grazing (N = 6) studies.

The 2019 case-controlled study conducted by Alemu et al. (2019) collected SOC
measurements over time, and provides sufficient data to calculate results in both a cross-sectional
and longitudinal manner (Fig 1). We incorporated both, to highlight the difference in apparent
SOC sequestration rates from MQE Cross-Sectional versus MQE Longitudinal study designs.

Fig. 1: Mean SOC sequestration rates in each regenerative grazing study (points and numbers) and
ranges (boxes). Point colors represent the maximum depth at which SOC measurements were

conducted. Studies with depth measurements of <20 cm or with only one depth stratum were classified as

having insufficient depth data (white points). Light gray boxes represent full ranges of data and black line

segments are medians for each category, excluding studies with insufficient depth data. Dark gray box

represents interquartile range for MQE Cross sectional category.

		All studies				Studies with sparse depth data excluded		
		SOC sequestration				SOC sequestration		
Study design	Studies	Range	Median (95% CI)	Mean (95% CI)	Studies	Range	Median (95% CI)	Mean (95% CI)
Time-Series Observational	3	2.29 – 7.1	3.59 (2.29 – 7.1)	4.33 (2.29 – 7.1)	3	2.29 – 7.1	3.59 (2.29 – 7.1)	4.33 (2.29 – 7.1)
Comparative Observational	5	-0.43 – 6.61	0.33 (-0.43 – 6.61)	2.42 (0.02 – 4.93)	3	-0.43 – 6.61	5.25 (-0.43 – 6.61)	3.81 (-0.43 – 6.61)
MQE Cross-Sectional	16	-0.58 – 3.37	0.09 (-0.05 – 0.42)	0.5 (0.06 – 1.05)	13	-0.58 – 3.37	0.12 (-0.05 – 0.49)	0.62 (0.06 – 1.24)
MQE Longitudinal	5	-0.3 – 0.89	-0.03 (-0.3 – 0.89)	0.13 (-0.18 – 0.52)	2	-0.3 – -0.13	-0.21 (-0.3 – -0.13)	-0.21 (-0.3 – -0.13)

Table 1: Summary statistics within each study design category of SOC sequestration in MgC ha^{-1} y^{-1} with

bootstrapped 95% confidence intervals (positive numbers = sequestration). Columns on the right were

calculated from data excluding studies with depth measurements that were deemed insufficient.

4.   Discussion

4.1. Carbon sequestration outcomes from regenerative grazing practices

Experimental designs produced different diverging conclusions about regenerative

management's influence on SOC. The highest-strength studies (MQE longitudinal) did not

demonstrate consistent evidence for regenerative grazing enhancing SOC sequestration. This

may be due to their small number (N = 5). Only two MQE longitudinal studies had suitable

depth measurements, neither of which found an additive effect of regenerative grazing upon SOC

over time. For the remaining studies, extrapolating their surface SOC measurements to the entire

soil profile would be specious because SOC can move throughout the soil column without

resulting in net SOC loss or gain (Hao et al. 2017).

Among MQE Cross-Sectional studies we found a positive *mean* SOC enhancement in

regenerative sites, which could suggest that regenerative grazing enhances SOC sequestration.

However, there are two important caveats to this finding. First, *median* SOC sequestration was

not significantly different from zero, meaning studies were approximately equally likely to find

SOC results in either direction. Second, without repeated measurements (i.e., longitudinal

studies) it is not possible to attribute SOC responses to regenerative grazing per se; measured

differences in SOC could be an artifact of different baselines from the start of the regenerative

grazing experiment.

Despite notable limitations in the evidentiary strength of multiple studies' design, studies

commonly included language that suggested SOC sequestration and/or enhancement was caused

by regenerative grazing, e.g. "AMP grazing can contribute to climate change mitigation through

SOC sequestration" (Stanley et al. 2018) or "...with significant [SOC] accumulation in the

improved grassland area" (Costa et al. 2022), across N = 12 studies. These claims overstate the

conclusions warranted by the experimental design, because measured differences may be due to

coincidental differences in baseline SOC, land management history, or other confounding

environmental variables that the study designs do not control for.

In all studies, differences in SOC could also arise from interventional or observational

biases. No studies included in our analysis used a randomized design. This introduces a potential

site selection bias if ranchers chose more productive pastures to try novel or risky management

methods, or the opposite.

4.2 Future recommendations

To substantiate claims about regenerative grazing enhancing or sequestering SOC, more

longitudinal studies are needed. Improvements to study design are readily feasible. Among MQE

Cross-Sectional study sites, additional work could be done to turn some into Longitudinal

studies; if management methods at these sites are presently the same as they were during

reported baseline measurements, a single additional set of soil measurements could be sufficient.

Ideally, multiple more years of measurement should occur at every site to demonstrate that any

net sequestration is not due to an anomalous year of climatic or ecological conditions.

Furthermore, additional studies should begin with randomized assignment to control for site

selection bias.

Observational studies are especially low-strength evidence that regenerative grazing

results in SOC sequestration because land use history is a confounding variable. While all three

Time-Series Observational studies reported positive SOC sequestration, their sites all had

histories of annual cropping (Machmuller et al. 2015; Stanley et al. 2018; Rowntree et al. 2020).

Transitioning to *any* system with perennial roots (e.g. conventional pasture, mowed meadow,

orchard, wetland, savanna, or forest) will result in SOC sequestration after tillage and other soil

disturbance are removed (Yang et al. 2020). By contrast, MQE Longitudinal designs control for

this history. Studies in this category show that additional SOC sequestration from regenerative

grazing is closer to and overlaps zero, with the only two studies that took sufficient depth

measurements (multiple and greater then 10 cm) showing SOC loss over time (Fig. 1; Table 1).

Despite this, observational studies used causal language suggesting that regenerative grazing

itself induced SOC sequestration.

We use conventional continuous grazing as a counterfactual for regenerative grazing

because it represents a directly comparable land use with the same fundamental goal of livestock

production. Other non-grazing land management practices—such as grazing exclusion, hay

harvesting, or wild grazing—were present in the literature (N = 11). These counterfactuals were

included in our auxiliary analysis (Figs. S6 & S7) because they are qualitatively different from a

production standpoint. Relaxed rotational grazing, defined by a lower frequency and/or stocking

rate than typical regenerative practices, was also deemed an inappropriate counterfactual for our

main analysis and was placed in the auxiliary analysis as it does not represent a distinct

alternative land use. When comparing regenerative grazing to these non-continuous grazing

counterfactuals, results were mixed. Additional SOC sequestration rates from regenerative

grazing still overlapped zero (Figs S6 & S7).

We found no significant difference in soil organic carbon (SOC) between sites grazed by

cattle and those grazed by other ruminants ($p = 0.09$). This indicates that livestock species may

not be a major confounding factor in our assessment. Although studies on cattle grazing

appeared to show slightly higher average SOC sequestration from regenerative sites, the limited

sample size for each study design category prevents us from making definitive claims about

species-specific SOC gains (Fig. S5, Table S1). Therefore, we did not find evidence that our

pooled results may be biased by the type of animal used.

5. Conclusion

Our analysis finds little robust support for the hypothesis that regenerative grazing

management can meaningfully sequester carbon in soil organic matter. Relatedly, there is little

evidence that regenerative grazing could potentially offset an appreciable quantity of ruminant

enteric fermentation methane or other greenhouse gas emissions. In order to test this hypothesis

more rigorously, more case-controlled longitudinal studies are needed. At a minimum, past

cross-sectional study sites would need only an additional soil measurement of similar quality, if

previous grazing management methods have continued, to provide higher-quality evidence that

SOC sequestration is occurring, and to attribute that SOC sequestration to regenerative grazing

itself. Ideally, however, multiple more years of measurement should be conducted.

Author Contributions

Docker Clark: data curation (lead); formal analysis (supporting); investigation (equal);

visualization (equal); writing - original draft preparation (equal). **Jasmine A. Dillon**:

methodology (supporting); visualization (supporting); writing - review and editing (equal).

Matthew N. Hayek: conceptualization (lead); data curation (supporting); formal analysis (lead);

funding acquisition (lead); investigation (equal); methodology (lead); project administration

(lead); supervision (lead); visualization (equal); writing - original draft preparation (equal);

writing - review and editing (equal). **Shelby C. McClelland**: formal analysis (supporting);

methodology (supporting); visualization (supporting); writing - review and editing (equal).

Funding Statement

This work was supported in part by a grant from NYU's Center for Environmental and

Animal Protection (CEAP) and a donation from Food System Innovations. The funders had no

influence over research design, methods, or findings.

Conflict of Interest Statement

The authors declare no conflicts of interest.

References

Alemu, A. W., Kröbel, R., McConkey, B. G., & Iwaasa, A. D. (2019). Effect of increasing

species diversity and grazing management on pasture productivity, animal performance, and

soil carbon sequestration of re-established pasture in canadian prairie. Animals, 9(4), 127.

https://doi.org/10.3390/ani9040127

Bartley, R., Abbott, B. N., Ghahramani, A., Ali, A., Kerr, R., Roth, C. H., & Kinsey-Henderson,

A. (2023). Do regenerative grazing management practices improve vegetation and soil

health in grazed rangelands? Preliminary insights from a space-for-time study in the Great

Barrier Reef catchments, Australia. The Rangeland Journal, 44(4), 221–246.

https://doi.org/10.1071/RJ22047

Bengtsson, H. (2025). _matrixStats: Functions that Apply to Rows and Columns of Matrices

(and to Vectors)_. (Version 1.5.0). https://CRAN.R-project.org/package=matrixStats

Broadbent, F. E. (1953). The Soil Organic Fraction. In Advances in Agronomy (Vol. 5, pp. 153–

183). Elsevier. https://doi.org/10.1016/S0065-2113(08)60229-1

Cattle, S. R., & Southorn, N. J. (2010). Macroporosity of pasture topsoils after three years of set-

stocked and rotational grazing by sheep. Soil Research, 48(1), 43.

https://doi.org/10.1071/SR09004

Chan, K. Y., Oates, A., Li, G. D., Conyers, M. K., Prangnell, R. J., Poile, G., Liu, D. L., &

Barchia, I. M. (2010). Soil carbon stocks under different pastures and pasture management

in the higher rainfall areas of south-eastern Australia. Soil Research, 48(1), 7.

https://doi.org/10.1071/SR09092

Chen, W., Huang, D., Liu, N., Zhang, Y., Badgery, W. B., Wang, X., & Shen, Y. (2015).

Improved grazing management may increase soil carbon sequestration in temperate steppe.

Scientific Reports, 5(1), 10892. https://doi.org/10.1038/srep10892

Conant, R. T., Six, J., & Paustian, K. (2003). Land use effects on soil carbon fractions in the

southeastern United States. I. Management-intensive versus extensive grazing. Biology and

Fertility of Soils, 38(6), 386–392. https://doi.org/10.1007/s00374-003-0652-z

Costa, C., Villegas, D. M., Bastidas, M., Matiz-Rubio, N., Rao, I., & Arango, J. (2022). Soil

carbon stocks and nitrous oxide emissions of pasture systems in Orinoquía region of

Colombia: Potential for developing land-based greenhouse gas removal projects. Frontiers

in Climate, 4, 916068. https://doi.org/10.3389/fclim.2022.916068

Díaz De Otálora, X., Epelde, L., Arranz, J., Garbisu, C., Ruiz, R., & Mandaluniz, N. (2021).

Regenerative rotational grazing management of dairy sheep increases springtime grass

production and topsoil carbon storage. Ecological Indicators, 125, 107484.

https://doi.org/10.1016/j.ecolind.2021.107484

Dillon, J., & Machmuller, M. (2021). Regenerative Grazing, Carbon, and Climate (pp. 1–16)

[White Paper]. Colorado State University.

https://pastureproject.org/publications/regenerative-grazing-carbon-and-climate/

Dong, L., Martinsen, V., Wu, Y., Zheng, Y., Liang, C., Liu, Z., & Mulder, J. (2021). Effect of

grazing exclusion and rotational grazing on labile soil organic carbon in north China.

European Journal of Soil Science, 72(1), 372–384. https://doi.org/10.1111/ejss.12952

Efron, B. (1979). Bootstrap Methods: Another Look at the Jackknife. The Annals of Statistics,

7(1). https://doi.org/10.1214/aos/1176344552

Eshel, G., Flamholz, A. I., Shepon, A., & Milo, R. (2025). US grass-fed

beef is as carbon intensive as industrial beef and ≈10-fold more

intensive than common protein-dense alternatives. Proceedings

of the National Academy of Sciences, 122(12), e2404329122.

https://doi.org/10.1073/pnas.2404329122

FAO. (2022). Global Livestock Environmental Assessment Model Description.

Gerber, P., Steinfeld, H., Henderson, B., Mottet, A., Opio, C., Dijkman, J., Falcucci, A., &

Tempio, G. (2013). Tackling climate change through livestock: a global assessment of

emissions and mitigation opportunities (Food and Agriculture Organization of the United

Nations (FAO), Ed.). FAO.

Giller, K. E., Hijbeek, R., Andersson, J. A., & Sumberg, J. (2021). Regenerative Agriculture: An

agronomic perspective. Outlook on Agriculture, 50(1), 13–25.

https://doi.org/10.1177/0030727021998063

Halpern, B. S., Frazier, M., Verstaen, J., Rayner, P.-E., Clawson, G., Blanchard, J. L., Cottrell,

R. S., Froehlich, H. E., Gephart, J. A., Jacobsen, N. S., Kuempel, C. D., McIntyre, P. B.,

Metian, M., Moran, D., Nash, K. L., Többen, J., & Williams, D. R. (2022). The

environmental footprint of global food production. Nature Sustainability, 5(12), 1027–1039.

https://doi.org/10.1038/s41893-022-00965-x

Hao, X., You, M., Han, X., Li, H., Zou, W., & Xing, B. (2017). Redistribution of Different

Organic Carbon Fractions in the Soil Profile of a Typical Chinese Mollisol with Land-Use

Change. Communications in Soil Science and Plant Analysis, 48(20), 2369–2380.

https://doi.org/10.1080/00103624.2017.1358741

Hillenbrand, M., Thompson, R., Wang, F., Apfelbaum, S., & Teague, R. (2019). Impacts of

holistic planned grazing with bison compared to continuous grazing with cattle in South

Dakota shortgrass prairie. Agriculture, Ecosystems & Environment, 279, 156–168.

https://doi.org/10.1016/j.agee.2019.02.005

Howard, P. J. A. (1965). The Carbon-Organic Matter Factor in Various Soil Types. Oikos, 15(2),

229. https://doi.org/10.2307/3565121

Khatri-Chhetri, U., Thompson, K. A., Quideau, S. A., Boyce, M. S., Chang, S. X., Bork, E. W.,

& Carlyle, C. N. (2024). Adaptive multi-paddock grazing increases mineral associated soil

carbon in Northern grasslands. Agriculture, Ecosystems & Environment, 369, 109000.

https://doi.org/10.1016/j.agee.2024.109000

Kurtz, D., Rey Montoya, S., & Ybarra, D. (2020). Impacto Del Pastoreo En Propiedades Fisico-

Quimicas De Un Psammacuent En Pastizales Del Nordeste Argentino. Revista Argentina

De Producción Animal, 40.

Leu, A. (2024, September 24). Response to Criticisms of Regenerative Grazing. Regeneration

International. https://regenerationinternational.org/2024/09/24/regenerative-grazing/

Lunt, H. A. (1931). THE CARBON-ORGANIC MATTER FACTOR IN FOREST SOIL

HUMUS: Soil Science, 32(1), 27–34. https://doi.org/10.1097/00010694-193107000-00003

Machmuller, M. B., Kramer, M. G., Cyle, T. K., Hill, N., Hancock, D., & Thompson, A. (2015).

Emerging land use practices rapidly increase soil organic matter. Nature Communications,

6(1), 6995. https://doi.org/10.1038/ncomms7995

McClelland, S. C., Paustian, K., & Schipanski, M. E. (2021). Management of cover crops in

temperate climates influences soil organic carbon stocks: a meta-analysis. Ecological

Applications, 31(3), e02278. https://doi.org/10.1002/eap.2278

McDonald, S. E., Reid, N., Waters, C. M., Smith, R., & Hunter, J. (2018). Improving ground

cover and landscape function in a semi-arid rangeland through alternative grazing

management. Agriculture, Ecosystems & Environment, 268, 8–14.

https://doi.org/10.1016/j.agee.2018.08.021

Mehre, J., Schneider, K., Jayasundara, S., Gillespie, A., & Wagner-Riddle, C. (2024). Adaptive
multi-paddock grazing increases soil carbon stocks and decreases the carbon footprint of
beef production in Ontario, Canada. Journal of Environmental Management, 371, 123255.
https://doi.org/10.1016/j.jenvman.2024.123255

Moore, J. M., Manter, D. K., & Maczko, K. A. (2023). Rotational Grazing Strategies Minimally
Impact Soil Microbial Communities and Carbon Dynamics—A Texas Case Study. Land,
12(8), 1517. https://doi.org/10.3390/land12081517

Mosier, S., Apfelbaum, S., Byck, P., Calderon, F., Teague, R., Thompson, R., & Cotrufo, M. F.
(2021). Adaptive multi-paddock grazing enhances soil carbon and nitrogen stocks and
stabilization through mineral association in southeastern U.S. grazing lands. Journal of
Environmental Management, 288, 112409. https://doi.org/10.1016/j.jenvman.2021.112409

Orgill, S. E., Condon, J. R., Conyers, M. K., Greene, R. S. B., Morris, S. G., & Murphy, B. W.
(2014). Sensitivity of soil carbon to management and environmental factors within
Australian perennial pasture systems. Geoderma, 214–215, 70–79.
https://doi.org/10.1016/j.geoderma.2013.10.001

Orgill, S. E., Waters, C. M., Melville, G., Toole, I., Alemseged, Y., & Smith, W. (2017).
Sensitivity of soil organic carbon to grazing management in the semi-arid rangelands of
south-eastern Australia. The Rangeland Journal, 39(2), 153. https://doi.org/10.1071/rj16020

Pelton, R. E. O., Kazanski, C. E., Keerthi, S., Racette, K. A., Gennet, S., Springer, N., Yacobson,
E., Wironen, M., Ray, D., Johnson, K., & Schmitt, J. (2024). Greenhouse gas emissions in
US beef production can be reduced by up to 30% with the adoption of selected mitigation
measures. Nature Food, 5(9), 787–797. https://doi.org/10.1038/s43016-024-01031-9

R Core Team. (2023). _R: A Language and Environment for Statistical Computing_ (Version
4.3.2). R Foundation for Statistical Computing. https://www.R-project.org/

Rowntree, J. E., Stanley, P. L., Maciel, I. C. F., Thorbecke, M., Rosenzweig, S. T., Hancock, D.
W., Guzman, A., & Raven, M. R. (2020). Ecosystem Impacts and Productive Capacity of a
Multi-Species Pastured Livestock System. Frontiers in Sustainable Food Systems, 4,
544984. https://doi.org/10.3389/fsufs.2020.544984

Sanderman, J., Reseigh, J., Wurst, M., Young, M.-A., & Austin, J. (2015). Impacts of Rotational
Grazing on Soil Carbon in Native Grass-Based Pastures in Southern Australia. PLOS ONE,
10(8), e0136157. https://doi.org/10.1371/journal.pone.0136157

Sanjari, G., Ghadiri, H., Ciesiolka, C. A. A., & Yu, B. (2008). Comparing the effects of

continuous and time-controlled grazing systems on soil characteristics in Southeast

Queensland. Soil Research, 46(4), 348. https://doi.org/10.1071/SR07220

Schatz, T., Ffoulkes, D., Shotton, P., & Hearnden, M. (2020). Effect of high-intensity rotational

grazing on the growth of cattle grazing buffel pasture in the Northern Territory and on soil

carbon sequestration. Animal Production Science, 60(15), 1814.

https://doi.org/10.1071/AN19552

Slowikowski, K. (2024). _ggrepel: Automatically Position Non-Overlapping Text Labels with

'ggplot2'_ (Version 0.9.6). https://CRAN.R-project.org/package=ggrepel

Stanley, P. L., Rowntree, J. E., Beede, D. K., DeLonge, M. S., & Hamm, M. W. (2018). Impacts

of soil carbon sequestration on life cycle greenhouse gas emissions in Midwestern USA

beef finishing systems. Agricultural Systems, 162, 249–258.

https://doi.org/10.1016/j.agsy.2018.02.003

Stanley, P., Roche, L., & Bowles, T. (2025). Amping up soil carbon: soil carbon stocks in

California rangelands under adaptive multi-paddock and conventional grazing management.

International Journal of Agricultural Sustainability, 23(1), 2461826.

https://doi.org/10.1080/14735903.2025.2461826

Subedi, A., Franklin, D., Cabrera, M., Dahal, S., Hancock, D., McPherson, A., & Stewart, L.

(2022). Extreme Weather and Grazing Management Influence Soil Carbon and Compaction.

Agronomy, 12(9), 2073. https://doi.org/10.3390/agronomy12092073

Teague, W. R., Dowhower, S. L., Baker, S. A., Haile, N., DeLaune, P. B., & Conover, D. M.

(2011). Grazing management impacts on vegetation, soil biota and soil chemical, physical

and hydrological properties in tall grass prairie. Agriculture, Ecosystems & Environment,

141(3–4), 310–322. https://doi.org/10.1016/j.agee.2011.03.009

Teutscherová, N., Vázquez, E., Sotelo, M., Villegas, D., Velásquez, N., Baquero, D., Pulleman,

M., & Arango, J. (2021). Intensive short-duration rotational grazing is associated with

improved soil quality within one year after establishment in Colombia. Applied Soil

Ecology, 159, 103835. https://doi.org/10.1016/j.apsoil.2020.103835

USDA NRCS. (2019). Soil Bulk Density/ Moisture / Aeration. Natural Resources Conservation

Service. https://www.nrcs.usda.gov/sites/default/files/2022-

10/Soil%20Bulk%20Density%20Moisture%20Aeration.pdf

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K.,

Ooms, J., Robinson, D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the

Tidyverse. Journal of Open Source Software, 4(43), 1686.

https://doi.org/10.21105/joss.01686

Yang, Y., Hobbie, S. E., Hernandez, R. R., Fargione, J., Grodsky, S. M., Tilman, D., Zhu, Y.-G.,

Luo, Y., Smith, T. M., Jungers, J. M., Yang, M., & Chen, W.-Q. (2020). Restoring

Abandoned Farmland to Mitigate Climate Change on a Full Earth. One Earth, 3(2), 176–

186. https://doi.org/10.1016/j.oneear.2020.07.019