Main Manuscript for

Offspring chemical control of adult reproductive transitions in a social insect

Baptiste Piqueret[1*], Jerrit Weissflog[2], Sandra Tretter[1,3], Tim Zetzsche[1,3], Daniel Veit[4], Stefan Bartram[5], Rayko Halitschke[2], Yuko Ulrich[1*]

Affiliations:

[1]Lise Meitner Research Group Social Behaviour, Max Planck Institute for Chemical Ecology, Jena, Germany

[2]Mass Spectrometry and Metabolomics, Max Planck Institute for Chemical Ecology, Jena, Germany

[3]Faculty of Biological Sciences, Friedrich Schiller University Jena, Germany

[4]Max Planck Institute for Chemical Ecology, Jena, Germany

[5]Department of Natural Product Biosynthesis, Max Planck Institute for Chemical Ecology, Jena, Germany

*Corresponding authors: Baptiste Piqueret; Yuko Ulrich

Email: bpiqueret@ice.mpg.de; yulrich@ice.mpg.de

Author Contributions: B.P.: conceptualization, investigation, formal analysis, methodology, visualization, writing—original draft and writing—review and editing; J.W.: investigation, methodology, formal analysis, writing—original draft, writing—review and editing; S.T.: investigation, writing—review and editing; T.Z.: investigation, writing—review and editing; D.V.: methodology; S.B.: formal analysis, writing—review and editing; R.H.: conceptualization, methodology, formal analysis, visualization, writing—original draft, writing—review and editing; Y.U: conceptualization, methodology, formal analysis, resources, project administration, supervision, writing—original draft, writing—review and editing. All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences - Ecology

Keywords: parental care, pheromone, clonal raider ant

This PDF file includes:

> Main Text
> Figures 1 to 3

Abstract

Parental care enhances offspring survival and growth but often entails a trade-off in which
caregivers temporarily suppress their own reproduction to invest in existing young. Such
reproductive transitions have independently evolved across diverse taxa, underscoring the
advantage of prioritising current over future reproduction. In vertebrates, the behavioural
sequence separating reproduction from parental care is controlled by offspring-derived cues.
While many insects display obligate parental care, the role of brood cues in regulating adult
reproduction remains unresolved outside advanced eusocial taxa, where reproductive plasticity
has largely been lost. Here, we investigate reversible reproductive transitions in the clonal raider
ant *Ooceraea biroi*, in which totipotent adults synchronously alternate between brood care and
egg laying under larval regulation. Using custom behavioural assays, we show that larvae inhibit
adult reproduction without physical contact, implicating volatile cues. Chemical analyses identified
a previously undescribed larval-specific compound, methyl 3-ethyl-2-hydroxy-4-methylpentanoate
(MEHMP), which is absent from other brood stages and ant species. Production of the compound
closely tracked larval development and colony cycles, linking its abundance to reproductive
synchrony. Exposure to synthetic MEHMP recapitulated the inhibitory effect of larval volatiles,
confirming its role as a pheromone that suppresses adult egg laying. To our knowledge, this is
the first identified brood pheromone that regulates reversible reproductive transitions in insects,
and the first brood pheromone described in ants. By providing a direct chemical link between
offspring presence and parental reproductive suppression, our findings highlight a convergent
strategy across distant evolutionary lineages.

Significance Statement
Parental care enhances offspring survival but induces profound physiological changes in
caregivers. A common feature across animals is that parents temporarily suppress their own
reproduction while caring for their young. The offspring cues enforcing these cycles remain poorly
understood, especially in insects. Using the clonal raider ant *Ooceraea biroi*, in which all females
naturally alternate between reproduction and brood care, we identify a larval pheromone that
suppresses adult egg laying. This pheromone is a previously undescribed chemical compound
produced only by larvae, and exposure to its synthetic version mimics the full biological effect of
larvae volatiles. By providing a direct chemical link between offspring presence and adult
reproductive suppression, our work highlights a general principle of offspring–parent
communication that shapes reproductive investment strategies across diverse taxa.

Main Text
34
35 **Introduction**
36
Parental care provides significant benefits by increasing offspring survival and growth, while
eliciting profound physiological and behavioural changes in caregivers. In species with offspring
that require extended care, parental care often induces a trade-off wherein caregivers temporarily
suppress their reproductive activity. This phenomenon has independently evolved across diverse
taxa, suggesting a selective advantage to prioritising existing offspring over new reproductive
attempts.

In vertebrates, the adaptive behavioural sequence that separates reproduction and brood care
is controlled by offspring-derived cues. For example, tactile cues associated with suckling
contribute to lactational infertility by promoting prolactin release in mammals (1, 2). Similarly,
tactile cues associated with egg incubation reduce egg laying by increasing prolactin secretion in
many birds (3–5). The temporary suppression of reproduction is lifted when these offspring-
derived cues decrease or disappear, e.g., at weaning, allowing parents to resume reproductive
activity.

Although many insects exhibit obligate parental care (6), the role of brood-derived cues in regulating reproductive transitions in adult caregivers remains unresolved. Current knowledge on insect brood-derived cues is largely based on advanced eusocial insects, in which queens monopolise reproduction and brood care is restricted to workers. In these derived systems, individual reproductive transitions no longer occur because egg laying and caregiving are performed by separate, morphologically specialised colony members, who have largely lost the ability to switch between reproductive and caregiving roles. Accordingly, the only brood-derived cues known to affect adult reproduction, the honeybee brood pheromones (*E*)-β-ocimene (7) and a mix of ten esters (8), reinforce reproductive division of labour by suppressing worker ovarian development but do not regulate individual reproductive transitions. Consistent with the requirement for permanent reproductive suppression of workers, even brood stages that do not require active care (e.g., capped brood) produce brood pheromones (9, 10), and several of its active components are also produced by queens (11, 12), suggesting inbuilt redundancy in the enforcement of worker sterility. Thus, even in the absence of brood-derived cues, honeybee workers fail to restore full reproductive potential, underscoring the extent to which reproductive plasticity has been lost in advanced social taxa. To uncover brood cues regulating adult reproductive transitions, it is therefore necessary to investigate systems that display this ancestral plasticity.

For this purpose, we used the clonal raider ant *Ooceraea biroi*. In this species, all individuals are totipotent and alternate between reproductive and brood care states. This cycle is regulated by larvae, which inhibit reproduction and induce brood care behaviour in adults (13–15). Because the presence of larvae in a colony is periodic, entire colonies alternate between brood care phases (larvae are present, workers do not lay eggs) and reproductive phases (larvae are absent, workers lay eggs). Furthermore, clonal raider ant colonies are naturally queenless, eliminating the confounding effects of queen pheromones, the primary mechanism of reproductive suppression in most social insects (16).

To investigate how larvae regulate adult reproductive transitions in *O. biroi*, we first developed a custom behavioural assay to ask whether physical contact and/or volatile cues mediate the inhibitory effect of larvae on adult egg laying. These assays strongly suggest a role for volatile chemical cues in controlling the switch between reproductive and brood care states. Using chemical analysis of headspace volatiles of all brood stages, we identify the compound methyl 3-ethyl-2-hydroxy-4-methylpentanoate (MEHMP), which is exclusively produced by *O. biroi* larvae. Behavioural assays show that chemically synthesised MEHMP inhibits adult egg laying to the same extent as larval volatiles. Collectively, these studies implicate a larval pheromone in the regulation of reproductive transitions.

Results

We designed a behavioural platform to quantitatively study the influence of brood on adult egg laying. This platform utilised multiplexed 2-chamber arenas with controlled airflow, thereby allowing the passage of volatile compounds, but preventing physical contact of the ants between the chambers (Fig. 1A). Using this assay, we first asked whether larvae can inhibit adult egg laying without physical contact. To this aim, we placed live larvae in the first chamber and a small colony of 12 workers in the second chamber ("larvae distance" treatment). As controls, workers were placed in physical contact with larvae ("larvae contact" treatment) in the second chamber (mimicking the brood care phase) or in physical contact with pupae ("no larvae" treatment) in the second chamber (mimicking the reproductive phase). Clonal raider ant stock colonies lay no eggs in the brood care phase and lay approximately one egg per ant in the reproductive phase. As expected, ants in physical contact with larvae laid no eggs (median (interquartile range IQR): 0, (0-0) eggs per worker; Fig. 1B) and, in contrast, ants in physical contact with pupae laid eggs (1.08 (1.00-1.17) eggs per worker). Thus, our custom assay recapitulates the phenotype of stock colonies in the two phases of their reproductive cycle. Although the presence of larvae in the first

chamber did not fully recapitulate the effect of contact with larvae (Dunn's test, larvae contact -
larvae distance: Z = -2.84, p < 0.01), this arrangement strongly reduced adult egg laying (0.46
(0.40-0.59) eggs per ant, ca. 58% reduction; larvae distance - no larvae: Z = -2.84, p < 0.01).
Thus, larvae reduce adult reproduction without physical contact, suggesting a role for volatile
chemical compounds in the regulation of transitions between reproductive and brood care states.

To identify candidate volatile compounds responsible for the observed effect on adults, we
collected volatiles in the headspace of different *O. biroi* brood development stages and of other
ant species using solid phase micro extraction (SPME) and analysed them using gas-
chromatography mass-spectrometry (GC-MS). We reasoned that any compound responsible for
the observed effect should be present in larvae headspace (the only brood stage inhibiting adult
egg laying) and absent in the headspace of other brood stages (i.e., eggs, prepupae, pupae) of
O. biroi. Because *O. biroi* feeds on other ant species (17), any candidate compound should also
be absent in other ant species, so that *O. biroi* reproduction would not be inhibited by its prey.
Using this approach, we identified a single larval-specific compound with a retention time of 8.8
min, which was produced only by *O. biroi* larvae (Fig.2A), but not by alien ant brood (*Tetramorium*
bicarinatum and *Iridomyrmex purpureus*, Fig. S1). The mass spectrum showed a strong *m/z* 90
base peak, which was used in extracted ion chromatograms for detailed comparisons and
compound quantification (Fig. 2A, Fig S1-S3). The *m/z* 90 ion is a diagnostic fragment of a
McLafferty rearrangement of α-hydroxycarboxylic acid methyl esters, and initial library searches
and further fragment analysis suggested a branched 8-carbon carboxylic acid methyl ester. We
synthesised several isomers with different branching patterns and identified the larval volatile
compound as methyl 3-ethyl-2-hydroxy-4-methylpentanoate (MEHMP), a previously undescribed
compound (Fig. 2B). The synthesised standard showed identical retention behaviour on different
GC stationary phases as the natural compound, as well as the same mass spectrum (Fig. 2C).
The GC-MS analyses suggest a stereospecific production of MEHMP by the larvae but the
absolute configuration of the pheromone compound could not be resolved yet, as the chemical
synthesis yielded a racemic mixture of the standard.

If MEHMP controls individual and colony phasic reproduction, we would expect its production to
correlate with natural brood development and colony phases. To test this, we quantified daily
MEHMP abundances in the headspace of colonies of 100 ants that naturally progressed from the
reproductive to the brood care phase, and back to the reproductive phase, using volatile
collection on polydimethyl siloxane (PDMS) tubing (Fig. 2D). We find that the abundance of the
compound closely mirrors cycle phases and brood development. It was undetectable in colonies
as adults laid eggs, appeared after eggs hatched into larvae, increased in abundance as larvae
grew over the course of the brood care phase, and waned shortly before larvae became
prepupae, which marks the cessation of larval feeding and the start of a new reproductive phase.
Thus, life stage specific (Fig. 2A) and colony-level, temporal (Fig. 2D) measurements show that
MEHMP is specific to larvae and tracks colony cycles, making it a viable candidate larval
pheromone controlling adult reproductive transitions and colony reproductive cycles.

Based on both sampling methods (SPME and PDMS), we estimate that a 4th instar *O. biroi* larva
produces approximately 1-2 pg of MEHMP per day (PDMS, day 19: mean ± s.d.:1.07 ± 0.63 pg
day^{-1}, n = 2; SPME: 2.13 ± 0.12 pg day^{-1}, n = 3; see Methods).

We next used our behavioural platform to assess whether MEHMP affects adult reproduction
(Fig. 3). Ant groups were exposed to 200 pg of synthetic MEHMP per day (ca. 100-200 larvae
equivalents) ("MEHMP") and compared to the same groups used previously, i.e. ants in physical
contact with larvae ("larvae contact"), ants without physical contact with larvae ("larvae distance"),
and ants with physical contact with pupae ("no larvae", Fig. 3A). As a minor adjustment, pupae
were included in all the treatments to control for potential pupal stimulation of egg laying.
Exposure to MEHMP alone induced a 33% reduction in egg laying compared to control (MEHMP:
0.67 (0.54-0.75) eggs per ant, no larvae: 1.00 (0.92-1.04), MEHMP - no larvae: Z = -4.39, p <
0.001; Fig. 3B) and recapitulated the effect of exposure to the full blend of larval volatiles (larvae

distance: 0.67 (0.50-0.77) eggs per ant; MEHMP – larvae distance: Z = -0.84, p = 0.40, Fig. 3B).
Latency to lay eggs did not vary across treatments in which eggs were laid (Kruskal-Wallis test:
χ^2 = 4.13, d.f. = 3, p = 0.25), indicating that egg laying was in fact inhibited and not merely
delayed.

Additionally, to test the specificity of the effect of MEHMP, we included a treatment with 200 pg
per day of its non-hydroxylated synthetic precursor, methyl 3-ethyl-4-methylpentanoate (MEMP),
which is absent from *O. biroi* larvae but was previously identified as a pheromone in another ant
(18). Exposure to MEHMP inhibited adult egg-laying more strongly than MEMP, indicating that
the response is compound-specific and the hydroxylation modification plays a role in the
pheromone function (Fig. S4; MEMP: 0.83 (0.67-0.94) eggs per ant, MEHMP - MEMP: Z = -2.38,
p < 0.05).

Discussion
164

Here, we show that a previously undescribed volatile compound, produced exclusively by larvae,
regulates reproductive transitions in adult *O. biroi* ants by inhibiting egg laying. Because larvae
occur only periodically in colonies, this pheromone likely contributes to reproductive synchrony,
coordinating colony-level transitions between brood care and reproductive phases. To our
knowledge, this represents the first identified larval pheromone that regulates reproductive
transitions in an insect, and the first brood pheromone reported in ants, whose existence has until
now remained uncertain (19, 20). While MEHMP was previously undescribed, the structurally
MEMP was identified in adult *Formica* and *Polyergus* ants (18, 21), though this compound likely
has a different function in these species. In our experiment, exposure to MEMP did not
recapitulate the effects of MEHMP, showing the effects on adult reproduction are specific to the
larval pheromone.

Past work in *O. biroi* has shown that larvae control adult ovarian activation by influencing adult
insulin signalling (22). The presence of larvae inhibits adult reproduction by suppressing *insulin-*
like peptide 2 (*ilp2*). Experimentally elevating ILP2 in adults overrides larval suppression of
worker reproduction, breaking the colony's reproductive cycle. The larval pheromone (MEHMP)
we have identified is a strong candidate as the proximate signal modulating ILP2 levels,
functioning upstream of insulin-controlled endocrine regulation. In this light, decreasing adult
sensitivity to this pheromone would likely disrupt the colony cycle, similar to the effect of artificially
elevating ILP2. This integration of pheromone signalling with insulin-mediated reproductive
control brings us closer to a complete mechanistic understanding of how a complex social
interaction, larval control of adult reproduction, is orchestrated in this system.

The volatility of MEHMP likely represents an important requirement for ensuring colony-wide
enforcement of reproductive synchrony. Like other social insects, *O. biroi* exhibits division of
labour (23), with some workers (e.g., foragers) having limited direct contact with larvae. If
reproductive inhibition relied solely on contact-based cues, such individuals could escape
suppression, potentially disrupting colony-wide synchrony. A volatile signal ensures that all
workers, regardless of their tasks, are exposed to larvae-derived inhibition and remain in the non-
reproductive state during the brood care phase. In this way, the pheromone provides a
mechanistic basis for how colonies maintain coordinated shifts between reproduction and brood
care.

The larval pheromone of *O. biroi* differs fundamentally from brood signals in advanced eusocial
insects, yet provides clues to the mechanisms by which reproductive division of labour could have
evolved. The honeybee brood pheromones act redundantly with queen pheromones to
permanently suppress worker reproduction, rather than to regulate reversible reproductive
transitions (7, 8). In *O. biroi*, the pheromone transiently suppresses worker reproduction, allowing
all individuals to alternate between brood care and egg laying. This dynamic is reminiscent of the

subsocial cycle thought to be the evolutionary precursor to eusociality, in which ancestral solitary wasps alternated between laying an egg and caring for the resulting larva until pupation (24). Natural variation in responsiveness to larval pheromones could have provided a foundation for the evolution of the reproductive division of labour that defines eusociality. Individuals more strongly inhibited by larval cues would have been less likely to reproduce and instead focused on brood care, whereas less sensitive individuals would have continued reproducing despite larval presence. Continued reproduction by less sensitive individuals would have maintained a constant presence of larvae, thereby reinforcing a shift toward reproductive monopolisation by a few individuals. Such asymmetry in reproductive outcomes could then have been stabilised and amplified by natural selection, ultimately giving rise to the fixed reproductive (queen) and non-reproductive (worker) castes characteristic of advanced eusocial insects. In this view, the cyclical, brood-regulated reproduction of *O. biroi* may represent a model of the ancestral condition from which strict reproductive division of labour evolved.

Although our study identifies a volatile larval pheromone as a central regulator of reproductive transitions in *O. biroi*, several questions remain. While the synthetic pheromone fully recapitulates the biological effect of larval volatiles, it only partially reproduces the effect of contact with larvae, suggesting additional contributions from other larval cues. Future work should determine whether other non-volatiles chemical cues or non-olfactory cues (e.g., tactile cues) contribute to reproductive suppression. At the mechanistic level, elucidating the sensory pathways by which the pheromone is perceived and how this perception is coupled to endocrine regulation of reproduction will be crucial for establishing a complete model. Finally, comparative studies in other ants, as well as subsocial and primitively eusocial species could reveal whether larval pheromones represent an ancestral regulator of adult reproductive transitions or a derived innovation of *O. biroi*. Larvae suppress worker reproduction in at least one primitively eusocial bee (25) and three ant species (26–28), making these promising systems in which to discover additional larval pheromones.

In summary, we identify a larval pheromone that regulates reproductive transitions in an insect, providing a direct link between brood care and reproductive suppression. Our discovery adds insects to the growing list of taxa in which offspring-derived cues temporarily suppress parental reproduction, highlighting a convergent solution to the fundamental trade-off between current and future reproduction. Just as offspring-derived tactile cues reduce parental fertility in mammals and birds, larval pheromones in *O. biroi* inhibit adult reproduction through chemical communication. This convergence underscores the central role of offspring signals in mediating parental investment strategies across distant evolutionary lineages.

Materials and Methods

Egg laying behavioural assay

We developed a platform to quantify the effects of brood, brood volatile organic compounds (VOCs), and synthetic compounds on adult egg laying behaviour. The platform consisted of multiplexed 2-chamber arenas (65 mm × 20 mm) coupled with an odorant delivery system with controlled airflow. Arenas were 3D-printed in polylactic acid and comprised an upstream round chamber (diameter: 12 mm) closed with an airtight metallic lid and a downstream rectangular chamber (28 x 11 mm) closed with a clear acrylic lid, allowing egg counting without colony disturbance. The chambers were separated by a metallic gate (diameter: 8.5 mm, width: 2 mm) that allowed air flow but prevented physical contact (including antennal contact) between ants and brood in the different chambers. Both chambers had a humidified plaster-of-Paris floor. The input and output air flows of each arena were set by analogous flow controllers (Key Instruments, USA). Air passed through two 250 mL bottles filled with distilled water and one 250 mL odorant delivery bottle (see below) before entering the upstream chamber and carrying any volatile compounds from that chamber to the downstream chamber.

Experiment 1 (Fig. 1A). We tested whether larvae can inhibit adult egg laying without physical
contact using three treatments: *i*) "larvae contact": 12 adults (16-day old, genetic lineage A (29))
were placed in direct contact with 12 larvae (6-day old, genetic lineage B) in the downstream
chamber; this treatment is analogous to the brood care phase, *ii*) "no larvae": 12 adults were
placed in direct contact with 12 pupae (5-day old, genetic lineage D) in the downstream chamber;
this treatment is analogous to the reproductive phase, *iii*) "larvae distance": 36 larvae were placed
in the upstream chamber, and 12 adults in the downstream chamber. Twelve replicate arenas
were used for each treatment. Batches of four arenas were connected to the same flow controller.
The natural larvae-to-worker ratio in *O. biroi* colonies is ca. 1:1; in (iii), we used a 3:1 larvae-to-
worker ratio to increase the likelihood of capturing an effect, if it existed. Air was pushed into each
arena at a 75 mL min^{-1}. For all arenas, a 4 mL vial of hexane (an organic solvent) was placed in
the odorant delivery bottle and exchanged daily. Hexane exposure was included to ensure it did
not prevent ant survival and egg laying, with future odorant-delivery experiments in mind.

Experiment 2 (Fig. 3). We tested whether the putative larval pheromone inhibits adult egg laying,
using a treatment with methyl 3-ethyl-2-hydroxy-4-methylpentanoate (MEHMP) delivery in
addition to the three treatments used previously. To test specificity of effects, we included a
treatment with methyl 3-ethyl-4-methylpentanoate (MEMP) delivery, a synthetic precursor of
MEHMP that is not produced by *O. biroi* larvae. As before, a 4 mL vial of hexane was placed in
the odorant delivery bottle of each arena and exchanged daily. In contrast to Experiment 1, pupae
were included in all groups to control for potential pupal stimulation of egg laying. Additionally, we
introduced three adults in the upstream chamber of all treatments to act as caregivers for the
larvae. The five treatments were: *i*) "larvae contact": 12 adults (11-day old, genetic lineage B)
were placed in direct contact with 12 larvae (6-day old, genetic lineage B) and 12 pupae (0-day
old, genetic lineage B) in the downstream chamber; *ii*) "no larvae": 12 adults were placed in direct
contact with 12 pupae in the downstream chamber; *iii*) "larvae distance": 12 larvae were placed in
the upstream chamber, and 12 adults were placed in the downstream chamber with 12 pupae; *iv*)
"MEHMP": 12 adults were placed in the downstream chamber with 12 pupae; 200 pg of methyl 3-
ethyl-2-hydroxy-4-methylpentanoate were added to the hexane vial, *v*) "MEMP": 12 adults were
placed in the downstream chamber with 12 pupae; 200 pg of methyl 3-ethyl-4-methylpentanoate
were added to the hexane vial. Sixteen replicate arenas were used per treatment, except for (*i*),
which had 4 replicates. Because ants in this treatment invariably laid no eggs in Experiment 1, we
included this treatment as a positive control but excluded it from statistical analyses in Experiment
2, to allow sufficient replication in the four other treatments. Air was pushed into each arena at 75
mL min^{-1}. Additionally, to minimise leaks of the synthetic compounds, air was actively pumped out
of each arena at 87.5 mL min^{-1} and filtered through activated carbon before release into the
experimental room.

In both experiments, the survival of adults, larvae, and pupae, as well as the number of eggs laid
was monitored daily. Dead larvae and pupae were replaced each day to maintain constant brood
exposure; dead adults were removed but not replaced. Arenas were humidified daily but received
no food to ensure that any differences in egg laying were not driven by food intake or foraging
behaviour. We calculated the median latency to lay eggs in the negative control ("no larvae") and
ended the experiments 3 days later, which occurred on day 8 in Experiment 1 and on day 9 in
Experiment 2.

In both experiments, adult survival was high (>93% in Experiment 1, >95% in Experiment 2).
Arenas in which >33% of adults died or escaped were excluded from further analyses, resulting in
the exclusion of no arenas in Experiment 1 and two arenas in Experiment 2.

Identification of the candidate pheromone

To identify candidate volatile compounds responsible for adult reproductive inhibition, we
collected volatiles in the headspace of different life stages using solid phase micro extraction
(SPME) and analysed them using gas chromatography-mass spectrometry (GC-MS). We
incubated 50-100 larvae (n = 18, mixed instars), 200 eggs (n = 3), 50-200 prepupae (n = 5) and

100 pupae (n = 7) in 1.5 mL glass vials with a SPME fibre (DVB/CAR/PDMS, Supelco, USA)
inserted through the septum cap (PTFE, Macherey-Nagel, Germany) and exposed them to the
brood headspace for 18 h at 28 °C. Additionally, we used the same method to collect headspace
extracts from the brood of two other ant species, *T. bicarinatum* (15 larvae, n = 1; 10 pupae, n =
3) and *I. purpureus* (3-10 larvae, n = 5; 2 pupae, n = 1), maintained at 25 °C in a separate room.
After sampling, the fibre was desorbed by splitless injection at 230 °C into a Shimadzu QP2010
GC-MS, equipped with a SLB-5MS GC column (30 m × 0.25 mm × 0.25 µm, Supelco, USA) and
helium as carrier gas (1.2 mL min^{-1}). The oven temperature was programmed at 50 °C for 1 min,
then increased to 260 °C at 10 °C min^{-1}, then to 320 °C at 20 °C min^{-1} and held for 5 min. Mass
spectra were recorded with electron impact ionisation at 70 eV. The software GCMSsolution
(v4.20, Shimadzu Corporation) was used for data analysis. Identification of the compounds was
accomplished by comparison of library databases, diagnostic ions and retention indices (RI)
calculated with a standard n-alkane solution (C7–C30, 99%, Sigma-Aldrich). The candidate
pheromone shows a RI of 1152 on the unipolar SLB-5ms and additional analyses on a polar
phase (ZB-WAX, Phenomenex) yielded a RI of 1635.

Chemical synthesis

Commercially available chemicals were used without further purification. All reactions were
carried out under an argon atmosphere. Preparative column chromatography was performed on
Silica gel 60 (230-400 mesh, Carl Roth GmbH) and TLC analysis on commercial Merck silica gel
60 F$_{254}$ plates. NMR spectra were measured on a Bruker Avance 400 NMR spectrometer.
Chemical shifts are reported in ppm downfield from TMS.

<u>Synthesis of methyl 3-ethyl-4-methylpentanoate (MEMP)</u>

A 250 mL three-necked flask, equipped with rubber septum and dropping funnel, was charged
with 1.18 g copper(I)-iodide (6.1 mmol) and put under an atmosphere of argon. The flask was
loaded with 70 mL anhydrous THF and 7 g of methyl (*Z*)-pent-2-enoate (61 mmol) were added.
Chlorotrimethylsilane (9.4 mL, 74 mmol) was then added via syringe. After stirring for 15 min at
20 °C, the mixture was cooled to -15°C in an ice/methanol bath. A 2 M solution of
isopropylmagnesium chloride in THF (36 mL, 74 mmol, Merck) was added dropwise over 2 hours,
while keeping the temperature below -10 °C. After stirring for an additional hour at -15 °C, the
reaction was slowly quenched by adding methanol (5 mL) and 2 M ammonium chloride solution
(100 mL). After separation of the liquid layers, the aqueous phase was extracted twice with ~75
mL diethyl ether. The organic phases were pooled and washed with water and brine (100 mL
each), dried with anhydrous MgSO$_4$, filtered and concentrated in vacuo. Vacuum distillation gave
MEMP (7.5 g, 76% yield) as a colourless liquid (B.p.110-120 °C at 50 mbar). ^{1}H-NMR: (400 MHz,
CDCl$_3$) d = 3.51 (s, 3H), 2.07 (qd, J_1 = 21.9 Hz, J_2 = 6.7 Hz, 2H), 1.50-1.60 (m, 2H), 1.01-1.20 (m,
2H), 0.89 (m, 9H) ppm; ^{13}C-NMR: (100 MHz, CDCl$_3$) d = 174.7, 51.4, 42.6, 35.6, 29.4, 23.7, 19.4,
18.5, 11.6 ppm. EI-MS: *m/z* (rel int.) = 127 (10), 115 (13), 101 (6), 87 (31), 85 (33), 83 (18), 74
(100), 70 (23), 59 (18), 57 (16), 43 (27).

<u>Synthesis of methyl 3-ethyl-2-hydroxy-4-methylpentanoate (MEHMP)</u>

MEMP (460 mg, 2.9 mmol) was dissolved in 6 mL anhydrous THF. The solution was cooled to -
78 °C and a 1 M solution of potassium bis(trimethylsilyl)amide (4 mL, 40 mmol) was added via
syringe. The mixture was allowed to warm to -40 °C and stirred for 1 h at this temperature before
being cooled to -78 °C again. Davis oxaziridine (3-phenyl-2-phenylsulfonyl-1,2-oxaziridine, 900
mg, 3.5 mmol) was dissolved in 2 mL anhydrous THF and added via syringe to the reaction
mixture. After stirring for 1 h at -78 °C, the mixture was quenched with 2 M ammonium chloride
solution (10 mL). The mixture was vigorously stirred at room temperature for 1 h, before
separation of the liquid layers. The aqueous phase was extracted with diethyl ether (2 x 10 mL)
and the combined organic phases were washed with water and brine (10 mL each), before being
dried with anhydrous MgSO$_4$, filtered and concentrated in vacuo. Column chromatography (silica
gel, 9:1 n-hexane/ethylacetate v/v as eluent) gave MEHMP (370 mg, 73% yield) as a colourless

oil. GC-MS-analyses (WAX-column) and NMR analysis showed that two pairs of diastereomers
are present in a ratio of 8:3.

Diastereomer A: [1]H-NMR: (400 MHz, CDCl$_3$): d = 4.34 (dd, J_1 = 4.9 Hz, J_2 = 2.2 Hz, 1H), 3.786 (s,
3H), 2.72 (d, J = 4.96 Hz, 1H, (OH)), 1.80-1.90 (m, 1H), 1.30-1.60 (m, 3H), 0.98 (d, J = 6.8 Hz,
6H), 0.86 (t, 3H) ppm; [13]C-NMR: (100 MHz, CDCl$_3$) d = 176.9, 71.2, 52.4, 49.3, 29.2, 20.6, 20.0,
19.4, 12.8 ppm. EI-MS: m/z (rel int.) = 115 (23), 97 (26), 90 (100), 85 (44), 73 (9), 71 (14), 59
(24), 57 (31), 55 (30), 43 (86), 41 (30).

Diastereomer B: [1]H-NMR: (400 MHz, CDCl$_3$): d = 4.30 (dd, J_1 = 4.7 Hz, J_2 = 3.8 Hz, 1H), 3.794 (s,
3H), 2.65 (d, J = 5.15 Hz, 1H, (OH)), 1.68-1.78 (m, 1H), 1.30-1.60 (m, 3H), 0.96 (t, J = 6.5 Hz,
3H), 0.91 (d, J = 6.5 Hz, 3H), 0.90 (t, 3H) ppm; [13]C-NMR: (100 MHz, CDCl$_3$) d = 176.7, 72.0,
52.37, 49.2, 28.4, 21.9, 19.38, 19.2, 12.5 ppm. EI-MS: m/z (rel int.) = 115 (23), 97 (26), 90 (100),
85 (44), 73 (9), 71 (14), 59 (24), 57 (31), 55 (30), 43 (86), 41 (30).

Time-course experiment and candidate compound quantification

Three groups of 100 adult ants (mixed age, genetic lineage B) were placed in circular, airtight
plastic cups (height: 4 cm, base diameter: 3 cm, top diameter: 4 cm) with a humidified plaster-of-
Paris floor and kept at 28 °C. Once ants began laying eggs, nest headspace was collected daily
using 5-mm pieces of polydimethyl siloxane (PDMS) tubing suspended in the nest on a pin for 24
h. As a control, headspace was collected daily from identical plaster-lined cups without ants (n =
2). When eggs hatched into larvae, marking the start of the brood care phase, each colony was
fed frozen brood of *T. bicarinatum* every 3-4 days. The experiment ended 2 days after the
appearance of the first prepupae. We estimated that ca. 70–80 larvae were present in each nest
during the experiment. PDMS headspace samples were inserted into a Thermal Desorption Unit
(TDU) sampling tube and analysed on a Shimadzu QP2010 GC-MS equipped with a TD-20
thermal desorption system and an SLB-5MS GC column (30 m × 0.25 mm × 0.25 µm; Supelco,
USA). The carrier gas was helium (1 mL min^{-1}). Injections were splitless (230 °C). The oven
program was: 50 °C for 1 min, ramp to 260 °C at 10 °C min^{-1}, ramp to 320 °C at 20 °C min^{-1} and
hold at 320 °C for 5 min. Mass spectra were recorded with electron-impact ionisation at 70 eV.

The amount of MEHMP collected in the headspace was quantified using calibration curves. 1 µL
of a hexane solution containing 0 pg, 50 pg, 100 pg, 500 pg, or 1 ng of MEHMP (n = 3 for each
concentration) was added to the bottom of a 1.5 mL glass vial, with 5-mm pieces of PDMS tubing
suspended on a pin through the cap for 18 h. Samples were analysed as above. A calibration
curve was generated by comparing peak areas of the synthetic standard with those from natural
samples (Fig. S2). This analysis indicated that each larva produced ca. 1 pg day^{-1} of MEHMP at
the peak of production (when larvae were 12–13 days old, on experimental day 19).

To confirm this estimate, we used SPME fibre sampled from 4[th] instar larvae, (n = 3, 100 larvae)
(see "Identification of the candidate pheromone" above) and an independent calibration. Here, 1
µL of a hexane solution containing 0 pg, 50 pg, 100 pg or 250 pg of MEHMP (n = 3 for each
concentration) was placed at the bottom of a 1.5 mL vial and exposed to a SPME fibre
(DVB/CAR/PDMS) for 18 h. This analysis yielded an estimate of 2.1 pg day^{-1} per larva (Fig. S3),
consistent with the PDMS-based quantification.

Statistical analyses

All analyses were performed in R v4.5.1 (30). Kruskal–Wallis tests were used to compare egg
laying across treatments on the last experimental day. To account for adult mortality, we
modelled the number of eggs per ant (eggs per arena divided by the number of live adults).
Pairwise comparisons between treatments of interest were performed using Dunn's tests
(*dunn.test* function from package *dunn.test* (31)) with Benjamini-Hochberg adjustment for multiple
comparisons. In Experiment 1, we tested whether the treatment "larvae distance" differed from
the two other treatments (i.e. "larvae contact" and "no larvae"). In Experiment 2, we tested
whether the treatment "MEHMP" differed from the treatments "larvae distance", "MEMP" and "no
larvae". Kruskal-Wallis tests were used to compare latency to lay eggs across treatments.

Acknowledgments
We thank Martin Kaltenpoth, Sarah O'Connor and Tobias Koellner for helpful discussions, Jonghyun Park for ant pictures, Sybille Lorenz for technical assistance with chemical analyses, Danny Kessler for access to climate chambers, and the whole Lise Meitner Research Group Social Behaviour for support with the experiments. This is Clonal Raider Ant Project paper no. 40.

References

1. P. Hwang, H. Guyda, H. Friesen, A Radioimmunoassay for Human Prolactin. *Proc. Natl. Acad. Sci.* **68**, 1902–1906 (1971).

2. E. C. R. Hackwell, S. R. Ladyman, R. S. E. Brown, D. R. Grattan, Mechanisms of Lactation-induced Infertility in Female Mice. *Endocrinology* **164** (2023).

3. H. Klomp, The Determination of Clutch-Size in Birds a Review. *Ardea* **55**, 1–124 (1970).

4. M. Massaro, A. N. Setiawan, L. S. Davis, Effects of artificial eggs on prolactin secretion, steroid levels, brood patch development, incubation onset and clutch size in the yellow-eyed penguin (Megadyptes antipodes). *Gen. Comp. Endocrinol.* **151**, 220–229 (2007).

5. M. R. Hall, External stimuli affecting incubation behavior and prolactin secretion in the duck (*Anas platyrhynchos*). *Horm. Behav.* **21**, 269–287 (1987).

6. J. W. Y. Wong, J. Meunier, M. Kölliker, The evolution of parental care in insects: the roles of ecology, life history and the social environment. *Ecol. Entomol.* **38**, 123–137 (2013).

7. A. Maisonnasse, *et al.*, A scientific note on E-β-ocimene, a new volatile primer pheromone that inhibits worker ovary development in honey bees. *Apidologie* **40**, 562–564 (2009).

8. A. Mohammedi, A. Paris, D. Crauser, Y. Le Conte, Effect of Aliphatic Esters on Ovary Development of Queenless Bees (*Apis mellifera* L.). *Naturwissenschaften* **85**, 455–458 (1998).

9. A. Noël, *et al.*, Detailed chemical analysis of honey bee (Apis mellifera) worker brood volatile profile from egg to emergence. *PLOS ONE* **18**, e0282120 (2023).

10. F. Mondet, D. Beslay, M. Cousin, Y. Le Conte, Impact of temporal and rearing environment variations on brood ester pheromone production in the honey bee Apis mellifera. *Comptes Rendus Chim.* **26**, 119–128 (2025).

11. C. I. Keeling, K. N. Slessor, H. A. Higo, M. L. Winston, New components of the honey bee (Apis mellifera L.) queen retinue pheromone. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 4486–4491 (2003).

12. D. C. Gilley, G. DeGrandi-Hoffman, J. E. Hooper, Volatile compounds emitted by live European honey bee (*Apis mellifera* L.) queens. *J. Insect Physiol.* **52**, 520–527 (2006).

13. F. Ravary, B. Jahyny, P. Jaisson, Brood stimulation controls the phasic reproductive cycle of the parthenogenetic ant *Cerapachys biroi*. *Insectes Sociaux* **53**, 20–26 (2006).

14. Y. Ulrich, D. Burns, R. Libbrecht, D. J. C. Kronauer, Ant larvae regulate worker foraging
behavior and ovarian activity in a dose-dependent manner. *Behav. Ecol. Sociobiol.* **70**,
1011–1018 (2016).

15. S. L. Jud, D. Knebel, Y. Ulrich, Intergenerational genotypic interactions drive collective
behavioural cycles in a social insect. *Proc Biol Sci* **289**, 20221273 (2022).

16. A. Van Oystaeyen, *et al.*, Conserved class of queen pheromones stops social insect
workers from reproducing. *Science* **343**, 287–90 (2014).

17. F. Ravary, P. Jaisson, The reproductive cycle of thelytokous colonies of Cerapachys biroi
Forel (Formicidae, Cerapachyinae). *Insectes Sociaux* **49**, 114–119 (2002).

18. L. Greenberg, *et al.*, Sex Attractant Pheromones of Virgin Queens of Sympatric Slave-
Making Ant Species in the Genus Polyergus, and their Possible Roles in Reproductive
Isolation. *J. Chem. Ecol.* **44**, 547–555 (2018).

19. L. Morel, R. K. V. Meer, Do Ant Brood Pheromones Exist? *Ann. Entomol. Soc. Am.* **81**,
705–710 (1988).

20. E. Schultner, U. Pulliainen, Brood recognition and discrimination in ants. *Insectes Sociaux*
67, 11–34 (2020).

21. W. Francke, J. Borchert, D. Klimetzek, Volatile Constituents of the Red Wood Ant Formica
rufa L. (Hymenoptera: Formicidae). *Z. Für Naturforschung C* **40**, 661–664 (1985).

22. V. Chandra, *et al.*, Social regulation of insulin signaling and the evolution of eusociality in
ants. *Science* **361**, 398–402 (2018).

23. Y. Ulrich, J. Saragosti, C. K. Tokita, C. E. Tarnita, D. J. C. Kronauer, Fitness benefits and
emergent division of labour at the onset of group living. *Nature* **560**, 635–638 (2018).

24. M. J. West-Eberhard, "Flexible strategy and social evolution" in *Animal Societies: Theories*
and Facts, Japan Scientific Societies Press, 1987, pp. 35–51.

25. J. Starkey, A. Brown, E. Amsalem, The road to sociality: brood regulation of worker
reproduction in the simple eusocial bee. *Anim. Behav.* **154**, 57–65 (2019).

26. J. D. Ebie, B. Hölldobler, J. Liebig, Larval regulation of worker reproduction in the
polydomous ant Novomessor cockerelli. *Sci. Nat.* **102** (2015).

27. I. Villalta, E. Angulo, S. Devers, X. Cerdá, R. Boulay, Regulation of worker egg laying by
larvae in a fission-performing ant. *Anim. Behav.* **106**, 149–156 (2015).

28. J. Heinze, B. Trunzer, P. S. Oliveira, B. Hölldobler, Regulation of reproduction in the
neotropical ponerine ant, Pachycondyla villosa. *J. Insect Behav.* **9**, 441–450 (1996).

29. W. Trible, S. K. McKenzie, D. J. C. Kronauer, Globally invasive populations of the clonal
raider ant are derived from Bangladesh: Source of the clonal raider ant invasion. *Biol. Lett.*
16, 1–5 (2020).

30. R Development Core Team, R: A Language and Environment for Statistical Computing.
(2008). Deposited 2008.

31.   Dinno, Alexis, dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums.
https://doi.org/10.32614/CRAN.package.dunn.test. Deposited 25 August 2014.

481
482   **Figures and Tables**
483
484

Figure 1: Larvae inhibit adult egg laying without direct contact. (A) 2-chamber
arenas with controlled airflow. Treatments from top to bottom: workers in physical
contact with larvae (red, "larvae contact"), workers without physical contact with larvae
(green, "larvae distance"), workers in physical contact with pupae (blue, "no larvae"). All
treatments were exposed to hexane to ensure comparability between experiments in this
study. **(B)** Number of eggs per ant on the last experimental day. Data points represent
replicate   arenas.   Black: median   ±   IQR.   **: p < 0.01.

Figure 2. Identification of MEHMP as a larva-specific candidate pheromone. (A) GC-MS chromatograms of SPME headspace extracts of different *O. biroi* brood stages (red: larvae). Top insert: 90 *m/z* trace. **(B)** Synthesis of MEHMP from methyl (*Z*)-pent-2-enoate. The reaction conditions for the subsequent 1,4-Michael-addition (using copper(I)-iodide and isopropylmagnesium chloride) and α-hydroxylation (using a Davis-oxidation) are shown and produced the target compound in 55% yield over the two steps. **(C)** Validation of the identification using MS spectra of the volatile produced by larvae (top) and MEHMP (bottom). **(D)** Daily abundance of MEHMP tracks the *O. biroi* colony cycle. Circles: replicate colonies of 100 workers in plaster nests (n = 3); triangles: control, empty plaster nests (n = 2). Dashed line: mean of ant colonies; grey: s.d. Top: transitions between reproductive phases (blue) and brood care phase (red).

506
507

Figure 3: MEHMP inhibits adult egg laying. (A) 2-chamber arenas with controlled airflow. Treatments from top to bottom: workers in physical contact with larvae (red, "larvae contact"), workers exposed to MEHMP (purple), workers without physical contact with larvae (green, "larvae distance"), workers in physical contact with pupae (blue, "no larvae"). All replicates were exposed to hexane to ensure comparability among treatments. **(B)** Number of eggs per ant on the last experimental day. Data points represent replicate arenas. Black: median ± IQR. (***: $p < 0.001$; "NS": non significant.