Evolution of crop phenotypic spaces through domestication

Arthur Wojcik[1,2], Harry Belcram[1], Agnès Rousselet[1], Manon Bouët[2], Marie Brault[2], Pierre

Serin[3], Augustin Desprez[1], Renaud Rincent[1], Andreas Peil[4], Karine Henry[5], Cécile Marchal[6],

Thierry Lacombe[7], Sylvain Glémin[8], Karine Alix[1], Pierre R. Gérard[1], Domenica Manicacci[1],

Yves Vigouroux[3], Catherine Dogimont[2], Maud I. Tenaillon[1]

[1] GQE - Le Moulon, Université Paris-Saclay, INRAE, CNRS, AgroParisTech, Gif-sur-Yvette,

France

[2] GAFL - Génétique et Amélioration des Fruits et Légumes, INRAE, Avignon, France

[3] UMR DIADE, Université de Montpellier, IRD, CIRAD, Montpellier, France

[4] Julius Kühn-Institut, Pillnitz, Germany

[5] Florimond-Desprez, Cappelle-en-Pévèle, France

[6] Grapevine Biological Resources Center, INRAE, Unité Expérimentale Domaine de Vassal,

University of Montpellier, 34340 Marseillan

[7] Institut Agro/Montpellier Supagro - UMR AGAP, Montpellier, France

[8] Laboratoire ECOBIO, UMR 6553 UR1-CNRS, Rennes, France

Corresponding author:

Maud I. Tenaillon, maud.tenaillon@inrae.fr, ORCID 0000-0002-0867-3678

Keywords: Domestication index, Divergent selection, Phenotypic convergence, Life-history

traits, Near infrared spectra, Trait correlations

Summary

- We used domestication as an *in vivo* replicated experiment to investigate how divergent selection has shaped the evolution of multivariate phenotypic spaces.

- We measured 11 to 57 qualitative and quantitative traits in 13 species, either unique or shared between species, and established a framework for cross-species comparisons.

- Our results revealed significant convergence that translated into a cross-species domestication syndrome. Most species exhibited a reduction of the multivariate phenotypic space during domestication. We demonstrated that Near Infrared spectra measured on leaves evolve through processes unrelated with domestication, enabling its use as a control for sampling effects across species. Building on this, we developed a multivariate Phenotypic Divergence Index (mPDI) to rank species by the extent of phenotypic divergence under domestication. We found a high disjunction of wild and domestic phenotypic spaces in all species. Neither the mPDI nor the relative size of wild versus domestic multivariate phenotypic spaces was influenced by the domestication timing or mating system. Lastly, we observed a progressive decoupling of trait correlations with increasing time since domestication.

- In addition to introducing a new index that can be applied across species for comparative purposes, our study uncovers recurring patterns shared among species, pointing to general principles underlying plant domestication.

Introduction

Domestication is a process through which organisms evolve in response to human-altered habitats (Lord et al., 2025). One of the outcomes of this process is phenotypic divergence between the forms, resulting from disruptive selection between wild populations shaped by natural selection in their native habitats and domestic populations shaped by the combined effects of natural selection and adaptation to anthropogenic niches (Glémin & Bataillon, 2009; Ross-Ibarra et al., 2007). From an evolutionary standpoint, plant domestication is a recent process that began during the neolithic revolution (Weisdorf, 2005) with the earliest known domestications occurring ~12,000 years ago in the Fertile Crescent (Barker, 2006). Although debates persist about the intentionality of early selection (Abbo & Gopher, 2020), archaeological evidence from the few crop species for which extensive temporal series are available supports a protracted model of domestication, with the slow fixation of domestic traits over thousands of generations as shown in einkorn wheat, rice and barley (Purugganan & Fuller, 2011). After domestication, crop expanded through trade and seed exchange, leading to the emergence of a genetic structuring into groups, well described in e.g. maize or pearl millet (Burgarella et al., 2018; Tenaillon & Charcosset, 2011).

Over 2500 plants species belonging to 173 botanical families have undergone cultivation or domestication — providing an upper bound for the number of domesticated plants (reviewed in Dirzo & Raven, 2003). Human selection has targeted a diversity of organs (e.g., flower, fruit, seed, tuber) and usages (e.g. food, fiber, ornamentals, medicine). Phenotypic changes that often arise through parallel evolution across species under domestication are collectively referred to as the phenotypic domestication syndrome. These changes include traits that facilitate cultivation, harvesting, and consumption and differ between crop-types e.g., between herbs and woody plants or between vegetatively and sexually propagated crops, for example (Fuller et al., 2023). Domestication traits are well-documented in annual crops. In cereals, for example, they include reduced tillering, loss of seed shattering, and increased ear and seed size (Glémin & Bataillon, 2009; Tenaillon & Manicacci, 2011). More rarely, they involve modifications of life-history traits such as an increase in selfing rate in sunflower, cacao or grapevine (Dempewolf et al., 2012). Compared to annual species, perennial crops are characterized by extended juvenile phases, fewer sexual generations, and a reliance on clonal propagation, which together make them younger domesticates (Gaut et al., 2015). Just like for annual seed crops, domestication traits also point to simplified plant structure, increased edible organs, loss of bitter compounds and reduced sexual reproduction for vegetatively reproduced

crop (Denham et al., 2020; Miller & Gross, 2011). Hence, convergence of domestication traits

across species within-crop types, well documented in cereals (Tenaillon & Manicacci, 2011),

seems to extend to more distant taxa (Meyer et al., 2012) but has yet to be tested.

The phenotypic domestication syndrome is accompanied by a molecular domestication

syndrome (Burban et al., 2022), primarily characterized by a loss of genetic diversity due to

bottlenecks and linked selection at key domestication genes as exemplified in maize (Wright

et al., 2005; Yamasaki et al., 2005) and other crops (reviewed in Gaut et al., 2015). Analysis

of maize landraces suggests that these combined processes have resulted in a loss of genetic

variance at traits targeted by selection (Yang et al., 2019). Thus, one might expect a reduction

of the domestic phenotypic space— that is, a reduction in the range of observable trait variation

expressed by cultivated forms—as divergence increases. This hypothesis requires formal

validation through a comparative analysis across multiple species, based on rigorous sampling

that limits the potential inflating effects of genetic structuring and local adaptation on domestic

phenotypic spaces following agriculture diffusion and modern breeding. Indeed, crop

expansion led to adaptation to novel environments, and selection for local preferences

(Bradshaw, 2016; Meyer & Purugganan, 2013), and thus to a range of new phenotypes, not

directly linked to domestication *per se*.

Another intriguing question arises from the concerted evolution of domestication traits

whose determinism involves complex gene regulatory networks (Studer et al., 2017). Several

studies indicate that domestication has led to a deep rewiring of these networks as shown by

the reorganization of co-expression networks (e.g., in cotton (Rapp et al., 2010), common bean

(Bellucci et al., 2014), and tomato (Sauvage et al., 2017)). Simulations suggest that the

combined effects of demography and selection during domestication result in a redistribution

of genetic correlations, leading to less distinct clusters and a moderate increase of these

correlations (Burban et al., 2022). Studies on maize and teosintes indeed showed an overall

change in the structure of the correlation matrix, but the findings were less clear for

reproductive traits directly affected by domestication, with some analyses indicating

conservation of correlations in the two forms. Contrary to the model predictions, teosintes

generally exhibit more significant and stronger correlations than maize (Yang et al., 2019).

Model predictions therefore need to be further tested across a wider range of species.

In sum, although individual domestication traits have now been described in many

systems, our understanding of the evolution of phenotypic spaces through the domestication

process, and the underlying key evolutionary mechanisms, remains superficial. This is at least

partly due to a lack of unifying statistical framework able to integrate data from different

species, where various and often distinct traits are experimentally measured, including traits of different types (quantitative and qualitative) and scales. The approach we propose here is to build on multivariate analysis to transform measured traits into statistically comparable objects across species; and to use a control trait to account for sampling heterogeneity across species providing a baseline of wild and domestic phenotypic divergence largely independent of domestication. From there, we derived a multivariate Phenotypic Divergence Index (mPDI), which quantifies the degree of phenotypic divergence between wild and domestic forms in each species.

By applying this unifying framework to 13 wild/domesticated pairs (13 species) including a range of domestication dates from ancient to more recent, different mating systems - outcrossers and selfers, and various life cycles from annual to perennial, we addressed the following questions: (1) Is there a significant phenotypic convergence across species? (2) How did phenotypic spaces evolve during domestication? (3) Do domestication dates and mating systems influence the mPDI? (4) Are there consistent patterns in trait correlations throughout the domestication process?

Material and Methods

Plant Material

Our study focused on 13 domestic species and the wild ancestors from which they were derived through domestication: African rice (*Oryza glaberrima*), apple (*Malus domestica*), cabbage (*Brassica oleracea*), common bean (*Phaseolus vulgaris*), eggplant (*Solanum melongena*), einkorn wheat (*Triticum monoccocum*), foxtail millet (*Setaria italica*), grape (*Vitis vinifera*), maize (*Zea mays*), melon (*Cucumis melo*), pearl millet (*Pennisetum glaucum*), sugar beet (*Beta vulgaris*) and tomato (*Solanum lycopersicum*) spanning eight botanical families (Fig. 1a, Table S1). These 13 wild/domestic pairs were chosen because they were all diploids and span a broad timeline of domestication, from the most ancient domestication of einkorn wheat around 9,500 years before present (YBP) to the most recent domestication of eggplant 2000 YBP (Table S1) that were converted in generations (Fig. 2a). Note that for species where two domestication centers were recorded, we focused on a single domestication center and the corresponding wild gene pool: South America for tomato, Mesoamerica for common bean, and Sudan for melon. For beet, we focused on sugar beet whose domestication happened later than for leaf beet (Table S1).

These 13 species furthermore exhibit a diversity of life-cycles from annual, biennial
(e.g., cabbage) to perennial (e.g., grapevine), with sugar beet transitioning from annual to
biennial during domestication (Table S1, Fig. 2a). They also display a range of reproductive
systems from highly self-fertilizing species (selfers) such as foxtail millet to obligate
outcrossers such as apple tree (Table S1). Most species have hermaphroditic flowers, except
for maize which is monoecious, grapevine that transitioned from dioecy to monoecy during
domestication, and melon from monoecy to andromonoecy (Table S1).

To facilitate a comparative analysis across species, we aimed to apply consistent
sampling criteria within each species. For wild/domestic pairs, we chose 17 to 20 wild
populations (accessions) to represent the wild gene pool and 16 to 22 landraces (accessions) to
represent the domestic gene pool (Table S2) except for melon and eggplant for which the wild
sampling encompassed 8 and 13 populations, respectively. For the domestic gene pool, we
focused on landraces in order to capture the domestication process itself while minimizing the
effects of more recent breeding. In addition, we minimized genetic structuring among
individuals to avoid confounding effects of genetic group in phenotypic analyses. To this end,
we relied on established domestication centers for sampling both wild populations and
landraces in nine of the systems (Table S1). Perennials were exceptions in this sampling design,
as we relied on existing and accessible tree collections. For grapevine, whose domestication
center is in South Caucasus, we focused on the western Mediterranean region, where a greater
number of wild and domestic accessions were available. For apple, the domestic sample was
confined to European traditional cultivars, while wild samples originate from Central Asia. For
cabbage, where the exact European domestication center is still debated (Mabry et al., 2021),
we focused on wild and domestic accessions from western Europe (France and United
Kingdom).
176

Plant phenotyping

Except for seed germination rate which required to evaluate many seeds per accession,
plant phenotyping was conducted using a three randomized-block design for all but four species
(melon, cabbage, apple, grape), with each block containing a single individual from each wild
population and each landrace for a total of maximum 120 individuals – 3x40 initial individuals
(Table S3). For melon and cabbage, we measured a single individual per population/landrace
(accession) in a randomized design (i.e., no blocks). For apples, we also evaluated a single
individual per accession in a conservation orchard with trees of different ages. For grapes, one
to five clones from a single individual per accession were phenotyped, and quantitative

variables were averaged across clones – qualitative variables being the same across clones. For a given accession, the clones were planted in a single row, so there were no environmental replicates (i.e, no blocks). Cabbage, grape, apple, einkorn wheat and foxtail millet were phenotyped in the field, while all other species were phenotyped in a greenhouse as they required day-length control for flowering (Table S3). Note that sugar beet was phenotyped in controlled chamber where plant cycles were synchronized to ensure simultaneous flowering of wild (annual) and domestic (biennial) individuals. All specific phenotyping conditions are detailed in Table S3.

The number of traits measured within species varied widely, ranging from 11 traits in apple to 57 traits in sugar beet (Fig. 1 and 3a), summing to 357 different traits across the 13 species (Fig. 3a). The traits are briefly described in Table S4 but a complete ontology can be found at https://doi.org/10.57745/QWEKVK. Traits were evaluated from germination to maturity and harvest. They included both common descriptors across most species, such as plant height or grain size, as well as species-specific traits, such as the acidity of grape must or the spiny or non-spiny leaves of eggplant (Fig. 1 and 3a).

The traits measured fell into three categories (1) continuous quantitative variables, such as stem circumferences, pH and the Brix index (an estimate of the sugar content); (2) discrete quantitative variables, such as the number of internodes or seeds; (3) qualitative variables, such as flower colour or the relative position of styles within the flower. We used imaging software to assess some of the traits including the number of pollen grains in grape flowers and several root traits in sugar beet (Bucksch et al., 2014).

NIRS acquisition and analysis

We acquired Near InfraRed Spectra (NIRS) on leaves for all species except apple. In addition, NIRS were collected on seeds for eight of the 13 species (Tables S4), specifically those for which sufficient seed material was available. In brief, NIR spectroscopy works by directing near-infrared light onto a sample and measuring the amount of light absorbed at different wavelengths, returning spectra whose shape are influenced by the chemical composition of tissues.

For NIRS phenotyping on seeds ($NIRS_{seed}$), we used a Perkin Elmer FT9700 NIR spectrometer. For samples with a limited number of seeds, the light source was directed at the seeds, and the acquisition was performed in complete darkness. To account for sample heterogeneity within species, we performed four technical replicates per sample, mixing the

seeds between each measurement. Each spectrum was obtained as the mean of 32 scans, and wavelengths ranged from 700 to 2500 nm with a step of 8 cm^{-1}, resulting on 1300 wavelengths.

For NIRS phenotyping of leaves, we harvested three leaves or leaf sections per individual. We then oven-dried them for 48h at 40°C in coffee filters as drying does not significantly affect spectra (Borraz-Martínez et al., 2019) and was more convenient for transport. NIRS measurements were taken using a LapSpec 2500 NIR spectrometer. We gathered three spectra per leaf, while carefully avoiding the main veins. Each spectrum was obtained as the mean of 10 scans, and wavelengths ranged from 350 to 2500 nm with a step of 1 nm, resulting on 2150 wavelengths.

We prepared the data as follows, using recommended procedures in Rincent et al. (2018) to control for additive and multiplicative effect of wavelength: (1) we loaded the spectra using the *asdreader* R package (https://github.com/pierreroudier/asdreader); (2) we discarded technical replicates from one individual whose deviation from the individual mean exceeded twice the standard deviation over at least 30% of the spectrum – apparent outlier replicates were also discarded; (3) we further normalized the spectra by sample, and computed their first derivative using a Savitsky-Golay filter (Savitzky & Golay, 1964) as implemented in the *signal* R package (https://cran.r-project.org/web/packages/signal/index.html) using a window size of 37 points and a filter order of two.

Because the wavelengths are numerous and not independent, we computed Principal Coordinate Analyses (PCoA) for leaf and seed samples separately. We used the projections of the individuals on the first two PCoA axes as two new independent traits. We evaluated whether clustering obtained with NIRS effectively grouped the technical replicates of the same individual. In order to do so, we conducted supervised hierarchical clustering of *g* groups, where *g* represents the number of individuals. The grouping of technical replicates was validated using the R package *flexclust* (Leisch, 2006) and the Rand index (Rand, 1971). The Rand index, which measures clustering similarity, ranges from zero to one, with the latter indicating perfect match between the observed and that predicted from individuals.

Domestication syndrome characterization

To describe the domestication syndrome in each species, we applied a trait-by-trait approach. For quantitative traits, we tested differences in mean between forms using a mixed linear model:

$X_{ijk} = \mu + B_i + F_j + A_k + (A \times F)_{jk} + \varepsilon_{ijk}$ (1)

where X_{ijk} represents the phenotypic value of one trait in block i, in form j and accession k; μ

is the phenotypic mean; B_i is the fixed effect of block I; F_j is the fixed effect of form j; A_k is

the random effect of accession k, with $A_k \sim \mathcal{N}(0, \sigma^2_k)$; $(A \times F)_{jk}$ is the random effect of the

interaction between form j and accession k, with $(A \times F)_{jk} \sim \mathcal{N}(0, \sigma^2_j)$, σ^2_j being the specific

variance for form j; ε_{ijk} is the residual error, with $\varepsilon_{ijk} \sim \mathcal{N}(0, \sigma2_E)$.

When there was no replicate, we used a simple linear model:

$X_j = \mu + F_j + \varepsilon_j$ (2)

Where X_j represent the phenotypic value of a trait in form j; μ is the phenotypic mean; F_j is the

fixed effect of form j; ε_j is the residual error, with $\varepsilon_j \sim \mathcal{N}(0, \sigma2_E)$. We used Student's t-tests as

implemented in the *lmertest* R package (Kuznetsova et al., 2017) to assess the significance of

fixed effects. For qualitative traits we used Fisher exact tests to assess significant differences

in the distribution. We calculated species-specific false discovery rate (FDR) q-values using

the Benjamini–Hochberg procedure (Benjamini & Hochberg, 1995) based on the p-values for

the form effect, applying a 5% threshold. We defined domestication-associated traits as those

showing a significant main effect of form, i.e. traits that differed between domestic and wild

forms. For graphical representation of trait values in each species, we constructed radarplot

using the R packages *ggplot2* (Wickham, 2009) and *ggradar* (https://github.com/ricardo-

bion/ggradar).

Domestication syndrome convergence

To test for convergence, we assessed whether species shared more domestication-

associated traits than expected by chance. First, we determined the observed number of

domestication-associated traits unique to a species, or shared by two, or three or more species.

Second, we applied a resampling procedure to generate a null distribution under the hypothesis

of random sharing of domestication-associated traits (i.e., no convergence). To account for the

structure of our data, we resampled traits using a design that mimicked both the number of

traits measured per species and the pattern of trait sharing across species. Of the 357 measured

traits, we identified *n* domestication-associated traits. Our procedure involved performing

10,000 resampling iterations in which *n* traits were randomly drawn from the *t* traits and

designated as domestication-associated traits. For each resampling, we generated a simulated

distribution that we compared to the observed values to compute p-values. Convergence was

supported if the number of traits shared across three or more species exceeded that expected

by chance, as determined by the simulated distribution. Note that because NIRS relies on the

first two axes of a PCoA, with potentially different wavelengths contributing unequally to each
axis across species, these axes are not homologous among species and were excluded from the
convergence analysis.
288

Computation of phenotypic spaces disjunction

We measured the extent of disjunction between the phenotypic spaces of wild and
domestic forms using the Pillai trace (Pillai, 1955) which quantifies how well grouping factors
separate observations in multivariate data. We first performed a Factorial Analysis of Mixed
Data (FAMD) on each dataset; we further considered all resulting axes as new multivariate
variables. Pillai trace was then calculated in a MANOVA using the form (wild or domestic) as
the explanatory variable and the FAMD axes as response variables. The resulting value ranged
from zero, indicating complete overlap of the spaces, to one, indicating complete disjunction
between the two forms.
Sampling heterogeneity between forms (wild and domestic) across species can bias the
estimates of the disjunction between phenotypic spaces. To account for sampling
heterogeneity—referring here to the fact that in some species wild and domesticated material
may have diverged more strongly than in others due to factors such as distance from the
domestication center (drift) or differing levels of crop improvement (selection) for example—
we used $NIRS_{leaf}$ as a control. $NIRS_{leaf}$ is assumed to capture variation primarily unrelated to
domestication. While certain wavelength may be shaped by selection during domestication,
using the full spectrum of wavelengths collectively provides a baseline for processes not
directly associated with domestication.
By computing Pillai trace on the two first axes of PCoA computed on $NIRS_{leaf}$, we were
able to estimate the disjunction unrelated with domestication ($Pillai_{control}$). We then corrected
the disjunction of other traits by subtracting $Pillai_{control}$ to obtain a corrected space disjunction.
In the following, we refer to the resulting index as the multivariate Phenotypic Divergence
Index (mPDI). We also computed the mPDI per unit of disjunction caused by processes
unrelated with domestication, by dividing mPDI by $Pillai_{control}$.
We assessed correlations between mPDI and archaeological estimates of the
domestication timing in generations (Table S1) using Kendall and Spearman correlation tests
and tested the effect of mating system on the mPDI using a Student t-test.
316
317
318

Computation of the of the multivariate phenotypic space

We aimed to compare the relative size of the multivariate phenotypic space between wild and domestic forms across all species pairs. As the determinant of the variance-covariance matrix describing a space is related to its size, we used it as a measurement of the phenotypic space hypervolume. To compute the variance-covariance matrix in each species, we (1) normalized the quantitative traits across all individuals; (2) transformed qualitative traits by decomposing them into their constituent levels (e.g., a trait with five levels became five binary traits), scoring each as present or absent. Considering each form (wild and domestic) within species, we further eliminated traits that were direct linear combinations of others. Finally, we computed the determinant of the variance-covariance matrix of traits for each form.

Since we removed linear combination of traits separately for wild and domestic forms, the sizes of the matrices were not necessarily equal, making direct comparisons of determinants impossible. To address this, we normalized the determinants by taking the r-root, where r is the highest dimension between the wild and domestic matrices. We then calculated the log ratio of the size of the multivariate domesticated phenotypic space to that of the wild multivariate phenotypic space. A ratio greater than one indicated that domesticated forms explore a broader phenotypic space than their wild counterparts, while a ratio smaller than one indicated the reverse. We tested the influence of the number of samples and the number of traits on the hypervolume size by using Kendall and Spearman correlation tests. We used the same tests to assess correlations between the log ratio of the phenotypic spaces and both, domestication timing in generations and mPDI (Table S1).

Because we assumed $NIRS_{leaf}$ primarily captures variation not driven by domestication, we hypothesized that the log ratio of domestic to wild space size based on $NIRS_{leaf}$ to correlate with the corresponding ratio of neutral genomic diversity. To test this hypothesis, we determined wild and domestic multivariate phenotypic spaces on $NIRS_{leaf}$ using all the axes of the corresponding PCoA as variables. We then tested correlations with ratio of genomic diversity taken from literature (Table S5) using Spearman and Kendall correlation tests.

Similarity of trait matrices between wild and domestic forms

For each form, we first calculated the mean value of each trait by averaging across replicates (when multiple replicates were available, Table S3). Second, we computed the pairwise phenotypic correlation matrices between traits using Spearman correlation coefficients and, phenotypic variance-covariance matrices. We used a Student t-test to compare the absolute mean values of correlations. We assessed the similarity of variance-covariance

matrices between forms using the random skewers method (Cheverud & Marroig, 2007), as

implemented in the *EvolQG* R package (Melo et al., 2016). This method is based on Lande's

equation (Lande, 1979), and evaluates how similarly two matrices respond to deformation. In

short, each matrix is multiplied by one random scaled deforming vector, producing two

matrices whose deformation can be described by a response vector. The correlation between

the two response vectors is measured. This process is repeated a large number of times (10,000

in our case), yielding the average response vector correlation. The advantage of the random

skewers method over a Mantel test, is that it provides a quantitative measure of the similarity

between the variance-covariance matrices of wild and domestic traits, enabling comparisons

across species.

We assessed correlations between the similarity of the matrices and archaeological

estimates of domestication timing in generations (Table S1) and mPDI with Kendall and

Spearman correlation tests.

Results

We measured 357 traits across 13 species, 189 of which were species-specific while 40

were shared by 2 to 10 species (Fig. 1a, Table S4). Such a diversity of species and traits poses

a statistical challenge for performing comparisons across species. In addition to trait-by-trait

analysis within species, we therefore developed a unifying statistical framework that includes

the establishment of a multivariate phenotypic divergence index (mPDI) and the use of specific

tools to compare the evolution of phenotypic spaces and trait correlations across species.

Domestication syndrome

We characterized the phenotypic domestication syndrome within species.

Domestication-associated traits were identified based on statistically significant differences in

trait values between wild and domestic forms. For example, in foxtail millet, we measured 15

traits, 12 of which were domestication-associated traits that differentiated wild and domestic

forms (Fig. 4a). Across systems, the percent of domestication-associated traits among all traits

ranged from 26% in melon and 35% in grapevine to 81% in maize and 89.7% in common bean

(Table S4; Figs S1-12).

In addition, as illustrated in foxtail millet for 8 out of 13 domestication-associated traits

(Fig. 4a), the domestic forms tended to display greater values than the wild forms, e.g., larger

leaves and spikes. This pattern was repeatedly observed across species extending to

domestication-associated traits related to plant and organ (leaf/spike/fruit/seed) size (Fig. 4a, Figs S1-S12). We also found a number of qualitative domestication-associated traits such as in common bean, where the colour, pattern and tendency for shattering in pods differed between wild and domestic forms (Fig. S4, FrCol, FrPat and FrTension). Our findings also highlighted new interesting domestication-associated traits, such as the length of flower petals in eggplant or the increase in the number of secondary stems in common bean (Figs S1-S12). We investigated seed/grain properties using NIRS, hypothesizing that domestication may have influenced their composition. Interestingly, we found significant differences in NIRS profiles between wild and domestic forms (either on the first axis of the PCoA – SNIRS1, or the second – SNIRS 2 or both) in melon, einkorn wheat, foxtail millet, cabbage, maize, African rice, common bean and pearl millet (Fig. 1b, Fig. 4b, Figs S1-12).

Convergence of domestication-associated traits

Among 78 species pairs, we found that 82% had at least one shared domestication-associated trait (Fig. 3b). The average number of pairwise shared traits across all species was 2.6 and ranged from 0 – between African rice and eggplant or African rice and grapevine and cabbage, to 7 – between maize and tomato or foxtail millet and maize (Fig. 3b). We formally tested phenotypic convergence for domestication-associated traits. While the number of unique domestication-associated traits to a given species did not significantly differ from expectation, we found a significant enrichment of domestication-associated traits (convergence) shared by three or more species (P-value=0.014, Fig. S13, Fig. S14) indicating convergence. Conversely, this was accompanied by a deficit of traits shared by exactly two species (P-value=0.014, Fig. S13, Fig. S14), consistent with the fixed total number of traits—where enrichment in one category entails a corresponding reduction in another.

Our dataset was strongly biased toward *Poaceae* (5 of the 13 species), a family known for phenotypic convergence of domestication-associated traits (Glémin & Bataillon, 2009). To ensure that observed convergence was not driven by this bias, we repeated the analysis including only one of the *Poaceae* species at a time. The results remained consistent (P-values<0.01) with an enrichment of traits shared by three or more species irrespective of the *Poaceae* species sampled. Altogether, our results indicate convergence in trait selection at a broad evolutionary scale, spanning multiple botanical families, including both monocots and dicots (Fig. 1).

The domestication traits displaying the greatest convergence are leaf size, leaf width and seed area shared respectively by 8, 7 and 9 species (Fig. 1b, LeafLen, LeafWid and SArea).

420   Notably the *Poaceae* family encompassed an important number of shared domestication-
421   associated traits (Fig. 1b), including grain width (SWid), the first axis of the PCoA based on
422   NIRS (SNIRS1) and the number of tillers (TllNo).
423
Index of phenotypic divergence
Previous studies have shown that NIR spectra are able to discriminate between species
(Lang et al., 2017; Meder et al., 2014) or varieties (Barbin et al., 2018), and can be used to
capture genetic similarity between genotypes (Rincent et al., 2018). Reasoning that leaf
chemical composition was less likely targeted by selection during domestication (except
perhaps in leaf crops such as cabbage), we proposed to use NIRS profiling on leaves ($NIRS_{leaf}$)
as a control to estimate the disjunction between wild and domestic forms that is unrelated with
domestication. To verify this assumption, we first verified that $NIRS_{leaf}$ captured a biological
signal. Rand index values ranged from 0.92 in common bean to 0.96 in pearl millet confirming
that $NIRS_{leaf}$ effectively captured phenotypic diversity among individuals, with high
consistency across replicates. Second, we verified that the values of the Pillai trace obtained
from $NIRS_{leaf}$ ($Pillai_{control}$) were much lower than the ones computed on other traits (≥0.71 on
other traits versus ≤0.16 for $NIRS_{leaf}$ excluding apple, Table S5). Consistently, we found no
distinction between wild and domesticated forms along the axes of the PCoA (Fig. 4c) in
contrast with what we obtained with NIRS profiling on seeds (SNIRS1 and SNIRS2) as
illustrated in foxtail millet (Fig. 4a-b) and all other species except apple (Figs S1, S3–S12). In
apple, $Pillai_{control}$ was high (0.70, Table S5) reflecting our sampling of distant Asiatic wild
accessions and European domestic landraces. Finally, we verified that the mPDI values were
not influenced by the number of traits measured in each species (Spearman: P-value= 0.57;
Kendall: P-value=0.56). Based on these results, we treated $NIRS_{leaf}$ as a control trait. We
further used $NIRS_{leaf}$ to calculate a multivariate phenotypic divergence index (mPDI). Because
the mPDI of apple was very close to zero (0.03) and the disjunction of the phenotypic space
was almost entirely due to divergence between forms unrelated to domestication, we decided
to exclude it from the rest of our analysis.
mPDI values ranged from 0.69 for sugar beet to 0.97 for melon with an average of 0.81
(Fig. 2b, Table S5). Note that mPDI values obtained for maize (0.77) were likely
underestimated due to suboptimal greenhouse conditions (Table S3), which hindered data
collection as many ears failed to produce seeds, resulting in poor characterization of the
reproductive organs. All together our results indicate that wild and domestic forms occupy
distinct phenotypic spaces with some variation across species: overlap between phenotypic

spaces was greater in sugar beet, tomato or pearl millet than in grape, common bean or melon (Fig. 2b). We found neither an effect of mating system on mPDI (Wilcoxon-Mann-Whitney: P-value= 0.76), nor a correlation between mPDI and the timing of domestication (Spearman: P-value= 0.64; Kendall: P-value= 0.58). We further tested the effect of mating system on the mPDI per unit of disjunction unrelated with domestication and found a tendency for selfers to be less disjoint than outcrossers (Fig. 2c, Wilcoxon-Mann-Whitney: P-value= 0.15).

Comparative analysis of the multivariate phenotypic spaces

Within each species, we calculated the log ratio of the determinant of the domesticated variance-covariance matrix over that of the wild matrix using all traits to quantify the relative expansion or contraction of multivariate phenotypic diversity in domesticated forms compared to wild forms. We did similarly for the control trait ($NIRS_{leaf}$). We verified that this ratio was neither influenced by the sampling size of the domestic (Spearman, P-value = 0.69; Kendall, P-value = 0.58) and the wild (Spearman, P-value = 0.69; Kendall, P-value = 0.58) forms, nor by the difference between the two (Spearman, P-value = 0.69; Kendall, P-value = 0.58).

Three out of the thirteen pairs —namely einkorn wheat, apple and maize— presented positive log ratio values, indicating a larger phenotypic space in domestic forms (Fig. 5). In most of the pairs (10 out of 13), the domestic forms exhibited smaller phenotypic spaces than their wild counterparts, indicating a tendency for shrinkage during domestication. The log ratio was neither affected by mating system (Wilcoxon-Mann-Whitney, P-value = 0.83), nor by the timing of domestication (Spearman, P-value = 0.69; Kendall, P-value = 0.58). When looking at the absolute values of the log ratio, the pairs covered a range from 0.07 in sugar beet, to 2.2 in grape that displayed the largest shrinkage of phenotypic space.

Finally, we computed the log ratio of phenotypic spaces using size of $NIRS_{leaf}$ and found a positive correlation (Spearman, P-value = 0.04; Kendall, P-value = 0.04, Fig S15) with the ratio of estimates of nucleotide diversity from the literature (Table S5) indicating that the evolution of the phenotypic space of $NIRS_{leaf}$ was mainly driven by genetic drift between wild and domesticated forms, further reinforcing its use as a control trait.

Comparative analysis of trait correlations

Common bean and tomato displayed significantly higher mean absolute correlations between traits than their wild counterparts. Conversely, cabbage, eggplant, einkorn wheat, foxtail millet, grape, melon and sugar beet displayed lower mean absolute correlation values in domestic forms than in wild forms. In the other five wild/domestic pairs — African rice, apple,

maize and pearl millet — we detected no difference in the average absolute correlation values
between domestic and wild form (Table S6).

Random skewers method was used to compute the similarity between domestic and
wild variance-covariance matrices of traits (Fig. 6). Similarity values ranged from 0.36 in
tomato to 0.99 in grape, spanning a continuous gradient. We found no impact of the mating
system (Wilcoxon-Mann-Whitney: P-value=0.09), and matrices similarity did not correlate
with phenotypic disjunction (Spearman: P-value= 0.72; Kendall: P-value= 0.85). However, it
did correlate negatively with domestication timing (Fig. 6; Spearman: P-value= 0.004; Kendall:
P-value= 0.01).

Discussion

Since Darwin (1868), domestication has served as an *in vivo* experiment for studying
phenotypic evolution, offering replicated experiments to explore how divergent selection has
shaped traits over time. However, most studies have focused within species, aiming to identify
domestication traits and their underlying genetic changes to infer mechanisms of adaptation to
human-made environments. Surprisingly, broader patterns of phenotypic shifts and the key
parameters driving their evolution remain poorly understood.

A major limitation has been the absence of a unified experimental and statistical
framework. Domestication traits vary across species, and, in the absence of standardized trait
ontologies, are often assessed using inconsistent methodologies. Additionally, standardizing
sampling across species poses significant challenges. To address these biases, we propose an
experimental design in which wild populations and landraces are sampled near the center of
domestication to minimize genetic structuring and avoid the confounding effects of recent
breeding for the domestic form. We phenotyped these landraces/populations using
standardized measurements across a diverse set of traits, representing both species-specific and
shared traits. As most of the traits were quantitative, we sought to minimize environmental
effects by using common gardens and standardized cultivation conditions (Table S3). Although
these conditions may not always optimal (e.g., for maize), plastic responses to cultivation
conditions is part of the domestication syndrome (Burban et al., 2022). In addition, we
customized a statistical framework to perform cross-species comparisons, and used NIRS
measured on leaves as a control trait to establish a multivariate Phenotypic Divergence Index
(mPDI).

Convergence of domestication-associated traits

The convergence of domestication-associated traits has been well documented in cereals for traits such as seed shattering. It has revealed notable cases of underlying genetic convergence where orthologous genes contribute to variation of the same trait across different species (Patterson et al., 2023). Recently, studies have explored convergence over deep evolutionary timescales in crops from distantly related families, once again highlighting cases of genetic convergence. An example is the *G* gene, which influences seed dormancy variation in rice, tomato, soybean, sorghum and maize (Rendón-Anaya & Herrera-Estrella, 2018). At the phenotypic level, however, meta-analyses of convergence across species struggle with the diversity of experimental designs and disparity of measurement methods. By grouping domestication syndrome traits of 203 crops in nine classes associated to function, Meyer, DuVal, et al., (2012) have hinted convergence in some functions like the production of secondary metabolites (66% of crops). Nevertheless, the statistical significance of these observations was not tested and remains to be established. This necessitates a customized test, as the number of measurable common traits between species decreases with increasing divergence across taxa.

By relying on a specific sampling strategy across species, the use of trait ontology in the MIAPPE format (Papoutsoglou et al., 2020) and a resampling test procedure, we demonstrated that when domestication-associated traits were shared, they tended to be shared by more than two species (Fig. S13). We were thus able to formally test for convergence of domestication-associated traits.

Convergence was found for a diversity of traits. In *Poaceae*, for example, we formally established the convergence observed for reduced tillering, spike and seed size increase (Glémin & Bataillon, 2009). Convergence extended to more distant taxa like common bean or melon as previously reported (Meyer et al., 2012). We showed increases in leaf length and plant height across eight and seven species, respectively (Fig. 1b), consistent with the well-documented trend of gigantism in crops (Milla & Matesanz, 2017). Noteworthy, seeds increased in size in nine species (Fig. 1b, SArea), a consequence of domestication known to extend to most crops. Indeed, seed enlargement during domestication is thought to have arisen initially as an indirect consequence of selection for deeper burial and increased interplant competition (Kluyver et al., 2013, 2017), and linked to the general increase of organ size by allometry (Milla & Matesanz, 2017). Thus, even though seeds are not always the consumed organ, convergence for this trait is found across taxa.

Interestingly, NIRS measured on seeds (SNIRS1, SNIRS2, Fig. 1b) was found to be a

domestication-associated trait for all systems on which it was measured, even on systems where

seeds were likely to have not been selected, e.g., cabbage and apple. For those crops, given that

NIRS is a predictor of grain quality, it implies that the changes observed following

domestication are not only relative to allometry but also to quality for a variety of species, as

previously described by chemical composition analysis in wheat or maize (Fang et al., 2020;

Zeibig et al., 2022). These changes may be related to the (partial) loss of seed dormancy during

domestication, as it is linked to changes in seed chemical composition (Penfield, 2017;

Prudente & Paiva, 2018).

The multivariate phenotypic divergence index

In order to better understand the evolutionary mechanisms driving phenotypic

divergence between wild and domesticated forms, it is essential to quantify this divergence

using a measure that allows cross-species comparisons. A previous attempt proposed a semi-

qualitative index, in which species were ranked according to their level of domestication from

low (1) to high (5) (Dempewolf et al., 2008). Here, we propose a multivariate divergence

quantitative index, the mPDI. Our results suggest that the mPDI is a reliable index. For

instance, the common bean, which is known for its significant phenotypic disjunction (García

et al., 1997; Peña-Valdivia et al., 2011) exhibits high mPDI (Fig. 2b). Conversely, sugar beet

a very recent to domestication displays one of the lowest mPDI value (Fig. 2b). Note that in

apple, mPDI was very close to zero, which can be explained by our sampling in this perennial

species. Indeed, our domestic landraces originated from West Europe, whereas wild

populations were sampled in Central Asia. The Pillai value computed on all traits was thus

almost equal to the value of the $Pillai_{control}$ obtained with $NIRS_{leaf}$, in line with the drift observed

during apple expansion to the west (Cornille et al., 2014).

Overall, we found a high disjunction of phenotypic spaces between wild and

domesticated forms across all species except apple (Pillai $\geq$ 0.69, Fig. 2b). Previous studies

focusing on two traits had proposed rather contradictory hypotheses, suggesting either a

disjunction (Glémin & Bataillon, 2009) or that domesticated forms represented a subset of the

wild phenotypic space (Milla et al., 2015). Our characterization, extended to multivariate

spaces, clearly supports the first hypothesis, confirming an expected outcome of divergent

selection.

Phenotypic divergence is influenced by both genetic drift and selection. We therefore

hypothesized that if it was primarily driven by drift, phenotypic divergence should increase

with the time since domestication. While one of the youngest domesticates, sugar beet, and the oldest domesticate, common bean, were indeed found at the opposite extremes of mPDI values (Fig. 2b), we did not find a significant trend. This suggests that selection dominates the observed pattern. This conclusion is supported by the lack of a relationship between the phenotypic divergence measured in our study and estimates of genomic differentiation between wild and domestic forms, estimated from previous studies (Table S5). This result contrasted to what we obtained from the control trait, for which Pillai values correlated with genomic differentiation (Fig. S15) reinforcing its use as a trait whose evolution was dominated by drift. Note that phenotypic divergence also reflects traits evolvability as measured by standing genetic variance (Opedal et al., 2023). We are thus capturing multiple effects with the mPDI, which are hard to decipher.

Because selfers can adapt faster than outcrossers (Ronfort & Glemin, 2013) potentially accelerating phenotypic divergence, we tested whether the mating system had an effect on the mPDI. We found no effect, and when the mPDI was expressed per unit of disjunction estimated from the control trait, we observed the opposite trend, namely that selfers tended to fall in the lower range of values (Fig. 2c). However, this result should be interpreted with caution due to the binary classification of species as either selfers or outcrossers. In practice, many selfing crops exhibit low levels of outcrossing, which can be sufficient to mix alleles and phenotypes (Morjan & Rieseberg, 2004). Conversely, in outcrossing species, barriers to gene flow between wild and domesticates exist – such as the *GA1* and *GA2* loci that prevent maize pollen from fertilizing teosintes (Chen et al., 2022; Evans & Kermicle, 2001) among other examples. Quantitative variation in the intensity of gene flow, rather than the mating system *per se*, may play a more critical role in determining mPDI rankings, but gene flow remains challenging to quantify.

As a measure of phenotypic divergence between wild and domestic forms, and thus of adaptation to anthropogenetic niches, mPDI could be correlated with the fitness of these forms in human-altered habitats or with their occurrence across environments differently affected by human activity. Following the domestication framework proposed by Lord et al. (2025), species with higher mPDI values are expected to be better adapted to human-mediated environments. Unlike other indices of phenotypic divergence, such as Darwin's (Haldane, 1949), Haldane's (Gingerich, 1993) or Qst (Spitze, 1993), mDPI offers a multivariate quantitative measure that allows cross-species comparisons and can be readily applied beyond domestication contexts—for instance, to assess local adaptation in populations (Fustier et al., 2019) or to study phenotypic evolution in experimental evolution assays (Desbiez-Piat et al.,

622 2023). Note that while easily applicable in distantly related plant taxa, the use of NIRS on
623 specific homologous organ as a control trait may be impractical for animal species.

625 **Evolution of the size of multivariate phenotypic spaces**

626    Domestication has led to a loss of genomic diversity due to both demographic and
627 selective processes (Wright et al., 2005; Yamasaki et al., 2005) that have reduced effective
628 population sizes, thereby decreasing standing variation. One consequence of this loss is a
629 depletion of genetic variance of traits under selection, as shown in maize (Yang et al., 2019), a
630 pattern that extends to traits that are not directly targeted by selection (Rolhauser et al., 2024).
631 Our results support the predicted loss of phenotypic variance, as evidenced by shrinkage of the
632 multivariate phenotypic space in domesticated forms compared to their wild counterparts in 10
633 of the 13 species. However, we found no correlation between this ratio and the corresponding
634 ratio of genomic diversity (Table S6; Spearman P-value=0.79; Kendall P-value=0.86),
635 consistent with studies indicating that domestication affects only a small fraction of the
636 genome—for example, 7.60% in maize (Hufford et al., 2012). Note that we found no effect of
637 the mating system on the ratio of phenotypic diversity between domestic and wild forms,
638 consistent with the hypothesis that selfing does not impair the response to directional selection
639 (Noël et al., 2017).

641 **Reorchestration of correlations between traits**

642    The concept of domestication syndrome not only refers to a series of traits that are
643 different between forms, but also to the correlations between them (Milla et al., 2015). The
644 impact of phenotypic shifts across multiple traits on the evolution of phenotypic spaces is still
645 not fully understood. Phenotypic evolution follows a genetic line of least resistance
646 corresponding to traits with high genetic variance (Mallard, Afonso, et al., 2023; Opedal et al.,
647 2023). The rate of this evolution is determined by the alignment between this line and the
648 direction of selection (Mallard, Afonso, et al., 2023), and therefore on the correlation between
649 traits. Classically, a genetic matrix of traits correlations (G-matrix) is computed. Here we relied
650 on the phenotypic matrix of traits correlations as a proxy for the G-matrix (Cheverud, 1988).

651    We found that traits correlations changed during domestication, although with no clear
652 pattern for increase or decrease across species. Previous studies predicted an increase in genetic
653 correlations between traits in domestic compared to wild forms (Burban et al., 2022), notably
654 because of the loss of genetic diversity in the domestic compartment and pleiotropic effect of
655 genes, e.g. *GL4* in African rice for reduced grain size and non-shattering (Wu et al., 2017) and

TB1 in maize for modification of plant and root architecture and spikelet development (Stitzer

& Ross-Ibarra, 2018). However, traits like seed shattering and grain size have been shown to

evolve concomitantly but at different rate in cereals (Fuller et al., 2012). Domestication have

also led to a decoupling of certain functions, such as root and leaf resource acquisition (Roucou

et al., 2018).

Interestingly, we showed that the more time has passed the more the uncoupling of

variance-covariance matrices, as indicated by the loss of similarity between wild and domestic

matrices (Fig. 6). This suggests that a general hallmark of domestication may involve a

reorganization of trait correlations, a pattern previously identified in different species (various

herbaceous crops, Milla et al., 2014; wheat, Roucou et al., 2018; maize, Yang et al., 2019). As

time passes, the interplay of mutation and selection reshapes the structure of the G-matrix

(Dugand et al., 2021; Houle et al., 2017; Mallard, Noble, et al., 2023), potentially leading to an

alignment to the adaptative landscape (Hohenlohe & Arnold, 2008; Hunt et al., 2007).

Consequently, as time since domestication increases, the G-matrices of wild and domestic

forms may diverge as they become aligned with their respective axes of selection. Altogether,

these observations echoes findings that recombination tends to increase during domestication

(Ross-Ibarra, 2004) or, at minimum, that the recombination landscape is reshaped along the

genome (Bursell et al., 2025). Such novel recombination dynamics can influence phenotypic

variation (Pan et al., 2016; Wagner, 2011) by breaking correlation between traits, potentially

accelerating G-matrix alignment with the direction of selection in the anthropogenic niche.

Conclusion

In this study, we present the first comparative analysis of plant domestication across

multiple species. We introduce a novel phenotypic divergence index (mPDI), applicable across

all domesticated species, providing a standardized framework for cross-species comparisons.

By characterizing the domestication syndrome in 13 species, we demonstrated that near-

infrared spectroscopy (NIRS) measurements on grains contribute to the domestication

syndrome in many species, highlighting the potential of the NIRS technology for both

calculating the mPDI and characterizing domestication-related changes.

Our findings support several key hypotheses of domestication: trait convergence,

disjunction between wild and domestic phenotypes, a relative reduction of the phenotypic space

in domesticated forms, and a reorganization of trait correlations. Importantly, we reveal the

emergence of shared patterns across species, independent of domestication timing or life-history traits, indicating the existence of general principles underlying plant domestication.

Acknowledgments

We are very grateful to Roberto Papa and Elena Bitocchi (Università Politecnica delle Marche, Ancona, Italy) for providing common bean seeds; Alain Label-Richardson (Université Rennes, France), Lorenzo Maggioni (European Cooperative Program for Plant Genetic Resources, Roma, Italy), Roland Von Bothmer (Swedish University of Agricultural Sciences, Alnarp, Sweden), and Charlotte Allender & Nick Fenby (University of Warwick, UK) for providing cabbage seeds; Monika Höfer (Julius Kühn Institute - Federal Research Institute for Cultivated Plants) for managing apple genetic ressource and to Amandine Cornille (NYU Abu Dhabi) for discussing the sampling design of apple trees. We thank the technical staff Virginie Heraudet, Nathalie Galic, Cécilia Zokly, Arnaud Leclerc, Victoria Rozon and Florie Legentil of IDEEV greenhouse for the maintenance of plants. We also thank Yaël Bourgain, Dominique Emmanuel, Laura Courreyan, Martin Simonoviez, Lucie Guerrin, Violette van der Horst and Theophile Thomas for their contribution to data collection. We thank Leila Zekraoui at IRD for her contribution to the discussion on the phenotyping strategy. We thank Martin Ecarnot at INRAE for the access and support at the NIRS plateform ARCAD in Montpellier. We thank Betty Gimenez, Odile Peyron, Sébastien Joannin and Laurent Bouby for the counting of grapevine pollen grains. We thank Alain Ghesquière for the photography of African rice (© IRD). We also deeply thank Mark Chapman and one anonymous reviewer for their insightful comments on the manuscript.

Competing interests

The authors declare no conflicts of interest

Data availability

Phenotypic data and NIR spectra are available on https://doi.org/10.57745/QWEKVK. R scripts are available on the INRAE Forge at https://forge.inrae.fr/gqe-gevad/domisol_phenotypic_spaces. Ontology of measured traits at the MIAPPE format is available at https://doi.org/10.57745/QWEKVK.

References

References

Abbo, S., & Gopher, A. (2020). Plant domestication in the Neolithic Near East: The humans-

plants liaison. *Quaternary Science Reviews*, *242*, 106412.

https://doi.org/10.1016/j.quascirev.2020.106412

Ahmed, H. I., Heuberger, M., Schoen, A., Koo, D.-H., Quiroz-Chavez, J., Adhikari, L.,

Raupp, J., Cauet, S., Rodde, N., Cravero, C., Callot, C., Lazo, G. R., Kathiresan, N.,

Sharma, P. K., Moot, I., Yadav, I. S., Singh, L., Saripalli, G., Rawat, N., … Krattinger,

S. G. (2023). Einkorn genomics sheds light on history of the oldest domesticated

wheat. *Nature*, *620*(7975), 830–838. https://doi.org/10.1038/s41586-023-06389-7

Arnoux, S., Fraïsse, C., & Sauvage, C. (2021). Genomic inference of complex domestication

histories in three Solanaceae species. *Journal of Evolutionary Biology*, *34*(2), 270–

283. https://doi.org/10.1111/jeb.13723

Barbin, D. F., Maciel, L. F., Bazoni, C. H. V., Ribeiro, M. da S., Carvalho, R. D. S., Bispo, E.

da S., Miranda, M. da P. S., & Hirooka, E. Y. (2018). Classification and compositional

characterization of different varieties of cocoa beans by near infrared spectroscopy

and multivariate statistical analyses. *Journal of Food Science and Technology*, *55*(7),

2457–2466. https://doi.org/10.1007/s13197-018-3163-5

Barker, G. (2006). *The Agricultural Revolution in Prehistory: Why did Foragers become*

Farmers? Oxford University Press.

https://doi.org/10.1093/oso/9780199281091.001.0001

Bartsch, D., Lehnen, M., Clegg, J., Pohl-Orf, M., Schuphan, I., & Ellstrand, N. C. (1999).

Impact of gene flow from cultivated beet on genetic diversity of wild sea beet

populations. *Molecular Ecology*, *8*(10), 1733–1741. https://doi.org/10.1046/j.1365-

294x.1999.00769.x

Bellucci, E., Bitocchi, E., Ferrarini, A., Benazzo, A., Biagetti, E., Klie, S., Minio, A., Rau, D.,

Rodriguez, M., Panziera, A., Venturini, L., Attene, G., Albertini, E., Jackson, S. A.,

Nanni, L., Fernie, A. R., Nikoloski, Z., Bertorelle, G., Delledonne, M., & Papa, R.

(2014). Decreased Nucleotide and Expression Diversity and Modified Coexpression

Patterns Characterize Domestication in the Common Bean. *The Plant Cell*, *26*(5),

1901–1912. https://doi.org/10.1105/tpc.114.124040

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society:*

Series B (Methodological), *57*(1), 289–300. https://doi.org/10.1111/j.2517-

6161.1995.tb02031.x

Bhishagratna, K. (1911). *An English translation of The Sushruta Samhita: Based on original*

Sanskrit text (Vol. 2).

Borraz-Martínez, S., Boqué, R., Simó, J., Mestre, M., & Gras, A. (2019). Development of a

methodology to analyze leaves from *Prunus dulcis* varieties using near infrared

spectroscopy. *Talanta*, *204*, 320–328. https://doi.org/10.1016/j.talanta.2019.05.105

Bradshaw, J. E. (2016). Domestication, Dispersion, Selection and Hybridization of Cultivated

Plants. In J. E. Bradshaw (Ed.), *Plant Breeding: Past, Present and Future* (pp. 3–38).

Springer International Publishing. https://doi.org/10.1007/978-3-319-23285-0_1

Bucksch, A., Burridge, J., York, L. M., Das, A., Nord, E., Weitz, J. S., & Lynch, J. P. (2014).

Image-Based High-Throughput Field Phenotyping of Crop Roots. *Plant Physiology*,

166(2), 470–486. https://doi.org/10.1104/pp.114.243519

Burban, E., Tenaillon, M. I., & Le Rouzic, A. (2022). Gene network simulations provide

testable predictions for the molecular domestication syndrome. *Genetics*, *220*(2),

iyab214. https://doi.org/10.1093/genetics/iyab214

Burgarella, C., Cubry, P., Kane, N. A., Varshney, R. K., Mariac, C., Liu, X., Shi, C., Thudi,

M., Couderc, M., Xu, X., Chitikineni, A., Scarcelli, N., Barnaud, A., Rhoné, B.,

Dupuy, C., François, O., Berthouly-Salazar, C., & Vigouroux, Y. (2018). A western

Sahara centre of domestication inferred from pearl millet genomes. *Nature Ecology &*

Evolution, *2*(9), 1377–1380. https://doi.org/10.1038/s41559-018-0643-y

Bursell, M., Rohilla, M., Ramirez, L., Cheng, Y., Schwarzkopf, E. J., Guerrero, R. F., &

Smukowski Heil, C. (2025). Mixed Outcomes in Recombination Rates After

Domestication: Revisiting Theory and Data. *Molecular Ecology*, *n/a*(n/a), e17773.

https://doi.org/10.1111/mec.17773

Cai, C., Bucher, J., Bakker, F. T., & Bonnema, G. (2022). Evidence for two domestication

lineages supporting a middle-eastern origin for Brassica oleracea crops from

diversified kale populations. *Horticulture Research*, *9*, uhac033.

https://doi.org/10.1093/hr/uhac033

Chen, Z., Zhang, Z., Zhang, H., Li, K., Cai, D., Zhao, L., Liu, J., & Chen, H. (2022). A pair of

non-Mendelian genes at the Ga2 locus confer unilateral cross-incompatibility in

maize. *Nature Communications*, *13*(1), 1993. https://doi.org/10.1038/s41467-022-

29729-z

Cheverud, J. M. (1988). A Comparison of Genetic and Phenotypic Correlations. *Evolution*,

42(5), 958–968. https://doi.org/10.1111/j.1558-5646.1988.tb02514.x

Cheverud, J. M., & Marroig, G. (2007). Research Article Comparing covariance matrices:

Random skewers method compared to the common principal components model.

Genetics and Molecular Biology, *30*, 461–469. https://doi.org/10.1590/S1415-

47572007000300027

Choi, J. Y., Zaidem, M., Gutaker, R., Dorph, K., Singh, R. K., & Purugganan, M. D. (2019). The complex geography of domestication of the African rice Oryza glaberrima. *PLOS Genetics*, *15*(3), e1007414. https://doi.org/10.1371/journal.pgen.1007414

Cornille, A., Giraud, T., Smulders, M. J. M., Roldán-Ruiz, I., & Gladieux, P. (2014). The domestication and evolutionary ecology of apples. *Trends in Genetics*, *30*(2), 57–65. https://doi.org/10.1016/j.tig.2013.10.002

Darwin, C. (1868). *The Variation of Animals and Plants Under Domestication* (H. Ritvo, Ed.; Vol. 1). Johns Hopkins University Press. https://doi.org/10.1017/CBO9780511709500

de Bock, T. S. M. (1986). The genus Beta: Domestication, taxonomy and interspecific hybridization for plant breeding. *Acta Horticulturae*, *38*(182), 335–343.

Dempewolf, H., Hodgins, K. A., Rummell, S. E., Ellstrand, N. C., & Rieseberg, L. H. (2012). Reproductive Isolation during Domestication. *The Plant Cell*, *24*(7), 2710–2717. https://doi.org/10.1105/tpc.112.100115

Dempewolf, H., Rieseberg, L. H., & Cronk, Q. C. (2008). Crop domestication in the Compositae: A family-wide trait assessment. *Genetic Resources and Crop Evolution*, *55*(8), 1141–1157. https://doi.org/10.1007/s10722-008-9315-0

Denham, T., Barton, H., Castillo, C., Crowther, A., Dotte-Sarout, E., Florin, S. A., Pritchard, J., Barron, A., Zhang, Y., & Fuller, D. Q. (2020). The domestication syndrome in vegetatively propagated field crops. *Annals of Botany*, *125*(4), 581–597. https://doi.org/10.1093/aob/mcz212

Desbiez-Piat, A., Ressayre, A., Marchadier, E., Noly, A., Remoué, C., Vitte, C., Belcram, H., Bourgais, A., Galic, N., Le Guilloux, M., Tenaillon, M. I., & Dillmann, C. (2023). Pervasive G × E interactions shape adaptive trajectories and the exploration of the phenotypic space in artificial selection experiments. *Genetics*, *225*(4), iyad186. https://doi.org/10.1093/genetics/iyad186

Dirzo, R., & Raven, P. H. (2003). Global State of Biodiversity and Loss. *Annual Review of*

Environment and Resources, *28*(Volume 28, 2003), 137–167.

https://doi.org/10.1146/annurev.energy.28.050302.105532

Duan, N., Bai, Y., Sun, H., Wang, N., Ma, Y., Li, M., Wang, X., Jiao, C., Legall, N., Mao, L.,

Wan, S., Wang, K., He, T., Feng, S., Zhang, Z., Mao, Z., Shen, X., Chen, X., Jiang, Y.,

… Chen, X. (2017). Genome re-sequencing reveals the history of apple and supports a

two-stage model for fruit enlargement. *Nature Communications*, *8*(1), 249.

https://doi.org/10.1038/s41467-017-00336-7

Dugand, R. J., Aguirre, J. D., Hine, E., Blows, M. W., & McGuigan, K. (2021). The

contribution of mutation and selection to multivariate quantitative genetic variance in

an outbred population of Drosophila serrata. *Proceedings of the National Academy of*

Sciences, *118*(31), e2026217118. https://doi.org/10.1073/pnas.2026217118

Dussert, Y., Snirc, A., & Robert, T. (2015). Inference of domestication history and

differentiation between early- and late-flowering varieties in pearl millet. *Molecular*

Ecology, *24*(7), 1387–1402. https://doi.org/10.1111/mec.13119

Evans, M. M. S., & Kermicle, J. L. (2001). Teosinte crossing barrier1, a locus governing

hybridization of teosinte with maize. *Theoretical and Applied Genetics*, *103*(2), 259–

265. https://doi.org/10.1007/s001220100549

Fang, H., Fu, X., Wang, Y., Xu, J., Feng, H., Li, W., Xu, J., Jittham, O., Zhang, X., Zhang, L.,

Yang, N., Xu, G., Wang, M., Li, X., Li, J., Yan, J., & Yang, X. (2020). Genetic basis of

kernel nutritional traits during maize domestication and improvement. *The Plant*

Journal, *101*(2), 278–292. https://doi.org/10.1111/tpj.14539

Fietta, A., Caramelli, D., Schlumbaum, A., & Vernesi, C. (2014). EvolvApple. Studying apple

domestication through the genome scan of ancient seeds. *6th International*

Symposium on Biomolecular Archaeology.

Fuller, D. Q., Asouti, E., & Purugganan, M. D. (2012). Cultivation as slow evolutionary

843   entanglement: Comparative data on rate and sequence of domestication. *Vegetation*

844   *History and Archaeobotany*, *21*(2), 131–145. https://doi.org/10.1007/s00334-011-

845   0329-8

846 Fuller, D. Q., Denham, T., & Allaby, R. (2023). Plant domestication and agricultural

847   ecologies. *Current Biology*, *33*(11), R636–R649.

848   https://doi.org/10.1016/j.cub.2023.04.038

849 Fustier, M.-A., Martínez-Ainsworth, N. E., Aguirre-Liguori, J. A., Venon, A., Corti, H.,

850   Rousselet, A., Dumas, F., Dittberner, H., Camarena, M. G., Grimanelli, D.,

851   Ovaskainen, O., Falque, M., Moreau, L., Meaux, J. de, Montes-Hernández, S.,

852   Eguiarte, L. E., Vigouroux, Y., Manicacci, D., & Tenaillon, M. I. (2019). Common

853   gardens in teosintes reveal the establishment of a syndrome of adaptation to altitude.

854   *PLOS Genetics*, *15*(12), e1008512. https://doi.org/10.1371/journal.pgen.1008512

855 García, E. H., Peña-valdivia, C. B., Rogelio aguirre, J. R., & Muruaga, J. S. M. (1997).

856   Morphological and Agronomic Traits of a Wild Population and an Improved Cultivar

857   of Common Bean (Phaseolus vulgarisL.). *Annals of Botany*, *79*(2), 207–213.

858   https://doi.org/10.1006/anbo.1996.0329

859 Gaut, B. S., Díez, C. M., & Morrell, P. L. (2015). Genomics and the Contrasting Dynamics of

860   Annual and Perennial Domestication. *Trends in Genetics*, *31*(12), 709–719.

861   https://doi.org/10.1016/j.tig.2015.10.002

862 Gingerich, P. D. (1993). Quantification and comparison of evolutionary rates. *American*

863   *Journal of Science*, *293*(A), 453–478.

864 Glémin, S., & Bataillon, T. (2009). A comparative view of the evolution of grasses under

865   domestication. *New Phytologist*, *183*(2), 273–290. https://doi.org/10.1111/j.1469-

866   8137.2009.02884.x

867 Gross, B. L., Henk, A. D., Richards, C. M., Fazio, G., & Volk, G. M. (2014). Genetic

868   diversity in Malus ×domestica (Rosaceae) through time in response to domestication.

869   *American Journal of Botany*, *101*(10), 1770–1779.

870   https://doi.org/10.3732/ajb.1400297

871 Haldane, J. B. S. (1949). Suggestions as to Quantitative Measurement of Rates of Evolution.

872   *Evolution*, *3*(1), 51–56. https://doi.org/10.2307/2405451

873 Hohenlohe, P. A., & Arnold, S. J. (2008). MIPoD: A Hypothesis-Testing Framework for

874   Microevolutionary Inference from Patterns of Divergence. *The American Naturalist*,

875   *171*(3), 366–385. https://doi.org/10.1086/527498

876 Houle, D., Bolstad, G. H., van der Linde, K., & Hansen, T. F. (2017). Mutation predicts 40

877   million years of fly wing evolution. *Nature*, *548*(7668), 447–450.

878   https://doi.org/10.1038/nature23473

879 Hufford, M. B., Xu, X., van Heerwaarden, J., Pyhäjärvi, T., Chia, J.-M., Cartwright, R. A.,

880   Elshire, R. J., Glaubitz, J. C., Guill, K. E., Kaeppler, S. M., Lai, J., Morrell, P. L.,

881   Shannon, L. M., Song, C., Springer, N. M., Swanson-Wagner, R. A., Tiffin, P., Wang,

882   J., Zhang, G., … Ross-Ibarra, J. (2012). Comparative population genomics of maize

883   domestication and improvement. *Nature Genetics*, *44*(7), 808–811.

884   https://doi.org/10.1038/ng.2309

885 Hunt, J., Blows, M. W., Zajitschek, F., Jennions, M. D., & Brooks, R. (2007). Reconciling

886   Strong Stabilizing Selection with the Maintenance of Genetic Variation in a Natural

887   Population of Black Field Crickets (Teleogryllus commodus). *Genetics*, *177*(2), 875–

888   880. https://doi.org/10.1534/genetics.107.077057

889 Kaplan, L. (1981). What is the origin of the common bean? *Economic Botany*, *35*(2), 240–

890   254. https://doi.org/10.1007/BF02858692

Kluyver, T. A., Charles, M., Jones, G., Rees, M., & Osborne, C. P. (2013). Did greater burial

depth increase the seed size of domesticated legumes? *Journal of Experimental*

Botany, *64*(13), 4101–4108. https://doi.org/10.1093/jxb/ert304

Kluyver, T. A., Jones, G., Pujol, B., Bennett, C., Mockford, E. J., Charles, M., Rees, M., &

Osborne, C. P. (2017). Unconscious selection drove seed enlargement in vegetable

crops. *Evolution Letters*, *1*(2), 64–72. https://doi.org/10.1002/evl3.6

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26.

https://doi.org/10.18637/jss.v082.i13

Lande, R. (1979). Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain:

Body Size Allometry. *Evolution*, *33*(1), 402–416. https://doi.org/10.2307/2407630

Lang, C., Almeida, D. R. A., & Costa, F. R. C. (2017). Discrimination of taxonomic identity

at species, genus and family levels using Fourier Transformed Near-Infrared

Spectroscopy (FT-NIR). *Forest Ecology and Management*, *406*, 219–227.

https://doi.org/10.1016/j.foreco.2017.09.003

Leisch, F. (2006). A toolbox for K-centroids cluster analysis. *Computational Statistics &*

Data Analysis, *51*(2), 526–544. https://doi.org/10.1016/j.csda.2005.10.006

Lord, K. A., Larson, G., Allaby, R. G., & Karlsson, E. K. (2025). A universally applicable

definition for domestication. *Proceedings of the National Academy of Sciences*,

122(22), e2413207122. https://doi.org/10.1073/pnas.2413207122

Mabry, M. E., Turner-Hissong, S. D., Gallagher, E. Y., McAlvay, A. C., An, H., Edger, P. P.,

Moore, J. D., Pink, D. A. C., Teakle, G. R., Stevens, C. J., Barker, G., Labate, J.,

Fuller, D. Q., Allaby, R. G., Beissinger, T., Decker, J. E., Gore, M. A., & Pires, J. C.

(2021). The Evolutionary History of Wild, Domesticated, and Feral Brassica oleracea

(Brassicaceae). *Molecular Biology and Evolution*, *38*(10), 4419–4434.

https://doi.org/10.1093/molbev/msab183

Maccaferri, M., Harris, N. S., Twardziok, S. O., Pasam, R. K., Gundlach, H., Spannagl, M.,

Ormanbekova, D., Lux, T., Prade, V. M., Milner, S. G., Himmelbach, A., Mascher, M.,

Bagnaresi, P., Faccioli, P., Cozzi, P., Lauria, M., Lazzari, B., Stella, A., Manconi, A.,

… Cattivelli, L. (2019). Durum wheat genome highlights past domestication

signatures and future improvement targets. *Nature Genetics*, *51*(5), 885–895.

https://doi.org/10.1038/s41588-019-0381-3

Maggioni, L., von Bothmer, R., Poulsen, G., & Härnström Aloisi, K. (2020). Survey and

genetic diversity of wild Brassica oleracea L. germplasm on the Atlantic coast of

France. *Genetic Resources and Crop Evolution*, *67*(7), 1853–1866.

https://doi.org/10.1007/s10722-020-00945-0

Mallard, F., Afonso, B., & Teotónio, H. (2023). Selection and the direction of phenotypic

evolution. *eLife*, *12*, e80993. https://doi.org/10.7554/eLife.80993

Mallard, F., Noble, L., Baer, C. F., & Teotónio, H. (2023). Variation in mutational

(co)variances. *G3 Genes|Genomes|Genetics*, *13*(2), jkac335.

https://doi.org/10.1093/g3journal/jkac335

Manning, K., Pelling, R., Higham, T., Schwenniger, J.-L., & Fuller, D. Q. (2011). 4500-Year

old domesticated pearl millet (*Pennisetum glaucum*) from the Tilemsi Valley, Mali:

New insights into an alternative cereal domestication pathway. *Journal of*

Archaeological Science, *38*(2), 312–322. https://doi.org/10.1016/j.jas.2010.09.007

Marrano, A., Micheletti, D., Lorenzi, S., Neale, D., & Grando, M. S. (2018). Genomic

signatures of different adaptations to environmental stimuli between wild and

cultivated Vitis vinifera L. *Horticulture Research*, *5*, 34.

https://doi.org/10.1038/s41438-018-0041-2

McGovern, P., Jalabadze, M., Batiuk, S., Callahan, M. P., Smith, K. E., Hall, G. R.,

Kvavadze, E., Maghradze, D., Rusishvili, N., Bouby, L., Failla, O., Cola, G., Mariani,

L., Boaretto, E., Bacilieri, R., This, P., Wales, N., & Lordkipanidze, D. (2017). Early

Neolithic wine of Georgia in the South Caucasus. *Proceedings of the National*

Academy of Sciences, *114*(48), E10309–E10318.

https://doi.org/10.1073/pnas.1714728114

Meder, R., Kain, D., Ebdon, N., Macdonell, P., & Brawner, J. T. (2014). Identifying

Hybridisation in Pinus Species Using near Infrared Spectroscopy of Foliage. *Journal*

of Near Infrared Spectroscopy, *22*(5), 337–345.

Melo, D., Garcia, G., Hubbe, A., Assis, A. P., & Marroig, G. (2016). EvolQG - An R package

for evolutionary quantitative genetics. *F1000Research*, *4*, 925.

https://doi.org/10.12688/f1000research.7082.3

Meyer, R. S., DuVal, A. E., & Jensen, H. R. (2012). Patterns and processes in crop

domestication: An historical review and quantitative analysis of 203 global food

crops. *New Phytologist*, *196*(1), 29–48. https://doi.org/10.1111/j.1469-

8137.2012.04253.x

Meyer, R. S., & Purugganan, M. D. (2013). Evolution of crop species: Genetics of

domestication and diversification. *Nature Reviews Genetics*, *14*(12), 840–852.

https://doi.org/10.1038/nrg3605

Milla, R. (2020). Crop Origins and Phylo Food: A database and a phylogenetic tree to

stimulate comparative analyses on the origins of food crops. *Global Ecology and*

Biogeography, *29*(4), 606–614. https://doi.org/10.1111/geb.13057

Milla, R., & Matesanz, S. (2017). Growing larger with domestication: A matter of physiology,

morphology or allocation? *Plant Biology*, *19*(3), 475–483.

https://doi.org/10.1111/plb.12545

Milla, R., Morente-López, J., Alonso-Rodrigo, J. M., Martín-Robles, N., & Stuart Chapin, F.

(2014). Shifts and disruptions in resource-use trait syndromes during the evolution of

herbaceous crops. *Proceedings of the Royal Society B: Biological Sciences*,

281(1793), 20141429. https://doi.org/10.1098/rspb.2014.1429

Milla, R., Osborne, C. P., Turcotte, M. M., & Violle, C. (2015). Plant domestication through

an ecological lens. *Trends in Ecology & Evolution*, *30*(8), 463–469.

https://doi.org/10.1016/j.tree.2015.06.006

Miller, A. J., & Gross, B. L. (2011). From forest to field: Perennial fruit crop domestication.

American Journal of Botany, *98*(9), 1389–1414. https://doi.org/10.3732/ajb.1000522

Morjan, C. L., & Rieseberg, L. H. (2004). How species evolve collectively: Implications of

gene flow and selection for the spread of advantageous alleles. *Molecular Ecology*,

13(6), 1341–1356. https://doi.org/10.1111/j.1365-294X.2004.02164.x

Murray, S. S. (2004). Searching for the origins of African rice domestication. *Antiquity*

Project Gallery78, *300*.

Noël, E., Jarne, P., Glémin, S., MacKenzie, A., Segard, A., Sarda, V., & David, P. (2017).

Experimental Evidence for the Negative Effects of Self-Fertilization on the Adaptive

Potential of Populations. *Current Biology*, *27*(2), 237–242.

https://doi.org/10.1016/j.cub.2016.11.015

Opedal, Ø. H., Armbruster, W. S., Hansen, T. F., Holstad, A., Pélabon, C., Andersson, S.,

Campbell, D. R., Caruso, C. M., Delph, L. F., Eckert, C. G., Lankinen, Å., Walter, G.

M., Ågren, J., & Bolstad, G. H. (2023). Evolvability and trait function predict

phenotypic divergence of plant populations. *Proceedings of the National Academy of*

Sciences, *120*(1), e2203228120. https://doi.org/10.1073/pnas.2203228120

Pan, Q., Li, L., Yang, X., Tong, H., Xu, S., Li, Z., Li, W., Muehlbauer, G. J., Li, J., & Yan, J.

(2016). Genome-wide recombination dynamics are associated with phenotypic

variation in maize. *New Phytologist*, *210*(3), 1083–1094.

https://doi.org/10.1111/nph.13810

Papoutsoglou, E. A., Faria, D., Arend, D., Arnaud, E., Athanasiadis, I. N., Chaves, I.,

Coppens, F., Cornut, G., Costa, B. V., Ćwiek-Kupczyńska, H., Droesbeke, B., Finkers,

R., Gruden, K., Junker, A., King, G. J., Krajewski, P., Lange, M., Laporte, M.-A.,

Michotey, C., … Pommier, C. (2020). Enabling reusability of plant phenomic datasets

with MIAPPE 1.1. *New Phytologist*, *227*(1), 260–273.

https://doi.org/10.1111/nph.16544

Patterson, E. L., Richardson, A., & Bartlett, M. (2023). Pushing the boundaries of organ

identity: Homology of the grass lemma. *American Journal of Botany*, *110*(4), e16161.

https://doi.org/10.1002/ajb2.16161

Peña-Valdivia, C. B., García-Nava, J. R., Aguirre R, J. R., Ybarra-Moncada, Ma. C., & López

H, M. (2011). Variation in Physical and Chemical Characteristics of Common Bean

(Phaseolus vulgaris L.) Grain along a Domestication Gradient. *Chemistry &*

Biodiversity, *8*(12), 2211–2225. https://doi.org/10.1002/cbdv.201100102

Penfield, S. (2017). Seed dormancy and germination. *Current Biology*, *27*(17), R874–R878.

https://doi.org/10.1016/j.cub.2017.05.050

Pillai, K. C. S. (1955). Some New Test Criteria in Multivariate Analysis. *The Annals of*

Mathematical Statistics, *26*(1), 117–121.

Piperno, D. R., & Flannery, K. V. (2001). The earliest archaeological maize (Zea mays L.)

from highland Mexico: New accelerator mass spectrometry dates and their

implications. *Proceedings of the National Academy of Sciences*, *98*(4), 2101–2103.

https://doi.org/10.1073/pnas.98.4.2101

Prudente, D. de O., & Paiva, R. (2018). Seed Dormancy and Germination: Physiological

Considerations. *Journal of Cell and Developmental Biology*, *2*(1).

https://www.imedpub.com/abstract/seed-dormancy-and-germination-physiological-considerations-22078.html

Purugganan, M. D., & Fuller, D. Q. (2011). Archeological data reveal slow rates of evolution during plant domestication. *Evolution*, *65*(1), 171-183. https://doi.org/10.1111/j.1558-5646.2010.01093.x

Rand, W. M. (1971). Objective Criteria for the Evaluation of Clustering Methods. *Journal of the American Statistical Association*, *66*(336), 846–850. https://doi.org/10.2307/2284239

Rapp, R. A., Haigler, C. H., Flagel, L., Hovav, R. H., Udall, J. A., & Wendel, J. F. (2010). Gene expression in developing fibres of Upland cotton (Gossypium hirsutum L.) was massively altered by domestication. *BMC Biology*, *8*(1), 139. https://doi.org/10.1186/1741-7007-8-139

Razifard, H., Ramos, A., Della Valle, A. L., Bodary, C., Goetz, E., Manser, E. J., Li, X., Zhang, L., Visa, S., Tieman, D., van der Knaap, E., & Caicedo, A. L. (2020). Genomic Evidence for Complex Domestication History of the Cultivated Tomato in Latin America. *Molecular Biology and Evolution*, *37*(4), 1118–1132. https://doi.org/10.1093/molbev/msz297

Rendón-Anaya, M., & Herrera-Estrella, A. (2018). The advantage of parallel selection of domestication genes to accelerate crop improvement. *Genome Biology*, *19*(1), 147. https://doi.org/10.1186/s13059-018-1537-7

Riaz, S., De Lorenzis, G., Velasco, D., Koehmstedt, A., Maghradze, D., Bobokashvili, Z., Musayev, M., Zdunic, G., Laucou, V., Andrew Walker, M., Failla, O., Preece, J. E., Aradhya, M., & Arroyo-Garcia, R. (2018). Genetic diversity analysis of cultivated and wild grapevine (Vitis vinifera L.) accessions around the Mediterranean basin and

Central Asia. *BMC Plant Biology*, *18*(1), 137. https://doi.org/10.1186/s12870-018-
1351-0

Rincent, R., Charpentier, J.-P., Faivre-Rampant, P., Paux, E., Le Gouis, J., Bastien, C., &
Segura, V. (2018). Phenomic Selection Is a Low-Cost and High-Throughput Method
Based on Indirect Predictions: Proof of Concept on Wheat and Poplar. *G3
Genes|Genomes|Genetics*, *8*(12), 3961–3972. https://doi.org/10.1534/g3.118.200760

Rolhauser, A. G., Isaac, M. E., Violle, C., Martin, A. R., Vasseur, F., Lemoine, T., Mahaut, L.,
Fort, F., Rotundo, J. L., & Vile, D. (2024). Phenotypic limits of crop diversity: A data
exploration of functional trait space. *New Phytologist*, *244*(2), 708–718.
https://doi.org/10.1111/nph.20050

Ronfort, J., & Glemin, S. (2013). Mating system, Haldane's sieve, and the domestication
process. *Evolution*, *67*(5), 1518–1526. https://doi.org/10.1111/evo.12025

Ross-Ibarra, J. (2004). The Evolution of Recombination under Domestication: A Test of Two
Hypotheses. *The American Naturalist*, *163*(1), 105–112.
https://doi.org/10.1086/380606

Ross-Ibarra, J., Morrell, P. L., & Gaut, B. S. (2007). Plant domestication, a unique
opportunity to identify the genetic basis of adaptation. *Proceedings of the National
Academy of Sciences*, *104*(suppl_1), 8641–8648.
https://doi.org/10.1073/pnas.0700643104

Roucou, A., Violle, C., Fort, F., Roumet, P., Ecarnot, M., & Vile, D. (2018). Shifts in plant
functional strategies over the course of wheat domestication. *Journal of Applied
Ecology*, *55*(1), 25–37. https://doi.org/10.1111/1365-2664.13029

Sauvage, C., Rau, A., Aichholz, C., Chadoeuf, J., Sarah, G., Ruiz, M., Santoni, S., Causse,
M., David, J., & Glémin, S. (2017). Domestication rewired gene expression and

nucleotide diversity patterns in tomato. *The Plant Journal*, *91*(4), 631–645.

https://doi.org/10.1111/tpj.13592

Savitzky, A., & Golay, M. J. E. (1964). Smoothing and Differentiation of Data by Simplified

Least Squares Procedures. *Analytical Chemistry*, *36*, 1627–1639.

https://doi.org/doi.org/10.1021/ac60319a045

Schmutz, J., McClean, P. E., Mamidi, S., Wu, G. A., Cannon, S. B., Grimwood, J., Jenkins, J.,

Shu, S., Song, Q., Chavarro, C., Torres-Torres, M., Geffroy, V., Moghaddam, S. M.,

Gao, D., Abernathy, B., Barry, K., Blair, M., Brick, M. A., Chovatia, M., … Jackson,

S. A. (2014). A reference genome for common bean and genome-wide analysis of dual

domestications. *Nature Genetics*, *46*(7), 707–713. https://doi.org/10.1038/ng.3008

Smith, B. D. (1998). *Emergence of Agriculture*.

Spitze, K. (1993). Population Structure in Daphnia Obtusa: Quantitative Genetic and

Allozymic Variation. *Genetics*, *135*(2), 367–374.

https://doi.org/10.1093/genetics/135.2.367

Stitzer, M. C., & Ross-Ibarra, J. (2018). Maize domestication and gene interaction. *New*

Phytologist, *220*(2), 395–408. https://doi.org/10.1111/nph.15350

Studer, A. J., Wang, H., & Doebley, J. F. (2017). Selection During Maize Domestication

Targeted a Gene Network Controlling Plant and Inflorescence Architecture. *Genetics*,

207(2), 755–765. https://doi.org/10.1534/genetics.117.300071

Tehseen, M. M., Wyatt, N. A., Bolton, M. D., Fugate, K. K., Preister, L. S., Yang, S.,

Ramachandran, V., Li, X., & Chu, C. (2024). Genetic drift, historic migration, and

limited gene flow contributing to the subpopulation divergence in wild sea beet (Beta

vulgaris ssp. Maritima (L.) Arcang). *PLOS ONE*, *19*(9), e0308626.

https://doi.org/10.1371/journal.pone.0308626

Tenaillon, M. I., & Charcosset, A. (2011). A European perspective on maize history. *Comptes*

Rendus. Biologies, *334*(3), 221–228. https://doi.org/10.1016/j.crvi.2010.12.015

Tenaillon, M. I., & Manicacci, D. (2011). Maize origins: An old question under the spotlights.

In *Advances in Maize* (Vol. 3, pp. 89–110). The Society for Experimental Biology.

van Zeist, W., & de Roller, G. J. (1993). Plant remains from Maadi, a predynastic site in

lower Egypt. *Vegetation History and Archaeobotany*, *2*(1), 1–14.

https://doi.org/10.1007/BF00191701

Varshney, R. K., Shi, C., Thudi, M., Mariac, C., Wallace, J., Qi, P., Zhang, H., Zhao, Y.,

Wang, X., Rathore, A., Srivastava, R. K., Chitikineni, A., Fan, G., Bajaj, P., Punnuri,

S., Gupta, S. K., Wang, H., Jiang, Y., Couderc, M., … Xu, X. (2017). Pearl millet

genome sequence provides a resource to improve agronomic traits in arid

environments. *Nature Biotechnology*, *35*(10), 969–976.

https://doi.org/10.1038/nbt.3943

Vavilov, N. (1926). *Studies on the origin of cultivated plants* (Vol. 16).

Vavilov, N. I. (1951). The Origin, Variation, Immunity and Breeding of Cultivated Plants.

Soil Science, *72*(6), 482.

Veltman, M. A., Flowers, J. M., Andel, T. R. van, & Schranz, M. E. (2019). Origins and

geographic diversification of African rice (Oryza glaberrima). *PLOS ONE*, *14*(3),

e0203508. https://doi.org/10.1371/journal.pone.0203508

Wagner, A. (2011). The low cost of recombination in creating novel phenotypes. *BioEssays*,

33(8), 636–646. https://doi.org/10.1002/bies.201100027

Wang, C., Chen, J., Zhi, H., Yang, L., Li, W., Wang, Y., Li, H., Zhao, B., Chen, M., & Diao,

X. (2010). Population genetics of foxtail millet and its wild ancestor. *BMC Genetics*,

11(1), 90. https://doi.org/10.1186/1471-2156-11-90

Weisdorf, J. L. (2005). From Foraging To Farming: Explaining The Neolithic Revolution.

Journal of Economic Surveys, *19*(4), 561–586. https://doi.org/10.1111/j.0950-

0804.2005.00259.x

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer.

https://doi.org/10.1007/978-0-387-98141-3

Wright, S. I., Bi, I. V., Schroeder, S. G., Yamasaki, M., Doebley, J. F., McMullen, M. D., &

Gaut, B. S. (2005). The Effects of Artificial Selection on the Maize Genome. *Science*,

308(5726), 1310–1314. https://doi.org/10.1126/science.1107891

Wu, W., Liu, X., Wang, M., Meyer, R. S., Luo, X., Ndjiondjop, M.-N., Tan, L., Zhang, J., Wu,

J., Cai, H., Sun, C., Wang, X., Wing, R. A., & Zhu, Z. (2017). A single-nucleotide

polymorphism causes smaller grain size and loss of seed shattering during African

rice domestication. *Nature Plants*, *3*(6), 1–7. https://doi.org/10.1038/nplants.2017.64

Yamasaki, M., Tenaillon, M. I., Vroh Bi, I., Schroeder, S. G., Sanchez-Villeda, H., Doebley, J.

F., Gaut, B. S., & McMullen, M. D. (2005). A Large-Scale Screen for Artificial

Selection in Maize Identifies Candidate Agronomic Loci for Domestication and Crop

Improvement. *The Plant Cell*, *17*(11), 2859–2872.

https://doi.org/10.1105/tpc.105.037242

Yang, C. J., Samayoa, L. F., Bradbury, P. J., Olukolu, B. A., Xue, W., York, A. M., Tuholski,

M. R., Wang, W., Daskalska, L. L., Neumeyer, M. A., Sanchez-Gonzalez, J. de J.,

Romay, M. C., Glaubitz, J. C., Sun, Q., Buckler, E. S., Holland, J. B., & Doebley, J. F.

(2019). The genetic architecture of teosinte catalyzed and constrained maize

domestication. *Proceedings of the National Academy of Sciences*, *116*(12), 5643–

5652. https://doi.org/10.1073/pnas.1820997116

Zeibig, F., Kilian, B., & Frei, M. (2022). The grain quality of wheat wild relatives in the

evolutionary context. *Theoretical and Applied Genetics*, *135*(11), 4029–4048.

https://doi.org/10.1007/s00122-021-04013-8

Zhao, G., Lian, Q., Zhang, Z., Fu, Q., He, Y., Ma, S., Ruggieri, V., Monforte, A. J., Wang, P.,

Julca, I., Wang, H., Liu, J., Xu, Y., Wang, R., Ji, J., Xu, Z., Kong, W., Zhong, Y.,

Shang, J., … Huang, S. (2019). A comprehensive genome variation map of melon

identifies multiple domestication events and loci influencing agronomic traits. *Nature*

Genetics, *51*(11), 1607–1615. https://doi.org/10.1038/s41588-019-0522-8

Zhao Z. (2014). The process of origin agriculture in China: Archeological evidence from

flotation results. *Quaternary Sciences*, *34*(1), 73–84.

https://doi.org/10.3969/j.issn.1001-7410.2014.10

Zohary, D., & Hopf, M. (1998). *Domestication of plants in the Old World. The origin and*

spread of cultivated plants in West Asia, Europe and the Nile Valley.

https://doi.org/10.5555/20013014838

1148

Tables and Figures

1150

Figure 1: Domestication-associated traits shared by multiple species. Phylogenetic

relationships of the 13 species (adapted from Milla et al., 2020) (a) and shared domestication-

associated traits in at least three species (b). The length of the tree branches is not to scale. *:

SNIRS1 and SNIRS2 cannot be considered homologous traits among species because NIRS

relies on the first two axes of a PCoA, with potentially different wavelengths contributing

unequally to each axis across species. Trait abbreviations can be found in Table S3.

1157

1158 Figure 2: Domestication timing and multivariate phenotypic divergence index (mPDI) for the

1159 13 species. Domestication timing (Table S1) is indicated in generations (a). Species are ranked

1160 by their mPDI (b) and the mPDI/$Pillai_{control}$ (c).

1161

Figure 3: Convergence in domestication-associated traits. The number of traits measured per species and shared between species is shown (a) along with the subset of domestication-associated traits (b).

Figure 4: Domestication syndrome in foxtail millet. Spider-plot for quantitative traits with significance (*) between wild and domestic forms at an FDR of 5% (a). First two axes of the PCoA computed from $NIRS_{seed}$ (b) and $NIRS_{leaf}$ (c). In (a) the dots represent the mean values, with the shaded area showing the standard error. Abbreviations for traits can be found in Table S3.

Figure 5: Relative size of domestic and wild multivariate phenotypic space. Log ratio of domestic over wild phenotypic space with positive/negative values indicating larger/smaller phenotypic space in domestic than in wild samples.

Figure 6: Correlation between similarity of wild and domestic variance-covariance matrices and domestication timing (Table S1). A matrix similarity of 1 indicates colinear responses to deformation.

Supporting Information

Figure S1: Domestication syndrome in African rice. Spider-plot for quantitative traits with significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of qualitative traits (b). First two axes of the PCoA computed from $NIRS_{seed}$ (c) and $NIRS_{leaf}$ (d). In (a) the dots represent the mean values, with the shaded area showing the standard error. Abbreviation meaning of traits can be found in table S3.

Figure S2: Domestication syndrome of apple. Spider-plot for quantitative traits with significance (*) between wild and domestic forms at an FDR of 5% (a). First two axes of the PCoA computed from $NIRS_{leaf}$ (b). In (a) the dots represent the mean values and the shaded areas denote standard error. Abbreviation meaning of traits can be found in table S3.

Figure S3: Domestication syndrome of cabbage. Spider-plot for quantitative traits with significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of

1195 qualitative traits (b). First two axes of the PCoA computed from $NIRS_{seed}$ (c) and $NIRS_{leaf}$ (d).

1196 In (a), the dots represent the mean values and the shaded areas denote standard error.

1197 Abbreviation meaning of traits can be found in table S3.

1198

1199 Figure S4: Domestication syndrome of common bean. Spider-plot for quantitative traits with

1200 significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of

1201 qualitative traits (b). First two axes of the PCoA computed from $NIRS_{seed}$ (c) and $NIRS_{leaf}$ (d).

1202 In (a), the dots represent the mean values and the shaded areas denote standard error.

1203 Abbreviation meaning of traits can be found in table S3.

1204

1205 Figure S5: Domestication syndrome of eggplant. Spider-plot for quantitative traits with

1206 significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of

1207 qualitative traits (b). First two axes of the PCoA computed from $NIRS_{leaf}$ (c). In (a), the dots

1208 represent the mean values and the shaded areas denote standard error. Abbreviation meaning

1209 of traits can be found in table S3.

1210

1211 Figure S6: Domestication syndrome of einkorn wheat. Spider-plot for quantitative traits with

1212 significance (*) between wild and domestic forms at an FDR of 5% (a). First two axes of the

1213 PCoA computed from $NIRS_{seed}$ (b) and $NIRS_{leaf}$ (c). In (a), the dots represent the mean values

1214 and the shaded areas denote standard error. Abbreviation meaning of traits can be found in

1215 table S3.

1216

1217 Figure S7: Domestication syndrome of melon. Spider-plot for quantitative traits with

1218 significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of

1219 qualitative traits (b). First two axes of the PCoA computed from $NIRS_{seed}$ (c) and $NIRS_{leaf}$ (d).

1220 In (a), the dots represent the mean values and the shaded areas denote standard error.

1221 Abbreviation meaning of traits can be found in table S3.

1222

1223 Figure S8: Domestication syndrome of grapevine. Spider-plot for quantitative traits with

1224 significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of

1225 qualitative traits (b). First two axes of the PCoA computed from $NIRS_{leaf}$ (c). In (a), the dots

1226 represent the mean values and the shaded areas denote standard error. Abbreviation meaning

1227 of traits can be found in table S3.

1228

1229   Figure S9: Domestication syndrome of maize. Spider-plot for quantitative traits with
1230   significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of
1231   qualitative traits (b). First two axes of the PCoA computed from $NIRS_{seed}$ (c) and $NIRS_{leaf}$ (d).
1232   In (a), the dots represent the mean values and the shaded areas denote standard error.
1233   Abbreviation meaning of traits can be found in table S3.

1234

1235   Figure S10: Domestication syndrome of pearl millet. Spider-plot for quantitative traits with
1236   significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of
1237   qualitative traits (b). First two axes of the PCoA computed from $NIRS_{seed}$ (c) and $NIRS_{leaf}$ (d).
1238   In (a), the dots represent the mean values and the shaded areas denote standard error.
1239   Abbreviation meaning of traits can be found in table S3.

1240

1241   Figure S11: Domestication syndrome of sugar beet. Spider-plot for quantitative traits with
1242   significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of
1243   qualitative traits (b). First two axes of the PCoA computed from $NIRS_{leaf}$ (c). In (a), the dots
1244   represent the mean values and the shaded areas denote standard error. Abbreviation meaning
1245   of traits can be found in table S3.

1246

1247   Figure S12: Domestication syndrome of tomato. Spider-plot for quantitative traits with
1248   significance (*) between wild and domestic forms at an FDR of 5% (a). Cleveland plot of
1249   qualitative traits (b). First two axes of the PCoA computed from $NIRS_{leaf}$ (c). In (a), the dots
1250   represent the mean values and the shaded areas denote standard error. Abbreviation meaning
1251   of traits can be found in table S3.

1252

1253   Figure S13: Testing for convergence of domestication-associated traits. Observed number of
1254   domestication traits either unique or shared by two, or three or more species are shown with
1255   the expected number of shared domestication traits as established by a resampling procedure.
1256   Significance was tested by Student t-test (P-values are indicated when significant; * P < 0.05;
1257   ** P <0.01).

1258

1259   Figure S14: Convergence of domestication traits. Pseudo-observed distribution and observed
1260   values for domestication-associated traits in our dataset (blue line), for traits either unique, or
1261   shared by two, or three or more species.

1262

1263   Figure S15: Correlation between the log ratio of domestic to wild NIRS$_{leaf}$ phenotypic space

1264   size with the ratio of domestic to wild genomic diversity (values from the literature, Table S5).

1265

1266   Table S1: Description of the 13 pairs of species with estimate of the domestication timing,

1267   domestication center used in this study, mating system and life cycle.

1268

1269   Table S2: Passport data of sampled accessions.

1270

1271   Table S3: Table S3: Number of traits measured and growing conditions during phenotyping in

1272   each species.

1273

1274   Table S4: List of measured traits for each species, and domestication-associated (DA) traits

1275   with corresponding q-values and percentage of DA-traits in each species.

1276

1277   Table S5: Ratio of domestic (D) over wild (W) multivariate phenotypic space, mDPI,

1278   Pillaicontrol, and estimates from the literature of the ratio (D/W) of genomic diversity and Fst.

1279

1280   Table S6: Average absolute pairwise correlation between traits, computed from all pairwise

1281   correlations in wild and domestic forms. P-values from Student t-tests.

(a)

Poaceae
- African rice
- Einkorn wheat
- Maize
- Foxtail millet
- Pearlmillet

Vitaceae
- Grapevine

Brassicaceae
- Cabbage

Curcurbitaceae
- Melon

Rosaceae
- Apple

Fabaceae
- Common bean

Chenopodiaceae
- Sugar beet

Solanaceae
- Tomato
- Eggplant

(b)

Fruit/Spike: FrCol, FrLen, FrNo, FrW, FrWid

Seeds: SArea, SLen, SWid, tSW, SNIRS1 *, SNIRS2 *

Vegetative part: Height, LeafArea, LeafLen, LeafWid, StemDiam, TilNo

Life cycle
- ○ Annual
- △ Biennial
- □ Perennial

Mating system
- ● Outcrosser
- ● Selfer

Number of measured traits
- 20
- 30
- 40
- 50

Domestication syndrome
- ● Associated
- ● Non-associated

(a)

Apple | Cabbage | Sugarbeet | Melon | Pearlmillet | Tomato | Common bean | Foxtail millet | Einkorn wheat

Present — 2500 — 5000 — 7500

Time since domestication (in generations)

Grapevine | Eggplant | African rice | Maize

(b)

0 — 0.25 — 0.5 — 0.75 — 1

Apple, Sugarbeet, Einkorn wheat, Pearlmillet, Tomato, Maize, Eggplant, African rice, Cabbage, Grapevine, Foxtail millet, Common bean, Melon

(c)

0 — 10 — 20 — 30 — 40

Apple, Tomato, Cabbage, African rice, Eggplant, Einkorn wheat, Common bean, Foxtail millet, Pearlmillet, Grape, Maize, Sugarbeet, Melon

Life cycle
○ Annual   △ Biennial   □ Perennial

Mating system
● Outcrosser   ● Selfer

Foxtail millet