# A protocol for chemical competence in phytopathogenic *Ralstonia*

**Author names**

Tabitha C. Cowell[1] (0009-0002-6055-6978)
Noah R. Guillome[1] (0009-0001-9720-9235)
Matthew L. Cope-Arguello[1] (0000-0002-2860-8179)
Neha N. Prasad[1] (0009-0002-7275-9406)
Tiffany M. Lowe-Power[1] (0000-0003-2681-3563)

**Affiliation**

[1]Department of Plant Pathology, University of California, Davis, Davis, CA, USA 95616

**Corresponding author and email address**

Tiffany M. Lowe-Power, tlowepower@ucdavis.edu

**Keywords**

Ralstonia, chemical competence, transformation, cloning, genetics

## Abstract

Here, we present a protocol for the transformation of *Ralstonia solanacearum* species complex (RSSC) strains by calcium chloride-induced chemical competence. We include a step-by-step protocol with media and solution recipes, quantitative results of the protocol's efficacy for four of five tested RSSC strains, and an explanation of the logic underlying the protocol development. The protocol involves sequentially treating overnight cultures with a cold calcium chloride solution and a cold calcium chloride/magnesium chloride/glycerol solution, transforming through alternating heat shocks and ice incubations, and recovering cells via an outgrowth in rich medium.

## Data summary

Supplemental File S1 is the preliminary version of this chemical competence protocol. Supplemental Figures S1 and S2 show transformation efficiency data for pUFR80 knockout plasmids, and heat shock number and temperature variation, respectively.

## Introduction

The *Ralstonia solanacearum* species complex, hereafter "*Ralstonia*", includes three species of plant pathogenic bacteria, divided into four phylotypes: *R. solanacearum* (phylotype II), *R. pseudosolanacearum* (phylotypes I and III), and *R. syzygii* (phylotype IV) (1). These *Ralstonia* species are in the Burkholderiaceae family and are model organisms for studying bacterial pathogenesis and physiology. Genetic tractability is a key requirement for model organisms. Currently, there are established methods for electroporation, natural transformation, and conjugation for *Ralstonia* (2,3). Here, we developed a protocol for *Ralstonia* transformation using calcium chloride-induced chemical competence. Like natural transformation, chemical competence methods are low-cost and accessible because they do not require specialized equipment and consumables like electroporators or electroporation cuvettes. Here, we present our chemical competence protocol, as well as protocol validation and logic underlying protocol development sections.

## Protocol

### Reagents

#### Media

**Table 1.** CPG rich medium (4).

| Final Concentration | Reagent | Manufacturers' Information |
|---|---|---|
| 1 g/L | Casamino acids | Research Products International, Cat. No. C45000-500.0 |
| 10 g/L | Peptone | Apex BioResearch Products, Cat. No. |

| | | 20-260 |
|---|---|---|
| 5 g/L | Glucose | Sigma-Aldrich, Cat. No. G8270 |
| 1 g/L | Yeast extract | Apex BioResearch Products, Cat. No. 20-254 |

52    For solid media (CPG+TZC agar), add:

| 15 g/L | Agar | Apex BioResearch Products, Cat. No. 20-248 |
|---|---|---|
| 2 mL/L | 1% (w/v) 2,3,5-triphenyl-2H-tetrazolium chloride (TZC) in water | Thermo Scientific, Cat. No. A10870.09 |

Media are sterilized by autoclaving. TZC and antibiotics should be added to molten CPG
agar cooled to ~57°C. Common antibiotics and final concentrations for genetically
engineering *Ralstonia* are kanamycin (25 µg/mL), gentamicin (10 µg/mL), tetracycline (30
µg/mL), and spectinomycin (40 µg/mL). *Ralstonia* isolates often have natural resistance to
the antibiotics chloramphenicol (5), penicillin (5), ampicillin (6), and trimethoprim
(unpublished data, Lowe-Power lab).

**Solutions**
**Table 2.** 100 mM CaCl$_2$.

| Final Concentration | Reagent | Manufacturers' Information |
|---|---|---|
| 100 mM | CaCl$_2$ · 2H$_2$O | Fisher Scientific, Cat. No. C79500 |

62    **Table 3.** Transformation salts with Glycerol (TG salt solution) (7).

| Final Concentration | Reagent | Manufacturers' Information |
|---|---|---|
| 75 mM | CaCl$_2$ · 2H$_2$O | Fisher Scientific, Cat. No. C79500 |
| 6 mM | MgCl$_2$ | Sigma-Aldrich, Cat. No. M8266 |
| 15% (v/v) | Glycerol | Thermo Scientific, Cat. No. A16205.0D |

Solutions should be filter-sterilized using filters with a 0.22 µm pore size and pre-chilled to
4°C prior to use.

**Procedure**

After the written protocol, there are notes and recommendations for steps 5, 6, 9, 12, 14, 19,
and 20.

**Culture Growth**
1. Streak out *Ralstonia* strain on CPG+TZC agar and incubate at 28°C for 2 days or
room temperature (20-22°C) for 3-4 days.

2. Inoculate a single colony into 5 mL CPG broth and incubate overnight in a shaking
incubator at 250 rpm and 28°C.

**Cell Treatment**

3. Perform all steps on ice except for centrifugations. Ensure that materials are pre-
chilled: solutions and microcentrifuge tubes (1.5 mL and 2.0 mL). Set a water bath to
45°C.
4. Incubate the overnight culture on ice for 10 min (make sure the culture tube is slightly
buried in ice so that the full culture volume is chilled).
5. Add 1 mL overnight culture to a chilled 1.5 mL microcentrifuge tube.
6. Centrifuge for 30 s at 13,000 x$g$ and pipette off the supernatant.
7. Resuspend cell pellet in 1 mL cold 100 mM $CaCl_2$ by pipetting to mix.
8. Incubate on ice for 20 min.
9. Centrifuge for 30 s at 13,000 x$g$ and pipette off the supernatant.
10. Resuspend cell pellet in 1 mL cold TG salt solution by pipetting to mix.
11. Incubate on ice for 15 min.
12. Centrifuge for 30 s at 13,000 x$g$ and pipette off the supernatant.
13. Resuspend cell pellet in 100 µL cold TG salt solution by pipetting to mix and keep the
chemically competent cell suspension on ice until use.

**Transformation**

14. Add plasmid DNA and 100 µL of the chemically competent cell suspension to a
chilled 2.0 mL microcentrifuge tube.
15. Incubate on ice for 15 min.
16. Heat shock in a water bath at 45°C for 2 min.
17. Incubate on ice for 15 min.
18. Repeat steps 16-17 twice for a total of three heat shocks and four ice incubations.
19. For the outgrowth, add 500 µL CPG broth and shake the tube horizontally for 4 hours
in a shaking incubator at 250 rpm and 28°C.
20. Plate 100 µL of the outgrowth culture onto prewarmed plates containing CPG+TZC
media with selective antibiotics. Incubate at 28°C for 2-5 days.

**Notes and Recommendations**

● Steps 5-6: More than 1 mL of the overnight culture can be pelleted if the user
requires a greater number of cells, or if the culture is harvested at a low cell density.
● Steps 6, 9, 12: Some cells will be lost after each centrifugation and supernatant
removal, but this is not a problem when working with high density overnight cultures
($A_{600}$ > 2). If cells are centrifuged for longer than 30 s, we recommend using a
temperature-controlled centrifuge set to 4°C to prevent the cells from warming during
centrifugation. We did not yield any transformants after six transformations from a
single attempt to scale up "Cell Treatment" (Steps 3-13) by centrifuging in 50 mL
conical tubes at 5,000 x$g$ for 10 min at 4°C.
● Step 14: When using plasmids that are 8 kb or larger, we recommend using large
amounts of plasmid DNA (≥ 1 µg) for higher rates of successful transformation. We
chose to move cells to 2.0 mL microcentrifuge tubes in Step 14 because these tubes
allow for better circulation of the culture during the outgrowth.

| 115 | • Step 19: The outgrowth can be carried out for as little as 2 hours. Prolonged
| 116 | outgrowths in excess of 5 hours should be avoided so that each colony on the
| 117 | selection plate is derived from an independent genetic event.
| 118 | • Step 20: The outgrowth culture can be diluted before plating, and a 1:10 dilution is
| 119 | usually sufficient. The outgrowth culture can also be concentrated by centrifuging and
| 120 | resuspending the cells in a smaller volume prior to plating.
| 121 |

## Protocol Validation

### Bacterial Strains, Plasmids, and Growth Conditions

We developed the chemical competence protocol using the phylotype I *R. pseudosolanacearum* strain GMI1000. The protocol was also tested on additional strains, including the phylotype II *R. solanacearum* strain IBSBF1503, the phylotype IV *R. syzygii* strain PSI07, and the phylotype III *R. pseudosolanacearum* strains UW386 and CMR15 (**Table 4**). All strains were stored as glycerol stocks at -80°C.

**Table 4.** *Ralstonia* strains used in the development of the chemical competence protocol.

| Strain | Phylotype | Species | Citation |
|---|---|---|---|
| GMI1000 | I | *R. pseudosolanacearum* | (8) |
| IBSBF1503 | II | *R. solanacearum* | (9,10) |
| PSI07 | IV | *R. syzygii* | (11) |
| UW386 | III | *R. pseudosolanacearum* | (12) |
| CMR15 | III | *R. pseudosolanacearum* | (11) |

We developed this protocol using pBBR1-MCS5 (13) and pSW002 (14) (**Table 5**). pBBR1-MCS5 is a 4.8 kb plasmid that can replicate in *Ralstonia* and confers gentamicin resistance. However, we caution against selecting pBBR1 to genetically engineer *Ralstonia solanacearum* species complex strains. We have anecdotally observed that plasmids with pBBR1 origins of replication (including pUFJ10) sometimes "make the strain sick" yielding colonies with an appearance of reduced exopolysaccharide production and elevated production of diffusible melanin. pSW002 is an 8.2 kb plasmid with a pVS1 origin of replication that can replicate in *Ralstonia* and confers tetracycline resistance. To date, we have not observed aberrant colony morphologies with pVS1-ori plasmids in *Ralstonia solanacearum* species complex strains. Plasmids were stored in *Escherichia coli* as glycerol stocks at -80°C.

**Table 5.** Plasmids used in the development of the chemical competence protocol.

| Plasmid | ori-R Type | Size (bp) | Antibiotic Resistance | Replicability | Citation |
|---|---|---|---|---|---|
| pBBR1-MCS5 | pBBR1 | 4,768 | Gentamicin | Replicable in *Ralstonia* | (13) |

| pSW002 | pVS1 | 8,248 | Tetracycline | Replicable in *Ralstonia* | (14) |

Plasmid extractions were performed according to the manufacturer's instructions with the
Zyppy Plasmid Miniprep Kit (Zymo Research, Cat. No. D4037), GeneJET Plasmid Miniprep
Kit (Thermo Scientific, Cat. No. K0503), or I-Blue Mini Plasmid Kit (IBI Scientific, Cat. No.
IB47172), using the kit's elution buffer or water to elute the plasmid DNA. Plasmid DNA
concentration and purity were assessed using a NanoDrop One (Thermo Scientific).

For growing *Ralstonia*, we used CPG media (4) (**Table 1**). Media were sterilized by
autoclaving. Liquid cultures were grown in a shaking incubator at 28°C and 250 rpm.
Cultures on agar plates were grown at 28°C. Concentrations and catalogs for antibiotics are
listed in "Media".

There were two media recipes developed for a preliminary chemical competence protocol
(Supplemental File S1). The first was a modified Super Optimal Broth (SOB) for *Ralstonia*
(ROB), which was composed of 1 g/L casamino acids (Research Products International, Cat.
No. C45000-500.0), 15 g/L peptone (Apex BioResearch Products, Cat. No. 20-260), 1 g/L
yeast extract (Apex BioResearch Products, Cat. No. 20-254), 12.5 mM KCl (EMD, Cat. No.
PX1405-1), 10.0 mM $MgCl_2$ (Sigma-Aldrich, Cat. No. M8266), and 10.0 mM $MgSO_4 \cdot 7H_2O$
(Fisher Scientific, Cat. No. M80-500). The second was a modified Super Optimal broth with
Catabolite Repression (SOC) for *Ralstonia* (ROC), which is ROB with glucose (Sigma-
Aldrich, Cat. No. G8270) added to a final concentration of 20 mM (or 0.4% (w/v)). Although
glucose induces catabolite repression in *E. coli*, a study recently demonstrated that
*Ralstonia* do not have glucose-inducible catabolite repression (15). Both ROB and ROC
were filter-sterilized using a 0.22 µm filter before use.

There were two solutions used in the chemical competence protocol. The first was 100 mM
$CaCl_2$ (**Table 2**). The second was Transformation salts with Glycerol (TG salt solution) (7)
(**Table 3**). Both solutions were filter sterilized using a 0.22 µm filter before use.

For growing *E. coli*, we used LB Broth powder (Apex BioResearch Products, Cat. No. 11-
121; 1.0% tryptone, 0.5% yeast extract, 1.0% sodium chloride), prepared as 25 g/L LB
powder, with 15 g/L agar (Apex BioResearch Products, Cat. No. 20-248) for solid media,
sterilized by autoclaving. Liquid cultures were grown in a shaking incubator at 37°C and 230
rpm. Cultures on agar plates were grown at 37°C. Concentrations and catalogs for
antibiotics are listed in "Media".

For cell density measurements in liquid, we used an Ultrospec 10 Cell Density Meter
(Amersham Biosciences) to measure the absorbance at 600 nm ($A_{600}$), based on the
approximate conversion of an $A_{600}$ value of 1 corresponding to a cell density of $5 \cdot 10^8$
cfu/mL for *Ralstonia*.

**Chemically Competent *Ralstonia***

We used the described protocol to transform the *Ralstonia* strains GMI1000, IBSBF1503,
PSI07, UW386, and CMR15 with the plasmids pBBR1-MCS5 and pSW002. For each strain,
we performed 5 to 19 transformations using the final version of the protocol and calculated
transformation efficiencies (**Figure 1**). For GMI1000 only, we also performed 12 to 28

transformations using the preliminary protocol, with both freshly prepared cells and cell
suspensions stored at -80°C for four weeks (**Figure 1**). We used plasmid amounts ranging
from 45.3 to 745.5 ng in volumes ranging from 0.8 to 15.0 µL.

## Electrocompetent *Ralstonia*

To compare our chemical competence protocol to an existing transformation method in
*Ralstonia*, we determined the transformation efficiency of electroporation of GMI1000 with
the plasmids pBBR1-MCS5 and pSW002. We performed three replicates for each plasmid in
two conditions: fresh cell suspensions and cell suspensions stored at -80°C for four weeks.

### Culture Growth
GMI1000 was struck out from a -80°C glycerol stock onto CPG+TZC agar and incubated at
28°C for two days. A single colony was inoculated into 5 mL of CPG broth and incubated
overnight in a shaking incubator at 28°C and 250 rpm. The $A_{600}$ of the overnight culture was
measured and diluted to an $A_{600}$ of 0.001. Then, two replicate subcultures were prepared,
each with 50 mL CPG broth and 1 mL of the $A_{600}$ 0.001 suspension, and incubated in a
shaking incubator at 28°C and 250 rpm for ~21 hours until the subcultures reached an $A_{600}$
of 0.48 and 0.52, respectively.

### Cell Treatment
The subcultures were placed on wet ice for 10 min, then poured into two chilled 50 mL
conical tubes. The subcultures were centrifuged at 5,000 x*g* for 10 min at 4°C, and the
supernatant was poured off. The first pellet was resuspended in 10 mL of cold (4°C) 10%
(v/v) glycerol by pipetting up and down. This glycerol-cell suspension was used to resuspend
the second pellet, in order to pool the cells into one tube. The cells were pelleted by
centrifugation and the supernatant was poured off. The cell pellet was resuspended in 6 mL
of cold 10% glycerol by pipetting to mix. The cells were centrifuged and the supernatant was
removed by pouring carefully to avoid disturbing the loose pellet. The cell pellet was
resuspended in the residual glycerol by pipetting. Cells were aliquoted in 80 µL portions in
chilled 1.5 mL microcentrifuge tubes on wet ice. At this point, the transformation steps were
carried out for half of the aliquots to acquire the data for fresh cells. The remaining aliquots
were moved to dry ice for 10 min before being moved to a -80°C freezer, where they were
stored for four weeks before performing the transformation steps. These were used to
acquire the data for stored cells.

### Transformation
34.6 ng of pBBR1-MCS5 or 46.2 ng of pSW002 was added to an 80 µL aliquot of
electrocompetent GMI1000 and mixed gently by pipetting, on ice. The mixture was
transferred to a 1 mm electroporation cuvette and shocked in an Electroporator 2510
(Eppendorf) set to 900 V. Researchers in the Lowe-Power lab have had success varying
electroporation from 900 to 1800 V, with anecdotes that higher voltages may increase
electrotransformation efficiency. Immediately after the shock, 1 mL of CPG broth was added
to the electroporation cuvette and mixed gently by pipetting up and down three times. The 1
mL mixture was transferred out of the electroporation cuvette and into a 2 mL
microcentrifuge tube for outgrowth. This tube was shaken horizontally in a shaking incubator
at 28°C and 250 rpm for 4 hours. Volumes (100 µL) of undiluted, 1:10 diluted, 1:100 diluted,
and 1:1000 diluted outgrowth culture were plated onto pre-warmed CPG+TZC+antibiotic

plates, spread using sterile glass beads. The plates were incubated for 3 days at 28°C.
Then, colonies were counted and used to calculate transformation efficiency.

### Calculations

Here, transformation efficiency is defined as the number of transformants per μg of plasmid
DNA. We calculated transformation efficiency (TE) using the following equation:

$$TE = \frac{colony\ count}{\mu g\ plasmid\ DNA} \cdot \frac{\mu L\ outgrowth}{\mu L\ plated}$$

Results were graphed using GraphPad Prism v10.5.0 for Windows.

### Results

Freshly prepared electrocompetent GMI1000 had a median transformation efficiency of
200,000 (range: 150,000-220,000) transformants per μg of plasmid DNA with pBBR1-MCS5,
and 31,000 (range: 29,000-31,000) transformants per μg of plasmid DNA with pSW002
(**Figure 1**). Stored electrocompetent GMI1000 had a median transformation efficiency of
6,600 (range: 2,300-22,000) transformants per μg of plasmid DNA with pBBR1-MCS5, and
2,200 (range: 650-8,700) transformants per μg of plasmid DNA with pSW002 (**Figure 1**).

Chemically competent GMI1000 that was freshly prepared using the preliminary protocol
(Supplemental File S1) had a median transformation efficiency of 270 (range: 0-6,100)
transformants per μg of plasmid DNA with pBBR1-MCS5, and 0 (range: 0-390)
transformants per μg of plasmid DNA with pSW002 (**Figure 1**). Chemically competent
GMI1000 that was stored after preparation using the preliminary protocol had a median
transformation efficiency of 0 (range: 0-130) transformants per μg of plasmid DNA with
pBBR1-MCS5 (**Figure 1**). There were no successful transformations with pSW002.

Chemically competent GMI1000 that was freshly prepared using the final protocol had a
median transformation efficiency of 700 (range: 230-2,300) transformants per μg of plasmid
DNA with pBBR1-MCS5, and 6.9 (range: 0-160) transformants per μg of plasmid DNA with
pSW002 (**Figure 1**). Chemically competent IBSBF1503 that was freshly prepared using the
final protocol had a median transformation efficiency of 70 (range: 28-420) transformants per
μg of plasmid DNA with pBBR1-MCS5 (**Figure 1**). There were no successful IBSBF1503
transformations with pSW002. Chemically competent PSI07 that was freshly prepared using
the final protocol had a median transformation efficiency of 7 (range: 0-35) transformants per
μg of plasmid DNA with pBBR1-MCS5, and 0 (range: 0-3) transformants per μg of plasmid
DNA with pSW002 (**Figure 1**). Chemically competent UW386 that was freshly prepared
using the final protocol had a median transformation efficiency of 125 (range: 14-410)
transformants per μg of plasmid DNA with pBBR1-MCS5, and 10.5 (range: 0-31)
transformants per μg of plasmid DNA with pSW002 (**Figure 1**). For chemically competent
CMR15 that was freshly prepared using the final protocol, there were no successful
transformations with pBBR1-MCS5 or pSW002 (**Figure 1**).

Additionally, we tested this protocol and the preliminary protocol in GMI1000 with ~10 kb,
pUFR80-based plasmids (16) that are not replicable in *Ralstonia*, aiming to perform

markerless gene knockouts through homologous recombination (Supplemental Figure S1).
We used plasmid amounts of 1 to 6.3 µg. Successful transformations had efficiencies of 0.6
to 96.0 transformants per µg of plasmid DNA.

**Figure 1.** Transformation efficiencies for electroporation and chemical competence in selected
*Ralstonia* strains (GMI1000, IBSBF1503, PSI07, UW386, and CMR15) using plasmids (**A**) pBBR1-
MCS5 and (**B**) pSW002. The symbols represent individual transformations with symbol shape
representing strain and symbol color representing the protocol variant. The dashed line indicates the
limit of detection.

278
279

## Logic Underlying Protocol Development

The *Ralstonia* chemical competence protocol is adapted from a chemical competence
protocol for *Pseudomonas aeruginosa* (7). The *P. aeruginosa* protocol treats cells with
divalent cations by washing cells from overnight cultures in a magnesium chloride solution
and then incubating, washing, and resuspending cells in the TG salt solution. The prepared
cells are transformed through a heat shock at 37°C, followed by an outgrowth recovery and
plating on selective media.

Due to a combination of literature review and iterative protocol testing, we adjusted the
concentrations of calcium chloride and magnesium chloride in the wash solutions, incubation
times of wash steps, and the temperature and number of heat shocks (17). For simplicity,
testing was limited to methods that used calcium chloride and/or magnesium chloride as the
inducer of chemical competence, but there are other chemicals (18–20) that have been
shown to induce competence in other bacteria. Future protocol development could test
whether alternative treatments like rubidium chloride (20), sepiolite (20), polyethylene glycol
(18), dimethyl sulfoxide (18), polyamines (25), and combinations thereof are more efficient.

When we tested the use of calcium chloride and magnesium chloride in the first wash step,
we found that calcium chloride alone was more effective than either magnesium chloride
alone or a 1:1 combination of calcium chloride and magnesium chloride. This is consistent
with existing data regarding chemical competence in other bacteria, which indicates that
calcium ions tend to be the most effective cation for chemical competence (17).

It has been previously determined that extended incubation, up to 24 hours, of cells in
calcium chloride improves transformation efficiency in *E. coli* (21). Since both wash solutions
in this protocol contain calcium chloride, we decided to incubate the cells during each wash
step. To keep the protocol within a reasonable time frame, the incubation times were limited
to 20 and 15 min, respectively. However, longer incubation times, such as an overnight
incubation, have the potential to further improve transformation efficiency.

Chemical competence protocols generally require one heat shock step; however, it has been
shown that multiple heat shock/ice incubation cycles can result in increased transformation
efficiency (22). We tested one, two, and three heat shock cycles for the chemical
competence of *Ralstonia*. We also tested heat shock temperatures of 37°C, 42°C, 45°C, and
48°C. Our results indicated that three heat shocks at 42°C, 45°C, and 48°C all had similar
transformation efficiencies; and one, two, and three heat shocks at 48°C all had similar
transformation efficiencies (Supplemental Figure S2).

These adjustments collectively resulted in the final version of our protocol, but we also tested
other factors, which are shown in our preliminary protocol (Supplemental File S1). The
preliminary protocol used magnesium-enriched growth media instead of CPG, cells
subcultured to exponential phase instead of overnight cultures, and took five days instead of
four. The differences in transformation efficiency between the preliminary and final protocols
were small enough that we opted to present the simpler and easier protocol above.

Previous studies have shown that the presence of magnesium ions in the media during
overnight growth increases the transformation efficiency of chemically competent *E. coli*
(23). Super Optimal Broth (SOB) and Super Optimal broth with Catabolite repression (SOC)
both contain magnesium ions and are commonly used in the preparation and outgrowth of
chemically competent *E. coli*, respectively (23). We adapted the SOB and SOC recipes to
develop *Ralstonia*-specific versions, which we named ROB and ROC. The main modification
was replacing the sodium chloride in SOB and SOC with the molar equivalent of potassium
chloride in ROB and ROC because *Ralstonia* is sensitive to sodium chloride (24).
Additionally, SOB and SOC have the same basic components of LB (Lysogeny Broth)
media, which is used to grow *E. coli*. Since *Ralstonia* is grown on CPG, we replaced the LB
components with the corresponding CPG components in the ROB and ROC recipes.
However, we determined that transformation efficiency with ROB/ROC media was similar to
standard CPG media. Thus, the final version of the protocol does not use ROB/ROC, but we

included these recipes because some of the data in Figure 1 use cells grown in ROB/ROC
("Chemically Competent (v1)"; Supplemental File S1).

The physiological state of cells is known to influence transformation efficiency, so we tested
*Ralstonia* from both mid-exponential phase subcultures and stationary phase overnight
cultures. In our electrocompetence protocol for *Ralstonia*, we use cells that were subcultured
and grown to mid-exponential phase. Similarly, when preparing chemically competent *E. coli*
it is common to use subcultured cells grown to mid-exponential phase. Based on this, our
preliminary chemical competence protocol used mid-exponential phase *Ralstonia* cells.
However, our testing showed that the subculturing step did not substantially impact
transformation efficiency, so it was excluded from the final protocol.

## Author statements

### Author contributions

345    TCC: Analysis, Experiments / Investigation, Methodology, Validation, Visualization, Writing –
346    original draft, NRG: Experiments / Investigation, Validation, Writing – review & editing,
347    MLCA: Experiments / Investigation, Validation, Writing – review & editing, NNP: Experiments
348    / Investigation, Validation, Writing – review & editing, TMLP: Conceptualization, Project
349    administration, Resources, Supervision, Writing – review & editing.

### Conflicts of interest

352    The authors declare that there are no conflicts of interest.

### Funding information

355    This work is supported by the joint NSF / USDA NIFA Plant Biotic Interactions program (NSF
356    award # 2336557 and NIFA award # 2024-67013-43303) to T. Lowe-Power, the USDA
357    Hatch Program (Project #1023861) to T. Lowe-Power, the USDA NIFA predoctoral
358    fellowship #2024-67011-42914 to M. Cope-Arguello, a Provost's Undergraduate Fellowship
359    to N. Prasad, and a Provost's Undergraduate Fellowship to T. Cowell.

### Acknowledgements

362    We would like to acknowledge the members of the Lowe-Power lab for valuable discussions.

## References

365    1.    Prior P, Ailloud F, Dalsing BL, Remenant B, Sanchez B, Allen C. Genomic and
366           proteomic evidence supporting the division of the plant pathogen *Ralstonia*
367           *solanacearum* into three species. *BMC Genomics*. 2016;17(1): 90.
368           https://doi.org/10.1186/s12864-016-2413-z.

2.    Bertolla F, Van Gijsegem F, Nesme X, Simonet P. Conditions for natural transformation

of *Ralstonia solanacearum*. *Applied and Environmental Microbiology*. 1997;63(12):
4965–4968. https://doi.org/10.1128/aem.63.12.4965-4968.1997.

3.  Perrier A, Barberis P, Genin S. Introduction of Genetic Material in *Ralstonia*
*solanacearum* Through Natural Transformation and Conjugation. In: Medina C, López-
Baena F (eds) *Host-Pathogen Interactions*. New York, NY: Humana Press; 2018. p.
201–207. https://doi.org/10.1007/978-1-4939-7604-1_16.

4.  Kelman A. The relationship of pathogenicity of *Pseudomonas solanacearum* to colony
appearance in a tetrazolium medium. *Phytopathology*. 1954;44(12): 693–695.

5.  Elphinstone JG, Hennessy J, Wilson JK, Stead DE. Sensitivity of different methods for
the detection of *Ralstonia solanacearum* in potato tuber extracts. *EPPO Bulletin*.
1996;26(3–4): 663–678. https://doi.org/10.1111/j.1365-2338.1996.tb01511.x.

6.  Tans Kersten J, Gay J, Allen C. *Ralstonia solanacearum* AmpD is required for
wild type bacterial wilt virulence. *Molecular Plant Pathology*. 2000;1(3): 179–185.
https://doi.org/10.1046/j.1364-3703.2000.00023.x.

7.  Chuanchuen R, Narasaki CT, Schweizer HP. Benchtop and Microcentrifuge
Preparation of *Pseudomonas aeruginosa* Competent Cells. *BioTechniques*. 2002;33(4):
760–763. https://doi.org/10.2144/02334bm08.

8.  Salanoubat M, Genin S, Artiguenave F, Gouzy J, Mangenot S, Arlat M, et al. Genome
sequence of the plant pathogen *Ralstonia solanacearum*. *Nature*. 2002;415: 497–502.
https://doi.org/10.1038/415497a.

9.  Ailloud F, Lowe T, Cellier G, Roche D, Allen C, Prior P. Comparative genomic analysis
of *Ralstonia solanacearum* reveals candidate genes for host specificity. *BMC*
*Genomics*. 2015;16(270). https://doi.org/10.1186/s12864-015-1474-8.

10.  Ailloud F, Lowe TM, Robène I, Cruveiller S, Allen C, Prior P. *In planta* comparative
transcriptomics of host-adapted strains of *Ralstonia solanacearum*. *PeerJ*. 2016;4:
e1549. https://doi.org/10.7717/peerj.1549.

11.  Remenant B, Coupat-Goutaland B, Guidot A, Cellier G, Wicker E, Allen C, et al.
Genomes of three tomato pathogens within the *Ralstonia solanacearum* species
complex reveal significant evolutionary divergence. *BMC Genomics*. 2010;11: 379.
https://doi.org/10.1186/1471-2164-11-379.

12.  Steidl OR, Truchon AN, Hayes MM, Allen C. Complete Genome Resources for
*Ralstonia* Bacterial Wilt Strains UW763 (Phylotype I); Rs5 and UW700 (Phylotype II);
and UW386, RUN2474, and RUN2279 (Phylotype III). *Molecular Plant-Microbe*
*Interactions*. 2021;34(10): 1212–1215. https://doi.org/10.1094/mpmi-04-21-0086-a.

13.  Kovach ME, Elzer PH, Steven Hill D, Robertson GT, Farris MA, Roop RM, et al. Four
new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different
antibiotic-resistance cassettes. *Gene*. 1995;166(1): 175–176.
https://doi.org/10.1016/0378-1119(95)00584-1.

14.  Wilton R, Ahrendt AJ, Shinde S, Sholto-Douglas DJ, Johnson JL, Brennan MB, et al. A
New Suite of Plasmid Vectors for Fluorescence-Based Imaging of Root Colonizing
Pseudomonads. *Frontiers in Plant Science*. 2018;8: 2242.
https://doi.org/10.3389/fpls.2017.02242.

15.   Baroukh C, Zemouri M, Genin S. Trophic preferences of the pathogen *Ralstonia*
*solanacearum* and consequences on its growth in xylem sap. *MicrobiologyOpen*.
2022;11(1): e1240. https://doi.org/10.1002/mbo3.1240.

16.   Castañeda A, Reddy JD, El-Yacoubi B, Gabriel DW. Mutagenesis of all Eight *avr*
Genes in *Xanthomonas campestris* pv. *campestris* Had No Detected Effect on
Pathogenicity, But One *avr* Gene Affected Race Specificity. *Molecular Plant-Microbe*
*Interactions®*. 2005;18(12): 1306–1317. https://doi.org/10.1094/MPMI-18-1306.

17.   Asif A, Mohsin H, Tanvir R, Rehman Y. Revisiting the Mechanisms Involved in Calcium
Chloride Induced Bacterial Transformation. *Frontiers in Microbiology*. 2017;8: 2169.
https://doi.org/10.3389/fmicb.2017.02169.

18.   Chung CT, Niemela SL, Miller RH. One-step preparation of competent *Escherichia coli*:
transformation and storage of bacterial cells in the same solution. *Proceedings of the*
*National Academy of Sciences*. 1989;86(7): 2172–2175.
https://doi.org/10.1073/pnas.86.7.2172.

19.   Karuna G, Munuswamy N. Induction of competency and comparison of transformation
efficiency in *Escherichia coli* using different salt solutions - Cations. *World Journal of*
*Pharmaceutical Sciences*. 2015;3(4): 732–736.

20.   Ren J, Lee H, Yoo SM, Yu MS, Park H, Na D. Combined chemical and physical
transformation method with RbCl and sepiolite for the transformation of various
bacterial species. *Journal of Microbiological Methods*. 2017;135: 48–51.
https://doi.org/10.1016/j.mimet.2017.02.001.

21.   Dagert M, Ehrlich SD. Prolonged incubation in calcium chloride improves the
competence of *Escherichia coli* cells. *Gene*. 1979;6(1): 23–28.
https://doi.org/10.1016/0378-1119(79)90082-9.

22.   Panja S, Aich P, Jana B, Basu T. How does plasmid DNA penetrate cell membranes in
artificial transformation process of *Escherichia coli*? *Molecular Membrane Biology*.
2008;25(5): 411–422. https://doi.org/10.1080/09687680802187765.

23.   Hanahan D. Studies on transformation of *Escherichia coli* with plasmids. *Journal of*
*Molecular Biology*. 1983;166(4): 557–580. https://doi.org/10.1016/S0022-
2836(83)80284-8.

24.   Jung H, Han G, Lee D, Jung HK, Kim YS, Kong HJ, et al. Understanding the Impact of
Salt Stress on Plant Pathogens Through Phenotypic and Transcriptomic Analysis.
*Plants*. 2025;14(1): 97. https://doi.org/10.3390/plants14010097.

25.   Thomas TJ, Tajmir-Riahi HA, Thomas T. Polyamine–DNA interactions and
development of gene delivery vehicles. *Amino Acids*. 2016;48(10): 2423–2431.
https://doi.org/10.1007/s00726-016-2246-8.

Transformation Efficiency

**A.** pBBR1-MCS, 4.8 kb
**B.** pSW002, 8.2 kb

- ● Electrocompetent phyl. I GMI1000 (Fresh)
- ● Electrocompetent phyl. I GMI1000 (Stored)
- ● Chemically Competent (v1) phyl. I GMI1000 (Fresh)
- ● Chemically Competent (v1) phyl. I GMI1000 (Stored)
- ● Chemically Competent (v2) phyl. I GMI1000 (Fresh)
- ▲ Chemically Competent (v2) phyl. II IBSBF1503 (Fresh)
- ■ Chemically Competent (v2) phyl. IV PSI07 (Fresh)
- ◆ Chemically Competent (v2) phyl. III UW386 (Fresh)
- ▼ Chemically Competent (v2) phyl. III CMR15 (Fresh)