deCYPher: Star Allele-Resolution Computational Framework of Pharmacogenes for

Haplotype-Resolved Long-Read Assemblies

Ting-Yu Chang[1], Yu-Shi Liu[1], Hong-Sheng Lai[2], Tsung-Kai Hung[1], Hsin-Fu Lin[1], Yu-Hui

Lin[3], Chia-Lang Hsu[1,4], Ya-Chien Yang[5,6], Chein-Yu Chen[2], Pei-Lung Chen[1,3,7,8,9], Jacob Shu-

Jui Hsu[1*]

Graduate Institute of Medical Genomics and Proteomics, College of Medicine, National

Taiwan University, Taipei, Taiwan[1]

Department of Biomechatronics Engineering, National Taiwan University, Taipei, Taiwan[2]

Department of Medical Genetics, National Taiwan University Hospital, Taipei, Taiwan[3]

Department of Medical Research, National Taiwan University Hospital, Taipei, Taiwan[4]

Department of Clinical Laboratory Sciences and Medical Biotechnology, College of Medicine,

National Taiwan University, Taipei, Taiwan[5]

Department of Laboratory Medicine, National Taiwan University Hospital, Taipei, Taiwan[6]

Genome and Systems Biology Degree Program, Academia Sinica and National Taiwan

University, Taipei, Taiwan[7]

Graduate Institute of Clinical Medicine, College of Medicine, National Taiwan University,

Taipei, Taiwan[8]

Division of Endocrinology and Metabolism, Department of Internal Medicine, National Taiwan

University Hospital, Taipei, Taiwan[9]

* Corresponding authors

J.S.H: jacobhsu@ntu.edu.tw

Running title

Pharmacogene Allele Profiling for all HPRC Genomes

Abstract

Although existing next-generation sequencing (NGS) tools, such as Aldy and Cyrius, have been applied for allele typing, they cannot achieve complete accuracy due to various genomic challenges including pseudogenes, structural variations, hybrid genes, copy number variations, and gene deletions. These complexities make accurate pharmacogene interpretation more challenging, despite the crucial role pharmacogenomics plays in precision medicine. We developed **deCYPher**, a tool that generates personalized pharmacogenomic reports from haplotype-resolved assemblies. The tool enables analysis of all PharmVar 1A level genes, such as *CYP2B6, CYP2C9, CYP2C19, CYP2D6, CYP3A5, CYP4F2, DPYD, NUDT15,* and *SLCO1B1*. Applied to all HPRC haplotypes (including both release 1 and release 2 data), deCYPher demonstrated high accuracy in resolving complex gene structures. In the case of CYP2D6, release 1 identified 6% gene multiplications, 6% full gene deletions, and 4% CYP2D6/CYP2D7 hybrids. By contrast, release 2 demonstrated an increased prevalence of multiplications (14%) and hybrids (11%), while the frequency of full gene deletions remained comparable at 5%. Comparison with pb-StarPhase revealed discrepancies in 12 of 94 assemblies in the release 1 dataset. For instance, in sample HG02257, Aldy, Cyrius, and deCYPher consistently identified the genotype as *2/*35, whereas pb-StarPhase reported *2/*2. Notably, the *35-defining variants were present in the BAM and VCF files in the pb-StarPhase pipeline, but the local read depth over the *35-specific region was only 5x in HG02257-p, suggesting that the misclassification likely resulted from insufficient coverage - a known limitation of pb-StarPhase under low-depth conditions.

Introduction

Pharmacogenomics investigates the genetic factors influencing drug efficacy and toxicity, with pharmacogenes identified as key contributors to therapeutic outcomes. The Pharmacogene Variation (PharmVar) Consortium curates star-allele definitions and structural-variant annotations to enable reproducible allele calling (Gaedigk et al., 2018), while CPIC and PharmGKB provide evidence-based drug-dosing and reporting guidance that depends on accurate genotype calls (PharmGkb, 2023; Relling & Klein, 2011). PharmVar currently catalogues pharmacogenes such as *CYP1A2, CYP2A6, CYP2A13, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP3A4, CYP3A5, CYP4F2, DPYD, NUDT15, NAT2*, and *SLCO1B1*, offering standardized allele definitions across major drug-metabolizing enzymes and transporters. Concurrently, CPIC has published clinical guidelines for an expanded array of genes—including *ABCG2, CACNA1S, CFTR, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP3A4, CYP3A5, CYP4F2, DPYD, G6PD, HLA-A, HLA-B, HLA-C, IFNL3, IFNL4, MT-RNR1, NUDT15, RYR1, SLCO1B1, TPMT, UGT1A1*, and *VKORC1*. Collectively, these complementary resources establish a molecular basis for the integration of pharmacogenomics into clinical practice.

This tool enables comprehensive analysis of all PharmVar level 1A genes. Within this group, cytochrome P450 (CYP) enzymes play a significant role, as they are involved in phase I oxidative metabolism for many commonly prescribed drugs and contribute to interindividual differences in pharmacokinetics and drug response. (Zanger & Schwab, 2013). Reviews of the CYP superfamily summarize how genetic variation and regulation of expression in these enzymes shape drug metabolism phenotypes and influence efficacy, adverse drug reactions, and drug–drug interactions.

Among CYP and related pharmacogenes, loci such as *CYP2D6*, *CYP2B6, CYP2C9,*

CYP2C19, and others are highly polymorphic. They are well-established predictors of clinically

actionable metabolic phenotypes (for example, poor, intermediate, normal, rapid, or ultrarapid

metabolizers) used in therapeutic decision-making. These genotype–phenotype translations

underpin numerous Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines

and PharmGKB clinical annotations that convert molecular test results into prescribing

recommendations. (Hicks et al., 2015; PharmGkb, 2023; Relling & Klein, 2011).

Accurate star-allele assignment for these genes is complicated by multiple genetic

variation types. In addition to single-nucleotide variants (SNVs) and small indels, clinically

important structural variation occurs frequently — including whole-gene deletions and

duplications, hybrid alleles created by gene–pseudogene conversions, and complex

rearrangements — all of which can have large effects on enzyme activity. *CYP2D6* serves as a

classic example; it is located in a region with related pseudogenes such as *CYP2D7* and features

numerous structural alterations. Approximately 5-10% of people have copy-number variants

like gene deletions or duplications, while *CYP2D6/CYP2D7* hybrid alleles are found in about

2-6%. Altogether, these variations make up more than 10% of clinically significant differences

in *CYP2D6* (Del Tredici et al., 2018; Gaedigk et al., 2017; Qiao et al., 2020).

Conventional short-read sequencing and basic variant-calling pipelines frequently

misidentify structural variants, leading to inaccurate star-allele calls and incorrect downstream

enzyme activity assignments. Consequently, substantial inter-laboratory and inter-algorithm

discrepancies in phenotype assignment often originate from differences in the underlying

genotype calls, particularly when structural variation is present (Del Tredici, Malhotra et al.,

2018; Gaedigk, Ingelman-Sundberg et al., 2018). A host of computational tools and

experimental workflows have been developed to tackle pharmacogene genotyping. Short-read–

based star-allele callers have achieved high concordance for common alleles (Chen et al., 2021;

Hari et al., 2023; Lee et al., 2019; Twesigomwe et al., 2020), but their effectiveness is limited

when complex structural rearrangements or high sequence homology to pseudogenes are

involved. Recent benchmarking studies indicate that sequence ensemble approaches combined

with specialized algorithms perform well. However, complete concordance among all allele

classes—including rare structural variants and hybrids—has not been consistently attained

(Chen et al., 2021; Qiao et al., 2020). Integrating multiple sequencing techniques, including

PacBio HiFi, ONT ultra-long reads, and Illumina Hi-C reads, to construct haplotype-resolved

assemblies has demonstrated the ability to resolve complex loci, detect novel structural variants,

and provide contiguous haplotypes that are essential for unambiguous allele definition (Cheng

et al., 2021; Liao et al., 2023; Nurk et al., 2022).

The pb-StarPhase demonstrates a phase-aware pipeline diploid typing of twenty-one

clinically important pharmacogenes utilizing PacBio HiFi reads, with the ability to correct or

refine historical genotype calls (Song et al., 2023; Wang et al., 2021). Although pb-StarPhase

have advanced allele resolution for complex pharmacogenes, their performance can still be

affected by sequencing coverage and read length, which may lead to incomplete or uncertain

genotype calls under suboptimal data quality. These limitations highlight the need for a targeted

and systematic framework that operates consistently across clinically relevant pharmacogenes,

integrates PharmVar allele definitions with CPIC and PharmGKB conventions, and delivers

high-confidence SNV, CNV, and hybrid allele calls directly from haplotype-resolved genome

assemblies.

We therefore developed **deCYPher**, a computational method for star-allelic resolution of

pharmacogenes listed in CPIC/PharmGKB/PharmVar using haplotype-resolved long-read

assemblies, particularly in genes with complex structural variations. We benchmarked

deCYPher on nine PharmVar 1A-level genes (*CYP2D6, CYP2B6, CYP2C9, CYP2C19,*

CYP3A5, CYP4F2, DPYD, NUDT15, and *SLCO1B1*) and compared its results with those of

Aldy, Cyrius, pb-StarPhase, and TypeAssembly. Compared to existing short-read or long-read

mapping–based approaches, deCYPher leverages fully assembled haplotypes to achieve high

accuracy and unambiguous allele definitions, thereby enhancing the reliability of clinical

pharmacogenomic reporting.

Results

<u>Method Overview</u>

The 47 phased diploid genome assemblies from the Human Pangenome Reference

Consortium (HPRC), were analyzed with our **deCYPher** pipeline (Fig.1). The deCYPher was

designed to achieve the highest resolution for allele assignments defined by the PharmVar

Consortium. The standardized CYP nomenclature — comprising the *CYP* root, family number,

subfamily letter, gene number, and a star allele designation — was used to ensure consistency

with established pharmacogenomic conventions.

For each haplotype-resolved assembly, deCYPher first identified the intervals of target

pharmacogenes and mapped curated reference sequences representing star alleles ("star-one"

references) to the phased contigs. Alignments were performed at full-sequence resolution to

capture both single-nucleotide variants (SNVs) and small indels. Alleles matching known

PharmVar definitions were directly assigned. When variant patterns did not match any existing

alleles, deCYPher flagged them as potential novel alleles, with particular attention to previously

unreported structural variants (SVs) such as novel hybrid genes or deletion genes.

We validated deCYPher-identified structural variants by comparing them to genotypes

from Aldy, Cyrius, and pb-StarPhase. Cross-comparison confirmed the structural variants.

Figure 1. Workflow of the deCYPher pipeline for CYP allele annotation in diploid genome assemblies.

The pipeline integrates the long-read alignment tool minimap2 (Li, 2018)with custom Python and Shell scripts to map allele sequences from the Pharmacogene Variation Consortium (PharmVar) database onto genome assemblies, including the HPRC Year-1 and Year-2 releases. This approach enables accurate identification of CYP gene alleles on phased contigs, achieving up to 95% accuracy in high-resolution allele assignment.

<u>Overview of the HPRC CYP Year-1 and Year-2 releases haplotypes and phenotypes</u>

We fully characterized the high-resolution alleles of CYP genes for all known CYP alleles

in Year-1 47 HPRC human genome assemblies and Year-2 232 HPRC human genome

assemblies (Fig.2A-B). According to the CYP genotype definitions from the Pharmacogene

Variation Consortium (PharmVar; <u>https://www.pharmvar.org/</u>), the CYP2D6 gene includes

185 star alleles, encompassing numerous hybrid and deletion alleles, as well as copy number

variants (CNVs). (Taylor et al., 2020). Our results show that, among the 94 CYP2D6 haplotypes

in the HPRC Year-1 dataset, 6 haplotypes (6.38%, 6/94) harbor one or more CNVs.

Additionally, 6 haplotypes (6.38%, 6/94) were identified as full gene deletions, as illustrated in

Figure 2C. Similarly, in the HPRC Year-2 dataset, 66 out of 464 CYP2D6 haplotypes (14.2%,

66/464) contained one or more CNVs, while 22 haplotypes (4.74%, 22/464) were classified as

full gene deletions, as shown in Figure 2C.

Regarding phenotype predictions, 24 samples (51.06%, 24/47) in the HPRC Year-1

assemblies were classified as Normal Metabolizers, whereas in the HPRC Year-2 assemblies,

121 samples (52.16%, 121/232) fell into this category (detailed data are provided in

Supplemental Tables 1 and 2). These findings address the limitations of previous technologies,

which could not resolve CYP gene structural variations at base-pair resolution across whole-

genome sequences, highlighting the importance of deCYPher for the comprehensive

characterization of pharmacogene genetic variation.

In this study, we denote paternal haplotypes as SampleID-p and maternal haplotypes as

SampleID-m, where SampleID corresponds to the individual sample. For example, HG002-p

and HG002-m represent the paternal and maternal haplotypes of sample HG002, respectively.

A.

B.

C.

Figure 2. Phenotypic profiles of pharmacogenes and structural variants (SVs) of *CYP2D6* identified in the 47 phased assemblies and 232 phased assemblies analyzed by deCYPher.

(A) Heatmap of predicted pharmacogene phenotypes across the HPRC Year-1 cohort.

(B) Heatmap of predicted pharmacogene phenotypes across the HPRC Year-2 cohort.

(C) Structural variant configurations of *CYP2D6*, including full-gene deletions, duplications, and gene–pseudogene hybrid arrangements, as detected in the HPRC Year-1,-2 assemblies.

Within the HPRC assemblies, in addition to *CYP2D6*, we also identified structural

variants in other pharmacogenes. For example, in HG01993-m we detected a *CYP2B6* hybrid

gene, while in HG00423-m we observed a full-gene deletion of *CYP3A5* — a variant that has

never been previously reported. Furthermore, in HG02615-m we identified a CYP4F2 copy

number variation (CNV), which is also novel, and in HG02615-p we found a full-gene deletion

of CYP4F2. Notably, all of these non-*CYP2D6* structural variants were discovered in the HPRC

Year-2 phased assemblies. This suggests that expanding the size of cohorts with PacBio HiFi

phased assemblies may enable the discovery of additional low allele frequency and novel

structural variants, further underscoring the importance of **deCYPher** for structural variant

analysis in pharmacogenes (Supplemental Tables 3 and 4).

<u>Distinctive variant positions defining the CYP2D6*68 hybrid allele</u>

We identified a *CYP2D6* hybrid allele, CYP2D6 *68, in five phased assemblies from the

HPRC dataset: HG00733-p, HG01258-p, HG01784-m, NA20805-p, and NA20827-m.

According to PharmVar, *68 is defined by a CYP2D7-to-CYP2D6 gene conversion spanning

exons 2 to 9. By contrast, *36 is characterized by a conversion limited to exon 9, and *82 by a

conversion restricted to exon 2. Consequently, any haplotype carrying the exon 2–9 conversion

of *68 will inherently meet the defining criteria of both *36 (exon 9 conversion) and *82 (exon

2 conversion). Based on this relationship, our initial strategy was to classify any assembly

carrying both *36 and *82 as *68 (Fig. 3A). Using this approach, we detected three *68 alleles

in the HPRC Year-1 phased assemblies—one in HG00733-p and two in HG01258-p—yielding

results that were fully concordant with calls from Aldy and Cyrius.

Although carrying both *36 and *82 as a *68 criterion successfully identified *68, it may

not capture all instances of this hybrid allele, as *68 involves a continuous exon 2–9 conversion.

To refine detection, we sought variant positions that could serve as a more complete signature

for *68. We first compiled all star alleles present across confirmed *68 assemblies and found

consistent co-occurrence of the following alleles: *10, *36, *39, *49, *53, *74, *82, *83, *86,

*88 (with one variant site optionally absent), *131, *132, *139, and *176. We then recorded

the variant positions corresponding to these alleles, excluding NG_008376.4:6032 T>C, to

generate a representative *68 variant profile (VCF) (Fig. 3B and Supplemental Table 5).

Figure 3. Detection and variant profile of CYP2D6*68.

CYP2D6*68 is characterized by a continuous CYP2D7-to-CYP2D6 conversion

spanning exons 2–9, in contrast to *36 (exon 9 only) and *82 (exon 2 only).

Assemblies harboring both *36 and *82 were classified as *68. Variant positions

defining the *68 hybrid allele across exons 2–9 are shown, with each marker indicating a characteristic nucleotide change.

<u>Fusion of *CYP2B6*</u>

Previous studies have shown that the **CYP2B6*29** allele shares complete sequence

identity with exons 1–4 of ***CYP2B7*** and exons 5–9 of ***CYP2B6***, and is associated with reduced

enzymatic activity. In the PharmVar database, CYP2B6*29 is annotated as a hybrid gene

between *CYP2B7* and *CYP2B6*. Furthermore, the GenBank database contains FASTA

sequences representing the *CYP2B7/CYP2B6* breakpoint junction, as documented in earlier

reports (Rotger *et al.* 2007; Martis *et al.* 2013; Gaedigk *et al.* 2018).

Although the genomic characterization of **CYP2B6*29** is already well documented,

current genotyping tools such as **Cyrius**, **Aldy**, and **pb-StarPhase** perform suboptimally in

detecting this allele. In both the HPRC Year-1 and Year-2 phased assemblies, no samples were

classified as CYP2B6*29 by these tools. However, when applying **deCYPher** to the HPRC

Year-2 phased assemblies, we identified **HG01993-m** as carrying the CYP2B6*29 allele.

deCYPher recognized this allele based on two key observations. First, we detected

abnormal mapping patterns when aligning the HG01993-m contigs to the **CYP2B6*1** reference:

read mapping with generally poor mapping quality, except for the primary mapping segment

spanning **exons 5–9**, where mapping quality was higher, whereas **exons 1–4** were

predominantly soft-clipped (Fig. 4A). This mapping profile is consistent with the known

structure of CYP2B6*29. Second, when aligning HG01993-m to the *CYP2B7/CYP2B6*

breakpoint junction sequence, we observed almost perfect alignment across the junction with

no evidence of soft or hard clipping, and only a few single-nucleotide variants (SNVs) (Fig.

4B). These Integrative Genomics Viewer(IGV)(Robinson et al., 2011)-based visualizations

provide orthogonal evidence that deCYPher correctly identified HG01993-m as CYP2B6*29.

Existing algorithms missed this call. This demonstrates deCYPher's unique ability to resolve

hybrid alleles in pharmacogenes that are often misclassified or overlooked by short-read–based

tools.

> **Figure 4. IGV visualization of CYP2B6*29 in HG01993-m.**
>
> (A) IGV alignment view of the HG01993-m contigs against the CYP2B6*1 reference. The region corresponding to exons 1–4 displayed extensive soft clipping and poor mapping quality, whereas alignment improved across exons 5–9, consistent with the CYP2B6*29 hybrid structure.

<u>Comparisons with pb-StarPhase in HPRC genomes</u>

pb-StarPhase (Wang & et al., 2021) provides an innovative approach for pharmacogene

genotyping using PacBio long-read whole-genome sequencing. To directly compare its

performance with deCYPher, we applied the recommended pb-StarPhase pipeline to the 47

HPRC Year-1 samples, starting from randomly selected long-read raw data. We then generated

diplotype calls for all samples and compared them against deCYPher results.

During this comparison, we observed that pb-StarPhase frequently produced

"NO_MATCH" calls. For the purpose of evaluation, such calls were treated as discordances

and provisionally considered errors on the pb-StarPhase side. For other discordant cases, we

examined upstream evidence (i.e., BAM and VCF files) to adjudicate correctness. Specifically,

if a critical variant was clearly present in the alignments and variant calls but missing from the

output, we classified the call as incorrect. Conversely, if a critical variant was clearly present

in the alignments and variant calls and also present in the output, we classified the call as correct.

In cases where both tools produced plausible but differing outputs supported by partial evidence,

we labeled the discrepancy as "Unknown", reflecting unresolved classification.

As an illustrative example, in HG03516, both the pbmm2-aligned bam file and the

Deepvariant vcf file contained the variant chr22:42129770 G>A, a defining variant of

CYP2D6*17 (Fig. 5). Nevertheless, pb-StarPhase assigned this sample as CYP2D6*2/*2,

missing the variant, whereas deCYPher correctly identified the diplotype as CYP2D6*17/*2.

Overall, deCYPher demonstrated superior accuracy for *CYP2D6*, achieving a precision of

94.68%, compared to 87.23% for pb-StarPhase — the largest discrepancy observed among the

nine pharmacogenes analyzed. Notably, both tools produced identical results for *CYP2C9* and

NUDT15 (Table 1).

These findings highlight the limitations of pb-StarPhase when allele-defining, while

underscoring the advantage of deCYPher in leveraging haplotype-resolved assemblies to

deliver unambiguous star-allele assignments. This accuracy is particularly critical for complex

loci such as *CYP2D6*, where reliable genotype-to-phenotype translation is essential for clinical

pharmacogenomics.

Figure 5. IGV visualization of the CYP2D*17-defining variant in HG03516.

This track shows clear evidence of the chr22:42129770 G>A substitution. Despite this, pb-StarPhase assigned the sample as CYP2D6*2/*2, whereas deCYPher correctly identified it as CYP2D6*17/*2.

Table 1. Comparison between deCYPher and pb-StarPhase.

	Total Differences	deCYPher Correct	Unknown	deCYPher Precision	pbStarPhase Precision
CYP2B6	6	6/6(100%)	0(0%)	94/94(100%)	84/94(89.36%)
CYP2C19	3	1/3(33.3%)	2/3(66.7%)	93/94(98.94%)	89/94(94.68%)
CYP2D6	12	7/12(58.3%)	5/12(41.7%)	89/94(94.68%)	82/94(87.23%)
CYP3A5	4	2/4(50%)	2/4(50%)	92/94(97.87%)	90/94(95.74%)
CYP4F2	2	2/2(100%)	0(0%)	94/94(100%)	92/94(97.87%)
DPYD	3	3/3(100%)	0(0%)	94/94(100%)	91/94(96.81%)
SLCO1B1	2	1/2(50%)	1/2(50%)	93/94(98.94%)	92/94(97.87%)
CYP2C9	0			94/94(100%)	94/94(100%)
NUDT15	0			94/94(100%)	94/94(100%)

Discussion

In this study, we developed and applied **deCYPher**, a computational framework for star

allele typing of pharmacogenes using haplotype-resolved assemblies. By leveraging long-read

contiguity and phasing, deCYPher enables accurate assignment of PharmVar-defined alleles,

detection of structural variants, and standardized diplotype-to-phenotype translation consistent

with CPIC conventions. Compared to existing genotyping tools such as Aldy, Cyrius, and pb-

StarPhase(Chen et al., 2021; Hari et al., 2023; Wang et al., 2021), deCYPher demonstrated

improved accuracy, particularly in loci with complex structural variation.

Hybrid alleles of *CYP2D6*, such as *36 and *82, have been well characterized in previous

studies (Gaedigk et al., 2010; Gaedigk et al., 2018). In line with these reports, we identified

CYP2D6*68, a complex hybrid allele defined in PharmVar as a gene conversion event spanning

exons 2–9 from *CYP2D7* (Gaedigk et al., 2018). Importantly, our analysis also revealed

structural variants outside of *CYP2D6*, including a CYP2B6*29 hybrid in HG01993-m, which,

to our knowledge, has not been reported in other tools before. These findings highlight the

strength of deCYPher in resolving both known and unclear structural variants across

pharmacogenes, particularly when complex scenarios involving CNVs, hybrid genes, or large

deletions complicate haplotype assignment.

These findings highlight a broader limitation of short-read pipelines. Previous work has

shown that genotype-to-phenotype assignments can differ substantially across laboratories and

algorithms, especially in the presence of structural variants(Twesigomwe et al., 2020). In such

cases, ambiguity often arises not because phenotype rules differ, but because the underlying

genotype calls are inconsistent. By integrating haplotype-resolved assemblies with PharmVar

allele definitions, deCYPher substantially reduces this discordance and produces more reliable

haplotype calls.

Our benchmarking against HPRC assemblies further supports this advantage. While pb-

StarPhase achieved a precision of 8% for *CYP2D6* calls, deCYPher achieved 94.7%,

representing the largest performance gap across the nine genes evaluated. In two genes

(*CYP2C9* and *NUDT15*), both tools produced identical results, indicating that the primary

benefit of deCYPher emerges in genes with extensive structural complexity. These results are

consistent with prior observations that structural variation is a major source of error in

pharmacogene genotyping (Twesigomwe et al., 2020; Zhou et al., 2017).

Finally, our work parallels advances in other structurally complex genomic loci, such as

the KIR region. The SKIRT pipeline has demonstrated how leveraging high-quality assemblies

enables the discovery of novel gene fusions and CNVs in KIR genes (Hung et al., 2024; Zhou

et al., 2024). By analogy, deCYPher provides a foundation for high-resolution pharmacogene

genotyping, paving the way for future studies in larger HiFi-based cohorts and the systematic

discovery of rare or novel pharmacogene SVs.

Methods

deCYPher - Algorithms of the Pharmacogenes annotation Pipeline

Read Mapping and Variant Detection

The deCYPher pipeline was developed to resolve PharmVar-defined star alleles and structural variation, including gene duplications and deletions. To capture multiple polymorphic alleles at each locus, we aligned PharmVar reference allele sequences to haplotype-resolved genome assemblies using **minimap2** (Li, 2018) optimized for PacBio HiFi data. Parameter settings and alignment configurations are provided in **Supplementary Note 2**.

To validate structural variants identified by deCYPher, we compared the results with genotypes obtained from established tools, including Aldy, Cyrius, and pb-StarPhase. The input data consisted of short-read Illumina whole-genome sequencing (WGS) from the 1KGP and HPRC cohorts, as well as long-read PacBio HiFi sequencing. After cross-comparing these results, we evaluated and confirmed the presence of structural variants.

Output and Visualization

deCYPher reports haplotype-level star allele calls in a structured JSON format, providing a standardized and machine-readable representation of nine PharmVar/CPIC-prioritized pharmacogenes. This format encodes PharmVar-defined star alleles together with copy number information and annotations of putative novel structural variants, thereby supporting downstream analysis, integration into pharmacogenomic pipelines, and clinical interpretation. Further details regarding the output structure and visualization are provided in **Supplementary Note 1**.

Data access

All data generated or analyzed during this study are included in this article, its Supplemental files, and Zenodo (https://doi.org/10.5281/zenodo.8094803, https://zenodo.org/records/14854401). deCYPher source code is available on GitHub (https://github.com/HolidayChang/deCYPher).

Competing interest statement

The authors declare no competing interests.

Acknowledgment

This work was supported in part by computational and storage resources provided by the National Center for High-performance Computing (NCHC), National Applied Research Laboratories (NARLabs), Taiwan. Funding informaiton

Author contributions

J.S.H., P-L.C., Y-S.L., and T-K.H. proposed the initial idea of annotating pharmacogene alleles using high-quality assemblies, initiated the project, and provided input during discussions.

T-Y.C. conceived the study, designed the analysis pipeline, acquired the HPRC data, performed the analyses, interpreted the results, and generated the figures and tables. T-Y.C. also managed the writing and organization of the manuscript, with suggestions and proofreading from J.S.H.

351   J.S.H. supervised the overall project.

352   All authors read and approved the final manuscript.

354   **Reference**

355   Chen, X., Shen, F., Gonzaludo, N., Malhotra, A., Rogert, C., Taft, R. J., Bentley,

356   D. R., & Eberle, M. A. (2021). Cyrius: accurate CYP2D6 genotyping using whole-

357   genome   sequencing   data.   *Pharmacogenomics   J*,   *21*(2),   251–261.

358   https://doi.org/10.1038/s41397-020-00205-5

359   Cheng, H., Concepcion, G. T., Faris, J. G., & et al. (2021). Haplotype-resolved de

360   novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2),

361   170–175. https://doi.org/10.1038/s41592-020-01056-5

362   Del Tredici, A. L., Malhotra, A., Dedek, M., & et al. (2018). Frequency of

363   CYP2D6 alleles including structural variants in the United States. *Frontiers in*

364   *Pharmacology*, *9*, 305. https://doi.org/10.3389/fphar.2018.00305

365   Gaedigk, A., Ingelman-Sundberg, M., Miller, N. A., & et al. (2018). The

366   Pharmacogene Variation (PharmVar) Consortium: Incorporation of the Human

367   Cytochrome P450 (CYP) Allele Nomenclature Database. *Clinical Pharmacology &*

368   *Therapeutics*, *103*(3), 399–401. https://doi.org/10.1002/cpt.910

369   Gaedigk, A., Sangkuhl, K., Whirl-Carrillo, M., Klein, T., & Leeder, J. S. (2017).

370   Prediction of CYP2D6 phenotype from genotype across world populations. *Genet Med*,

371   *19*(1), 69–76. https://doi.org/10.1038/gim.2016.80

372   Hari, A., Zhou, Q., Gonzaludo, N., Harting, J., Scott, S. A., Qin, X., Scherer, S.,

Sahinalp, S. C., & Numanagić, I. (2023). An efficient genotyper and star-allele caller

for   pharmacogenomics.   *Genome   Res*,   *33*(1),   61–70.

https://doi.org/10.1101/gr.277075.122

Hicks, J. K., Sangkuhl, K., Swen, J. J., & et al. (2015). Clinical Pharmacogenetics

Implementation Consortium guideline (CPIC) for CYP2D6 and CYP2C19 genotypes

and dosing of selective serotonin reuptake inhibitors. *Clinical Pharmacology &*

Therapeutics, *98*(2), 127–134. https://doi.org/10.1002/cpt.147

Hung, T. K., Liu, W. C., Lai, S. K., Chuang, H. W., Lee, Y. C., Lin, H. Y., Hsu,

C. L., Chen, C. Y., Yang, Y. C., Hsu, J. S., & Chen, P. L. (2024). Genetic complexity

of killer-cell immunoglobulin-like receptor genes in human pangenome assemblies.

Genome Res, *34*(8), 1211–1223. https://doi.org/10.1101/gr.278358.123

Lee, S. B., Wheeler, M. M., Patterson, K., McGee, S., Dalton, R., Woodahl, E. L.,

Gaedigk, A., Thummel, K. E., & Nickerson, D. A. (2019). Stargazer: a software tool

for calling star alleles from next-generation sequencing data using CYP2D6 as a model.

Genet Med, *21*(2), 361–372. https://doi.org/10.1038/s41436-018-0054-0

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences.

Bioinformatics, *34*(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Liao, W. W., Asri, M., Wang, T., & et al. (2023). A draft human pangenome

reference. *Nature*, *617*(7960), 312–324. https://doi.org/10.1038/s41586-023-05896-x

Nurk, S., Koren, S., Rhie, A., & et al. (2022). The complete sequence of a human

genome. *Science*, *376*(6588), 44–53. https://doi.org/10.1126/science.abj6987

PharmGkb. (2023). Gene-Specific Information Tables.

Qiao, W., Yang, Y., Sebra, R., & et al. (2020). Long-read single molecule real-

time full gene sequencing of cytochrome P450-2B6. *Pharmacogenetics and Genomics*,

30(2), 37–46. https://doi.org/10.1097/FPC.0000000000000388

Relling, M. V., & Klein, T. E. (2011). CPIC: Clinical Pharmacogenetics

Implementation Consortium of the Pharmacogenomics Research Network. *Clinical*

Pharmacology & Therapeutics, *89*(3), 464–467. https://doi.org/10.1038/clpt.2010.279

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S.,

Getz, G., & Mesirov, J. P. (2011). Integrative genomics viewer. *Nature Biotechnology*,

29(1), 24–26. https://doi.org/10.1038/nbt.1754

Song, W., & et al. (2023). Benchmarking pharmacogene genotyping accuracy

across   sequencing   technologies.   *Pharmacogenomics   Journal*.

https://doi.org/10.1038/s41397-023-00310-2

Taylor, C., Crosby, I., Yip, V., Maguire, P., Pirmohamed, M., & Turner, R. M.

(2020). A Review of the Important Role of CYP2D6 in Pharmacogenomics. *Genes

(Basel)*, *11*(11). https://doi.org/10.3390/genes11111295

Twesigomwe, D., Wright, G. E. B., Drögemöller, B. I., da Rocha, J., Lombard, Z.,

& Hazelhurst, S. (2020). A systematic comparison of pharmacogene star allele calling

bioinformatics algorithms: a focus on CYP2D6 genotyping. *NPJ Genom Med*, *5*, 30.

https://doi.org/10.1038/s41525-020-0135-2

Wang, J., & et al. (2021). StarPhase: haplotype-aware variant calling and

diplotype   assignment   from   long-read   assemblies.   *bioRxiv*.

https://doi.org/10.1101/2021.12.15.472864

Zanger, U. M., & Schwab, M. (2013). Cytochrome P450 enzymes in drug

metabolism: regulation of gene expression, enzyme activities, and impact of genetic

variation.   *Pharmacological   Therapeutics*,   *138*(1),   103–141.

https://doi.org/10.1016/j.pharmthera.2012.12.007

Zhou, Y., Ingelman-Sundberg, M., & Lauschke, V. M. (2017). Worldwide

Distribution of Cytochrome P450 Alleles: A Meta-analysis of Population-scale

Sequencing   Projects.   *Clin   Pharmacol   Ther*,   *102*(4),   688–700.

https://doi.org/10.1002/cpt.690

Zhou, Y., Song, L., & Li, H. (2024). Full resolution HLA and KIR genes

annotation   for   human   genome   assemblies.   *bioRxiv*.

https://doi.org/10.1101/2024.01.20.576452