Scanning and active sampling behaviours emerge from conserved insect neural circuits

Cody A Freas[1,2], Antoine Wystrach[1]

[1]Research Center on Animal Cognition (CRCA), Center for Integrative Biology (CBI), CNRS, University of Toulouse, Toulouse, France

[2]School of Natural Sciences, Macquarie University, Sydney, NSW Australia

* Corresponding Author

Address for correspondence:
Cody A Freas
School of Natural Sciences
Macquarie University
Sydney, NSW 2113 Australia
Email: cody.freas@mq.edu.au

Abstract

Navigating insects often pause and rotate to sample their surroundings, behaviours termed scanning. These and other active sampling behaviours embody navigational uncertainty, and are key for spatial learning, yet their neural basis remains unclear and existing models impose scanning behaviours rather than explaining its emergence. Here, we show that desert ants' scanning dynamics can emerge spontaneously from the same conserved neural circuits used for goal-directed navigation, without requiring a specialized scanning module. We built a biologically grounded model combining central complex (CX) steering and lateral accessory lobe (LAL) oscillators, and added a downstream stochastic inhibition of forward speed. This minimal system produced diverse, realistic scan dynamics; saccades, fixations and reversals, whose features were qualitatively compared to high-speed video recordings of *Melophorus bagoti* scanning. Detailed analysis of these natural scans confirmed model predictions, including how scan structure depends on oscillator phase, goal-heading deviation, and navigational uncertainty. Furthermore, the model reveals that simple modulation of forward speed unifies a broad range of behaviors across ant species, from dashes to smooth oscillatory trajectories to pirouettes and voltes. Crucially, it establishes a distributed control principle, where forward speed acts as a single adjustable parameter, for both individuals and through evolution, to regulate the balance between goal-driven exploitation and information-seeking exploration.

Keywords: Central complex; Lateral Accessory Lobes; Distributed motor control; Central pattern generator (CPG); Oscillations

Background

Insects will often interrupt their forward movement and rotate to sample visual information from multiple body orientations. These behaviours appear widespread across both walking, crawling and flying insects (as well as other taxa), and take various forms and names such as 'scans', 'volte', 'pirouette', 'turn back' 'peering', 'sweep' or 'dance' (Baird et al., 2012; Gomez-Marin and Louis, 2014; Deeti et al., 2023a; Fleischmann et al., 2017; Graham and Collett, 2002; Lehrer, 1993; Mouritsen et al., 2004; Müller and Wehner, 2010; Stürzl et al., 2016; Tarsitano and Andrew, 1999; Ugolini, 2006; Wallace, 1959; Wehner et al., 1992; Wystrach et al., 2014; Zeil and Fleischmann, 2019). Despite its prevalence across taxa, the neural mechanisms underlying these seemingly highly structured behaviors remain unclear.

Scans are especially prevalent in visually guided ants (Deeti et al., 2023a; Fleischmann et al., 2017; Freas et al., 2018; Müller and Wehner, 2010; Nicholson et al., 1999; Wehner et al., 1996, 1996; Wystrach et al., 2014; Zeil and Fleischmann, 2019). Ants occasionally cease forward motion, and the ensuing scanning behaviours consist of distinct periods: saccades, when the navigator is rotating, and fixations, when the ant pauses all movement (Figure 1A-C; Video 1,2, Müller and Wehner, 2010; Wystrach *et al.*, 2014; Zeil and Fleischmann, 2019; Deeti, Cheng, *et al.*, 2023; Freas and Cheng, 2025). Multiple saccades and fixations make up a singular scan, which may contain one or more rotational reversals (right then left/left then right) before forward movement resumes. The probability of scanning, while stochastic (Deeti, Cheng, *et al.*, 2023), increases with uncertainty, when views are unfamiliar or unexpected (Deeti et al., 2023a; Freas et al., 2018; Schwarz et al., 2020b; Wystrach et al., 2014) such as during learning walks (Müller and Wehner, 2010; Zeil and Fleischmann, 2019) and novel route formation (Freas and Cheng, 2025, 2022; Haalck et al., 2023), or when views have been associated with aversive events (Freas et al., 2022; Wystrach et al., 2020a).

Functionally, scans are associated with information sampling, helping ants choose headings when cues are unfamiliar or in conflicts (Wystrach et al., 2014; Wystrach, Buehlmann, *et al.*, 2020; Deeti, Cheng, *et al.*, 2023; Freas and Cheng, 2025). Additionally, their prevalence during both learning-walks and early route formation indicates a role in view acquisition (Freas and Cheng, 2025; Müller and Wehner, 2010; Wystrach, 2023; Wystrach et al., 2014; Zeil, 2023; Zeil and Fleischmann, 2019).

Our mechanistic understanding of insect navigation has improved tremendously in the last decades - thanks to a combination of behavioural, neurobiological and modelling studies (Honkanen et al., 2019; Webb and Wystrach, 2016) - but current models do not explain the emergence and natural dynamics of scanning behaviours. Instead, 'scanning routines' are sometimes force-added in models to help the agent or robot sample views across directions (Baddeley et al., 2012; Gattaux et al., 2023; Husbands et al., 2021; Knight et al., 2019; Le Möel and Wystrach, 2020; Murray et al., 2019; Wystrach et al., 2013).

Here we ask whether and how ants' scanning dynamics could emerge from neural
structures typically implicated in insect navigation: the central complex (CX) and the
lateral accessory lobes (LALs). While the circuitry of these structures is detailed
primarily in a few insect model species, it is sufficiently conserved to allow for neural
modelling of insect navigation in various contexts (Adden et al., 2022; Clément et al.,
2023; Collett et al., 2025; Goulard et al., 2023; Shiu et al., 2024; Steinbeck et al., 2020;
Stone et al., 2017; Webb and Wystrach, 2016; Wystrach et al., 2020b). Both CX and
LAL regions are well established to integrate multiple directional information and
control steering commands in navigating insects (Franconville et al., 2018; Heinze,
2017; Honkanen et al., 2019; Hulse et al., 2021; Li et al., 2020; Namiki and Kanzaki,
2016; Pfeiffer and Homberg, 2014).

Previous studies have established notably how goal headings are set in the central
complex (CX) (Fisher, 2022; Green and Maimon, 2018) and modelling efforts have
shown that this circuitry can naturally accommodate innate and learnt guidance such
as path integration, learn vectors, visual route following or homing as observed in ants
and bees (Goulard et al., 2021; Honkanen et al., 2019; Stone et al., 2017 Le Moël et
al. 2019; Wystrach, 2023; Wystrach et al., 2020b). In parallel, oscillatory dynamics in
the lateral accessory lobes (LALs), produced by reciprocal inhibition across both
hemispheres and conveyed by so-called descending flip-flopping neurons, were
shown to drive the spontaneous zigzags displayed by moths upon losing their
pheromone plume (Kanzaki and Mishima, 1996; Mishima and Kanzaki, 1998, 1999;
Wada and Kanzaki, 2005; Kanzaki et al., 2005; Iwano et al., 2010). Here also,
subsequent modelling efforts have shown how these circuits can equally support the
continuous lateral oscillations displayed by a wide range of insect species (Adden et
al., 2022; Kanzaki et al., 2005; Namiki and Kanzaki, 2016; Steinbeck et al., 2020;
Wystrach et al., 2016) including ants (Clément et al., 2023; Dauzere-Peres et al.,
2026). Building on these elements, the present work addresses a distinct question that
current navigation models do not resolve: whether discrete active sampling behaviours
such as ant scanning require specialised control mechanisms, or instead emerge from
interactions within these conserved navigation circuits.

To investigate this, we analysed high-speed recordings of natural scanning behaviours
in *Melophorus bagoti* foragers and asked whether their detailed dynamics could be
captured by modelling the interactions between the CX and the LALs, under the
condition of occasional halted forward movement. We developed a biologically
constrained neural model of a CX steering output that modulates a downstream
intrinsic LALs oscillator, under the assumption that forward speed may be transiently
gated by a stochastic extrinsic signal. Finally, we added the physical constraint that
forward velocity increasingly limits rotation. Without introducing a dedicated scanning
module, this model is sufficient to reproduce a diversity key qualitative features of
natural scans, including the relative durations and amplitudes of saccades, fixations,

directional reversals, and the emergence of rare full-loop scans, which arise as a
consequence of CX–LAL and forward-angular speed interactions.

In addition to behaviors typically categorized as 'scans,' the model produces
information sampling behaviors often described using different terminology; including
'pirouettes' observed in ants (Fleischmann et al., 2017; Zeil and Fleischmann, 2019),
dung beetle 'dances' (Baird et al., 2012), and *Drosophila* 'reorientations' (Demir et al.,
2020; Gomez-Marin et al., 2011). Furthermore, by reducing rather than terminating
forward speed, the model also captures continuous-movement behaviors such as
voltes and large oscillating paths (Fleischmann et al., 2017; Zeil and Fleischmann,
2019; Clément, Schwarz and Wystrach, 2023). These findings point to a conserved
neural control strategy where forward speed inhibition gates the expression of angular
motor output, allowing for shifting seamlessly between information gathering and goal-
oriented progression. The few novel assumptions added here enable us to form
predictions for future work.

Neural substrates of insect navigation

To interpret these results, we briefly summarise the neural substrates that motivated
the model. The CX tracks the insects' heading relative to the world, transforms
egocentric sensory information into allocentric current and goals directions, and
compares both to output steering commands. Within the CX's substructures, the
current heading direction is represented neurally as a bump of activity shifting along
the protocerebral bridge (PB) and the Ellipsoid Body (EB), while goal directions are
represented in the fan-shaped body (FB) (Honkanen et al., 2019; Kim et al., 2019;
Pfeiffer and Homberg, 2014; Seelig and Jayaraman, 2015; Stone et al., 2017). Current
and goal direction representations are compared within the CX and project left/right
steering commands to the LALs (Green et al., 2019; Honkanen et al., 2019; Iwano et
al., 2010; Mussells Pires et al., 2024; Westeinde et al., 2024; Figure 2A). The goal
direction in the CX can be updated by various pathways from different sensory
modalities (Honkanen et al., 2019), such as wind and odours (Buehlmann et al., 2020;
Knaden and Graham, 2016; Müller and Wehner, 2007; Steck et al., 2011), visual
familiarity via the Mushroom bodies (Wystrach et al., 2020b) or path integration (Lu et
al., 2022; Lyu et al., 2022; Stone et al., 2017). In experienced ants, these different
sources of directional information are integrated and usually act in synergy to form a
goal directed either at the nest, or towards a known food source (Buehlmann et al.,
2020; Knaden and Graham, 2016; Müller and Wehner, 2007; Steck et al., 2011;
Wehner et al., 2016; Wystrach et al., 2015; Wystrach and Schwarz, 2013). In the
current modelling work we assumed the existence of such a goal direction in the FB
of the CX.

The LALs are premotor centers that act as a bottleneck, integrating information from
multiple higher processing centres and sensory inputs, including outputs from the

central complex, and relay these signals via descending neurons that transmit motor
commands to the thorax (Namiki and Kanzaki, 2016; Shih et al., 2015; Steinbeck et
al., 2020). The LALs also produce long-lasting, intrinsic alternation between left and
right turn via so-called flip-flop neurons (Berni, 2015; Iwano et al., 2010; Kanzaki,
2005; Namiki and Kanzaki, 2016; Namiki et al., 2014, Steinbeck et al., 2020), which
can result in the regular lateral oscillations observed in many insects (Freas and
Cheng, 2022; Izquierdo and Lockery, 2010; Kanzaki et al., 1992; Kuenen and Baker,
1983; Namiki and Kanzaki, 2016; Olberg, 1983; Wystrach et al., 2016), notably in ants
(Video 4, (Clément et al., 2023; Collet et al., 2025; Deeti and Cheng, 2025; Haalck et
al., 2023; Hangartner, 1967). The generation of oscillatory movements allows for
continuous spatial sampling of information, and modeling studies demonstrate that
when these oscillations are modulated by odor or visual cues, they offer a highly
effective strategy for odour plume or gradient tracking, visual route following or goal
pinpointing (Adden et al., 2022; Kodzhabashev and Mangan, 2015; Le Möel and
Wystrach, 2020; Wystrach et al., 2016).

Together, this CX–LAL circuitry is well suited to control continuous steering and
sampling during locomotion (Collett et al., 2025). However, until now it is unknown if
this same circuitry underlies the discrete structures of ant scanning behaviour without
invoking a dedicated control routine.

229
Figure 1. Image composites of fixations (pauses – typically 50-150ms) during three different examples
of scanning behaviours in *Melophorus bagoti*, all taken within 1m of the nest entrance, in inexperienced

ants forming their route to a feeder. Images were extracted and compiled from highspeed video taken
at 600fps at 1080p using a Chronos 2.1HD camera (field of view, 30 cm × 17 cm). For each image,
fixations are indicated by consecutively numbered black arrows denoting the ant's orientation (except
reversal). Rotational movement between fixations, saccades, are classified as either away (pink) or
towards (green) the goal direction (~90°). When a fixation precedes a change in turning (from left to
right in these examples), this fixation is classified as a reversal (orange). (**A**) Denotes a scanning
example where it reverses direction once but then turns in a complete loop with no reversal (fixations
6-16). (**B**) Shows a scan which contains multiple reversals (two). (**C**) Illustrates a scan with one reversal,
~180° from the goal direction. Black dotted arrows denote pre/post scan forward movement.

241
Figure 2. Diagrams of the brain region connections and outputs using the neural circuit model.
(**A**) The central complex (CX), a mid-brain region whose subregions compare representations of the
goal heading with a representation of the agent's compass-based current heading. The CX outputs
bilateral signals to turn the left (red line) and right (blue line) to the Lateral Accessory Lobes (LALs). (**B**)

Within the LALs oscillator, left (L) and right (R) neurons reciprocally inhibit one another (red and blue
connections), while attempting to maintain a basal firing rate via internal feedback (circular black
arrows), forming an oscillator which outputs a stable anti-phasic oscillatory activity between the L and
R neurons across time (red and blue lines). (**C**) A steering signal is outputted from the LALs and results
in the agent's forward and angular movement. This steering signal stopped via an external inhibitory
'freeze signal', that breaks the angular and forward speed central pattern generators (CPG), initiating
the scan. A threshold was implemented for the underlying angular drive to restart the CPG after each
fixation period, resulting in a saccade whose magnitude was determined by this accumulated angular
drive. The agent's angular speed was normalised by the forward speed (linear normalisation - angular
speed = angular drive/(forward speed + 0.1)) so the agent produces larger saccades magnitudes during
scans. Threshold was determined to roughly mirror saccade magnitudes in real world ants. Scan
duration was implemented as an exponential duration distribution through a dice-roll at each simulation
step. (**D**) Characteristics of an example path, with a single scanning bout, generated by the model.
Black dots represent the agent's head position at each simulated step coupled with coloured bars which
indicate both the forward speed (arbitrary scale) and heading direction of each step. During the
scanning bout, the agent's forward speed is zero and the model produces several fixations in separate
directions. The fixations of this scanning bout are zoomed and separated into a sequence of fixations.
Coloured arrows indicate saccades, rotational movements between fixations which were defined as
either away (pink) or towards (green) the goal direction. (**E**) The modelled CX's turning signal output
towards the goal direction, based on if the agent's heading direction is to the left or right of the goal. (**F**)
The oscillatory activity between the R and L neurons in the LALs during the example path. The agent's
(**G**) heading direction, (**H**) angular speed, and (**I**) forward at each step during the path.

Results and Discussion

A simple model for scanning behaviour

We first constructed a simple model of the insect central complex (CX; Figure 2A),
following standard approaches (Goulard et al., 2023; Stone et al., 2017; Wystrach,
2023; Wystrach et al., 2020b). This circuit compares the agent's current heading to a
stored goal heading (which, in our simulations, is directionally fixed) and generates
lateralised steering signals to reduce heading error. Unlike previous models (but as in
Adden *et al.*, 2022) the CX output here does not steer the agent directly, but instead
modulates an intrinsic oscillator that generates alternating turns (Figure 2B), a
mechanism consistent with the continuous oscillatory patterns known to underlie ant
navigation (Clément, Schwarz and Wystrach, 2023).

This oscillator, which is widespread in insects (Cheng, 2024; Kanzaki, 2005; Kanzaki
and Mishima, 1996), resides in the lateral accessory lobes (LAL), a pre-motor region
that receives, among others, CX outputs (Heinze, 2017). We modelled the oscillator
as a system of reciprocal inhibition between the left and right hemispheres, producing
alternating steering signals (Kanzaki and Mishima, 1996; Steinbeck, Adden and
Graham, 2020; Clément, Schwarz and Wystrach, 2023). The difference between these
left and right signals - such as conveyed by descending neurons (DN) to the thoracic
ganglia (Büschges and Ache, 2025) - determines the agent's angular speed, thereby
generating the rhythmic turning dynamics akin to those observed in ants (Clément,
Schwarz and Wystrach, 2023). In combination with the CX modulation, this oscillator
produces a continuous oscillatory trajectory generally oriented toward the goal
direction set in the CX (which is always to the right in our examples; Figure 2D).

To generate scanning behaviour (Figure 1A–C), we added a mechanism to
intermittently interrupt forward movement, as observed in desert ants, where pauses
occur randomly (Deeti et al., 2023a). B*ecause we did not want to form pre-*
assumptions for how such a 'freeze signal' could be implemented in the insect nervous

system; in our model this was achieved using a simple external signal that halts
forward motion at random intervals. Since scan duration in real ants follows a Poisson
like distribution (Deeti et al., 2023a), we approximated this by performing at each
simulation step a dice-roll with equal probability (calibrated as 1/mean of ants
observed scan duration) to restart forward motion, producing the desired Poisson
distribution of scans' duration.
306
During scanning, real ants display rotational saccades of variable duration and angular
magnitude (Figure 1A–C). To replicate this, we introduced a threshold-based
mechanism: after each fixation (i.e., zero angular and forward speed), the underlying
angular steering signal accumulates until surpassing a threshold, triggering a saccade.
The resulting angular magnitude of the saccade corresponds to the sum of the angular
drive accumulated during the fixation. Here also we stuck to a non-neural, straight-
forward algorithmic level, as we did not want to make assumptions about how such a
cumulate-and-release mechanism could be neurally implemented in the insect brain
(see discussion for potential implementations).
316
With this, the agent now scans, but the heading deviation during scanning remains
within the same limited range as during forward movement, highly constrained toward
the goal direction. This does not reflect real scans, which involve much larger angular
deviations (Deeti et al., 2023a; Fleischmann et al., 2017; Freas et al., 2019; Freas and
Cheng, 2025; Müller and Wehner, 2010; Wystrach et al., 2014). We hypothesized that
this limited turning during forward movement is due in part to biomechanical
constraints: fast forward motion impedes sharp turns, while slowing down allows
easier rotation. When ants stop, they are free to execute sharp turns via leg
coordination that would be otherwise impossible (e.g., inner legs moving backward
and outer legs forward), enabling high rotation during scans (Figure 1). To implement
this constraint, we normalized the agent's angular speed by the inverse of forward
speed: the slower the agent moves, the more it turns for a given steering signal. As a
result, the agent now produces larger saccades and greater heading deviations during
scans, while remaining more constrained during forward movement. To match the
saccade amplitudes observed in real ants, we adjusted the threshold for breaking
fixation so that the resulting distribution of saccade angles (Figure 3A) approximates
those seen in ants (Figure 3B). Higher thresholds yield less frequent but larger
saccades.
335
336 Overall, the interaction between the CX output (Figure 2A), oscillator phase (Figure
337 2B), and the scan control mechanism (Figure 2C) now produces complex, nonlinear
338 dynamics, giving rise to emergent behaviours, which we explored and compared to
339 real ants in the following subsections.
340
341 *CX steering influences saccade magnitude*
342
343 A key prediction of our model is that the CX steering mechanism operates
344 continuously, including during scans, and its influence should be detectable in the
345 scan's structure. Specifically, the strength of the CX's corrective signal toward the goal
346 increases with the agent's angular deviation from that goal (Figure 2A).
347

If guided solely by the CX, the agent would barely turn away from the goal but due to
noise. However, actual steering is governed by the downstream oscillator in the LAL,
which ultimately determines the direction of each saccade during scans. Saccades
toward the goal occur when the oscillator is in phase with the CX corrective signal,
resulting in stronger angular drive and therefore larger saccades. Conversely,
saccades away from the goal arise when the oscillator and CX corrective signals are
in conflict, the later restraining the former, producing weaker angular drive and thus
smaller saccades. Since the CX's corrective signal roughly scales with angular
deviation (Figure 2A), this difference in saccade amplitude (toward vs. away from
goal) should be minimal when the agent is facing the goal direction but becomes more
pronounced as it is facing away (Figure 3C).

This predicted pattern closely matches real ant data (Figure 3D). Across all datasets,
including inexperienced and experienced ants, whether homing or foraging (treated as
random error in our LME), saccade amplitude was significantly explained by both the
saccade direction (towards and away from the goal) ($F_{(1,1890)}$ = 5.08, p = 0.024) and
angular deviation ($F_{(1,1890)}$ = 18.69, p < 0.001) and by and large explained by their
interaction indeed ($F_{(1,1890)}$ = 43.04, p < 0.001). Specifically, saccades toward the goal
were larger than those turning away from it, and this difference increased as the ants
faced away from their goal, supporting the model's prediction. Together, this strongly
supports that both the oscillator and CX are at play during scanning, with the later
continuously pulling the ants towards its goal heading.

Our statistical analysis on saccade amplitude also revealed significant fixed effects for
the individuals (LRstat =200.25; p < 0.001) and conditions (LRstat =49.08, p < 0.001)
(see Statistical Methods for details) as well as an interaction between heading
orientation (*angular deviation* from goal) and whether the previous fixation was a
reversal ($F_{(1,1890)}$ = 6.86, p = 0.009; Figure 4A), which we will address in the next
sections.

375
Figure 3. Saccade angle distribution and the effect of CX steering guidance on saccade angle
magnitude. Saccade angle distributions in (**A**) the modelled agent and (**B**) real ants with all scan
conditions combined. After each fixation, the subsequent next saccade angle is plotted against the
fixation's angular divergence from the goal direction in both (**C**) the model agent and (**D**) real ants with
all conditions combined. The general trends of the model are classified through coloured arrows (away-
green, towards-pink) representing the increasing next saccade angle towards the goal direction and the
decreasing next saccade angle away from the goal direction of the modelled agent when fixation
orientation from the goal direction was large. This represents the effect of the changing strength of the
CX steering signal with increasing divergence from the goal direction; easier to turn towards goal and
thus large saccades, while harder to turn further away and small saccades. This pattern is replicated
and significant in real world ants. For the box and whisker plots in panels C and D, the box spans the
inter quartile range while the horizontal line indicates the mean. Whiskers extend to the INr X 1.5 while
outliers beyond this range are shown as '+' symbols. The indentation of the bar around the mean
indicates the 95% confidence interval.
390
391 *The oscillator phase influences fixation duration*
392
Having explored saccade amplitude, we next asked whether the model also accounts
for fixation duration. In our model, fixations during scanning are periods of fully
inhibited locomotion. A fixation ends when the angular drive (generated by the
oscillator) crosses a threshold of angular drive (left or right), breaking the inhibition
and triggering a saccade (Figure 4A–C). Since the angular drive (L - R activity of the
LAL) fluctuates cyclically with the oscillator phase, the time required to reach this
threshold naturally varies.

Notably, the models' fixations can be especially prolonged during reversal phases;
when the oscillator transitions from left to right, or vice versa. In extreme cases, the
angular drive initially builds toward the direction of the previous saccade (e.g., left),
but before crossing the threshold, the oscillator reverses phase. The drive then shifts
in the opposite direction (e.g., right) and must accumulate again in that new direction
before triggering a saccade (Figure 4B, yellow period of second reversal). As a result,
reversal fixations have a tendency to last longer than other fixations (Figure 4A,D).
Remarkably, ant behavior supports the model's prediction (Figure 4E). In our linear
mixed-effects model (LME), reversal fixation status was the only significant predictor
of fixation duration ($F_{(1,1894)}$ = 76.01; p < 0.001).

The model also predicts a subtler pattern: fixation duration gradually decreases with
distance from a reversal fixation (Figure 4D). This occurs because angular drive is on
average strongest mid-cycle (i.e., in the middle of a left or right phase) and weakest
near phase transitions. When we replaced the binary "reversal/non-reversal" variable
with this sequential phase information (Figure 4A) in our statistical model, the effect
on fixation duration remained significant, but only as an interaction with the upcoming
saccade's direction relative to the goal ($F_{(1,1894)}$ = 4.65; p = 0.031). Specifically,
because the CX upregulates the oscillator on the side toward the goal, the angular
drive raises quicker and fixation duration tends to be shorter for saccades in that
direction.

Together, these results support the idea that fixation duration arises from a threshold-
crossing mechanism and signal governed by the oscillator's phase and its interaction
with CX input. Despite outward stillness during fixation, oscillator and CX are
continuously active and shaping scan dynamics.
424
425 *Influence of neural noise and thresholds on fixation duration*
426
Our model predicts a general negative correlation between fixation duration and the
amplitude of the subsequent saccade (Figure 4F, red). This arises because a short
fixation - that is, a rapid threshold crossing - implies a strong angular drive, and
stronger angular drive typically results in a larger saccade.

However, we were initially surprised to observe in our model that this negative
correlation breaks down for the shortest fixations (Q1): in this regime the agents exhibit
a positive correlation instead (Figure 4F, blue). This pattern is an indirect consequence
of neural noise, which is systematically introduced in the model. Specifically, noise in
the inhibitory neuron freezing movement causes variability in the fixation-breaking
threshold (illustrated as a grey band in Figure 4B). When this threshold is
stochastically lowered due to noise, it can be reached more quickly, producing very
short fixations if the current angular drive is strong. However, because the threshold
is lower, the accumulated angular drive at threshold crossing is necessarily reduced,
limiting the magnitude of the resulting saccade. This mechanism imposes an upper

bound on the accumulated drive, and therefore on saccade amplitude, for the shortest
fixations, creating a hard limit in the correlation plot (grey region, Figure 4F). The
outcome is a positive correlation between fixation duration and saccade amplitude for
the shortest fixations (Figure 4F, blue).

While this effect was not initially anticipated, its predictions are supported by
behavioral data. Ants' fixation duration is positively correlated with saccade amplitude
for the shortest fixations (Figure 4G, blue, Quartile 1: $F_{(1, 281)}$ = 11.00, p = 0.001; R^2 =
0.038; $t_{(281)}$ = 3.32, p = 0.001), and negatively correlated for longer fixations (Figure
4G, pink, Quartiles 2-4: $F_{(1, 795)}$ = 5.97, p = 0.015; R^2 = 0.007; $t_{(795)}$ = –2.44, p = 0.015).

The emergence of this complex pattern from the model, despite not being explicitly
designed to capture it, supports the hypothesis that fixations are governed by an
internal accumulation of excitatory activity (angular drive), which is released upon
reaching a threshold. This mechanism is consistent with integration-to-bound neural
models that have been proposed for self-initiated actions in other species (e.g.
Murakami *et al.*, 2014).

458
Figure 4. Relationship between fixation duration and reversal. (**A**) Sequence of fixation directions
and their duration for an example scan containing multiple reversals (orange), showcasing that the
longest fixations occur when the ant reverses directions on the next saccade (fixation duration
continuum, blue-min; yellow-max). (**B**) shows changes in the cumulative turning drive over the course
of the example scan. To initiate a saccade, the drive must surpass a threshold (line). This threshold
possesses some level of noise which leads to variance in when a saccade appears as well as to the
variance in saccade magnitude. Reversals occur when the oscillatory cycle changes phase (±), which
are associated with longer times for turning drive to accumulate beyond threshold (orange). (**C**) Angular
speed through the simulation, spikes represent saccades to the left and right and the change from
positive to negative illustrates reversal periods (orange). Relationship between fixation duration and
oscillatory cycle changes (orange) and subsequently reversals, occur in (**D**) modelled and (**E**) real ants.
In both, the longest fixation duration typically occurs at the reversal (0), increasing as the reversal
approaches (-), while decreasing post reversal (+). For these box and whisker plots, the box spans the
inter quartile range while the horizontal line indicates the mean. Whiskers extend to the INr X 1.5 while
outliers beyond this range are shown as '+' symbols. The indentation of the bar around the mean

indicates the 95% confidence interval. Correlation between fixation duration and the next saccade angle
in (**F**) modelled and (**G**) real ants. Data are split into the lower 25% (Q1, Blue) and upper 75% (Q2-Q4,
Pink) of the distribution. In both, the general tendency in Q1 is significantly positive (real ant; Linear
regression model; $F_{(1, 281)}$ = 11.00, p = 0.001), while beyond this (Q2-Q4) the tendency is significantly
negative ($F_{(1, 795)}$ = 5.97, p = 0.015). The 'unlikely' zone in grey depicts the area of high saccade
magnitudes following very short fixations, which should be highly unlikely given the low threshold drive
must accumulate needed to break fixation in these instances.

Random Timing of Scan Starts and Stops

Previous work on scanning behavior in ants (*Deeti et al., 2023*) found that the number
of saccades within each scanning bout follows a Poisson-like distribution (Figure 5B).
From this, they inferred that the initiation and termination of scans are governed by a
random-rate process.

In our model, this stochasticity is implemented by a probabilistic trigger: scan initiation
is triggered at a random time, and at each moment, scanning can end with a probability
proportional to (*1 / mean ants' observed scan duration*), producing the desired poisson
distribution of number of saccades (Figure 5A). Because of this, scans can begin or
end at arbitrary points in the oscillatory cycle. This leads to a specific prediction: in
scans containing multiple reversals (direction changes in turning), the first and last
sweeps, defined here as a bout of sequential saccades in the *same* direction, should
often be truncated compared to "full" sweeps (e.g., the second or penultimate sweeps),
which start and end exactly at reversals. In other words, sweeps bounded by two
reversal should reflect a full (one-sided) oscillatory cycle, while sweeps bounded by
the beginning or the end of a scan may not, due to the stochastic start or end of scans.
Therefore the latter should tend to be shorter, and thus contain less saccades, than
the former.

To test this, we excluded scans with one or zero reversals, focusing on bouts where
we could compare truncated sweeps (first and last) with adjacent full sweeps (second
and penultimate). The model predicts, and the data confirm, that truncated sweeps
contain fewer saccades than their adjacent full sweeps (Figures 5A–D). Specifically,
the first sweep is significantly shorter than the second (Figure 5C, z = 4.47; p <
0.0001), and the last sweep is significantly shorter than the penultimate (Figure 5D, z
= -3.30; p = 0.001).

These results thus support the stochasticity of scan initiation and termination. The
inhibition of forward motion seems to occur independently of the oscillatory phase.
Also, it confirms that within scans, saccade direction follows the underlying oscillatory
cycle, rather than being purely random.

Figure 5. Scans start and end irrespective of oscillator state and saccade number distributions. In the modelled agent, (**A**) the number of saccades within a scanning sweep (before first reversal) was compared with the post-reversal sweep number showing a general upward trend. (**B**) the number of saccades within the penultimate sweep compared to the final sweep of the scanning bout, showing a downward trend. In real ants, (**C**) comparison of the number of saccades in the first and second sweep and (**D**) the penultimate and final scanning sweeps. Both model and real ant sweep comparisons only contain scans which contained two plus reversals, in order to compare the starting/stopping sweeps with a full, within reversals sweep (second/penultimate). Statistical comparisons were made using Wilcoxon tests. The distribution of saccade counts in each scanning bout within (**E**) the model, showing a poisson distribution and in (**F**) real ants, showing a 'poisson-like' distribution.

CX strength links navigational uncertainty to scan structure

We next examined how navigational uncertainty, which can be modelled as the strength of the CX's steering signal, influences scan structure. In the model, a heavily weighted CX signal, representing high certainty in the goal direction, produces greater corrective steering, thus constraining the agent heading toward the goal. As this signal

weakens, so does its corrective influence, allowing for larger deviations away from the
goal direction. Thus the model predicts an inverse relationship between CX strength
(i.e., navigational certainty) and this divergence, which we quantified by measuring the
rotation away from the goal covered by the scanning ant before performing a first
reversal (Figure 6A, arrow). Ant behavioural data echoes this pattern: experienced
ants - which we assume have a high certainty in their goal direction - showed smaller
divergences from the goal on their first reversal than inexperienced ants (Figure 6B;
$F_{(3, 274)}$ = 3.20, p = 0.0239).

The model also predicts a positive association between CX strength and the agent's
saccade angle (Figure 6C, arrow). While initially counter intuitive, as a strongly
weighted CX signal constrains angular divergence away from the goal, strong CX
steering also produces much larger corrective saccades when it aligns with the
oscillator phase (as shown in Figure 3C). This pattern aligns well with our real ant
testing conditions, with significantly larger saccades in experienced ants vs.
inexperienced ants (Figure 6D; $F_{(1, 2137)}$ = 587.6, p < 0.001).

These results support the CX's corrective steering input role in modulating the
structure of scanning, tuning how tightly ants orient to their goal as well as how wide
their saccades are during scanning, especially when in phase with the oscillator.

551
Figure 6. The strength of the CX's corrective turn signal, and its inverse relationship with
navigational uncertainty. The CX's signal signal strength on the oscillator would be predicted to be
strong when navigational uncertainty is low, such as when the ant is highly experienced. Conversely,
CX signals would be weak when navigational uncertainty is high, like during route formation. The model
predicts that under high CX signal strength, (**A**) the first reversal direction (which is likely to follow a
sweep away from the goal) should be constrained angularly to closer to the goal direction. Additionally,
this metric should rise, the CX signal strength decreases. (**B**) These trends closely mirror the real ant
data, with highly experienced foragers, showing first reversal directions which were more constrained
angularly to closer to the goal direction compared to inexperienced forager conditions. The model also
predicts, somewhat initially continued intuitively, that (**C**) saccade angle should also be inversely
associated with CX signal strength (grey arrow), which mirrors (**D**) the reduction in saccade angle
amplitude in high uncertainty conditions in inexperienced foragers. Initially one might theorise low CX
strength should mean larger saccades, yet it is under high corrective CX steering strength that we see
large turns back towards the goal (See back to Figure 3C,D), resulting in this association. For all box
and whisker plots, the box spans the inter quartile range while the horizontal line indicates the mean.
Whiskers extend to the INr X 1.5 while outliers beyond this range are shown as '+' symbols. The
indentation of the bar around the mean indicates the 95% confidence interval.

Rare 'Full Loop' Scans reveal CX–Oscillator Interactions
571
572   Despite its simplicity, our model's closed-loop dynamics generate a surprising diversity
573   of scan forms. One striking example is the emergence of rare 'Full Loop' scans (Figure
574   7A,B; Video 4,5), in which the agent completes a long series of saccades in the same
575   direction, producing a full loop before resuming normal movement.
576
In the model, these occur when a strong central complex (CX) corrective steering
signal coincides with a critical point in the oscillator's cycle, effectively shifting the
oscillator's phase by producing a rebound of activity before the current cycle ends
(Figure 7A,B, black arrows). Instead of reversing direction at the usual reversal points,
the agent's saccade angle decreases (as usual) but then increases again in the same
turning direction, producing a complete rotation.
583
584   Such Full Loop scans are not unique to the model; they also occur, albeit rarely, in
585   real ants, and have been reported in multiple species (e.g., *M. bagoti* Figure 1A, Video
586   1; *Cataglyphis cursor,* Video 2; (Fleischmann et al., 2017; Freas et al., 2019; Freas
587   and Cheng, 2025; Zeil and Fleischmann, 2019).
588
Mechanistically, the occurrence of these full loop scans indicates that the CX
modulates the oscillator to enable a rebound within the same phase. If, by contrast,
the CX and oscillator acted purely in parallel (with their outputs summed
independently), the oscillator would produce a reversal in angular gain as usual,
conflicting with the CX output and preventing a fast completion of the loop. Consistent
with the model, real ants show re-acceleration in the second half of the 'Full Loop' scan
(real ants: Figure 1A, fixations 12–16; Videos 1–3; model: Figure 7A,B), supporting
the idea that CX output is indeed branched upstream of the oscillators, and can shift
its phase.
598
599   These events are too rare in our current dataset for statistical analyses, and our simple
600   model is not designed for quantitative comparisons. Nonetheless, such rare behaviors
601   may offer valuable insights into the coupling between oscillatory motor control and
602   upstream mechanisms, and could be a promising focus for future studies.
603

604
Figure 7. Diverse behaviours emerge from interactions between the central complex's (CX)
steering signal and the modulation of the agent's forward speed. (**A**) A single 'Full Loop' scan
(Video 5), the agent terminates forward speed and exhibits a scanning bout with both fixations and
saccades. Here, the CX's corrective steering excites the oscillator and reverses its phase (black arrow),
resulting in the agent continuing the loop rotation rather than reversing. These full loop behaviours are
rare within scanning (most reverse) but widely observed in ants (Video 1-3). (**B**) Shows a double 'Full
Loop' scan example where the agent performs two full loop rotations in a scan. As in panel A,
(Simulations - Video 6). Here, the oscillator's phase is reversed by the CX twice (black arrows). (**C**) A
'volte' is a loop without stopping (Simulation - Video 7). A common behaviour in *Cataglyphis* desert ants
(Fleischmann et al., 2017). This behaviour arises in the agent when forward speed is low but not
stopped, boosting angular speed and allowing the CX steering output to reverse oscillator phase without
fixations. (**D**) An example of a '*Myrmecia*-like' path in the agent, characterized by moderate forward
speed and larger lateral oscillations, reminiscent of real world Myrmecia ants (Clément, Schwarz and
Wystrach, 2023; e.g. Video - 4). Here, moderate forward speed allows for larger alternating turns and
lateral displacement.

620
621
622
623

Graded Forward Speed Control Generates Voltes and Oscillatory Patterns

In our initial implementation, forward speed inhibition was binary: the agent ran or stopped completely, as if the central pattern generator controlling leg movement were switched off at a random time. Insects, however, show more nuanced control. For example, desert ants can slow down as their path integration vector decreases (Buehlmann et al., 2018), or accelerate when on fully familiar routes (Clément et al., 2023; Haalck et al., 2023).

We therefore modified the model to allow forward speed inhibition to work along a continuum, with low inhibition merely slowing down the insect rather than stopping it. Because we assumed that forward speed biomechanically impedes angular speed, a sudden but partial reduction in forward speed produces sharper turns; and in some cases complete loops (Figure 7C) resembling so-called 'volte' that have been described in the ant literature (Fleischmann et al., 2017; Zeil and Fleischmann, 2019).

Furthermore, it is known that the oscillator phase itself modulates forward speed, which can be modeled simply as the sum of the oscillator's left and right activities (Clément et al., 2023). This coupling accelerates the agent when facing its travel direction and slows it when oriented sideways, producing efficient trajectories to cover ground while looking on the sides. The expression of this control varies among species and is particularly strong in *Myrmecia croslandi* (Clément, Schwarz and Wystrach, 2023; Video 4; see also Video 3 for *M. nigriceps*). In the model, continuously operating at lower forward speed, releases the constraint on angular speed and naturally produces the large, regular oscillations seen in these ants (Figure 7D) (See Supplemental Figure 1 for example paths).

Interestingly, in such slower agents, increasing the strength of the CX's influence on the oscillator produces not only more goal-oriented, faster trajectories, but also increases oscillation frequencies while decreasing their regularity (Supplemental Figure 1A,B). These four covarying properties mirrors the behavior of *M. croslandi* ants on familiar routes compared to unfamiliar terrain (Clément et al., 2023), consistent with the idea that visual familiarity generates strong CX goal-heading signals (Wystrach, 2023).

Finally, in strongly oscillating situations, full loops can also arise without apparent external inhibitory signal. Instead, they can emerge naturally from the continuous oscillatory control on forward speed, which slows the animal when facing away from its goal and thus enhances the rotation at the end of a sweep (e.g., *Myrmecia nigriceps*, Video 3). This suggests that the forward–angular speed coupling is a general property of the locomotor system, not solely tied to an external "stop" phase, though whether it is purely biomechanical or neurally reinforced remains to be tested.

Overall, introducing graded forward speed control shows that the same CX–oscillator interactions, originally modeled for scanning, can also explain patterns observed in continuous navigation (periods of goal-directed forward movement), linking route familiarity, speed modulation, scanning behaviours, CX and oscillatory turning dynamics.

General Discussion

This study shows that diverse movement behaviors in ants, including scanning,
pirouettes, voltes, and goal-directed runs, can all emerge from a single, conserved
neural architecture. By combining biologically grounded models of central complex
(CX) steering, intrinsic lateral accessory lobe (LAL) oscillators, and assuming a
physical constraint of forward speed onto angular speed, our models show that
modulation of forward speed can capture fine-grained scanning dynamics that match
high-speed behavioral data. Importantly, these behaviors do not require a dedicated
scanning module; instead, scans -together with a panel of observed intermediates
behaviours- arise naturally from continuous control mechanisms already implicated in
insect navigation. More broadly, these findings suggest that scan-like behaviours need
not be interpreted as specialised routines for acquiring nest-aligned views.

CX and Oscillator Interactions

In our model, the central complex (CX) outputs modulate the oscillator circuit in the
lateral accessory lobes (LAL), producing goal-oriented oscillatory paths (Figure 2D;
Figure 7A-D, for all paths goal direction to the right). While the CX–LAL pathways
have been modelled (Adden et al., 2022) in the context of plume-tracking in moths,
their potential implication in ant navigation remains unexplored. The scanning
dynamics observed here constrain possible architectures and offer insights into how
the CX output may shape motor control via the LAL.

CX steering gain increases with angular deviation

Our data show that corrective saccade amplitude increases with the agent's angular
deviation from the goal (Figure 3) which is consistent with behavioural data in ants and
flies during continuous navigation (Lent et al., 2010; Westeinde et al., 2024) and
suggests that the CX output signal increase with angular deviation. In Drosophila, this
signature has been proposed to involve PFL2 neurons, which respond maximally when the fly
faces away from the goal, and modulate steering gain by converging with PFL3 neurons
(which drive left or right turns) onto downstream descending neurons (Westeinde et al., 2024).
Here, for the sake of simplicity, we only modelled the two PFL3 populations (PFL-L
and PFL-R), tuned to both ±135° from the goal (i.e., ±45° from the anti-goal), which
also mimics CX steering cells observed in monarch butterflies (see Fig.4G of Beetz,
Kraus and el Jundi, 2023), and result in similar steering dynamics as observed in
Drosophila (compare Figure 2A and Westeinde *et al.*, 2024). Whatever the exact
implementation, the preserved relation between amplitude and angle away from the
goal during scan (Figure 3) supports the idea of a conserved CX output function that
encodes angular deviation from the goal with increasing strength, and is at play during
both scanning and continuous navigation.

CX directly modulates the LAL oscillator

A key question is whether the CX directly modulates the LAL oscillator (serial control),
or whether both act independently in parallel.

In moths, descending neurons in the LALs exhibit characteristic 'flip-flop' activity
patterns that correlate with zigzagging maneuvers (Olberg, 1983; Kanzaki and Ikeda,
1994). Computational models suggest that having these LAL neurons modulated by
the CX output can explain aspects of the moths' plume-tracking behaviour (Adden et
al., 2022). In ants, behavioural studies show that strong directional drives elicited by
the path integrator or visual familiarity do not only gain behavioural weights and
sharpen directional accuracy (Wehner et al., 2016; Wystrach et al. 2015, Legge et al.
2014) but also increase the ants' oscillation frequency (Haalck et al., 2023, Clément
et al., 2023). Assuming that the path integrator and visual familiarity modulate goal
signals in the CX - as modelled here and elsewhere (Wystrach et al., 2020b, Stone et
al., 2017)- and that the intrinsic oscillator is in the LAL (Clément et al., 2023, Steinbeck
et al., 2020), this frequency increase suggests that CX output modulates the intrinsic
oscillatory activity of the LAL. Similar increases in ants oscillation frequency also occur
in response to optic-flow prediction errors, although these signal may bypass the CX
(Dauzere-Peres and Wystrach, 2024).

Our findings offer a qualitatively different form of evidence for a direct modulation of
the LAL oscillator by the CX. The rare 'full loop' scans - 360° turns without reversal -
only emerge in the model because the CX can shift the oscillator's phase mid-cycle
(Figure 3A,B). This requires tight coupling between CX output and LAL dynamics, and
would not occur if both acted independently. Together, present and past results
provide strong support for a direct, serial modulation of the LAL oscillator by the CX in
ants.

An opponent process in the CX output

In our model, the output of the CX to the LALs (via PFL neurons) provides both
ipsilateral excitatory connections and a contralateral inhibitory connection (Figure 2A).
In insects, the latter may be indirect, via the stimulation of inhibitory neurons such as
the protocerebral bilateral neurons in the LAL (Kanzaki et al., 2004; Mishima and
Kanzaki, 1999). Functionally, this contralateral inhibition ensures that only one side of
the LAL is activated by the CX at a time, avoiding simultaneous bilateral activation,
which in the model would produce inappropriate bursts of forward motion, particularly
when the agent is misoriented.

More generally, this opponent organisation between the left and right hemisphere
converts CX directional information into a normalised steering signal that reflects
angular deviation. Similar opponent processes have been proposed for mushroom
body output neurons encoding approach vs. avoidance, or left vs. right scene
familiarity signals (Le Möel and Wystrach, 2020; Murray et al., 2019; Wystrach, 2023),
suggesting that lateralised, opponent coding is a general principle in insect
navigational circuits for ensuring stable and interpretable motor outputs.

A Simple Control Principle: Forward Speed Gates Exploration

753
754 A key insight from this work is the importance of the assumed constraint of forward
755 speed on the expression of body rotation. This effectively enables a one-dimensional
756 control to adjust the expression of active sampling along the tradeoff between
757 exploitation and exploration. At one extreme, forward speed is fully inhibited, halting
758 leg CPG output, and making angular drive maximally expressed, producing what we

term scans. At the other, high forward speed suppresses angular turns, yielding straight, fast goal-directed trajectories. By simply adjusting this one parameter, the model generates a spectrum of behaviors, from sweeping oscillations to full rotational scans, mirroring inter- and intra-species variation observed in real ants (Figure 8). This spectrum of apparently distinct behaviours does not reflect different mechanisms, but the same distributed control process. The diversity arises from the dynamic interactions between the CX, the oscillator, and thoracic CPGs, and, crucially, the mechanical constraint of forward speed on rotation.

Figure 8. Summary of the behavioural spectrum produced by the modelled agent as a function of forward speed inhibition facilitating angular speed. High forward speed results in an agent that 'sprints', with its straight paths resembling desert ants (*Melophorrs bagoti and Cataglyphis*). Decreasing forward speed progressively increases angular speed facilitation, leading to the large oscillations of Myrmecia under moderate forward speed (Clément, Schwarz and Wystrach, 2023; Video 4), 'voltes' under low forward speed, and ultimately scanning behaviours when forward speed reaches zero. Here, angular facilitation plateaus, the central pattern generator (CPG) is disrupted and the agent scans, with the choreography of this scan, are dictated by interactions between the CX signal strength and the oscillator is observed in ants (Real Ants - Video 1-3; Simulations - Video 5,6)

A continuum of movement modes across context and species

The continuum also helps explain interspecific variation. Desert ants (*Melophorus*
bagoti, multiple *Cataglyphis species, Occymyrmex robustior, etc…*), adapted to
thermophilic foraging, favour high forward speed, which thus limit oscillations
amplitude, and rely on brief, stochastic stops for visual sampling (Clément et al., 2023;
Deeti and Cheng, 2025; Muser et al., 2005); figure 8). *Myrmecia* ants, by contrast,
forage at lower temperature and can afford to operate at slower speeds, allowing large,
regular oscillations to coexist with forward progression (Clément et al., 2023); Figure
8). Our model can shift seamlessly from one style to the other by simply adjusting
forward speed.

An individual can also shift along the continuum to adapt to the current context. For
instance, experienced *Cataglyphis velox* running along their familiar route, where
navigational certainty -which can be implemented as a strong goal heading in the CX
(Figure 6)- favour exploitation by constraining angular deviation, increasing forward
speed and thus diminishing the occurrence scans. Similarly, *Myrmecia* can stop and
scan under high uncertainty (Freas et al., 2018; Islam et al., 2020) or accelerate and
straighten to escape adverse situations (Clément et al., 2023; Deeti et al., 2023b).

The emergence of such different-looking trajectories suggests that the modulation of
forward speed may represent an ancestral and fundamental control strategy in ants.
This simple control is used at both individual decision and evolutionary time scales :
to adjust individuals' behaviour to the current context, as well as ant species to their
ecology.

An ancestral design? Striking parallels with crawling larvae

This forward–angular coupling is not unique to ants. Similar mechanisms appear
across taxa (even in bacterial run–tumble cycles), highlighting a shared control
strategy (Cheng, 2024). In *Drosophila* larvae, angular reorientation also arises from a
continuous body-bend oscillator. The expression of body bending is constrained by
forward crawling peristalsis (Wystrach et al., 2016), which can be transiently
suppressed by descending neurons (e.g., PDM-DN), producing a stop and unmasking
the oscillator expression by enabling strong body rotation (Tastekin et al., 2018; Berni
et al., 2012; Pulver et al., 2015), akin to ant scans.

Larvae's stops also appear stochastic, and their frequency can be modulated by
negative sensory evidence such as going down attractive odour gradient (Berni et al.,
2012; Gomez-Marin and Louis, 2014), just as scanning in ants is more likely under
high navigational uncertainty (Deeti et al., 2023a; Freas et al., 2022, 2018; Freas and
Cheng, 2025; Schwarz et al., 2020a; Wystrach et al., 2020a, 2014).

Finally, in both ants (Buehlmann et al., 2018; Haalck et al., 2023) and larvae (Gomez-
Marin and Louis, 2014; Luo et al., 2010), forward speed control is not binary, but
graded, modulated smoothly by sensory evidence.

Although not fully developed in fly larvae, the CX and LAL are conserved structures,
ancestral to arthropods (Kanzaki, 2005; Kanzaki and Mishima, 1996; Pfeiffer and
Homberg, 2014; Heinze, 2024). Whether the descending neurons that inhibit forward
movements in *Drosophila* (Rayshubskiy et al., 2025; Tastekin et al., 2018) are similarly
conserved across arthropods, or represent convergent evolutions in some taxa,
remains to be seen.

In adult *Drosophila* and other insects, multiple descending neurons modulating forward
speed have been identified (Büschges and Ache, 2025). Some (BPNs), promote
acceleration (Bidaye et al., 2020), whereas others trigger strong speed reductions, or
halting responses (Sapkal et al., 2024). Furthermore, left-right firing rate differences
between descending neurons (DNa-1/DNa02) can predict rotational velocity driving
pivoting behaviour (Büschges and Ache, 2025; Rayshubskiy et al., 2025; Yang et al.,
2024); providing a circuit-level analogue to our modelled angular speed modulation.

Scanning in flying hymenopterans
834
Flying hymenopterans such as bees and wasps share much of their navigational
toolkit with ants (Collett et al., 2013; Collett and Hempel de Ibarra, 2023; Zeil, 2012).
As in ants, their scene recognition and path integration systems are thought to
generate a goal heading within the Central Complex (Stone et al., 2017; Honkanen
et al., 2019) and they also exhibit regular oscillations and scanning behaviours,
particularly during learning flights or when approaching a goal (Philippides et al.,
2013; Collett et al., 2013, 2023; Zeil et al., 1996). However, these behaviours are
expressed differently from those of ants, notably through sideways motion, arcs, and
loops, rather than by simply stopping and rotating on the spot.
844
The bodily constraints assumed in the present model are intended to reflect those of
a typical walking ant: the agent cannot move sideways. Ants are capable of
sideways or backward motion, but apparently only when no alternative is available,
such as when dragging a heavy item (Schwarz et al., 2017). It would be interesting
to determine whether scanning properties such as those observed in bees and
wasps could emerge from the model proposed here under different physical
constraints adapted to flight, while keeping the same neural mechanisms. Novel
behaviours may indeed evolve through changes in the body rather than in the brain
(Dudley, 2000).
854
855   *Model limitations and future directions*
856
While our model successfully captures fine-scale scan dynamics, several limitations
remain. First, we assumed a single, static goal direction represented in the fan-shaped
body of the CX. While this suffices to simulate realistic scanning of ants navigating
toward a known goal, real insects often juggle multiple directional cues, from path
integration, visual landmarks, wind, and olfaction, and may maintain simultaneous goal
representations. These goals may be employed sequentially or weighted adaptively
within the CX (Freas et al., 2020; Le Moël et al., 2019; Stone et al., 2017; Sun et al.,
2020; Wehner et al., 2016; Wystrach et al., 2015). Extending the model to simulate
multi-goal conflict or cue integration would allow exploration of how insects handle
navigational uncertainty, and how the latter impacts scanning.
867
In our simulations, the CX goal representation remained fixed in both direction and
strength throughout each trial. This simplification allowed us to isolate and compare
the effects of different CX strengths on scanning behaviour (Figure 6). However, goal
headings in the CX are likely to be updated continuously (including during scans
themselves), for instance via novel inputs from visual recognition in the MB (Goulard
et al., 2021; Sun et al., 2020). This would affect saccades direction and duration.
Exploring such dynamics lies beyond the scope of the present study but would

represent an interesting direction for future work. Notably, our proposed CX-LAL-Body
relationship could be implemented downstream of an existing path integration or
visual-based model (or both) to form predictions about the occurrence and dynamic of
scans along the path, as well as their impact on the emerging trajectories.
879
The model also does not currently explain the phenomena of nest-directed fixations.
These so-called "pirouettes" or "lookbacks", where longer fixations happen when
looking towards the nest have been reported in ants during learning walks or initial
route formation (Collett et al., 2023; Collett and Hempel De Ibarra, 2023; Fleischmann
et al., 2017; Freas and Cheng, 2018, 2025; Müller and Wehner, 2010; Robert et al.,
2018; Stürzl et al., 2016; Zeil and Fleischmann, 2019). While in our data set scans did
rarely extend beyond 150° away from the goal (Figure 6), our model predicts that when
this goal is weak - such as in entirely naive ants - reversal, smaller saccades and
longer fixations will tend towards 180° from the goal, which might coincide with the
origin, that is, the nest direction. Alternatively, the nest direction could actually be
represented as a goal heading. This could be addressed by incorporating either (a) a
dual-goal representation (e.g., one for the feeder and one for the nest) with modulated
weighting, or (b) a dynamic switch of goal orientation during the scan, perhaps based
on distance from home. This raises several empirical predictions. For example, if ants
perform scans while navigating along two-leg outbound routes where the goal is at 90°
to the nest's direction, then fixation patterns during scans may reveal whether multiple
goals influence scanning structure. If the CX indeed holds multiple co-active directional
representations, scan fixations should systematically cluster around these multiple
directions.
899
Finally, scan termination is currently modeled as a stochastic process, implemented
through a random-rate "freeze-release" mechanism to reproduce the observed
Poisson-like distribution of scan durations (Deeti et al., 2023a). However, behavioural
evidence suggests that scan duration is not entirely random. Scans tend to last longer
in contexts of high uncertainty, such as during learning walks or early route formation,
and become shorter or absent on well-learned routes (Wystrach et al., 2014). This
implies that the probability of termination likely depends on an internal variable,
perhaps an accumulating "forward drive" or confidence estimate, that builds over time
and releases the motor system from inhibition once a threshold is reached. This
mechanism would parallel the angular drive threshold that governs saccade initiation
in our current model (Figure 4) and resembles integration-to-bound processes
proposed in other species, including mammals (Murakami et al., 2014).
While our model does not include detailed biomechanics of leg coordination,
proprioceptive feedback, or thoracic pattern generators (CPGs), it makes testable
predictions about how angular and forward drive might interact at the motor level. For
instance, the assumption that angular drive -and forward drive- accumulates
continuously, even during motionless fixations predicts the existence of neurons or
motor circuits that integrate excitatory input over time and release movement when
thresholds are crossed, paralleling the control logic found in *Drosophila* larvae
(Rayshubskiy et al., 2025; Tastekin et al., 2018). Whether such a putative integration
over time is achieved at the level of descending neurons, or downstream in the thoracic
CPG themselves remains to be seen.
922
923 *Conclusions*
924

This study demonstrates that the diverse movement patterns of ants, from brief scans
to full pirouettes and sweeping oscillations, can arise from a single, conserved neural
architecture: the interaction between central complex (CX) steering and an intrinsic
oscillator in the lateral accessory lobes (LAL). By introducing minimal, biologically
plausible additions - a stochastic inhibition of forward speed, a saccade-initiation
threshold, and forward–angular speed coupling - the model reproduces a panel of fine-
scale scanning dynamics seen in high-speed recordings of *Melophorus bagoti*, and
generalizes across species and behavioural contexts.

A key insight is that forward speed acts as a distributed control dial, gating the
expression of angular movements and enabling smooth transitions between goal-
driven progression and exploratory sampling. This unifying principle offers a simple
yet powerful mechanism for balancing exploitation and exploration, adaptable across
ecological niches and timescales.

Together, these directions open exciting opportunities to link brain models of
sensorimotor control to peripheral control such as thoracic ganglia, refine our
understanding of decision timing, and extend the model to richer multi-goal or
multisensory environments.

942
943 **Materials and Methods**
944

Model Overview

We developed a computational model to test whether scanning behaviour dynamics
can emerge from the interaction between central complex (CX) steering and an
intrinsic lateral accessory lobe (LAL) oscillator. The model simulates single runs
where the virtual agent possesses a fixed goal direction in the CX and forward speed
will be inhibited at a predetermined time and released stochastically to trigger a
single scanning bout. The model was implemented in MATLAB R2016b, and code is
openly available at https://github.com/antnavteam/CX_oscillator_scans. The code is
extensively commented to allow readers to examine all implementation details,
parameter choices, and computational logic.

The model comprises three interacting components: (1) a CX steering circuit that
compares current heading with a goal direction to generate corrective signals
(adapted from Wystrach et al., 2020b), (2) an intrinsic LAL oscillator that produces
rhythmic left-right activity (adapted from Clément et al., 2023), and (3) motor output
rules that convert neural activity into angular and forward velocity.

Central Complex Steering Component

The CX steering architecture follows the computational framework established in
previous insect navigation models (Stone et al., 2017; Wystrach et al., 2020b), where
current and goal direction are encoded as bumps of activity across rings of 8
neurons, and compared via shifted representations to generate steering commands.

The CX encodes heading information as activity patterns across eight sectors with
fixed angular references every 45°: $\theta refs \; (i) \; = \; \{0, \pi/4, \pi/2, 3\pi/4, \pi, 5\pi/4, 3\pi/2, 7\pi/4\}$.
$4\}$.

Current heading is represented by a bump of activity across eight EPG (Ellipsoid
body-Protocerebral bridge-Gall) neurons, which track the agent heading (TH). The
goal direction is represented by a bump of activity across eight goal cells (here
termed HDB), which remains fixed at $\theta\_goal = \mathbf{0}$:

$$EPG(i, t) \; = \; ((\pi \; - \; |\pi_2\pi(\theta ref(i) \; - \; TH(t))|)^\wedge p) \, / \, \pi^\wedge p$$

$$HDB(i) \; = \; ((\pi \; - \; |\pi_2\pi(\theta ref(i) \; - \; \theta\_goal)|)^\wedge p) \, / \, \pi^\wedge p$$

In these equations, TH(t) represents the agent's current heading, θ_goal is the fixed
goal direction (which we set to 0), p (= 3) controls the width of the activity bump, and
$\pi_2\pi()$ wraps angles to [-π, +π]. Both EPG and HDB produce activity bumps centered
on their respective directions, with activity falling off smoothly away from the peak.

Two PFL (Protocerebral bridge-Fan-shaped body-Lateral accessory lobe)
populations compare spatially shifted copies of the heading representation with the
goal representation. Each PFL population computes the difference between the goal
signal and a shifted heading signal at each PB sector. Because neural activity under
inhibition is bounded at zero, negative values are prevented and set to zero:

$$PFL\_R(i, t) \; = \; max(\mathbf{0}, HDB(i) \; - \; EPG(i - \mathbf{1}, t))$$

$$PFL\_L(i, t) \; = \; max(\mathbf{0}, HDB(i) \; - \; EPG(i + \mathbf{1}, t))$$

where the indices i-1 and i+1 represent spatial shifts of ±1 sector (±45°). When the
goal representation is stronger than the (shifted) heading representation at a given
position, the PFL neuron becomes active; otherwise it is silent. This generates
asymmetric activity patterns: PFL_R becomes active when the goal lies to the right
of the agent's current heading, and PFL_L when the goal lies to the left of the
agent's heading. The summed activity across all sectors produces the bilateral CX
steering output:

$$CX\_out\_R(t) \; = \; \Sigma_i \, PFL\_R(i, t)$$

$$CX\_out\_L(t) \; = \; \Sigma_i \, PFL\_L(i, t)$$

The net difference between these outputs indicate whether the goal lies rather to the
right or the left of the current heading. Evidence in butterflies shows that corrective
output signals of the CX peaks when the insect is oriented 135° away from its goal
(Beetz et al., 2023). We implemented this by simply reversing the CX steering signal
(see below): evidence for the goal on the right will steer the agent towards the left
(and reciprocally), so that the agent will tend to move in the opposite direction (180°

away) from his CX goal representation, which effectively, acts here as an anti-goal.
As a result, the comparison of ±1 sector (±45°) shifted representation makes the
magnitude of CX_out peaks when current heading is shifted by 45° from the CX
goal, that is 135° (180° - 45°) from the agent goal (Figure 3). Note that this is
equivalent to a shift of ±3 sectors (±135°) with a non-reversed steering control. As a
result, the CX produces a stronger corrective steering signal when the agent is more
misoriented, a property consistent with behavioural data in ants and flies (Lent et al.,
2010; Westeinde et al., 2024).

To produce unambiguous unilateral steering signals, the two CX outputs go through
a cross hemispheric reciprocal inhibition (Figure 3), and then modulates the oscillator
via ipsilateral activation: CX_input_R for the right oscillator neuron R, and CX_input_L
for the left oscillator neuron L.

$$CX\_input\_R(t) = max(\mathbf{0}, \quad CX\_out\_R(t) - CX\_input\_L(t)) \cdot g\_CX$$

$$CX\_input\_L(t) = max(\mathbf{0}, \quad CX\_out\_L(t) - CX\_input\_R(t)) \cdot g\_CX$$

g_CX represents the CX gain parameter, that is, the strength of the CX impact on
the oscillator, which remains constant throughout each run, and whose impact on
scanning was investigated (Figure 6)

Intrinsic LAL Oscillator

Two reciprocally inhibiting neurons (R and L) generate oscillatory activity through
mutual inhibition and negative feedback representing synaptic exhaustion (Clément
et al., 2023):

$$R(t) = R(t-\mathbf{1}) - R\_NF(t-\mathbf{1}) - \alpha \cdot L(t-\mathbf{1}) + CX\_input\_R(t) + \xi\_osc$$

$$L(t) = L(t-\mathbf{1}) - L\_NF(t-\mathbf{1}) - \alpha \cdot R(t-\mathbf{1}) + CX\_input\_L(t) + \xi\_osc$$

α = 0.1 represents the reciprocal inhibition strength and ξ_osc ~ N(0, 0.05) is
Gaussian noise. The negative feedback (synaptic exhaustion) evolves as:

$$R\_NF(t) = R\_NF(t-\mathbf{1}) \cdot (\mathbf{1}-\beta) + \beta \cdot (R(t) - (R\_NF(t-\mathbf{1}) + s))$$

$$L\_NF(t) = L\_NF(t-\mathbf{1}) \cdot (\mathbf{1}-\beta) + \beta \cdot (L(t) - (L\_NF(t-\mathbf{1}) + s))$$

β = 0.01 represents the exhaustion rate and s = 0.5 is the steady-state firing rate.
Neural activity is constrained to [0, 2] to prevent runaway firing activity.

1029

Motor Output

Angular velocity (ω) and forward velocity (v) are derived from oscillator activity:

$$\omega(t) = (L(t) - R(t)) \cdot g\_ang \cdot (\mathbf{1}/(v(t - \mathbf{1}) + \mathbf{0.15})) + \xi\_motor$$

$$v(t) = ((L(t) + R(t))^\wedge k + \xi\_motor) \cdot g\_fwd$$

g_ang = 0.03 represents angular motor gain, g_fwd represents forward motor gain, k
controls how much forward speed varies with oscillator output, and ξ_motor ~ N(0,
0.1) is motor noise. The term (1/(v + 0.15)) implements the coupling between forward
and angular speed: faster forward movement reduces turning ability. +0.15 prevents
an infinite value when v=0.

To simulate species differences, we used two parameter sets:

Desert ants (e.g., *Cataglyphis* and *Melophorus*): k = 0.5, g_fwd = 1.0 (faster, smaller
impact of the oscillator on forward speed)

Myrmecia ants: k = 1.5, g_fwd = 0.2 (slower, larger impact of the oscillator on
forward speed)

The agent's heading and position update at each timestep:

$$TH(t) = TH(t - \mathbf{1}) + \omega(t)$$

$$X(t) = X(t - \mathbf{1}) + v(t) \cdot cos(TH(t))$$

$$Y(t) = Y(t - \mathbf{1}) + v(t) \cdot sin(TH(t))$$

Scan Generation via Forward Speed Inhibition

Forward speed can be inhibited to trigger scanning behaviour. We implemented two
inhibition modes:

Full inhibition (fwd_speed_inhibition = 0): Forward velocity is set to zero, halting
the locomotor central pattern generator (CPG). During this period, angular drive
$(L(t) - R(t))$ accumulates :

$$cumul\_drive(t) = cumul\_drive(t - \mathbf{1}) + (L(t) - R(t))$$

The agent remains in fixation (ω(t) = 0) until accumulated drive exceeds a threshold
to break the CPG inhibition θ_CPG = 2.0, then releases a saccade using the usual
formula for rotation:

$$\omega(t) = cumul\_drive(t) \cdot g\_ang \cdot (\mathbf{1}/(v(t) + \mathbf{0.15})) + \xi\_motor$$

$$cumul\_drive(t) = \mathbf{0}$$

This release resets the cumulated drive to zero, that is below CPG desinhibition
threshold, making the agent freeze and the angular drive cumulation resume. This

produces the characteristic scan structure of alternating fixations and saccades
observed in ants.

Partial inhibition (fwd_speed_inhibition > 0): Forward speed is reduced but not
halted (e.g., set to 0.2), and angular control proceeds normally without accumulation.
This generates broad oscillations or "voltes" rather than discrete scans.

Scan termination is stochastic: at each timestep during inhibition, forward speed
resumes with probability p_stop = 0.5. This produces exponentially distributed scan
durations matching the Poisson-like distribution observed in *M. bagoti* (Deeti et al.,
2023a).

Model Parameters

Parameter	Symbol	Desert Ants	Myrmecia	Justification
CX Parameters				
Bump width exponent	p	3	3	Set to produce realistic bump width (e.g., Kim et al., 2019)
CX output gain	g_CX	0.5	0.5	Effect studied in Figure 6
Oscillator Parameters				
Reciprocal inhibition	α	0.1	0.1	See Supp. Figure 2
Exhaustion rate	β	0.01	0.01	See Supp. Figure 2
Steady state	s	0.5	0.5	Neural oscillation require intrinsic activity
Oscillator noise	ξ_osc	0.05	0.05	Add realistic neural noise
Neural activity bounds	-	[0, 2]	[0, 2]	Prevents non-biologically realistic runaway activity
Motor Parameters				

Angular motor gain	g_ang	0.03	0.03	Fitted to match oscillation amplitude in continuous walking
Forward motor gain	g_fwd	1.0	0.2	Set to produce realistic speed difference across species
Forward oscillator control	k	0.5	1.5	Species-specific; controls oscillator's impacts on speed
Motor noise	ξ_motor	0.1	0.1	Adds realistic path variability

Scan Parameters

CPG angular threshold	θ_CPG	2.0	2.0	Fitted to produce realistic saccade duration
Scan termination probability	p_stop	0.05/step	0.05/step	Produces scan durations matching *M. bagoti* distribution
Forward speed during scan	v_scan	0	0	Full inhibition of forward speed

Simulation Protocol

Each simulation ran for 400 timesteps representing (given our oscillatory parameter
choice; (see Supplemental figure 2) around 6 to 10 oscillation cycles (depending on
CX strength). This would take ants around 5s to 25s (depending on species,
individual, temperature and other conditions), which is sufficient to encompass one
realistic scan duration. Initial conditions were: R(0) = 1, L(0) = 0, TH(0) = 0, position
(0, 0). Forward speed inhibition (i.e., scanning) began at timestep 200 (±50
timesteps random variation to ensure that scan onset occurs at various moments
along the oscillatory cycle). The model generated time series of:

● Position (X, Y) and heading (TH)
● Neural activities (R, L, CX_outputs)
● Velocities (angular ω, forward v)
● During scans: fixation durations, body orientations, saccade amplitudes

Model Outputs and behavioural Comparisons

From scanning periods (when v = 0), we extracted:

Fixation metrics: Duration (time between saccades), body orientation relative to
goal direction

Saccade metrics: Angular amplitude, direction (toward/away from goal), reversal
events (change in turning direction)

Scan structure: Total duration, number of fixations per scan, cumulative orientation
at first reversal

These were compared qualitatively against empirical data from *M. bagoti* using the
same metrics extracted from high-speed videos (see behavioural Experiments
section). We did not perform quantitative parameter optimization, as our goal was to
assess whether scanning properties naturally emerged from the proposed circuit
architecture rather than to fit the model precisely to data.

1098
1099   *Behavioural Experiments*

1100
1101   *Species and site*

1102
Natural scans were recorded using the red honeypot ant *Melophorus bagoti,* during
the Australian summer from December to February (experienced foragers -
Familiar/Unfamiliar conditions: 2010, see Deeti et al. 2023; inexperienced foragers –
Outbound/Inbound route formation conditions: 2023/2024 summer season) on a field
site (23°45′28.12″S, 133°52′59.77″E) located at the Centre for Appropriate Technology
campus, south of Alice Springs, Northern Territory, Australia. *M. bagoti* inhabit a
visually cluttered environment, exemplified by large amounts of buffel grass tussocks
(*Pennisetum cennchroides*), with scattered eucalyptus trees and bushes.

1111
Desert ants navigate alone using multiple concurrent, primarily visual, strategies
(Cheng et al., 2009; Collett and Collett, 2002; Freas and Spetch, 2018, 2023; Wehner,
2009; Wystrach et al., 2012; Zeil, 2023); including path-integration (Cheng et al., 2009;
Collett and Collett, 2002; Wehner and Srinivasan, 2003) and learned views around the
nest and along foraging routes (Collett, 2010; Freas et al., 2017; Wystrach et al., 2012;
Zeil, 2023; Zeil and Fleischmann, 2019).

Two data sets of high speed videos of scanning bouts were used in this project. The
first set of high speed scanning bout videos were collected using inexperienced
foragers, as individuals formed a straight-line route between a feeder and the nest. In
these ants, scanning bouts are known to have a high occurrence near the nest, in both
outbound and inbound ants. Videos were collected both at the onset of their *outbound*
journey or during the *inbound* trip just before reaching the goal. The second was a
group of highly experienced ants which were released either along or near their
established foraging route (familiar) or at a distant (unfamiliar) site (previously
collected in Deeti et al. 2023).

1127
1128   *Inexperienced forager procedure*

1129

High speed video recordings of inexperienced ants as they scanned while forming
their stereotypical route between and nest and feeder conducted on a single *M. bagoti*
nest in a closed arena with 10cm high walls enclosing the nest and feeder (Same
arena as Freas and Cheng, 2025, with a separate cohort of ants). A stocked feeder
was sunk into the ground 7m from the nest. As we were only interested in
inexperienced foragers with little knowledge of the area beyond the nest, prior to video
recording all foragers that emerged from the nest entrance were marked as
experienced using enamel paint (Tamiya™) and these experienced individuals were
excluded. On day six, non-painted foragers were allowed to exit the nest entrance and
find the feeder. Extending 0.2-1.0m from the nest entrance we spread a thin layer of
white sand along the ground. This sand helped the red ants stand out from the red
ground during filming and pose estimation analysis. Above this site, we positioned a
downward facing Chronos 2.1-HD highspeed camera (KRON Tech), 50cm above the
ground (1920×1080pixels, 600fps) with a field of view of 30cm×17cm.

1144

For five days prior to recording, one exposure to the feeder encouraged foragers to
leave the nest entrance in the general direction of the camera's field of view allowing
us to record any scanning behaviours during the first few foraging trips, as the
stereotypical route formed. After collecting ~25 individuals' outbound scans, we
switched to focusing on recently emerged foragers on their inbound trip collecting any
inbound scans, occurring near the nest. Once a forager completed a scan, or multiple
scans within the recording frame, they were collected after they left the area and
marked as completed to prevent repeated recording.

1153

1154   *Experienced foragers procedure*

1155

Experienced forager scans dynamics were extracted and analysed from a data set
published in Detti et al. (2023), which focused specifically at spanning metric
distributions; the following summary is further explained there. Two stocked feeders
(cookie pieces) were sunk into the ground 5m from the nest, separated 120° (Right
and Left sites). Ants exited each feeder via a 1m long, 10cm wide channel towards the
nest, slopped up to ground level where ants exited onto a 60 cm × 120 cm 'scanning
platform' (wooden board with white paper surface). Similar set-ups (minus the feeder)
were placed at three other local sites (Middle, Opposite, and Far).

1164

First, only the Right feeder was stocked with cookies and foragers were allowed to
train along this route, with each visiting individual marked upon their first visit.
Individuals were allowed to train along this route for two days prior to testing,
repeatedly returning to the feeder and leaving via the channel with food. Once highly
experienced, inbound foragers were tested by collection just before entering the nest
and released at one of the four familiar test sites. As each individual reached the
scanning platform they were video recorded using a high-speed camera (Casio EX-
F1, FOV: 30×30 cm at 300 fps). This testing procedure was then repeated with a
second group of foragers trained to the Left feeder. A final set up was placed in a far
(40m), unfamiliar site where a separate group of experienced foragers was tested.

1175

1176   *Pose extraction*

1177

For both sets of scan video datasets, we extracted the body orientations of ants during fixations to calculate their angle compared to the goal direction (e.g., nest or feeder) and their duration. Fixations were defined as periods when the ant's head and body remained stationary between video frames, with no forward or rotational movement. Scanning can be composed of multiple scanning bouts, with little to no forward movement within a bout, while the ant rotates in place (saccades), or fixating in different directions, with a scanning bout ceasing once the ant resumes forward motion (Deeti et al., 2023a). By collecting body orientation during fixations, we extracted, the duration of each fixation, the ant's *body orientation* relative to the goal, the angular change between each fixation (saccade angle) and duration, if the saccade was 'towards' or 'away' from the goal direction and when the fixation was followed by saccade that was a reversal in turning direction.

In the Experienced individuals (300 frames/s), orientation was extracted using a custom MATLAB code (see Deeti et al., 2023 for full description), with fixations identified as stationary periods between frames with body orientation estimated using points at the head and pronotum. In inexperienced forager *f* scans (600frames/s), body orientation during fixations were determined manually using SLEAP software (Pereira et al., 2022). Fixations 'started' when the head stopped moving between frames and ended with the onset of either forward or rotational movement. Orientation was calculated using two body landmarks: the front centre of the head and the head/body connection (pronotum). As multiple scanning bouts can occur within a single path, each scanning bout began with the first, within frame fixation and ended after forward movement resumed for at least two full leg cycles. Additional fixations within this two-step window were still counted as within the same scanning bout.

*Statistical analysis (*real world *Melophorus bagoti scans)*

    Grouping

For experienced foragers, scan data from all familiar release sites (Left, Right, Middle, Opposite) were pooled into a single familiar condition, while scans from the distant site were treated as unfamiliar. For inexperienced foragers, scans were separated into two groups based on if the scan occurred during the outbound and inbound portion of the foraging trip.

    Linear mixed-effects models (LMEs)

LME models (Matlab2016) were used to assess the qualitative predictions of the model compared to our real ant data, rather than to optimize model fit. Accordingly, we report F-values derived from ANOVA tables, which test the significance of fixed effects rather than relying on t-values associated with individual parameter estimates. Importantly, this approach fits our qualitative focus. We first run models with interaction between factors. If no interaction was significant, we re-run the model for additive effect only.

Effects on *saccade aptitude* and *fixation duration* were analysed using Linear mixed-effect models with both *individual* and *test conditions* as random effects and with *angular deviation* from the goal direction as a continuous fixed effect and with reversal

1227 and Towards/Away (for fixations this corresponds with the upcoming saccade) as
1228 categorical fixed effects.
1229
To assess the effect of experience on cumulative orientation vs. the goal during the
first reversal of each scan, we conducted LME with the *cumulative orientation* as the
dependent variable, with *condition* as the fixed effect and *individual* as a random effect
to account for repeated measures across individuals. We ran a second LME to assess
the effect of *experience* on the *saccade amplitude*, with *condition* as the fixed effect
and *individual* as the random effect to account for repeated measures across
individuals.
1237
1238   *Fixation duration & saccade amplitude - linear regression*
1239
1240 To assess the relationship between fixation duration and saccade amplitude, we first
1241 plotted the entire dataset. After observing the general trends, we separated shorter
1242 fixation durations (Quartile 1) from the rest of the dataset (Quartile 2-4) and performed
1243 separate linear regressions for both subsets. Regression slopes were then tested for
1244 significance using *t*-tests on the fitted model coefficients.
1245
1246   *Start/stop anytime during oscillation cycle*
1247
1248 To compare the duration via the number of saccades within a sweep between distinct
1249 sweeps (e.g., first vs. second sweep; penultimate vs. final sweep), we used the
1250 Wilcoxon signed-rank tests for paired data.
1251
1252   *First reversal/longest fixation - 'binomial' chi square association*
1253
1254 To assess if reversal and longest fixation duration were associated we tested them
1255 using the chi-square test of independence (χ^2), and odds ratios with 95% confidence
1256 intervals were calculated.
1257
1258

1259 **Acknowledgements**
1260 We are grateful to the Centre for Appropriate Technology for permission to work on
1261 site and access to the nests. We thank Paul Graham for providing the experienced
1262 forager scan dataset, and Leo Clément and Gabriel G. Gattaux for contributing videos
1263 of ant behaviour.
1264
1265 **Funding Statement**
1266 The funders had no role in study design, data collection and interpretation, or the
1267 decision to submit the work for publication.
1268
1269 **Funding Information**
1270 This project was funded by a Macquarie University Research Fellowship
1271 (MQRF0001094) and by the European Research Council (RESIL-ANT - 101125881).
1272
1273 **Competing interests**
1274 No competing interests declared.

1275
1276 **Contributions**

1277 C.A.F. - Conceptualization, Data curation, Formal analysis, Funding acquisition,
1278 Validation, Investigation, Visualization, Methodology, Writing.

1279 A.W.   -Conceptualization, Data curation, Formal analysis, Modelling, Funding
1280 acquisition, Validation, Investigation, Visualization, Methodology, Writing.

Data Availability Statement

1282 All data, documentation and code is made available online at:
1283 (https://github.com/antnavteam/CX_oscillator_scans')

References

Adden A, Stewart TC, Webb B, Heinze S. 2022. A Neural Model for Insect
Steering Applied to Olfaction and Path Integration. *Neural Comput* **34**:2205–
2231. doi:10.1162/neco_a_01540

Baddeley B, Graham P, Husbands P, Philippides A. 2012. A Model of Ant
Route Navigation Driven by Scene Familiarity. *PLoS Comput Biol*
8:e1002336. doi:10.1371/journal.pcbi.1002336

Baird E, Byrne MJ, Smolka J, Warrant EJ, Dacke M. 2012. The Dung Beetle
Dance: An Orientation Behaviour? *PLOS ONE* **7**:e30211.
doi:10.1371/journal.pone.0030211

Beetz MJ, Kraus C, el Jundi B. 2023. Neural representation of goal direction
in the monarch butterfly brain. *Nat Commun* **14**:5859. doi:10.1038/s41467-
023-41526-w

Berni J. 2015. Genetic Dissection of a Regionally Differentiated Network for
Exploratory Behavior in *Drosophila* Larvae. *Curr Biol* **25**:1319–1326.
doi:10.1016/j.cub.2015.03.023

Berni J, Pulver SR, Griffith LC, Bate M. 2012. Autonomous Circuitry for
Substrate Exploration in Freely Moving *Drosophila* Larvae. *Curr Biol* **22**:1861–
1870. doi:10.1016/j.cub.2012.07.048

Bidaye SS, Laturney M, Chang AK, Liu Y, Bockemühl T, Büschges A, Scott K.
2020. Two Brain Pathways Initiate Distinct Forward Walking Programs in
Drosophila. *Neuron* **108**:469-485.e8. doi:10.1016/j.neuron.2020.07.032

Buehlmann C, Fernandes ASD, Graham P. 2018. The interaction of path integration and terrestrial visual cues in navigating desert ants: what can we learn from path characteristics? *J Exp Biol* **221**:jeb167304. doi:10.1242/jeb.167304

Buehlmann C, Mangan M, Graham P. 2020. Multimodal interactions in insect navigation. *Anim Cogn* **23**:1129–1141. doi:10.1007/s10071-020-01383-2

Büschges A, Ache JM. 2025. Motor control on the move: from insights in insects to general mechanisms. *Physiol Rev* **105**:975–1031. doi:10.1152/physrev.00009.2024

Cheng K. 2024. Oscillators and servomechanisms in navigation and orientation. *Commun Integr Biol* **17**:2293268. doi:10.1080/19420889.2023.2293268

Cheng K, Narendra A, Sommer S, Wehner R. 2009. Traveling in clutter: navigation in the Central Australian desert ant Melophorus bagoti. *Behav Processes* **80**:261–268.

Clément L, Schwarz S, Wystrach A. 2023. An intrinsic oscillator underlies visual navigation in ants. *Curr Biol* **33**:411–422. doi:10.1016/j.cub.2022.11.059

Collett M. 2010. How desert ants use a visual landmark for guidance along a habitual route. *Proc Natl Acad Sci USA* **107**:11638–11643. doi:10.1073/pnas.1001401107

Collett T, Collett M. 2002. Memory use in insect visual navigation. *Nat Rev Neurosci* **3**:542–52. doi:10.1038/nrn872

Collett T, Graham P, Heinze S. 2025. The neuroethology of ant navigation. *Curr Biol* **35**:R110–R124. doi:10.1016/j.cub.2024.12.034

Collett TS, Hempel De Ibarra N. 2023. An 'instinct for learning': the learning flights and walks of bees, wasps and ants from the 1850s to now. *J Exp Biol* **226**:jeb245278. doi:10.1242/jeb.245278

Collett, TS, Hempel De Ibarra N, Riabinina O, Philippides A. 2013. Coordinating compass-based and nest-based flight directions during bumblebee learning and return flights. Journal of Experimental Biology, **216**(6), 1105-1113.

Collett TS, Robert T, Frasnelli E, Philippides A, Hempel De Ibarra N. 2023. How bumblebees coordinate path integration and body orientation at the start of their first learning flight. *J Exp Biol* **226**:jeb245271. doi:10.1242/jeb.245271

Dauzere-Peres O, Wystrach A. 2024. Ants integrate proprioception as well as visual context and efference copies to make robust predictions. *Nat Commun* **15**:10205. doi:10.1038/s41467-024-53856-4

Dauzere-Peres O, De Wever S, Wystrach A. 2026. Predictive coding and oscillations underlie the optomotor response in distant insect lineages. bioRxiv, 2026-03.

Deeti S, Cheng K. 2025. Desert ants (*Melophorus bagoti*) oscillate and scan more in navigation when the visual scene changes. *Anim Cogn* **28**:15. doi:10.1007/s10071-025-01936-3

Deeti S, Cheng K, Graham P, Wystrach A. 2023a. Scanning behaviour in ants: an interplay between random-rate processes and oscillators. *J Comp Physiol A* **209**:625–639. doi:10.1007/s00359-023-01628-8

Deeti S, Islam M, Freas C, Murray T, Cheng K. 2023b. Intricacies of running a route without success in night-active bull ants (*Myrmecia midas*). *J Exp Psychol Anim Learn Cogn* **49**:111–126. doi:10.1037/xan0000350

Demir M, Kadakia N, Anderson HD, Clark DA, Emonet T. 2020. Walking *Drosophila* navigate complex plumes using stochastic decisions biased by the timing of odor encounters. *eLife* **9**:e57524. doi:10.7554/eLife.57524

Dudley R. 2000. The Biomechanics of Insect Flight: Form, Function, Evolution. Princeton University Press.

Fisher YE. 2022. Flexible navigational computations in the *Drosophila* central complex. *Curr Opin Neurobiol*, **73**:102514.

Fleischmann PN, Grob R, Wehner R, Rössler W. 2017. Species-specific differences in the fine structure of learning walk elements in Cataglyphis ants. *J Exp Biol* **220**:2426–2435. doi:10.1242/jeb.158147

Franconville R, Beron C, Jayaraman V. 2018. Building a functional connectome of the *Drosophila* central complex. *eLife* **7**:e37017. doi:10.7554/eLife.37017

Freas CA, Cheng K. 2018. Landmark learning, cue conflict, and outbound
view sequence in navigating desert ants. *Journal of Experimental Psychology:*
Animal Learning and Cognition, **44**(4): 409–421.

Freas CA, Cheng K. 2025. Visual learning, route formation and the
choreography of looking back in desert ants, *Melophorus bagoti*. *Anim Behav*
222:123125.

Freas CA, Cheng K. 2022. The Basis of Navigation Across Species. *Annu*
Rev Psychol **73**:217–241. doi:10.1146/annurev-psych-020821-111311

Freas CA, Congdon JV, Plowes NJR, Spetch ML. 2020. Pheromone cue
triggers switch between vectors in the desert harvest ant, Veromessor
pergandei. *Anim Cogn* **23**:1087–1105. doi:10.1007/s10071-020-01354-7

Freas CA, Fleischmann PN, Cheng K. 2019. Experimental ethology of
learning in desert ants: Becoming expert navigators. *Behav Processes*
158:181–191. doi:10.1016/j.beproc.2018.12.001

Freas CA, Spetch ML. 2019. Terrestrial cue learning and retention during the
outbound and inbound foraging trip in the desert ant, Cataglyphis velox.
Journal of Comparative Physiology A. **205**(2):177-89.

Freas CA, Spetch ML. 2023. Varieties of visual navigation in insects. *Anim*
Cogn **26**:319–342. doi:10.1007/s10071-022-01720-7

Freas CA, Whyte C, Cheng K. 2017. Skyline retention and retroactive
interference in the navigating Australian desert ant, Melophorus bagoti. *J*
Comp Physiol A Neuroethol Sens Neural Behav Physiol **203**:353–367.
doi:10.1007/s00359-017-1174-8

Freas CA, Wystrach A, Narendra A, Cheng K. 2018. The View from the Trees:
Nocturnal Bull Ants, Myrmecia midas, Use the Surrounding Panorama While
Descending from Trees. *Front Psychol* **9**. doi:10.3389/fpsyg.2018.00016

Freas CA, Wystrach A, Schwarz S, Spetch ML. 2022. Aversive view
memories and risk perception in navigating ants. *Sci Rep* **12**:2899.

Green J, Maimon G. 2018. Building a heading signal from anatomically
defined neuron types in the *Drosophila* central complex. Current opinion in
neurobiology, 52, 156-164.

Gattaux G, Vimbert R, Wystrach A, Serres JR, Ruffier F. 2023. Antcar: Simple
Route Following Task with Ants-Inspired Vision and Neural Model.

Gomez-Marin A, Louis M. 2014. Multilevel control of run orientation in
Drosophila larval chemotaxis. *Front Behav Neurosci* **8**.
doi:10.3389/fnbeh.2014.00038

Gomez-Marin A, Stephens GJ, Louis M. 2011. Active sampling and decision
making in *Drosophila* chemotaxis. *Nat Commun* **2**:441.
doi:10.1038/ncomms1455

Goulard, R, Buehlmann, C, Niven, JE, Graham, P, Webb, B. 2021. A unified
mechanism for innate and learned visual landmark guidance in the insect
central complex. PLoS computational biology, **17**(9): e1009383.

Goulard R, Heinze S, Webb B. 2023. Emergent spatial goals in an integrative
model of the insect central complex. *PLOS Comput Biol* **19**:e1011480.
doi:10.1371/journal.pcbi.1011480

Graham P, Collett TS. 2002. View-based navigation in insects: how wood ants
(*Formica rufa* L.) look at and are guided by extended landmarks. *J Exp Biol*
205:2499–2509. doi:10.1242/jeb.205.16.2499

Green J, Vijayan V, Mussells Pires P, Adachi A, Maimon G. 2019. A neural
heading estimate is compared with an internal goal to guide oriented
navigation. *Nat Neurosci* **22**:1460–1468. doi:10.1038/s41593-019-0444-x

Haalck L, Mangan M, Wystrach A, Clement L, Webb B, Risse B. 2023.
CATER: Combined Animal Tracking & Environment Reconstruction. *Sci Adv*
9:eadg2094. doi:10.1126/sciadv.adg2094

Hangartner W. 1967. Spezifitat und Inaktivierung des Spurpheromons von
Lasius fuliginosus Latr. und Orientierung der Arbeiterinnen im Duftfeld. *Z Vgl*
Physiol **57**:103–136. doi:10.1007/BF00303068

Heinze S. 2017. Unraveling the neural basis of insect navigation. *Curr Opin*
Insect Sci **24**:58–67. doi:10.1016/j.cois.2017.09.001

Honkanen A, Adden A, da Silva Freitas J, Heinze S. 2019. The insect central
complex and the neural basis of navigational strategies. *J Exp Biol*
222:jeb188854. doi:10.1242/jeb.188854

Hulse BK, Haberkern H, Franconville R, Turner-Evans D, Takemura S, Wolff T, Noorman M, Dreher M, Dan C, Parekh R, Hermundstad AM, Rubin GM, Jayaraman V. 2021. A connectome of the *Drosophila* central complex reveals network motifs suitable for flexible navigation and context-dependent action selection. *eLife* **10**:e66039. doi:10.7554/eLife.66039

Husbands P, Shim Y, Garvie M, Dewar A, Domcsek N, Graham P, Knight J, Nowotny T, Philippides A. 2021. Recent advances in evolutionary and bio-inspired adaptive robotics: Exploiting embodied dynamics. *Appl Intell* **51**:6467–6496. doi:10.1007/s10489-021-02275-9

Islam M, Freas CA, Cheng K. 2020. Effect of large visual changes on the navigation of the nocturnal bull ant, Myrmecia midas | Animal Cognition. *Anim Cogn* **23**:1071–1080.

Iwano M, Hill ES, Mori A, Mishima Tatsuya, Mishima Tsuneko, Ito K, Kanzaki R. 2010. Neurons associated with the flip-flop activity in the lateral accessory lobe and ventral protocerebrum of the silkworm moth brain. *J Comp Neurol* **518**:366–388. doi:10.1002/cne.22224

Izquierdo EJ, Lockery SR. 2010. Evolution and Analysis of Minimal Neural Circuits for Klinotaxis in *Caenorhabditis elegans*. *J Neurosci* **30**:12908–12917. doi:10.1523/JNEUROSCI.2606-10.2010

Kanzaki R. 2005. Neural Basis of Odor-source Searching Behavior in Insect Brain Systems Evaluated with a Mobile Robot. *Chem Senses* **30**:i285–i286. doi:10.1093/chemse/bjh226

Kanzaki, R., & Ikeda, A. (1994). Morphology and physiology of pheromone-triggered flip-flopping descending interneurons of the male silkworm moth, Bombyx mori. In *Olfaction and Taste XI: Proceedings of the 11th International Symposium on Olfaction and Taste and of the 27th Japanese Symposium on Taste and Smell. Joint Meeting held at Kosei-nenkin Kaikan, Sapporo, Japan, July 12–16, 1993* (pp. 851-851). Tokyo: Springer Japan.

Kanzaki R, Mishima T. 1996. Pheromone-Triggered 'Fiipflopping' Neural Signals Correlate with Activities of Neck Motor Neurons of a Male Moth, Bombyx mori. *Zoolog Sci* **13**:79–87. doi:10.2108/zsj.13.79

Kanzaki R, Nagasawa S, Shimoyama I. 2004. Neural Basis of Odor-Source Searching Behavior in insect Microbrain Systems Evaluated with a Mobile Robot In: Kato N, Ayers J, Morikawa H, editors. Bio-Mechanisms of Swimming and Flying. Tokyo: Springer Japan. pp. 155–170. doi:10.1007/978-4-431-53951-3_12

Kanzaki R, Sugi N, Shibuya T. 1992. Self-generated Zigzag Turning of Bombyx mori Males during Pheromone-mediated Upwind Walking(Physology). *Zoolog Sci* **9**:515–527.

Kim SS, Hermundstad AM, Romani S, Abbott LF, Jayaraman V. 2019. Generation of stable heading representations in diverse visual scenes. *Nature* **576**:126–131. doi:10.1038/s41586-019-1767-1

Knaden M, Graham P. 2016. The Sensory Ecology of Ant Navigation: From Natural Environments to Neural Mechanisms. *Annu Rev Entomol* **61**:63–76. doi:10.1146/annurev-ento-010715-023703

Knight JC, Sakhapov D, Domcsek N, Dewar ADM, Graham P, Nowotny T, Philippides A. 2019. Insect-Inspired Visual Navigation On-Board an Autonomous Robot: Real-World Routes Encoded in a Single Layer NetworkThe 2019 Conference on Artificial Life. Presented at the The 2019 Conference on Artificial Life. Newcastle, United Kingdom: MIT Press. pp. 60–67. doi:10.1162/isal_a_00141

Kodzhabashev A, Mangan M. 2015. Route Following Without Scanning In: Wilson SP, Verschure PFMJ, Mura A, Prescott TJ, editors. Biomimetic and Biohybrid Systems. Cham: Springer International Publishing. pp. 199–210. doi:10.1007/978-3-319-22979-9_20

Kuenen LPS, Baker TC. 1983. A non-anemotactic mechanism used in pheromone source location by flying moths. *Physiol Entomol* **8**:277–289. doi:10.1111/j.1365-3032.1983.tb00360.x

Legge, E. L., Wystrach, A., Spetch, M. L., & Cheng, K. (2014). Combining sky and earth: desert ants (Melophorus bagoti) show weighted integration of celestial and terrestrial cues. Journal of Experimental Biology, 217(23), 4159-4166.

Le Moël F, Stone T, Lihoreau M, Wystrach A, Webb B. 2019. The Central Complex as a Potential Substrate for Vector Based Navigation. *Front Psychol* **10**:690. doi:10.3389/fpsyg.2019.00690

Le Möel F, Wystrach A. 2020. Opponent processes in visual memories: A model of attraction and repulsion in navigating insects' mushroom bodies. *PLOS Comput Biol* **16**:e1007631. doi:10.1371/journal.pcbi.1007631

Lehrer M. 1993. Why do bees turn back and look? *J Comp Physiol A* **172**:549–563. doi:10.1007/BF00213678

Lent DavidD, Graham P, Collett TS. 2010. Image-matching during ant navigation occurs through saccade-like body turns controlled by learned visual features. *Proc Natl Acad Sci* **107**:16348–16353. doi:10.1073/pnas.1006021107

Li F, Lindsey JW, Marin EC, Otto N, Dreher M, Dempsey G, Stark I, Bates AS, Pleijzier MW, Schlegel P, Nern A, Takemura S, Eckstein N, Yang T, Francis A, Braun A, Parekh R, Costa M, Scheffer LK, Aso Y, Jefferis GS, Abbott LF, Litwin-Kumar A, Waddell S, Rubin GM. 2020. The connectome of the adult *Drosophila* mushroom body provides insights into function. *eLife* **9**:e62576. doi:10.7554/eLife.62576

Lu J, Behbahani AH, Hamburg L, Westeinde EA, Dawson PM, Lyu C, Maimon G, Dickinson MH, Druckmann S, Wilson RI. 2022. Transforming representations of movement from body- to world-centric space. *Nature* **601**:98–104. doi:10.1038/s41586-021-04191-x

Luo L, Gershow M, Rosenzweig M, Kang K, Fang-Yen C, Garrity PA, Samuel ADT. 2010. Navigational Decision Making in *Drosophila* Thermotaxis. *J Neurosci* **30**:4261–4272. doi:10.1523/JNEUROSCI.4090-09.2010

Lyu C, Abbott LF, Maimon G. 2022. Building an allocentric travelling direction signal via vector computation. *Nature* **601**:92–97. doi:10.1038/s41586-021-04067-0

Mishima T, Kanzaki R. 1999. Physiological and morphological characterization of olfactory descending interneurons of the male silkworm moth, Bombyx mori. *J Comp Physiol [A]* **184**:143–160. doi:10.1007/s003590050314

Mouritsen H, Feenders G, Liedvogel M, Kropp W. 2004. Migratory Birds Use Head Scans to Detect the Direction of the Earth's Magnetic Field. *Curr Biol* **14**:1946–1949. doi:10.1016/j.cub.2004.10.025

Müller M, Wehner R. 2010. Path Integration Provides a Scaffold for Landmark Learning in Desert Ants. *Curr Biol* **20**:1368–1371. doi:10.1016/j.cub.2010.06.035

Müller M, Wehner R. 2007. Wind and sky as compass cues in desert ant navigation. *Naturwissenschaften* **94**:589–594. doi:10.1007/s00114-007-0232-4

Murakami M, Vicente MI, Costa GM, Mainen ZF. 2014. Neural antecedents of self-initiated actions in secondary motor cortex. *Nat Neurosci* **17**:1574–1582. doi:10.1038/nn.3826

Murray T, Kocsi Z, Dahmen H, Narendra A, Le Möel F, Wystrach A, Zeil J. 2019. The role of attractive and repellent scene memories in ant homing (*Myrmecia croslandi*). *J Exp Biol* jeb.210021. doi:10.1242/jeb.210021

Muser B, Sommer S, Wolf H, Wehner R. 2005. Foraging ecology of the thermophilic Australian desert ant, Melophorus bagoti. *Aust J Zool* **53**:301. doi:10.1071/ZO05023

Mussells Pires P, Zhang L, Parache V, Abbott LF, Maimon G. 2024. Converting an allocentric goal into an egocentric steering signal. *Nature* **626**:808–818. doi:10.1038/s41586-023-07006-3

Namiki S, Iwabuchi S, Pansopha Kono P, Kanzaki R. 2014. Information flow through neural circuits for pheromone orientation. *Nat Commun* **5**:5919. doi:10.1038/ncomms6919

Namiki S, Kanzaki R. 2016. The neurobiological basis of orientation in insects: insights from the silkmoth mating dance. *Curr Opin Insect Sci* **15**:16–26. doi:10.1016/j.cois.2016.02.009

Nicholson DJ, Judd SPD, Cartwright BA, Collett TS. 1999. Learning walks and landmark guidance in wood ants (Formica rufa). *J Exp Biol* **202**:1831–1838. doi:10.1242/jeb.202.13.1831

Olberg RM. 1983. Pheromone-triggered flip-flopping interneurons in the ventral nerve cord of the silkworm moth,Bombyx mori. *J Comp Physiol A* **152**:297–307. doi:10.1007/BF00606236

Pereira TD, Tabris N, Matsliah A, Turner DM, Li J, Ravindranath S, Papadoyannis ES, Normand E, Deutsch DS, Wang ZY, McKenzie-Smith GC, Mitelut CC, Castro MD, D'Uva J, Kislin M, Sanes DH, Kocher SD, Wang SS-H, Falkner AL, Shaevitz JW, Murthy M. 2022. SLEAP: A deep learning system for multi-animal pose tracking. *Nat Methods* **19**:486–495. doi:10.1038/s41592-022-01426-1

Pfeiffer K, Homberg U. 2014. Organization and Functional Roles of the Central Complex in the Insect Brain. *Annu Rev Entomol* **59**:165–184. doi:10.1146/annurev-ento-011613-162031

Philippides, A, de Ibarra, NH, Riabinina, O, Collett, TS, 2013. Bumblebee calligraphy: the design and control of flight motifs in the learning and return flights of *Bombus terrestris*. *Journal of Experimental Biology,* **216**(6), 1093-1104.

Rayshubskiy A, Holtz SL, Bates AS, Vanderbeck QX, Serratosa Capdevila L, Rockwell V, Wilson R. 2025. Neural circuit mechanisms for steering control in walking *Drosophila. eLife* **13**:RP102230. doi:10.7554/eLife.102230.3

Robert T, Frasnelli E, De Ibarra NH, Collett TS. 2018. Variations on a theme: bumblebee learning flights from the nest and from flowers. *J Exp Biol* jeb.172601. doi:10.1242/jeb.172601

Sapkal N, Mancini N, Kumar DS, Spiller N, Murakami K, Vitelli G, Bargeron B, Maier K, Eichler K, Jefferis GSXE, Shiu PK, Sterne GR, Bidaye SS. 2024. Neural circuit mechanisms underlying context-specific halting in *Drosophila. Nature* **634**:191–200. doi:10.1038/s41586-024-07854-7

Schwarz S, Clement L, Gkanias E, Wystrach A. 2020a. How do backward-walking ants (Cataglyphis velox) cope with navigational uncertainty? *Anim Behav* **164**:133–142. doi:10.1016/j.anbehav.2020.04.006

Schwarz S, Mangan M, Webb B, Wystrach A. 2020b. Route-following ants respond to alterations of the view sequence. *J Exp Biol* **223**:jeb218701. doi:10.1242/jeb.218701

Seelig JD, Jayaraman V. 2015. Neural dynamics for landmark orientation and angular path integration. *Nature* **521**:186–191. doi:10.1038/nature14446

Shih C-T, Sporns O, Yuan S-L, Su T-S, Lin Y-J, Chuang C-C, Wang T-Y, Lo C-C, Greenspan RJ, Chiang A-S. 2015. Connectomics-Based Analysis of Information Flow in the *Drosophila* Brain. *Curr Biol* **25**:1249–1258. doi:10.1016/j.cub.2015.03.021

Shiu PK, Sterne GR, Spiller N, Franconville R, Sandoval A, Zhou J, Simha N, Kang CH, Yu Seongbong, Kim JS, Dorkenwald S, Matsliah A, Schlegel P, Yu Szi-chieh, McKellar CE, Sterling A, Costa M, Eichler K, Bates AS, Eckstein N, Funke J, Jefferis GSXE, Murthy M, Bidaye SS, Hampel S, Seeds AM, Scott K. 2024. A *Drosophila* computational brain model reveals sensorimotor processing. *Nature* **634**:210–219. doi:10.1038/s41586-024-07763-9

Steck K, Hansson BS, Knaden M. 2011. Desert ants benefit from combining visual and olfactory landmarks. *J Exp Biol* **214**:1307–1312. doi:10.1242/jeb.053579

Steinbeck F, Adden A, Graham P. 2020. Connecting brain to behaviour: a role
for general purpose steering circuits in insect orientation? *J Exp Biol*
223:jeb212332. doi:10.1242/jeb.212332

Stone T, Webb B, Adden A, Weddig NB, Honkanen A, Templin R, Wcislo W,
Scimeca L, Warrant E, Heinze S. 2017. An Anatomically Constrained Model
for Path Integration in the Bee Brain. *Curr Biol* **27**:3069-3085.e11.
doi:10.1016/j.cub.2017.08.052

Stürzl W, Zeil J, Boeddeker N, Hemmi JM. 2016. How Wasps Acquire and
Use Views for Homing. *Curr Biol* **26**:470–482. doi:10.1016/j.cub.2015.12.052

Sun X, Yue S, Mangan M. 2020. A decentralised neural model explaining
optimal integration of navigational strategies in insects. *eLife* **9**:e54026.
doi:10.7554/eLife.54026

Tarsitano MS, Andrew R. 1999. Scanning and route selection in the jumping
spider *Portia labiata*. *Anim Behav* **58**:255–265. doi:10.1006/anbe.1999.1138

Tastekin I, Khandelwal A, Tadres D, Fessner ND, Truman JW, Zlatic M,
Cardona A, Louis M. 2018. Sensorimotor pathway controlling stopping
behavior during chemotaxis in the *Drosophila melanogaster* larva. *eLife*
7:e38740. doi:10.7554/eLife.38740

Ugolini A. 2006. Equatorial sandhoppers use body scans to detect the earth's
magnetic field. *J Comp Physiol A* **192**:45–49. doi:10.1007/s00359-005-0046-9

Wallace GK. 1959. Visual Scanning in the Desert Locust Schistocerca
Gregaria Forskål. *J Exp Biol* **36**:512–525. doi:10.1242/jeb.36.3.512

Webb B, Wystrach A. 2016. Neural mechanisms of insect navigation. *Curr
Opin Insect Sci* **15**:27–39. doi:10.1016/j.cois.2016.02.011

Wehner R. 2009. The architecture of the desert ant's navigational toolkit
(Hymenoptera: Formicidae). *Myrmecol News* **12**:85–96.

Wehner R, Fukushi T, Wehner S. 1992. Rotatory components of movement in
high speed desert ants, Cataglyphis bombycina. Presented at the
Proceedings of the 20th Göttingen Neurobiology Conference.

Wehner R, Hoinville T, Cruse H, Cheng K. 2016. Steering intermediate
courses: desert ants combine information from various navigational routines. *J
Comp Physiol A* **202**:459–472. doi:10.1007/s00359-016-1094-z

Wehner R, Michel B, Antonsen P. 1996. Visual Navigation in Insects: Coupling of Egocentric and Geocentric Information. *J Exp Biol* **199**:129–140. doi:10.1242/jeb.199.1.129

Wehner R, Srinivasan MV. 2003. Path Integration in Insects. Oxford University Press.

Westeinde EA, Kellogg E, Dawson PM, Lu J, Hamburg L, Midler B, Druckmann S, Wilson RI. 2024. Transforming a head direction signal into a goal-oriented steering command. *Nature* **626**:819–826. doi:10.1038/s41586-024-07039-2

Wystrach A. 2023. Neurons from pre-motor areas to the Mushroom bodies can orchestrate latent visual learning in navigating insects. doi:10.1101/2023.03.09.531867

Wystrach A, Beugnon G, Cheng K. 2012. Ants might use different view-matching strategies on and off the route. *J Exp Biol* **215**:44–55. doi:10.1242/jeb.059584

Wystrach A, Buehlmann C, Schwarz S, Cheng K, Graham P. 2020a. Rapid Aversive and Memory Trace Learning during Route Navigation in Desert Ants. *Curr Biol* **30**:1927-1933.e2. doi:10.1016/j.cub.2020.02.082

Wystrach A, Lagogiannis K, Webb B. 2016. Continuous lateral oscillations as a core mechanism for taxis in *Drosophila* larvae. *eLife* **5**:e15504. doi:10.7554/eLife.15504

Wystrach A, Mangan M, Philippides A, Graham P. 2013. Snapshots in ants? New interpretations of paradigmatic experiments. *J Exp Biol* jeb.082941. doi:10.1242/jeb.082941

Wystrach A, Mangan M, Webb B. 2015. Optimal cue integration in ants. *Proc Biol Sci* **282**:20151484. doi:10.1098/rspb.2015.1484

Wystrach A, Moël FL, Clement L, Schwarz S. 2020b. A lateralised design for the interaction of visual memories and heading representations in navigating ants. doi:10.1101/2020.08.13.249193

Wystrach A, Philippides A, Aurejac A, Cheng K, Graham P. 2014. Visual scanning behaviours and their role in the navigation of the Australian desert ant Melophorus bagoti. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **200**:615–626. doi:10.1007/s00359-014-0900-8

Wystrach A, Schwarz S. 2013. Ants use a predictive mechanism to
compensate for passive displacements by wind. *Curr Biol* **23**:R1083–R1085.
doi:10.1016/j.cub.2013.10.072

Yang HH, Brezovec BE, Serratosa Capdevila L, Vanderbeck QX, Adachi A,
Mann RS, Wilson RI. 2024. Fine-grained descending control of steering in
walking *Drosophila*. *Cell* **187**:6290-6308.e27. doi:10.1016/j.cell.2024.08.033

Zeil J. 2023. Visual navigation: properties, acquisition and use of views. *J
Comp Physiol A* **209**:499–514. doi:10.1007/s00359-022-01599-2

Zeil J, Fleischmann PN. 2019. The learning walks of ants (Hymenoptera:
Formicidae). *Myrmecol News* **29**:93–110.
doi:10.25849/MYRMECOL.NEWS_029:093

Zeil J, Kelber A, Voss R. 1996. Structure and function of learning flights in
bees and wasps. Journal of experimental biology, 199(1), 245-252.