Scanning and active sampling behaviours emerge from conserved insect neural circuits

Cody A Freas[1,2], Antoine Wystrach[1]

[1]Research Center on Animal Cognition (CRCA), Center for Integrative Biology (CBI), CNRS, University of Toulouse, Toulouse, France
[2]School of Natural Sciences, Macquarie University, Sydney, NSW Australia

* Corresponding Author

Address for correspondence:
Cody A Freas
School of Natural Sciences
Macquarie University
Sydney, NSW 2113 Australia
Email: cody.freas@mq.edu.au

Abstract

Navigating insects often pause and rotate to sample their surroundings, behaviours termed scanning. These and other active sampling behaviours embody navigational uncertainty, and are key for spatial learning, yet their neural basis remains unclear and existing models impose scanning behaviours rather than explaining its emergence. Here, we show that scanning can emerge spontaneously from the same conserved neural circuits used for goal-directed navigation, without requiring a specialized scanning module. We built a biologically grounded model combining central complex (CX) steering, lateral accessory lobe (LAL) oscillators, and stochastic inhibition of forward speed. This minimal system produced diverse, realistic scan dynamics; saccades, fixations and reversals, whose features were qualitatively compared to high-speed video recordings of *Melophorus bagoti* scanning. Detailed analysis of these natural scans confirmed model predictions, including how scan structure depends on oscillator phase, goal-heading deviation, and navigational uncertainty. Furthermore, the model reveals that simple modulation of forward speed unifies a broad range of behaviors across ant species, from dashes to smooth oscillatory trajectories to pirouettes and voltes. Crucially, it establishes a general distributed control principle; where forward speed acts as a single adjustable parameter regulating the balance between goal-driven exploitation and information-seeking exploration, without requiring centralized decision-making processes.

Keywords: Central complex; Lateral Accessory Lobes; Distributed motor control; Central pattern generator (CPG); Oscillations

Background

Insects will often interrupt their forward movement and rotate in place, sampling visual information from multiple body orientations. These behaviours appear widespread across both walking and flying insects (as well as other taxa), and take various forms and names such as 'scans', 'volte', 'pirouette', 'turn back' 'peering' or 'dance' (Baird et al., 2012; Deeti et al., 2023a; Fleischmann et al., 2017; Graham and Collett, 2002; Lehrer, 1993; Mouritsen et al., 2004; Müller and Wehner, 2010; Stürzl et al., 2016; Tarsitano and Andrew, 1999; Ugolini, 2006; Wallace, 1959; Wehner et al., 1992; Wystrach et al., 2014; Zeil and Fleischmann, 2019). Despite its prevalence across taxa, the neural mechanisms underlying these seemingly highly structured behaviors remains unclear.

Scans are especially prevalent in visually guided ants (Deeti et al., 2023a; Fleischmann et al., 2017; Freas et al., 2018; Müller and Wehner, 2010; Nicholson et al., 1999; Wehner et al., 1996, 1996; Wystrach et al., 2014; Zeil and Fleischmann, 2019). An ant scanning behaviour consists of distinct periods: saccades, when the navigator is rotating, and fixations, when the ant pauses all movement (Figure 1A-C; Video 1,2, Müller and Wehner, 2010; Wystrach *et al.*, 2014; Zeil and Fleischmann, 2019; Deeti, Cheng, *et al.*, 2023; Freas and Cheng, 2025). Multiple saccades and fixations make up a singular scan, which may contain one or more rotational reversals (right then left/left then right) before forward movement resumes.The probability of scanning, while stochastic (Deeti, Cheng, *et al.*, 2023), increases with uncertainty, such as during learning walks (Müller and Wehner, 2010; Zeil and Fleischmann, 2019), route formation (Freas and Cheng, 2025, 2022; Haalck et al., 2023), when views are unfamiliar or unexpected (Deeti et al., 2023a; Freas et al., 2018; Schwarz et al., 2020b; Wystrach et al., 2014) or when views have been associated with aversive events (Freas et al., 2022; Wystrach et al., 2020a).

Functionally, scans are associated with information sampling, helping ants choose headings when cues are unfamiliar or in flux (Wystrach et al., 2014; Wystrach, Buehlmann, *et al.*, 2020; Deeti, Cheng, *et al.*, 2023; Freas and Cheng, 2025). Additionally, their prevalence during both learning-walks and early route formation indicates a role in view acquisition (Freas and Cheng, 2025; Müller and Wehner, 2010; Wystrach, 2023; Wystrach et al., 2014; Zeil, 2023; Zeil and Fleischmann, 2019).

Our mechanistic understanding of insect navigation has improved tremendously in the last decades - thanks to a combination of behavioural, neurobiological and modelling studies (Honkanen et al., 2019; Webb and Wystrach, 2016) - but current models do not explain the emergence and natural dynamics of scanning behaviours. Instead, 'scanning routines' are sometimes force-added in models to help the agent or robot sample views across directions (Baddeley et al., 2012; Gattaux et al., 2023; Husbands et al., 2021; Knight et al., 2019; Le Möel and Wystrach, 2020; Murray et al., 2019; Wystrach et al., 2013).

Here we ask whether and how ants' scanning dynamics could emerge from neural
structures typically implicated in insect navigation: the central complex (CX) and the
lateral accessory lobes (LALs). While the circuitry of these structures is detailed
primarily in a few insect model species, it is sufficiently conserved to allow for neural
modelling of insect navigation in various contexts (Adden et al., 2022; Clément et al.,
2023; Collett et al., 2025; Goulard et al., 2023; Shiu et al., 2024; Steinbeck et al., 2020; Stone
et al., 2017; Webb and Wystrach, 2016; Wystrach et al., 2020b). Both CX and LAL regions
are well established to integrate multiple directional information and control steering
commands in navigating insects (Franconville et al., 2018; Heinze, 2017; Honkanen et al.,
2019; Hulse et al., 2021; Li et al., 2020; Namiki and Kanzaki, 2016; Pfeiffer and Homberg,
2014).

Previous studies have established many key components of insect navigation,
including how goal headings are set in the central complex (CX) (Fisher, 2022;
Green and Maimon, 2018; Wystrach et al., 2020b; Stone et al., 2017) and the
intrinsic oscillatory dynamics in the lateral accessory lobes (LALs) that support
continuous zigzagging locomotion (Clément et al., 2023; Kanzaki, 2005; Namiki and
Kanzaki, 2016; Steinbeck et al., 2020). Building on these elements, the present work
addresses a distinct question that current navigation models do not resolve: whether
discrete active sampling behaviours such as ant scanning require specialised control
mechanisms, or instead emerge from interactions within the conserved navigation
circuitry itself.

To investigate this, we analysed high-speed recordings of natural scanning
behaviour in *Melophorus bagoti* foragers and asked whether their detailed dynamics
could be captured by the interactions between the CX and the LALs, under the
condition of occasional halted forward movement. We developed a biologically
constrained neural model by producing a CX steering output that modulates a
downstream intrinsic LALs oscillator, under the assumption that forward speed may
be transiently gated by a stochastic extrinsic signal. Without introducing a dedicated
scanning module or behavioural state switch, this model reproduces key qualitative
features of natural scans, including saccades, fixations, directional reversals, and
rare full-loop scans, which arise as a consequence of CX–LAL and forward-angular
speed interactions.

In addition to scans, an emergent property of this model is that by reducing instead
of terminating forward speed, the virtual agent produces additional behaviours,
typically categorised as non-scanning, observed in ants and other insects (Baird et
al., 2012; Demir et al., 2020; Gomez-Marin et al., 2011); including 'voltes' and full
pirouettes observed in some desert ant species (Fleischmann et al., 2017; Zeil and
Fleischmann, 2019), or the large oscillating paths observed in *Myrmecia* species
(Clément, Schwarz and Wystrach, 2023). These findings point to a conserved neural
control strategy where forward speed inhibition gates the expression of angular
motor output, allowing for shifting seamlessly between information gathering and

goal-oriented progression. The few novel assumptions added here enable us to form
predictions for future work.

Neural substrates of insect navigation

To interpret these results, we briefly summarise the neural substrates that motivated
the model. The CX tracks the insects' heading relative to the world, transforms
egocentric sensory information into allocentric current and goals directions, and
compares both to output steering commands. Within the CX's substructures, the
current heading direction is represented neurally as a bump of activity shifting along
the protocerebral bridge (PB) and the Ellipsoid Body (EB), while goal directions are
represented in the fan-shaped body (FB) (Honkanen et al., 2019; Kim et al., 2019;
Pfeiffer and Homberg, 2014; Seelig and Jayaraman, 2015; Stone et al., 2017). Current and
goal direction representations are compared within the CX and project left/right
steering commands to the LALs (Green et al., 2019; Honkanen et al., 2019; Iwano et al.,
2010; Mussells Pires et al., 2024; Westeinde et al., 2024); Figure 2A). The goal direction in
the CX can be updated by various pathways from different sensory modalities
(Honkanen et al., 2019), such as wind and odours ((Buehlmann et al., 2020; Knaden and
Graham, 2016; Müller and Wehner, 2007; Steck et al., 2011), visual familiarity via the
Mushroom bodies (Wystrach et al., 2020b) or path integration (Lu et al., 2022; Lyu et al.,
2022; Stone et al., 2017). In experienced ants, these different sources of directional
information are integrated and usually act in synergy to form a goal directed either at
the nest, or towards a known food source (Buehlmann et al., 2020; Knaden and Graham,
2016; Müller and Wehner, 2007; Steck et al., 2011; Wehner et al., 2016; Wystrach et al.,
2015; Wystrach and Schwarz, 2013). In the current modelling work we assumed the
existence of such a goal direction in the FB of the CX.

The LALs are premotor centers that act as a bottleneck, integrating information from
multiple higher processing centres and sensory inputs, including outputs from the
central complex, and relay these signals via descending neurons that transmit motor
commands to the thorax (Namiki and Kanzaki, 2016; Shih et al., 2015; Steinbeck et al.,
2020). The LALs also produce alternation between left and right turn via so-called
flip-flop neurons (Berni, 2015; Iwano et al., 2010; Kanzaki, 2005; Namiki and Kanzaki,
2016; Steinbeck et al., 2020), and present intrinsic activity which can result in the
regular lateral oscillations observed in many insects (Freas and Cheng, 2022; Iwano et
al., 2010; Izquierdo and Lockery, 2010; Kanzaki et al., 1992; Kuenen and Baker, 1983;
Namiki et al., 2014; Namiki and Kanzaki, 2016; Olberg, 1983; Wystrach et al., 2016),
notably in ants (Video 4, (Clément et al., 2023; Deeti and Cheng, 2025; Haalck et al., 2023;
Hangartner, 1967). The intrinsic generation of oscillatory movements allows for
continuous spatial sampling of information, and modeling studies demonstrate that
when these oscillations are modulated by odor or visual cues, they offer a highly
effective strategy for odour plume or gradient tracking, visual route following or goal
pinpointing (Adden et al., 2022; Kodzhabashev and Mangan, 2015; Le Möel and Wystrach,
2020; Wystrach et al., 2016).

Together, this CX–LAL circuitry is well suited to control continuous steering and sampling during locomotion. However, until now it is unknown if this same circuitry is sufficient to produce the discrete structures of ant scanning behaviour without invoking dedicated control routines.

Figure 1. Image composites of fixations (pauses – typically 50-150ms) during three different examples of scanning behaviours in *Melophorus bagoti*, all taken within 1m of the nest entrance, in inexperienced ants forming their route to a feeder. Images were extracted and compiled from highspeed video taken at 600fps at 1080p using a Chronos 2.1HD camera (field of view, 30 cm × 17 cm). For each image, fixations are indicated by consecutively numbered black arrows denoting the ant's orientation (except reversal). Rotational movement between fixations, saccades, are classified as either away (pink) or towards (green) the goal direction (~90°). When a fixation precedes a change in turning (from left to right in these examples), this fixation is classified as a reversal (orange). (**A**) Denotes a scanning example where it reverses direction once but then turns in a complete loop with no reversal (fixations 6-16). (**B**) Shows a scan which contains multiple reversals (two). (**C**) Illustrates a scan with one reversal, ~180° from the goal direction. Black dotted arrows denote pre/post scan forward movement.

228
Figure 2. Diagrams of the brain region connections and outputs using the neural circuit
model. (**A**) The central complex (CX), a mid-brain region whose subregions compare representations
of the goal heading with a representation of the agent's compass-based current heading. The CX
outputs bilateral signals to turn the left (red line) and right (blue line) to the Lateral Accessory Lobes
(LALs). (**B**) Within the LALs oscillator, left (L) and right (R) neurons reciprocally inhibit one another
(red and blue connections), while attempting to maintain a basal firing rate via internal feedback
(circular black arrows), forming an oscillator which outputs a stable anti-phasic oscillatory activity
between the L and R neurons across time (red and blue lines). (**C**) A steering signal is outputted from
the LALs and results in the agent's forward and angular movement. This steering signal stopped via
an external inhibitory 'freeze signal', that breaks the angular and forward speed central pattern
generators (CPG), initiating the scan. A threshold was implemented for the underlying angular drive to
restart the CPG after each fixation period, resulting in a saccade whose magnitude was determined
by this accumulated angular drive. The agent's angular speed was normalised by the forward speed
(linear normalisation - angular speed = angular drive/(forward speed + 0.1)) so the agent produces
larger saccades magnitudes during scans. Threshold was determined to roughly mirror saccade
magnitudes in real world ants. Scan duration was implemented as an exponential duration distribution
through a dice-roll at each simulation step. (**D**) Characteristics of an example path, with a single
scanning bout, generated by the model. Black dots represent the agent's head position at each
simulated step coupled with coloured bars which indicate both the forward speed (arbitrary scale) and

heading direction of each step. During the scanning bout, the agent's forward speed is zero and the
model produces several fixations in separate directions. The fixations of this scanning bout are
zoomed and separated into a sequence of fixations. Coloured arrows indicate saccades, rotational
movements between fixations which were defined as either away (pink) or towards (green) the goal
direction. (**E**) The modelled CX's turning signal output towards the goal direction, based on if the
agent's heading direction is to the left or right of the goal. **(F)** The oscillatory activity between the R
and L neurons in the LALs during the example path. The agent's (**G**) heading direction, (**H**) angular
speed, and (**I**) forward at each step during the path.

Results and Discussion

A simple model for scanning behaviour

We first constructed a simple model of the insect central complex (CX; Figure 2A), following standard approaches (Goulard et al., 2023; Stone et al., 2017; Wystrach, 2023; Wystrach et al., 2020b). This circuit compares the agent's current heading to a stored goal heading (which, in our simulations, is directionally fixed) and generates lateralised steering signals to reduce heading error. However, unlike previous models (but as in Adden *et al.*, 2022) the CX output here does not steer the agent directly, but instead modulates an intrinsic oscillator that generates alternating turns (Figure 2B), a mechanism consistent with the continuous oscillatory patterns known to underlie ant navigation (Clément, Schwarz and Wystrach, 2023).

This oscillator, which is widespread in insects (Cheng, 2024; Kanzaki, 2005; Kanzaki and Mishima, 1996), resides in the lateral accessory lobes (LAL), a pre-motor region that receives, among others, CX outputs (Heinze, 2017). We modelled the oscillator as a system of reciprocal inhibition between the left and right hemispheres, producing alternating steering signals (Kanzaki and Mishima, 1996; Steinbeck, Adden and Graham, 2020; Clément, Schwarz and Wystrach, 2023). The difference between these left and right signals - such as conveyed by descending neurons (DN) to the thoracic ganglia (Büschges and Ache, 2025) - determines the agent's angular speed, thereby generating the rhythmic turning dynamics akin to those observed in ants (Clément, Schwarz and Wystrach, 2023). In combination with the CX modulation, this oscillator produces a continuous oscillatory trajectory generally oriented toward the goal direction set in the CX (which is always to the right in our examples; Figure 2D).

To generate scanning behaviour (Figure 1A–C), we implemented a mechanism to intermittently interrupt forward movement, as observed in real ants, where pauses occur randomly (Deeti et al., 2023a). In our model this was achieved using a simple external inhibitory 'freeze signal' that halts forward motion at random intervals. Since scan duration in real ants follows a Poisson like distribution (Deeti et al., 2023a), we approximated this by performing at each simulation step a dice-roll with equal probability (calibrated as 1/mean of ants observed scan duration) to restart forward motion, producing the desired Poisson distribution of scans' duration.

During scanning, real ants display rotational saccades of variable duration and angular magnitude (Figure 1A–C). To replicate this, we introduced a threshold-based mechanism: after each fixation (i.e., zero angular and forward speed), the underlying angular steering signal accumulates until surpassing a threshold, triggering a

298 saccade. The resulting angular magnitude of the saccade corresponds to the sum of
299 the angular drive accumulated during the fixation.
300
With this, the agent now scans, but the heading deviation during scanning remains
within the same limited range as during forward movement, highly constrained
toward the goal direction. This does not reflect real scans, which involve much larger
angular deviations (Deeti et al., 2023a; Fleischmann et al., 2017; Freas et al., 2019; Freas
and Cheng, 2025; Müller and Wehner, 2010; Wystrach et al., 2014). We hypothesized that
this limited turning during forward movement is due in part to biomechanical
constraints: fast forward motion impedes sharp turns, while slowing down allows
easier rotation. When ants stop, they are free to execute sharp turns via leg
coordination that would be otherwise impossible (e.g., inner legs moving backward
and outer legs forward), enabling high rotation during scans (Figure 1). To implement
this constraint, we normalized the agent's angular speed by the inverse of forward
speed: the slower the agent moves, the more it turns for a given steering signal. As a
result, the agent now produces larger saccades and greater heading deviations
during scans, while remaining more constrained during forward movement. To match
the saccade amplitudes observed in real ants, we adjusted the threshold for breaking
fixation so that the resulting distribution of saccade angles (Figure 3A) approximates
those seen in ants (Figure 3B). Higher thresholds yield less frequent but larger
saccades.
319
320 Overall, the interaction between the CX output (Figure 2A), oscillator phase (Figure
321 2B), and the scan control mechanism (Figure 2C) produces complex, nonlinear
322 dynamics, giving rise to emergent behaviours, which we explored and compared to
323 real ants in the following subsections.
324
325 *CX steering influences saccade magnitude*
326
327 A key prediction of our model is that the CX steering mechanism operates
328 continuously, including during scans, and its influence should be detectable in the
329 scan's structure. Specifically, the strength of the CX's corrective signal toward the
330 goal increases with the agent's angular deviation from that goal (Figure 2A).
331
If guided solely by the CX, the agent would barely turn away from the goal but due to
noise. However, actual steering is governed by the downstream oscillator in the LAL,
which ultimately determines the direction of each saccade during scans. Saccades
toward the goal occur when the oscillator is in phase with the CX corrective signal,
resulting in stronger angular drive and therefore larger saccades. Conversely,
saccades away from the goal arise when the oscillator and CX corrective signals are
in conflict, the later restraining the former, producing weaker angular drive and thus
smaller saccades. Since the CX's corrective signal roughly scales with angular
deviation (Figure 2A), this difference in saccade amplitude (toward vs. away from
goal) should be minimal when the agent is facing the goal direction but becomes
more pronounced as it is facing away (Figure 3C).
343
344 This predicted pattern closely matches real ant data (Figure 3D). Across all datasets,
345 including inexperienced and experienced ants, whether homing or foraging (treated

as random error in our LME), saccade amplitude was significantly explained by both
the saccade direction (towards and away from the goal) ($F_{(1,1890)}$ = 5.08, p = 0.024)
and angular deviation ($F_{(1,1890)}$ = 18.69, p < 0.001) and by and large explained by
their interaction indeed ($F_{(1,1890)}$ = 43.04, p < 0.001). Specifically, saccades toward
the goal were larger than those turning away from it, and this difference increased as
the ants faced away from their goal, supporting the model's prediction. Together, this
strongly supports that both the oscillator and CX are at play during scanning, with the
later continuously pulling the ants towards its goal heading.

Our statistical analysis on saccade amplitude also revealed significant fixed effects
for the individuals (LRstat =200.25; p < 0.001) and conditions (LRstat =49.08, p <
0.001) (see Statistical Methods for details) as well as an interaction between heading
orientation (*angular deviation* from goal) and whether the previous fixation was a
reversal ($F_{(1,1890)}$ = 6.86, p = 0.009; Figure 4A), which we will address in the next
sections.

359
Figure 3. Saccade angle distribution and the effect of CX steering guidance on saccade angle
magnitude. Saccade angle distributions in (**A**) the modelled agent and (**B**) real ants with all scan
conditions combined. After each fixation, the subsequent next saccade angle is plotted against the
fixation's angular divergence from the goal direction in both (**C**) the model agent and (**D**) real ants with
all conditions combined. The general trends of the model are classified through coloured arrows
(away-green, towards-pink) representing the increasing next saccade angle towards the goal direction
and the decreasing next saccade angle away from the goal direction of the modelled agent when
fixation orientation from the goal direction was large. This represents the effect of the changing
strength of the CX steering signal with increasing divergence from the goal direction; easier to turn
towards goal and thus large saccades, while harder to turn further away and small saccades. This
pattern is replicated and significant in real world ants. For the box and whisker plots in panels C and
D, the box spans the inter quartile range while the horizontal line indicates the mean. Whiskers extend

to the INr X 1.5 while outliers beyond this range are shown as '+' symbols. The indentation of the bar
around the mean indicates the 95% confidence interval.
374
375   *The oscillator phase influences fixation duration*
376
Having explored saccade amplitude, we next asked whether the model also
accounts for fixation duration. In our model, fixations during scanning are periods of
fully inhibited locomotion. A fixation ends when the angular drive (generated by the
oscillator) crosses a threshold of angular drive (left or right), breaking the inhibition
and triggering a saccade (Figure 4A–C). Since the angular drive (L - R activity of the
LAL) fluctuates cyclically with the oscillator phase, the time required to reach this
threshold naturally varies.

Notably, the models' fixations can be especially prolonged during reversal phases;
when the oscillator transitions from left to right, or vice versa. In extreme cases, the
angular drive initially builds toward the direction of the previous saccade (e.g., left),
but before crossing the threshold, the oscillator reverses phase. The drive then shifts
in the opposite direction (e.g., right) and must accumulate again in that new direction
before triggering a saccade (Figure 4B, yellow period of second reversal). As a
result, reversal fixations have a tendency to last longer than other fixations (Figure
4A,D). Remarkably, ant behavior supports the model's prediction (Figure 4E). In our
linear mixed-effects model (LME), reversal fixation status was the only significant
predictor of fixation duration ($F_{(1,1894)} = 76.01$; $p < 0.001$).

The model also predicts a subtler pattern: fixation duration gradually decreases with
distance from a reversal fixation (Figure 4D). This occurs because angular drive is
on average strongest mid-cycle (i.e., in the middle of a left or right phase) and
weakest near phase transitions. When we replaced the binary "reversal/non-
reversal" variable with this sequential phase information (Figure 4A) in our statistical
model, the effect on fixation duration remained significant, but only as an interaction
with the upcoming saccade's direction relative to the goal ($F_{(1,1894)} = 4.65$; $p = 0.031$).
Specifically, because the CX upregulates the oscillator on the side toward the goal,
the angular drive raises quicker and fixation duration tends to be shorter for
saccades in that direction.

Together, these results support the idea that fixation duration arises from a
threshold-crossing mechanism and signal governed by the oscillator's phase and its
interaction with CX input. Despite outward stillness during fixation, oscillator and CX
are continuously active and shaping scan dynamics.
408
409   *Influence of neural noise and thresholds on fixation duration*
410
Our model predicts a general negative correlation between fixation duration and the
amplitude of the subsequent saccade (Figure 4F, red). This arises because a short

fixation - that is, a rapid threshold crossing - implies a strong angular drive, and
stronger angular drive typically results in a larger saccade.

However, we were initially surprised to observe in our model that this negative
correlation breaks down for the shortest fixations (Q1): in this regime the agents
exhibit a positive correlation instead (Figure 4F, blue). This pattern is an indirect
consequence of neural noise, which is systematically introduced in the model.
Specifically, noise in the inhibitory neuron freezing movement causes variability in
the fixation-breaking threshold (illustrated as a grey band in Figure 4B). When this
threshold is stochastically lowered due to noise, it can be reached more quickly,
producing very short fixations if the current angular drive is strong. However,
because the threshold is lower, the accumulated angular drive at threshold crossing
is necessarily reduced, limiting the magnitude of the resulting saccade. This
mechanism imposes an upper bound on the accumulated drive, and therefore on
saccade amplitude, for the shortest fixations, creating a hard limit in the correlation
plot (grey region, Figure 4F). The outcome is a positive correlation between fixation
duration and saccade amplitude for the shortest fixations (Figure 4F, blue).

While this effect was not initially anticipated, its predictions are supported by
behavioral data. Ants' fixation duration is positively correlated with saccade
amplitude for the shortest fixations (Figure 4G, blue, Quartile 1: $F_{(1, 281)}$ = 11.00, p =
0.001; R^2 = 0.038; $t_{(281)}$ = 3.32, p = 0.001), and negatively correlated for longer
fixations (Figure 4G, pink, Quartiles 2-4: $F_{(1, 795)}$ = 5.97, p = 0.015; R^2 = 0.007; $t_{(795)}$ =
−2.44, p = 0.015).

The emergence of this complex pattern from the model, despite not being explicitly
designed to capture it, supports the hypothesis that fixations are governed by an
internal accumulation of excitatory activity (angular drive), which is released upon
reaching a threshold. This mechanism is consistent with integration-to-bound neural
models that have been proposed for self-initiated actions in other species (e.g.
Murakami *et al.*, 2014).

443
Figure 4. Relationship between fixation duration and reversal. (**A**) Sequence of fixation directions
and their duration for an example scan containing multiple reversals (orange), showcasing that the
longest fixations occur when the ant reverses directions on the next saccade (fixation duration
continuum, blue-min; yellow-max). (**B**) shows changes in the cumulative turning drive over the course
of the example scan. To initiate a saccade, the drive must surpass a threshold (line). This threshold
possesses some level of noise which leads to variance in when a saccade appears as well as to the
variance in saccade magnitude. Reversals occur when the oscillatory cycle changes phase (±), which
are associated with longer times for turning drive to accumulate beyond threshold (orange). (**C**)
Angular speed through the simulation, spikes represent saccades to the left and right and the change
from positive to negative illustrates reversal periods (orange). Relationship between fixation duration
and oscillatory cycle changes (orange) and subsequently reversals, occur in (**D**) modelled and (**E**) real
ants. In both, the longest fixation duration typically occurs at the reversal (0), increasing as the
reversal approaches (-), while decreasing post reversal (+). For these box and whisker plots, the box
spans the inter quartile range while the horizontal line indicates the mean. Whiskers extend to the INr
X 1.5 while outliers beyond this range are shown as '+' symbols. The indentation of the bar around the
mean indicates the 95% confidence interval. Correlation between fixation duration and the next
saccade angle in (**F**) modelled and (**G**) real ants. Data are split into the lower 25% (Q1, Blue) and
upper 75% (Q2-Q4, Pink) of the distribution. In both, the general tendency in Q1 is significantly
positive (real ant; Linear regression model; $F_{(1, 281)} = 11.00$, p = .001), while beyond this (Q2-Q4) the
tendency is significantly negative ($F_{(1, 795)} = 5.97$, p = 0.015). The 'unlikely' zone in grey depicts the
area of high saccade magnitudes following very short fixations, which should be highly unlikely given
the low threshold drive must accumulate needed to break fixation in these instances.
466
467 *Random Timing of Scan Starts and Stops*
468
Previous work on scanning behavior in ants (*Deeti et al., 2023*) found that the
number of saccades within each scanning bout follows a Poisson-like distribution
(Figure 5B). From this, they inferred that the initiation and termination of scans are
governed by a random-rate process.

In our model, this stochasticity is implemented by a probabilistic trigger: scan
initiation is triggered at a random time, and at each moment, scanning can end with
a probability proportional to (*1 / mean ants' observed scan duration*), producing the
desired poisson distribution of number of saccades (Figure 5A). Because of this,
scans can begin or end at arbitrary points in the oscillatory cycle. This leads to a
specific prediction: in scans containing multiple reversals (direction changes in
turning), the first and last sweeps, defined here as a bout of sequential saccades in
the *same* direction, should often be truncated compared to "full" sweeps (e.g., the
second or penultimate sweeps), which start and end exactly at reversals. In other
words, sweeps bounded by two reversal should reflect a full (one-sided) oscillatory

cycle, while sweeps bounded by the beginning or the end of a scan may not, due to
the stochastic start or end of scans. Therefore the latter should tend to be shorter,
and thus contain less saccades, than the former.

To test this, we excluded scans with one or zero reversals, focusing on bouts where
we could compare truncated sweeps (first and last) with adjacent full sweeps
(second and penultimate). The model predicts, and the data confirm, that truncated
sweeps contain fewer saccades than their adjacent full sweeps (Figures 5A–D).
Specifically, the first sweep is significantly shorter than the second (Figure 5C, z =
4.47; $p < 0.0001$), and the last sweep is significantly shorter than the penultimate
(Figure 5D, $z = -3.30$; $p = 0.001$).

These results thus support the stochasticity of scan initiation and termination. The
inhibition of forward motion seems to occur independently of the oscillatory phase.
Also, it confirms that within scans, saccade direction follows the underlying
oscillatory cycle, rather than being purely random.

499
Figure 5. Scans start and end irrespective of oscillator state and saccade number
distributions. In the modelled agent, (**A**) the number of saccades within a scanning sweep (before
first reversal) was compared with the post-reversal sweep number showing a general upward trend.
(**B**) the number of saccades within the penultimate sweep compared to the final sweep of the
scanning bout, showing a downward trend. In real ants, (**C**) comparison of the number of saccades in
the first and second sweep and (**D**) the penultimate and final scanning sweeps. Both model and real
ant sweep comparisons only contain scans which contained two plus reversals, in order to compare
the starting/stopping sweeps with a full, within reversals sweep (second/penultimate). Statistical
comparisons were made using Wilcoxon tests. The distribution of saccade counts in each scanning
bout within (**E**) the model, showing a poisson distribution and in (**F**) real ants, showing a 'poisson-like'
distribution.
511
512 *CX strength links navigational uncertainty to scan structure*
513
We next examined how navigational uncertainty, which can be modelled as the
strength of the CX's steering signal, influences scan structure. In the model, a
heavily weighted CX signal, representing high certainty in the goal direction,
produces greater corrective steering, thus constraining the agent heading toward the
goal. As this signal weakens, so does its corrective influence, allowing for larger
deviations away from the goal direction. Thus the model predicts an inverse
relationship between CX strength (i.e., navigational certainty) and this divergence,

which we quantified by measuring the rotation away from the goal covered by the
scanning ant before performing a first reversal (Figure 6A, arrow). Ant behavioural
data echoes this pattern: experienced ants - which we assume have a high certainty
in their goal direction - showed smaller divergences from the goal on their first
reversal than inexperienced ants (Figure 6B; $F_{(3, 274)}$ = 3.20, p = 0.0239).

The model also predicts a positive association between CX strength and the agent's
saccade angle (Figure 6C, arrow). While initially counter intuitive, as a strongly
weighted CX signal constrains angular divergence away from the goal, strong CX
steering also produces much larger corrective saccades when it aligns with the
oscillator phase (as shown in Figure 3C). This pattern aligns well with our real ant
testing conditions, with significantly larger saccades in experienced ants vs.
inexperienced ants (Figure 6D; $F_{(1, 2137)}$ = 587.6, p < 0.001).

These results support the CX's corrective steering input role in modulating the
structure of scanning, tuning how tightly ants orient to their goal as well as how wide
their saccades are during scanning, especially when in phase with the oscillator.

536
Figure 6. The strength of the CX's corrective turn signal, and its inverse relationship with
navigational uncertainty. The CX's signal signal strength on the oscillator would be predicted to be
strong when navigational uncertainty is low, such as when the ant is highly experienced. Conversely,
CX signals would be weak when navigational uncertainty is high, like during route formation. The
model predicts that under high CX signal strength, (**A**) the first reversal direction (which is likely to
follow a sweep away from the goal) should be constrained angularly to closer to the goal direction.
Additionally, this metric should rise, the CX signal strength decreases. (**B**) These trends closely mirror
the real ant data, with highly experienced foragers, showing first reversal directions which were more
constrained angularly to closer to the goal direction compared to inexperienced forager conditions.
The model also predicts, somewhat initially continued intuitively, that (**C**) saccade angle should also
be inversely associated with CX signal strength (grey arrow), which mirrors (**D**) the reduction in
saccade angle amplitude in high uncertainty conditions in inexperienced foragers. Initially one might
theorise low CX strength should mean larger saccades, yet it is under high corrective CX steering
strength that we see large turns back towards the goal (See back to Figure 3C,D), resulting in this
association. For all box and whisker plots, the box spans the inter quartile range while the horizontal
line indicates the mean. Whiskers extend to the INr X 1.5 while outliers beyond this range are shown
as '+' symbols. The indentation of the bar around the mean indicates the 95% confidence interval.

554
555 *Rare 'Full Loop' Scans reveal CX–Oscillator Interactions*
556

557 Despite its simplicity, our model's closed-loop dynamics generate a surprising
558 diversity of scan forms. One striking example is the emergence of rare 'Full Loop'
559 scans (Figure 7A,B; Video 4,5), in which the agent completes a long series of
560 saccades in the same direction, producing a full loop before resuming normal
561 movement.
562

In the model, these occur when a strong central complex (CX) corrective steering
signal coincides with a critical point in the oscillator's cycle, effectively shifting the
phase by producing a rebound of activity before the current cycle ends (Figure 7A,B,
black arrows). Instead of reversing direction at the usual reversal points, the agent's
saccade angle decreases (as usual) but then increases again in the same turning
direction, producing a complete rotation.
569

570 Such Full Loop scans are not unique to the model; they also occur, albeit rarely, in
571 real ants, and have been reported in multiple species (e.g., *M. bagoti* Figure 1A,
572 Video 1; *Cataglyphis cursor,* Video 2; (Fleischmann et al., 2017; Freas et al., 2019; Freas
573 and Cheng, 2025; Zeil and Fleischmann, 2019).
574

Mechanistically, our results suggest that the CX modulates the oscillator to enable a
rebound within the same phase. If, by contrast, the CX and oscillator acted purely in
parallel (with their outputs summed independently), the oscillator would produce a
reversal in angular gain as usual, conflicting with the CX output and preventing a fast
completion of the loop. Consistent with the model, real ants show re-acceleration in
the second half of the 'Full Loop' scan (real ants: Figure 1A, fixations 12–16; Videos
1–3; model: Figure 7A,B), supporting the idea that CX output is indeed branched
upstream of the oscillators, and can shift its phase.
583

584 These events are too rare in our current dataset for statistical analyses, and our
585 simple model is not designed for quantitative comparisons. Nonetheless, such rare
586 behaviors may offer valuable insights into the coupling between oscillatory motor
587 control and upstream mechanisms, and could be a promising focus for future
588 studies.

589
590
Figure 7. Diverse behaviours emerge from interactions between the central complex's (CX)
steering signal and the modulation of the agent's forward speed. (**A**) A single 'Full Loop' scan
(Video 5), the agent terminates forward speed and exhibits a scanning bout with both fixations and
saccades. Here, the CX's corrective steering excites the oscillator and reverses its phase (black
arrow), resulting in the agent continuing the loop rotation rather than reversing. These full loop
behaviours are rare within scanning (most reverse) but widely observed in ants (Video 1-3). (**B**)
Shows a double 'Full Loop' scan example where the agent performs two full loop rotations in a scan.
As in panel A, (Simulations - Video 6). Here, the oscillator's phase is reversed by the CX twice (black
arrows). (**C**) A 'volte' is a loop without stopping (Simulation - Video 7). A common behaviour in
Cataglyphis desert ants (Fleischmann et al., 2017). This behaviour arises in the agent when
forward speed is low but not stopped, boosting angular speed and allowing the CX steering output to
reverse oscillator phase without fixations. (**D**) An example of a '*Myrmecia*-like' path in the agent,
characterized by moderate forward speed and larger lateral oscillations, reminiscent of real world
Myrmecia ants (Clément, Schwarz and Wystrach, 2023; e.g. Video - 4). Here, moderate forward
speed allows for larger alternating turns and lateral displacement.
606
607
608
609

Graded Forward Speed Control Generates Voltes and Oscillatory Patterns

In our initial implementation, forward speed inhibition was binary: the agent ran or stopped completely, as if the central pattern generator controlling leg movement were switched off at a random time. Insects, however, show more nuanced control. For example, desert ants can slow down as their path integration vector decreases (Buehlmann et al., 2018), or accelerate when on fully familiar routes (Clément et al., 2023; Haalck et al., 2023).

We therefore modified the model to allow forward speed inhibition to work along a continuum, with low inhibition merely slowing down the insect rather than stopping it. Because we assumed that forward speed biomechanically impedes angular speed, a sudden but partial reduction in forward speed produces sharper turns; and in some cases complete loops (Figure 7C) resembling so-called 'volte' that have been described in the ant literature (Fleischmann et al., 2017; Zeil and Fleischmann, 2019).

Furthermore, it is known that the oscillator phase itself modulates forward speed, which can be modeled simply as the sum of the oscillator's left and right activities (Clément et al., 2023). This coupling accelerates the agent when facing its travel direction and slows it when oriented sideways, producing efficient search trajectories. The effect of this control varies among species and is particularly strong in *Myrmecia croslandi* (Clément, Schwarz and Wystrach, 2023; Video 4; see also Video 3 for *M. nigriceps*). In the model, continuously operating at lower forward speed, releases the constraint on angular speed and naturally produces the large, regular oscillations seen in these ants (Figure 7D), regardless of CX signal strength (See Supplemental Figure 1 for example paths under weak and strong CX signal strength).

Interestingly, as a result of this control apparent in such slower agents, increasing the strength of the CX's influence on the oscillator produces not only more goal-oriented trajectories, but also increase forward speed and increase the oscillations frequencies while decreasing their regularity (Supplemental Figure 1A,B). These four covarying proprieties mirrors the behavior of *M. croslandi* ants on familiar routes compared to unfamiliar terrain (Clément et al., 2023), consistent with the idea that visual familiarity generates strong CX goal-heading signals (Wystrach, 2023).

Finally, in strongly oscillating situations, full loops also arise without apparent external inhibitory signal. Instead, they can emerge naturally from the continuous oscillatory control on forward speed, which slows the animal when facing away from its goal and thus enhances the rotation at the end of a sweep (e.g., *Myrmecia nigriceps*, Video 3). This suggests that the forward–angular speed coupling is a general property of the locomotor system, not solely tied to an external "stop" phase, though whether it is purely biomechanical or neurally reinforced remains to be tested.

Overall, introducing graded forward speed control shows that the same CX–oscillator interaction, originally modeled for scanning, can also explain patterns observed in continuous navigation (periods of goal-directed forward movement), linking route familiarity, speed modulation, scanning behaviours, CX and oscillatory turning dynamics.

General Discussion

This study shows that diverse movement behaviors in ants, including scanning, pirouettes, voltes, and goal-directed runs, can all emerge from a single, conserved neural architecture. By combining biologically grounded models of central complex (CX) steering, intrinsic lateral accessory lobe (LAL) oscillators, and minimal modulation of forward speed, we capture fine-grained scanning dynamics that match high-speed behavioral data. Importantly, these behaviors do not require dedicated scanning modules; instead, they arise naturally from continuous control mechanisms already implicated in insect navigation.

CX and Oscillator Interactions

In our model, the central complex (CX) outputs modulate the oscillator circuit in the lateral accessory lobes (LAL), producing goal-oriented oscillatory paths (Figure 2D; Figure 7A-D, for all paths goal direction to the right). While the anatomical pathways mediating this CX–LAL connection have been explored in species like moths (Adden et al., 2022), their implementation in ants remains unknown. The scanning dynamics observed here constrain possible architectures and offer insights into how the CX output shapes motor control via the LAL.

CX steering gain increases with angular deviation

Our data show that corrective saccade amplitude increases with the agent's angular deviation from the goal (Figure 3) which is consistent with behavioural data in ants and flies during continuous navigation (Lent et al., 2010; Westeinde et al., 2024) and suggests that the CX output signal increase with angular deviation . In *Drosophila*, the CX outputs PFL2 neurons respond maximally when the fly faces directly away from the goal, enhancing activity in PFL3L and PFL3R neurons that drive left or right turns. Here, we simply modelled two PFL populations (PFL-L and PFL-R), tuned to both ±135° from the goal (i.e., ±45° from the anti-goal), which mimics CX steering cells observed in monarch butterflies (see Fig.4G of Beetz, Kraus and el Jundi, 2023), and result in similar steering dynamics as observed in *drosophila* (compare Figure 2A and Westeinde *et al.*, 2024). Whatever the exact implementation, evidence supports the idea of a conserved CX output function that encodes angular deviation from the goal with increasing strength, and is at play during both scanning and continuous navigation.

CX directly modulates the LAL oscillator

 A key question is whether the CX directly modulates the LAL oscillator (serial control), or whether both act independently in parallel. In plume-tracking moths, CX output has been shown to modulate LAL flip-flop neurons driving zigzagging (Adden et al., 2022). In ants, strong goal signals in the CX - whether elicited by the path integrator or visual familiarity (Wehner et al., 2016; Wystrach et al., 2020b, 2015) do not only sharpen directional accuracy but also increase oscillation frequency (Clément et al., 2023), showing that CX output modulates the LAL intrinsic oscillatory activity .

Similar frequency increases occur in response to optic-flow prediction errors,
although these likely bypass the CX (Dauzere-Peres and Wystrach, 2024).

Our findings offer a qualitatively different form of evidence for a direct modulation of
the LAL oscillator by the CX. The rare 'full loop' scans - 360° turns without reversal -
only emerge in the model because the CX can shift the oscillator's phase mid-cycle
(Figure 3A,B). This requires tight coupling between CX output and LAL dynamics,
and would not occur if both acted independently. Together, present and past results
provide strong support for a direct, serial modulation of the LAL oscillator by the CX
in ants.

An opponent process in the CX output

In our model, the output of the CX to the LALs (via PFL neurons) provides both
ipsilateral excitatory connections and a contralateral inhibitory connection (Figure
2A). In insects, the latter may be indirect, via the stimulation of inhibitory neurons
such as the protocerebral bilateral neurons in the LAL (Adden et al., 2022; Kanzaki et
al., 2004; Mishima and Kanzaki, 1999). Functionally, this contralateral inhibition ensures
that only one side of the LAL is activated by the CX at a time, avoiding simultaneous
bilateral activation, which in the model would produce inappropriate bursts of forward
motion, particularly when the agent is misoriented.

More generally, this opponent organisation between the left and right hemisphere
converts CX directional information into a normalised steering signal that reflects
angular deviation . Similar opponent processes have been proposed for mushroom
body output neurons encoding approach vs. avoidance, or left vs. right scene
familiarity signals (Le Möel and Wystrach, 2020; Murray et al., 2019; Wystrach, 2023),
suggesting that lateralised, opponent coding is a general principle in insect
navigational circuits for ensuring stable and interpretable motor outputs.

A Simple Control Principle: Forward Speed Gates Exploration

727
The key insight from this work is that forward speed modulates the expression of
angular motor output, effectively acting as a behavioral switch between exploitation
and exploration. When forward motion is suppressed, angular drive is released,
producing scanning behaviors. When forward speed is high, angular deviations are
dampened, yielding straight, goal-directed trajectories. By simply adjusting this one
parameter, the model generates a spectrum of behaviors, from sweeping oscillations
to full rotational scans, mirroring inter- and intra-species variation observed in real
ants. Forward speed modulation of active sampling reflects a distributed control
process, arising from the dynamic interactions between the CX, oscillator, and
thoracic CPGs, rather than requiring high-level modulation from the brain.

738
Figure 8. Summary of the behavioural spectrum produced by the modelled agent as a function
of forward speed inhibition facilitating angular speed. High forward speed results in an agent that
'sprints', with its straight paths resembling desert ants (*Melophorrs bagoti and Cataglyphis*).
Decreasing forward speed progressively increases angular speed facilitation, leading to the large
oscillations of Myrmecia under moderate forward speed (Clément, Schwarz and Wystrach, 2023;
Video 4), 'voltes' under low forward speed, and ultimately scanning behaviours when forward speed
reaches zero. Here, angular facilitation plateaus, the central pattern generator (CPG) is disrupted and
the agent scans, with the choreography of this scan, are dictated by interactions between the CX
signal strength and the oscillator is observed in ants (Real Ants - Video 1-3; Simulations - Video 5,6)
748

A continuum of movement modes across context and species

At one extreme, forward speed is fully inhibited, halting leg CPG output, and making
angular drive maximally expressed, producing what we term scans. At the other,
high forward speed suppresses angular turns, yielding straight, fast trajectories.
Between these extremes lies a spectrum of behaviors, such as broad oscillations or
the occasional 'volte' loops, that emerge when forward speed is only partially
reduced (Figure 8).

This continuum reflects a functional trade-off between exploration (high angular
motion) and exploitation (high forward speed), which can simply be adjusted by a
single parameter, forward speed.

This continuum also helps explain interspecific variation. Desert ants (*Melophorus*
bagoti, multiple *Cataglyphis species, Occymyrmex robustior etc...*), adapted to
thermophilic foraging, favour high forward speed, which thus limit oscillations
amplitude, and rely on brief, stochastic stops for visual sampling (Clément et al., 2023;

Deeti and Cheng, 2025; Muser et al., 2005); figure 8). *Myrmecia* ants, by contrast, forage
at lower temperature and can afford to operate at slower speeds, allowing large,
regular oscillations to coexist with forward progression (Clément et al., 2023); Figure
8). Our model can shift seamlessly from one style to the other by simply adjusting
forward speed.

An individual can also shift along the continuum to adapt to the current context. For
instance, experienced *Cataglyphis velox* running along their familiar route, where
navigational certainty -which can be implemented as a strong goal heading in the
CX (Figure 6)- favour exploitation by constraining angular deviation, increasing
forward speed and thus diminishing the occurrence scans. Similarly, *Myrmecia* can
stop and scan under high uncertainty (Freas et al., 2018; Islam et al., 2020) or
accelerate and straighten to escape adverse situations (Clément et al., 2023; Deeti et
al., 2023b).

The emergence of such different-looking trajectories suggests that the modulation of
forward speed may represent an ancestral and fundamental control strategy in ants.
This simple control is used at both individual decision and evolutionary time scales :
to adjust individuals' behavior to the current context, as well as ant species to their
ecology.

An ancestral design? Parallels across taxa

This forward–angular coupling is not unique to ants. Similar mechanisms appear
across taxa (even in bacterial run–tumble cycles), highlighting a shared control
strategy (Cheng, 2024). In drosophila larvae, angular reorientation also arises from a
continuous body-bend oscillator. The expression of body bending is constrained by
forward crawling peristalsis (Wystrach et al., 2016), which can be transiently
suppressed by descending neurons (e.g., PDM-DN), unmasking the oscillator and
producing sharp turns (Tastekin et al., 2018; Berni et al., 2012; Pulver et al., 2015),
akin to ant scans.

In adult drosophila and other insects, multiple descending neurons have been
identified that modulate forward speed (Büschges and Ache, 2025). Some (BPNs),
promote acceleration (Bidaye et al., 2020), whereas others trigger strong speed
reductions, or halting responses (Sapkal et al., 2024). Furthermore, left-right firing rate
differences between descending neurons (DNa-1/DNa02) can predict rotational
velocity driving pivoting behaviour (Büschges and Ache, 2025; Rayshubskiy et al., 2025;
Yang et al., 2024); providing a circuit-level analogue to our modelled angular speed
modulation.

Larvae's stops also appear stochastic, and their frequency can be modulated by
negative sensory evidence such as going down attractive odour gradient (Berni et al.,
2012; Gomez-Marin and Louis, 2014), just as scanning in ants is more likely under high
navigational uncertainty (Deeti et al., 2023a; Freas et al., 2022, 2018; Freas and Cheng,
2025; Schwarz et al., 2020a; Wystrach et al., 2020a, 2014).

Finally, in both ants (Buehlmann et al., 2018; Haalck et al., 2023) and larvae (Gomez-
Marin and Louis, 2014; Luo et al., 2010), forward speed control is not binary, but
graded, modulated smoothly by sensory evidence.

Although not fully developed in larvae, the CX and LAL are conserved structures,
ancestral to arthropods (Kanzaki, 2005; Kanzaki and Mishima, 1996; Pfeiffer and
Homberg, 2014). Whether the descending neurons that inhibit forward movements in
Drosophila (Rayshubskiy et al., 2025; Tastekin et al., 2018) are similarly conserved
across arthropods, or represent convergent evolutions in some taxa, remains to be
seen.

Model limitations and future directions
813
While our model successfully captures fine-scale scan dynamics, several limitations
remain. First, we assumed a single, static goal direction represented in the fan-
shaped body of the CX. While this suffices to simulate realistic scanning of ants
navigating toward a known goal, real insects often juggle multiple directional cues,
from path integration, visual landmarks, wind, and olfaction, and may maintain
simultaneous goal representations. These goals may be employed sequentially or
weighted adaptively within the CX (Freas et al., 2020; Le Moël et al., 2019; Stone et al.,
2017; Sun et al., 2020; Wehner et al., 2016; Wystrach et al., 2015). Extending the model to
simulate multi-goal conflict or cue integration would allow exploration of how insects
handle navigational uncertainty, and how the latter impacts scanning.

824
Second, the model does not currently explain nest-directed fixations. These so-
called "pirouettes" or "lookbacks", where longer fixations happen when looking
towards the nest have been reported in ants during learning walks or initial route
formation (Collett et al., 2023; Collett and Hempel De Ibarra, 2023; Fleischmann et al.,
2017; Freas and Cheng, 2025; Müller and Wehner, 2010; Robert et al., 2018; Stürzl et al.,
2016; Zeil and Fleischmann, 2019). While in our data set scans did rarely extend
beyond 150° away from the goal (Figure 6), our model predicts that when this goal is
weak - such as in entirely naive ants - reversal, smaller saccades and longer
fixations will tend towards 180° from the goal, which might coincide with the origin.
Alternatively, this could be addressed by incorporating either (a) a dual-goal
representation (e.g., one for the feeder and one for the nest) with modulated
weighting, or (b) a dynamic switch of goal orientation during the scan, perhaps based
on distance from home. This raises several empirical predictions. For example, if
ants perform scans while navigating along two-leg outbound routes where the goal is
at 90° to the nest's direction, then fixation patterns during scans may reveal whether
multiple goals influence scanning structure. If the CX indeed holds multiple co-active
directional representations, scan fixations should systematically cluster around both
vectors under uncertainty.

843
Finally, scan termination is currently modeled as a stochastic process, implemented
through a random-rate "freeze release" mechanism to reproduce the observed
Poisson-like distribution of scan durations (Deeti et al., 2023a). However, behavioral
evidence suggests that scan duration is not entirely random. Scans tend to last
longer in contexts of high uncertainty, such as during learning walks or early route
formation, and become shorter or absent on well-learned routes (Wystrach et al.,
2014). This implies that termination likely depends on an internal variable, perhaps
an accumulating "forward drive" or confidence estimate, that builds over time and
releases the motor system from inhibition once a threshold is reached. This
mechanism would parallel the angular drive threshold that governs saccade initiation

in our current model (Figure 4) and resembles integration-to-bound processes
proposed in other species, including mammals (Murakami et al., 2014).
While our model does not include detailed biomechanics of leg coordination,
proprioceptive feedback, or thoracic pattern generators (CPGs), it makes testable
predictions about how angular and forward drive might interact at the motor level.
For instance, we propose that angular drive accumulates continuously, even during
motionless fixations, and that a separate, forward drive determines when locomotion
resumes. This predicts the existence of neurons or motor circuits that integrate
excitatory input over time and release movement when thresholds are crossed,
paralleling the control logic found in *Drosophila* larvae (Rayshubskiy et al., 2025;
Tastekin et al., 2018). Whether this integration over time is achieved at the level of
descending neurons, or downstream in the thoracic CPG themselves remains to be
seen.
867
868 *Conclusions*
869
This study demonstrates that the diverse movement patterns of ants, from brief
scans to full pirouettes and sweeping oscillations, can arise from a single, conserved
neural control architecture: the interaction between central complex (CX) steering
and an intrinsic oscillator in the lateral accessory lobes (LAL). By introducing only
minimal, biologically plausible additions, stochastic inhibition of forward speed, a
saccade-initiation threshold, and forward–angular speed coupling, the model
reproduces fine-scale scanning dynamics seen in high-speed recordings of
Melophorus bagoti, and generalizes across species and behavioral contexts.

A key insight is that forward speed acts as a distributed control dial, gating the
expression of angular movements and enabling smooth transitions between goal-
driven progression and exploratory sampling. This unifying principle offers a simple
yet powerful mechanism for balancing exploitation and exploration, adaptable across
ecological niches and timescales.

Together, these directions open exciting opportunities to link brain models of
sensorimotor control to peripheral control such as thoracic ganglia, refine our
understanding of decision timing, and extend the model to richer multi-goal or
multisensory environments.

887
888 **Materials and Methods**
889
890 *Computational model*
891
The computational model is presented in Figure 1. It was built and run using Matlab®
R2016b and its code is open access and available at:
(https://github.com/antnavteam/CX_oscillator_scans). This step-based model
leverages a model of the CX steering output (Wystrach et al., 2020b) and an intrinsic
oscillator model (Clément et al., 2023), simulating interactions to randomly cease
forward speed to produce a 'scan-like' behaviour. The code's logic is fully
commented and provides a description of the parameters.
899
900 *Behavioural Experiments*

901
902   *species and site*
903
Natural scans were recorded using the red honeypot ant *Melophorus bagoti,* during
the Australian summer from December to February (experienced foragers -
Familiar/Unfamiliar conditions: 2010, see Deeti et al. 2023; inexperienced foragers –
Outbound/Inbound route formation conditions: 2023/2024 summer season) on a field
site (23°45′28.12″S, 133°52′59.77″E) located at the Centre for Appropriate
Technology campus, south of Alice Springs, Northern Territory, Australia. *M. bagoti*
inhabit a visually cluttered environment, exemplified by large amounts of buffel grass
tussocks (*Pennisetum cennchroides*), with scattered eucalyptus trees and bushes.
912
Desert ants navigate alone using multiple concurrent, primarily visual, strategies
(Cheng et al., 2009; Collett and Collett, 2002; Freas and Spetch, 2023; Wehner, 2009;
Wystrach et al., 2012; Zeil, 2023); including path-integration (Cheng et al., 2009; Collett
and Collett, 2002; Wehner and Srinivasan, 2003) and learned views around the nest and
along foraging routes (Collett, 2010; Freas et al., 2017; Wystrach et al., 2012; Zeil, 2023;
Zeil and Fleischmann, 2019).
Two data sets of high speed videos of scanning bouts were used in this project. The
first set of high speed scanning bout videos were collected using inexperienced
foragers, as individuals formed a straight-line route between a feeder and the nest.
In these ants, scanning bouts are known to have a high occurrence near the nest, in
both outbound and inbound ants. Videos were collected both at the onset of their
outbound journey or during the *inbound* trip just before reaching the goal.   The
second was a group of highly experienced ants which were released either along or
near their established foraging route (familiar) or at a distant (unfamiliar) site
(previously collected in Deeti et al. 2023).
928
929   *inexperienced forager procedure*
930
High speed video recordings of inexperienced ants as they scanned while forming
their stereotypical route between and nest and feeder conducted on a single *M.*
bagoti nest in a closed arena with 10cm high walls enclosing the nest and feeder
(Same arena as Freas and Cheng, 2025, with a separate cohort of ants). A stocked
feeder was sunk into the ground 7m from the nest. As we were only interested in
inexperienced foragers with little knowledge of the area beyond the nest, prior to
video recording all foragers that emerged from the nest entrance were marked as
experienced using enamel paint (Tamiya™) and these experienced individuals were
excluded. On day six, non-painted foragers were allowed to exit the nest entrance
and find the feeder. Extending 0.2-1.0m from the nest entrance we spread a thin
layer of white sand along the ground. This sand helped the red ants stand out from
the red ground during filming and pose estimation analysis. Above this site, we
positioned a downward facing Chronos 2.1-HD highspeed camera (KRON Tech),
50cm above the ground (1920×1080pixels, 600fps) with a field of view of
30cm×17cm.
946
For five days prior to recording, One exposure to the feeder, encouraged foragers to
leave the nest entrance in the general direction of the camera's field of view allowing
us to record any scanning behaviours during the first few foraging trips, as the

stereotypical route formed. After collecting ~25 individuals' outbound scans, we
switched to focusing on recently emerged foragers on their inbound trip collecting
any inbound scans, occurring near the nest. Once a forager completed a scan, or
multiple scans within the recording frame, they were collected after they left the area
and marked as completed to prevent repeated recording.
955
956 *experienced foragers procedure*
957
Experienced forager scans dynamics were extracted and analysed from a data set
published in Detti et al. (2023), which focused specifically at spanning metric
distributions; the following summary is further explained there. Two stocked feeders
(cookie pieces) were sunk into the ground 5m from the nest, separated 120° (Right
and Left sites). Ants exited each feeder via a 1m long, 10cm wide channel towards
the nest, slopped up to ground level where ants exited onto a 60 cm × 120 cm
'scanning platform' (wooden board with white paper surface). Similar set-ups (minus
the feeder) were placed at three other local sites (Middle, Opposite, and Far).
966
First, only the Right feeder was stocked with cookies and foragers were allowed to
train along this route, with each visiting individual marked upon their first visit.
Individuals were allowed to train along this route for two days prior to testing,
repeatedly returning to the feeder and leaving via the channel with food. Once highly
experienced, inbound foragers were tested by collection just before entering the nest
and released at one of the four familiar test sites. As each individual reached the
scanning platform they were video recorded using a high-speed camera (Casio EX-
F1, FOV: 30×30 cm at 300 fps). This testing procedure was then repeated with a
second group of foragers trained to the Left feeder. A final set up was placed in a far
(40m), unfamiliar site where a separate group of experienced foragers was tested.
977
978
979
980 *pose extraction*
981
For both sets of scan video datasets, we extracted the body orientations of ants
during fixations to calculate their angle compared to the goal direction (e.g., nest or
feeder) and their duration. Fixations were defined as periods when the ant's head
and body remained stationary between video frames, with no forward or rotational
movement. Scanning can be composed of multiple scanning bouts, with little to no
forward movement within a bout, while the ant rotates in place (saccades), or fixating
in different directions, with a scanning bout ceasing once the ant resumes forward
motion (Deeti et al., 2023a). By collecting body orientation during fixations, we
extracted, the duration of each fixation, the ant's *body orientation* relative to the goal,
the angular change between each fixation (saccade angle) and duration, if the
saccade was 'towards' or 'away' from the goal direction and when the fixation was
followed by saccade that was a reversal in turning direction.
994
995 In the Experienced individuals (300 frames/s), orientation was extracted using a
996 custom MATLAB code (see Deeti et al., 2023 for full description), with fixations
997 identified as stationary periods between frames with body orientation estimated
998 using points at the head and pronotum. In inexperienced forager *f* scans
999 (600frames/s), body orientation during fixations were determined manually using

SLEAP software (Pereira et al., 2022). Fixations 'started' when the head stopped moving between frames and ended with the onset of either forward or rotational movement. Orientation was calculated using two body landmarks: the front centre of the head and the head/body connection (pronotum). As multiple scanning bouts can occur within a single path, each scanning bout began with the first, within frame fixation and ended after forward movement resumed for at least two full leg cycles. Additional fixations within this two-step window were still counted as within the same scanning bout.

*Statistical analysis (*real world *Melophorus bagoti scans)*

    grouping

For experienced foragers, scan data from all familiar release sites (Left, Right, Middle, Opposite) were pooled into a single familiar condition, while scans from the distant site were treated as unfamiliar. For inexperienced foragers, scans were separated into two groups based on if the scan occurred during the outbound and inbound portion of the foraging trip.

    linear mixed-effects models (LMEs)

LME models (Matlab2016) were used to assess the qualitative predictions of the model compared to our real ant data, rather than to optimize model fit. Accordingly, we report F-values derived from ANOVA tables, which test the significance of fixed effects rather than relying on t-values associated with individual parameter estimates. Importantly, this approach fits our qualitative focus. We first run models with interaction between factors. If no interaction was significant, we re-run the model for additive effect only.

Effects on *saccade aptitude* and *fixation duration* were analysed using Linear mixed-effect models with both *individual* and *test conditions* as random effects and with *angular deviation* from the goal direction as a continuous fixed effect and with reversal and Towards/Away (for fixations this corresponds with the upcoming saccade) as categorical fixed effects.

To assess the effect of experience on cumulative orientation vs. the goal during the first reversal of each scan, we conducted LME with the *cumulative orientation* as the dependent variable, with *condition* as the fixed effect and *individual* as a random effect to account for repeated measures across individuals. We ran a second LME to assess the effect of *experience* on the *saccade amplitude*, with *condition* as the fixed effect and *individual* as the random effect to account for repeated measures across individuals.

    fixation duration & saccade amplitude - linear regression

To assess the relationship between fixation duration and saccade amplitude, we first plotted the entire dataset. After observing the general trends, we separated shorter fixation durations (Quartile 1) from the rest of the dataset (Quartile 2-4) and performed separate linear regressions for both subsets. Regression slopes were then tested for significance using *t*-tests on the fitted model coefficients.

1050
1051 *start/stop anytime during oscillation cycle*

1052
1053 To compare the duration via the number of saccades within a sweep between
1054 distinct sweeps (e.g., first vs. second sweep; penultimate vs. final sweep), we used
1055 the Wilcoxon signed-rank tests for paired data.

1056
1057 *first reversal/longest fixation - 'binomial' chi square association*

1058
1059 To assess if reversal and longest fixation duration were associated we tested them
1060 using the chi-square test of independence (χ^2), and odds ratios with 95% confidence
1061 intervals were calculated.

1062
1063

1064 **Acknowledgements**
1065 We are grateful to the Centre for Appropriate Technology for permission to work on
1066 site and access to the nests. We thank Paul Graham for providing the experienced
1067 forager scan dataset, and Leo Clément and Gabriel G. Gattaux for contributing
1068 videos of ant behaviour.

1069
1070 **Funding Statement**
1071 The funders had no role in study design, data collection and interpretation, or the
1072 decision to submit the work for publication.

1073
1074 **Funding Information**
1075 This project was funded by a Macquarie University Research Fellowship
1076 (MQRF0001094) and by the European Research Council (RESIL-ANT - 101125881).

1077
1078 **Competing interests**
1079 No competing interests declared.

1080
Contributions
C.A.F. - Conceptualization, Data curation, Formal analysis, Funding acquisition,
Validation, Investigation, Visualization, Methodology, Writing.

A.W.   -Conceptualization, Data curation, Formal analysis, Modelling, Funding
acquisition, Validation, Investigation, Visualization, Methodology, Writing.

Data Availability Statement

All data, documentation and code is made available online at:
(https://github.com/antnavteam/CX_oscillator_scans')

References

Adden A, Stewart TC, Webb B, Heinze S. 2022. A Neural Model for Insect Steering
Applied to Olfaction and Path Integration. *Neural Comput* **34**:2205–2231.
doi:10.1162/neco_a_01540

Baddeley B, Graham P, Husbands P, Philippides A. 2012. A Model of Ant Route
Navigation Driven by Scene Familiarity. *PLoS Comput Biol* **8**:e1002336.
doi:10.1371/journal.pcbi.1002336

Baird E, Byrne MJ, Smolka J, Warrant EJ, Dacke M. 2012. The Dung Beetle Dance:
An Orientation Behaviour? *PLOS ONE* **7**:e30211. doi:10.1371/journal.pone.0030211

Beetz MJ, Kraus C, el Jundi B. 2023. Neural representation of goal direction in the
monarch butterfly brain. *Nat Commun* **14**:5859. doi:10.1038/s41467-023-41526-w

Berni J. 2015. Genetic Dissection of a Regionally Differentiated Network for
Exploratory Behavior in Drosophila Larvae. *Curr Biol* **25**:1319–1326.
doi:10.1016/j.cub.2015.03.023

Berni J, Pulver SR, Griffith LC, Bate M. 2012. Autonomous Circuitry for Substrate
Exploration in Freely Moving Drosophila Larvae. *Curr Biol* **22**:1861–1870.
doi:10.1016/j.cub.2012.07.048

Bidaye SS, Laturney M, Chang AK, Liu Y, Bockemühl T, Büschges A, Scott K.
2020. Two Brain Pathways Initiate Distinct Forward Walking Programs in
Drosophila. *Neuron* **108**:469-485.e8. doi:10.1016/j.neuron.2020.07.032

Buehlmann C, Fernandes ASD, Graham P. 2018. The interaction of path integration
and terrestrial visual cues in navigating desert ants: what can we learn from path
characteristics? *J Exp Biol* **221**:jeb167304. doi:10.1242/jeb.167304

Buehlmann C, Mangan M, Graham P. 2020. Multimodal interactions in insect
navigation. *Anim Cogn* **23**:1129–1141. doi:10.1007/s10071-020-01383-2

Büschges A, Ache JM. 2025. Motor control on the move: from insights in insects to
general mechanisms. *Physiol Rev* **105**:975–1031. doi:10.1152/physrev.00009.2024

Cheng K. 2024. Oscillators and servomechanisms in navigation and orientation.
Commun Integr Biol **17**:2293268. doi:10.1080/19420889.2023.2293268

Cheng K, Narendra A, Sommer S, Wehner R. 2009. Traveling in clutter: navigation in
the Central Australian desert ant Melophorus bagoti. *Behav Processes* **80**:261–268.

Clément L, Schwarz S, Wystrach A. 2023. An intrinsic oscillator underlies visual
navigation in ants. *Curr Biol* **33**:411–422. doi:10.1016/j.cub.2022.11.059

Collett M. 2010. How desert ants use a visual landmark for guidance along a habitual
route. *Proc Natl Acad Sci USA* **107**:11638–11643. doi:10.1073/pnas.1001401107

Collett T, Collett M. 2002. Memory use in insect visual navigation. *Nat Rev Neurosci*
3:542–52. doi:10.1038/nrn872

Collett T, Graham P, Heinze S. 2025. The neuroethology of ant navigation. *Curr Biol*
35:R110–R124. doi:10.1016/j.cub.2024.12.034

Collett TS, Hempel De Ibarra N. 2023. An 'instinct for learning': the learning flights
and walks of bees, wasps and ants from the 1850s to now. *J Exp Biol* **226**:jeb245278.
doi:10.1242/jeb.245278

Collett TS, Robert T, Frasnelli E, Philippides A, Hempel De Ibarra N. 2023. How
bumblebees coordinate path integration and body orientation at the start of their first
learning flight. *J Exp Biol* **226**:jeb245271. doi:10.1242/jeb.245271

Dauzere-Peres O, Wystrach A. 2024. Ants integrate proprioception as well as visual
context and efference copies to make robust predictions. *Nat Commun* **15**:10205.
doi:10.1038/s41467-024-53856-4

Deeti S, Cheng K. 2025. Desert ants (Melophorus bagoti) oscillate and scan more in
navigation when the visual scene changes. *Anim Cogn* **28**:15. doi:10.1007/s10071-
025-01936-3

Deeti S, Cheng K, Graham P, Wystrach A. 2023a. Scanning behaviour in ants: an
interplay between random-rate processes and oscillators. *J Comp Physiol A* **209**:625–
639. doi:10.1007/s00359-023-01628-8

Deeti S, Islam M, Freas C, Murray T, Cheng K. 2023b. Intricacies of running a route
without success in night-active bull ants (Myrmecia midas). *J Exp Psychol Anim*
Learn Cogn **49**:111–126. doi:10.1037/xan0000350

Demir M, Kadakia N, Anderson HD, Clark DA, Emonet T. 2020. Walking Drosophila
navigate complex plumes using stochastic decisions biased by the timing of odor
encounters. *eLife* **9**:e57524. doi:10.7554/eLife.57524

Fisher, Y. E. (2022). Flexible navigational computations in the Drosophila central
complex. Current opinion in neurobiology, 73, 102514.

Fleischmann PN, Grob R, Wehner R, Rössler W. 2017. Species-specific differences in the fine structure of learning walk elements in Cataglyphis ants. *J Exp Biol* **220**:2426–2435. doi:10.1242/jeb.158147

Franconville R, Beron C, Jayaraman V. 2018. Building a functional connectome of the Drosophila central complex. *eLife* **7**:e37017. doi:10.7554/eLife.37017

Freas CA, Cheng K. 2025. Visual learning, route formation and the choreography of looking back in desert ants, Melophorus bagoti. *Anim Behav* **222**:123125.

Freas CA, Cheng K. 2022. The Basis of Navigation Across Species. *Annu Rev Psychol* **73**:217–241. doi:10.1146/annurev-psych-020821-111311

Freas CA, Congdon JV, Plowes NJR, Spetch ML. 2020. Pheromone cue triggers switch between vectors in the desert harvest ant, Veromessor pergandei. *Anim Cogn* **23**:1087–1105. doi:10.1007/s10071-020-01354-7

Freas CA, Fleischmann PN, Cheng K. 2019. Experimental ethology of learning in desert ants: Becoming expert navigators. *Behav Processes* **158**:181–191. doi:10.1016/j.beproc.2018.12.001

Freas CA, Spetch ML. 2023. Varieties of visual navigation in insects. *Anim Cogn* **26**:319–342. doi:10.1007/s10071-022-01720-7

Freas CA, Whyte C, Cheng K. 2017. Skyline retention and retroactive interference in the navigating Australian desert ant, Melophorus bagoti. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **203**:353–367. doi:10.1007/s00359-017-1174-8

Freas CA, Wystrach A, Narendra A, Cheng K. 2018. The View from the Trees: Nocturnal Bull Ants, Myrmecia midas, Use the Surrounding Panorama While Descending from Trees. *Front Psychol* **9**. doi:10.3389/fpsyg.2018.00016

Freas CA, Wystrach A, Schwarz S, Spetch ML. 2022. Aversive view memories and risk perception in navigating ants. *Sci Rep* **12**:2899.

Green, J., & Maimon, G. (2018). Building a heading signal from anatomically defined neuron types in the Drosophila central complex. Current opinion in neurobiology, 52, 156-164.

Gattaux G, Vimbert R, Wystrach A, Serres JR, Ruffier F. 2023. Antcar: Simple Route Following Task with Ants-Inspired Vision and Neural Model.

Gomez-Marin A, Louis M. 2014. Multilevel control of run orientation in Drosophila
larval chemotaxis. *Front Behav Neurosci* **8**. doi:10.3389/fnbeh.2014.00038

Gomez-Marin A, Stephens GJ, Louis M. 2011. Active sampling and decision making
in Drosophila chemotaxis. *Nat Commun* **2**:441. doi:10.1038/ncomms1455

Goulard R, Heinze S, Webb B. 2023. Emergent spatial goals in an integrative model
of the insect central complex. *PLOS Comput Biol* **19**:e1011480.
doi:10.1371/journal.pcbi.1011480

Graham P, Collett TS. 2002. View-based navigation in insects: how wood ants (
Formica rufa L.) look at and are guided by extended landmarks. *J Exp Biol*
205:2499–2509. doi:10.1242/jeb.205.16.2499

Green J, Vijayan V, Mussells Pires P, Adachi A, Maimon G. 2019. A neural heading
estimate is compared with an internal goal to guide oriented navigation. *Nat Neurosci*
22:1460–1468. doi:10.1038/s41593-019-0444-x

Haalck L, Mangan M, Wystrach A, Clement L, Webb B, Risse B. 2023. CATER:
Combined Animal Tracking & Environment Reconstruction. *Sci Adv* **9**:eadg2094.
doi:10.1126/sciadv.adg2094

Hangartner W. 1967. Spezifitat und Inaktivierung des Spurpheromons von Lasius
fuliginosus Latr. und Orientierung der Arbeiterinnen im Duftfeld. *Z Vgl Physiol*
57:103–136. doi:10.1007/BF00303068

Heinze S. 2017. Unraveling the neural basis of insect navigation. *Curr Opin Insect Sci*
24:58–67. doi:10.1016/j.cois.2017.09.001

Honkanen A, Adden A, da Silva Freitas J, Heinze S. 2019. The insect central complex
and the neural basis of navigational strategies. *J Exp Biol* **222**:jeb188854.
doi:10.1242/jeb.188854

Hulse BK, Haberkern H, Franconville R, Turner-Evans D, Takemura S, Wolff T,
Noorman M, Dreher M, Dan C, Parekh R, Hermundstad AM, Rubin GM, Jayaraman
V. 2021. A connectome of the Drosophila central complex reveals network motifs
suitable for flexible navigation and context-dependent action selection. *eLife*
10:e66039. doi:10.7554/eLife.66039

Husbands P, Shim Y, Garvie M, Dewar A, Domcsek N, Graham P, Knight J,
Nowotny T, Philippides A. 2021. Recent advances in evolutionary and bio-inspired
adaptive robotics: Exploiting embodied dynamics. *Appl Intell* **51**:6467–6496.
doi:10.1007/s10489-021-02275-9

Islam M, Freas CA, Cheng K. 2020. Effect of large visual changes on the navigation
of the nocturnal bull ant, Myrmecia midas | Animal Cognition. *Anim Cogn* **23**:1071–
1080.

Iwano M, Hill ES, Mori A, Mishima Tatsuya, Mishima Tsuneko, Ito K, Kanzaki R.
2010. Neurons associated with the flip-flop activity in the lateral accessory lobe and
ventral protocerebrum of the silkworm moth brain. *J Comp Neurol* **518**:366–388.
doi:10.1002/cne.22224

Izquierdo EJ, Lockery SR. 2010. Evolution and Analysis of Minimal Neural Circuits
for Klinotaxis in *Caenorhabditis elegans*. *J Neurosci* **30**:12908–12917.
doi:10.1523/JNEUROSCI.2606-10.2010

Kanzaki R. 2005. Neural Basis of Odor-source Searching Behavior in Insect Brain
Systems Evaluated with a Mobile Robot. *Chem Senses* **30**:i285–i286.
doi:10.1093/chemse/bjh226

Kanzaki R, Mishima T. 1996. Pheromone-Triggered 'Fiipflopping' Neural Signals
Correlate with Activities of Neck Motor Neurons of a Male Moth, Bombyx mori.
Zoolog Sci **13**:79–87. doi:10.2108/zsj.13.79

Kanzaki R, Nagasawa S, Shimoyama I. 2004. Neural Basis of Odor-Source Searching
Behavior in insect Microbrain Systems Evaluated with a Mobile Robot In: Kato N,
Ayers J, Morikawa H, editors. Bio-Mechanisms of Swimming and Flying. Tokyo:
Springer Japan. pp. 155–170. doi:10.1007/978-4-431-53951-3_12

Kanzaki R, Sugi N, Shibuya T. 1992. Self-generated Zigzag Turning of Bombyx mori
Males during Pheromone-mediated Upwind Walking(Physology). *Zoolog Sci* **9**:515–
527.

Kim SS, Hermundstad AM, Romani S, Abbott LF, Jayaraman V. 2019. Generation of
stable heading representations in diverse visual scenes. *Nature* **576**:126–131.
doi:10.1038/s41586-019-1767-1

Knaden M, Graham P. 2016. The Sensory Ecology of Ant Navigation: From Natural
Environments to Neural Mechanisms. *Annu Rev Entomol* **61**:63–76.
doi:10.1146/annurev-ento-010715-023703

Knight JC, Sakhapov D, Domcsek N, Dewar ADM, Graham P, Nowotny T,
Philippides A. 2019. Insect-Inspired Visual Navigation On-Board an Autonomous
Robot: Real-World Routes Encoded in a Single Layer NetworkThe 2019 Conference
on Artificial Life. Presented at the The 2019 Conference on Artificial Life. Newcastle,
United Kingdom: MIT Press. pp. 60–67. doi:10.1162/isal_a_00141

Kodzhabashev A, Mangan M. 2015. Route Following Without Scanning In: Wilson SP, Verschure PFMJ, Mura A, Prescott TJ, editors. Biomimetic and Biohybrid Systems. Cham: Springer International Publishing. pp. 199–210. doi:10.1007/978-3-319-22979-9_20

Kuenen LPS, Baker TC. 1983. A non□anemotactic mechanism used in pheromone source location by flying moths. *Physiol Entomol* **8**:277–289. doi:10.1111/j.1365-3032.1983.tb00360.x

Le Moël F, Stone T, Lihoreau M, Wystrach A, Webb B. 2019. The Central Complex as a Potential Substrate for Vector Based Navigation. *Front Psychol* **10**:690. doi:10.3389/fpsyg.2019.00690

Le Möel F, Wystrach A. 2020. Opponent processes in visual memories: A model of attraction and repulsion in navigating insects' mushroom bodies. *PLOS Comput Biol* **16**:e1007631. doi:10.1371/journal.pcbi.1007631

Lehrer M. 1993. Why do bees turn back and look? *J Comp Physiol A* **172**:549–563. doi:10.1007/BF00213678

Lent DavidD, Graham P, Collett TS. 2010. Image-matching during ant navigation occurs through saccade-like body turns controlled by learned visual features. *Proc Natl Acad Sci* **107**:16348–16353. doi:10.1073/pnas.1006021107

Li F, Lindsey JW, Marin EC, Otto N, Dreher M, Dempsey G, Stark I, Bates AS, Pleijzier MW, Schlegel P, Nern A, Takemura S, Eckstein N, Yang T, Francis A, Braun A, Parekh R, Costa M, Scheffer LK, Aso Y, Jefferis GS, Abbott LF, Litwin-Kumar A, Waddell S, Rubin GM. 2020. The connectome of the adult Drosophila mushroom body provides insights into function. *eLife* **9**:e62576. doi:10.7554/eLife.62576

Lu J, Behbahani AH, Hamburg L, Westeinde EA, Dawson PM, Lyu C, Maimon G, Dickinson MH, Druckmann S, Wilson RI. 2022. Transforming representations of movement from body- to world-centric space. *Nature* **601**:98–104. doi:10.1038/s41586-021-04191-x

Luo L, Gershow M, Rosenzweig M, Kang K, Fang-Yen C, Garrity PA, Samuel ADT. 2010. Navigational Decision Making in Drosophila Thermotaxis. *J Neurosci* **30**:4261–4272. doi:10.1523/JNEUROSCI.4090-09.2010

Lyu C, Abbott LF, Maimon G. 2022. Building an allocentric travelling direction signal via vector computation. *Nature* **601**:92–97. doi:10.1038/s41586-021-04067-0

Mishima T, Kanzaki R. 1999. Physiological and morphological characterization of olfactory descending interneurons of the male silkworm moth, Bombyx mori. *J Comp Physiol [A]* **184**:143–160. doi:10.1007/s003590050314

Mouritsen H, Feenders G, Liedvogel M, Kropp W. 2004. Migratory Birds Use Head Scans to Detect the Direction of the Earth's Magnetic Field. *Curr Biol* **14**:1946–1949. doi:10.1016/j.cub.2004.10.025

Müller M, Wehner R. 2010. Path Integration Provides a Scaffold for Landmark Learning in Desert Ants. *Curr Biol* **20**:1368–1371. doi:10.1016/j.cub.2010.06.035

Müller M, Wehner R. 2007. Wind and sky as compass cues in desert ant navigation. *Naturwissenschaften* **94**:589–594. doi:10.1007/s00114-007-0232-4

Murakami M, Vicente MI, Costa GM, Mainen ZF. 2014. Neural antecedents of self-initiated actions in secondary motor cortex. *Nat Neurosci* **17**:1574–1582. doi:10.1038/nn.3826

Murray T, Kocsi Z, Dahmen H, Narendra A, Le Möel F, Wystrach A, Zeil J. 2019. The role of attractive and repellent scene memories in ant homing (*Myrmecia croslandi*). *J Exp Biol* jeb.210021. doi:10.1242/jeb.210021

Muser B, Sommer S, Wolf H, Wehner R. 2005. Foraging ecology of the thermophilic Australian desert ant, Melophorus bagoti. *Aust J Zool* **53**:301. doi:10.1071/ZO05023

Mussells Pires P, Zhang L, Parache V, Abbott LF, Maimon G. 2024. Converting an allocentric goal into an egocentric steering signal. *Nature* **626**:808–818. doi:10.1038/s41586-023-07006-3

Namiki S, Iwabuchi S, Pansopha Kono P, Kanzaki R. 2014. Information flow through neural circuits for pheromone orientation. *Nat Commun* **5**:5919. doi:10.1038/ncomms6919

Namiki S, Kanzaki R. 2016. The neurobiological basis of orientation in insects: insights from the silkmoth mating dance. *Curr Opin Insect Sci* **15**:16–26. doi:10.1016/j.cois.2016.02.009

Nicholson DJ, Judd SPD, Cartwright BA, Collett TS. 1999. Learning walks and landmark guidance in wood ants (Formica rufa). *J Exp Biol* **202**:1831–1838. doi:10.1242/jeb.202.13.1831

Olberg RM. 1983. Pheromone-triggered flip-flopping interneurons in the ventral nerve cord of the silkworm moth,Bombyx mori. *J Comp Physiol A* **152**:297–307. doi:10.1007/BF00606236

Pereira TD, Tabris N, Matsliah A, Turner DM, Li J, Ravindranath S, Papadoyannis ES, Normand E, Deutsch DS, Wang ZY, McKenzie-Smith GC, Mitelut CC, Castro MD, D'Uva J, Kislin M, Sanes DH, Kocher SD, Wang SS-H, Falkner AL, Shaevitz JW, Murthy M. 2022. SLEAP: A deep learning system for multi-animal pose tracking. *Nat Methods* **19**:486–495. doi:10.1038/s41592-022-01426-1

Pfeiffer K, Homberg U. 2014. Organization and Functional Roles of the Central Complex in the Insect Brain. *Annu Rev Entomol* **59**:165–184. doi:10.1146/annurev-ento-011613-162031

Rayshubskiy A, Holtz SL, Bates AS, Vanderbeck QX, Serratosa Capdevila L, Rockwell V, Wilson R. 2025. Neural circuit mechanisms for steering control in walking Drosophila. *eLife* **13**:RP102230. doi:10.7554/eLife.102230.3

Robert T, Frasnelli E, De Ibarra NH, Collett TS. 2018. Variations on a theme: bumblebee learning flights from the nest and from flowers. *J Exp Biol* jeb.172601. doi:10.1242/jeb.172601

Sapkal N, Mancini N, Kumar DS, Spiller N, Murakami K, Vitelli G, Bargeron B, Maier K, Eichler K, Jefferis GSXE, Shiu PK, Sterne GR, Bidaye SS. 2024. Neural circuit mechanisms underlying context-specific halting in Drosophila. *Nature* **634**:191–200. doi:10.1038/s41586-024-07854-7

Schwarz S, Clement L, Gkanias E, Wystrach A. 2020a. How do backward-walking ants (Cataglyphis velox) cope with navigational uncertainty? *Anim Behav* **164**:133–142. doi:10.1016/j.anbehav.2020.04.006

Schwarz S, Mangan M, Webb B, Wystrach A. 2020b. Route-following ants respond to alterations of the view sequence. *J Exp Biol* **223**:jeb218701. doi:10.1242/jeb.218701

Seelig JD, Jayaraman V. 2015. Neural dynamics for landmark orientation and angular path integration. *Nature* **521**:186–191. doi:10.1038/nature14446

Shih C-T, Sporns O, Yuan S-L, Su T-S, Lin Y-J, Chuang C-C, Wang T-Y, Lo C-C, Greenspan RJ, Chiang A-S. 2015. Connectomics-Based Analysis of Information Flow in the Drosophila Brain. *Curr Biol* **25**:1249–1258. doi:10.1016/j.cub.2015.03.021

Shiu PK, Sterne GR, Spiller N, Franconville R, Sandoval A, Zhou J, Simha N, Kang
CH, Yu Seongbong, Kim JS, Dorkenwald S, Matsliah A, Schlegel P, Yu Szi-chieh,
McKellar CE, Sterling A, Costa M, Eichler K, Bates AS, Eckstein N, Funke J, Jefferis
GSXE, Murthy M, Bidaye SS, Hampel S, Seeds AM, Scott K. 2024. A Drosophila
computational brain model reveals sensorimotor processing. *Nature* **634**:210–219.
doi:10.1038/s41586-024-07763-9

Steck K, Hansson BS, Knaden M. 2011. Desert ants benefit from combining visual
and olfactory landmarks. *J Exp Biol* **214**:1307–1312. doi:10.1242/jeb.053579

Steinbeck F, Adden A, Graham P. 2020. Connecting brain to behaviour: a role for
general purpose steering circuits in insect orientation? *J Exp Biol* **223**:jeb212332.
doi:10.1242/jeb.212332

Stone T, Webb B, Adden A, Weddig NB, Honkanen A, Templin R, Wcislo W,
Scimeca L, Warrant E, Heinze S. 2017. An Anatomically Constrained Model for Path
Integration in the Bee Brain. *Curr Biol* **27**:3069-3085.e11.
doi:10.1016/j.cub.2017.08.052

Stürzl W, Zeil J, Boeddeker N, Hemmi JM. 2016. How Wasps Acquire and Use
Views for Homing. *Curr Biol* **26**:470–482. doi:10.1016/j.cub.2015.12.052

Sun X, Yue S, Mangan M. 2020. A decentralised neural model explaining optimal
integration of navigational strategies in insects. *eLife* **9**:e54026.
doi:10.7554/eLife.54026

Tarsitano MS, Andrew R. 1999. Scanning and route selection in the jumping spider
Portia labiata. *Anim Behav* **58**:255–265. doi:10.1006/anbe.1999.1138

Tastekin I, Khandelwal A, Tadres D, Fessner ND, Truman JW, Zlatic M, Cardona A,
Louis M. 2018. Sensorimotor pathway controlling stopping behavior during
chemotaxis in the Drosophila melanogaster larva. *eLife* **7**:e38740.
doi:10.7554/eLife.38740

Ugolini A. 2006. Equatorial sandhoppers use body scans to detect the earth's
magnetic field. *J Comp Physiol A* **192**:45–49. doi:10.1007/s00359-005-0046-9

Wallace GK. 1959. Visual Scanning in the Desert Locust Schistocerca Gregaria
Forskål. *J Exp Biol* **36**:512–525. doi:10.1242/jeb.36.3.512

Webb B, Wystrach A. 2016. Neural mechanisms of insect navigation. *Curr Opin*
Insect Sci **15**:27–39. doi:10.1016/j.cois.2016.02.011

Wehner R. 2009. The architecture of the desert ant's navigational toolkit
(Hymenoptera: Formicidae). *Myrmecol News* **12**:85–96.

Wehner R, Fukushi T, Wehner S. 1992. Rotatory components of movement in high
speed desert ants, Cataglyphis bombycina. Presented at the Proceedings of the 20th
Göttingen Neurobiology Conference.

Wehner R, Hoinville T, Cruse H, Cheng K. 2016. Steering intermediate courses:
desert ants combine information from various navigational routines. *J Comp Physiol*
A **202**:459–472. doi:10.1007/s00359-016-1094-z

Wehner R, Michel B, Antonsen P. 1996. Visual Navigation in Insects: Coupling of
Egocentric and Geocentric Information. *J Exp Biol* **199**:129–140.
doi:10.1242/jeb.199.1.129

Wehner R, Srinivasan MV. 2003. Path Integration in Insects. Oxford University Press.

Westeinde EA, Kellogg E, Dawson PM, Lu J, Hamburg L, Midler B, Druckmann S,
Wilson RI. 2024. Transforming a head direction signal into a goal-oriented steering
command. *Nature* **626**:819–826. doi:10.1038/s41586-024-07039-2

Wystrach A. 2023. Neurons from pre-motor areas to the Mushroom bodies can
orchestrate latent visual learning in navigating insects.
doi:10.1101/2023.03.09.531867

Wystrach A, Beugnon G, Cheng K. 2012. Ants might use different view-matching
strategies on and off the route. *J Exp Biol* **215**:44–55. doi:10.1242/jeb.059584

Wystrach A, Buehlmann C, Schwarz S, Cheng K, Graham P. 2020a. Rapid Aversive
and Memory Trace Learning during Route Navigation in Desert Ants. *Curr Biol*
30:1927-1933.e2. doi:10.1016/j.cub.2020.02.082

Wystrach A, Lagogiannis K, Webb B. 2016. Continuous lateral oscillations as a core
mechanism for taxis in Drosophila larvae. *eLife* **5**:e15504. doi:10.7554/eLife.15504

Wystrach A, Mangan M, Philippides A, Graham P. 2013. Snapshots in ants? New
interpretations of paradigmatic experiments. *J Exp Biol* jeb.082941.
doi:10.1242/jeb.082941

Wystrach A, Mangan M, Webb B. 2015. Optimal cue integration in ants. *Proc Biol*
Sci **282**:20151484. doi:10.1098/rspb.2015.1484

Wystrach A, Moël FL, Clement L, Schwarz S. 2020b. A lateralised design for the
interaction of visual memories and heading representations in navigating ants.
doi:10.1101/2020.08.13.249193

Wystrach A, Philippides A, Aurejac A, Cheng K, Graham P. 2014. Visual scanning
behaviours and their role in the navigation of the Australian desert ant Melophorus
bagoti. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **200**:615–626.
doi:10.1007/s00359-014-0900-8

Wystrach A, Schwarz S. 2013. Ants use a predictive mechanism to compensate for
passive displacements by wind. *Curr Biol* **23**:R1083–R1085.
doi:10.1016/j.cub.2013.10.072

Yang HH, Brezovec BE, Serratosa Capdevila L, Vanderbeck QX, Adachi A, Mann
RS, Wilson RI. 2024. Fine-grained descending control of steering in walking
Drosophila. *Cell* **187**:6290-6308.e27. doi:10.1016/j.cell.2024.08.033

Zeil J. 2023. Visual navigation: properties, acquisition and use of views. *J Comp*
Physiol A **209**:499–514. doi:10.1007/s00359-022-01599-2

Zeil J, Fleischmann PN. 2019. The learning walks of ants (Hymenoptera:
Formicidae). *Myrmecol News* **29**:93–110.
doi:10.25849/MYRMECOL.NEWS_029:093