HIF1α-dependent induction of the T-Type calcium channel CaV3.2 mediates hypoxia-induced neuronal hyperexcitability

Anna R. Tröscher [1,2,3], Despina Tsortouktzidis [1,4,5], Franziska Ammer-Pickhardt [2,3], Martin Aichholzer [3,6], Philip-Rudolf Rauch [3,6], Tobias Rossmann [3,6], Nico Stroh-Holly [3,6], Baran Alti[3,6], Andreas Gruber [3,6], Raimund Helbok [2,3], Jan Haubold [2,7], Christian Thome [3,7], Maren Engelhardt [3,7], Tim J. von Oertzen[9], Susanne Schoch[1], Albert J. Becker [1,§], Karen M.J. van Loo [8,10,§]

Institute of Cellular Neurosciences II (IZN II), Medical Faculty, University of Bonn, Bonn, Germany
Department of Neurology, Johannes Kepler University Linz, Kepler University Hospital, Linz, Austria
Clinical Research Institute for Neurosciences, Johannes Kepler University Linz and Kepler University Hospital, Linz, Austria
Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), 53175 Bonn, Germany
University of Cologne, Institute of Molecular and Behavioral Neuroscience, 50937 Cologne, Germany
Department of Neurosurgery, Johannes Kepler University Linz, Kepler University Hospital, Linz, Austria
Institute of Anatomy and Cell Biology, Johannes Kepler University, Linz, Austria
Department of Epileptology and Neurology, RWTH Aachen University, Aachen, Germany
Medical Directorate, University Hospital Würzburg, Würzburg, Germany
Department of Neurosurgery, RWTH Aachen University, Aachen, Germany

§ equally contributed

Correspondence should be addressed to:

Anna Tröscher, PhD
Krankenhausstraße 5
4020 Linz
Austria
Tel.: +43 732 24688755
Email: anna.troescher-boehm@jku.at

Funding

For this project, Anna Tröscher received funding from the Alexander von Humboldt Foundation.

Acknowledgements

 We thank Pia Trebing, Sabine Opitz and Tobias Herbinger for technical assistance.

Author contributions

ART, TJvO, KMJvL, SS and AJB contributed to the conception and design of the study. ART, DT, FAP, JH, CT, KMJvL contributed to the acquisition and analysis of data. ART and JH contributed formal analysis and software. MA, PRR, TR, NSH, BA, AG contributed to sample collection. KMJvL and ART contributed to draft preparation. ME, RH, AJB contributed to reviewing and editing.

Conflict of Interest

TJvO received speaker fees from Angelini-Pharma, Eisai, and UCB, ad board reimbursements from Angelini-Pharma, Jazz Pharma, and Zogenix, all outside of the present study. All other authors have no conflict of interest.

Abstract

Post-stroke epilepsy (PSE) is a major cause of acquired epilepsy in adults, yet the molecular mechanisms linking post-ischemic hypoxia to neuronal hyperexcitability remain poorly understood. The transcription factor hypoxia-inducible factor 1α (HIF1α) is a central mediator of the cellular response to hypoxia and may contribute to epileptogenesis by regulating ion channel expression. Here, we identify the T-type calcium channel CaV3.2 (encoded by *Cacna1h*) as a direct transcriptional target of HIF1α and demonstrate its role in hypoxia-induced network hyperexcitability. Using neuronal cell lines, primary cortical neurons, and organotypic brain slice cultures (OTCs) from mouse and human tissue, we show that HIF1α activation, either through hypoxia or HIF1α overexpression, consistently increases *Cacna1h* expression. In NS20Y cells, overexpression of a normoxia-stable HIF1α variant increased *Cacna1h* promoter activity in both fluorescent and dual-luciferase reporter assays. The same effect was observed in primary cortical neurons, where HIF1α overexpression also elevated network activity measured by multielectrode array recordings. In murine and human OTCs, hypoxia led to marked increase of HIF1α and Cacna1h expression at both transcript and protein levels. Furthermore, oxygen-glucose deprivation followed by reoxygenation (OGD/R) induced a persistent increase in neuronal firing rate, which was recapitulated by HIF1α overexpression alone. Together, these results establish HIF1α as a key transcriptional regulator of CaV3.2 in neurons and reveal a conserved hypoxia-HIF1α-CaV3.2 pathway that enhances neuronal excitability. This mechanism may underlie hypoxia-induced network hyperactivity and contribute to the pathogenesis of post-ischemic epileptogenesis.

Introduction

Post-stroke epilepsy (PSE) is a common complication among stroke survivors and represents a major cause of acquired epilepsy in adults. It is characterized by the occurrence of unprovoked seizures more than seven days after a stroke, affecting approximately 20% of patients following severe ischemic stroke (Beghi et al., 2010; Galovic et al., 2018; Tanaka et al., 2024). Despite its clinical significance, the molecular mechanisms underlying hypoxia-induced epileptogenesis remain poorly understood.

Multiple processes contribute to PSE development, including permanent structural alterations, such as gliotic scarring, neurovascular dysfunction, maladaptive plasticity, and increased neuroinflammation (Galovic et al., 2018; Gruber et al., 2023; Tröscher et al., 2021). In addition, dynamic transcriptional alterations may drive the transition to an epileptogenic state. For example, the transcription factor early growth response protein 1 (Egr1) has been identified as a potential "master switch" of epileptogenesis, promoting neuronal hyperexcitability through the induction of channelopathies such as enhanced T-type calcium channel activity (Becker et al., 2008; van Loo et al., 2019, 2012).

The transcription factor hypoxia-inducible factor 1α (HIF1α) is a key mediator of the cellular response to hypoxia. Under normoxic conditions, HIF1α is continuously produced but rapidly degraded; during hypoxia, inhibition of hydroxylation stabilizes HIF1α, allowing it to dimerize with HIF1β and activate transcription of genes involved in cell death, energy metabolism, and angiogenesis (Sharp and Bernaudin, 2004). HIF1α may also contribute to post-ischemic epileptogenesis by directly regulating ion channel expression. In particular, it may bind to promoter regions of T-type calcium channel genes and initiate an epileptogenic cascade in neurons within the stroke-affected penumbra region (Sellak et al., 2014). Among these channels, CaV3.2 has been repeatedly implicated in chronic epilepsy models (Becker et al., 2008; van Loo et al., 2019, 2012). It is upregulated up to three-fold during the early latent phase of epileptogenesis, leading to increased intrinsic bursting in hippocampal neurons (Sanabria et al., 2001; van Loo et al., 2012).

Outside the brain, hypoxia-induced upregulation of CaV3.2 has been demonstrated in glomus cells, PC12 cells, and cardiomyocytes (Del Toro et al., 2003; Makarenko et al., 2016; Morishima et al., 2021), as well as in pulmonary smooth muscle cells where HIF1α-mediated CaV3.2 expression enhances T-type channel currents (Sellak et al., 2014).

To date, the role of HIF1α in regulating CaV3.2 expression in the brain, and its potential contribution to epileptogenesis, remains unknown. To address this, we investigated the relationship between HIF1α activation and CaV3.2 expression in NS20Y cells, primary neuronal cultures, and organotypic brain slice cultures from both human and mouse tissue. We demonstrate that CaV3.2 expression is increased following hypoxia as well as after HIF1α overexpression alone. Moreover, multiple electrode array recordings revealed a corresponding increase in network excitability, reflected by a higher weighted mean firing rate upon HIF1α activation.

Materials and Methods

Cell Culture, Transfection and Luciferase Assay

NS20Y cells (Sigma, #08062517) were maintained at 37°C, 5% CO_2 in DMEM (Sigma, D6546) supplemented with 10 % heat inactivated FBS (Gibco #10270106), 2 mM L-Glutamine (Gibco #11539876) and 100 units/mL penicillin/streptomycin (Gibco #15140122). Cells were transfected in 24-well plates using Lipofectamine (Invitrogen, #11668030) with the following DNA concentrations per well: pAAV-hSyn-HIF1 300ng or CAG-GFP 300ng in controls, Cacna1h-Luciferase 100ng and Renilla 25 ng (luciferase experiments), or Cacna1h-mRuby 200 ng and CAG-GFP 75 ng (imaging experiments). 48 h after transfection, cells were imaged or collected for luciferase assays. Dual luciferase reporter assays (Promega, #E1910) were performed according to the manufacturers' instructions and firefly/Renilla luciferase ratios were determined using the Glomax Luminometer (Promega).

Cell Imaging

Images were obtained using an inverted phase contrast fluorescent microscope (Zeiss Axio Observer A1 with objectives 20X, LD A-Plan) and processed using Fiji. The cell surface area, set by GFP fluorescent signal, was determined using the Weka trainable segmentation plugin. Integrated density, measured by mRuby fluorescent signal, was normalized to the cell surface area to obtain reported values of integrated density/cell surface.

Neuronal Cultures

All animals were handled according to government regulations as approved by local authorities (LANUV Recklinghausen) and all procedures were performed according to the guidelines of the University Hospital Bonn Animal-Care-Committee as well as the European Directive (2010/63&EU).

Pregnant C57BL/6 mice were deeply anesthetized at embryonic day 16 and sacrificed by decapitation. Abdominal skin and uterus were opened, embryos extracted and kept in PBS (Gibco #14190094) on ice. After decapitation of embryos, brains were removed and divided in sagittal fashion, meninges and cerebellum were removed, and cortex was dissected and placed in cold HBSS (Gibco #14170138). Tissue was digested with Trypsin (Gibco #15090046, 1:10 in HBSS) for 20 min. Next, cells were dissociated with DNase I (Roche #112484932001) and BME (Gibco #41010109), supplemented with FBS. Cell culture plates were coated for 1h at room temperature with Poly-D-lysine (Sigma-Aldrich #11284932001). 70.000 Cells were seeded per well in a 24-well plate. Cells were kept in BME medium (Gibco #21010046), supplemented with 1 % FBS, 0.5 mM L-Glutamine, 0.5 % glucose (Gibco #15384895) and incubated at 37 °C, 5% CO_2. For multielectrode array (MEA) experiments, 40.000 neurons were seeded per well in a 24-well Cytoview-MEA plate (Axion #M384-tMEA-24W).

Constructs

The pAAV-syn-mHif1a-T2A-EGFP plasmid was generated by a two fragment in-fusion cloning (FC; Takara Bio Europe/Clontech). pcDNA3-mHIF-1α-MYC (P402A/P577A/N813A) (Addgene #44028, PMID: 17804822), a plasmid with a normoxic stable variant of HIF1 α, was used as a PCR template for the mHIF1α fragment (Forward primer: 5'-cgcatcgattgaattatggagggcgccgg-3', reverse primer: 5'-tcgctagccagatcctcttctgagatgagtttttgtt-3'), and the pLenti-U6- sgRNALacZ-Syn-dCas9-KRAB-T2A-eGFP plasmid (PMID: 35069110) was used as a PCR template for the T2A-EGFP fragment (forward primer: 5'-ggatctggctagcgagggcagagg-3', reverse primer: 5'-ccgctcggtccgcacgcggggaggcggc-3'). Both fragments were subsequently inserted in the *EcoR*I/*Pml*I sites of the pAAV-hSyn-MCS plasmid (PMID: 35069110).

Viral Production and Neuronal Transduction

AAV1/2 viruses were produced in HEK293-AAV cells (Agilent, #240073) by triple $CaPO_4$ transfection as described previously (van Loo et al., 2012). Transduction of primary neurons was performed at days DIV4 in 24 well plates. For luciferase analyses cells were collected at DIV12-13.

Mouse organotypic brain slice cultures

Nine week old C57BL/J6 mice (males/females) were euthanized by cervical dislocation and brains were quickly extracted. The whole brain was cut coronally in 300 μm sections on a vibratome in oxygenated aCSF (124mM NaCl, 26 mM $NaHCO_3$, 10mM D-Glucose, 3 mM KCl, 1.25 mM NaH_2PO_4, 2 mM $CaCl_2$, 1 mM $MgSO_4$, 10 mM HEPES). Brain sections were split into single hemispheres and put on filter membranes (BRAND®, #782720) in 12-well plates and incubated with organotypic slice medium (66 % Neurobasal-A (Gibco #10888-022), 33 % HBSS (Gibco, #14025-092), 0.625 % D-Glucose (Gibco, #A2494001), 0.01 M HEPES (Gibco, #15630049), 2 mM L-Glutamine (Gibco, #25030-081), 0.5 X B27 (Gibco, #A35828-01), 2 % Anti-Anti (Merck, #A5955) at 35 °C and 5 % CO_2. One hemisphere was used for normoxic control and the other hemisphere for hypoxia. On DIV3, half the slices were put in a hypoxia cell culture system (BioSpherix X3 Xvivo System) with 3 % O_2. After four days in hypoxic conditions, hypoxic and normoxic samples were used for RNA isolation.

RNA isolation, cDNA synthesis and qPCR analysis

RNA from mouse OTCs was isolated with the RNeasy mini kit (Qiagen, #74104) according to the manufacturer's instructions. Subsequently, 40 ng of RNA was reverse transcribed using the Luna RT Master mix (New Englad Biolab, #E3010L). qPCR was performed with the Luna Universal qPCR master mix (New England Biolabs, #M300S) according to the manufacturer's instructions. In short, a 20 µl PCR reaction was used for each sample, including 0.25 µM forward and reverse primers and 1 µl cDNA. Cycling was performed with a 60 sec initial denaturation, followed by 40 rounds of 15 seconds at 95 °C and 45 seconds at 60 °C. Standard melting curve was performed after qPCR amplification. Synaptophysin was used as housekeeping gene for normalization. Primers used were the following: Synaptophysin: 5'-TATCAACCCGATTACGGGCA-3' and 5'-TGGGCTTCACTGACCAGATT-3'; HIF1-alpha: 5'-CCATTTTCAACTCAGGACACTG-3' and 5'-GTGCTCATACTTGGAGGGCT-3'; CaV3.2: 5'-ATGTCATCACCATGTCCATGGA-3' and 5'-ACGTAGTTGCAGTACTTAAGGGCC-3'.

Human organotypic brain slice cultures

All patients gave informed written consent and the study was approved by the Ethics commission of the Johannes Kepler University, under the number EK1250/2023. Patient details can be found in Supplementary Table 1. After surgical resection, human tissue from the access path of tumor resections or epilepsy surgery was quickly transported to the lab in aCSF, where blood vessels and meninges were removed, before it was cut into 300 µm sections in oxygenated aCSF at the vibratome. Sections were transferred to filter membranes in 6-well plates and incubated in organotypic slice medium at 35 °C and 5 % CO_2. At DIV3, half of the human tissue was transferred to the oxygen-controlled incubator and incubated at 3 % O_2 for 4 days (until DIV7). From each patient, one section from each condition was directly fixed for 90 min in 2 % PFA and another section was used for western blot analysis.

Western Blot analysis

Tissue was lysed in RIPA buffer substituted with Halt Protease and Phosphatase Inhibitor Cocktail (100X, Thermo Scientific #78440). For each 1 mg tissue 10 µl buffer was used and tissue was subjected to 3 cycles of freezing and thawing to break up cell membranes. The amount of protein isolated was quantified with the Pierce BCA Protein assay (Thermo Scientific #A65453) and 20 ug were loaded on a 10 % SDS-PAGE gel. Proteins were transferred to a nitrocellulose membrane (Bio-Rad #1620115) and Ponceau S (ThermoFisher #A4000279) staining was performed to visualize total protein content. After imaging of the Ponceau S-stained membranes and destaining, unspecific antibody binding was blocked by 2% fish block (2% fish skin gelatin (Sigma #G7041-100G) in PBS-T). Primary antibodies against HIF1α (Proteintech #20960-1-AP) were used in a concentration of 1:1000 in fish block over night at 4 °C. Biotinylated secondary antibodies (Goat anti-Rabbit IgG (h+L) Cross-Absorbed Secondary Antibody, HRP, #G-21234, 1:5000) were used for 45 min at room temperature. The membrane was then incubated for 5 min in the Immobilon Western HRP Substrate (Millipore #WBKLS0500) and imaged at the Bio-Rad ChemiDoc MP Imaging System. Quantification of bands was performed with the Image Lab software (Image Lab 6.1) from Biorad and whole protein normalization was used.

Immunofluorescence Staining and Imaging

Human OTCs, which showed an upregulation of HIF1α in the western blot analysis, were used for free floating immunofluorescence stainings. To this end, fixed sections were incubated in fish block (1 % BSA (aliquoted/), 0.2 % fish skin gelatin (Sigma #G7041-100G), 0.01 % triton X-100(Merck #102419574) for 1 h at room temperature, followed by overnight incubation at 4 °C with primary antibodies against

NeuN (1:1000, Merck #4182374) and CaV3.2 (1:1000, Invitrogen #8M1106). Secondary antibody incubation was performed with an anti-chicken 647 (1:5000, Invitrogen #2335730) and an anti-rabbit 488 (1:5000, Invitrogen #A-11001) antibody. Sections were mounted on glass slides, covered in ROTI Mount FluorCare (Carl Roth # HP19.1). Sections were imaged at a confocal microscope (Leica Stellaris 5) with the same intensity settings for each paired sample (normoxic and hypoxic). For the quantification of CaV3.2 content, the percentage of the image covered by the positive CaV3.2 signal, using the same threshold settings for each matched sample, was calculated using Fiji software.

Oxygen-glucose deprivation/re-perfusion model (OGD/R)

Primary neuronal cultures were grown under standard conditions until days in vitro (DIV) 11/12 to enable neuronal maturation and network formation. For OGD/R, HBSS was bubbled with a 95 % N_2/5 % CO_2 gas for 5 min with 8-10L/min to deoxygenate the incubation medium. Standard neuronal culturing medium was removed and collected. Normoxic or deoxygenated HBSS were added to control and OGD/R cultures, respectively. OGD/R cultures were placed in the hypoxia incubation chamber (Stemmcell technologies, #27310) and the chamber was filled with a 95 % N_2/5 % CO_2 gas for 4 min with a flowrate of 20L/min. All cells were incubated for 15 min in the incubator, then OGD/R cultures were removed from the chamber and HBSS was removed from all cultures and replaced with the original culturing medium.

MEA Recordings

Between DIV13/14 to DIV 22/24 of neuronal cultures, electrical activity of the cultures was measured using the Maestro Edge MEA system of Axion Biosystems every two days in 24-well plates including 16 electrodes per well (4x4 grid, recording area: 1.1mmx1.1mm, electrode spacing: 350 µm). Plates were kept at 37 °C and 5 % CO_2 during the whole time of the recording. Plates were put in the apparatus, left to equilibrate for 10 min, followed by 10 min recording of spontaneous activity with the Axis Navigator software (version 2.0.4.21). Raw recording data were analyzed with the same software with the following settings: Spike detection: 6 x SD from voltage baseline, minimal burst criterion: minimum of 5 spikes, maximum inter-spike interval: 100 ms. Spikes traces were generated by the Axion Neural Metric Tool (version 3.2.25) and voltage traces using the Axion Data Export Tool (version 3.4.5.). For analyses, only wells with at least 15 active electrodes at the first measurement were used. The weighted mean firing rate was calculated (mean firing rate per well, corrected for the number of active electrodes) and normalized for each well separately to the activity at the first measurement. For analysis, measurements from DIV13-14 (Bin 1), DIV15-17 (Bin 2), DIV 18/19 (Bin 3) and DIV22-24 (Bin 4) were pooled.

Artificial Neural Network Analysis of MEA Data

To predict how closely the behavior of HIF1α-positive and HIF1α-negative cultures resembles that of wild-type cultures under hypoxic or normoxic conditions, an artificial neural network was constructed. As input, we used the raw MEA recordings, which were initially partitioned by well and subsequently divided into training, validation, and test sets. The training and validation sets contained only samples from the classes HYPOXIA and NORMOXIA_WILD_TYPE, whereas the test set comprised NORMOXIA_HIF_NEG and NORMOXIA_HIF_POS samples. The resulting datasets were further segmented into 100 ms intervals, individually classified by the model.
A CNN–LSTM architecture was employed, based on the rationale that the CNN component extracts relevant features, while the LSTM captures temporal dependencies. The detailed model definition can

be seen in the supplementary information. To improve generalization, several strategies were applied, including class reweighting, weight decay, and learning rate scheduling. Importantly, only a randomly selected subset of the total input data was used for training, to exclude overfitting. Importantly, a separate model was trained for each time-bin.

<u>Statistics</u>

Two-way repeated measures ANOVA with Sidaks multiple comparison correction was performed for analysis of MEA recordings (weighted mean firing rate: WMFR) over time. Results from qPCR, western blot and immunofluorescent stainings were analyzed with paired T tests. Analysis of fluorescent reporter activity in NS20Y cells was performed by unpaired T tests. Luciferase dual reporter assay analyses were analyzed using one-sample T Tests. An alpha value of below 0.05 was considered to be statistically significant.

Results

<u>HIF1α regulates CaV3.2 expression in neuronal cell lines</u>

To determine whether HIF1α might act as a transcriptional regulator of the T-type calcium channel CaV3.2 and whether this pathway contributes to hypoxia-induced neuronal hyperexcitability, we first performed an *in silico* analysis of the human *Cacna1h* promoter region. Using the JASPAR database with a relative profile score threshold of 80%, we screened the 2135 nucleotides upstream of the transcriptional start site (ATG) for potential HIF1α binding motifs. This analysis revealed 44 putative HIF1α binding sites within the *Cacna1h* promoter region (Supplementary Table 2), suggesting that *Cacna1h* may be a direct transcriptional target of HIF1α. To determine whether HIF1α directly regulates *Cacna1h* expression in the brain, we used the neuronal cell line NS20Y. Cells were transfected with either a control plasmid (CAG–GFP) or a plasmid encoding a normoxia-stable HIF1α variant (Syn–HIF1α–GFP), together with a fluorescent reporter under the *Cacna1h* promoter (Tsortouktzidis et al., 2022). Cells expressing HIF1α displayed a significant increase in mRuby reporter fluorescence, indicating enhanced *Cacna1h* promoter activity (p = 0.013; Fig. 1A).

To validate these findings, we repeated the experiment using a dual-luciferase reporter assay (Tsortouktzidis et al., 2022). HIF1α overexpression again significantly increased *Cacna1h* promoter-driven luciferase activity compared with controls (p = 0.037; Fig. 1B). These results demonstrate that HIF1α is sufficient to activate *Cacna1h* transcription in neuronal cell lines.

Fig 1: CaV3.2 promoter activation upon overexpression of HIF1α: NS20Y cells transfected with HIF1α show a significant upregulation of CaV3.2 in the (A) fluorescent reporter assay (Two-sided T Test, p = 0.013) (B) as well as in the dual Luciferase reporter assay (one-sample T Test, p = 0.037).

<u>HIF1α induces CaV3.2 promoter activity and increases excitability in primary neurons</u>

We next examined whether this regulatory relationship is maintained in primary neurons. Murine primary cortical cultures were co-transfected with the dual-luciferase reporter system and either a control plasmid or the HIF1α construct. Consistent with the results in NS20Y cells, HIF1α expression significantly increased *Cacna1h* promoter activity as surrogate marker for CaV3.2 mRNA expression in primary neurons (p = 0.016; Fig. 2A).

To determine whether this transcriptional activation translates into altered neuronal activity, we assessed network dynamics by multielectrode array (MEA) recordings following adeno-associated viral (AAV) transduction with either GFP (rAAV-hSyn–GFP) or HIF1α (rAAV-hSyn–HIF1α–GFP). Overexpression of HIF1α led to a significant increase in the weighted mean firing rate (WMFR) compared with controls (two-way ANOVA HIF+/HIF–: F = 4.17, p = 0.049; Fig. 2B,C).

Application of the previously trained CNN–LSTM classifier (developed for hypoxia classification) to this dataset showed that HIF1α-overexpressing samples were predominantly categorized as displaying "hypoxia-like" activity patterns, whereas controls were classified as "normoxia-like," except for bin 4 (Fig. 2D). These findings indicate that HIF1α overexpression alone is sufficient to induce both *Cacna1h* transcription and network hyperexcitability, functionally mimicking hypoxia-induced neuronal activity changes.

Fig 2: Upregulation of CaV3.2 upon HIF1α transduction induces increased WMFR: (A) primary neurons transduced with HIF1α show a significant upregulation of CaV3.2 in the Dual Luciferase reporter assay (one-sample T-Test, p = 0.016). (B) Primary neurons transduced with AAV-hSyn-HIF1α show a significantly increased WMFR compared to primary neurons transduced with a control virus (AAV-hSyn-GFP, two-way ANOVA HIF+/HIF–: F = 4.17, p = 0.049). (C) shows representative images of recordings of AAV-hSyn-GFP controls and AAV-hSyn-HIF1α with activity of every electrode shown in the left panel and bursting behavior on the right panel. (D) Artificial neural network analysis confirmed more hypoxia-like behavior of cells transduced with AAV-hSyn- HIF1α compared to control virus.

Hypoxia increases HIF1α and CaV3.2 expression in organotypic brain slice cultures

To extend these findings to a more physiologically relevant system, we analyzed murine OTCs exposed to hypoxia or maintained under normoxic conditions. Hypoxia led to a fourfold increase in *Hif1a* expression, confirming successful induction of a hypoxic cellular response (p = 0.05; Fig. 3A). In parallel, CaV3.2 expression increased 3.5-fold compared with normoxic controls (p = 0.009; Fig. 3B), indicating that hypoxia enhances both Hif1α and CaV3.2 expression in brain tissue.

We next assessed whether this response is conserved in human OTCs. Western blot analysis confirmed a significant increase in HIF1α protein levels under hypoxic conditions (p = 0.04; Fig. 3C). Immunofluorescence staining further revealed a significant expansion of the CaV3.2-positive area following hypoxia (p = 0.03; Fig. 3D). Representative images illustrate enhanced CaV3.2 immunoreactivity in dendrites of NeuN-positive neurons in hypoxia-treated slices compared with normoxic controls (Fig. 3E).

Together, these results demonstrate that hypoxia induces HIF1α stabilization and CaV3.2 upregulation across cell lines, primary neurons, and organotypic slice cultures, indicating a conserved mechanism by which hypoxic signaling enhances T-type calcium channel expression in neurons.

Fig 3: Upregulation of CaV3.2 after hypoxia in OTCs: In murine OTCs (A) HIF1α shows a significant upregulation upon incubation in hypoxia in qPCR analysis compared to OTCs kept at normoxic conditions (two-sided paired T-Test, p = 0.05). (B) qPCR for Cacna1h revealed a drastic upregulation upon hypoxia compared to OTCs incubated in ambient oxygen concentration (two-sided paired T-Test, p = 0.009). (C) Human OTCs showed a significant upregulation of HIF1α upon exposure to hypoxia compared to OTCs kept at normoxic condition in western blot analysis (two-sided paired T-test, p = 0.04). Representative images of bands used for quantification are shown below (Ponceau S staining for WPN and HIF1α band at 120 kDa). (D) Analysis immunofluorescent stainings of CaV3.2 revealed a higher area fraction of CaV3.2 upon hypoxia compared to slices kept at normoxic conditions (two-sided paired T-Test, p = 0.03). (E) Representative images of immunofluorescent stainings of NeuN positive cells (neurons) and CaV3.2 staining. In normoxic conditions very little CaV3.2 expression is present whereas upregulation of CaV3.2 is visible on the apical dendrite of shown neuron. Scale bar corresponds to 10 μm.

Hypoxia-induced HIF1α activation increases neuronal firing rate

Finally, we investigated the functional consequences of hypoxia-induced HIF1α activation on neuronal network activity. MEA recordings were performed on primary cortical neurons maintained under normoxia or exposed to 15 min of hypoxia followed by reoxygenation (OGD/R). Over the subsequent 10 days, normoxic control cultures showed a mild, maturation-related increase in WMFR, whereas OGD/R-treated cultures exhibited a significantly greater and sustained increase (two-way ANOVA hypoxia × normoxia: F = 7.66, p = 0.009; Fig. 4A,B). In parallel, deep learning analysis using a CNN–LSTM architecture achieved an average ROC–AUC of 0.70 and an F1-score of 0.69 across all validation bins (Fig. 4C), confirming robust classification of hypoxia-induced network hyperactivity. These findings

support a mechanistic link between hypoxia, HIF1α stabilization, and CaV3.2 upregulation, culminating in increased neuronal excitability *in vitro*.

Across complementary experimental systems, from neuronal cell lines to primary neurons and organotypic brain slice cultures, HIF1α activation consistently upregulates CaV3.2 expression and enhances neuronal firing. These data identify HIF1α as a key transcriptional regulator of *Cacna1h* and suggest that HIF1α-driven CaV3.2 induction contributes to hypoxia-induced network hyperexcitability relevant to post-ischemic epileptogenesis.

Fig 4: OGD/R leads to increased neural firing rate: (A) Primary neurons exposed to the OGD/R model showed a significantly increased WMFR compared to neurons kept at normoxic conditions throughout the experiment (two-way ANOVA hypoxia × normoxia: F = 7.66, p = 0.009). (B) shows representative images of recordings of normoxic controls and hypoxia exposed OGD/R cells with activity of every electrode shown in the left panel and bursting behavior on the right panel. (C) Artificial neural network analysis confirmed similar behavior of hypoxia-exposed neurons to HIF+ (AAV-hSyn-HIF1α) cells, whereas neurons kept at normoxic conditions behaved rather like HIF- (AAV-hSyn-GFP) cells.

Discussion

PSE is a severe and frequent complication of ischemic stroke, affecting up to 20% of patients and currently lacking preventive treatment options (Beghi et al., 2010; Galovic et al., 2018; Tanaka et al., 2024). At present, patients are treated only after the first seizure, emphasizing the urgent need to identify molecular mechanisms that could be targeted to prevent epileptogenesis in high-risk individuals. PSE is a multifactorial disorder involving gliotic scar formation, chronic neuroinflammation, and neurovascular dysfunction (Galovic et al., 2018; Gruber et al., 2023; Tröscher et al., 2021). However, cellular and molecular changes that precede network hyperexcitability remain poorly characterized.

Here, we identify HIF1α as a transcriptional regulator of the T-type calcium channel CaV3.2 and demonstrate that this pathway contributes to hypoxia-induced neuronal hyperexcitability. Using complementary models, including NS20Y cells, primary neuronal cultures, and OTCs from mouse and human, we show that activation of HIF1α, either by transient hypoxia or direct overexpression, leads to *Cacna1h* upregulation and increased neuronal firing rates. These findings establish a direct molecular link between hypoxic signaling and the development of hyperexcitable neuronal networks, providing a potential mechanistic basis for post-ischemic epileptogenesis.

HIF1α is a central transcription factor orchestrating cellular adaptation to low oxygen by regulating genes involved in metabolism, angiogenesis, and survival (Sharp and Bernaudin, 2004). In neurons, accumulating evidence suggests additional roles beyond metabolic adaptation, including modulation of synaptic activity and ion channel expression (Arias-Cavieres et al., 2020; Sobrido-Cameán et al., 2025; Wang et al., 2015; Zhang et al., 2023). We demonstrate that *Cacna1h*, which encodes the T-type calcium channel CaV3.2, is transcriptionally activated by HIF1α. Using reporter assays, we show that HIF1α alone, under normoxic conditions, induces CaV3.2 expression in NS20Y cells and primary neurons. This finding aligns with bioinformatic predictions of HIF1α binding motifs within the *Cacna1h* promoter and reveals a previously unknown regulatory mechanism in the central nervous system.

Previous studies have shown that *CaV3.2* is transiently upregulated following pilocarpine-induced status epilepticus and contributes to epileptogenesis by promoting intrinsic bursting and network synchronization (Becker et al., 2008; van Loo et al., 2019, 2012). Similarly, increased *Cacna1h* expression has been reported in peripheral cell types exposed to hypoxia, including glomus cells and cardiomyocytes (Del Toro et al., 2003; Makarenko et al., 2016; Morishima et al., 2021). However, its regulation in neurons during hypoxia has not been demonstrated until now. Our results show that both murine and human OTCs increase HIF1α and CaV3.2 upon hypoxia, establishing this mechanism as a conserved hypoxia-responsive pathway in neuronal tissue.

Functionally, we demonstrate that both hypoxia and HIF1α overexpression increase network activity over time. MEA recordings revealed that neurons exposed to OGD/R develop a persistent increase in the WMFR over 10 days, consistent with long-term changes in excitability rather than transient metabolic effects. This time frame parallels the latent period before clinical seizure onset in PSE, typically defined as seizures occurring more than one week after stroke (Samake et al., 2023). Notably, HIF1α overexpression alone recapitulated this sustained increase in WMFR, confirming that HIF1α activation is sufficient to induce electrophysiological changes characteristic of hypoxia-treated neurons. Artificial neural network analyses further distinguished these conditions, demonstrating consistent classification of HIF1α-overexpressing or hypoxia-treated cultures as "hyperexcitable" or "hypoxia-like." Increased CaV3.2 expression likely contributes to these electrophysiological changes.

CaV3.2 channels mediate low-threshold calcium spikes that promote burst firing and oscillatory activity in cortical and thalamic circuits (Cain et al., 2018; Dumenieu et al., 2018; Harding and Zamponi, 2022; Tringham et al., 2012). Thus, HIF1α-driven upregulation of *Cacna1h* may enhance calcium influx, facilitate activity-dependent transcription, and initiate long-term plasticity programs associated with epileptogenesis.

Our findings suggest that HIF1α–CaV3.2 signaling represents an early molecular event linking hypoxia to network hyperexcitability. In the ischemic penumbra, neurons experience reduced oxygen and glucose supply but remain viable, undergoing structural and transcriptional remodeling that may predispose them to aberrant excitability (Ginsberg, 2003; Stroemer et al., 1995). Persistent HIF1α activation in these neurons could drive CaV3.2-dependent calcium entry and secondary signaling cascades, contributing to maladaptive plasticity and, ultimately, seizure generation. Because approximately 20 % of patients with severe strokes develop PSE (Galovic et al., 2018), the HIF1α–CaV3.2 axis may act in a subset of vulnerable neurons or patients.

In summary, we identify a novel hypoxia–HIF1α–CaV3.2 signaling axis that drives neuronal hyperexcitability and may contribute to PSE. By demonstrating that HIF1α alone is sufficient to induce *Cacna1h* expression and increase neuronal firing, this study expands the known repertoire of HIF1α target genes and establishes CaV3.2 as a mechanistic link between hypoxia and long-term excitability changes. Because both HIF1α and CaV3.2 are pharmacologically targetable, this pathway may represent a promising therapeutic target to prevent or mitigate epileptogenesis following ischemic injury.

References

Arias-Cavieres, A., Khuu, M.A., Nwakudu, C.U., Barnard, J.E., Dalgin, G., Garcia, A.J., 2020. A HIF1a-Dependent Pro-Oxidant State Disrupts Synaptic Plasticity and Impairs Spatial Memory in Response to Intermittent Hypoxia. eNeuro 7, ENEURO.0024-20.2020. https://doi.org/10.1523/ENEURO.0024-20.2020

Becker, A.J., Pitsch, J., Sochivko, D., Opitz, T., Staniek, M., Chen, C.-C., Campbell, K.P., Schoch, S., Yaari, Y., Beck, H., 2008. Transcriptional Upregulation of Cav3.2 Mediates Epileptogenesis in the Pilocarpine Model of Epilepsy. Journal of Neuroscience 28, 13341–13353. https://doi.org/10.1523/JNEUROSCI.1421-08.2008

Beghi, E., Carpio, A., Forsgren, L., Hesdorffer, D.C., Malmgren, K., Sander, J.W., Tomson, T., Hauser, W.A., 2010. Recommendation for a definition of acute symptomatic seizure. Epilepsia 51, 671–675. https://doi.org/10.1111/j.1528-1167.2009.02285.x

Cain, S.M., Tyson, J.R., Choi, H.B., Ko, R., Lin, P.J.C., LeDue, J.M., Powell, K.L., Bernier, L.-P., Rungta, R.L., Yang, Y., Cullis, P.R., O'Brien, T.J., MacVicar, B.A., Snutch, T.P., 2018. CaV3.2 drives sustained burst-firing, which is critical for absence seizure propagation in reticular thalamic neurons. Epilepsia 59, 778. https://doi.org/10.1111/EPI.14018

Del Toro, R., Levitsky, K.L., López-Barneo, J., Chiara, M.D., 2003. Induction of T-type calcium channel gene expression by chronic hypoxia. J Biol Chem 278, 22316–24. https://doi.org/10.1074/jbc.M212576200

Dumenieu, M., Senkov, O., Mironov, A., Bourinet, E., Kreutz, M.R., Dityatev, A., Heine, M., Bikbaev, A., Lopez-Rojas, J., 2018. The Low-Threshold Calcium Channel Cav3.2 Mediates Burst Firing of

Mature Dentate Granule Cells. Cereb Cortex 28, 2594–2609. https://doi.org/10.1093/CERCOR/BHY084

Galovic, M., Döhler, N., Erdélyi-Canavese, B., Felbecker, A., Siebel, P., Conrad, J., Evers, S., Winklehner, M., von Oertzen, T.J., Haring, H.P., Serafini, A., Gregoraci, G., Valente, M., Janes, F., Gigli, G.L., Keezer, M.R., Duncan, J.S., Sander, J.W., Koepp, M.J., Tettenborn, B., 2018. Prediction of late seizures after ischaemic stroke with a novel prognostic model (the SeLECT score): a multivariable prediction model development and validation study. Lancet Neurol 17, 143. https://doi.org/10.1016/S1474-4422(17)30404-0

Ginsberg, M.D., 2003. Adventures in the pathophysiology of brain ischemia: Penumbra, gene expression, neuroprotection: The 2002 Thomas Willis lecture. Stroke. https://doi.org/10.1161/01.STR.0000048846.09677.62

Gruber, J., Gattringer, T., Mayr, G., Schwarzenhofer, D., Kneihsl, M., Wagner, J., Sonnberger, M., Deutschmann, H., Haidegger, M., Fandler-Höfler, S., Ropele, S., Enzinger, C., von Oertzen, T., 2023. Frequency and predictors of poststroke epilepsy after mechanical thrombectomy for large vessel occlusion stroke: results from a multicenter cohort study. J Neurol 270, 6064–6070. https://doi.org/10.1007/S00415-023-11966-X

Harding, E.K., Zamponi, G.W., 2022. Central and peripheral contributions of T-type calcium channels in pain. Mol Brain 15, 1–13. https://doi.org/10.1186/S13041-022-00923-W/FIGURES/3

Makarenko, V. V, Ahmmed, G.U., Peng, Y.-J., Khan, S.A., Nanduri, J., Kumar, G.K., Fox, A.P., Prabhakar, N.R., 2016. CaV3.2 T-type Ca^{2+} channels mediate the augmented calcium influx in carotid body glomus cells by chronic intermittent hypoxia. J Neurophysiol 115, 345–54. https://doi.org/10.1152/jn.00775.2015

Morishima, M., Fujita, T., Osagawa, S., Kubota, H., Ono, K., 2021. Enhanced bdnf actions following acute hypoxia facilitate hif-1α-dependent upregulation of cav3-t-type ca2+ channels in rat cardiomyocytes. Membranes (Basel) 11. https://doi.org/10.3390/MEMBRANES11070470,

Samake, B., Houot, M., Zavanone, C., Vassilev, K., Thivard, L., Herlin, B., Dupont, S., 2023. Late but not early seizures impact negatively early post stroke recovery: A case-control study. Eur Stroke J 8, 784–791. https://doi.org/10.1177/23969873231182493

Sanabria, E.R.G., Su, H., Yaari, Y., 2001. Initiation of network bursts by Ca^{2+}-dependent intrinsic bursting in the rat pilocarpine model of temporal lobe epilepsy. J Physiol 532, 205–216. https://doi.org/10.1111/j.1469-7793.2001.0205g.x

Sellak, H., Zhou, C., Liu, B., Chen, H., Lincoln, T.M., Wu, S., 2014. Transcriptional regulation of α1H T-type calcium channel under hypoxia. Am J Physiol Cell Physiol 307, C648–C656. https://doi.org/10.1152/ajpcell.00210.2014

Sharp, F.R., Bernaudin, M., 2004. HIF1 and oxygen sensing in the brain. Nature Reviews Neuroscience 2004 5:6 5, 437–448. https://doi.org/10.1038/nrn1408

Sobrido-Cameán, D., Coulson, B., Miller, M., Oswald, M.C.W., Pettini, T., Bailey, D.M.D., Baines, R.A., Landgraf, M., 2025. Mitochondrial ROS and HIF-1α signaling mediate synaptic plasticity in the critical period. PLoS Biol 23, e3003338. https://doi.org/10.1371/JOURNAL.PBIO.3003338

Stroemer, R.P., Kent, T.A., Hulsebosch, C.E., 1995. Neocortical neural sprouting, synaptogenesis, and behavioral recovery after neocortical infarction in rats. Stroke 26, 2135–2144. https://doi.org/10.1161/01.STR.26.11.2135,

Tanaka, T., Ihara, M., Fukuma, K., Mishra, N.K., Koepp, M.J., Guekht, A., Ikeda, A., 2024. Pathophysiology, Diagnosis, Prognosis, and Prevention of Poststroke Epilepsy: Clinical and Research Implications. Neurology 102, e209450. https://doi.org/10.1212/WNL.0000000000209450

Tringham, E., Powell, K.L., Cain, S.M., Kuplast, K., Mezeyova, J., Weerapura, M., Eduljee, C., Jiang, X., Smith, P., Morrison, J.L., Jones, N.C., Braine, E., Rind, G., Fee-Maki, M., Parker, D., Pajouhesh, H., Parmar, M., O'Brien, T.J., Snutch, T.P., 2012. T-type calcium channel blockers that attenuate thalamic burst firing and suppress absence seizures. Sci Transl Med 4. https://doi.org/10.1126/SCITRANSLMED.3003120

Tröscher, A.R., Gruber, J., Wagner, J.N., Böhm, V., Wahl, A.S., von Oertzen, T.J., 2021. Inflammation Mediated Epileptogenesis as Possible Mechanism Underlying Ischemic Post-stroke Epilepsy. Front Aging Neurosci 13, 895. https://doi.org/10.3389/FNAGI.2021.781174/BIBTEX

van Loo, K.M.J., Rummel, C.K., Pitsch, J., Müller, J.A., Bikbaev, A.F., Martinez-Chavez, E., Blaess, S., Dietrich, D., Heine, M., Becker, A.J., Schoch, S., 2019. Calcium Channel Subunit $\alpha2\delta4$ Is Regulated by Early Growth Response 1 and Facilitates Epileptogenesis. J Neurosci 39, 3175–3187. https://doi.org/10.1523/JNEUROSCI.1731-18.2019

van Loo, K.M.J., Schaub, C., Pernhorst, K., Yaari, Y., Beck, H., Schoch, S., Becker, A.J., 2012. Transcriptional regulation of T-type calcium channel CaV3.2: bi-directionality by early growth response 1 (Egr1) and repressor element 1 (RE-1) protein-silencing transcription factor (REST). J Biol Chem 287, 15489–501. https://doi.org/10.1074/jbc.M111.310763

Wang, G., Bochorishvili, G., Chen, Y., Salvati, K.A., Zhang, P., Dubel, S.J., Perez-Reyes, E., Snutch, T.P., Stornetta, R.L., Deisseroth, K., Erisir, A., Todorovic, S.M., Luo, J.H., Kapur, J., Beenhakker, M.P., Zhu, J.J., 2015. CaV3.2 calcium channels control NMDA receptor-mediated transmission: a new mechanism for absence epilepsy. Genes Dev 29, 1535. https://doi.org/10.1101/GAD.260869.115

Zhang, Y., Pan, Y. da, Zheng, W. ying, Li, H. yu, Zhu, M. zhen, Ou yang, W. jie, Qian, Y., Turecki, G., Mechawar, N., Zhu, X. hong, 2023. Enhancing HIF-1α–P2X2 signaling in dorsal raphe serotonergic neurons promotes psychological resilience. Redox Biol 69, 103005. https://doi.org/10.1016/J.REDOX.2023.103005