PERINATAL ORGANOPHOSPHATE FLAME RETARDANT EXPOSURE ALTERS

ADULT HPA AXIS FUNCTION AND AVOIDANCE BEHAVIOR IN A SEX-SPECIFIC

MANNER IN MICE

Catherine M. Rojas[a], Julia DeLucca[b], Caylee A, Brown[a], Ali Yasrebi[b], Savannah Chiou[b],

Nicholas T. Bello[b], Troy A. Roepke[a,b]

Affiliations:

[a] *Joint Graduate Program in Toxicology, Rutgers, The State University of New Jersey,*

Piscataway, NJ 08854, USA

[b] *Department of Animal Sciences, School of Environmental & Biological Sciences, Rutgers, The*

State University of New Jersey, New Brunswick, NJ 08901, USA

Corresponding Author:

T.A. Roepke. Department of Animal Sciences, School of Environmental and Biological Sciences,

Rutgers, The State University of New Jersey, 84 Lipman Drive, Bartlett Hall, New Brunswick, NJ,

08901, USA. E-mail address: ta.roepke@rutgers.edu. Phone: (848) 932-9454

Keywords:

flame retardants, endocrine disruption, acute stress, HPA axis

Abbreviations

EPM, Elevated Plus Maze; GD, Gestational Day; LDB, Light-Dark Box; OFT, Open Field Test;

OPFR, Organophosphate Flame Retardants; PBDE, Polybrominated Diphenyl Ether; PND,

postnatal day; TCP, Tricresyl Phosphate; TDCPP, Tris(1,3-dichloro-2-propyl) phosphate; TPP,

Triphenyl Phosphate.

Abstract

Organophosphate flame retardants (OPFRs) are ubiquitous flame-retardant additives with endocrine-disrupting properties. Despite increasing evidence that OPFRs impact neurodevelopment, their effects on the neuroendocrine stress response remain poorly understood. To examine their long-term impact on stress regulation, we treated pregnant C57Bl/6J dams to a mixture of tris(1,3-dichloro-2-propyl) phosphate (TDCPP), triphenyl phosphate (TPP), and tricresyl phosphate (TCP; 1 mg/kg each) from gestational day (GD) 7 through postnatal day (PND) 14. Adult offspring (8-9 weeks of age) were then challenged with acute stressors, including 1 h restraint or a 6-day acute variable stress (AVS) paradigm. Perinatal OPFR exposure produced persistent, sex-specific alterations in the hypothalamic–pituitary–adrenal (HPA) axis and stress-related neurocircuitry. Following 1 h restraint, OPFR-treated females showed heightened serum corticosterone. In addition, gene expression analysis revealed sex-dependent disruptions in key stress-regulatory pathways after OPFR treatment and 1 h restraint in the hypothalamus (*Crhr1, Crhr2, Ptpn5*) and pituitary (*Crhr1, Pomc, Nr3c1*). Females demonstrated more differences in adrenal gene expression related to steroidogenesis (*Mc2r, Cyp11b2*) and catecholamine biosynthesis (*Dbh, Pnmt*), with OPFR-treated groups having blunted responses. OPFR AVS females displayed reduced corticosterone and downregulated *Pacap/Pac1r* expression in the bed nucleus of the stria terminalis (BNST), accompanied by increased behavioral avoidance and immobility. In males, OPFR exposure led to increased BNST *Pacap* and *Pac1r*, expression, along with hyperactivity and avoidance behaviors. Together, these findings demonstrate that early-life OPFR exposure induces lasting, sex-specific dysregulation of the HPA axis and associated stress circuits, highlighting OPFRs as developmental neuroendocrine disruptors with implications for mood and stress-related disorders.

Introduction

The worldwide production of flame retardants has increased to more than 2.39 million tons in 2019 (1). There has been increased usage of flame retardants to adhere to fire safety regulations. While many organohalogen flame retardants, such as polybrominated diphenyl ethers (PBDEs) have been phased out due to overt toxicity and bioaccumulation, organophosphate flame retardants (OPFRs) continue to be used as a "regrettable substitution" (2). OPFRs are a class of endocrine-disrupting chemicals (EDCs) that have demonstrated deleterious effects on neural development, reproduction, and brain and thyroid function (3-6). OPFRs have low degradation potential and persist in household dust since they are not chemically bound to products like furniture, mattresses, and children's products (7). As a result, vulnerable populations are exposed to OPFRs, particularly children who have had 4-5x higher levels of urinary OPFR metabolites compared to their mother which has been associated with adverse neurological outcomes (8,9).

Disruptions in neurological development have been linked to mood disorders and maladaptive behaviors (10). Perinatal exposure to OPFRs have shown to affect neurological development through disrupting neurogenesis, synaptogenesis, and neurotransmission (11). Epidemiological studies have shown that combined exposure to OPFRs is significantly associated with the increased risk of depression (12). In addition, positive associations of prenatal exposure to organophosphate esters (OPEs) has been associated with greater behavioral problem scores in children from mothers with pregnancy-related anxiety (13). There is limited research on how OPFRs specifically affect the neuronal pathways responsible for regulating stress responses. No studies to date have looked at how perinatal OPFR treatment in mice can alter the development of the major stress response pathway, the hypothalamic-pituitary-adrenal (HPA) axis.

Upon activation of the HPA axis by external stressors, the paraventricular nucleus of the

hypothalamus (PVN) synthesizes and releases corticotrophin releasing hormone (CRH) at the

median eminence which enters the hypophyseal portal blood system and reaches the anterior

pituitary (14). CRH binds and activates CRH-receptor 1 (CRHR1), the main receptor responsible

for mediating HPA axis activation at the level of the pituitary. CRHR1 increases pro-

opiomelanocortin (POMC) transcription which is post-translationally cleaved to form

Adrenocorticotropin Hormone (ACTH). ACTH is secreted from the anterior pituitary corticotrope

cells and directly binds to melanocortin 2 receptor (MC2R) located on the surface of the adrenal

zona fasciculata cells. ACTH signaling stimulates an increase in precursor cholesterol for

steroidogenesis by: increasing *de novo* cellular cholesterol synthesis, mobilizing cholesterol esters

stored in lipid droplets, and increasing low-density-lipoprotein (LDL) receptor-mediated

cholesterol uptake (15). Most importantly, ACTH signaling initiates the activation of the single

steroidogenic rate limiting step, the conversion of free cholesterol to pregnenolone by CYP11A1.

The adrenal glands are responsible for producing and releasing hormones during the stress

response. Glucocorticoid production in the adrenal zona fasciculata is species-specific; while

humans primarily produce cortisol, rats and mice predominantly produce corticosterone (CORT)

due to poor expression of CYP17 (15). In parallel, the adrenal medulla's chromaffin cells rapidly

synthesize and release catecholamines as part of the Sympathetic-Adreno-Medullary (SAM) axis.

In response to external stressors detected by the brainstem, sympathetic preganglionic neurons in

the thoracic spinal cord that innervate the adrenal medulla trigger the release of norepinephrine

and epinephrine (adrenaline) (16). Glucocorticoids further amplify this response, as cortisol/CORT

enhances phenylethanolamine-N-methyltransferase (*Pnmt*) activity, an enzyme that converts

norepinephrine to epinephrine (17). Therefore, both axes are essential for mounting a proper stress

response through creating a positive bidirectional feedback loop.

An essential mechanism of the stress response is the system of negative feedback loops

that restore HPA activity to pre-stress conditions. Long feedback loops involve CORT-mediated

inhibition of corticotropin-releasing hormone (CRH) synthesis and secretion in the hypothalamus,

as well as suppression of adrenocorticotropic hormone (ACTH) synthesis in the pituitary. Various

short feedback loops also exist, such as ACTH-mediated inhibition of CRH synthesis in the adrenal

glands (14,18). Glucocorticoid receptors (GRs) further regulate these feedback mechanisms,

facilitating termination of the stress response (19). Organophosphate flame retardants (OPFRs)

have been shown to antagonize GR activity and disrupt steroidogenesis in human adrenocortical

carcinoma (H295R) cells (20-22). Rodent studies likewise report alterations in adrenal size and

function following OPFR exposure (23-26), suggesting that OPFRs may interfere with HPA axis

regulation. However, no studies have examined how perinatal OPFR exposure affects HPA activity

in adulthood. Dysregulation of the stress response is linked to the pathophysiology of stress-related

mood disorders, which are more prevalent in females (27,28). We therefore hypothesized that

perinatal OPFR exposure would disrupt the adult HPA axis response to acute stressors, with female

mice having increased susceptibility.

Acute stressors, such as immobilization or foot shock, are well documented to elevate

circulating corticosterone in rodents (29). Furthermore, acute or subchronic variable stress

paradigms are widely used to model anxiety- and depression-like behaviors in rodent studies (30).

Notably, sex differences in these responses have been documented, likely reflecting the distinct

roles of androgens and estrogens in hypothalamic–pituitary–adrenal (HPA) axis regulation and the

overlapping distribution of gonadal and steroid receptors within stress-related neural circuits

(28,31). Our laboratory has likewise observed sex-dependent effects of chronic stress on behavior,

neuronal activity, and the central stress transcriptome in males and in females dependent on estrous

stage (32-34). Given that organophosphate flame retardants (OPFRs) can disrupt both neural

development and hormonal signaling, these findings raise the question of whether perinatal OPFR

exposure similarly alters central stress responses in a sex-specific manner.

Materials and Methods

Animal Care

All animal treatments and procedures were completed in accordance with National Institutes of

Health guidelines, the Rutgers Institutional Animal Care and Use Committee, and the ARRIVE

guidelines for reporting animal research. Wild-type (WT) C57Bl/6J (Jackson Laboratory), both

female and male, were bred in-house and maintained under controlled conditions: temperature

(25°C), humidity (30-70%), and a 12 h light/dark cycle. Mice had *ad libitum* access to water and

a standard, low-phytoestrogen chow diet (Lab Diets, 5V75). Females were identified by the

presence of a vagina and an estrous cycle, and males were identified by the presence of a penis

and scrotal sac and larger anogenital distance.

Flame retardants

The OPFR mixture we have used previously (35,36) includes tricresyl phosphate (TCP,

AccuStandard, New Haven, CT, CAS#:1330-78-5, 99%), tris(1,3-dichloro-2-propyl)phosphate

(TDCPP, Sigma Aldrich, St. Louis, MO, CAS#:13674-87-8, 95.6%), and triphenyl phosphate

(TPP, Sigma Aldrich, St. Louis, MO; CAS#:115-86-6, 99%). For the mixture, 100 mg of each

OPFR were dissolved in 1 ml of acetone (Sigma-Aldrich), then 100 µl of acetone stock solution

was added to 10 ml sesame oil and allowed to vent for 48 h.

Experimental Design

Females were pair-housed with males for 1 week. Gestational day (GD) 0 was confirmed by the

presence of a vaginal plug. Dams were weighed every 3 days to adjust dosing concentrations and

were randomly assigned to treatment groups: sesame oil vehicle combined with powdered peanut

butter (PB2®, PB2 Foods) or OPFR mixture (see below for preparation of mixture) in sesame oil

combined with powdered peanut butter. Dosing was orally administered at 1 mg of each

OPFR/kg bodyweight/day daily from GD 7 to post-natal day (PND) 14.

Due to limited knowledge of OPFR effects on the stress response, the first set of

experiments that were conducted utilized a single acute stressor. From each litter, 2 male and 2

female pups were weaned at PND 21. One male or female were assigned to a control

(unstressed) group, and their littermates to the restraint group. Control and restrained mice were

housed separately, regardless of maternal treatment. All offspring were maintained on a standard,

low-phytoestrogen chow diet (Lab Diets, 5V75) and housed *with ad libitum* access to food and

water. At 7-8 weeks of age, both control and restrained mice were handled daily for 5 min during

the week prior to sacrifice. At 8-9 weeks of age, restrained mice were subjected to 1 h of restraint

stress using a 50 mL conical tube and euthanized immediately afterwards. All mice were

euthanized by decapitation between 1000-1130 h, with control mice being euthanized before

restrained mice. The 1 h restraint study included the following experimental groups and animals

used: male:oil:con- 13, male:oil:restrained- 11, male:OPFR:con- 11, male:OPFR:restrained- 11,

female:oil:con- 11, female:oil:restrained- 11, female:OPFR:con- 14, and

female:OPFR:restrained- 12.

183   The next set of experiments utilized an acute variable stress (AVS) paradigm. Pups from

each litter (2 males and 2 females) were weaned at PND 21. One male or female were assigned

to a control (unstressed) group, and their single littermate to the acute variable stress (AVS)

group. Control and AVS mice were housed separately, regardless of maternal treatment. All

offspring were maintained on a standard, low-phytoestrogen chow diet (Lab Diets, 5V75) and

housed with *ad libitum* access to food and water. At 7-8 weeks of age, control mice were handled

daily for 5 min for 6 days leading up to behavior testing. Experimental mice were exposed to a 6-

day AVS paradigm (described below). Behavior testing was conducted from Day 7 – Day 10 in

the following order: open field test (OFT), elevated plus maze (EPM), light/dark box emergence

test (LDB), and novelty suppressed feeding (NSF) (see below for more details). On Day 11, all

mice were euthanized by decapitation between 1000-1130 h, with control mice being euthanized

before AVS mice. This study included the following experimental groups and animals used:

male:oil:con- 13, male:oil:AVS- 12, male:OPFR:con- 12, male:OPFR:AVS- 12, female:oil:con-

12, female:oil:AVS- 12, female:OPFR:con- 15, and female:OPFR:AVS- 15. One oil-treated AVS

female mouse died during the last stressor in the startle box chamber. Their weight was

consistent with the previous week, and their right eye was cloudy, thus, their death was most

likely stemmed from other internal complications. All other mice were in good health (according

to IACUC guidelines) and did not lose weight a significant amount of weight until NSF testing.

Acute Variable Stress Paradigm

We reviewed the literature for physiological stressors that were commonly used in acute stress or

fear-inducing paradigms to create our novel paradigm (30,37,38). Day 1 was 1 h restraint using a

50mL conical tube which was placed in their home cage. Day 2 was 1 h cold stress at 4°C in a

fridge only intended for animal use. Day 3 was 1 h platform shake at 100 rpm using a modified

orbital plate shaker. Day 4 was 1 h restraint and day 5 was 1 h platform shake. Day 6 was 30 min

of sound stress using a startle box (MedAssociates; 12,000Hz/105dB/30 ms). The stressors were

repeated randomly so that the same stressor was not done two days in a row. The last stressor

was sound stress since we wanted to collect startle response data to observe differences in

treatment and sex (Fig. S6).

Behavior assays

Adult male offspring were tested before female offspring each day of testing. Mice were

acclimated to the behavior room for at least 24 h prior to the first test. OFT was conducted first

as it is the least stressful, EPM was conducted second to allow for washout between the OFT and

LDB since they are conducted in the same apparatus, and NSF was conducted last to avoid

confounding effects of food deprivation on other tests. The OFT apparatus measures 40 cm long

x 40 cm wide x 40 cm high with an open top and a 64-square grid floor. The mice were placed on

the bottom left corner of the arena at the start of the test and the ANY-maze software measured

activity in the perimeter, corners, 20 cm center, and 10 cm center for 10 min. The EPM apparatus

has 4 arms which are 30 cm long x 5 cm wide which all intersect at the 5 cm square center. The

walls that enclose 2 of the arms are 15 cm tall with open tops. The mice were placed into the

center of the apparatus at the start of the test and the ANY-maze software measured activity in

open arms, closed arms, and center.

The LDB test is conducted in the OFT apparatus with a black insert that is 20 cm long x

40 cm wide x 40 cm tall. The mice are placed on the bottom left corner of the arena and the

ANY-maze software measured activity in the light zone, transition zone, and dark zone. Although

there is no camera inside the dark portion of the apparatus, the software can still measure entries,

exits, and time spent. The NSF is conducted in a white rectangular box with an open top (30 cm

long x 50 cm wide x 16 cm tall) and fresh bedding. The center of the arena contains a pedestal

(10 cm diameter) with a fresh food pellet attached. The mice were placed into the bottom left

corner of the arena and the ANY-maze software measured how long it took for the mice to eat.

Once the mice eat, or 10 minutes have elapsed without eating, the mice are then placed in their

home cage for 5 min.

The OFT, EPM, and LDB chambers were cleaned with MB-10 (Quip labs) spray in

between each animal to ensure that behavior was not influenced by scent. The novel area for

NSF was raked in between each animal, and the apparatus was cleaned with MB-10 and fresh

bedding was added before female mice were tested. The NSF pedestal was also cleaned in

between each animal, with the paper disk and pellet being changed for female mice. All tests

were recorded and analyzed by the ANY-Maze behavior monitoring software (ANY-Maze,

Version 6, Stoelting, USA). Vaginal cytology was performed on the female offspring after each

test to confirm estrous stage.

Exclusion criteria for each test included mice that did not enter all zones at least once or

if the test did not reach full duration. We excluded one mouse from OFT because there were

technical issues where the test ended early by accident. We excluded one OPFR AVS female

mouse in EPM because they only entered one closed arm and stayed in the same spot for the

duration of the test (this same mouse also did not eat in the novel arena). We excluded one mouse

for LDB because there were technical issues where the test ended early by accident. We excluded

one cohort of mice (12 mice) from NSF because their cages were not changed the day before

which caused most of the mice to not eat in the novel area or home cage. Although their food

was removed the day prior, the mice most likely ate their feces or small pellets of food that fell

onto the bottom of their cage.

Tissue preparation and quantitative real-time PCR

Sections of the paraventricular nucleus of the hypothalamus (PVN) and anterodorsal bed nucleus

of the stria terminalis (adBNST) were microdissected from adult offspring under a dissecting

microscope as previously described (35). The pituitary (both anterior and posterior) and both

adrenal glands were collected for RNA extraction as well. All tissues were stored in RNALater

(Life Technologies, Grand Island, NY) at -80°C, and RNA was extracted using RNAqueous-

Micro Kits, Life Technologies. RNA quantity and quality were assessed using a NanoDrop-2000

spectrophotometer (ThermoFisher, Waltham, MA) and RIN was determined using an RNA 6000

Nano kit on an Agilent Bioanalyzer. cDNA was synthesized as previously described (39) and 4

µl of cDNA was amplified by either Power SYBR Green (ThermoFisher) or SSO Advanced

(BioRad, Hercules, CA) Master Mix. Refer to supplemental tables 1 and 2 for the complete list

of primer sequences used in these experiments. Relative gene expression was determined using

the δδCT method calculated by the geomean of reference genes *Actb*, *Gapdh,* and *Hprt*.

Corticosterone Assay

Trunk blood was collected immediately following decapitation and kept at room temperature for

at least 30 min. Serum was then isolated, collected, and stored at -80°C for CORT level analysis.

CORT levels were measured using an ELISA kit (RTC002R; BioVendor) with the lower limit of

detection being 6.1 ng/mL. The intra-assay coefficient of variation range was 5.9%–8.9% and the

inter-assay coefficient of variation range was 7.2%–7.5%. For analysis, a logarithmic regression

was used to convert absorbance values to concentrations in ng/ml.

Data Analysis

Data were stratified by sex and analyzed by 2-way ANOVA (maternal treatment and adult

exposure) followed by Fisher's LSD post-hoc comparison as we were only interested in within

treatment differences using GraphPad Prism 10 (San Diego, CA). Kaplan-Meier survival curves

were analyzed by Log-rank test. All gene expression data were normalized to the oil control

group within each sex. Outliers were determined using the Grubbs test, and values that exceeded

3 SDs ($\alpha<0.01$) above or below the group mean were excluded. Results were considered

statistically significant at $\alpha<0.05$ and data are represented as mean $\pm$ SEM.

Results

For the following findings, all treatment effects reported are effects of perinatal OPFR treatment

and all exposure effects reported are effects of adult restraint or AVS exposure.

Corticosterone and epinephrine measurement after 1 h restraint

CORT analysis revealed an effect of exposure [$F_{(1,27)}=11.33$, $P=0.0023$] where OPFR

restrained females had increased levels compared to OPFR control females ($P<0.01$) (Fig. 1A).

There was also an exposure effect in male mice [$F_{(1,25)}=93.91$, $P<0.0001$] where restrained

mice had higher levels of CORT compared to control mice across treatment groups ($P<0.0001$)

(Fig. 1C). No differences in serum epinephrine concentrations were observed across treatment or

exposure groups (Fig. 1B and 1D). Overall, perinatal OPFR treatment and restraint stress

increased circulating corticosterone levels in both sexes

OPFR treatment altered stress-related genes in the PVN

We selected genes that are part of the central stress response pathway. In females, there was an

exposure effect for *Crh* expression [$F_{(1,27)}=4.685$, $P=0.0394$] (Fig. 2A). A treatment effect was

seen for *Crhr1* expression in females [$F_{(1,28)}=5.729$, $P=0.0236$] (Fig. 2B). An exposure effect

was seen in female *Crhr2* expression [$F_{(1, 28)}=7.198$, $P=0.0121$], with an increase in OPFR

restrained group compared to OPFR control group ($P<0.05$) and a trending decrease in OPFR

control mice compared to oil control ($P=0.0526$; Fig. 2C). No differences were noted for *Pacap*

expression across all groups (Fig. 2D). *Pac1r* expression was affected by treatment

[$F_{(1,26)}=7.692$, $P=0.0101$], with a decrease in expression in OPFR restrained females compared

to oil restrained females ($P<0.05$; Fig. 2E). In addition, *Ptpn5* expression was affected by

treatment [$F_{(1,27)}=8.846$, $P=0.0061$] where there was an increase in expression in OPFR control

females compared to oil control females ($P<0.01$; Fig. 2F).

In males, no differences were seen in *Crh, Crhr2,* and *Pac1r* expression (Figs. 2G, 2I,

2K). We observed treatment [$F_{(1,25)}=33.72$, $P<0.0001$] and exposure [$F_{(1,25)}=9.377$, $P=0.0052$]

effects, as well as an interaction [$F_{(1,25)}=13.02$, $P=0.0013$] effect in *Crhr1* expression with an

increase in OPFR control males compared to OPFR restrained (~2-fold; $P<0.001$) and oil control

($P<0.0001$) males (Fig. 2H). In males, *Pacap* expression was affected by exposure

[$F_{(1,24)}=4.374$, $P=0.0473$] where there was a decrease in OPFR restrained mice compared to

OPFR control mice ($P<0.05$; Fig. 2J). OPFR treatment increased *Ptpn5* expression in control

male mice ($P<0.01$) with a treatment effect of [$F_{(1,26)}=13.47$, $P=0.0011$] (Fig. 2L). Collectively,

these findings show that perinatal OPFR exposure alters the expression of central stress-related

genes in a sex-dependent manner, particularly affecting *Crhr1*, *Crhr2*, and *Ptpn5* in both control

and stressed conditions.

OPFR treatment altered pituitary gene expression independent of sex or exposure

We examined CRH-related genes in the pituitary. In females, *Crhr1* expression was affected by

treatment [$F_{(1,26)}=9.472$, $P=0.0049$] with increased levels in OPFR control females compared to

oil control females ($P<0.05$; Fig. 3A). Treatment effects were more pronounced in male *Crhr1*

expression [$F_{(1,24)}=36.56$, $P<0.0001$] where both OPFR groups were ~2-fold higher compared

to oil groups ($P<0.01$; $P<0.0001$; Fig. 3D). *Pomc* (ACTH) expression was affected by treatment

in females [$F_{(1,24)}=28.70$, $P<0.0001$] and males [$F_{(1,23)}=20.37$, $P=0.0002$] where OPFR

decreased levels in both exposure groups in both sexes ($P<0.01$; Fig. 3B and 3E). In females,

Nr3c1 (GR) expression was affected by treatment [$F_{(1,26)}=5.406$, $P=0.0281$] and exposure

[$F_{(1,26)}=6.724$, $P=0.0154$] (Fig. 3C). In males, *Nr3c1* expression was affected by treatment

[$F_{(1,24)}=23.60$, $P<0.0001$] where OPFR increased expression in both exposure groups ($P<0.01$;

Fig. 3F). These data indicate that OPFR treatment disrupts pituitary expression of key stress-axis

genes, including *Crhr1*, *Pomc*, and *Nr3c1*, in both males and females, independent of stress

exposure.

Perinatal OPFR treatment led to blunted adrenal gene expression in female offspring

Genes along the catecholamine pathway in the adrenals from adult offspring were analyzed. In

females, *Dbh* expression was affected by treatment [$F_{(1,23)}=11.01$, $P=0.0030$] and exposure

[F(1,23)=6.619, P=0.0170] with a ~2-fold decrease in the oil restrained group compared to oil

control and OPFR restrained group (P<0.01; Fig. 4A). *Pnmt* expression was affected by

treatment [F(1,24)=22.39, P<0.0001] and exposure [F(1,24)=4.881, P=0.0370] with an increase

in oil restrained females compared to oil control (P<0.05) and OPFR restrained (P<0.001)

females (Fig. 4B). In addition, OPFR treatment decreased *Pnmt* expression in control females

(P<0.05). No differences were seen for male *Dbh* expression (Fig. 4C). In males, *Pnmt*

expression was affected by exposure [F(1,24)=6.194, P=0.0202] with an increase in oil restrained

mice compared to oil control mice (P<0.01; Fig. 4D).

The steroidogenic pathway (Fig. S1) was analyzed in adrenals collected from adult

offspring. In females, an exposure effect was noted for *Mc2r* expression [F(1,24)=5.793,

P=0.0241] with a decrease in the oil restrained group compared to oil control group (P<0.01; Fig.

S1A). No differences were found for the following genes in females: *Cyp11a1*, *Cyp21a1*, and

Cyp11b1 (Figs. S1B-1D). In females, *Cyp11b2* expression was lower in OPFR control mice

compared to oil control mice (P<0.05; Fig. S1E). In males, no differences in expression were

observed for steroidogenic genes: *Mc2r*, *Cyp11a1*, *Cyp21a1*, *Cyp11b1,* or *Cyp11b2* (Figs. S1F-

1J). These data provide evidence that OPFR exposure and stress affect adrenal gene expression

in females more than males, especially in genes related to catecholamine biosynthesis (*Dbh*,

Pnmt), while steroidogenic gene expression was largely unaltered in males.

AVS exposure affected HPA axis response in OPFR-treated female offspring

CORT analysis revealed an effect of exposure [F(1,28)=6.082, P=0.0200] where OPFR AVS

females had decreased levels compared to OPFR control females (P<0.05) (Fig. 5A). There was

also a trending increase (P=0.0581) in CORT serum in OPFR control females compared to oil

control females which could account for the significant difference within the OPFR group. No

effects were seen for CORT levels in males (Fig. 5B). Overall, OPFR-treated females exhibited

altered HPA axis responsivity, with decreased corticosterone after AVS, while male

corticosterone levels remained unaffected.

Perinatal OPFR treatment has a greater impact on PVN gene expression than AVS exposure

We selected genes that are part of the central stress response pathway and a marker of

inflammation. In females, no effects were seen for *Crh* or *Crhr2* expression (Fig. S2A and S2B).

A treatment effect was seen for *Crhr1* expression in females [$F_{(1,19)}=8.095$, $P=0.0103$], with a

decrease in OPFR control group compared to oil control group ($P<0.05$; Fig. 6A). Treatment

effects were seen in female *Pacap* and *Pac1r* expression [$F_{(1,19)}=27.44$, $P<0.0001$;

$F_{(1,19)}=28.03$, $P<0.0001$], with a ~2-fold increase in OPFR groups compared to oil groups for

both genes ($P<0.01$; Fig. 6B and 6C). *Ikk,* a regulator of the inflammatory NF-kB pathway, was

affected by exposure [$F_{(1,17)}=10.04$, $P=0.0056$], including an increase in oil AVS females

compared to oil control females ($P<0.05$; Fig. S2C). No effects were seen in *Ptpn5* expression

for females or males (not shown).

In males, no effects were seen in *Crh, Crhr2,* or *Ikk* expression (Figs. S2D-S2F). We

observed a treatment effect for *Crhr1* expression [$F_{(1,19)}=15.20$, $P=0.0010$] with a decrease in

OPFR AVS males compared to oil AVS males ($P<0.001$; Fig. 6D). Similar to females, treatment

effects were seen in male *Pacap* and *Pac1r* expression [$F_{(1,20)}=37.51$, $P<0.0001$;

$F_{(1,20)}=43.39$, $P<0.0001$], with a ~2-fold increase in OPFR groups compared to oil groups for

both genes ($P<0.001$; Fig. 6E and 6F). Collectively, these findings show that perinatal OPFR

treatment alters expression of stress-related and inflammatory genes in the PVN, with increased

Pacap pathway expression in both sexes and sex-specific effects on *Crhr1* and *Ikk*.

Perinatal OPFR treatment and AVS exposure disrupted PACAP pathway in the BNST of female

offspring

We also investigated the expression of stress-related genes in the adBNST since GABAergic

projections from the adBNST to the PVN are involved in negative feedback of the HPA axis

(31). In females, no differences were seen in *Crh* or *Crhr2* (Fig. S3A and S3B). *Crhr1* expression

was affected by treatment [$F_{(1,24)}=4.835$, $P=0.0378$] where OPFR control females had increased

levels compared to oil control females ($P<0.05$; Fig. 7A). We observed treatment, exposure, and

interaction effects in female *Pacap* expression [$F_{(1,25)}=4.381$, $P=0.0467$; $F_{(1,25)}=50.20$,

$P<0.0001$; $F_{(1,25)}=5.899$, $P=0.0227$], where OPFR treatment increased expression by ~3-fold

($P<0.01$) and AVS decreased expression within the OPFR group ($P<0.01$; Fig. 7B). *Pac1r*

expression was affected by treatment [$F_{(1,24)}=177.1$, $P<0.0001$], where OPFR treatment

increased expression by ~2.5-fold and AVS decreased expression within the OPFR group

($P<0.05$; Fig. 7C). *Ikk* expression was affected by treatment and exposure [$F_{(1,25)}=5.390$,

$P=0.0287$; $F_{(1,25)}=5.636$, $P=0.0256$], with an upregulation in OPFR control and oil AVS females

compared to oil control females ($P<0.01$; Fig. S3C).

In males, no pairwise differences were seen in *Crh* or *Crhr1* (Fig. S3D and S3E);

however, there was a treatment effect for *Crhr1* [$F_{(1,25)}=4.275$, $P=0.0492$]. We observed a

decrease in *Crhr2* expression after AVS exposure in OPFR males ($P<0.05$; Fig. 7D). *Pacap* and

Pac1r expression was affected by treatment [$F_{(1,24)}=55.18$, $P<0.0001$; $F_{(1,26)}=65.17$,

$P<0.0001$], where OPFR treatment increased expression by ~3-fold ($P<0.001$; Fig. 7E and 7F).

Ikk expression was affected by treatment [$F_{(1,26)}=87.77$, $P<0.0001$], where OPFR increased

overall expression ($P<0.0001$; Fig. S3F). These data provide evidence that OPFR exposure

dysregulates stress and inflammatory signaling in the adBNST in a sex-dependent manner,

particularly enhancing *Pacap* pathway and *Ikk* expression, and altering *Crhr1* and *Crhr2*

expression.

Sex specific response to OFT after OPFR treatment and AVS exposure

The OFT assesses exploratory and avoidant behavior. In females, overall locomotion was not

affected (Fig. S4A and S4B). In males, locomotion was affected by AVS exposure, with an

increase in distance traveled ($F_{(1,44)}=30.00$, $P<0.0001$) and mean speed ($F_{(1,44)}=30.51$,

$P<0.0001$) regardless of treatment ($P<0.01$; Fig. S4A and S4B). In females, there was no effects

seen for time immobile in the perimeter (Fig. 8A), however there was a significant effect of

exposure [$F_{(1,47)}=7.185$, $P=0.0101$] where OPFR AVS mice spent a longer time immobile in the

corners than OPFR control mice ($P<0.05$; Fig. 8B). In males, an exposure effect [$F_{(1,44)}=6.274$,

$P=0.0160$] was noted for time immobile in the perimeter, where OPFR AVS mice spent less time

immobile in the perimeter compared to OPFR control mice ($P<0.05$; Fig. 8C). We also observed

effects on treatment [$F_{(1,39)}=8.729$, $P=0.0053$] and exposure [$F_{(1,39)}=4.416$, $P=0.0421$] for

time immobile in the corners, with OPFR AVS mice spending less time immobile than oil AVS

mice ($P<0.05$; Fig. 8D). These findings demonstrate that AVS increases avoidance behavior and

alters locomotion in a sex-specific manner, with females showing increased corner behavior and

males displaying heightened activity and mobility.

AVS increased activity in EPM in control male offspring

The EPM is another test that evaluates avoidance behavior. In males, exposure effects were noted

for distance and mean speed [$F_{(1,45)}$=7.650, P=0.0082; $F_{(1,45)}$=7.594, P=0.0084], where AVS

increased the locomotion of oil males (P<0.05; Fig. 9E and 9F). Additional exposure effects were

observed for closed arms entries and distance traveled [$F_{(1,45)}$=7.024, P=0.0111; $F_{(1,44)}$=12.96,

P=0.0008]. Specifically, AVS increased closed arm entries and distance traveled in oil males, and

increased distance traveled of OPFR males (P<0.05; Fig. 9G and 9H). Together, these results

suggest that AVS enhances avoidance behavior in males, particularly increasing closed-arm

activity and overall locomotion in oil- and OPFR-treated mice.

Perinatal OPFR treatment and AVS exposure led to sex-specific responses in LDB

The LDB examines exploratory and avoidance behavior. In females, significant effects of

exposure were noted for light zone entries and exits [$F_{(1,45)}$=7.351, P=0.0095; $F_{(1,45)}$=7.941,

P=0.0072], with AVS decreasing both entries and exits in OPFR groups (P<0.01; Fig. 10A and

10B). The opposite was observed for light zone entries and exits in OPFR males (P<0.05; Fig.

10E and 10F). In females, exposure effects were noted for dark zone entries and exits

[$F_{(1,46)}$=4.938, P=0.0312; $F_{(1,46)}$=4.359, P=0.0424], with AVS decreasing both entries and

exits in OPFR groups (P<0.05; Fig. 10C and 10D). In males, exposure effects were also noted for

dark zone entries and exits [$F_{(1,45)}$=4.103, P=0.0488; $F_{(1,45)}$=4.195, P=0.0464]; however, AVS

increased entries and exits in OPFR groups (P<0.05; Fig. 10G and 10H). These results show that

AVS exposure modulates light/dark zone transitions in a sex- and treatment-dependent manner,

decreasing exploratory behavior in OPFR females and increasing it in OPFR males.

Perinatal OPFR treatment decreases latency to feed in males only

The NSF is a conflict-based test which assesses how long it takes for the mouse to eat the pellet

in the center of a brightly lit novel area, with longer latency to eat indicating increased avoidance

behavior. In males, we observed an exposure effect for amount of pellet eaten [$F_{(1,39)}=9.672$,

P=0.0035], where oil control males ate more than oil AVS and OPFR control males (P<0.05; Fig.

S5D). Latency to eat in the novel arena was affected by treatment [$F_{(1,39)}=5.647$, P=0.0225],

with OPFR decreasing latency in control males (P<0.05; Fig. 11E). The probability to eat as

shown by the Kaplan-Meier curve (Fig. 11F), had an effect of treatment in the novel arena where

OPFR control males were more likely to eat than oil control males (P= 0.0179). These data

suggest that OPFR alters conflict-based avoidance behavior in males, with decreased latency to

feed and increased likelihood of eating under anxiogenic conditions.

Discussion

While the global use of brominated flame retardants has declined, the use of OPFRs has

increased in countries such as the US and China (40). A report from Albany, New York, USA,

estimates daily indoor OPFR exposure via inhalation of air, dermal sorption, and dust ingestion

at 149, 279, and 390 ng/kg bw/day, respectively (41). Although the combined effects of perinatal

OPFR treatment and acute stress have not been studied, *in vitro* evidence has shown that PBDEs

increase cortisol and aldosterone production, while TPP decreases their production (26,42). Rat

studies using various flame retardants have also shown mixed results, with some reporting

increased adrenal weight and CORT levels (24,26), while others report a dose-dependent

decrease in adrenal weight and CORT in males only (23). Human data have linked increased

OPE diester concentrations with 18%-41% higher cortisol and cortisone levels (43). In our study,

we observed sex-dependent disruption of the HPA axis by perinatal OPFR treatment.

Both females and males demonstrated exposure effects for CORT levels that were

elevated after 1 h restraint. However, only OPFR-treated females exhibited a significant increase

in CORT, while oil-treated females did not. This suggests that perinatal OPFR treatment may

exacerbate stress response in females. Electrophysiological recordings have shown that treatment

with 17β-estradiol (E2) increase PVN CRH neuronal activity and correlate with CORT levels,

therefore it is possible that OPFR treatment is eliciting similar effects to endogenous hormones

(44). The higher variability in CORT levels among oil control females, coupled with a slightly

lower (though not significant) average in OPFR controls, may contribute to this treatment-

specific difference. Previous studies have shown that female rodents have more pronounced

stress responses than males which may also account for these effects (45,46). Sex differences in

CORT levels have also been noted in humans, where women with depression have been reported

to have higher cortisol levels than non-depressed women (47). No differences were noted for

epinephrine levels between groups, likely because levels had returned to baseline after 1 h of

restraint. Previous work in male mice shows epinephrine peaking at ~40 minutes post-stress (48),

which aligns with this finding.

In the PVN, an exposure effect was noted for *Crh* and *Crh2* in females, with there being a

significant increase in *Crhr2* following stress in the OPFR group. CRH R2 is important for

ACTH production and recovery of the HPA axis recovery, as previously shown in *Crhr2*

knockout mice (49). This response may be exacerbated in OPFR-treated females due to trending

decrease in OPFR-control expression of *Crhr2*. We also observed a treatment-related decrease in

Pac1r, which encodes a receptor for PACAP, a neuropeptide that activates CRH neurons and

stimulates the stress axis (50). The observed increase in *Crhr2* and decrease in *Pac1r* may reflect

HPA axis recovery processes occurring during the restraint period, which aligns with prior

reports showing CORT peaking 30–60 minutes into restraint and recovery initiating while the

stressor remains present (51). In females and males, increased *Ptpn5* expression in OPFR control

mice suggests long-term effects on synaptic plasticity in the PVN resulting from perinatal OPFR

exposure.

In males, we saw pronounced differences in *Crhr1* expression, with elevated levels in

OPFR-control animals compared to both oil control males and OPFR-restrained males. While no

EDC studies have previously reported increased *Crhr1* expression in the PVN, one study of

gestational intermittent hypoxia (GIH) found increased protein levels of CRHR1 in the PVN of

embryonic day (E) 19 male mice, but not female mice (52). Interestingly, they associated GIH

reprogramming of CRH R1 signaling in the PVN to anxiogenic effects in adult rat offspring. Our

previous work found anxiogenic effects of perinatal OPFR treatment in the open field test and

anxiolytic effects in the elevated plus maze in unrestrained male offspring (36). The elevated

Crhr1 expression observed here may reflect fetal programming that sensitizes the pituitary to

CRH. Decreases in both *Crhr1* and *Pacap* in OPFR-restrained males may indicate negative

feedback mechanisms within the central stress response pathway.

In the pituitary, significant treatment effects were observed in females and males across

all genes analyzed. Perinatal OPFR treatment increased *Crhr1* expression but decreased *Pomc*

expression, likely due to elevated *Nr3c1* (GR) expression, which negatively regulates *Pomc*. GRs

are activated by high CORT levels and partake in the negative feedback loop that terminates the

stress response (53). They are known targets of EDCs, including flame retardants (20,54-56). A

developmental study conducted in zebrafish larvae showed dose-dependent increases in GR

mRNA expression following TDCPP and TPP treatment (57). In our study, *Nr3c1* expression was

affected by both treatment and exposure in females, while treatment effects were more prominent

in males with pairwise differences. These increases in *Nr3c1* may alter the regulation of negative

feedback in the HPA axis. Future studies could also evaluate GR expression in the hypothalamus,

where its role is to reduce *Crh* expression. The OPFR-induced effects in the pituitary were

consistent across sexes.

In the adrenal glands, no steroidogenic changes were observed in males, whereas a few

were found in females. Specifically, *Mc2r* (ACTH receptor) expression decreased in oil

restrained females, but not in OPFR-treated counterparts, suggesting that perinatal OPFR

treatment may impair adrenal sensitivity to ACTH. Additionally, *Cyp11b2* expression was

reduced in OPFR control females, possibly reflecting upstream functional disruptions, as a slight

decrease in OPFR control female CORT was also observed. Human adrenal cell studies (H295R,

female) have shown increased expression of *Cyp11b1* and *Cyp11b2* following flame retardant

exposure (21,22). Although another developmental study using a mixture of EDCs (phthalates,

pesticides, and bisphenol A) showed a dose-dependent decrease in *Cyp11b1* adrenal expression

in female adult offspring that underwent behavioral testing (58). It is important to note that

steroidogenic pathways in rodents differ from humans in that 1) rodents mainly synthesize CORT

due to poor Cyp17 expression, 2) rodents utilize HDL cholesterol via the SR-B1 pathway while

humans utilize LDL cholesterol via the endocytic pathway as steroid precursors, and 3) mice do

not have a zona reticularis and as a result, do not produce androgens (15).

Despite the differences between mouse and human adrenal cortices, adrenal medullary

function is conserved across both species. During development, neuroendocrine cells from the

neural crest invade the adrenal cortex by embryonic day (E) 11 and differentiate into chromaffin

cells, completing development by E15 (59). Upon activation of the sympathetic-adrenal-

medullary (SAM) axis, these cells release norepinephrine (noradrenaline) and epinephrine

(adrenaline), otherwise known as the "fight-or-flight" hormones. These catecholamines enhance

cardiovascular output and mobilize energy stores (60). *Dbh* converts dopamine into

norepinephrine and *Pnmt* then converts norepinephrine to epinephrine. In females, 1-h restraint

decreased *Dbh* expression, but this was blunted by OPFR treatment. Similarly, *Pnmt* expression

increased with stress in oil-treated females, but this response was also blunted in OPFR-treated

mice. While *Dbh* expression for OPFR-treated and oil control females are at similar levels, *Pnmt*

expression of OPFR-treated females fell below that of controls. This could relate to slightly

reduced serum CORT levels in OPFR females, as CORT regulates PNMT activity. These

treatment-related changes in females likely stem from disrupted embryonic development of the

adrenal medulla. In males, we observed increased *Pnmt* following stress, as expected, though the

response was blunted in OPFR-treated males, although not statistically significant. These

patterns suggest medullary development may also be impacted in males, though to a lesser

degree.

Utilizing a 6-day AVS paradigm, we revealed that perinatal OPFR treatment can lead to

deficits in central stress response and behavior in adulthood. In OPFR AVS females, we observed

blunted levels of CORT, BNST *Pacap* and *Pac1r* mRNA expression, as well as increased

avoidance and immobility suggesting a weakened HPA axis response. In males, we saw that

overall perinatal OPFR treatment increased *Pacap*, *Pac1r*, and *Ikk* in the BNST. For male

behavior, AVS increased avoidance while OFPR increased hyperactivity. Overall, our study

demonstrated striking sex differences in response to adult acute stress after perinatal OPFR

treatment.

For CORT levels, we observed sex-specific vulnerability, with OPFR AVS females

showing reduced CORT. Additionally, there was a trending increase in CORT in OPFR control

females relative to oil controls, which is driving this exposure effect. Other studies have also

reported elevated CORT following prenatal EDC treatment (61). It is important to note that these

samples were collected 5 days after the last stressor, during the morning when CORT levels are

lowest in rodents. Future studies may therefore benefit from measuring CORT immediately after

the last stressor through submandibular bleeds to determine immediate effects on the stress

response. Especially since other studies have typically seen increases in CORT in females after

acute stress paradigms (30).

For the neural stress circuitry, treatment differences were noted in both females and

males, especially with *Pacap* and its receptor. We saw ~2-fold increases in the PVN and ~3-fold

increases in the BNST after OPFR treatment which shows excitability of PACAP pathways in the

central stress response. The upregulation in this pathway indicates increased avoidance, which

was noted in both groups for behavior. PACAP-deficient mice have demonstrated attenuation of

stress-induced responses, including diminished CORT release and reduced activation of the HPA

axis (62,63). This further supports the idea that PACAP is crucial for promoting the stress

response. Interestingly, we observed a decrease in *Pacap* and its receptor after AVS exposure in

OPFR-treated females. This reduction is consistent with the observed decrease in CORT levels in

females and aligns with existing literature linking PACAP signaling to HPA axis regulation (50).

Notably, this exposure-related change was not observed in males within the BNST, supporting

both our previous findings and other research demonstrating BNST sex-specific response to

stress (34,64).

Perinatal OPFR treatment also led to increases in the inflammatory mediator *Ikk*, a key

component of the NF-κB pathway. Specifically, we observed elevated *Ikk* expression in oil AVS

females in both the PVN and BNST, aligning with existing literature that highlights the role of

proinflammatory mediators in the stress response, particularly following chronic restraint stress

(65). Importantly, we did not observe these exposure-related increases in Ikk in males,

suggesting a sexually dimorphic stress response which has previously documented in other 6-day

stress models, where female mice show greater susceptibility to depression-like behaviors

(30,66). Interestingly, we saw significant treatment differences on *Ikk* in males and females only

in the BNST. OPFRs are known to induce neurotoxicity through inflammatory pathways (67-70).

In addition, NHANES epidemiologic data have demonstrated a significant positive association

between urinary OPFR concentrations, elevated inflammatory markers, and the risk of

depression in humans (12). An additional mouse model of *in utero* OPFR exposure also

demonstrated placenta inflammation, disrupted neuronal development and synaptic transmission,

and persistent avoidant behaviors in adulthood (71). Together, these findings suggest that early-

life OPFR exposure may produce long-lasting neurobehavioral effects by promoting persistent

inflammation. More inflammatory markers further down the Ikk pathway, such as IL-1, should

be explored in future studies to elucidate this mechanism of OPFR-induced neurotoxicity.

The behavior tests that were conducted demonstrated pronounced sex-specific effects in

response to perinatal OPFR treatment and AVS exposure. In the OFT, OPFR AVS females had

increased immobility in the corners, indicative of increased avoidant behavior. This may be due

to hypoactivation of the HPA axis, although more measurements of CORT levels would need to

be taken immediately after stressors to confirm. For males, we saw an increase in locomotion

and corner time after AVS. Perinatal OPFR treatment induced hyperactivity, as we also observed

a trending increase in time mobile in OPFR AVS males. The combined effects of OPFR and AVS

made males more hyperactive in the perimeter, noted by reduced immobility in the perimeter and

corners and pattern of movement in the heat maps. We have previously also seen that perinatal

OPFR treatment leads to hyperactivity in males (36).

In the EPM, males showed increased locomotion, closed arm entries and distance after

AVS, confirming hyperactivity and avoidance seen in OFT. In the LDB, OPFR AVS females had

reduced dark and light entries/exits whereas OPFR AVS males had the opposite effect. Male

OPFR controls had a trending decreasing in distance traveled in the light zone demonstrating

avoidance, however AVS abolished this effect to levels comparable to oil AVS males. In the NSF,

we saw treatment differences where OPFR males had reduced latency to eat, reflecting reduced

avoidance. AVS exposure eliminated this treatment difference, supporting previous tests of

increased avoidance in OPFR AVS groups.

Overall, the behavior tests indicate that OPFR AVS females display increased avoidance

and reduced activity. Increased immobility may reflect a passive reaction coping mechanism,

often associated with depressive-like behavior, and similar increases in immobility have been

observed in females exposed to a low-dose mixture of EDCs (61). Therefore, future studies

should investigate whether perinatal OPFR treatment combined with AVS exposure affects

motivation-related behaviors, particularly given that the BNST is implicated in effort-based

decision and motivation behaviors (72). Future work could determine whether OPFR treatment

and AVS exposure increase freezing behavior. During the final stressor, which involved acoustic

stimuli from a startle box, no group differences were found in startle amplitude. However,

treatment- and sex-specific differences emerged during the "null periods" between stimuli:

OPFR females exhibited delayed movement following stimulus cessation compared to oil

females and OPFR males. These findings highlight a potential sex-specific vulnerability to

OPFR and AVS exposure, suggesting that OPFR-treated females may exhibit heightened

behavioral inhibition and altered stress response due to HPA dysregulation. It is possible that

perinatal OPFR treatment alters glucocorticoid or mineralocorticoid related feedback

mechanisms which may account for these differences (28,31).

Conclusion

This study demonstrates that perinatal exposure to organophosphate flame retardants (OPFRs)

leads to long-lasting, sex-specific disruptions in the neuroendocrine stress response. Female

OPFR-treated offspring exhibited heightened corticosterone release following restraint stress.

These changes were accompanied by significant modulation of gene expression in the

paraventricular nucleus (PVN), pituitary, and adrenal glands, particularly in genes involved in

CRH signaling, glucocorticoid feedback, and catecholamine synthesis. The lower basal CORT

levels in OPFR AVS females points to dysregulation of mechanisms within the HPA axis. These

effects were particularly pronounced in stress-sensitive brain regions such as the BNST and

PVN, further supporting a sex-dependent vulnerability. Given that OPFRs interfere with

hormone receptor function, these findings may implicate disrupted receptor-mediated signaling

during critical periods of brain development. Collectively, these results provide compelling

evidence that early-life OPFR treatment reprograms central stress circuits, with long-lasting

consequences on behavioral and endocrine responses to stress. These alterations may increase

susceptibility to mood and anxiety disorders, particularly in females, underscoring the need for

further investigation into sex-specific mechanisms and long-term neurobehavioral outcomes.

Author Contributions: Conceptualization, T.A.R, C.M.R; investigation, C.M.R, A.Y, C.B, J.D,

S.C; analysis, C.M.R, N.B; resources, T.A.R, N.B; writing-original draft preparation, C.M.R;

writing- review and editing, C.M.R, T.A.R, N.B; supervision, T.A.R; funding acquisition, T.A.R.

Funding: This research was supported by funds from National Institutes of Health

(R01MH123544; R21ES035848; P30ES005022). C.M.R. was previously supported by and C.B.

is currently supported by the National Institute of Environmental Health Sciences

(T32ES007148).

Disclosures: The authors have nothing to disclose.

Data Availability: The data that support the findings of this study are available in the Materials

and Methods and/or Results section of this article. Supplementary Materials are available at

(editor will insert link??)

References

1.   de Boer J, Harrad S, Sharkey M. The European Regulatory Strategy for flame retardants– The right direction but still a risk of getting lost. *Chemosphere*. 2024;347:140638.

2.   Blum A, Behl M, Birnbaum LS, et al. Organophosphate ester flame retardants: are they a regrettable substitution for polybrominated diphenyl ethers? *Environmental Science & Technology Letters*. 2019;6(11):638-649.

3.   van Melis LVJ, Zimnik KN, Persad AR, et al. Exposure to organophosphate flame retardants and phthalates alters neuronal activity and network development. *Toxicology*. Aug 2025;515:154168. doi:10.1016/j.tox.2025.154168

4.   Yan Z, Feng C, Jin X, et al. Organophosphate esters cause thyroid dysfunction via multiple signaling pathways in zebrafish brain. *Environ Sci Ecotechnol*. Oct 2022;12:100198. doi:10.1016/j.ese.2022.100198

5.   Percy Z, Vuong AM, Xu Y, et al. Maternal Urinary Organophosphate Esters and Alterations in Maternal and Neonatal Thyroid Hormones. *American Journal of Epidemiology*. 2021;190(9):1793-1802. doi:10.1093/aje/kwab086

6.   Li Y, Zhao M, Shen X, et al. Adverse Effects of Aryl Organophosphate Esters on Reproduction. *Environ Sci Technol*. Jun 3 2025;59(21):10122-10135. doi:10.1021/acs.est.5c01992

7.   Van den Eede N, Dirtu AC, Neels H, Covaci A. Analytical developments and preliminary assessment of human exposure to organophosphate flame retardants from indoor dust. *Environ Int*. Feb 2011;37(2):454-61. doi:10.1016/j.envint.2010.11.010

8.   Butt CM, Congleton J, Hoffman K, Fang M, Stapleton HM. Metabolites of organophosphate flame retardants and 2-ethylhexyl tetrabromobenzoate in urine from paired mothers and toddlers. *Environ Sci Technol*. Sep 2 2014;48(17):10432-8. doi:10.1021/es5025299

9.   Doherty BT, Hoffman K, Keil AP, et al. Prenatal exposure to organophosphate esters and behavioral development in young children in the Pregnancy, Infection, and Nutrition Study. *Neurotoxicology*. 2019;73:150-160.

10.   Dubovický M. Neurobehavioral manifestations of developmental impairment of the brain. *Interdiscip Toxicol*. Jun 2010;3(2):59-67. doi:10.2478/v10102-010-0012-4

11.   Rojas CM, Wiersielis KR, Roepke TA. Mechanisms of Flame Retardant Toxicity and Their Impacts on Anxiety and Cognition in the Brain. *Endocrinology*. Apr 15 2025;doi:10.1210/endocr/bqaf074

12.   Chen Y-X, Zheng J, Zhang X-F. Association analysis between organophosphorus flame retardants exposure and the risk of depression: Data from NHANES 2017–2018. *Journal of Affective Disorders*. 2024/06/15/ 2024;355:385-391. doi:https://doi.org/10.1016/j.jad.2024.04.004

13.   Lu M, Han F, Liu Y, et al. Pregnancy-related anxiety modifies the effects of maternal exposure to organophosphate esters on preschoolers' behavioral development. *Environmental Research*. 2025/04/15/ 2025;271:121081. doi:https://doi.org/10.1016/j.envres.2025.121081

14.   Panagiotakopoulos L, Neigh GN. Development of the HPA axis: where and when do sex differences manifest? *Front Neuroendocrinol*. Aug 2014;35(3):285-302. doi:10.1016/j.yfrne.2014.03.002

15. Shen W-J, Azhar S, Kraemer FB. ACTH Regulation of Adrenal SR-B1. Review.
Frontiers in Endocrinology. 2016-May-13 2016;Volume 7 - 2016doi:10.3389/fendo.2016.00042
16. Bhagavan NV, Ha C-E. Chapter 30 - Endocrine Metabolism III: Adrenal Glands. In:
Bhagavan NV, Ha C-E, eds. *Essentials of Medical Biochemistry (Second Edition)*. Academic
Press; 2015:559-575.
17. Osinga TE, van den Eijnden MHA, Kema IP, et al. Unilateral and Bilateral
Adrenalectomy for Pheochromocytoma Requires Adjustment of Urinary and Plasma
Metanephrine Reference Ranges. *The Journal of Clinical Endocrinology & Metabolism*.
2013;98(3):1076-1083. doi:10.1210/jc.2012-3418
18. Sawchenko PE, Arias C. Evidence for Short-Loop Feedback Effects of ACTH on CRF
and Vasopressin Expression in Parvocellular Neurosecretory Neurons. *Journal of*
Neuroendocrinology. 1995;7(9):721-731. doi:https://doi.org/10.1111/j.1365-
2826.1995.tb00814.x
19. Harris AP, Holmes MC, de Kloet ER, Chapman KE, Seckl JR. Mineralocorticoid and
glucocorticoid receptor balance in control of HPA axis and behaviour.
Psychoneuroendocrinology. May 2013;38(5):648-58. doi:10.1016/j.psyneuen.2012.08.007
20. Zhang Q, Wang J, Zhu J, Liu J, Zhao M. Potential Glucocorticoid and Mineralocorticoid
Effects of Nine Organophosphate Flame Retardants. *Environmental Science & Technology*.
2017/05/16 2017;51(10):5803-5810. doi:10.1021/acs.est.7b01237
21. Zhang Q, Yu C, Fu L, Gu S, Wang C. New Insights in the Endocrine Disrupting Effects
of Three Primary Metabolites of Organophosphate Flame Retardants. *Environmental Science &*
Technology. 2020/04/07 2020;54(7):4465-4474. doi:10.1021/acs.est.9b07874
22. Li Z, Robaire B, Hales BF. The Organophosphate Esters Used as Flame Retardants and
Plasticizers Affect H295R Adrenal Cell Phenotypes and Functions. *Endocrinology*. Aug 1
2023;164(9)doi:10.1210/endocr/bqad119
23. Li Z, Hales BF, Robaire B. Impact of exposure to a mixture of organophosphate esters on
the adrenal glands of Sprague Dawley rats. *Toxicological Sciences*. 2024;204(1):43-56.
doi:10.1093/toxsci/kfae154
24. Wade MG, Kawata A, Rigden M, Caldwell D, Holloway AC. Toxicity of Flame Retardant
Isopropylated Triphenyl Phosphate: Liver, Adrenal, and Metabolic Effects. *Int J Toxicol*. Jul/Aug
2019;38(4):279-290. doi:10.1177/1091581819851502
25. Akimoto T, Kobayashi S, Nakayama A, et al. Toxicological effects of Tris (1,3-dichloro-
2-propyl) phosphate exposure in adult male rats differ depending on the history of exposure in
the neonatal period. *J Appl Toxicol*. Sep 2022;42(9):1503-1509. doi:10.1002/jat.4316
26. Dungar BM, Schupbach CD, Jacobson JR, Kopf PG. Adrenal Corticosteroid Perturbation
by the Endocrine Disruptor BDE-47 in a Human Adrenocortical Cell Line and Male Rats.
Endocrinology. 2021;162(11)doi:10.1210/endocr/bqab160
27. Chrousos GP, Gold PW. The concepts of stress and stress system disorders: overview of
physical and behavioral homeostasis. *Jama*. 1992;267(9):1244-1252.
28. Handa RJ, Weiser MJ. Gonadal steroid hormones and the hypothalamo-pituitary-adrenal
axis. *Front Neuroendocrinol*. Apr 2014;35(2):197-220. doi:10.1016/j.yfrne.2013.11.001
29. Gong S, Miao Y-L, Jiao G-Z, et al. Dynamics and Correlation of Serum Cortisol and
Corticosterone under Different Physiological or Stressful Conditions in Mice. *PLOS ONE*.
2015;10(2):e0117503. doi:10.1371/journal.pone.0117503

30.   Hodes GE, Pfau ML, Purushothaman I, et al. Sex Differences in Nucleus Accumbens
Transcriptome Profiles Associated with Susceptibility versus Resilience to Subchronic Variable
Stress. *J Neurosci*. Dec 16 2015;35(50):16362-76. doi:10.1523/jneurosci.1392-15.2015
31.   Lebow MA, Chen A. Overshadowed by the amygdala: the bed nucleus of the stria
terminalis emerges as key to psychiatric disorders. *Molecular Psychiatry*. 2016/04/01
2016;21(4):450-463. doi:10.1038/mp.2016.1
32.   Degroat TJ, Paladino SE, Denney K, Moran KM, Samuels BA, Roepke TA. Chronic
social instability stress differentially affects the behavior and the transcriptome of the
anterodorsal bed nucleus of the stria terminalis between male and female mice. *bioRxiv*. Jan 21
2025;doi:10.1101/2025.01.21.634142
33.   Degroat TJ, Paladino SE, Samuels BA, Roepke TA. Chronic variable mild stress alters
the transcriptome and signaling properties of the anterodorsal bed nucleus of the stria terminalis
in a sex-dependent manner. *J Neuroendocrinol*. Aug 2025;37(8):e70041. doi:10.1111/jne.70041
34.   Degroat TJ, Wiersielis K, Denney K, et al. Chronic stress and its effects on behavior,
RNA expression of the bed nucleus of the stria terminalis, and the M-current of NPY neurons.
Psychoneuroendocrinology. 2024/03/01/ 2024;161:106920.
doi:https://doi.org/10.1016/j.psyneuen.2023.106920
35.   Adams S, Wiersielis K, Yasrebi A, et al. Sex- and age-dependent effects of maternal
organophosphate flame-retardant exposure on neonatal hypothalamic and hepatic gene
expression. *Reprod Toxicol*. Jun 2020;94:65-74. doi:10.1016/j.reprotox.2020.04.001
36.   Wiersielis KR, Adams S, Yasrebi A, Conde K, Roepke TA. Maternal exposure to
organophosphate flame retardants alters locomotor and anxiety-like behavior in male and female
adult offspring. *Horm Behav*. Jun 2020;122:104759. doi:10.1016/j.yhbeh.2020.104759
37.   Daldrup T, Remmes J, Lesting J, et al. Expression of freezing and fear-potentiated startle
during sustained fear in mice. *Genes, Brain and Behavior*. 2015;14(3):281-291.
doi:https://doi.org/10.1111/gbb.12211
38.   Koenig JI, Elmer GI, Shepard PD, et al. Prenatal exposure to a repeated variable stress
paradigm elicits behavioral and neuroendocrinological changes in the adult offspring: potential
relevance to schizophrenia. *Behav Brain Res*. Jan 30 2005;156(2):251-61.
doi:10.1016/j.bbr.2004.05.030
39.   Krumm EA, Patel VJ, Tillery TS, et al. Organophosphate flame-retardants alter adult
mouse homeostasis and gene expression in a sex-dependent manner potentially through
interactions with ERα. *Toxicological sciences*. 2018;162(1):212-224.
40.   Shi S, Feng Q, Zhang J, et al. Global patterns of human exposure to flame retardants
indoors. *Science of The Total Environment*. 2024/02/20/ 2024;912:169393.
doi:https://doi.org/10.1016/j.scitotenv.2023.169393
41.   Kim U-J, Wang Y, Li W, Kannan K. Occurrence of and human exposure to
organophosphate flame retardants/plasticizers in indoor air and dust from various
microenvironments in the United States. *Environment International*. 2019/04/01/ 2019;125:342-
349. doi:https://doi.org/10.1016/j.envint.2019.01.065
42.   Li Y, Wang X, Zhu Q, et al. Organophosphate Flame Retardants in Pregnant Women:
Sources, Occurrence, and Potential Risks to Pregnancy Outcomes. *Environmental Science &*
Technology. 2023/05/09 2023;57(18):7109-7128. doi:10.1021/acs.est.2c06503
43.   Ji Y, Yao Y, Duan Y, et al. Association between urinary organophosphate flame retardant
diesters and steroid hormones: A metabolomic study on type 2 diabetes mellitus cases and
controls. *Sci Total Environ*. Feb 20 2021;756:143836. doi:10.1016/j.scitotenv.2020.143836

44.   Hu P, Liu J, Yasrebi A, et al. Gq Protein-Coupled Membrane-Initiated Estrogen Signaling
Rapidly Excites Corticotropin-Releasing Hormone Neurons in the Hypothalamic Paraventricular
Nucleus in Female Mice. *Endocrinology*. Sep 2016;157(9):3604-20. doi:10.1210/en.2016-1191
45.   Rivier C. Gender, sex steroids, corticotropin-releasing factor, nitric oxide, and the HPA
response to stress. *Pharmacology biochemistry and behavior*. 1999;64(4):737-751.
46.   Viau V, Bingham B, Davis J, Lee P, Wong M. Gender and puberty interact on the stress-
induced activation of parvocellular neurosecretory neurons and corticotropin-releasing hormone
messenger ribonucleic acid expression in the rat. *Endocrinology*. 2005;146(1):137-146.
47.   Kokras N, Hodes GE, Bangasser DA, Dalla C. Sex differences in the hypothalamic–
pituitary–adrenal axis: An obstacle to antidepressant drug development? *British Journal of
Pharmacology*. 2019;176(21):4090-4106. doi:https://doi.org/10.1111/bph.14710
48.   Jeong KH, Jacobson L, Pacák K, Widmaier EP, Goldstein DS, Majzoub JA. Impaired
basal and restraint-induced epinephrine secretion in corticotropin-releasing hormone-deficient
mice. *Endocrinology*. Mar 2000;141(3):1142-50. doi:10.1210/endo.141.3.7370
49.   Coste SC, Kesterson RA, Heldwein KA, et al. Abnormal adaptations to stress and
impaired cardiovascular function in mice lacking corticotropin-releasing hormone receptor-2.
Nat Genet. Apr 2000;24(4):403-9. doi:10.1038/74255
50.   Ebner K, Fontebasso V, Ferro F, Singewald N, Hannibal J. PACAP regulates
neuroendocrine and behavioral stress responses via CRF-containing neurons of the rat
hypothalamic paraventricular nucleus. *Neuropsychopharmacology*. Feb 2025;50(3):519-530.
doi:10.1038/s41386-024-02016-9
51.   Herman JP, McKlveen JM, Ghosal S, et al. Regulation of the Hypothalamic-Pituitary-
Adrenocortical Stress Response. *Comprehensive Physiology*. 2016:603-621.
52.   Fan J-M, Wang X, Hao K, Yuan Y, Chen X-Q, Du J-Z. Upregulation of PVN CRHR1 by
gestational intermittent hypoxia selectively triggers a male-specific anxiogenic effect in rat
offspring. *Hormones and Behavior*. 2013/01/01/ 2013;63(1):25-31.
doi:https://doi.org/10.1016/j.yhbeh.2012.11.005
53.   Kolber BJ, Wieczorek L, Muglia LJ. Hypothalamic-pituitary-adrenal axis dysregulation
and behavioral analysis of mouse mutants with altered glucocorticoid or mineralocorticoid
receptor function. *Stress*. Sep 2008;11(5):321-38. doi:10.1080/10253890701821081
54.   Zhang J, Yang Y, Liu W, Schlenk D, Liu J. Glucocorticoid and mineralocorticoid
receptors and corticosteroid homeostasis are potential targets for endocrine-disrupting chemicals.
Environ Int. Dec 2019;133(Pt A):105133. doi:10.1016/j.envint.2019.105133
55.   Liu QS, Sun Z, Ren X, et al. Chemical Structure-Related Adipogenic Effects of
Tetrabromobisphenol A and Its Analogues on 3T3-L1 Preadipocytes. *Environmental Science &
Technology*. 2020/05/19 2020;54(10):6262-6271. doi:10.1021/acs.est.0c00624
56.   Graceli JB, Dettogni RS, Merlo E, et al. The impact of endocrine-disrupting chemical
exposure in the mammalian hypothalamic-pituitary axis. *Molecular and Cellular Endocrinology*.
2020/12/01/ 2020;518:110997. doi:https://doi.org/10.1016/j.mce.2020.110997
57.   Liu C, Wang Q, Liang K, et al. Effects of tris(1,3-dichloro-2-propyl) phosphate and
triphenyl phosphate on receptor-associated mRNA expression in zebrafish embryos/larvae.
Aquatic Toxicology. 2013/03/15/ 2013;128-129:147-157.
doi:https://doi.org/10.1016/j.aquatox.2012.12.010
58.   Repouskou A, Papadopoulou A-K, Panagiotidou E, et al. Long term transcriptional and
behavioral effects in mice developmentally exposed to a mixture of endocrine disruptors

associated with delayed human neurodevelopment. *Scientific Reports*. 2020/06/09 2020;10(1):9367. doi:10.1038/s41598-020-66379-x

59.   Sass B. Embryology, Adrenal Gland, Mouse. In: Jones TC, Capen CC, Mohr U, eds. *Endocrine System*. Springer Berlin Heidelberg; 1996:381-386.

60.   Ubuka T. Subchapter 131B - Noradrenaline/adrenaline. In: Ando H, Ukena K, Nagata S, eds. *Handbook of Hormones (Second Edition)*. Academic Press; 2021:1041-1044.

61.   Kaimal A, Hooversmith JM, Cherry AD, et al. Prenatal exposure to bisphenol A and/or diethylhexyl phthalate alters stress responses in rat offspring in a sex- and dose-dependent manner. Original Research. *Frontiers in Toxicology*. 2023-December-13 2023;Volume 5 - 2023doi:10.3389/ftox.2023.1264238

62.   Stroth N, Eiden LE. Stress hormone synthesis in mouse hypothalamus and adrenal gland triggered by restraint is dependent on pituitary adenylate cyclase-activating polypeptide signaling. *Neuroscience*. Feb 17 2010;165(4):1025-30. doi:10.1016/j.neuroscience.2009.11.023

63.   Lehmann ML, Mustafa T, Eiden AM, Herkenham M, Eiden LE. PACAP-deficient mice show attenuated corticosterone secretion and fail to develop depressive behavior during chronic social defeat stress. *Psychoneuroendocrinology*. May 2013;38(5):702-15. doi:10.1016/j.psyneuen.2012.09.006

64.   Hammack SE, Braas KM, May V. Chemoarchitecture of the bed nucleus of the stria terminalis: Neurophenotypic diversity and function. *Handb Clin Neurol*. 2021;179:385-402. doi:10.1016/b978-0-12-819975-6.00025-x

65.   Wang X, Weng G, Gao Y, Wang Y, Zhang C. Quercetin ameliorates chronic restraint stress- and LPS-induced anxiety-like behaviors by modulating neuroinflammation in the lateral hypothalamus. *Psychopharmacology*. 2025/04/10 2025;doi:10.1007/s00213-025-06784-0

66.   Brancato A, Bregman D, Ahn HF, et al. Sub-chronic variable stress induces sex-specific effects on glutamatergic synapses in the nucleus accumbens. *Neuroscience*. May 14 2017;350:180-189. doi:10.1016/j.neuroscience.2017.03.014

67.   Liu X, Zhao X, Wang Y, et al. Triphenyl phosphate permeates the blood brain barrier and induces neurotoxicity in mouse brain. *Chemosphere*. 2020/08/01/ 2020;252:126470. doi:https://doi.org/10.1016/j.chemosphere.2020.126470

68.   Wang X, Song F. The neurotoxicity of organophosphorus flame retardant tris (1, 3-dichloro-2-propyl) phosphate (TDCPP): main effects and its underlying mechanisms. *Environmental Pollution*. 2024;346:123569.

69.   Yun K, Jeon H, Kho Y, Ji K. Potential adverse outcome pathway of neurodevelopmental toxicity, inflammatory response, and oxidative stress induction mediated by three alkyl organophosphate flame retardants in zebrafish larvae. *Chemosphere*. 2024/05/01/ 2024;356:141901. doi:https://doi.org/10.1016/j.chemosphere.2024.141901

70.   Zhong X, Wu J, Ke W, et al. Neonatal exposure to organophosphorus flame retardant TDCPP elicits neurotoxicity in mouse hippocampus via microglia-mediated inflammation in vivo and in vitro. *Archives of Toxicology*. 2020/02/01 2020;94(2):541-552. doi:10.1007/s00204-019-02635-y

71.   Hong J, Lu X, Wang J, et al. Triphenyl phosphate disturbs placental tryptophan metabolism and induces neurobehavior abnormal in male offspring. *Ecotoxicology and Environmental Safety*. 2022/09/15/ 2022;243:113978. doi:https://doi.org/10.1016/j.ecoenv.2022.113978

72.   Maita I, Bazer A, Chae K, et al. Chemogenetic activation of corticotropin-releasing factor-expressing neurons in the anterior bed nucleus of the stria terminalis reduces effortful

907 motivation behaviors. *Neuropsychopharmacology*. Jan 2024;49(2):377-385. doi:10.1038/s41386-
908 023-01646-9
909

910 **Figures**

912 **Figure 1.** Serum concentrations of corticosterone and epinephrine in (A/B) female and (C/D)

913 male mice. Data are represented as mean ± SEM and dots represent the sample size (number of

914 litters) per treatment and exposure (**= P<0.01, ****= P<0.0001).

Figure 2. Paraventricular nucleus of the hypothalamus (PVN) mRNA expression in female/male

adult offspring. (A/G) *Crh*; (B/H) *Crhr1*; (C/I) *Crhr2*; (D/J) *Pacap*; (E/K) *Pac1r*; and (F/L)

Ptpn5. Data are represented as mean ± SEM and dots represent the sample size (number of

litters) per treatment and exposure. (*= P<0.05, **= P<0.01 , ***= P<0.001, ****= P<0.0001).

Figure 3. Pituitary mRNA expression in female/male adult offspring: (A/D) *Crhr1*; (B/E) *Pomc*;

and (C/F) *Nr3c1*. Data are represented as mean ± SEM and dots represent the sample size

(number of litters) per treatment and exposure. (*= P<0.05, **= P<0.01, ***= P<0.001, ****=

P<0.0001).

Figure 4. Adrenal mRNA expression of genes in the catecholamine pathway. Female/male adult

offspring: (A/C) *Dbh* and (B/D) *Pnmt.* Data are represented as mean ± SEM and dots represent

the sample size (number of litters) per treatment and exposure. (*= P<0.05, **= P<0.01, ***=

P<0.001).

Figure 5. Corticosterone concentrations in serum of (A) female and (B) male mice. Data are

represented as mean ± SEM and dots represent the sample size (number of litters) per treatment

and exposure (*= P<0.05).

Figure 6. Paraventricular nucleus of the hypothalamus (PVN) mRNA expression in female/male

adult offspring. (A/D) *Crhr1*; (B/E) *Pacap*; and (C/F) *Pac1r*. Data are represented as mean ±

SEM and dots represent the sample size (number of litters) per treatment and exposure. (*=

P<0.05, **= P<0.01, ***= P<0.001, ****= P<0.0001).

Figure 7. Bed nucleus stria terminalis (BNST) mRNA expression. Female adult offspring: (A)

Crhr1; (B) *Pacap*; and (C) *Pac1r*. Male adult offspring: (A) *Crhr2*; (B) *Pacap*; and (C) *Pac1r*.

Data are represented as mean ± SEM and dots represent the sample size (number of litters) per

treatment and exposure. (*= P<0.05, **= P<0.01, ***= P<0.001, ****= P<0.0001).

Figure 8. Open Field Test (OFT)- time immobile in the perimeter and corners for female (A and B) and male (C and D) adult offspring. Data are represented as mean ± SEM and dots represent the sample size (number of litters) per treatment and exposure. (*= P<0.05). Heat maps for each treatment and exposure group were generated by ANY-maze based on the animal's center point for the entire duration of the test, with the hottest point (in red) indicating maximum amount of time spent in an area.

1002

1003

1004

1005

1008

Figure 9. Elevated Plus Maze (EPM) results for female/male adult offspring: (A/E) Distance;

(B/F) Mean speed; (C/G) Entries into closed arms; and (D/H) distance traveled in closed arms.

Data are represented as mean ± SEM and dots represent the sample size (number of litters) per

treatment and exposure. (*= P<0.05, **= P<0.01).

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

Figure 10. Light/dark box (LDB) emergence test- light and dark zone activity. Female/male

adult offspring: (A/E) Entries into light zone; (B/F) Exits from light zone; (C/G) Entries into

dark zone; (D/H) Exits from dark zone. Data are represented as mean ± SEM and dots represent

the sample size (number of litters) per treatment and exposure. (*= P<0.05, **= P<0.01).

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1044

Figure 11. Novelty-suppressed feeding (NSF) for female/male adult offspring: (A/E) Latency to

eat in the novel arena; (B/F) Kaplan-Meier survival curve for the probability to eat in the novel

arena; (C/G) Latency to eat in the home cage; (D/H) Kaplan-Meier survival curve for the

probability to eat in the home cage. Data are represented as mean ± SEM and dots represent the

sample size (number of litters) per treatment and exposure.

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062   **Supplemental Tables/Figures**

1063

1064   **Table S1.** List of primers for qPCR.

Gene	Accession #	Forward Primer	Reverse Primer
Actb	NM_007393.3	GCCCTGAGGCTCTTTTCCA	TAGTTTCATGGATGCCACAGGA
Crh	NM_205769.2	AGGAGGCATCCTGAGAGAAGT	CATGTTAGGGGCGCTCTC
Crhr1	NM_001430903.1	CGCAAGTGGATGTTCGTCT	GGGGCCCTGGTAGATGTAGT
Crhr2	NM_001288620.1	AAGCTGGTGATTTGGTGGAC	GGTGGATGCTCGTAACTTCG
Gapdh	NM_008084.2	TGACGTGCCGCCTGGAGAAA	AGTGTAGCCCAAGATGCCCTTCAG
Hprt	NM_013556	GCTTGCTGGTGAAAAGGACCTCTCGAAG	CCCTGAAGTACTCATTATAGTCAAGGGCAT
Ikk	NM_001159774.1	CCATATCCTGGCTGTCACCT	GGCACCTTGGATGACCTAGA
Pacap	NM_001409528.1	CAGAAGGCCAGTCACCTCTGTC	GAGAGTAGGCGAGCAGCCAAAG
Pac1r	NM_133511	AACGACCTGATGGGACTAAAC	CGGAAGCGGCACAAGATGACC
Pomc	NM_008895	GGAAGATGCCGAGATTCTGC	TCCGTTGCCAGGAAACAC
Ptpn5	NM_013643.2	GCACTGTGGCCGACTTTTG	CTGCACGGTGATCTCCACG

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075   **Table S2.** List of primers for qPCR ordered from Bio-Rad.

Gene	Assay ID
Cyp11a1	qMmuCID0005308
Cyp11b1	qMmuCID0019988
Cyp11b2	qMmuCED0045612
Cyp21a1	qMmuCED0001670
Dbh	qMmuCID0018644
Mc2r	qMmuCID0007182
Nr3c1	qMmuCID0016200
Pnmt	qMmuCED0048695

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090 **Figure S1.** Adrenal mRNA expression- steroidogenic pathway. Female/male adult offspring:

1091 (A/F) *Mc2r*; (B/G) *Cyp11a1*; (C/H) *Cyp21a1;* (D/I) *Cyp11b1*; and (E/J) *Cyp11b2*. Data are

1092 represented as mean ± SEM and dots represent the sample size (number of litters) per treatment

1093 and exposure. (*= P<0.05, **= P<0.01).

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107 **Figure S2.** PVN mRNA expression. Female/male adult offspring: (A/D) *Crh*; (B/E) *Crhr2*; and

1108 (C/F) *Ikk.* Data are represented as mean ± SEM and dots represent the sample size (number of

1109 litters) per treatment and exposure. (*= P<0.05).

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121 **Figure S3.** BNST mRNA expression. Female adult offspring: (A) *Crh*; (B) *Crhr2*; and (C) *Ikk.*

1122 Male adult offspring: (D) *Crh*; (E) *Crhr1*; and (F) *Ikk* Data are represented as mean ± SEM and

1123 dots represent the sample size (number of litters) per treatment and exposure. (**= P<0.01,

1124 ****= P<0.0001).

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135 **Figure S4.** Behavior tests- locomotion. OFT- female/male adult offspring: (A/C) distance and

1136 (B/D) mean speed. LDB- female/male adult offspring: (E/G) distance and (F/H) mean speed.

1137 Data are represented as mean ± SEM and dots represent the sample size (number of litters) per

1138 treatment and exposure. (**= P<0.01, ****= P<0.0001).

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152 **Figure S5.** NSF- female/male adult offspring: (A/C) weight loss percentage and (B/D) amount of

1153 pellet eaten. Data are represented as mean ± SEM and dots represent the sample size (number of

1154 litters) per treatment and exposure. (*= P<0.05, ***= P<0.001).

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164 **Figure S6.** Startle box activity in null and S2 phase. (A) Null latency; (B) Null duration; (C) S2-

1165 null duration. Data are represented as mean ± SEM and dots represent the sample size (number

1166 of litters) per treatment and exposure. (*= P<0.05, ***= P<0.001).

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181