Longitudinal intergenerational hyperscanning indexes changes in social connection

Longitudinal intergenerational hyperscanning and social connection

Ryssa Moffat[1], Guillaume Dumas[2,3], Emily S. Cross[1]

[1] Social Brain Sciences Lab, ETH Zurich, Switzerland
[2] CHU Sainte-Justine Azrieli Research Center, Department of Psychiatry, University of Montreal, Quebec, Canada
[3] Mila – Quebec Artificial Intelligence Institute, Quebec, Canada

* Correspondence:
Ryssa Moffat
rmoffat@ethz.ch

Emily Cross
ecross@ethz.ch

Abstract

Loneliness is globally acknowledged as a severe and burgeoning health risk, fuelling interest in helping people of all ages form meaningful social connections. One promising approach consists of intergenerational social programs. While behavioural and qualitative evidence derived from such programs promise health and wellbeing benefits, the physiological consequences of repeated intergenerational encounters remain unknown. Insight into physiological changes will shed light on the mechanisms of social connection and can inform program design choices. We charted changes in interpersonal neural synchrony (INS) in 31 intergenerational (older/younger adult) and 30 same generation (younger adult) dyads across a six-session creative drawing program. At each session, dyads completed self-report measures, drew together and alone, and had their cortical activation recorded with fNIRS. In both groups, INS was greater while dyads drew together than alone. Across sessions, intergenerational dyads' INS decreased and same generation dyads' INS increased. INS in RIFG~RTPJ and RIFG~RIFG were predictive of loneliness levels and feelings of social closeness, respectively. The research reinforces the multi-faceted nature of INS dynamics as social connections are forged.

1 Introduction

Experiences of loneliness can have serious negative consequences for physical and mental health. As a result, widespread efforts are being made to combat loneliness (1). According to meta-analyses of behavioural studies, community programs providing opportunities for repeated contact between generations can improve older adults' (65+ years) physical, social, and cognitive health (2), while also reducing younger adults' stereotypes about aging (3,4). Moreover, physicians are increasingly prescribing attendance at such programs–a practice called social prescribing (5)–further highlighting the value the medical field attributes to such programs. To date the existing evidence base for intergenerational community programs consists of behavioural and qualitative studies. While offering promising results in terms of self-reported benefits and social engagement, such studies offer little insight into how or why intergenerational relationships develop on a physiological level. In this study, we sought to provide a physiological perspective on the trajectory of intergenerational relationship development using a longitudinal hyperscanning approach, comparing intergenerational and same generation dyads who completed a six-week creative drawing program. Specifically, we tracked the trajectories of dyads' interpersonal neural synchrony (INS) and changes in social connectedness.

The following sections briefly describe the benefits of intergenerational social interactions, what is known about INS in intergenerational interactions, and the benefits of longitudinal perspectives on INS. Following this, we detail the aims and hypotheses of the present study.

1.1 Intergenerational social interactions combat loneliness by offering social connections

Loneliness can be defined as "a negative, subjective emotional state resulting from a discrepancy between one's desired and actual experiences of connection" (6). Put more succinctly, loneliness can be thought of as *perceived* social isolation. Policy makers and researchers have long emphasised the severe risk that loneliness poses to physical and mental health (1,7,8). Public figures and media outlets go so far as to compare the adverse health impacts (e.g., mortality) of loneliness to those of smoking cigarettes (9). Considering everyday functioning, longitudinal studies show that loneliness and social isolation can precipitate declines in cognitive functioning including verbal ability, numeracy, and short-term memory (10,11). Loneliness is also associated with reduced perceptual speed (12) and biases toward negative evaluations of social information (13–15). At the brain level,

chronic loneliness and episodes of social isolation have been associated with altered structure
(typically reduction in tissue volume) and altered function of prefrontal and posterior superior
temporal cortices, as well as the insula, hippocampus, and amygdala (16–18). Attentional, visual, and
default mode networks are also reported to show stronger functional connectivity in lonely
individuals, which is believed to be associated with the bias toward processing social information
negatively and a tendency toward increased rumination (18–20).

To combat loneliness, policy makers, researchers (1,21,22), and health practitioners (5) promote the
creation of opportunities for meaningful social interactions for all members of society, and especially
for those from populations most susceptible to loneliness. This typically involves initiating
community programs that provide a platform for regular socialising opportunities (6,23–25). Some
community programs are intentionally designed to bring together members of different age groups,
i.e., to be intergenerational. One reason for this is that people of all ages experience loneliness (26–
28). Another is that intergenerational social interactions can improve social and physical wellbeing
for older adults, as well as increase empathy and feelings of social closeness toward older adults from
younger participants (2). The observed improvements seem to stem from repeated intergenerational
contact (29–31). In this context, repeated intergenerational contact requires regular engagement in
intergenerational community programs, which is most likely to be achieved by ensuring that
programs offer meaningful social interactions. One approach that has been shown to be particularly
successful in this domain is arts-based and creative activities (32). Indeed, a growing body of
evidence documents the social and physical health benefits of intergenerational arts programs for
participants of all ages (30,33–35).

Despite the mounting behavioural and qualitative evidence that repeated intergenerational
interactions yield social benefits, including reduced loneliness, little is known about the physiological
aspects that contribute to *how* and *why* these intergenerational programs offer such benefits. This is
important because the neurophysiological signatures of budding intergenerational relationships have
the potential to be harnessed as tools to promote the formation of new social connections, yet they
remain unknown (36).

1.2   Interpersonal neural synchrony (INS) during collaborative intergenerational interactions

One source of physiological information is brain activity. Recording brain activity from two or more
people while they interact is commonly referred to as 'hyperscanning' and offers a measure of
interpersonal neural synchrony (INS; the extent to which patterns of brain activity are temporally
aligned). Mobile functional near-infrared spectroscopy (fNIRS) and electroencephalography (EEG)
are particularly well suited to recording brain activity during social interactions in real-world settings
(36,37). Hyperscanning with non-mobile techniques, such as functional magnetic resonance imaging
(fMRI) and magneto-encephalography (MEG), is possible, though it offers less realistic social
interactions and is less logistically feasible (38,39).

1.2.1 INS in social brain networks differs between solo and dyadic tasks

In studies comparing INS while dyads perform tasks collaboratively and alone, INS is consistently
greater for collaborative performance than solo performance. This pattern has been observed for
finger tapping (40), cognitive tasks such as n-back and object detection tasks (41,42), singing (43),
tracing shapes on a screen using computer buttons or a stylus (44,45), drawing specific verbs as in
Pictionary (46), among other activities. A meta-analysis of fNIRS hyperscanning studies by
Czeszumski et al. (47) reports that during collaboration, INS is commonly observed between both
interactants' prefrontal cortex (PFC), inferior frontal cortex (IFG) or temporoparietal junction (TPJ).

INS in the prefrontal and inferior frontal cortices is proposed to reflect initiation and reception of
joint attention (48–50), a social competence that depends on cognitive control (51,52). INS in the
TPJ, a key node of the Theory of Mind (ToM) network (53,54), is believed to reflect perspective
taking and integration of social information (47). Complementary findings from EEG hyperscanning
studies suggest that the strength and nature of collaboration shape the levels of INS recorded over
frontal and prefrontal brain regions for alpha, beta, and theta frequency bands (55–59). Specifically,
the alpha band may index joint attention and visual processing, while beta and theta bands may
reflect joint encoding of sensorimotor information (37,55,56). In combination, the findings from
fNIRS and EEG hyperscanning studies indicate that INS levels during collaboration are shaped by
processes such as joint attention and perspective taking.

Many hyperscanning studies on collaboration can be considered intergenerational, in that they
examine collaboration between parent–child/infant or student–teacher dyads (for reviews, see:
,60,61). Though these studies offer the field of social neuroscience a valuable starting point for
understanding intergenerational collaboration, two gaps in the literature have yet to be filled: First,
the hyperscanning literature on collaboration distinctly lacks any representation of older adults.
Second, the hyperscanning literature more generally offers very few longitudinal perspectives into
INS exceeding two time points. We now briefly explore the importance of addressing these gaps.

1.2.2 Older adults are underrepresented in hyperscanning research

To date, older adults' representation in the hyperscanning literature is limited to one hyperscanning
study (62), one study simulating INS (63), and one position piece arguing for the inclusion of older
adults in such studies (36). It is critical to address this knowledge gap as the real-world settings in
which older adults interact with younger adults and children (with their children and grandchildren,
in assisted living, when seeking medical care, in community programs or through volunteer work) are
numerous. Moreover, the proportion of older adults in the world's population is projected to continue
growing until the end of this century (64), meaning that filling this knowledge gap will benefit a
steadily increasing number of people.

In the one study to date exploring intergenerational INS involving older adults, Dikker et al. (63)
employed a neurocognitive model to simulate EEG-based INS during intergenerational social
interactions between grandparents, parents, and grandchildren. In this seminal study, the authors
found greater simulated INS between grandparent-grandchild dyads and grandparent-parent-child
triads, relative to grandparent-parent and parent-child dyads. The authors report that the increase in
simulated grandparent-child INS may be driven by age-related changes in the spectrum of brain
activity. The authors also encourage the collection of real-world data to validate these simulations
(63). These findings underscore how including people of all ages, particularly underrepresented older
adults, in intergenerational hyperscanning will enhance our understanding of INS and a wider array
of types of social connections.

1.2.3 INS trajectories across days, weeks, and years remain uncharted

Most hyperscanning studies offer snapshots of INS in a single moment in time, and compare INS
between contexts, tasks, or mental states. The *trajectory* in INS over days or weeks remains
unknown, despite the value of granularity insights for understanding changes in behaviour related to
the investigated contexts, task, and mental states. Longitudinal hyperscanning studies typically
include two time points, as opposed to a denser sampling of the trajectory of INS. Numerous reviews
from different domains of social neuroscience (e.g., developmental, clinical, communication etc.)
recognise the lack of longitudinal insights into INS and call for longitudinal hyperscanning studies to

be conducted (e.g., 65–71). The discrepancy between the number of calls for studies tracking INS
trajectories and the paucity of studies reporting them substantiates the critical need for additional
multi-session hyperscanning studies. Concretely, multi-session longitudinal designs can fill in the
blanks in our understanding of the dynamics of forming and maintaining social connections.

To the best of our knowledge, five hyperscanning studies have recorded INS at multiple sessions and
subsequently analysed the aggregated time points, as opposed to reporting changes in INS across
sessions (55,72–75). These studies are reviewed in greater detail in a position piece by Moffat et al.
(36). Only one fNIRS hyperscanning study offers insights into the trajectory of INS during
relationship formation (76). Sened et al.'s (76) proof-of-concept study tracked the trajectory of INS
over bilateral IFG in 8 clinician-patient dyads at the first, third and fifth sessions of a 6-session
intervention, where dyads completed imagery and cognitive behavioural training procedures. Sened
et al. report increasing INS across sessions, as well as an association between increasing INS and
reduced anxiety symptoms, improved wellbeing and perceived quality of session, but not the
therapeutic alliance (i.e., the patient's rating of quality of the collaborative relationship between
clinician and patient). One further study takes a unique approach to understanding INS in the context
of relationship formation, though only encompassing two time points (77). INS was recorded from 11
dyads at two sessions, the first within 5 days of the dyad becoming flatmates, and the second after 7
months of co-living. Wang et al. report that social closeness can be predicted from INS with
approximately 69% accuracy, using a deep neural network model and EEG hyperscanning recordings
from frontotemporal brain regions. These two studies (76,77) provide preliminary evidence that INS
between dyads increases as dyads grow more familiar with each other, however a larger sample size
and number of time points would provide a richer picture of INS trajectories in developing
relationships.

1.2.4 Cross-sectional research suggests that social closeness modulates INS

Taking a cross-sectional approach to understand social closeness and relationship development, some
studies compare romantic couples, friends and strangers at single moments in time. Such studies
suggest that the nature of a dyad's relationship influences INS during forms of interaction including
joint motor action, conversation, and classroom learning (45,62,72,78–80). Djalovski et al. (78)
recruited romantic couples, friends and strangers to complete a finger tapping task as dyads, and a
conversation-based empathy-giving task while brain activity was measured using EEG. During the
finger tapping task, INS was greatest between romantic couples, followed by friends, and lowest
between strangers. The authors suggest that cumulative joint experience with a specific partner
enhances prediction of the partners movements, resulting in greater INS. The empathy-giving task
revealed the opposite gradient with romantic couples showing the least INS, friends showing more,
and strangers showing the most INS. Djalovski et al. propose that strangers monitor each other for
potential negative responses more closely than couples to avoid the consequences of a potential
misunderstanding. In another study examining verbal communication, Speer et al. (80) recruited
friends and strangers to engage in prompted conversations with demarcated turns. The authors report
that across the span of conversation, friends showed decreasing INS and strangers showed increasing
INS, with the reductions in INS corresponding to more exploratory conversations and increases in
INS seeking to converge on common ground. Though phrased differently, Speer et al.'s interpretation
is closely aligned to Djalovski et al.'s interpretation: The less well a dyad knows each other, the
greater the INS during verbal communication. To explore this further, we highlight that a central
difference between the motor and verbal communication tasks is that the motor task (i.e., drawing
specific objects with an 'Etch A Sketch') is highly specified, setting both dyad members'
expectations clearly and symmetrically, whereas the verbal communication tasks (i.e., describing a

troubling or distressing event and conversing with demarcated turns) are temporally asymmetrical
and inherently unpredictable. We propose that the degree of predictability of behaviour induced by
the task contributes to the opposite INS gradients.

Considered together, these studies suggest that when dyads made up of strangers meet for the first
time and engage in a creative, unconstrained activity, such as drawing, INS is likely to be low.
Across repeated encounters, INS levels are likely to increase. Drawing on Dikker's model simulating
intergenerational INS, one would expect that generational constellations may impact INS levels and
may interact with the trajectory of INS as dyads become socially closer.

1.3   Present study: Tracking intergenerational INS longitudinally, across repeated
collaborative interactions

In this preregistered study (https://osf.io/hz6tm), we had three aims:

1. To characterise INS in intergenerational dyads made up of one young and one older adult.
2. To chart the trajectory of INS within same generation and intergenerational dyads, as
relationships form across repeated social interactions.
3. To characterise the relationship between INS and differing levels of loneliness and social
closeness, and varying attitudes toward other generations.

To achieve these aims, we recruited 31 intergenerational and 30 same generation dyads to complete a
six-session creative drawing program. At each session, we collected measures of participants'
loneliness and attitudes toward other generations, then recorded dyads' brain activity using fNIRS
while they drew independently and collaboratively with oil pastels. Participants then reported how
socially close they felt to their drawing partner. We calculated INS using wavelet transform
coherence (WTC) for all pairs of four regions of interest (ROIs), which were the bilateral IFG and
TPJ. According to fNIRS best practices, we report both fNIRS signals (oxygenated haemoglobin
[HbO] and deoxygenated haemoglobin [HbR]), with HbO results presented in the main manuscript
and HbR results presented in the Supplementary Materials. We employed a Bayesian statistical
analysis approach, according to which we make inferences based on 95% confidence intervals called
95% HPDs (highest posterior density region) rather than p-values. Further methodological details are
presented in Methods. Detailed results tables with estimates and 95% HPD are presented in
Supplementary Materials, separately from results text for ease of reading.

Our findings pertaining to INS in intergenerational and same generation dyads, tracked across 6
sessions in a real-world context resembling a community art program, deliver first insights into INS
involving older adults, multi-session trajectories of INS, as well as the relationship between INS and
measures of social connection.

2   Results

2.1   Self-reported loneliness, social closeness, and attitudes toward generations

We first investigated how self-reported loneliness, social closeness within dyads, as well as attitudes
towards generations differ between intergenerational and same generation dyads (estimates with 95%
HPD in Table 1; visualisations in Supplementary Figures 0 and 1).

Loneliness. Intergenerational dyads reported feeling substantially less lonely than same generation
dyads. Feelings of loneliness decreased substantially for both groups across sessions, with no

difference in rate. **Social closeness (OIS).** Intergenerational dyads reported substantially greater
feelings of social closeness, compared to same generation dyads. Feelings of social closeness
increased substantially across sessions for both groups, with no difference in rate. **Attitudes toward**
generations (allophilia). Same generation dyads reported feeling substantially closer to their own
generation than the intergenerational dyads, who reported feeling equally close to their own and the
other generation. Allophilia scores stayed stable across sessions (i.e., no evidence of change
observed).

Table 1. Unstandardised estimates computed per self-reported measure with 95% HPD in square brackets. Point
estimates for all six sessions combined are presented first, followed by the estimates of the slope of change across
sessions in the lower section. Self-report measures are loneliness, social closeness (OIS), and attitudes toward generations
(allophilia). Visualisations of distributions for each measure and trajectory across sessions by age (i.e., older or younger
adult) and group in Supplementary Figures 0 and 1.

	Loneliness	Social Closeness	Allophilia
Sessions combined			
Intergenerational	1.28 [1.02, 1.54]	4.08 [3.76, 4.39]	-1.19 [-3.45, 1.05]
Same gen	1.89 [1.62, 2.16]	3.52 [3.19, 3.38]	
Intergen-Same gen	-0.61 [-0.99, -0.25]	0.56 [0.10, 1.00]	4.48 [1.29, 7.56]
Change across six sessions			
Intergenerational	-0.06 [-0.10, -0.02]	0.24 [0.19, 0.27]	-0.21 [-0.55, 0.12]
Same gen	-0.07 [-.11, - 0.03]	0.25 [0.20, 0.30]	-0.02 [-0.37, 0.30]
Intergen-Same gen	0.01 [-0.05, 0.07]	-0.01 [-0.08, 0.07]	-0.18 [-0.66, 0.28]

Relationships between measures of social connection. As presented in Table 2, greater social
closeness was substantially associated with reduced feelings of loneliness for same generation dyads
and a matching trend was observed for intergenerational dyads. These associations did not differ
between groups. Greater social closeness was substantially associated with more positive attitudes
toward other generations for intergeneration dyads but not same generation dyads. This association
showed a trending difference between groups. Greater loneliness was substantially associated with
more positive attitudes toward other generations for same generation dyads but not intergenerational
dyads. This association did not differ between groups.

Table 2. Unstandardised estimates of relationship between self-reported measures per group with 95% HPD in square brackets. Self-report measures are loneliness, social closeness (OIS), and attitudes toward generations (allophilia).

	Loneliness	Social Closeness
Social Closeness (Intergen)	-0.13 [-0.27, 0.01]	
Social Closeness (Samegen)	-0.16 [-0.28, -0.04]	
Allophilia (Intergen)	0.01 [-0.00, 0.02]	0.02 [0.01, 0.04]
Allophilia (Samegen)	0.02 [0.01, 0.03]	0.01 [-0.01, 0.2]

2.2 INS

2.2.1 Real dyads vs. pseudo dyads

The first step in our analysis of INS was to perform contrasts between real and pseudo dyads for INS estimates reflecting all sessions combined and those reflecting change in INS across sessions per combination of ROI pair, task, and group. Contrasts serve to assess whether the level of observed INS (or the slope of change in INS) differs meaningfully from a null distribution of INS levels (or slopes) made of dyads who never interacted with one another. Further statistical analyses can then be performed on the ROI pairs where real dyads INS differed substantially from pseudo dyads' INS. We report estimates and contrasts with 95% HPD in Supplementary Tables 1-4.

INS levels for all sessions combined. We found that all substantial differences corresponded to greater INS for real than pseudo dyads (black squares in Figure 1A; solid lines in Figure 1B; Supplementary Table 2). The ROI pairs showing substantial differences were RIFG~RIFG for intergenerational dyads drawing alone, LIFG~LIFG, LTPJ~LTPJ, RIFG~RIFG, LIFG~RIFG, LIFG~LTPJ, RIFG~LTPJ for intergenerational dyads drawing together, no ROI pairs for same-generation dyads drawing alone, and LIFG~RIFG and RIFG~RTPJ for same-generation dyads drawing together. HbR analyses were similar, in that LTPJ~LTPJ, LIFG~RIFG, and LIFG~LTPJ showed substantial differences between real and pseudo dyads for intergenerational dyads drawing together (reported in full in Supplementary Materials).

Change in INS levels across sessions. We observed a more positive slope of change for all ROI pairs that differed substantially between real and pseudo dyads, with one exception (black squares Figure 2A; solid lines in Figure 2B; Supplementary Table 4). The ROI pairs showing substantial differences were LIFG~RTPJ for intergenerational dyads drawing alone, LTPJ~LPTJ and RIFG~RTPJ for intergenerational dyads drawing together, no ROI pairs for same-generation dyads drawing alone, and RTPJ~RTPJ and LIFG~LTPJ for same-generation dyads drawing together. HbR analyses showed little similarity (reported in full in Supplementary Materials).

Figure 1. A) INS for all ROI pairs per task and group; black squares indicate that INS levels differ between real and pseudo dyads substantially (i.e., 95% HPD do not overlap). B) Visualisation of all possible ROI pairs (n=10). Dashed lines indicate ROI pairs where INS did not differ substantially between real and pseudo dyads. Solid lines indicate ROI pairs where INS differed substantially between real and pseudo dyads. Black solid lines indicate ROI pairs where task, group, and task*group contrasts showed substantial differences. Grey solid lines indicate ROI pairs where contrasts showed no substantial differences. C) INS levels with 95% HPD for all ROI pairs that differed between real and pseudo dyads, for which task, group, and task*group contrasts were computed. LIFG~LIFG: No differences. RIFG~RIFG: No differences. LTPJ~LTPJ: Task difference, drawing together > alone. LIFG~RIFG: Task difference, drawing together > alone. Interaction, drawing together > alone for intergenerational dyads. LIFG~LTPJ: No differences. RIFG~LTPJ: Task difference, drawing together > alone. RIFG~RTPJ: Task difference, drawing together > alone. Interaction, drawing together > alone for same generation dyads. Full list of estimates and contrasts with 95% HPD available in Supplementary Tables 1-2 for HbO. HbR analyses visualised in Supplementary Figure 2, showed the same pattern of results for LIFG~RIFG and RIFG~LTPJ.

Figure 2. A) Slope of change in INS across six sessions; black squares indicate that INS levels differ between real and pseudo dyads substantially (i.e., 95% HPD do not overlap). B) Visualisation of all possible ROI pairs (n=10). Dashed lines indicate ROI pairs where the slope of change in INS did not differ substantially between real and pseudo dyads. Solid lines indicate ROI pairs where the slope of change in INS differed substantially between real and pseudo dyads. Black solid lines indicate ROI pairs where task, group, and task*group contrasts showed substantial differences. C) Slope of change in INS across sessions for all ROI pairs that differed between real and pseudo dyads, for which task, group, and task*group contrasts were computed. Solid lines indicate INS~Session relationship was substantial (i.e., 95% HPD does not overlap with 0). Dashed lines indicate non-substantial INS~Session relationships. Grey shading shows 95%, 89%, and 50% intervals of the posterior predictive distributions. LTPJ~LTPJ: Interaction effect, drawing together > alone for intergenerational dyads. RTPJ~RTPJ: Interaction effects, drawing together < alone for intergenerational dyads and intergenerational < same generation dyads for drawing together. LIFG~LTPJ: Group effect, intergenerational < same generation dyads. LIFG~RTPJ: Interaction effects, drawing together < alone for intergenerational dyads, intergenerational > same generation dyads for drawing alone, and intergenerational < same generation dyads for drawing together. RIFG~RTPJ: Interaction effects, drawing together < alone for intergenerational dyads, intergenerational > same generation dyads for drawing alone, and intergenerational < same generation dyads for drawing together. Full list of estimates and contrasts with HPD available in Supplementary Tables 3-4 for HbO. HbR analyses, visualised in Supplementary Figure 3, showed the same pattern of results for RTPJ~RTPJ and the same interaction effect of drawing together < alone for intergenerational dyads.

2.2.2 Composition of non-homologous ROI pairs in intergenerational dyads

For the intergenerational dyads specifically, we examined whether INS differed as a function of the generational composition of the ROI pairs. In other words, whether an ROI belonging to an older or younger member of a dyad influenced INS levels for all sessions or slopes across sessions. Full table of estimates and contrasts with HPD available in Supplementary Table 5.

INS levels for all sessions combined. When intergenerational dyads drew together, there was a trend of greater INS when the LIFG belonged to the older adult and the LTPJ belonged to the younger adult than vice versa. We observed no substantial differences in INS for any composition of any ROI pair for drawing alone. **Change in INS levels across sessions.** No substantial differences in the INS slopes across sessions for any composition of any ROI pair, for drawing alone or together.

2.2.3 Contrasting drawing conditions and groups

INS levels for all sessions combined. We observed no substantial differences between groups (i.e. group effect) but did observe task effects. We observed substantially greater INS for drawing together than alone in LTPJ~LTPJ, LIFG~RIFG, RIFG~LTPJ and RIFG~RTPJ (i.e., task effect). With respect to interaction between group and task, we found that in LTPJ~LTPJ, INS was substantially greater for drawing together than alone for intergenerational dyads. In RIFG~RTPJ, INS was substantially greater for drawing together than alone for same generation dyads. Parameter estimates per ROI pair, task and group visualised in Figure 1C. Full list of contrasts with 95% HPD supplied in Supplementary Table 6.

Change in INS levels across sessions. We observed no substantial difference between drawing alone and together (i.e., no task effects) but did observe the following group effect: We observed a substantially more positive INS slope for the same generation than the intergenerational dyads in LIFG~LTPJ. We also found several interactions effects. Intergenerational dyads showed a substantially more positive INS slope for drawing together than alone in LTPJ~LPTJ, and substantially more positive INS slopes for drawing alone than together in RTPJ~RTPJ, LIFG~RTPJ, and RIFG~RTPJ. This difference was substantial for drawing together, whereby the INS slope was more positive for same-generation dyads, relative to intergenerational dyads. Same-generation dyads further showed substantially more positive INS slopes than intergenerational dyads for drawing together in RTPJ~RTPJ, LIFG~RTPJ and RIFG~RTPJ. Intergenerational dyads showed substantially more positive INS slopes than same generation dyads when drawing alone in LIFG~RTPJ and RIFG~RTPJ. Parameter estimates per ROI pair, task and group are visualised in Figure 2C. Full list of contrasts with 95% HPD supplied in Supplementary Table 7.

2.2.4 Relationship between INS and self-report measures

Loneliness. *Summed.* We observed no substantial main, task, or group effect on the relationship between INS and summed loneliness scores. We observed an interaction effect in RIFG~RTPJ, for the intergenerational dyads, whereby the INS~summed loneliness relationship was substantially more negative for drawing alone than together (Figure 3; Supplementary Tables 10-11). *Difference.* No substantial main effect or group effects were observed. We observed a substantial effect of task for RIFG~RIFG, whereby the INS~loneliness difference relationship was substantially more negative for drawing together than alone. We observed substantial interaction effects for RIFG~RIFG, RIFG~RTPJ, and LTPJ~LTPJ. For RIFG~RIFG, we observed a substantial task difference for the intergenerational dyads only. For LTPJ~LPTJ, we observed a substantial task difference for intergenerational, but not same generation dyads and a substantial difference between groups for

drawing alone but not together. For RIFG~RTPJ, we observed a substantial task difference for same generation dyads only (Figure 3; Supplementary Tables 12-13).

Social closeness (OIS). *Summed*. We observed a substantial positive relationship between summed social closeness scores and INS in RIFG~RTPJ (main effect). In LIFG~LIFG, we observed a substantial effect of task, whereby there was a substantially more positive relationship between INS and summed social closeness for drawing together than alone. We observed a group effect for RIFG~RIFG, whereby intergenerational dyads showed a substantially more positive relationship than same generation dyads, and RIFG~RTPJ where intergenerational dyads showed a substantially more negative relationship than same generation dyads. Substantial interaction effects observed for LIFG~LIFG and RIFG~RIFG, RIFG~RTPJ. In LIFG~LIFG there was a substantial task difference for same generation dyads only. In RIFG~RIFG the interaction effect showed a substantial group difference for drawing alone only. In RIFG~RTPJ, we observed a substantial task difference for the same generation dyads only and a substantial group difference only when dyads drew alone (Figure 3; Supplementary Tables 14-15). *Difference*. No substantial main, task, or group effect on the relationship between INS and social closeness difference scores was observed. We did find a substantial interaction effect for RIFG~RIFG: There was a substantial task difference for both groups and a substantial group difference when drawing alone only (Figure 3; Supplementary Tables 16-17).

Allophilia. *Summed*. We observed a substantial negative relationship between summed allophilia scores and INS in LTPJ~LPTJ and LIFG~RIFG (main effect). We also found a substantial group effect for RIFG~RTPJ, where the relationship between INS and the summed allophilia scores was substantially more positive for the intergenerational dyads than same generation dyads. A substantial interaction effect was observed in LIFG~LTPJ, with a substantial task difference for intergenerational dyads only (Figure 3; Supplementary Table 18-19). *Difference*. No substantial main or group effects were observed. We observed a substantial task effect in RIFG~RIFG, where the relationship between INS and the difference in allophilia scores was substantially more positive for drawing together than alone. We found a substantial interaction effect for RIFG~RIFG, where we observed a substantial task difference for same generation dyads only (Figure 3; Supplementary Tables 20-21).

Figure 3. A) Slope of change in INS across levels of self-report measures for ROI pairs with at least one substantial relationship or substantial difference revealed by contrasts (for all ROI pairs see Supplementary Figures 5A-F; estimates and contrasts with 95 % HPD in Supplementary Tables 10-21). Lines indicate whether INS~Self-report measure relationship was substantial (solid) or non-substantial (dashed). Shading shows 95%, 89%, and 50% intervals for the posterior predictive distributions. Results of contrasts between task, group, and task*group contrasts for the panels above are as follows. *LIFG~LIFG Social closeness (Sum):* Task effect, negative relationship for drawing alone and drawing alone < together. Interaction effect, together > alone for same generation. *RIFG~RIFG Loneliness (Dif):* Task effect, drawing together < alone. *RIFG~RIFG Social Closeness (Sum):* Group effect, intergenerational > same generation dyads.

Interaction effect, intergenerational > same generation for drawing alone. *RIFG~RIFG Social Closeness (Dif):* Interaction
effects, drawing together > alone for intergenerational dyads, drawing together < alone for same generation dyads, and
intergenerational > same generation for drawing alone. *RIFG~RIFG Allophilia (Dif):* Task effect, positive relationship for
drawing together and drawing together > alone. Interaction effect, drawing together > alone for same generation dyads.
LTPJ~LTPJ Loneliness (Dif): Interaction effect, intergenerational < same generation dyads for drawing alone.
LTPJ~LTPJ Allophilia (Sum): Main effect, negative relationship. Group effect, negative relationship for intergenerational
dyads only. Task effect, negative relationship for drawing alone and together. *LIFG~RIFG Allophilia (Sum):* Main effect,
negative relationship. Group effect, negative relationship for same generation dyads only. *LIFG~LTPJ Allophilia (Sum):*
Interaction effect, drawing together < alone for intergenerational dyads. *RIFG~RTPJ Loneliness (Sum):* No differences.
RIFG~RTPJ Loneliness (Dif): Interaction effect, drawing together < alone for same generation dyads only. *RIFG~RTPJ*
Social Closeness (Sum): Main effect, positive relationship. Group effect, positive relationship for same generation dyads
and intergenerational < same generation. Task effect, positive relationship for drawing alone. Interaction effects, drawing
together < alone for same generation dyads, and intergenerational < same generation for drawing alone. *RIFG~RTPJ*
Allophilia (Sum): Group effect, intergenerational > same generation dyads. B) Visualisation of all possible ROI pairs
(n=10). Dashed lines indicate ROI pairs where i) neither INS for all sessions combines nor ii) the slope of change in INS
differed substantially between real and pseudo dyads. Solid lines indicate ROI pairs where INS (for sessions combined or
slope of change across sessions) differed substantially between real and pseudo dyads. Black solid lines indicate ROI
pairs where task, group, and task*group contrasts showed substantial differences. Analyses of HbR, visualised in
Supplementary Figure 4, showed similar pattern of results for Social Closeness (Sum) and Allophilia (Dif) in
RIFG~RIFG, as well as Loneliness (Dif) in LTPJ~LTPJ.

3   Discussion

We invited intergenerational and same generation dyads to complete a six-sessions of creative
drawing (resembling a community art program), and charted dyads' levels of loneliness, social
closeness, attitudes toward generations and INS as they formed social connections across 6 sessions.
Our first aim was to characterise INS within intergenerational dyads comprising one young and one
older adult. In a large sample (sessions combined), we show that INS among intergenerational dyads
is largely comparable to same generation dyads. Specifically, both dyad types show greater INS in
LTPJ~LTPJ, LIFG~RIFG, RIFG~LTPJ, and RIFG~RTPJ for drawing together than alone, with the
greatest differences in LIFG~RIFG for intergeneration dyads and RIFG~RTPJ for same generation
dyads. Our second aim was to chart the trajectory of INS in same generation and intergenerational
dyads, across repeated social interactions. While drawing together, ROI pairs that include the right
TPJ (RTPJ~RTPJ, LIFG~RTPJ, and RIFG~RTPJ) showed decreasing INS for intergenerational
dyads and increasing INS for same generation dyads. Our third aim was to characterise the
relationship between INS and social connectedness, via loneliness, social closeness, and attitudes
toward generations. We found that the relationship between loneliness and INS in RIFG~RIFG
differed when dyads drew alone and together: Greater similarity in dyadic levels of loneliness
predicted greater INS when dyads drew together and lower INS when they drew alone. Greater social
closeness predicted increasing INS in RIFG~RTPJ for all dyads and greater INS in RIFG~RIFG for
intergenerational than same generation dyads. With respect to dyadic attitudes towards generations,
we observed that INS in LIFG~RIFG and LTPJ~LTPJ was elevated among dyads who felt further
from other generations (i.e., closer to their own). Below, we explore the implications of these
findings.

Individual's levels of loneliness and social closeness decreased and increased, respectively, across
sessions, for both generational constellations. These findings match the target outcomes of the
community programs proposed by policy makers, researchers, and health practitioners (1,5,21,22)
and indicate that social connections were indeed formed.

Considering all sessions together, we observed that the generational constellation (i.e.,
intergenerational vs. same generation) predicted differences in individual's levels of loneliness, social
closeness and attitudes toward other generations. For example, simply being in the presence of a

similar or quite differently aged person may alter one's perceptions of one's loneliness. If the other
person is of a similar age, one might feel more lonely than if the person is older, in which case one
would self-report feeling less lonely – perhaps by virtue of social comparison (81). This finding
should be validated in future work by comparing intergenerational dyads with same generation dyads
composed of two older adults, as well as two younger adults. Greater feelings of social closeness
among intergenerational, as compared to same generation, dyads may relate to overcoming perceived
generational differences or generational stereotypes (82). Notably, the distributions of difference in
social closeness did not differ between groups, suggesting that the increased social closeness
observed for intergenerational dyads was not driven by older or younger adults' responses alone, but
rather by both age groups. Intergenerational dyads held more charitable attitudes toward other
generations than same generation dyads, aligning with findings that contact with older adults reduces
younger adults' negative stereotypes of and anxiety regarding older adults (83). Among
intergenerational dyads, individuals' attitudes toward generations were linked to their social
closeness ratings, highlighting possible bidirectional interplay between attitudes and social closeness.
Moreover, among same generation dyads, greater loneliness was associated with a weaker preference
towards one's 'own' (younger) generation, highlighting the window of opportunity for
intergenerational social activities in combatting loneliness.

This study offers novel insights into INS during collaboration in dyads composed of older and
younger adults, adding rich empirical data to the two existing studies that involve older adults
(62,63). The fNIRS hyperscanning literature on collaboration focuses on the prefrontal and inferior
frontal cortices and the TPJ (47). The bilateral IFG play a role in the initiation and reception of joint
attention (48–50) and the bilateral TPJ are central to the integration of social information (53,54).
Moreover, with some exceptions (e.g., 62), hyperscanning studies typically only report homologous
pairs of brain areas (e.g., LIFG~LIFG), for which the interpretation of heightened INS may be
simpler. We preregistered and now report analyses of INS that include non-homologous ROI pairs
(e.g., LIFG~RIFG) and offer possible reasoning for observations of heightened INS.

In our large sample (all sessions combined), we show that INS in intergenerational dyads is largely
comparable to same generation dyads. That is, we observed greater INS in LTPJ~LTPJ, LIFG~RIFG,
RIFG~LTPJ, and RIFG~RTPJ for drawing together than alone. This aligns with the existing findings
that collaboration co-occurs with greater INS in brain regions that process social information. We
interpret these findings as evidence that collaborative drawing results in heightened integration of
social information (LTPJ~LTPJ), joint attention (LIFG~RIFG), and perhaps integration of the other
person's maintenance of joint attention as social information (RIFG~LTPJ and RIFG~RTPJ). We
also observed robust differences in INS while drawing together and alone in separate ROI pairs for
each group: LIFG~RIFG for the intergenerational dyads and RIFG~RTPJ for the same generation
dyads. We cautiously speculate that intergenerational dyads' level of cognitive control while drawing
together is particularly high and aligned within dyads, and that same generation dyads may evaluate
and integrate information regarding the strength or continuity of their partner's joint attention. This
interpretation requires further investigation using paradigms that can quantify joint attention and
cognitive control.

INS in IFG is proposed by some researchers to be unique to verbal communication, though a few
studies show that INS in IFG may also play a role in non-verbal communication (47). With our very
large sample, we can provide strong evidence that INS involving the IFG occurs during non-verbal
communication, such as silent collaborative drawing.

Cross-sectional studies suggest that INS levels during collaboration decrease with higher levels of
familiarity (such as friendship or stable romantic relationships) that require repeated social
interactions to establish (45,62,72,78–80). We document INS trajectories during collaboration and
independent drawing by two (initial) strangers with the highest temporal granularity to date – 6
consecutive time points. Our findings provide clear evidence of INS changes from a dyad's first
encounter to the final recorded session, entailing ~6 hours of time spent together and ~90 minutes of
collaborative activity spread across 6 weeks.

We observed that while dyads drew together, ROI pairs that include right TPJ (RTPJ~RTPJ,
LIFG~RTPJ, and RIFG~RTPJ) showed decreasing INS among intergenerational dyads but increasing
INS among same generation dyads. This may suggest that, in the context of collaborative drawing,
intergenerational dyads' monitoring and integration of their partner's levels of joint attention may
decrease with increasing familiarity, whereas same generation dyads may monitor each other's levels
of joint attention more closely as familiarity increases. Across sessions, INS increased in RTPJ~RTPJ
for same generation dyads and LTPJ~LTPJ for intergenerational dyads. This difference in
lateralisation may be related to the integration of "matched" or "mismatched" predictions. Whereas
the right TPJ is associated with identifying "mismatched" predictions, the left TPJ encodes both
"matched" and "mismatched" predictions (84). Accordingly, it is possible that same generation dyads
become increasingly sensitive to "mismatched" predictions with increasing familiarity while
intergenerational dyads may encode more "matched" predictions with increasing familiarity.
Alternatively for intergenerational dyads, the frequency or absolute amount of mismatched
predictions may decrease with increasing familiarity.

We found that the relationship between loneliness and INS in RIFG~RIFG differed when dyads drew
alone and together: Greater similarity in dyadic levels of loneliness predicted greater INS when dyads
drew together and lower INS when they drew alone. This may point to how differences in loneliness
can shape joint attention. Experiences of loneliness predict changes in cognition, such as perceptual
speed, memory, and negatively biased perceptions of social information (10,11,13–15). It follows
that these changes in cognition may co-occur with changes in joint attention or may alter the facility
with which individuals engage in joint attention (85). It is possible that lonelier individuals may
require more numerous neural resources to maintain joint attention or alternatively present
hypervigilant tendencies, both of which could result in elevated INS in RIFG~RIFG (19,86,87).

To understand INS beyond the changes in social connectedness incurred by repeated social
interactions, we examined the extent to which dyads' self-reported feelings of social closeness were
reflected in INS. We observed that greater reported social closeness predicted greater INS in
RIFG~RTPJ for all dyads and in RIFG~RIFG for the intergenerational dyads. We interpret the
increasing INS in RIFG~RTPJ as meaning that dyads who feel close to one another monitor and
integrate each other's joint attention more than dyads who feel less socially close. The increasing INS
for intergenerational dyads in RIFG~RIFG may reflect joint attention as well as heightened arousal
or the facilitation of joint attention via heightened arousal (62,79,88,89).

With respect to dyadic attitudes towards generations, we observed that INS in LIFG~RIFG and
LTPJ~LTPJ decreased as feelings of closeness to other generations increased. As this analysis is
particularly exploratory in nature, with no empirical precedent, we offer a tentative suggestion that
similarity in attitudes towards generations may give dyads a more internally consistent basis upon
which they form predictions about each other's behaviour (i.e., elsewhere termed "common ground";
,80). These findings can serve as a point of departure for research examining the impact of different
sources of 'common ground' on INS. For example, studies could explore whether cumulative life

experience (e.g., number of years lived) gives older adults an advantage for finding common ground
and/or whether similarities in socialisation (e.g., culture, age, socio-economic status, hobbies) play
bigger role in establishing common ground that could shape INS.

As recommended by Simons et al. (90), we make explicit the constraints on generality of this work.
Our findings only generalise to the targeted brain regions (i.e., bilateral IFG and TPJ). Though our
study offers typical coverage for fNIRS hyperscanning studies, we encourage the use of high-density
fNIRS optode arrays in future research on this topic (91). We observed overlap between HbO and
HbR results for analyses of all sessions together and INS trajectories across sessions, with some
overlap in HbO and HbR results for relationships between INS and self-reported measures. Our
adequate pairing of sample size and conservative Bayesian approach notwithstanding, we encourage
other researchers to replicate these findings. Moreover, the INS self-report measures are not
necessarily generalisable across cultures, due to differences in formation of relationships (92,93).
Though our findings pertaining to changes in loneliness and social closeness across sessions are in
line with findings from behavioural and qualitative studies performed as part of more elaborate and
enriching community art programs, our sessions were attended by one dyad at a time, with partners
assigned based on availability. Future studies could expand our paradigm to small group sessions
with multiple dyads and free partner choice to reflect community programs more closely. We have no
reason to believe that the results depend on other characteristics of the participants, materials, or
context.

One limitation of this study is that we did not have the resources to include a group of same
generation dyads composed of two older adults, though this would have been the optimal design.
Without such a group, we cannot be sure that differences between groups are purely related to the
generational constellation and not age-related differences in neural processing. Future research
should compare same generation dyads across different age ranges. A second limitation of this study
is that three optodes of one Photon Cap device unexpectedly became defective partway through data
collection (and were replaced as rapidly as possible). This resulted in the exclusion of several
participants' right TPJ channels for the affected sessions. Thus, the number of data points for ROI
pairs involving right TPJ, particularly RTPJ~RPTJ, is lower than for other ROI pairs (Supplementary
Figure 7). Our Bayesian approach accounted for this difference in sample size per ROI pair by partial
pooling (for an excellent tutorial, see: ,94).

4   Conclusion

In this study, we documented INS in intergenerational dyads including younger and older adults and
charted the trajectory of INS during the formation of social connections facilitated by collaborative
and creative activity. Our main findings were that, in same generation and intergenerational dyads,
INS was greater while dyads drew together than alone. Across sessions, intergenerational dyads' INS
decreased and same generation dyads' INS increased. Heightened feelings of social closeness were
associated with greater INS in RIFG~RTPJ, and greater differences in loneliness levels between dyad
members predicted lower INS in RIFG~RIFG. This longitudinal, intergenerational study reinforces
that the dynamics of INS during the formation of social connections are multifaceted and merit
further empirical attention.

5   Methods

5.1   Participants

We recruited 122 community-dwelling participants from Zurich, Switzerland. Of these, 31 were older
adults (aged 69+ years) and 91 were younger adults (aged 18–35 years). We assigned participants to
intergenerational dyads (n = 31) and same generation dyads (n = 30) based on availability for
sessions (e.g., matching people available at the same day/time for 6 consecutive weeks). Due to the
logistical challenge posed by recruiting for and scheduling a multi-session study of this scale, gender
pairings were not prioritised (see Supplementary Figures 8 and 9 for summary of gender pairings and
session spacing). Intergenerational dyads consisted of an older adult (76 ± 4 years); 18 female, 13
male) with a younger adult (24 ± 4 years; 19 female, 11 male, 1 other). Same generation dyads
consisted of two younger adults (22.5 ± 3.6 years; 36 female, 24 male). All dyads started the study as
strangers. To be included, participants had no known history of neurological or psychological
disorders (e.g., stroke, concussion, ADHD, autism, schizophrenia, depression) and were fluent in
German.

We recruited participants via the University of Zurich's Healthy Longevity Centre, ETH Zurich's
DESCIL student pool, clubs and organizations for senior citizens (e.g., theatre groups and choirs),
and social media. We obtained written informed consent from all participants. Ethical approval was
obtained from the Ethics Committee of the Canton Zurich (Ref: 2023-01073). The study was
conducted according to the Declaration of Helsinki.

5.2   Measures

5.2.1 Self-report measures

Loneliness. Participants completed the *6-Item Scale for Overall, Emotional, and Social Loneliness*
(95) at the beginning of each session.

Social closeness. Participants indicated how socially close they felt to the other member of their dyad
using the *Inclusion of Other in the Self Scale* (96) at the end of each session. This measure uses Venn
diagrams with incremental increases from no overlap to complete overlap as visual analogue
representation of social relationships.

Attitudes toward other generations. The *Allophilia Scale* (97) is typically used to characterise
participants' attitudes toward an outgroup (e.g., people of another ethnicity or political leaning). We
adapted this scale in three variants: a) "younger people", "older people", and "people my age". At the
beginning of each session all participants completed the scale with "people my age". Then older
adults completed the scale with "younger people" and younger adults completed the scale with "older
people". To obtain baseline-corrected attitudes towards other generations, we subtracted each
participant's score for "people my age" from their score for the age group they did not belong to.

5.2.2 fNIRS recordings

We recorded fNIRS signals using two Cortivision Photons Cap (Cortivision sp. z o.o., Lublin,
Poland) and Cortiview software. Each device had 16 LED sources, which emitted wavelengths of 760
and 850 nm, and 10 detectors (26 optodes in total). The sample rate was 4-6 Hz. We positioned
optodes according to the 10-5 EEG positions (98) over bilateral IFG and TPJ based on MNI-space
coordinates derived from the Neurosynth database (https://neurosynth.org/). Specifically, we used the
search terms 'ifg' and 'tpj' and recorded the coordinates of the area with the highest activation in
each the left and right hemisphere, defining the 'centre' of our ROIs. There were the following: *left*
IFG: -48, 22, 8; *right IFG:* 50, 22, 16; *left TPJ:* -56, -54, 20; and *right TPJ:* 56, -52, 18. See
Supplementary Figure 10 for visualisation of ROI centre coordinates and Supplementary Figure 11
for optode arrangement.

Digitisation of optodes. After placement of the cap according to fiducial markers and before attaching optodes, we recorded a 3D image of each participant's head using Structure Sensor Pro (model SA39) attached to an Apple iPad Pro (v. 10). The 3D head scan was processed using MATLAB (v. 2023b) and the Fieldtrip toolbox (99). Labels for all detectors and sensors and fiducial reference points were assigned to the 3D head scan and then co-registered to MNI space. Next, we obtained the interoptode distance per channel, the coordinates of the point of measurement between optodes making up each channel, as well as distance from the point of measurement to the centre of the ROI. See Supplementary Figure 12 for visualisation of optode distances from ROI centres.

5.3 Procedure

Dyads attended six sessions, with approximately one week between sessions (Supplementary Figure 9 shows schedule of sessions and number of days between sessions per dyad). Sessions were led by a female lead-researcher who provided instructions following a script (available on OSF: https://osf.io/zms2v) and answered participants' technical and scientific queries. A second female researcher was present to aid with technical steps and did not speak unless directly addressed. Both researchers politely deflected participants' social inquiries.

The order of events at each session is visualised in Figure 4A. At the beginning of each session, dyads completed self-report measures of loneliness and attitudes towards generations (allophilia) on an iPad, with a felt divider between participants providing privacy. Then the fNIRS caps were placed on participants' heads and the second researcher recorded 3D models of each participant's head with cap on (participants instructed to sit very still while researcher circled participant with 3D scanner). Next, the lead researcher reminded that participants could converse at any time except while drawing. The researchers attached optodes to the caps, ensuring that optodes came in direct contact with the scalp by clearing hair. Then fNIRS devices were turned on, paired with recording laptops (n = 2) and fNIRS signals were verified for best possible calibration. Video recording was initiated (one GoPro11 Black was positioned with a direct view of participants as shown in Figure 4B, to allow for future motion tracking analyses).

Next, participants drew three drawings with oil pastels, without talking. Participants were instructed that they would have 5 min to draw anything they liked, and that they would complete the first drawing alone and the second and third together on a single piece of paper. No further instructions were provided (e.g., no suggestions for motifs). While participants drew alone, they were separated by a felt divider, ensuring that they could not see each other. The divider was removed when they drew together. In cases where participants conversed to plan their drawings, the researchers waited for a natural pause to begin the 5 min timer. Each drawing was labelled and digitised after the session.

After completing the three drawings, the participants completed a short collaborative activity (different at each session to keep participants motivated). Then, fNIRS recordings were stopped, and still wearing the fNIRS caps, participants completed a self-report measure of social closeness (OIS) and provided written commentary on their experience in the session (Figure 4A). After the final session, the lead researcher held a semi-structured interview with each participant individually to gain qualitative insights into the participant's experience.

Self-report measures were administered via RedCap®. fNIRS recordings were synchronised using triggers sent from PsychoPy (v. 2022.2.5; ,100) marking the beginning of each drawing and the collaborative activity.

Alone　　　　　　　　　　　**Together**

Figure 4. A) Flowchart visualising the protocol followed in each session. B) Left: An intergenerational dyad drawing alone on separate pages with a grey felt divider between them, blocking their view of each other. Right: The same intergenerational dyad drawing together on one page. The overlay of colourful lines shows the 2D motion tracking, which was captured, but is not reported here.

5.4 Data analysis

5.4.1 Interbrain synchrony

5.4.1.1 Individual level

Channel selection. We aimed to record from the identical position on participants heads at each session, however due to the stretchy nature of neoprene caps and the inevitability of hair growth/haircuts, optode positions varied from session to session. To overcome this challenge, we prioritise proximity to the centre of each ROI (101). That is, for each participant at each session, we selected the best channel per ROI. First, we excluded channels without visible cardiac rhythms (~1 Hz), with a scalp-coupling index (SCI; 102) of <.7 for frequencies between 0.7-1.35 Hz, and with an interoptode distance <20 mm or >40 mm. Next, we ranked the remaining channels by distance from the centre of the ROI (closest to furthest) and selected the closest channel as the 'best channel' (Supplementary Figure 13). See Supplementary Table 0 for description of number of recordings and sessions excluded, as well as reasons with for exclusion.

Preprocessing. We preprocessed the fNIRS signals using MNE (version 1.71; ,103,104), MNE-NIRS (version 0.71; ,105). As summarised in Supplementary Figure 13, the signal was converted from raw intensity to optical density. Motion artefacts were corrected using the temporal derivative distribution repair algorithm (106) and short channel regression was applied using the short channel with the highest SCI (105). Next, the signal was converted to concentrations of HbO and HbR using the Modified Beer–Lambert law (107,108) using an age-specific partial pathlength factor to account for age-related differences in the biological tissues probed by NIR light (109,110). This algorithm is suitable for participants aged 0-70 years. For participants aged >70 years, we specified the age parameter as 70. Last, a bandpass filter from 0.015 to 0.4 Hz was applied.

5.4.1.2 Dyad level

Wavelet transform coherence (WTC). We computed INS values for all possible ROI pairs (N = 10; homologous ROI pairs = LIFG~LIFG, RIFG~RIFG, LTPJ~LTPJ, RTPJ~RTPJ; non homologous ROI pairs = LIFG~RIFG, LIFG~LTPJ, LIFG~RTPJ, LTPJ~RTPJ, RIFG~LTPJ, RIFG~RTPJ) for each real dyad and pseudo dyad (Supplementary Figure 13). Pseudo dyads are dyads made by pairing participants who never interacted, but align with our group types (e.g., both younger adults, or one older and one younger adult). We computed INS values for pseudo dyads to obtain a null distribution against which to compare INS in real dyads. Differences in INS between real and pseudo dyads for specific ROI pairs indicate that INS levels are the result of the real interactions of real pairs and do not occur spuriously between any two people completing the same tasks in the same setting. WTC was computed using HyPyP (version 0.5.0b11; ,111) with a complex Morlet wavelet, for which the period range was defined as 5-50 seconds (0.02-0.2 Hz).

5.4.1.3 Group level

For our preregistered statistical analyses, we followed a Bayesian approach to multi-level regression (112). We fit multi-level models using the brms package (version 22.2; ,113) in the R language (version 4.5.1; ,114) within the RStudio IDE (version 2025.05.1+513; ,115). From these models, we report and interpret the posterior distribution of key parameters using a 95% credible interval, calculated via the 95% highest posterior density region (HPD) method (112). For readers more accustomed to a Frequentist approach that presents p-values, we recommend perusing Kruschke & Liddell (116), and we offer the following (simplified) heuristic for interpreting 95% HPDs: Parameter and comparison estimates can be said to be *substantial* when the 95% HPD does not contain zero and to be trends when the tip of an 95% HPD-tail overlaps with zero.

Self-report measures. To assess the extent to which loneliness, social closeness, and attitudes toward generations change across the six sessions, we fit the following model for each self-report measure separately, using individual scores (i.e., not dyadic sum or difference scores): *Score ~ 1 + Group * Session + (1|DyadID/ParticipantID)*. We subsequently computed contrasts between groups (intergenerational vs. same generation).

INS for sessions combined. We first sought to establish which, if any, ROI pairs showed substantially different INS levels for real dyads relative to pseudo dyads. We fit separate models for each ROI pair (n = 10), task (n = 2; drawing alone or drawing together), and group (n = 2; intergenerational or same generation): *INS ~ Dyad Type * Session*, where *Dyad Type* refers to real and pseudo dyads. Contrasts between real and pseudo dyads per model identified the ROI pairs for further analysis (Supplementary Figure 13).

Changes in INS across sessions. We first sought to establish which, if any, ROI pairs showed substantially different INS trajectories across sessions for real dyads relative to pseudo dyads. We fit separate models for each ROI pair (n = 10), task (n = 2; drawing alone or drawing together), and group (n = 2; intergenerational or same generation): *INS ~ Dyad Type * Session*, where *Dyad Type* refers to real and pseudo dyads. Contrasts between real and pseudo dyads per model identified the ROI pairs for further analysis (Supplementary Figure 13).

Further analyses with identified ROI pairs. We first fit models to assess whether intergenerational dyads show any directionality for non-homologous ROI pairs (e.g., high INS for older adult's left IFG and younger adult's right IFG, but not the inverse). Next, we fit one model for all sessions combined and one for changes across sessions including only real dyads and the respectively

identified ROI pairs (models shown second tier of Supplementary Figure 13). We computed planned
contrasts to examine the effect of task (drawing alone vs. drawing together), group (intergenerational
vs. same generation), and the interaction between task and group. Finally, we fit a model (shown in
second-last tier of Supplementary Figure 13) to all ROI pairs identified (sessions combined and
across sessions) to examine the extent to which self-report measures of loneliness, social closeness
(OIS) and attitudes toward generations (allophilia) predict levels of INS. To capture dyadic aspects of
each measure, we included summed and absolute difference scores per dyad (117,118). We report
HbO findings in the main report and HbR findings in the Supplementary Material, with commentary
on commonalities throughout the main report.

Author contributions

743 RM: Conceptualization, equipment donation, data curation, formal analysis, Investigation,
744 Visualization, Writing – original draft. GD: Methodology, Software, Writing – review & editing.
745 ESC: Conceptualization, funding acquisition, Project administration, Writing – review & editing.

Acknowledgements

From the Social Brain Science Lab at ETH Zurich, we thank Tessa Portier and Medea Häuselmann
for their assistance with data collection, Linda Fanconi for scheduling participants' sessions, Alistair
Gadola for coding participant behaviour, Fanny Mougel for digitising the drawings. We thank
Courtney Casale, Chantal Nagel, and Tessa Portier for proofreading. From the CHU Sainte-Justine
Azrieli Research Center of the University of Montreal, we thank Patrice Fortin for collaborating to
make HyPyP suitable for fNIRS hyperscanning data and providing hands-on assistance while
package functions were in development. We also thank the survey center operated by UZH Healthy
Longevity Center, University of Zurich, for their assistance with recruitment of older adults from the
Zurich community.

Conflict of interest

757 The authors declare no potential conflict of interest.

Data availability statement

759 These data were collected after the submission of the following preregistration: https://osf.io/hz6tm.
760 The data collected and analysed for this study can be found in our repository entitled 'Longitudinal
761 perspectives on intergenerational inter-brain synchrony' on OSF: https://osf.io/xcgp6/.

Funding

Two Cortivision Photon Cap devices were provided to RM through the Cortivision Pathfinder
Program (grant number CPP-2023/09/01). Oil pastels were provided to RM by Caran D'Ache
(agreement number 712517). RM and ESC were supported by the Social Brain Sciences Lab at ETH
Zurich. GD was supported by the Institute for Data Valorization, Montreal and the Canada First
Research Excellence Fund (IVADO; CF00137433), the Fonds de recherche du Québec (FRQ;
285289), and the Natural Sciences and Engineering Research Council of Canada (NSERC; DGECR-
2023-00089).

References

1.   WHO. Social isolation and loneliness among older people: Advocacy brief [Internet]. World
Health Organization; 2021. Available from:
https://www.who.int/publications/i/item/9789240030749

2.   Petersen J. A meta-analytic review of the effects of intergenerational programs for youth and
older adults. Educational Gerontology [Internet]. 2023 Mar 4 [cited 2024 Mar 21];49(3):175–
89. Available from: https://doi.org/10.1080/03601277.2022.2102340

3.   Bartlett SP, Solomon P, Gellis Z. Comparative effectiveness of intergenerational service-
learning programs on student outcomes of knowledge, attitude, and ageism. Educational
Gerontology [Internet]. 2021 Dec 2 [cited 2023 Jun 13];47(12):559–73. Available from:
https://www.tandfonline.com/doi/full/10.1080/03601277.2021.2011596

4.   Gonzales E, Morrow-Howell N, Gilbert P. Changing medical students' attitudes toward older
adults. Gerontology & Geriatrics Education [Internet]. 2010 Aug 26 [cited 2024 Mar
27];31(3):220–34. Available from: https://doi.org/10.1080/02701960.2010.503128

5.   Scarpetti G, Shadowen H, Williams GA, Winkelmann J, Kroneman M, Groenewegen PP, et al.
A comparison of social prescribing approaches across twelve high-income countries. Health
Policy [Internet]. 2024 Apr [cited 2025 Feb 11];142:104992. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0168851024000022

6.   WHO. From loneliness to social connection [Internet]. World Health Organization; 2025.
Available from: https://www.who.int/publications/i/item/978240112360

7.   House JS, Landis KR, Umberson D. Social relationships and health. Science [Internet]. 1988 Jul
29 [cited 2025 Aug 26];241(4865):540–5. Available from:
https://www.science.org/doi/10.1126/science.3399889

8.   Wang F, Gao Y, Han Z, Yu Y, Long Z, Jiang X, et al. A systematic review and meta-analysis of
90 cohort studies of social isolation, loneliness and mortality. Nat Hum Behav [Internet]. 2023
Jun 19 [cited 2023 Jun 23]; Available from: https://www.nature.com/articles/s41562-023-
01617-6

9.   The Lancet. Loneliness as a health issue. The Lancet [Internet]. 2023 Jul [cited 2025 Aug
25];402(10396):79. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0140673623014113

10. Ayalon L, Shiovitz-Ezra S, Roziner I. A cross-lagged model of the reciprocal associations of
loneliness and memory functioning. Psychology and Aging [Internet]. 2016 May [cited 2025
Aug 28];31(3):255–61. Available from: https://doi.apa.org/doi/10.1037/pag0000075

11. Cachón-Alonso L, Hakulinen C, Jokela M, Komulainen K, Elovainio M. Loneliness and
cognitive function in older adults: Longitudinal analysis in 15 countries. Psychology and Aging
[Internet]. 2023 Dec [cited 2025 Aug 28];38(8):778–89. Available from:
https://doi.apa.org/doi/10.1037/pag0000777

12. Drewelies J, Windsor TD, Duezel S, Demuth I, Wagner GG, Lindenberger U, et al. Age
trajectories of perceptual speed and loneliness: Separating between-person and within-person
associations. Neupert S, editor. The Journals of Gerontology: Series B [Internet]. 2022 Jan 11

[cited 2025 Aug 28];77(1):118–29. Available from:
https://academic.oup.com/psychsocgerontology/article/77/1/118/6424221

13. Cacioppo JT, Hawkley LC. Perceived social isolation and cognition. Trends in Cognitive
Sciences [Internet]. 2009 Oct [cited 2025 Aug 28];13(10):447–54. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1364661309001478

14. Okruszek Ł, Piejka A, Krawczyk M, Schudy A, Wiśniewska M, Żurek K, et al. Owner of a
lonely mind? Social cognitive capacity is associated with objective, but not perceived social
isolation in healthy individuals. Journal of Research in Personality [Internet]. 2021 Aug [cited
2025 Aug 28];93:104103. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0092656621000404

15. Spithoven AWM, Bijttebier P, Goossens L. It is all in their mind: A review on information
processing bias in lonely individuals. Clinical Psychology Review [Internet]. 2017 Dec [cited
2025 Aug 28];58:97–114. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0272735816303336

16. Cacioppo S, Capitanio JP, Cacioppo JT. Toward a neurology of loneliness. Psychological
Bulletin [Internet]. 2014 [cited 2025 Aug 28];140(6):1464–504. Available from:
https://doi.apa.org/doi/10.1037/a0037618

17. Düzel S, Drewelies J, Gerstorf D, Demuth I, Steinhagen-Thiessen E, Lindenberger U, et al.
Structural Brain Correlates of Loneliness among Older Adults. Sci Rep [Internet]. 2019 Sep 19
[cited 2025 Aug 28];9(1):13569. Available from: https://www.nature.com/articles/s41598-019-
49888-2

18. Lam JA, Murray ER, Yu KE, Ramsey M, Nguyen TT, Mishra J, et al. Neurobiology of
loneliness: A systematic review. Neuropsychopharmacol [Internet]. 2021 Oct [cited 2025 Aug
28];46(11):1873–87. Available from: https://www.nature.com/articles/s41386-021-01058-7

19. Brilliant D, Takeuchi H, Nouchi R, Yokoyama R, Kotozaki Y, Nakagawa S, et al. Loneliness
inside of the brain: evidence from a large dataset of resting-state fMRI in young adult. Sci Rep
[Internet]. 2022 May 12 [cited 2025 Aug 28];12(1):7856. Available from:
https://www.nature.com/articles/s41598-022-11724-5

20. Lieberz J, Shamay-Tsoory SG, Saporta N, Esser T, Kuskova E, Stoffel-Wagner B, et al.
Loneliness and the social brain: How perceived social isolation impairs human interactions.
Advanced Science [Internet]. 2021 Nov [cited 2025 Aug 28];8(21):2102076. Available from:
https://onlinelibrary.wiley.com/doi/10.1002/advs.202102076

21. Holt-Lunstad J, Perissinotto C. Social isolation and loneliness as medical issues. N Engl J Med
[Internet]. 2023 Jan 19 [cited 2025 Aug 26];388(3):193–5. Available from:
http://www.nejm.org/doi/10.1056/NEJMp2208029

22. Wigfield A, Turner R, Alden S, Green M, Karania VK. Developing a new conceptual
framework of meaningful interaction for understanding social isolation and loneliness. Social
Policy and Society [Internet]. 2022 Apr [cited 2025 Aug 26];21(2):172–93. Available from:
https://www.cambridge.org/core/product/identifier/S147474642000055X/type/journal_article

23.   Bessaha ML, Sabbath EL, Morris Z, Malik S, Scheinfeld L, Saragossi J. A systematic review of
850   loneliness interventions among non-elderly adults. Clin Soc Work J [Internet]. 2020 Mar [cited
851   2025 Aug 26];48(1):110–25. Available from: http://link.springer.com/10.1007/s10615-019-
852   00724-0

853   24.   Noone C, Yang K. Community-based responses to loneliness in older people: A systematic
854   review of qualitative studies. Health & Social Care in the Community [Internet]. 2022 [cited
855   2025 Aug 26];30(4):e859–73. Available from:
856   https://onlinelibrary.wiley.com/doi/abs/10.1111/hsc.13682

857   25.   O'Rourke HM, Collins L, Sidani S. Interventions to address social connectedness and loneliness
858   for older adults: a scoping review. BMC Geriatr [Internet]. 2018 Dec [cited 2025 Aug
859   26];18(1):214. Available from: https://bmcgeriatr.biomedcentral.com/articles/10.1186/s12877-
860   018-0897-x

861   26.   Cigna US. Loneliness Index [Internet]. Cigna US; 2018. Available from:
862   https://www.multivu.com/players/English/8294451-cigna-us-loneliness-
863   survey/#:~:text=The%20survey%20of%20more%20than,people%20who%20really%20understa
864   nd%20them.

865   27.   Hawkley LC, Buecker S, Kaiser T, Luhmann M. Loneliness from young adulthood to old age:
866   Explaining age differences in loneliness. International Journal of Behavioral Development
867   [Internet]. 2022 Jan [cited 2024 Jul 16];46(1):39–49. Available from:
868   http://journals.sagepub.com/doi/10.1177/0165025420971048

869   28.   Qualter P, Vanhalst J, Harris R, Van Roekel E, Lodder G, Bangee M, et al. Loneliness across
870   the life span. Perspect Psychol Sci [Internet]. 2015 Mar [cited 2025 Aug 26];10(2):250–64.
871   Available from: https://journals.sagepub.com/doi/10.1177/1745691615568999

872   29.   Lee K, Jarrott SE, Juckett LA. Documented outcomes for older adults in intergenerational
873   programming: A scoping review: research. Journal of Intergenerational Relationships [Internet].
874   2020 Apr 2 [cited 2025 Aug 26];18(2):113–38. Available from:
875   https://www.tandfonline.com/doi/full/10.1080/15350770.2019.1673276

876   30.   Lokon E, Li Y, Kunkel S. Allophilia: Increasing college students' "liking" of older adults with
877   dementia through arts-based intergenerational experiences. Gerontology & Geriatrics Education
878   [Internet]. 2020 Oct 1 [cited 2023 Mar 17];41(4):494–507. Available from:
879   https://www.tandfonline.com/doi/full/10.1080/02701960.2018.1515740

880   31.   Newman S, Ward C, Smith T, Wilson J, McCrea J. Intergenerational programs: Past, present
881   and future. Rev. ed. New York: Routledge; 2014. 264 p.

882   32.   WHO. What is the evidence on the role of the arts in improving health and well-being? A
883   scoping review [Internet]. Copenhagen: World Health Organization; 2019. Available from:
884   https://iris.who.int/handle/10665/329834

885   33.   Anderson S, Fast J, Keating N, Eales J, Chivers S, Barnet D. Translating knowledge: Promoting
886   health through intergenerational community arts programming. Health Promotion Practice
887   [Internet]. 2017 Jan [cited 2023 Mar 21];18(1):15–25. Available from:
888   http://journals.sagepub.com/doi/10.1177/1524839915625037

34. Jenkins LK, Farrer R, Aujla IJ. Understanding the impact of an intergenerational arts and health
project: A study into the psychological well-being of participants, carers and artists. Public
Health [Internet]. 2021 May [cited 2023 Mar 21];194:121–6. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0033350621000883

35. Rubin SE, Gendron TL, Wren CA, Ogbonna KC, Gonzales EG, Peron EP. Challenging
gerontophobia and ageism through a collaborative intergenerational art program. Journal of
Intergenerational Relationships [Internet]. 2015 Jul 3 [cited 2023 Mar 21];13(3):241–54.
Available from: http://www.tandfonline.com/doi/full/10.1080/15350770.2015.1058213

36. Moffat R, Casale CE, Cross ES. Mobile fNIRS for exploring inter-brain synchrony across
generations and time. Frontiers in Neuroergonomics [Internet]. 2024 [cited 2024 Jan 4];4.
Available from: https://www.frontiersin.org/articles/10.3389/fnrgo.2023.1260738

37. Liu D, Liu S, Liu X, Zhang C, Li A, Jin C, et al. Interactive Brain Activity: Review and
Progress on EEG-Based Hyperscanning in Social Interactions. Frontiers in Psychology.
2018;9:1862.

38. Hirata M, Ikeda T, Kikuchi M, Kimura T, Hiraishi H, Yoshimura Y, et al. Hyperscanning MEG
for understanding mother-child cerebral interactions. Front Hum Neurosci [Internet]. 2014 Mar
4 [cited 2023 Jun 7];8. Available from:
http://journal.frontiersin.org/article/10.3389/fnhum.2014.00118/abstract

39. Misaki M, Kerr KL, Ratliff EL, Cosgrove KT, Simmons WK, Morris AS, et al. Beyond
synchrony: the capacity of fMRI hyperscanning for the study of human social interaction. Social
Cognitive and Affective Neuroscience [Internet]. 2021 Jan 18 [cited 2024 Sep 6];16(1–2):84–
92. Available from: https://academic.oup.com/scan/article/16/1-2/84/5940486

40. Holper L, Scholkmann F, Wolf M. Between-brain connectivity during imitation measured by
fNIRS. NeuroImage. 2012;63(1):212–22.

41. Cui X, Bryant DM, Reiss AL. NIRS-based hyperscanning reveals increased interpersonal
coherence in superior frontal cortex during cooperation. NeuroImage. 2012;59(3):2430–7.

42. Dommer L, Jäger N, Scholkmann F, Wolf M, Holper L. Between-brain coherence during joint
n-back task performance: A two-person functional near-infrared spectroscopy study.
Behavioural Brain Research [Internet]. 2012 Oct [cited 2023 Jun 20];234(2):212–22. Available
from: https://linkinghub.elsevier.com/retrieve/pii/S0166432812004378

43. Osaka N, Minamoto T, Yaoi K, Azuma M, Shimada YM, Osaka M. How two brains make one
synchronized mind in the inferior frontal cortex: fNIRS-based hyperscanning during
cooperative singing. Frontiers in Psychology [Internet]. 2015 [cited 2023 Oct 25];6. Available
from: https://www.frontiersin.org/articles/10.3389/fpsyg.2015.01811

44. Li L, Wang H, Luo H, Zhang X, Zhang R, Li X. Interpersonal neural synchronization during
cooperative behavior of basketball players: a fNIRS-based hyperscanning study. Frontiers in
Human Neuroscience [Internet]. 2020 [cited 2023 Oct 25];14. Available from:
https://www.frontiersin.org/articles/10.3389/fnhum.2020.00169

45.　Sun B, Xiao W, Lin S, Shao Y, Li W, Zhang W. Cooperation with partners of differing social
experience: An fNIRS-based hyperscanning study. Brain and Cognition [Internet]. 2021 Nov
[cited 2025 Sep 3];154:105803. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0278262621001238

46.　Xie H, Karipidis II, Howell A, Schreier M, Sheau KE, Manchanda MK, et al. Finding the neural
correlates of collaboration using a three-person fMRI hyperscanning paradigm. Proc Natl Acad
Sci USA [Internet]. 2020 Sep 15 [cited 2025 Jun 11];117(37):23066–72. Available from:
https://pnas.org/doi/full/10.1073/pnas.1917407117

47.　Czeszumski A, Liang SHY, Dikker S, König P, Lee CP, Koole SL, et al. Cooperative behavior
evokes interbrain synchrony in the prefrontal and temporoparietal cortex: A systematic review
and meta-analysis of fNIRS hyperscanning studies. eNeuro [Internet]. 2022 Mar [cited 2023 Jun
28];9(2):ENEURO.0268-21.2022. Available from:
https://www.eneuro.org/lookup/doi/10.1523/ENEURO.0268-21.2022

48.　Koechlin E, Ody C, Kouneiher F. The architecture of cognitive control in the human prefrontal
cortex. Science [Internet]. 2003 Nov 14 [cited 2025 Aug 28];302(5648):1181–5. Available
from: https://www.science.org/doi/10.1126/science.1088545

49.　Redcay E, Kleiner M, Saxe R. Look at this: The neural correlates of initiating and responding to
bids for joint attention. Front Hum Neurosci [Internet]. 2012 [cited 2025 Aug 28];6. Available
from: http://journal.frontiersin.org/article/10.3389/fnhum.2012.00169/abstract

50.　Yoshioka A, Tanabe HC, Sumiya M, Nakagawa E, Okazaki S, Koike T, et al. Neural substrates
of shared visual experiences: A hyperscanning fMRI study. Social Cognitive and Affective
Neuroscience [Internet]. 2021 Dec 30 [cited 2023 Jun 7];16(12):1264–75. Available from:
https://academic.oup.com/scan/article/16/12/1264/6310483

51.　Ding K, Wang H, Wang Q, Li H, Li C. Inhibitory control associated with the neural mechanism
of joint attention in preschoolers: An fNIRS evidence. International Journal of
Psychophysiology [Internet]. 2023 Oct [cited 2025 Sep 11];192:53–61. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0167876023004981

52.　Mundy P. A review of joint attention and social-cognitive brain systems in typical development
and autism spectrum disorder. Eur J of Neuroscience [Internet]. 2018 Mar [cited 2025 Sep
11];47(6):497–514. Available from: https://onlinelibrary.wiley.com/doi/10.1111/ejn.13720

53.　Jacoby N, Bruneau E, Koster-Hale J, Saxe R. Localizing pain matrix and theory of mind
networks with both verbal and non-verbal stimuli. NeuroImage [Internet]. 2016 Feb [cited 2023
Dec 13];126:39–48. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1053811915010472

54.　Saxe R, Kanwisher N. People thinking about thinking people: The role of the temporo-parietal
junction in "theory of mind." NeuroImage [Internet]. 2003 Aug;19(4):1835–42. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1053811903002301

55.　Dikker S, Wan L, Davidesco I, Kaggen L, Oostrik M, McClintock J, et al. Brain-to-brain
synchrony tracks real-world dynamic group interactions in the classroom. Current Biology

[Internet]. 2017 May [cited 2023 Jun 13];27(9):1375–80. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0960982217304116

56. Dumas G, Nadel J, Soussignan R, Martinerie J, Garnero. Inter-brain synchronization during
social interaction. PLOS ONE. 2010;5(8):e12166.

57. Labonté-LeMoyne É, Léger PM, Resseguier B, Bastarache-Roberge MC, Fredette M, Sénécal
S, et al. Are we in flow neurophysiological correlates of flow states in a collaborative game. In:
Proceedings of the 2016 CHI Conference Extended Abstracts on Human Factors in Computing
Systems [Internet]. San Jose California USA: ACM; 2016 [cited 2025 Aug 28]. p. 1980–8.
Available from: https://dl.acm.org/doi/10.1145/2851581.2892356

58. Sinha N, Maszczyk T, Wanxuan Z, Tan J, Dauwels J. EEG hyperscanning study of inter-brain
synchrony during cooperative and competitive interaction. In: 2016 IEEE International
Conference on Systems, Man, and Cybernetics (SMC) [Internet]. 2016 [cited 2025 Aug 28]. p.
004813–8. Available from: https://ieeexplore.ieee.org/document/7844990/

59. Toppi J, Borghini G, Petti M, He EJ, De Giusti V, He B, et al. Investigating Cooperative
Behavior in Ecological Settings: An EEG Hyperscanning Study. Yao D, editor. PLoS ONE
[Internet]. 2016 Apr 28 [cited 2025 Aug 28];11(4):e0154236. Available from:
https://dx.plos.org/10.1371/journal.pone.0154236

60. De Felice S, Hamilton AFDC, Ponari M, Vigliocco G. Learning from others is good, with
others is better: The role of social interaction in human acquisition of new knowledge. Phil
Trans R Soc B [Internet]. 2023 Feb 13 [cited 2023 Jun 22];378(1870):20210357. Available
from: https://royalsocietypublishing.org/doi/10.1098/rstb.2021.0357

61. Nguyen T, Bánki A, Markova G, Hoehl S. Studying parent-child interaction with
hyperscanning. In: Progress in Brain Research [Internet]. Elsevier; 2020 [cited 2023 Jun 7]. p.
1–24. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0079612320300455

62. Zhang Q, Liu Z, Qian H, Hu Y, Gao X. Interpersonal competition in elderly couples: A
functional near-infrared spectroscopy hyperscanning study. Brain Sciences [Internet]. 2023 Mar
31 [cited 2023 Jun 7];13(4):600. Available from: https://www.mdpi.com/2076-3425/13/4/600

63. Dikker S, Brito NH, Dumas G. It takes a village: A multi-brain approach to studying
multigenerational family communication. Developmental Cognitive Neuroscience [Internet].
2024 Feb [cited 2023 Dec 18];65:101330. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1878929323001354

64. United Nations. World population prospects 2024: Summary of results. New York: United
Nations; 2024.

65. Adel L, Moses L, Irvine E, Greenway KT, Dumas G, Lifshitz M. A systematic review of
hyperscanning in clinical encounters. Neuroscience & Biobehavioral Reviews [Internet]. 2025
Sep [cited 2025 Jun 17];176:106248. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0149763425002490

66. Bi X, Cui H, Ma Y. Hyperscanning Studies on Interbrain Synchrony and Child Development: A
Narrative Review. Neuroscience [Internet]. 2023 Oct [cited 2025 Aug 28];530:38–45. Available
from: https://linkinghub.elsevier.com/retrieve/pii/S0306452223004049

67. De Felice S, Chand T, Croy I, Engert V, Goldstein P, Holroyd CB, et al. Relational
neuroscience: Insights from hyperscanning research. Neuroscience & Biobehavioral Reviews
[Internet]. 2025 Feb [cited 2025 Aug 28];169:105979. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0149763424004482

68. Hoehl S, Bánki A, Brzozowska A, Carollo A, Kostorz K, Nguyen T, et al. A developmental
framework of interpersonal neural synchrony [Internet]. PsyArXiv; 2025 [cited 2025 Aug 28].
Available from: https://osf.io/m7uyq_v1

69. Kelsen BA, Sumich A, Kasabov N, Liang SHY, Wang GY. What has social neuroscience
learned from hyperscanning studies of spoken communication? A systematic review.
Neuroscience & Biobehavioral Reviews [Internet]. 2022 Jan [cited 2023 Jun 7];132:1249–62.
Available from: https://linkinghub.elsevier.com/retrieve/pii/S0149763420305650

70. Noreika V, Georgieva S, Wass S, Leong V. 14 challenges and their solutions for conducting
social neuroscience and longitudinal EEG research with infants. Infant Behavior and
Development [Internet]. 2020 Feb [cited 2023 Jun 7];58:101393. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0163638319301638

71. Turk E, Vroomen J, Fonken Y, Levy J, Van Den Heuvel MI. In sync with your child: The
potential of parent–child electroencephalography in developmental research. Developmental
Psychobiology [Internet]. 2022 Apr [cited 2023 Jun 7];64(3). Available from:
https://onlinelibrary.wiley.com/doi/10.1002/dev.22221

72. Bevilacqua D, Davidesco I, Wan L, Chaloner K, Rowland J, Ding M, et al. Brain-to-brain
synchrony and learning outcomes vary by student–teacher dynamics: Evidence from a real-
world classroom electroencephalography study. Journal of Cognitive Neuroscience [Internet].
2019 Mar 1 [cited 2023 Jun 13];31(3):401–11. Available from:
https://direct.mit.edu/jocn/article/31/3/401/28957/Brain-to-Brain-Synchrony-and-Learning-
Outcomes

73. Ellingsen DM, Isenburg K, Jung C, Lee J, Gerber J, Mawla I, et al. Brain-to-brain mechanisms
underlying pain empathy and social modulation of pain in the patient-clinician interaction. Proc
Natl Acad Sci USA [Internet]. 2023 Jun 27 [cited 2023 Jun 26];120(26):e2212910120.
Available from: https://pnas.org/doi/10.1073/pnas.2212910120

74. Grahl A, Anzolin A, Isenburg K, Lee J, Ellingsen DM, Jung C, et al. Brain and behavioral
correlates of the patient-clinician relationship: A longitudinal fMRI hyper-scanning study of
chronic pain patients. The Journal of Pain [Internet]. 2021 May [cited 2023 Jun 20];22(5):602.
Available from: https://linkinghub.elsevier.com/retrieve/pii/S1526590021001310

75. Grahl A, Anzolin A, Ellis S, Barton-Zuckerman M, Lee J, Jung C, et al. The patient-clinician
relationship, expectancy and prior experience can modulate fibromyalgia treatment outcomes: a
longitudinal fmrihyperscan study. The Journal of Pain [Internet]. 2023 Apr [cited 2023 Jun
23];24(4):74–5. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1526590023002560

76. Sened H, Gorst Kaduri K, Nathan Gamliel H, Rafaeli E, Zilcha-Mano S, Shamay-Tsoory S.
Inter-brain plasticity as a mechanism of change in psychotherapy: A proof of concept focusing
on test anxiety. Psychotherapy Research [Internet]. 2025 Jan 20 [cited 2025 Aug 28];1–15.
Available from: https://www.tandfonline.com/doi/full/10.1080/10503307.2025.2451798

77. Wang D, Ren Y, Chen W. Relationship evolution shapes inter-brain synchrony in affective
sharing: The role of self-expansion. Brain Struct Funct [Internet]. 2024 Jul 25 [cited 2025 Aug
28];229(9):2269–83. Available from: https://link.springer.com/10.1007/s00429-024-02841-0

78. Djalovski A, Dumas G, Kinreich S, Feldman R. Human attachments shape interbrain synchrony
toward efficient performance of social goals. NeuroImage [Internet]. 2021 Feb [cited 2023 Jun
22];226:117600. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1053811920310855

79. Long Y, Chen C, Wu K, Zhou S, Zhou F, Zheng L, et al. Interpersonal conflict increases
interpersonal neural synchronization in romantic couples. Cerebral Cortex [Internet]. 2022 Jul
21 [cited 2023 Jun 22];32(15):3254–68. Available from:
https://academic.oup.com/cercor/article/32/15/3254/6445112

80. Speer SPH, Mwilambwe-Tshilobo L, Tsoi L, Burns SM, Falk EB, Tamir DI. Hyperscanning
shows friends explore and strangers converge in conversation. Nat Commun [Internet]. 2024
Sep 5 [cited 2025 Jan 5];15(1):7781. Available from: https://www.nature.com/articles/s41467-
024-51990-7

81. Arnold AJ, Kappes HB, Klinenberg E, Winkielman P. The role of comparisons in judgments of
loneliness. Front Psychol [Internet]. 2021 Mar 24 [cited 2025 Sep 9];12:498305. Available
from: https://www.frontiersin.org/articles/10.3389/fpsyg.2021.498305/full

82. Galinsky AD, Ku G, Wang CS. Perspective-taking and self-other overlap: Fostering social
bonds and facilitating social coordination. Group Processes & Intergroup Relations [Internet].
2005 Apr [cited 2025 Sep 9];8(2):109–24. Available from:
https://journals.sagepub.com/doi/10.1177/1368430205051060

83. Liao T, Zhuoga C, Chen X. Contact with grandparents and young people's explicit and implicit
attitudes toward older adults. BMC Psychol [Internet]. 2023 Sep 26 [cited 2024 Jul
18];11(1):289. Available from:
https://bmcpsychology.biomedcentral.com/articles/10.1186/s40359-023-01344-7

84. Doricchi F, Lasaponara S, Pazzaglia M, Silvetti M. Left and right temporal-parietal junctions
(TPJs) as "match/mismatch" hedonic machines: A unifying account of TPJ function. Physics of
Life Reviews [Internet]. 2022 Sep [cited 2023 Oct 5];42:56–92. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1571064522000434

85. Corbin IM, Dhand A. Unshared minds, decaying worlds: towards a pathology of chronic
loneliness. The Journal of Medicine and Philosophy: A Forum for Bioethics and Philosophy of
Medicine [Internet]. 2024 Jul 11 [cited 2025 Sep 10];49(4):354–66. Available from:
https://academic.oup.com/jmp/article/49/4/354/7685476

86. Hampshire A, Chamberlain SR, Monti MM, Duncan J, Owen AM. The role of the right inferior
frontal gyrus: inhibition and attentional control. NeuroImage [Internet]. 2010 Apr [cited 2022

Sep 13];50(3):1313–9. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S1053811909013986

87.   Saporta N, Scheele D, Lieberz J, Nevat M, Kanterman A, Hurlemann R, et al. Altered activation
in the action observation system during synchronization in high loneliness individuals. Cerebral
Cortex [Internet]. 2022 Dec 20 [cited 2024 Aug 8];33(2):385–402. Available from:
https://academic.oup.com/cercor/article/33/2/385/6537595

88.   Bast N, Polzer L, Raji N, Schnettler L, Kleber S, Lemler C, et al. Early intervention increases
reactive joint attention in autistic preschoolers with arousal regulation as mediator. Eur Child
Adolesc Psychiatry [Internet]. 2025 May 10 [cited 2025 Sep 11]; Available from:
https://link.springer.com/10.1007/s00787-025-02738-1

89.   Lachat F, Hugueville L, Lemaréchal JD, Conty L, George N. Oscillatory brain correlates of live
joint attention: a dual-EEG study. Front Hum Neurosci [Internet]. 2012 [cited 2025 Sep 11];6.
Available from: http://journal.frontiersin.org/article/10.3389/fnhum.2012.00156/abstract

90.   Simons DJ, Shoda Y, Lindsay DS. Constraints on Generality (COG): A Proposed Addition to
All Empirical Papers. Perspect Psychol Sci [Internet]. 2017 Nov [cited 2025 Sep
29];12(6):1123–8. Available from:
https://journals.sagepub.com/doi/10.1177/1745691617708630

91.   Shuo G, Li Y, Geng Y, Li D, Xu Q, Yang D, et al. Advancing inter-brain synchrony
measurement: A comparative hyperscanning study of high-density diffuse optical tomography
and functional near-infrared spectroscopy [Internet]. bioRxiv; 2025. Available from:
https://www.biorxiv.org/content/10.1101/2025.07.02.662877v1.abstract

92.   Goodwin R. Personal relationships across cultures [Internet]. 1st ed. Routledge; 2013 [cited
2025 Sep 12]. Available from: https://www.taylorfrancis.com/books/9780203434161

93.   Lu P, Oh J, Leahy KE, Chopik WJ. Friendship importance around the world: Links to cultural
factors, health, and well-being. Front Psychol [Internet]. 2021 Jan 18 [cited 2025 Sep
12];11:570839. Available from:
https://www.frontiersin.org/articles/10.3389/fpsyg.2020.570839/full

94.   Nalborczyk L, Batailler C, Vilain A, Bürkner PC. An introduction to Bayesian multilevel
models using brms: A case study of gender effects on vowel variability in standard indonesian.
Journal of Speech, Language, and Hearing Research. 2019;62:1225–42.

95.   De Jong Gierveld J, van Tilburg T. A 6-item scale for overall, emotional, and social loneliness:
Confirmatory tests on survey data. Res Aging [Internet]. 2006 Sep [cited 2023 Mar
20];28(5):582–98. Available from: http://journals.sagepub.com/doi/10.1177/0164027506289723

96.   Aron A, Aron EN, Smollan D. Inclusion of other in the self scale and the structure of
interpersonal closeness. Journal of Personality and Social Psychology. 1992;63(4):17.

97.   Pittinsky TL, Rosenthal SA, Montoya RM. Measuring positive attitudes toward outgroups:
Development and validation of the Allophilia Scale. In: Tropp LR, Mallett RK, editors. Moving
beyond prejudice reduction: Pathways to positive intergroup relations [Internet]. Washington:

American Psychological Association; 2011 [cited 2023 Mar 23]. p. 41–60. Available from: http://content.apa.org/books/12319-002

98. Oostenveld R, Praamstra P. The five percent electrode system for high-resolution EEG and ERP measurements. Clinical Neurophysiology. 2001;112(4):713–9.

99. Oostenveld R, Fries P, Maris E, Schoffelen JM. Fieldtrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Computational Intelligence and Neuroscience [Internet]. 2011 [cited 2024 Jul 9];2011:1–9. Available from: http://www.hindawi.com/journals/cin/2011/156869/

100. Peirce JW, Hirst RJ, MacAskill MR. Building Experiments in PsychoPy [Internet]. 2nd ed. London: Sage; 2022. Available from: https://uk.sagepub.com/en-gb/eur/building-experiments-in-psychopy/book273700

101. De Felice S, Hakim U, Gunasekara N, Pinti P, Tachtsidis I, Hamilton A. Having a chat and then watching a movie: how social interaction synchronises our brains during co-watching. Oxford Open Neuroscience [Internet]. 2024 Feb 1 [cited 2024 May 17];3:kvae006. Available from: https://academic.oup.com/oons/article/doi/10.1093/oons/kvae006/7634764

102. Pollonini L, Olds C, Abaya H, Bortfeld H, Beauchamp MS, Oghalai JS. Auditory cortex activation to natural speech and simulated cochlear implant speech measured with functional near-infrared spectroscopy. Hearing Research [Internet]. 2014;309:84–93. Available from: http://dx.doi.org/10.1016/j.heares.2013.11.007

103. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MEG and EEG data analysis with MNE-Python. Frontiers in Neuroscience. 2013;7:1–13.

104. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MNE software for processing MEG and EEG data. NeuroImage [Internet]. 2014;86:446–60. Available from: http://dx.doi.org/10.1016/j.neuroimage.2013.10.027

105. Luke R, Larson E, Shader MJ, Innes-Brown H, Van Yper L, Lee AKC, et al. Analysis methods for measuring passive auditory fNIRS responses generated by a block-design paradigm. Neurophotonics. 2021;8(2):025008.

106. Santosa H, Huppert TJ, Fishburn F, Zhai X. Investigation of the sensitivity-specificity of canonical- and deconvolution-based linear models in evoked functional near-infrared spectroscopy. Neurophotonics [Internet]. 2019;6(2):1–10. Available from: https://www.spiedigitallibrary.org/journals/neurophotonics/volume-6/issue-02/025009/Investigation-of-the-sensitivity-specificity-of-canonical--and-deconvolution/10.1117/1.NPh.6.2.025009.full

107. Delpy DT, Cope M, Van Der Zee P, Arridge S, Wray S, Wyatt J. Estimation of optical pathlength through tissue from direct time of flight measurement. Physics in Medicine and Biology. 1988;33(12):1433–42.

108. Kocsis L, Herman P, Eke A. The modified Beer-Lambert law revisited. Physics in Medicine and Biology. 2006;51(5):N91–8.

109. Garcia-Castro G. Implementing the Differential Pathlength Factor (DPF) in Python: The Scholkmann Method [Internet]. Available from: https://gongcastro.github.io/blog/dpf-scholkmann/dpf-scholkmann.html

110. Scholkmann F, Wolf M. General equation for the differential pathlength factor of the frontal human head depending on wavelength and age. J Biomed Opt [Internet]. 2013 Oct 11 [cited 2023 Jun 28];18(10):105004. Available from: http://biomedicaloptics.spiedigitallibrary.org/article.aspx?doi=10.1117/1.JBO.18.10.105004

111. Ayrolles A, Brun F, Chen P, Djalovski A, Beauxis Y, Delorme R, et al. HyPyP: a Hyperscanning Python Pipeline for inter-brain connectivity analysis. Social Cognitive and Affective Neuroscience [Internet]. 2021 Jan 18 [cited 2023 Jun 7];16(1–2):72–83. Available from: https://academic.oup.com/scan/article/16/1-2/72/5919711

112. McElreath R. Statistical Rethinking; A Bayesian Course with Examples in R and Stan. 2nd Edition. New York: Chapman and Hall/CRC; 2020. 612 p.

113. Bürkner PC. brms: An R Package for Bayesian Multilevel Models Using Stan. J Stat Soft [Internet]. 2017 [cited 2022 Apr 6];80(1):1–28. Available from: http://www.jstatsoft.org/v80/i01/

114. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2022. Available from: http://www.r-project.org/

115. RStudio Team. RStudio: Integrated Development for R [Internet]. Boston, MA: RStudio, Inc.; 2020. Available from: http://www.rstudio.com/

116. Kruschke JK, Liddell TM. The Bayesian New Statistics: Hypothesis testing, estimation, meta-analysis, and power analysis from a Bayesian perspective. Psychon Bull Rev [Internet]. 2018 Feb [cited 2022 Feb 1];25(1):178–206. Available from: http://link.springer.com/10.3758/s13423-016-1221-4

117. Boukarras S, Placidi V, Rossano F, Era V, Aglioti SM, Candidi M. Interpersonal physiological synchrony during dyadic joint action is increased by task novelty and reduced by social anxiety. Psychophysiology [Internet]. 2025 Mar [cited 2025 Aug 25];62(3):e70031. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11913774/

118. Kenny DA, Kashy DA, Cook WA. Using multilevel modeling to study dyads. In: Dyadic Data Analysis, [Internet]. Guilford Publications; 2006. p. 78–99. Available from: https://ebookcentral.proquest.com/lib/ethz/detail.action?docID=362550