Longitudinal intergenerational hyperscanning reveals indices of relationship formation and loneliness

Ryssa Moffat[1], Guillaume Dumas[2,3], Emily S. Cross[1]

[1] Social Brain Sciences Lab, ETH Zurich, Switzerland
[2] CHU Sainte-Justine Azrieli Research Center, Department of Psychiatry, University of Montreal, Quebec, Canada
[3] Mila – Quebec Artificial Intelligence Institute, Quebec, Canada

* Correspondence:
Ryssa Moffat
rmoffat@ethz.ch

Emily Cross
ecross@ethz.ch

Keywords: interbrain synchrony, fNIRS, intergenerational, older adults, loneliness, community art program

Abstract
Loneliness is globally acknowledged as a severe and burgeoning health risk, fuelling interest in helping people of all ages form meaningful social connections. One promising approach consists of intergenerational social programs. While behavioural and qualitative evidence derived from such programs promise health and wellbeing benefits, the physiological consequences of repeated intergenerational encounters remain unknown. Insight into physiological changes will shed light on the mechanisms of social connection and inform program design choices. We charted changes in interpersonal neural synchrony (INS) in 31 intergenerational (older/younger adult) and 30 same generation (younger adult) dyads across a six-session art program. At each session, dyads completed self-report measures, drew together and alone, and had their cortical activation recorded with fNIRS. In both groups, INS was greater while dyads drew together than alone. Across sessions, intergenerational dyads' INS decreased and same generation dyads' INS increased. Further findings highlight the promise of INS as a biomarker for changes in loneliness and the development of social relationships.

1   Introduction

Experiences of loneliness can have serious negative consequences for physical and mental health. As a result, widespread efforts are being made to combat loneliness (WHO, 2021). According to meta-analyses of behavioural studies, community programs providing opportunities for repeated contact between generations can improve older adults' (65+ years) physical, social, and cognitive health (Petersen, 2023), while also reducing younger adults' stereotypes about aging (Bartlett et al., 2021; Gonzales et al., 2010). Moreover, physicians are increasingly prescribing attendance at such programs–a practice called social prescribing (Scarpetti et al., 2024)–further highlighting the value the medical field attributes to such programs. To date the existing evidence base for intergenerational community programs consists of behavioural and qualitative studies. While offering promising results in terms of self-reported benefits and social engagement, such studies offer little insight into how or why intergenerational relationships develop on a physiological level. In this study, we sought

to provide a physiological perspective on the trajectory of intergenerational relationship development
using a longitudinal hyperscanning approach, comparing intergenerational and same generation
dyads who completed a six-week art program. Specifically, we tracked the trajectories of dyads'
social wellbeing, abilities to collaborate, and levels of interpersonal neural synchrony (INS).

The following sections briefly describe the benefits of intergenerational social interactions, what is
known about INS in intergenerational interactions, and the benefits of longitudinal perspectives on
INS. Following this, we detail the aims and hypotheses of the present study.

1.1 Intergenerational social interactions to combat loneliness

Loneliness can be defined as "a negative, subjective emotional state resulting from a discrepancy
between one's desired and actual experiences of connection" (WHO, 2025). Put more succinctly,
loneliness can be thought of as *perceived* social isolation. Policy makers and researchers have long
emphasised the severe risk that loneliness poses to physical and mental health (House et al., 1988; F.
Wang et al., 2023; WHO, 2021). Public figures and media outlets go so far as to compare the adverse
health impacts (e.g., mortality) of loneliness to those of smoking cigarettes (The Lancet, 2023).
Considering everyday functioning, longitudinal studies show that loneliness and social isolation can
precipitate declines in cognitive functioning including verbal ability, numeracy, and short-term
memory (Ayalon et al., 2016; Cachón-Alonso et al., 2023). Loneliness is also associated with
reduced perceptual speed (Drewelies et al., 2022) and biases toward negative evaluations of social
information (J. T. Cacioppo & Hawkley, 2009; Okruszek et al., 2021; Spithoven et al., 2017). At the
brain level, chronic loneliness and episodes of social isolation have been associated with altered
structure (typically reduction in tissue volume) and altered function of prefrontal and posterior
superior temporal cortices, as well as the insula, hippocampus, and amygdala (S. Cacioppo et al.,
2014; Düzel et al., 2019; Lam et al., 2021). Attentional, visual, and default mode networks are also
reported to show stronger functional connectivity in lonely individuals, which is believed to be
associated with the bias toward processing social information negatively and a tendency toward
increased rumination (Brilliant et al., 2022; Lam et al., 2021; Lieberz et al., 2021).

To combat loneliness, policy makers, researchers (Holt-Lunstad & Perissinotto, 2023; WHO, 2021;
Wigfield et al., 2022), and health practitioners (Scarpetti et al., 2024) promote the creation of
opportunities for meaningful social interactions for all members of society, and especially for those
from populations most susceptible to loneliness. This typically involves initiating community
programs that provide a platform for regular socialising opportunities (Bessaha et al., 2020; Noone &
Yang, 2022; O'Rourke et al., 2018; WHO, 2025). Some community programs are intentionally
designed to bring together members of different age groups, i.e., to be intergenerational. One reason
for this is that people of all ages experience loneliness (Cigna US, 2018; Hawkley et al., 2022;
Qualter et al., 2015). Another is that intergenerational social interactions can improve social and
physical wellbeing for older adults, as well as increase empathy and feelings of social closeness
toward older adults from younger participants (Petersen, 2023). The observed improvements seem to
stem from repeated intergenerational contact (Lee et al., 2020; Lokon et al., 2020; Newman et al.,
2014). In this context, repeated intergenerational contact requires regular engagement in
intergenerational community programs, which is most likely to be achieved by ensuring that
programs offer meaningful social interactions. One approach that has been shown to be particularly
successful in this domain is arts-based and creative activities (WHO, 2019). Indeed, a growing body
of evidence documents the social and physical health benefits of intergenerational arts programs for
participants of all ages (Anderson et al., 2017; Jenkins et al., 2021; Lokon et al., 2020; Rubin et al.,
2015).

Despite the mounting behavioural and qualitative evidence that repeated intergenerational
interactions yield social benefits, including reduced loneliness, little is known about the physiological
aspects that contribute to *how* and *why* these intergenerational programs offer such benefits. This is
important because the physiological signatures of budding intergenerational relationships have the
potential to be harnessed as tools to mitigate loneliness, yet they remain unknown (Moffat et al.,
2024).

1.2   Interpersonal neural synchrony (INS) during collaborative intergenerational interactions

One source of physiological information is brain activity. Recording brain activity from two or more
people while they interact is commonly referred to as 'hyperscanning' and offers a measure of
interpersonal neural synchrony (INS; the extent to which patterns of brain activity are temporally
aligned). Mobile functional near-infrared spectroscopy (fNIRS) and electroencephalography (EEG)
are particularly well suited to recording brain activity during social interactions in real-world settings
(Liu et al., 2018; Moffat et al., 2024). Hyperscanning with non-mobile techniques, such as functional
magnetic resonance imaging (fMRI) and magneto-encephalography (MEG), is possible, though it
offers less realistic social interactions and is less logistically feasible (Hirata et al., 2014; Misaki et
al., 2021).

1.2.1 INS in social brain networks differs between solo and dyadic tasks

In studies comparing INS while dyads perform tasks collaboratively and alone, INS is consistently
greater for collaborative performance than solo performance. This pattern has been observed for
finger tapping (Holper et al., 2012), cognitive tasks such as n-back and object detection tasks (Cui et
al., 2012; Dommer et al., 2012), singing (Osaka et al., 2015), tracing shapes on a screen using
computer buttons or a stylus (Li et al., 2020; Sun et al., 2021), drawing specific verbs as in Pictionary
(Xie et al., 2020), among other activities. A meta-analysis of fNIRS hyperscanning studies by
Czeszumski et al. (2022) reports that during collaboration, INS is commonly observed between both
interactants' prefrontal cortex (PFC), inferior frontal cortex (IFG) or temporoparietal junction (TPJ).
INS in the prefrontal and inferior frontal cortices is proposed to reflect initiation and reception of
joint attention (Koechlin et al., 2003; Redcay et al., 2012; Yoshioka et al., 2021), a social competence
that depends on cognitive control (Ding et al., 2023; Mundy, 2018). INS in the TPJ, a key node of the
Theory of Mind (ToM) network (Jacoby et al., 2016; Saxe & Kanwisher, 2003), is believed to reflect
perspective taking and integration of social information (Czeszumski et al., 2022). Complementary
findings from hyperscanning studies using EEG suggest that the strength and nature of collaboration
shape the levels of INS recorded over frontal and prefrontal brain regions for alpha, beta, and theta
frequency bands (Dikker et al., 2017; Dumas et al., 2010; Labonté-LeMoyne et al., 2016; Sinha et al.,
2016; Toppi et al., 2016). In the context of INS in EEG signals, the alpha band may index joint
attention and visual processing, while beta and theta bands may reflect joint encoding of
sensorimotor information (Dikker et al., 2017; Dumas et al., 2010; Liu et al., 2018). In combination,
the findings from fNIRS and EEG hyperscanning studies indicate that INS levels during
collaboration are shaped by processes such as joint attention and perspective taking.

Many hyperscanning studies on collaboration can be considered intergenerational, in that they
examine collaboration between parent-child/infant or student-teacher dyads (for reviews, see: De
Felice et al., 2023; Nguyen et al., 2020). Though these studies offer the field of social neuroscience a
valuable starting point for understanding intergenerational collaboration, two gaps in the literature
have yet to be filled: First, the hyperscanning literature on collaboration distinctly lacks any
representation of older adults. Second, the hyperscanning literature more generally offers very few

longitudinal perspectives into INS exceeding two time points. We now briefly explore the importance
of addressing these gaps.

1.2.2 Older adults are underrepresented in hyperscanning research

To date, older adults' representation in the hyperscanning literature is limited to one hyperscanning
study (Zhang et al., 2023), one study simulating INS (Dikker et al., 2024), and one position piece
arguing for the inclusion of older adults in such studies (Moffat et al., 2024). It is critical to address
this knowledge gap as the real-world settings in which older adults interact with younger adults and
children (with their children and grandchildren, in assisted living, when seeking medical care, in
community programs or through volunteer work) are numerous. Moreover, the proportion of older
adults in the world's population is projected to continue growing until the end of this century. With
an estimated 25% of populations across a number of global regions, including Europe and North
America, expected to be older than 65 years in 2050 (United Nations, 2024), filling this knowledge
gap will benefit a steadily increasing number of people.

In the one study to date exploring intergenerational INS involving older adults, Dikker et al. (2024)
employed a neurocognitive model to simulate EEG-based INS during intergenerational social
interactions between grandparents, parents, and grandchildren. In this seminal study, the authors

found greater simulated INS between grandparent-grandchild dyads and grandparent-parent-child
triads, relative to grandparent-parent and parent-child dyads. The authors report that the increase in
simulated grandparent-child INS may be driven by age-related changes in the spectrum of brain
activity. The authors also encourage the collection of real-world data to validate these simulations
(Dikker et al., 2024). These findings underscore how including people of all ages, particularly
underrepresented older adults, in intergenerational hyperscanning will enhance our understanding of
INS across a broader range of everyday social interactions across the lifespan.

1.2.3 INS trajectories across days, weeks, and years remain uncharted

Most hyperscanning studies offer snapshots of INS in a single moment in time, and compare INS
between contexts, tasks, or mental states. The *trajectory* in INS over days or weeks remains
unknown, despite the value of granularity insights for understanding changes in behaviour related to
the investigated contexts, task, and mental states. Longitudinal hyperscanning studies typically
include two time points, as opposed to a denser sampling of the trajectory of INS. Numerous reviews
from different domains of social neuroscience (e.g., developmental, clinical, communication etc.)
recognise the lack of longitudinal insights into INS and call for longitudinal hyperscanning studies to
be conducted (e.g., Adel et al., 2025; Bi et al., 2023; De Felice et al., 2025; Hoehl et al., 2025; Kelsen
et al., 2022; Noreika et al., 2020; Turk et al., 2022). The discrepancy between the number of calls for
studies tracking INS trajectories and the paucity of studies reporting them substantiates the critical
need for additional multi-session hyperscanning studies. Concretely, multi-session longitudinal
designs can fill in the blanks in our understanding of INS dynamics. A more fine-grained
understanding of how the shared neural mechanisms underpinning social interactions change over
time may help health practitioners maximize the social or cognitive outcomes of interventions (e.g.,
improving intergenerational communication, mitigating experiences of loneliness, identifying social
difficulties, and general validation of interventions).

To the best of our knowledge, five hyperscanning studies have recorded INS at multiple sessions and
subsequently analysed the aggregated time points, as opposed to reporting changes in INS across
sessions (Bevilacqua et al., 2019; Dikker et al., 2017; Ellingsen et al., 2023; Grahl et al., 2021, 2023).

These studies are reviewed in greater detail in a position piece by Moffat et al. (2024). Only one
fNIRS hyperscanning study offers insights into the trajectory of INS during relationship formation
(Sened et al., 2025). Sened et al.'s (2025) proof-of-concept study tracked the trajectory of INS over
bilateral IFG in 8 clinician-patient dyads at the first, third and fifth sessions of a 6-session
intervention, where dyads completed imagery and cognitive behavioural training procedures. Sened
et al. report increasing INS across sessions, as well as an association between increasing INS and
reduced anxiety symptoms, improved wellbeing and perceived quality of session, but not the
therapeutic alliance (i.e., the patient's rating of quality of the collaborative relationship between
clinician and patient). One further study takes a unique approach to understanding INS in the context
of relationship formation, though only encompassing two time points (D. Wang et al., 2024). INS
was recorded from 11 dyads at two sessions, the first within 5 days of the dyad becoming flatmates,
and the second after 7 months of co-living. Wang et al. report that social closeness can be predicted
from INS with approximately 69% accuracy, using a deep neural network model and EEG
hyperscanning recordings from frontotemporal brain regions. These two studies (Sened et al., 2025;
D. Wang et al., 2024) provide preliminary evidence that INS between dyads increases as dyads grow
more familiar with each other, however a larger sample size and number of time points would
provide a richer picture of INS trajectories in developing relationships.

1.2.4 Cross-sectional research suggests that social closeness modulates INS

Taking a cross-sectional approach to understand social closeness and relationship development, some
studies compare romantic couples, friends and strangers at single moments in time. Such studies
suggest that the nature of a dyad's relationship influences INS during forms of interaction including
joint motor action, conversation, and classroom learning (Bevilacqua et al., 2019; Djalovski et al.,
2021; Long et al., 2022; Speer et al., 2024; Sun et al., 2021; Zhang et al., 2023). Djalovski et al.
(2021) recruited romantic couples, friends and strangers to complete a finger tapping task as dyads,
and a conversation-based empathy-giving task while brain activity was measured using EEG. During
the finger tapping task, INS was greatest between romantic couples, followed by friends, and lowest
between strangers. The authors suggest that cumulative joint experience with a specific partner
enhances prediction of the partners movements, resulting in greater INS. The empathy-giving task
revealed the opposite gradient with romantic couples showing the least INS, friends showing more,
and strangers showing the most INS. Djalovski et al. propose that strangers monitor each other for
potential negative responses more closely than couples to avoid the consequences of a potential
misunderstanding. In another study examining verbal communication, Speer et al. (2024) recruited
friends and strangers to engage in prompted conversations with demarcated turns. The authors report
that across the span of conversation, friends showed decreasing INS and strangers showed increasing
INS, with the reductions in INS corresponding to more exploratory conversations and increases in
INS seeking to converge on common ground. Though phrased differently, Speer et al.'s interpretation
is closely aligned to Djalovski et al.'s interpretation: The less well a dyad knows each other, the
greater the INS during verbal communication. To explore this further, we highlight that a central
difference between the motor and verbal communication tasks is that the motor task (i.e., drawing
specific objects with an 'Etch A Sketch') is highly specified, setting both dyad members'
expectations clearly and symmetrically, whereas the verbal communication tasks (i.e., describing a
troubling or distressing event and conversing with demarcated turns) are temporally asymmetrical
and inherently unpredictable. We propose that the degree of predictability of behaviour induced by
the task contributes to the opposite INS gradients.

Other cross-sectional hyperscanning studies provide evidence that social closeness can shape the
arousal that people experience while interacting, as well as INS. In a conversation study, Long et al.

(2022) recorded brain activity with fNIRS while participants engaged in supportive and conflictual
conversation. The authors observed that romantic couples showed greater INS than control dyads
when engaging in debate. INS was associated with dyads' arousal ratings, suggesting the dynamic
range of arousal in conflict-related conversations with a romantic partner moderates INS. Further,
Zhang et al. (2023) compared INS during a competitive button-pressing task performed by older
romantic couples and older strangers and found that romantic couples were more competitive and
showed greater INS than strangers (note: this is to our knowledge the only hyperscanning study that
includes older adults). These studies suggest that dyads who are socially closer explore more of the
'dynamic range' of arousal.

Considered together, these studies suggest that when dyads made up of strangers meet for the first
time and engage in a creative, unconstrained activity, such as drawing, INS is likely to be low.
Across repeated encounters, INS levels are likely to increase. Drawing on Dikker's model simulating
intergenerational INS, one would expect that generational constellations may impact INS levels and
may interact with the trajectory of INS as dyads become socially closer.

1.3   Present study: Tracking intergenerational INS longitudinally, across repeated
collaborative interactions

In this preregistered study (https://osf.io/hz6tm), we had three aims:

1. To characterise INS in intergenerational dyads made up of one young and one older adult.
2. To chart the trajectory of INS within same generation and intergenerational dyads, as
relationships form across repeated social interactions.
3. To characterise the relationship between INS and differing levels of loneliness, as well as
social closeness, and attitudes toward generations (i.e., one's own vs. other generations).

To achieve these aims, we recruited 31 intergenerational and 30 same generation dyads to complete a
six-session art program. At each session, we collected measures of participants' loneliness and
attitudes toward other generations, then recorded dyads' brain activity using fNIRS while they drew
independently and collaboratively with oil pastels. We calculated INS using wavelet transform
coherence (WTC) for all pairs of four regions of interest (ROIs), which were the bilateral IFG and
TPJ. According to fNIRS best practices, we report both fNIRS signals (oxygenated haemoglobin
[HbO] and deoxygenated haemoglobin [HbR]), with HbO results presented in the main manuscript
and HbR results presented in the Supplementary Materials. We employed a Bayesian statistical
analysis approach, according to which we make inferences based on 95% confidence intervals called
95% HPDs (highest posterior density region) rather than p-values. See Methods and Supplementary
Materials for additional details including complete tables with estimates and 95% HPD, which we
present separately from results text for ease of reading.

Our findings pertaining to INS in intergenerational and same generation dyads, tracked across 6
sessions in a real-world context resembling a community art program, deliver first insights into INS
involving older adults, multi-session trajectories of INS, as well as INS' potential as a biomarker for
tracking changes in loneliness and social closeness in budding relationships.

2   Results

2.1   Self-reported loneliness, social closeness, and attitudes toward generations

We first investigated how self-reported loneliness, social closeness within dyads, as well as attitudes
towards generations differ between intergenerational and same generation dyads (Figure 1; Table 1).

Loneliness. Intergenerational dyads reported feeling substantially less lonely than same generation
dyads. Feelings of loneliness decreased substantially for both groups across sessions, with no
difference in rate. **Social closeness (OIS).** Intergenerational dyads reported substantially greater
feelings of social closeness, compared to same generation dyads. Feelings of social closeness
increased substantially across sessions for both groups, with no difference in rate. **Attitudes toward**
generations (allophilia). Same generation dyads reported feeling substantially closer to their own
generation than the intergenerational dyads, who reported feeling equally close to their own and the
other generation. Allophilia scores stayed stable across sessions (i.e., no change observed).

Relationships between measures. Greater social closeness was substantially associated with
reduced feelings of loneliness for same generation dyads and a matching trend was observed for
intergenerational dyads. These associations did not differ between groups. Greater social closeness
was substantially associated with more positive attitudes toward other generations for intergeneration
dyads but not same generation dyads. This association showed a trending difference between groups.
Greater loneliness was substantially associated with more positive attitudes toward other generations
for same generation dyads but not intergenerational dyads. This association did not differ between
groups.

Figure 1. Top: Distributions of self-reported loneliness, social closeness (OIS), and attitudes toward generations
(allophilia) of individual participants. Middle: Distribution of summed scores per dyad. Bottom: Distribution of
difference scores per dyads.

Table 1. Unstandardised estimates computed per self-reported measure with 95% HPD in square brackets. Point
estimates for all six sessions combined are presented first, followed by the estimates of the slope of change across
sessions in the lower section. Self-report measures are loneliness, social closeness (OIS), and attitudes toward generations
(allophilia). Visualisation by age (i.e., older or younger adult) and group in Supplementary Figure 1.

	Loneliness	Social Closeness	Allophilia
Sessions combined			
Intergenerational	1.28 [1.02, 1.54]	4.08 [3.76, 4.39]	-1.19 [-3.45, 1.05]
Same gen	1.89 [1.62, 2.16]	3.52 [3.19, 3.38]	
Intergen-Same gen	-0.61 [-0.99, -0.25]	0.56 [0.10, 1.00]	4.48 [1.29, 7.56]
Change across six sessions			
Intergenerational	-0.06 [-0.10, -0.02]	0.24 [0.19, 0.27]	-0.21 [-0.55, 0.12]
Same gen	-0.07 [-.11, - 0.03]	0.25 [0.20, 0.30]	-0.02 [-0.37, 0.30]
Intergen-Same gen	0.01 [-0.05, 0.07]	-0.01 [-0.08, 0.07]	-0.18 [-0.66, 0.28]

286 **Table 2.** Unstandardised estimates of relationship between self-reported measures per group with 95% HPD in square
287 brackets. Self-report measures are loneliness, social closeness (OIS), and attitudes toward generations (allophilia).

	Loneliness	Social Closeness
Social Closeness (Intergen)	-0.13 [-0.27, 0.01]	
Social Closeness (Samegen)	-0.16 [-0.28, -0.04]	
Allophilia (Intergen)	0.01 [-0.00, 0.02]	0.02 [0.01, 0.04]
Allophilia (Samegen)	0.02 [0.01, 0.03]	0.01 [-0.01, 0.2]

2.2 INS

2.2.1 Real dyads vs. pseudo dyads

The first step in our analysis of INS was to perform contrasts between real and pseudo dyads for INS
estimates reflecting all sessions combined and those reflecting change in INS across sessions per
combination of ROI pair, task, and group. Contrasts serve to assess whether the level of observed
INS (or the slope of change in INS) differs meaningfully from a null distribution of INS levels (or
slopes) made of dyads who never interacted with one another. Further statistical analyses can then be
performed on the ROI pairs where real dyads INS differed substantially from pseudo dyads' INS. We
report estimates and contrasts with 95% HPD in Supplementary Tables 1-4.

INS levels for all sessions combined. We found that all substantial differences corresponded to
greater INS for real than pseudo dyads (black squares in Figure 2A; solid lines in Figure 2B;
Supplementary Table 2). The ROI pairs showing substantial differences were RIFG~RIFG for
intergenerational dyads drawing alone, LIFG~LIFG, LTPJ~LTPJ, RIFG~RIFG, LIFG~RIFG,
LIFG~LTPJ, RIFG~LTPJ for intergenerational dyads drawing together, no ROI pairs for same-
generation dyads drawing alone, and LIFG~RIFG and RIFG~RTPJ for same-generation dyads
drawing together. HbR analyses were similar, in that LTPJ~LTPJ, LIFG~RIFG, and LIFG~LTPJ
showed substantial differences between real and pseudo dyads for intergenerational dyads drawing
together (reported in full in Supplementary Materials).

Change in INS levels across sessions. We observed a more positive slope of change for all ROI pairs that differed substantially between real and pseudo dyads, with one exception (black squares Figure 3A; solid lines in Figure 3B; Supplementary Table 4). The ROI pairs showing substantial differences were LIFG~RTPJ for intergenerational dyads drawing alone, LTPJ~LPTJ and RIFG~RTPJ for intergenerational dyads drawing together, no ROI pairs for same-generation dyads drawing alone, and RTPJ~RTPJ and LIFG~LTPJ for same-generation dyads drawing together. HbR analyses showed little similarity (reported in full in Supplementary Materials).

Figure 2. A) INS for all ROI pairs per task and group; black squares indicate that INS levels differ between real and pseudo dyads substantially (i.e., 95% HPD do not overlap). B) Visualisation of all possible ROI pairs (n=10). Dashed lines indicate ROI pairs where INS did not differ substantially between real and pseudo dyads. Solid lines indicate ROI pairs where INS differed substantially between real and pseudo dyads. Black solid lines indicate ROI pairs where task, group, and task*group contrasts showed substantial differences. Grey solid lines indicate ROI pairs where contrasts showed no substantial differences. C) INS levels with 95% HPD for all ROI pairs that differed between real and pseudo dyads, for which task, group, and task*group contrasts were computed. LIFG~LIFG: No differences. RIFG~RIFG: No differences. LTPJ~LTPJ: Task difference, drawing together > alone. LIFG~RIFG: Task difference, drawing together > alone. Interaction, drawing together > alone for intergenerational dyads. LIFG~LTPJ: No differences. RIFG~LTPJ: Task difference, drawing together > alone. RIFG~RTPJ: Task difference, drawing together > alone. Interaction, drawing together > alone for same generation dyads. Full list of estimates and contrasts with 95% HPD available in Supplementary Tables 1-2 for HbO. HbR analyses visualised in Supplementary Figure 2, showed the same pattern of results for LIFG~RIFG and RIFG~LTPJ.

Figure 3. A) Slope of change in INS across six sessions; black squares indicate that INS levels differ between real and pseudo dyads substantially (i.e., 95% HPD do not overlap). B) Visualisation of all possible ROI pairs (n=10). Dashed lines indicate ROI pairs where the slope of change in INS did not differ substantially between real and pseudo dyads. Solid lines indicate ROI pairs where the slope of change in INS differed substantially between real and pseudo dyads. Black solid lines indicate ROI pairs where task, group, and task*group contrasts showed substantial differences. C) Slope of change in INS across sessions for all ROI pairs that differed between real and pseudo dyads, for which task, group, and task*group contrasts were computed. Solid lines indicate INS~Session relationship was substantial (i.e., 95% HPD does not overlap with 0). Dashed lines indicate non-substantial INS~Session relationships. Grey shading shows 95%, 89%, and 50% intervals of the posterior predictive distributions. LTPJ~LTPJ: Interaction effect, drawing together > alone for intergenerational dyads. RTPJ~RTPJ: Interaction effects, drawing together < alone for intergenerational dyads and intergenerational < same generation dyads for drawing together. LIFG~LTPJ: Group effect, intergenerational < same generation dyads. LIFG~RTPJ: Interaction effects, drawing together < alone for intergenerational dyads, intergenerational > same generation dyads for drawing alone, and intergenerational < same generation dyads for drawing together. RIFG~RTPJ: Interaction effects, drawing together < alone for intergenerational dyads, intergenerational > same generation dyads for drawing alone, and intergenerational < same generation dyads for drawing together. Full list of estimates and contrasts with HPD available in Supplementary Tables 3-4 for HbO. HbR analyses, visualised in Supplementary Figure 3, showed the same pattern of results for RTPJ~RTPJ and the same interaction effect of drawing together < alone for intergenerational dyads.

2.2.2 Composition of non-homologous ROI pairs in intergenerational dyads

For the intergenerational dyads specifically, we examined whether INS differed as a function of the generational composition of the ROI pairs. In other words, whether an ROI belonging to an older or younger member of a dyad influenced INS levels for all sessions or slopes across sessions. Full list of estimates and contrasts with HPD available in Supplementary Table 5.

INS levels for all sessions combined. When intergenerational dyads drew together, there was a trend of greater INS when the LIFG belonged to the older adult and the LTPJ belonged to the younger adult than vice versa. We observed no substantial differences in INS for any composition of any ROI pair for drawing alone. **Change in INS levels across sessions.** No substantial differences in the INS slopes across sessions for any composition of any ROI pair, for drawing alone or together.

2.2.3 Contrasting drawing conditions and groups

INS levels for all sessions combined. We observed no substantial differences between groups (i.e. group effect) but did observe task effects. We observed substantially greater INS for drawing together than alone in LTPJ~LTPJ, LIFG~RIFG, RIFG~LTPJ and RIFG~RTPJ (i.e., task effect). With respect to interaction between group and task, we found that in LTPJ~LTPJ, INS was substantially greater for drawing together than alone for intergenerational dyads. In RIFG~RTPJ, INS was substantially greater for drawing together than alone for same generation dyads. Parameter estimates per ROI pair, task and group visualised in Figure 2C. Full list of contrasts with 95% HPD supplied in Supplementary Table 6.

Change in INS levels across sessions. We observed no substantial difference between drawing alone and together (i.e., no task effects) but did observe the following group effect: We observed a substantially more positive INS slope for the same generation than the intergenerational dyads in LIFG~LTPJ. We also found several interactions effects. Intergenerational dyads showed a substantially more positive INS slope for drawing together than alone in LTPJ~LPTJ, and substantially more positive INS slopes for drawing alone than together in RTPJ~RTPJ, LIFG~RTPJ, and RIFG~RTPJ. This difference was substantial for drawing together, whereby the INS slope was more positive for same-generation dyads, relative to intergenerational dyads. Same-generation dyads further showed substantially more positive INS slopes than intergenerational dyads for drawing together in RTPJ~RTPJ, LIFG~RTPJ and RIFG~RTPJ. Intergenerational dyads showed substantially more positive INS slopes than same generation dyads when drawing alone in LIFG~RTPJ and RIFG~RTPJ. Parameter estimates per ROI pair, task and group are visualised in Figure 3C. Full list of contrasts with 95% HPD supplied in Supplementary Table 7.

2.2.4 Relationship between INS and self-report measures

Loneliness. *Summed.* We observed no substantial main, task, or group effect on the relationship between INS and summed loneliness scores. We observed an interaction effect in RIFG~RTPJ, for the intergenerational dyads, whereby the INS~summed loneliness relationship was substantially more negative for drawing alone than together (Figure 4; Supplementary Tables 10-11). *Difference.* No substantial main effect or group effects were observed. We observed a substantial effect of task for RIFG~RIFG, whereby the INS~loneliness difference relationship was substantially more negative for drawing together than alone. We observed substantial interaction effects for RIFG~RIFG, RIFG~RTPJ, and LTPJ~LTPJ. For RIFG~RIFG, we observed a substantial task difference for the intergenerational dyads only. For LTPJ~LPTJ, we observed a substantial task difference for intergenerational, but not same generation dyads and a substantial difference between groups for drawing alone but not together. For RIFG~RTPJ, we observed a substantial task difference for same generation dyads only (Figure 4; Supplementary Tables 12-13).

Social closeness (OIS). *Summed.* We observed a substantial positive relationship between summed social closeness scores and INS in RIFG~RTPJ (main effect). In LIFG~LIFG, we observed a substantial effect of task, whereby there was a substantially more positive relationship between INS and summed social closeness for drawing together than alone. We observed a group effect for RIFG~RIFG, whereby intergenerational dyads showed a substantially more positive relationship than same generation dyads, and RIFG~RTPJ where intergenerational dyads showed a substantially more negative relationship than same generation dyads. Substantial interaction effects observed for LIFG~LIFG and RIFG~RIFG, RIFG~RTPJ. In LIFG~LIFG there was a substantial task difference for same generation dyads only. In RIFG~RIFG the interaction effect showed a substantial group difference for drawing alone only. In RIFG~RTPJ, we observed a substantial task difference for the

same generation dyads only and a substantial group difference only when dyads drew alone (Figure 4; Supplementary Tables 14-15). *Difference.* No substantial main, task, or group effect on the relationship between INS and social closeness difference scores was observed. We did find a substantial interaction effect for RIFG~RIFG: There was a substantial task difference for both groups and a substantial group difference when drawing alone only (Figure 4; Supplementary Tables 16-17).

Allophilia. *Summed*. We observed a substantial negative relationship between summed allophilia scores and INS in LTPJ~LPTJ and LIFG~RIFG (main effect). We also found a substantial group effect for RIFG~RTPJ, where the relationship between INS and the summed allophilia scores was substantially more positive for the intergenerational dyads than same generation dyads. A substantial interaction effect was observed in LIFG~LTPJ, with a substantial task difference for intergenerational dyads only (Figure 4; Supplementary Table 18-19). *Difference*. No substantial main or group effects were observed. We observed a substantial task effect in RIFG~RIFG, where the relationship between INS and the difference in allophilia scores was substantially more positive for drawing together than alone. We found a substantial interaction effect for RIFG~RIFG, where we observed a substantial task difference for same generation dyads only (Figure 4; Supplementary Tables 20-21).

Figure 4. A) Slope of change in INS across levels of self-report measures for ROI pairs with at least one substantial
relationship or substantial difference revealed by contrasts (for all ROI pairs see Supplementary Figures 5A-F; estimates
and contrasts with 95 % HPD in Supplementary Tables 10-21). Lines indicate whether INS~Self-report measure
relationship was substantial (solid) or non-substantial (dashed). Shading shows 95%, 89%, and 50% intervals for the
posterior predictive distributions. Results of contrasts between task, group, and task*group contrasts for the panels above
are as follows. *LIFG~LIFG Social closeness (Sum):* Task effect, negative relationship for drawing alone and drawing
alone < together. Interaction effect, together > alone for same generation. *RIFG~RIFG Loneliness (Dif):* Task effect,
drawing together < alone. *RIFG~RIFG Social Closeness (Sum):* Group effect, intergenerational > same generation dyads.

Interaction effect, intergenerational > same generation for drawing alone. *RIFG~RIFG Social Closeness (Dif):* Interaction
effects, drawing together > alone for intergenerational dyads, drawing together < alone for same generation dyads, and
intergenerational > same generation for drawing alone. *RIFG~RIFG Allophilia (Dif):* Task effect, positive relationship for
drawing together and drawing together > alone. Interaction effect, drawing together > alone for same generation dyads.
LTPJ~LTPJ Loneliness (Dif): Interaction effect, intergenerational < same generation dyads for drawing alone.
LTPJ~LTPJ Allophilia (Sum): Main effect, negative relationship. Group effect, negative relationship for intergenerational
dyads only. Task effect, negative relationship for drawing alone and together. *LIFG~RIFG Allophilia (Sum):* Main effect,
negative relationship. Group effect, negative relationship for same generation dyads only. *LIFG~LTPJ Allophilia (Sum):*
Interaction effect, drawing together < alone for intergenerational dyads. *RIFG~RTPJ Loneliness (Sum):* No differences.
RIFG~RTPJ Loneliness (Dif): Interaction effect, drawing together < alone for same generation dyads only. *RIFG~RTPJ*
Social Closeness (Sum): Main effect, positive relationship. Group effect, positive relationship for same generation dyads
and intergenerational < same generation. Task effect, positive relationship for drawing alone. Interaction effects, drawing
together < alone for same generation dyads, and intergenerational < same generation for drawing alone. *RIFG~RTPJ*
Allophilia (Sum): Group effect, intergenerational > same generation dyads. B) Visualisation of all possible ROI pairs
(n=10). Dashed lines indicate ROI pairs where i) neither INS for all sessions combines nor ii) the slope of change in INS
differed substantially between real and pseudo dyads. Solid lines indicate ROI pairs where INS (for sessions combined or
slope of change across sessions) differed substantially between real and pseudo dyads. Black solid lines indicate ROI
pairs where task, group, and task*group contrasts showed substantial differences. Analyses of HbR, visualised in
Supplementary Figure 4, showed similar pattern of results for Social Closeness (Sum) and Allophilia (Dif) in
RIFG~RIFG, as well as Loneliness (Dif) in LTPJ~LTPJ.

3   Discussion

We invited intergenerational and same generation dyads to complete a six-sessions, resembling a
community art program, and charted dyads' levels of loneliness, social closeness, attitudes toward
generations and INS as they became acquainted across 6 sessions. Our first aim was to characterise
INS within intergenerational dyads comprising one young and one older adult. In a large sample
(sessions combined), we show that INS among intergenerational dyads is largely comparable to same
generation dyads. Specifically, both dyad types show greater INS in LTPJ~LTPJ, LIFG~RIFG,
RIFG~LTPJ, and RIFG~RTPJ for drawing together than alone, with the greatest differences in
LIFG~RIFG for intergeneration dyads and RIFG~RTPJ for same generation dyads. Our second aim
was to chart the trajectory of INS in same generation and intergenerational dyads, across repeated
social interactions. While drawing together, ROI pairs that include the right TPJ (RTPJ~RTPJ,
LIFG~RTPJ, and RIFG~RTPJ) showed decreasing INS for intergenerational dyads and increasing
INS for same generation dyads. Our third aim was to characterise the relationship between INS and
differing levels of loneliness, social closeness, and attitudes toward generations. We found that the
relationship between loneliness and INS in RIFG~RIFG differed when dyads drew alone and
together: Greater similarity in dyadic levels of loneliness predicted greater INS when dyads drew
together and lower INS when they drew alone. Greater social closeness predicted increasing INS in
RIFG~RTPJ for all dyads and greater INS in RIFG~RIFG for intergenerational than same generation
dyads. With respect to dyadic attitudes towards generations, we observed that INS in LIFG~RIFG
and LTPJ~LTPJ was elevated among dyads who felt further from other generations (i.e., closer to
their own). Below, we explore the implications of these findings.

Individual's levels of loneliness and social closeness decreased and increased, respectively, across
sessions, for both generational constellations. These findings match the target outcomes of the
community programs proposed by policy makers, researchers, and health practitioners (Holt-Lunstad
& Perissinotto, 2023; Scarpetti et al., 2024; WHO, 2021; Wigfield et al., 2022).

Considering all sessions together, we observed that the generational constellation (i.e.,
intergenerational vs. same generation) predicted differences in individual's levels of loneliness, social
closeness and attitudes toward other generations. For example, simply being in the presence of a
similar or quite differently aged person can alter one's perceptions of one's loneliness. If the other

person is of a similar age, one might feel more lonely than if the person is older, in which case one would self-report feeling less lonely – perhaps by virtue of social comparison (Arnold et al., 2021). This finding should be validated in future work by comparing intergenerational dyads with same generation dyads composed of two older adults, as well as two younger adults. Greater feelings of social closeness among intergenerational, as compared to same generation, dyads may relate to overcoming perceived generational differences or generational stereotypes (Galinsky et al., 2005). Notably, the distributions of difference in social closeness did not differ between groups (Figure 1), suggesting that the increased social closeness observed for intergenerational dyads was not driven by older or younger adults' responses alone, but rather by both age groups. Intergenerational dyads held more charitable attitudes toward other generations than same generation dyads (Figure 1), aligning with findings that contact with older adults reduces younger adults' negative stereotypes of and anxiety regarding older adults (Liao et al., 2023). Among intergenerational dyads, individuals' attitudes toward generations were linked to their social closeness ratings, highlighting possible bidirectional interplay between attitudes and social closeness. Moreover, among same generation dyads, greater loneliness was associated with a weaker preference towards one's 'own' (younger) generation, highlighting the window of opportunity for intergenerational social activities in combatting loneliness.

This study offers novel insights into INS during collaboration in dyads composed of older and younger adults, adding rich empirical data to the two existing studies that involve older adults (Dikker et al., 2024; Zhang et al., 2023). The fNIRS hyperscanning literature on collaboration focuses on the prefrontal and inferior frontal cortices and the TPJ (Czeszumski et al., 2022). The bilateral IFG play a role in the initiation and reception of joint attention (Koechlin et al., 2003; Redcay et al., 2012; Yoshioka et al., 2021) and the bilateral TPJ are central to the integration of social information (Jacoby et al., 2016; Saxe & Kanwisher, 2003). Moreover, with some exceptions (e.g., Zhang et al., 2023), hyperscanning studies typically only report homologous pairs of brain areas (e.g., LIFG~LIFG), for which the interpretation of heightened INS may be simpler. We preregistered and now report analyses of INS that include non-homologous ROI pairs (e.g., LIFG~RIFG) and offer possible reasoning for observations of heightened INS.

In our large sample (all sessions combined), we show that INS in intergenerational dyads is largely comparable to same generation dyads. That is, we observed greater INS in LTPJ~LTPJ, LIFG~RIFG, RIFG~LTPJ, and RIFG~RTPJ for drawing together than alone. This aligns with the existing findings that collaboration co-occurs with greater INS in brain regions that process social information. We interpret these findings as evidence that collaborative drawing results in heightened integration of social information (LTPJ~LTPJ), joint attention (LIFG~RIFG), and perhaps integration of the other person's maintenance of joint attention as social information (RIFG~LTPJ and RIFG~RTPJ). We also observed robust differences in INS while drawing together and alone in separate ROI pairs for each group: LIFG~RIFG for the intergenerational dyads and RIFG~RTPJ for the same generation dyads. We cautiously speculate that intergenerational dyads' level of cognitive control while drawing together is particularly high and aligned within dyads, and that same generation dyads may evaluate and integrate information regarding the strength or continuity of their partner's joint attention. This interpretation requires further investigation using paradigms that can quantify joint attention and cognitive control.

INS in IFG is proposed by some researchers to be unique to verbal communication, while a few studies show that INS in IFG may also play a role in non-verbal communication (Czeszumski et al., 2022). With our very large sample, we can provide strong evidence that INS involving the IFG occurs during non-verbal communication, such as silent collaborative drawing.

Cross-sectional studies suggest that INS levels during collaboration decrease with higher levels of familiarity (such as friendship or stable romantic relationships) that require repeated social interactions to establish (e.g., (Bevilacqua et al., 2019; Djalovski et al., 2021; Long et al., 2022; Speer et al., 2024; Sun et al., 2021; Zhang et al., 2023). Sened et al.'s (2025) proof-of-concept study reports the INS trajectories of eight dyads at three (in a series of six) therapeutic sessions. Sened et al. find increasing INS in bilateral IFG across the three sessions, while clinician-patient dyads completed imagery and cognitive behavioural training procedures. We document INS trajectories during collaboration and independent drawing by two (initial) strangers with the highest temporal granularity to date – 6 consecutive time points. Our findings provide clear evidence of INS changes from a dyad's first encounter to the final recorded session, entailing ~6 hours of time spent together and ~90 minutes of collaborative activity spread across 6 weeks.

We observed that while dyads drew together, ROI pairs that include right TPJ (RTPJ~RTPJ, LIFG~RTPJ, and RIFG~RTPJ) showed decreasing INS among intergenerational dyads but increasing INS among same generation dyads. This may suggest that, in the context of collaborative drawing, intergenerational dyads' monitoring and integration of their partner's levels of joint attention may decrease with increasing familiarity, whereas same generation dyads may monitor each other's levels of joint attention more closely as familiarity increases. Across sessions, INS increased in RTPJ~RTPJ for same generation dyads and LTPJ~LTPJ for intergenerational dyads. This difference in lateralisation may be related to the integration of "matched" or "mismatched" predictions. Whereas the right TPJ is associated with identifying "mismatched" predictions, the left TPJ encodes both "matched" and "mismatched" predictions (Doricchi et al., 2022). Accordingly, it is possible that same generation dyads become increasingly sensitive to "mismatched" predictions with increasing familiarity while intergenerational dyads may encode more "matched" predictions with increasing familiarity. Alternatively for intergenerational dyads, the frequency or absolute amount of mismatched predictions may decrease with increasing familiarity.

We found that the relationship between loneliness and INS in RIFG~RIFG differed when dyads drew alone and together: Greater similarity in dyadic levels of loneliness predicted greater INS when dyads drew together and lower INS when they drew alone. This may point to how differences in loneliness can shape joint attention. Experiences of loneliness predict changes in cognition, such as perceptual speed, memory, and negatively biased perceptions of social information (Ayalon et al., 2016; Cachón-Alonso et al., 2023; J. T. Cacioppo & Hawkley, 2009; Okruszek et al., 2021; Spithoven et al., 2017). It follows that these changes in cognition may co-occur with changes in joint attention or may alter the facility with which individuals engage in joint attention (Corbin & Dhand, 2024). It is possible that lonelier individuals may require more numerous neural resources to maintain joint attention or alternatively present hypervigilant tendencies, both of which could result in elevated INS in RIFG~RIFG (Brilliant et al., 2022; Hampshire et al., 2010; Saporta et al., 2022). Further investigation should establish the extent to which this differential pattern of INS in RIFG~RIFG during collaborative and independent tasks can be used as a biomarker of loneliness in research and real community programs aiming to combat loneliness (Ahkter-Kahn & Au, 2020; Pan & Cheng, 2020).

To understand INS beyond the changes in familiarity incurred by repeated social interactions, we examined the extent to which dyads' self-reported feelings of social closeness were reflected in INS. We observed that greater reported social closeness predicted greater INS in RIFG~RTPJ for all dyads and in RIFG~RIFG for the intergenerational dyads. We interpret the increasing INS in RIFG~RTPJ as meaning that dyads who feel close to one another monitor and integrate each other's joint attention more than dyads who feel less socially close. The increasing INS for intergenerational dyads in

RIFG~RIFG may reflect joint attention as well as heightened arousal or the facilitation of joint
attention via heightened arousal (Bast et al., 2025; Lachat et al., 2012; Long et al., 2022; Zhang et al.,
2023). INS may be useful as a biomarker for changes in social closeness across repeated social
interactions for all adults of all ages, though potentially more robust for same generation dyads.

With respect to dyadic attitudes towards generations, we observed that INS in LIFG~RIFG and
LTPJ~LTPJ decreased as feelings of closeness to other generations increased. As this analysis is
particularly exploratory in nature, with no empirical precedent, we offer a tentative suggestion that
similarity in attitudes towards generations may give dyads a more internally consistent basis upon
which they form predictions about each other's behaviour (i.e., elsewhere termed "common ground";
Speer et al., 2024). These findings can serve as a point of departure for research examining the
impact of different sources of 'common ground' on INS. For example, studies could explore whether
cumulative life experience (e.g., number of years lived) gives older adults an advantage for finding
common ground and/or whether similarities in socialisation (e.g., culture, age, socio-economic status,
hobbies) play bigger role in establishing common ground that could shape INS.

As recommended by Simons et al. (2017), we make explicit the constraints on generality of this
work. Our findings only generalise to the targeted brain regions (i.e., bilateral IFG and TPJ). Though
our study offers typical coverage for fNIRS hyperscanning studies, we encourage the use of high-
density fNIRS optode arrays in future research on this topic (Shuo et al., 2025). We observed overlap
between HbO and HbR results for analyses of all sessions together and INS trajectories across
sessions, with some overlap in HbO and HbR results for relationships between INS and self-reported
measures. Our adequate pairing of sample size and conservative Bayesian approach notwithstanding,
we encourage other researchers to replicate these findings. Moreover, the INS self-report measures
are not necessarily generalisable across cultures, due to differences in formation of relationships
(Goodwin, 2013; Lu et al., 2021). Though our findings pertaining to changes in loneliness and social
closeness across sessions are in line with findings from behavioural and qualitative studies performed
as part of more elaborate and enriching community art programs, our sessions were attended by one
dyad at a time, with partners assigned based on availability. Future studies could expand our
paradigm to small group sessions with multiple dyads and free partner choice to reflect community
programs more closely. We have no reason to believe that the results depend on other characteristics
of the participants, materials, or context.

One limitation of this study is that we did not have the resources to include a group of same
generation dyads composed of two older adults, though this would have been the optimal design.
Without such a group, we cannot be sure that differences between groups are purely related to the
generational constellation and not age-related differences in neural processing. Future research
should compare same generation dyads across different age ranges. A second limitation of this study
is that three optodes of one Photon Cap device unexpectedly became defective partway through data
collection (and were replaced as rapidly as possible). This resulted in the exclusion of several
participants' right TPJ channels for the affected sessions. Thus, the number of data points for ROI
pairs involving right TPJ, particularly RTPJ~RPTJ, is lower than for other ROI pairs (Supplementary
Figure 7). Our Bayesian approach accounted for this difference in sample size per ROI pair by partial
pooling (for an excellent tutorial, see: Nalborczyk et al., 2019).

4   Conclusion

The study met its overarching aim of developing the evidence base for intergenerational community
programs aiming to combat loneliness. Specifically, we documented INS in intergenerational dyads

including older adults, charted the trajectory of INS from first meeting to greater levels of familiarity while intergenerational and same generational dyads engaged in collaborative and creative activity, thus demonstrating the promise of INS as a biomarker for predicting changes in loneliness.

5   Methods

5.1   Participants

We recruited 122 community-dwelling participants from Zurich, Switzerland. Of these, 31 were older adults (aged 69+ years) and 91 were younger adults (aged 18–35 years). We assigned participants to intergenerational dyads (n = 31) and same generation dyads (n = 30) based on availability for sessions (e.g., matching people available at the same day/time for 6 consecutive weeks). Due to the logistical challenge posed by recruiting for and scheduling a multi-session study of this scale, gender pairings were not prioritised (see Supplementary Figures 8 and 9 for summary of gender pairings and session spacing). Intergenerational dyads consisted of an older adult (76 ± 4 years); 18 female, 13 male) with a younger adult (24 ± 4 years; 19 female, 11 male, 1 other). Same generation dyads consisted of two younger adults (22.5 ± 3.6 years; 36 female, 24 male). All dyads started the study as strangers. To be included, participants had no known history of neurological or psychological disorders (e.g., stroke, concussion, ADHD, autism, schizophrenia, depression) and were fluent in German.

We recruited participants via the University of Zurich's Healthy Longevity Centre, ETH Zurich's DESCIL student pool, clubs and organizations for senior citizens (e.g., theatre groups and choirs), and social media. We obtained written informed consent from all participants. Ethical approval was obtained from the Ethics Committee of the Canton Zurich (Ref: 2023-01073). The study was conducted according to the Declaration of Helsinki.

5.2   Measures

5.2.1 Self-report measures

Loneliness. Participants completed the *6-Item Scale for Overall, Emotional, and Social Loneliness* (De Jong Gierveld & van Tilburg, 2006) at the beginning of each session.

Social closeness. Participants indicated how socially close they felt to the other member of their dyad using the *Inclusion of Other in the Self Scale* (Aron et al., 1992) at the end of each session. This measure uses Venn diagrams with incremental increases from no overlap to complete overlap as visual analogue representation of social relationships.

Attitudes toward other generations. The *Allophilia Scale* (Pittinsky et al., 2011) is typically used to characterise participants' attitudes toward an outgroup (e.g., people of another ethnicity or political leaning). We adapted this scale in three variants: a) "younger people", "older people", and "people my age". At the beginning of each session all participants completed the scale with "people my age". Then older adults completed the scale with "younger people" and younger adults completed the scale with "older people". To obtain baseline-corrected attitudes towards other generations, we subtracted each participant's score for "people my age" from their score for the age group they did not belong to.

5.2.2 fNIRS recordings

We recorded fNIRS signals using two Cortivision Photons Cap (Cortivision sp. z o.o., Lublin, Poland) and Cortiview software. Each device had 16 LED sources, which emitted wavelengths of 760 and 850 nm, and 10 detectors (26 optodes in total). The sample rate was 4-6 Hz. We positioned optodes according to the 10-5 EEG positions (Oostenveld & Praamstra, 2001) over bilateral IFG and TPJ based on MNI-space coordinates derived from the Neurosynth database (https://neurosynth.org/). Specifically, we used the search terms 'ifg' and 'tpj' and recorded the coordinates of the area with the highest activation in each the left and right hemisphere, defining the 'centre' of our ROIs. There were the following: *left IFG:* -48, 22, 8; *right IFG:* 50, 22, 16; *left TPJ:* -56, -54, 20; and *right TPJ:* 56, -52, 18. See Supplementary Figure 10 for visualisation of ROI centre coordinates and Supplementary Figure 11 for optode arrangement.

Digitisation of optodes. After placement of the cap according to fiducial markers and before attaching optodes, we recorded a 3D image of each participant's head using Structure Sensor Pro (model SA39) attached to an Apple iPad Pro (v. 10). The 3D head scan was processed using MATLAB (v. 2023b) and the Fieldtrip toolbox (Oostenveld et al., 2011; Appendix 4). Labels for all detectors and sensors and fiducial reference points were assigned to the 3D head scan and then co-registered to MNI space. Next, we obtained the interoptode distance per channel, the coordinates of the point of measurement between optodes making up each channel, as well as distance from the point of measurement to the centre of the ROI. See Supplementary Figure 12 for visualisation of optode distances from ROI centres.

5.3   Procedure

Dyads attended six sessions, with approximately one week between sessions (Supplementary Figure 9 shows schedule of sessions and number of days between sessions per dyad). Sessions were led by a female lead-researcher who provided instructions following a script (available on OSF: https://osf.io/zms2v) and answered participants' technical and scientific queries. A second female researcher was present to aid with technical steps and did not speak unless directly addressed. Both researchers politely deflected participants' social inquiries.

The order of events at each session is visualised in Figure 5A. At the beginning of each session, dyads completed self-report measures of loneliness and attitudes towards generations (allophilia) on an iPad, with a felt divider between participants providing privacy. Then the fNIRS caps were placed on participants' heads and the second researcher recorded 3D models of each participant's head with cap on (participants instructed to sit very still while researcher circled participant with 3D scanner). Next, the lead researcher reminded that participants could converse at any time except while drawing. The researchers attached optodes to the caps, ensuring that optodes came in direct contact with the scalp by clearing hair. Then fNIRS devices were turned on, paired with recording laptops (n = 2) and fNIRS signals were verified for best possible calibration. Video recording was initiated (one GoPro11 Black was positioned with a direct view of participants as shown in Figure 5B, to allow for future motion tracking analyses).

Next, participants drew three drawings with oil pastels, without talking. Participants were instructed that they would have 5 min to draw anything they liked, and that they would complete the first drawing alone and the second and third together on a single piece of paper. No further instructions were provided (e.g., no suggestions for motifs). While participants drew alone, they were separated by a felt divider, ensuring that they could not see each other. The divider was removed when they drew together. In cases where participants conversed to plan their drawings, the researchers waited

for a natural pause to begin the 5 min timer. Each drawing was labelled and digitised after the
session.

Following the three drawings, the participants completed a short collaborative activity (different at
each session to keep participants motivated). Then, fNIRS recordings were stopped, and still wearing
the fNIRS caps, participants completed a self-report measure of social closeness (OIS) and provided
written commentary on their experience in the session (Figure 5A). After the final session, the lead
researcher held a semi-structured interview with each participant individually to gain qualitative
insights into the participant's experience.

Self-report measures were administered via RedCap®. fNIRS recordings were synchronised using
triggers sent from PsychoPy (v. 2022.2.5; Peirce et al., 2022) marking the beginning of each drawing
and the collaborative activity.

706   **Alone**   **Together**

Figure 5. A) Flowchart visualising the protocol followed in each session. B) Left: An intergenerational dyad drawing
alone on separate pages with a grey felt divider between them, blocking their view of each other. Right: The same
intergenerational dyad drawing together on one page. The overlay of colourful lines shows the 2D motion tracking, which
was captured, but is not reported here.

5.4 Data analysis

5.4.1 Interbrain synchrony

5.4.1.1 Individual level

Channel selection. We aimed to record from the identical position on participants heads at each
session, however due to the stretchy nature of neoprene caps and the inevitability of hair
growth/haircuts, optode positions varied from session to session. To overcome this challenge, we
prioritise proximity to the centre of each ROI (De Felice et al., 2024). That is, for each participant at
each session, we selected the best channel per ROI. First, we excluded channels without visible
cardiac rhythms (~1 Hz), with a scalp-coupling index (SCI; Pollonini et al., 2014) of <.7 for

frequencies between 0.7-1.35 Hz, and with an interoptode distance <20 mm or >40 mm. Next, we
ranked the remaining channels by distance from the centre of the ROI (closest to furthest) and
selected the closest channel as the 'best channel' (Figure 6A). See Supplementary Table 0 for
description of number of recordings and sessions excluded, as well as reasons with for exclusion.

Preprocessing. We preprocessed the fNIRS signals using MNE (version 1.71; Gramfort et al., 2013,
2014), MNE-NIRS (version 0.71; Luke et al., 2021). As summarised in Figure 6A, the signal was
converted from raw intensity to optical density. Motion artefacts were corrected using the temporal
derivative distribution repair algorithm (Santosa et al., 2019) and short channel regression was
applied using the short channel with the highest SCI (Luke et al., 2021). Next, the signal was
converted to concentrations of HbO and HbR using the Modified Beer–Lambert law (Delpy et al.,
1988; Kocsis et al., 2006) using an age-specific partial pathlength factor to account for age-related
differences in the biological tissues probed by NIR light (Garcia-Castro, n.d.; Scholkmann & Wolf,
2013). This algorithm is suitable for participants aged 0-70 years. For participants aged >70 years, we
specified the age parameter as 70. Last, a bandpass filter from 0.015 to 0.4 Hz was applied.

5.4.1.2 Dyad level

Wavelet transform coherence (WTC). We computed INS values for all possible ROI pairs (N = 10;
homologous ROI pairs = LIFG~LIFG, RIFG~RIFG, LTPJ~LTPJ, RTPJ~RTPJ; non homologous
ROI pairs = LIFG~RIFG, LIFG~LTPJ, LIFG~RTPJ, LTPJ~RTPJ, RIFG~LTPJ, RIFG~RTPJ) for
each real dyad and pseudo dyad (Figure 6A). Pseudo dyads are dyads made by pairing participants
who never interacted, but align with our group types (e.g., both younger adults, or one older and one
younger adult). We computed INS values for pseudo dyads to obtain a null distribution against which
to compare INS in real dyads. Differences in INS between real and pseudo dyads for specific ROI
pairs indicate that INS levels are the result of the real interactions of real pairs and do not occur
spuriously between any two people completing the same tasks in the same setting. WTC was
computed using HyPyP (version 0.5.0b11; Ayrolles et al., 2021) with a complex Morlet wavelet, for
which the period range was defined as 5-50 seconds (0.02-0.2 Hz).

5.4.1.3 Group level

For our preregistered statistical analyses, we followed a Bayesian approach to multi-level regression
(McElreath, 2020). We fit multi-level models using the brms package (version 22.2; Bürkner, 2017)
in the R language (version 4.5.1; R Core Team, 2022) within the RStudio IDE (version
2025.05.1+513; RStudio Team, 2020). From these models, we report and interpret the posterior
distribution of key parameters using a 95% credible interval, calculated via the 95% highest posterior
density region (HPD) method (McElreath, 2020). For readers more accustomed to a Frequentist
approach that presents p-values, we recommend perusing Kruschke & Liddell (2018), and we offer
the following (simplified) heuristic for interpreting 95% HPDs: Parameter and comparison estimates
can be said to be *substantial* when the 95% HPD does not contain zero and to be trends when the tip
of an 95% HPD-tail overlaps with zero.

Self-report measures. To assess the extent to which loneliness, social closeness, and attitudes
toward generations change across the six sessions, we fit the following model for each self-report
measure separately, using individual scores (i.e., not dyadic sum or difference scores): *Score ~ 1 +*
*Group * Session + (1|DyadID/ParticipantID)*. We subsequently computed contrasts between groups
(intergenerational vs. same generation).

INS for sessions combined. We first sought to establish which, if any, ROI pairs showed
substantially different INS levels for real dyads relative to pseudo dyads. We fit separate models for
each ROI pair (n = 10), task (n = 2; drawing alone or drawing together), and group (n = 2;
intergenerational or same generation): *INS ~ Dyad Type * Session*, where *Dyad Type* refers to real
and pseudo dyads. Contrasts between real and pseudo dyads per model identified the ROI pairs for
further analysis (Figure 6B).

Changes in INS across sessions. We first sought to establish which, if any, ROI pairs showed
substantially different INS trajectories across sessions for real dyads relative to pseudo dyads. We fit
separate models for each ROI pair (n = 10), task (n = 2; drawing alone or drawing together), and
group (n = 2; intergenerational or same generation): *INS ~ Dyad Type * Session*, where *Dyad Type*
refers to real and pseudo dyads. Contrasts between real and pseudo dyads per model identified the
ROI pairs for further analysis (Figure 6B).

Further analyses with identified ROI pairs. We first fit models to assess whether intergenerational
dyads show any directionality for non-homologous ROI pairs (e.g., high INS for older adult's left
IFG and younger adult's right IFG, but not the inverse). Next, we fit one model for all sessions
combined and one for changes across sessions including only real dyads and the respectively
identified ROI pairs (models shown second tier of Figure 6B). We computed planned contrasts to
examine the effect of task (drawing alone vs. drawing together), group (intergenerational vs. same
generation), and the interaction between task and group. Finally, we fit a model (shown in second-
last tier of Figure 6B) to all ROI pairs identified (sessions combined and across sessions) to examine
the extent to which self-report measures of loneliness, social closeness (OIS) and attitudes toward
generations (allophilia) predict levels of INS. To capture dyadic aspects of each measure, we
included summed and absolute difference scores per dyad (Boukarras et al., 2025; Kenny et al.,
2006). We report HbO findings in the main report and HbR findings in the Supplementary Material,
with commentary on commonalities throughout the main report.

Figure 6. A) Flowchart visualising the procedure for selecting which channels to include in analysis for individual
participants, preprocessing steps for individual participants and calculation of WTC for dyads. B) Flowchart visualising
the order in which group-level analyses were performed in the R package brms, and showing the models fit.

Author contributions

RM: Conceptualization, equipment donation, data curation, formal analysis, Investigation, Visualization, Writing – original draft. GD: Methodology, Software, Writing – review & editing. ESC: Conceptualization, funding acquisition, Project administration, Writing – review & editing.

Acknowledgements

From the Social Brain Science Lab at ETH Zurich, we thank Tessa Portier and Medea Häuselmann for their assistance with data collection, Linda Fanconi for scheduling participants' sessions, Alistair Gadola for coding participant behaviour, Fanny Mougel for digitising the drawings. We thank Courtney Casale, Chantal Nagel, and Tessa Portier for proofreading. From the CHU Sainte-Justine Azrieli Research Center of the University of Montreal, we thank Patrice Fortin for collaborating to make HyPyP suitable for fNIRS hyperscanning data and providing hands-on assistance while package functions were in development. We also thank the survey center operated by UZH Healthy Longevity Center, University of Zurich, for their assistance with recruitment of older adults from the Zurich community.

Conflict of interest

The authors declare no potential conflict of interest.

Data availability statement

These data were collected after the submission of the following preregistration: https://osf.io/hz6tm. The data collected and analysed for this study can be found in our repository entitled 'Longitudinal perspectives on intergenerational inter-brain synchrony' on OSF: https://osf.io/xcgp6/.

Funding

Two Cortivision Photon Cap devices were provided to RM through the Cortivision Pathfinder Program (grant number CPP-2023/09/01). Oil pastels were provided to RM by Caran D'Ache (agreement number 712517). RM and ESC were supported by the Social Brain Sciences Lab at ETH Zurich. GD was supported by the Institute for Data Valorization, Montreal and the Canada First Research Excellence Fund (IVADO; CF00137433), the Fonds de recherche du Québec (FRQ; 285289), and the Natural Sciences and Engineering Research Council of Canada (NSERC; DGECR-2023-00089).

References

Adel, L., Moses, L., Irvine, E., Greenway, K. T., Dumas, G., & Lifshitz, M. (2025). A systematic review of hyperscanning in clinical encounters. *Neuroscience & Biobehavioral Reviews*, *176*, 106248. https://doi.org/10.1016/j.neubiorev.2025.106248

Ahkter-Kahn, S., & Au, R. (2020). Why loneliness interventions are unsuccessful: A call for precision health. *Advances in Geriatric Medicine and Research*, *2*(3), e200016. https://doi.org/10.20900/agmr20200016

Anderson, S., Fast, J., Keating, N., Eales, J., Chivers, S., & Barnet, D. (2017). Translating knowledge: Promoting health through intergenerational community arts programming. *Health Promotion Practice*, *18*(1), 15–25. https://doi.org/10.1177/1524839915625037

Arnold, A. J., Kappes, H. B., Klinenberg, E., & Winkielman, P. (2021). The role of comparisons in judgments of loneliness. *Frontiers in Psychology*, *12*, 498305. https://doi.org/10.3389/fpsyg.2021.498305

Aron, A., Aron, E. N., & Smollan, D. (1992). Inclusion of other in the self scale and the structure of interpersonal closeness. *Journal of Personality and Social Psychology*, *63*(4), 17. https://doi.org/10.1037/0022-3514.63.4.596

Ayalon, L., Shiovitz-Ezra, S., & Roziner, I. (2016). A cross-lagged model of the reciprocal associations of loneliness and memory functioning. *Psychology and Aging*, *31*(3), 255–261. https://doi.org/10.1037/pag0000075

Ayrolles, A., Brun, F., Chen, P., Djalovski, A., Beauxis, Y., Delorme, R., Bourgeron, T., Dikker, S., & Dumas, G. (2021). HyPyP: A Hyperscanning Python Pipeline for inter-brain connectivity analysis. *Social Cognitive and Affective Neuroscience*, *16*(1–2), 72–83. https://doi.org/10.1093/scan/nsaa141

Bartlett, S. P., Solomon, P., & Gellis, Z. (2021). Comparative effectiveness of intergenerational service-learning programs on student outcomes of knowledge, attitude, and ageism. *Educational Gerontology*, *47*(12), 559–573. https://doi.org/10.1080/03601277.2021.2011596

Bast, N., Polzer, L., Raji, N., Schnettler, L., Kleber, S., Lemler, C., Kitzerow-Cleven, J., Kim, Z., Schaer, M., & Freitag, C. M. (2025). Early intervention increases reactive joint attention in autistic preschoolers with arousal regulation as mediator. *European Child & Adolescent Psychiatry*. https://doi.org/10.1007/s00787-025-02738-1

Bessaha, M. L., Sabbath, E. L., Morris, Z., Malik, S., Scheinfeld, L., & Saragossi, J. (2020). A systematic review of loneliness interventions among non-elderly adults. *Clinical Social Work Journal*, *48*(1), 110–125. https://doi.org/10.1007/s10615-019-00724-0

Bevilacqua, D., Davidesco, I., Wan, L., Chaloner, K., Rowland, J., Ding, M., Poeppel, D., & Dikker, S. (2019). Brain-to-brain synchrony and learning outcomes vary by student–teacher dynamics: Evidence from a real-world classroom electroencephalography study. *Journal of Cognitive Neuroscience*, *31*(3), 401–411. https://doi.org/10.1162/jocn_a_01274

Bi, X., Cui, H., & Ma, Y. (2023). Hyperscanning Studies on Interbrain Synchrony and Child Development: A Narrative Review. *Neuroscience*, *530*, 38–45. https://doi.org/10.1016/j.neuroscience.2023.08.035

Boukarras, S., Placidi, V., Rossano, F., Era, V., Aglioti, S. M., & Candidi, M. (2025). Interpersonal physiological synchrony during dyadic joint action is increased by task novelty and reduced by social anxiety. *Psychophysiology*, *62*(3), e70031. https://doi.org/10.1111/psyp.70031

Brilliant, D., Takeuchi, H., Nouchi, R., Yokoyama, R., Kotozaki, Y., Nakagawa, S., Hanawa, S., Sekiguchi, A., Ikeda, S., Sakaki, K., Kawata, K. H. D. S., Nozawa, T., Yokota, S., Magistro, D., & Kawashima, R. (2022). Loneliness inside of the brain: Evidence from a large dataset of resting-state fMRI in young adult. *Scientific Reports*, *12*(1), 7856. https://doi.org/10.1038/s41598-022-11724-5

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, *80*(1), 1–28. https://doi.org/10.18637/jss.v080.i01

Cachón-Alonso, L., Hakulinen, C., Jokela, M., Komulainen, K., & Elovainio, M. (2023). Loneliness and cognitive function in older adults: Longitudinal analysis in 15 countries. *Psychology and Aging*, *38*(8), 778–789. https://doi.org/10.1037/pag0000777

Cacioppo, J. T., & Hawkley, L. C. (2009). Perceived social isolation and cognition. *Trends in Cognitive Sciences*, *13*(10), 447–454. https://doi.org/10.1016/j.tics.2009.06.005

Cacioppo, S., Capitanio, J. P., & Cacioppo, J. T. (2014). Toward a neurology of loneliness. *Psychological Bulletin*, *140*(6), 1464–1504. https://doi.org/10.1037/a0037618

Cigna US. (2018). *Loneliness Index*. Cigna US. https://www.multivu.com/players/English/8294451-cigna-us-loneliness-survey/#:~:text=The%20survey%20of%20more%20than,people%20who%20really%20understand%20them.

Corbin, I. M., & Dhand, A. (2024). Unshared minds, decaying worlds: Towards a pathology of chronic loneliness. *The Journal of Medicine and Philosophy: A Forum for Bioethics and Philosophy of Medicine*, *49*(4), 354–366. https://doi.org/10.1093/jmp/jhae020

Cui, X., Bryant, D. M., & Reiss, A. L. (2012). NIRS-based hyperscanning reveals increased interpersonal coherence in superior frontal cortex during cooperation. *NeuroImage*, *59*(3), 2430–2437. https://doi.org/10.1016/j.neuroimage.2011.09.003

Czeszumski, A., Liang, S. H.-Y., Dikker, S., König, P., Lee, C.-P., Koole, S. L., & Kelsen, B. (2022). Cooperative behavior evokes interbrain synchrony in the prefrontal and temporoparietal cortex: A systematic review and meta-analysis of fNIRS hyperscanning studies. *Eneuro*, *9*(2), ENEURO.0268-21.2022. https://doi.org/10.1523/ENEURO.0268-21.2022

De Felice, S., Chand, T., Croy, I., Engert, V., Goldstein, P., Holroyd, C. B., Kirsch, P., Krach, S., Ma, Y., Scheele, D., Schurz, M., Schweinberger, S. R., Hoehl, S., & Vrticka, P. (2025). Relational neuroscience: Insights from hyperscanning research. *Neuroscience & Biobehavioral Reviews*, *169*, 105979. https://doi.org/10.1016/j.neubiorev.2024.105979

De Felice, S., Hakim, U., Gunasekara, N., Pinti, P., Tachtsidis, I., & Hamilton, A. (2024). Having a chat and then watching a movie: How social interaction synchronises our brains during co-watching. *Oxford Open Neuroscience*, *3*, kvae006. https://doi.org/10.1093/oons/kvae006

De Felice, S., Hamilton, A. F. D. C., Ponari, M., & Vigliocco, G. (2023). Learning from others is good, with others is better: The role of social interaction in human acquisition of new knowledge. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *378*(1870), 20210357. https://doi.org/10.1098/rstb.2021.0357

De Jong Gierveld, J., & van Tilburg, T. (2006). A 6-item scale for overall, emotional, and social loneliness: Confirmatory tests on survey data. *Research on Aging*, *28*(5), 582–598. https://doi.org/10.1177/0164027506289723

Delpy, D. T., Cope, M., Van Der Zee, P., Arridge, S., Wray, S., & Wyatt, J. (1988). Estimation of optical pathlength through tissue from direct time of flight measurement. *Physics in Medicine and Biology*, *33*(12), 1433–1442. https://doi.org/10.1088/0031-9155/33/12/008

Dikker, S., Brito, N. H., & Dumas, G. (2024). It takes a village: A multi-brain approach to studying multigenerational family communication. *Developmental Cognitive Neuroscience*, *65*, 101330. https://doi.org/10.1016/j.dcn.2023.101330

Dikker, S., Wan, L., Davidesco, I., Kaggen, L., Oostrik, M., McClintock, J., Rowland, J., Michalareas, G., Van Bavel, J. J., Ding, M., & Poeppel, D. (2017). Brain-to-brain synchrony tracks real-world dynamic group interactions in the classroom. *Current Biology*, *27*(9), 1375–1380. https://doi.org/10.1016/j.cub.2017.04.002

Ding, K., Wang, H., Wang, Q., Li, H., & Li, C. (2023). Inhibitory control associated with the neural mechanism of joint attention in preschoolers: An fNIRS evidence. *International Journal of Psychophysiology*, *192*, 53–61. https://doi.org/10.1016/j.ijpsycho.2023.08.006

Djalovski, A., Dumas, G., Kinreich, S., & Feldman, R. (2021). Human attachments shape interbrain synchrony toward efficient performance of social goals. *NeuroImage*, *226*, 117600. https://doi.org/10.1016/j.neuroimage.2020.117600

Dommer, L., Jäger, N., Scholkmann, F., Wolf, M., & Holper, L. (2012). Between-brain coherence during joint n-back task performance: A two-person functional near-infrared spectroscopy study. *Behavioural Brain Research*, *234*(2), 212–222. https://doi.org/10.1016/j.bbr.2012.06.024

Doricchi, F., Lasaponara, S., Pazzaglia, M., & Silvetti, M. (2022). Left and right temporal-parietal junctions (TPJs) as "match/mismatch" hedonic machines: A unifying account of TPJ function. *Physics of Life Reviews*, *42*, 56–92. https://doi.org/10.1016/j.plrev.2022.07.001

Drewelies, J., Windsor, T. D., Duezel, S., Demuth, I., Wagner, G. G., Lindenberger, U., Gerstorf, D., & Ghisletta, P. (2022). Age trajectories of perceptual speed and loneliness: Separating between-person and within-person associations. *The Journals of Gerontology: Series B*, *77*(1), 118–129. https://doi.org/10.1093/geronb/gbab180

Dumas, G., Nadel, J., Soussignan, R., Martinerie, J., & Garnero. (2010). Inter-brain synchronization during social interaction. *PLOS ONE*, *5*(8), e12166.

Düzel, S., Drewelies, J., Gerstorf, D., Demuth, I., Steinhagen-Thiessen, E., Lindenberger, U., & Kühn, S. (2019). Structural Brain Correlates of Loneliness among Older Adults. *Scientific Reports*, *9*(1), 13569. https://doi.org/10.1038/s41598-019-49888-2

Ellingsen, D.-M., Isenburg, K., Jung, C., Lee, J., Gerber, J., Mawla, I., Sclocco, R., Grahl, A., Anzolin, A., Edwards, R. R., Kelley, J. M., Kirsch, I., Kaptchuk, T. J., & Napadow, V. (2023). Brain-to-brain mechanisms underlying pain empathy and social modulation of pain in the patient-clinician interaction. *Proceedings of the National Academy of Sciences*, *120*(26), e2212910120. https://doi.org/10.1073/pnas.2212910120

Galinsky, A. D., Ku, G., & Wang, C. S. (2005). Perspective-taking and self-other overlap: Fostering social bonds and facilitating social coordination. *Group Processes & Intergroup Relations*, *8*(2), 109–124. https://doi.org/10.1177/1368430205051060

Garcia-Castro, G. (n.d.). *Implementing the Differential Pathlength Factor (DPF) in Python: The Scholkmann Method.* https://gongcastro.github.io/blog/dpf-scholkmann/dpf-scholkmann.html

Gonzales, E., Morrow-Howell, N., & Gilbert, P. (2010). Changing medical students' attitudes toward older adults. *Gerontology & Geriatrics Education*, *31*(3), 220–234. https://doi.org/10.1080/02701960.2010.503128

Goodwin, R. (2013). *Personal relationships across cultures* (1st ed.). Routledge. https://doi.org/10.4324/9780203434161

Grahl, A., Anzolin, A., Ellis, S., Barton-Zuckerman, M., Lee, J., Jung, C., Isenburg, K., Ellingsen, D.-M., Kelley, J., Kaptchuk, T., & Napadow, V. (2023). The patient-clinician relationship, expectancy and prior experience can modulate fibromyalgia treatment outcomes: A longitudinal fmrihyperscan study. *The Journal of Pain*, *24*(4), 74–75. https://doi.org/10.1016/j.jpain.2023.02.217

Grahl, A., Anzolin, A., Isenburg, K., Lee, J., Ellingsen, D.-M., Jung, C., Zuckerman, M. B., Gerber, J., Kelley, J., Kirsch, I., Kaptchuk, T., & Napadow, V. (2021). Brain and behavioral correlates of the patient-clinician relationship: A longitudinal fMRI hyper-scanning study of chronic pain patients. *The Journal of Pain*, *22*(5), 602. https://doi.org/10.1016/j.jpain.2021.03.097

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Goj, R., Jas, M., Brooks, T., Parkkonen, L., & Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, *7*, 1–13. https://doi.org/10.3389/fnins.2013.00267

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Parkkonen, L., & Hämäläinen, M. S. (2014). MNE software for processing MEG and EEG data. *NeuroImage*, *86*, 446–460. https://doi.org/10.1016/j.neuroimage.2013.10.027

Hampshire, A., Chamberlain, S. R., Monti, M. M., Duncan, J., & Owen, A. M. (2010). The role of the right inferior frontal gyrus: Inhibition and attentional control. *NeuroImage*, *50*(3), 1313–1319. https://doi.org/10.1016/j.neuroimage.2009.12.109

Hawkley, L. C., Buecker, S., Kaiser, T., & Luhmann, M. (2022). Loneliness from young adulthood to old age: Explaining age differences in loneliness. *International Journal of Behavioral Development*, *46*(1), 39–49. https://doi.org/10.1177/0165025420971048

Hirata, M., Ikeda, T., Kikuchi, M., Kimura, T., Hiraishi, H., Yoshimura, Y., & Asada, M. (2014). Hyperscanning MEG for understanding mother-child cerebral interactions. *Frontiers in Human Neuroscience*, *8*. https://doi.org/10.3389/fnhum.2014.00118

Hoehl, S., Bánki, A., Brzozowska, A., Carollo, A., Kostorz, K., Nguyen, T., Pletti, C., Reisner, S., Schäfer, V., Schaetz, C., & Tünte, M. R. (2025). *A developmental framework of interpersonal neural synchrony*. PsyArXiv. https://doi.org/10.31234/osf.io/m7uyq_v1

Holper, L., Scholkmann, F., & Wolf, M. (2012). Between-brain connectivity during imitation measured by fNIRS. *NeuroImage*, *63*(1), 212–222. https://doi.org/10.1016/j.neuroimage.2012.06.028

Holt-Lunstad, J., & Perissinotto, C. (2023). Social isolation and loneliness as medical issues. *New England Journal of Medicine*, *388*(3), 193–195. https://doi.org/10.1056/NEJMp2208029

House, J. S., Landis, K. R., & Umberson, D. (1988). Social relationships and health. *Science*, *241*(4865), 540–545. https://doi.org/10.1126/science.3399889

Jacoby, N., Bruneau, E., Koster-Hale, J., & Saxe, R. (2016). Localizing pain matrix and theory of mind networks with both verbal and non-verbal stimuli. *NeuroImage*, *126*, 39–48. https://doi.org/10.1016/j.neuroimage.2015.11.025

Jenkins, L. K., Farrer, R., & Aujla, I. J. (2021). Understanding the impact of an intergenerational arts and health project: A study into the psychological well-being of participants, carers and artists. *Public Health*, *194*, 121–126. https://doi.org/10.1016/j.puhe.2021.02.029

Kelsen, B. A., Sumich, A., Kasabov, N., Liang, S. H. Y., & Wang, G. Y. (2022). What has social neuroscience learned from hyperscanning studies of spoken communication? A systematic review. *Neuroscience & Biobehavioral Reviews*, *132*, 1249–1262. https://doi.org/10.1016/j.neubiorev.2020.09.008

Kenny, D. A., Kashy, D. A., & Cook, W. A. (2006). Using multilevel modeling to study dyads. In *Dyadic Data Analysis,* (pp. 78–99). Guilford Publications. https://ebookcentral.proquest.com/lib/ethz/detail.action?docID=362550

Kocsis, L., Herman, P., & Eke, A. (2006). The modified Beer-Lambert law revisited. *Physics in Medicine and Biology*, *51*(5), N91–N98. https://doi.org/10.1088/0031-9155/51/5/N02

Koechlin, E., Ody, C., & Kouneiher, F. (2003). The architecture of cognitive control in the human prefrontal cortex. *Science*, *302*(5648), 1181–1185. https://doi.org/10.1126/science.1088545

Kruschke, J. K., & Liddell, T. M. (2018). The Bayesian New Statistics: Hypothesis testing, estimation, meta-analysis, and power analysis from a Bayesian perspective. *Psychonomic Bulletin & Review*, *25*(1), 178–206. https://doi.org/10.3758/s13423-016-1221-4

Labonté-LeMoyne, É., Léger, P.-M., Resseguier, B., Bastarache-Roberge, M.-C., Fredette, M., Sénécal, S., & Courtemanche, F. (2016). Are we in flow neurophysiological correlates of flow states in a collaborative game. *Proceedings of the 2016 CHI Conference Extended Abstracts on Human Factors in Computing Systems*, 1980–1988. https://doi.org/10.1145/2851581.2892356

Lachat, F., Hugueville, L., Lemaréchal, J.-D., Conty, L., & George, N. (2012). Oscillatory brain correlates of live joint attention: A dual-EEG study. *Frontiers in Human Neuroscience*, *6*. https://doi.org/10.3389/fnhum.2012.00156

Lam, J. A., Murray, E. R., Yu, K. E., Ramsey, M., Nguyen, T. T., Mishra, J., Martis, B., Thomas, M. L., & Lee, E. E. (2021). Neurobiology of loneliness: A systematic review. *Neuropsychopharmacology*, *46*(11), 1873–1887. https://doi.org/10.1038/s41386-021-01058-7

Lee, K., Jarrott, S. E., & Juckett, L. A. (2020). Documented outcomes for older adults in intergenerational programming: A scoping review: research. *Journal of Intergenerational Relationships*, *18*(2), 113–138. https://doi.org/10.1080/15350770.2019.1673276

Li, L., Wang, H., Luo, H., Zhang, X., Zhang, R., & Li, X. (2020). Interpersonal neural synchronization during cooperative behavior of basketball players: A fNIRS-based hyperscanning study. *Frontiers in Human Neuroscience*, *14*. https://www.frontiersin.org/articles/10.3389/fnhum.2020.00169

Liao, T., Zhuoga, C., & Chen, X. (2023). Contact with grandparents and young people's explicit and implicit attitudes toward older adults. *BMC Psychology*, *11*(1), 289. https://doi.org/10.1186/s40359-023-01344-7

Lieberz, J., Shamay-Tsoory, S. G., Saporta, N., Esser, T., Kuskova, E., Stoffel-Wagner, B., Hurlemann, R., & Scheele, D. (2021). Loneliness and the social brain: How perceived social isolation impairs human interactions. *Advanced Science*, *8*(21), 2102076. https://doi.org/10.1002/advs.202102076

Liu, D., Liu, S., Liu, X., Zhang, C., Li, A., Jin, C., Chen, Y., Wang, H., & Zhang, X. (2018). Interactive Brain Activity: Review and Progress on EEG-Based Hyperscanning in Social Interactions. *Frontiers in Psychology*, *9*, 1862. https://doi.org/10.3389/fpsyg.2018.01862

Lokon, E., Li, Y., & Kunkel, S. (2020). Allophilia: Increasing college students' "liking" of older adults with dementia through arts-based intergenerational experiences. *Gerontology & Geriatrics Education*, *41*(4), 494–507. https://doi.org/10.1080/02701960.2018.1515740

Long, Y., Chen, C., Wu, K., Zhou, S., Zhou, F., Zheng, L., Zhao, H., Zhai, Y., & Lu, C. (2022). Interpersonal conflict increases interpersonal neural synchronization in romantic couples. *Cerebral Cortex*, *32*(15), 3254–3268. https://doi.org/10.1093/cercor/bhab413

Lu, P., Oh, J., Leahy, K. E., & Chopik, W. J. (2021). Friendship importance around the world: Links to cultural factors, health, and well-being. *Frontiers in Psychology*, *11*, 570839. https://doi.org/10.3389/fpsyg.2020.570839

Luke, R., Larson, E., Shader, M. J., Innes-Brown, H., Van Yper, L., Lee, A. K. C., Sowman, P. F., & McAlpine, D. (2021). Analysis methods for measuring passive auditory fNIRS responses generated by a block-design paradigm. *Neurophotonics*, *8*(2), 025008. https://doi.org/10.1117/1.nph.8.2.025008

McElreath, R. (2020). *Statistical Rethinking; A Bayesian Course with Examples in R and Stan* (2nd Edition). Chapman and Hall/CRC.

Misaki, M., Kerr, K. L., Ratliff, E. L., Cosgrove, K. T., Simmons, W. K., Morris, A. S., & Bodurka, J. (2021). Beyond synchrony: The capacity of fMRI hyperscanning for the study of human social interaction. *Social Cognitive and Affective Neuroscience*, *16*(1–2), 84–92. https://doi.org/10.1093/scan/nsaa143

Moffat, R., Casale, C. E., & Cross, E. S. (2024). Mobile fNIRS for exploring inter-brain synchrony across generations and time. *Frontiers in Neuroergonomics*, *4*. https://doi.org/10.3389/fnrgo.2023.1260738

Mundy, P. (2018). A review of joint attention and social-cognitive brain systems in typical development and autism spectrum disorder. *European Journal of Neuroscience*, *47*(6), 497–514. https://doi.org/10.1111/ejn.13720

Nalborczyk, L., Batailler, C., Vilain, A., & Bürkner, P.-C. (2019). An introduction to Bayesian multilevel models using brms: A case study of gender effects on vowel variability in standard indonesian. *Journal of Speech, Language, and Hearing Research*, *62*, 1225–1242. https://doi.org/10.1044/2018_JSLHR-S-18-0006

Newman, S., Ward, C., Smith, T., Wilson, J., & McCrea, J. (2014). *Intergenerational programs: Past, present and future*
(Rev. ed.). Routledge.

Nguyen, T., Bánki, A., Markova, G., & Hoehl, S. (2020). Studying parent-child interaction with hyperscanning. In
Progress in Brain Research (Vol. 254, pp. 1–24). Elsevier. https://doi.org/10.1016/bs.pbr.2020.05.003

Noone, C., & Yang, K. (2022). Community-based responses to loneliness in older people: A systematic review of
qualitative studies. *Health & Social Care in the Community*, *30*(4), e859–e873.
https://doi.org/10.1111/hsc.13682

Noreika, V., Georgieva, S., Wass, S., & Leong, V. (2020). 14 challenges and their solutions for conducting social
neuroscience and longitudinal EEG research with infants. *Infant Behavior and Development*, *58*, 101393.
https://doi.org/10.1016/j.infbeh.2019.101393

Okruszek, Ł., Piejka, A., Krawczyk, M., Schudy, A., Wiśniewska, M., Żurek, K., & Pinkham, A. (2021). Owner of a
lonely mind? Social cognitive capacity is associated with objective, but not perceived social isolation in healthy
individuals. *Journal of Research in Personality*, *93*, 104103. https://doi.org/10.1016/j.jrp.2021.104103

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). Fieldtrip: Open source software for advanced analysis of
MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*, 1–9.
https://doi.org/10.1155/2011/156869

Oostenveld, R., & Praamstra, P. (2001). The five percent electrode system for high-resolution EEG and ERP
measurements. *Clinical Neurophysiology*, *112*(4), 713–719. https://doi.org/10.1016/S1388-2457(00)00527-7

O'Rourke, H. M., Collins, L., & Sidani, S. (2018). Interventions to address social connectedness and loneliness for older
adults: A scoping review. *BMC Geriatrics*, *18*(1), 214. https://doi.org/10.1186/s12877-018-0897-x

Osaka, N., Minamoto, T., Yaoi, K., Azuma, M., Shimada, Y. M., & Osaka, M. (2015). How two brains make one
synchronized mind in the inferior frontal cortex: fNIRS-based hyperscanning during cooperative singing.
Frontiers in Psychology, *6*. https://www.frontiersin.org/articles/10.3389/fpsyg.2015.01811

Pan, Y., & Cheng, X. (2020). Two-person approaches to studying social interaction in psychiatry: Uses and clinical
relevance. *Frontiers in Psychiatry*, *11*, 301. https://doi.org/10.3389/fpsyt.2020.00301

Peirce, J. W., Hirst, R. J., & MacAskill, M. R. (2022). *Building Experiments in PsychoPy* (2nd ed.). Sage.
https://uk.sagepub.com/en-gb/eur/building-experiments-in-psychopy/book273700

Petersen, J. (2023). A meta-analytic review of the effects of intergenerational programs for youth and older adults.
Educational Gerontology, *49*(3), 175–189. https://doi.org/10.1080/03601277.2022.2102340

Pittinsky, T. L., Rosenthal, S. A., & Montoya, R. M. (2011). Measuring positive attitudes toward outgroups:
Development and validation of the Allophilia Scale. In L. R. Tropp & R. K. Mallett (Eds.), *Moving beyond*
prejudice reduction: Pathways to positive intergroup relations. (pp. 41–60). American Psychological
Association. https://doi.org/10.1037/12319-002

Pollonini, L., Olds, C., Abaya, H., Bortfeld, H., Beauchamp, M. S., & Oghalai, J. S. (2014). Auditory cortex activation to
natural speech and simulated cochlear implant speech measured with functional near-infrared spectroscopy.
Hearing Research, *309*, 84–93. https://doi.org/10.1016/j.heares.2013.11.007

Qualter, P., Vanhalst, J., Harris, R., Van Roekel, E., Lodder, G., Bangee, M., Maes, M., & Verhagen, M. (2015).
Loneliness across the life span. *Perspectives on Psychological Science*, *10*(2), 250–264.
https://doi.org/10.1177/1745691615568999

R Core Team. (2022). *R: A language and environment for statistical computing* [Computer software]. R Foundation for
Statistical Computing. http://www.r-project.org/

Redcay, E., Kleiner, M., & Saxe, R. (2012). Look at this: The neural correlates of initiating and responding to bids for
joint attention. *Frontiers in Human Neuroscience*, *6*. https://doi.org/10.3389/fnhum.2012.00169

RStudio Team. (2020). *RStudio: Integrated Development for R* [Computer software]. RStudio, Inc.
http://www.rstudio.com/

Rubin, S. E., Gendron, T. L., Wren, C. A., Ogbonna, K. C., Gonzales, E. G., & Peron, E. P. (2015). Challenging
gerontophobia and ageism through a collaborative intergenerational art program. *Journal of Intergenerational*
Relationships, *13*(3), 241–254. https://doi.org/10.1080/15350770.2015.1058213

Santosa, H., Huppert, T. J., Fishburn, F., & Zhai, X. (2019). Investigation of the sensitivity-specificity of canonical- and
deconvolution-based linear models in evoked functional near-infrared spectroscopy. *Neurophotonics*, *6*(2), 1–10.
https://doi.org/10.1117/1.NPh.6.2.025009

Saporta, N., Scheele, D., Lieberz, J., Nevat, M., Kanterman, A., Hurlemann, R., & Shamay-Tsoory, S. G. (2022). Altered
activation in the action observation system during synchronization in high loneliness individuals. *Cerebral*
Cortex, *33*(2), 385–402. https://doi.org/10.1093/cercor/bhac073

Saxe, R., & Kanwisher, N. (2003). People thinking about thinking people: The role of the temporo-parietal junction in
"theory of mind." *NeuroImage*, *19*(4), 1835–1842. https://doi.org/10.1016/S1053-8119(03)00230-1

Scarpetti, G., Shadowen, H., Williams, G. A., Winkelmann, J., Kroneman, M., Groenewegen, P. P., De Jong, J. D.,
Fronteira, I., Augusto, G. F., Hsiung, S., Slade, S., Rojatz, D., Kallayova, D., Katreniakova, Z., Nagyova, I.,
Kylänen, M., Vracko, P., Jesurasa, A., Wallace, Z., … Van Ginneken, E. (2024). A comparison of social
prescribing approaches across twelve high-income countries. *Health Policy*, *142*, 104992.
https://doi.org/10.1016/j.healthpol.2024.104992

Scholkmann, F., & Wolf, M. (2013). General equation for the differential pathlength factor of the frontal human head
depending on wavelength and age. *Journal of Biomedical Optics*, *18*(10), 105004.
https://doi.org/10.1117/1.JBO.18.10.105004

Sened, H., Gorst Kaduri, K., Nathan Gamliel, H., Rafaeli, E., Zilcha-Mano, S., & Shamay-Tsoory, S. (2025). Inter-brain
plasticity as a mechanism of change in psychotherapy: A proof of concept focusing on test anxiety.
Psychotherapy Research, 1–15. https://doi.org/10.1080/10503307.2025.2451798

Shuo, G., Li, Y., Geng, Y., Li, D., Xu, Q., Yang, D., Zhang, Y., & Li, R. (2025). *Advancing inter-brain synchrony*
measurement: A comparative hyperscanning study of high-density diffuse optical tomography and functional
near-infrared spectroscopy. bioRxiv. https://doi.org/10.1101/2025.07.02.662877

Simons, D. J., Shoda, Y., & Lindsay, D. S. (2017). Constraints on Generality (COG): A Proposed Addition to All
Empirical Papers. *Perspectives on Psychological Science*, *12*(6), 1123–1128.
https://doi.org/10.1177/1745691617708630

Sinha, N., Maszczyk, T., Wanxuan, Z., Tan, J., & Dauwels, J. (2016). EEG hyperscanning study of inter-brain synchrony
during cooperative and competitive interaction. *2016 IEEE International Conference on Systems, Man, and*
Cybernetics (SMC), 004813–004818. https://doi.org/10.1109/SMC.2016.7844990

Speer, S. P. H., Mwilambwe-Tshilobo, L., Tsoi, L., Burns, S. M., Falk, E. B., & Tamir, D. I. (2024). Hyperscanning
shows friends explore and strangers converge in conversation. *Nature Communications*, *15*(1), 7781.
https://doi.org/10.1038/s41467-024-51990-7

Spithoven, A. W. M., Bijttebier, P., & Goossens, L. (2017). It is all in their mind: A review on information processing
bias in lonely individuals. *Clinical Psychology Review*, *58*, 97–114. https://doi.org/10.1016/j.cpr.2017.10.003

Sun, B., Xiao, W., Lin, S., Shao, Y., Li, W., & Zhang, W. (2021). Cooperation with partners of differing social
experience: An fNIRS-based hyperscanning study. *Brain and Cognition*, *154*, 105803.
https://doi.org/10.1016/j.bandc.2021.105803

The Lancet. (2023). Loneliness as a health issue. *The Lancet*, *402*(10396), 79. https://doi.org/10.1016/S0140-
6736(23)01411-3

Toppi, J., Borghini, G., Petti, M., He, E. J., De Giusti, V., He, B., Astolfi, L., & Babiloni, F. (2016). Investigating
Cooperative Behavior in Ecological Settings: An EEG Hyperscanning Study. *PLOS ONE*, *11*(4), e0154236.
https://doi.org/10.1371/journal.pone.0154236

Turk, E., Vroomen, J., Fonken, Y., Levy, J., & Van Den Heuvel, M. I. (2022). In sync with your child: The potential of
parent–child electroencephalography in developmental research. *Developmental Psychobiology*, *64*(3).
https://doi.org/10.1002/dev.22221

United Nations. (2024). *World population prospects 2024: Summary of results*. United Nations.

Wang, D., Ren, Y., & Chen, W. (2024). Relationship evolution shapes inter-brain synchrony in affective sharing: The
role of self-expansion. *Brain Structure and Function*, *229*(9), 2269–2283. https://doi.org/10.1007/s00429-024-
02841-0

Wang, F., Gao, Y., Han, Z., Yu, Y., Long, Z., Jiang, X., Wu, Y., Pei, B., Cao, Y., Ye, J., Wang, M., & Zhao, Y. (2023). A
systematic review and meta-analysis of 90 cohort studies of social isolation, loneliness and mortality. *Nature*
Human Behaviour. https://doi.org/10.1038/s41562-023-01617-6

WHO. (2019). *What is the evidence on the role of the arts in improving health and well-being? A scoping review*. World
Health Organization. https://iris.who.int/handle/10665/329834

WHO. (2021). *Social isolation and loneliness among older people: Advocacy brief*. World Health Organization.
https://www.who.int/publications/i/item/9789240030749

WHO. (2025). *From loneliness to social connection*. World Health Organization.
https://www.who.int/publications/i/item/978240112360

Wigfield, A., Turner, R., Alden, S., Green, M., & Karania, V. K. (2022). Developing a new conceptual framework of
meaningful interaction for understanding social isolation and loneliness. *Social Policy and Society*, *21*(2), 172–
193. https://doi.org/10.1017/S147474642000055X

Xie, H., Karipidis, I. I., Howell, A., Schreier, M., Sheau, K. E., Manchanda, M. K., Ayub, R., Glover, G. H., Jung, M.,
Reiss, A. L., & Saggar, M. (2020). Finding the neural correlates of collaboration using a three-person fMRI
hyperscanning paradigm. *Proceedings of the National Academy of Sciences*, *117*(37), 23066–23072.
https://doi.org/10.1073/pnas.1917407117

Yoshioka, A., Tanabe, H. C., Sumiya, M., Nakagawa, E., Okazaki, S., Koike, T., & Sadato, N. (2021). Neural substrates
of shared visual experiences: A hyperscanning fMRI study. *Social Cognitive and Affective Neuroscience*, *16*(12),
1264–1275. https://doi.org/10.1093/scan/nsab082

Zhang, Q., Liu, Z., Qian, H., Hu, Y., & Gao, X. (2023). Interpersonal competition in elderly couples: A functional near-
infrared spectroscopy hyperscanning study. *Brain Sciences*, *13*(4), 600.
https://doi.org/10.3390/brainsci13040600

1147