Title: Effects of color-enhancing filters on color salience in normal trichromats

Authors: Camilla Simoncelli[1], Don McPherson[2], Kenneth Knoblauch[3], Michael Webster[1]

[1] Department of Psychology and Graduate Program in Integrative Neuroscience, University of Nevada Reno, US

[2] Chief Science Officer, EnChroma, 2547 8th. St., Ste. 35 Berkeley, CA 94710

[3] Univ Lyon, Université Claude Bernard Lyon 1, Inserm U1208, Stem Cell and Brain Research Institute, Bron, France

Abstract

Notch filters can alter color contrasts by selectively filtering different spectral bands of the stimulus and have been developed to enhance reddish-greenish contrasts for color-deficient observers with anomalous trichromacy. We examined the effects of such filters on color salience for normal trichromats, using a visual search paradigm where the task was to locate a color target superimposed on a variegated chromatic background, similar to foraging for fruits among foliage. Background colors varied along a bluish-yellowish or purpliish to yellow-green (short-wave cone isolating) axis, roughly spanning the range of dominant color variations in arid or lush environments. Target colors sampled a wide range of hue angles and contrasts. Testing was conducted on a computer monitor, with the filter effects simulated by calculating corresponding chromaticities with or without the filter for naturalistic (Munsell) reflectances. The filter evaluated (Enchroma SuperX® glasses) was designed to increase color contrast along a magenta-green axis. Consistent with this, search times for targets on the blue-yellow background were significantly faster for the filter condition, because the filter increased the target-background color difference. Alternatively, overall differences in search times were not observed for the S-cone background. The differences on the two backgrounds could be qualitatively accounted for by the relative salience of the stimuli predicted by a perceptual color space (CIELAB). Our results demonstrate the efficacy of the filters for enhancing visual performance for normal trichromats and naturalistic tasks, and illustrate how these effects depend on the potential color characteristics of the environment.

Introduction

Normal human color vision depends on sampling the light spectrum by three types of cone photoreceptors maximally sensitive to short (S), medium (M), or long (L) wavelengths in the visible spectrum. Information about the spectral content of the light is contained in the differences or relative responses of the different cones, manifested in the underlying cone and color opponency in post-receptoral color processing. In a fixed context, larger differences correspond to stronger color contrasts or roughly to more saturated colors, while the type of difference (e.g. which cone is stimulated more) corresponds to the chromatic direction or roughly to hue.

The spectrum of a visual stimulus depends on both the characteristics of the illumination and the surface materials (Foster, 2011). Color contrasts in a scene could be enhanced by varying the spectrum emitted by the illuminant and is the principle employed in wide color gamut lighting and displays, which use spectrally narrow primaries to increase the range of spectral differences (Feng et al., 2016; Ilic et al., 2022; Wen at al., 2021; Zhu et al., 2015;). Alternatively, the spectra can also be shaped by filtering the light reaching the eye. Notch filters block or transmit different portions of the visible spectrum, and have been developed to increase the relative differences in the signals of the L and M cones as an aid for anomalous trichromacy (the most common form of color deficiency, resulting from a reduced separation in the peaks of the two longer-wavelength sensitive cones) (Somers & Bosten, 2024). A number of recent studies have evaluated the efficacy of the filters for color perception and performance for individuals with color deficiencies, and have documented conditions under which the filters do or do not have significant impact (Gómez-Robledo et al., 2018; Lynch & Ng, 2024 ; Marques et al., 2023; Moreland et al., 2022; Nascimento & Foster, 2022; Patterson et al., 2022; Pattie et al., 2022; Rabin et al., 2022; Somers & Bosten, 2024; Somers et al., 2024; Werner et al., 2020).

In the present study, we instead examine the potential impact of spectral notch filters on *normal* color vision, focusing on how the changes in color contrast could impact the salience or conspicuousness of different colors. To assess this, we used a color search task in which observers had to locate a color target embedded in a color-varying background. Visual search is a ubiquitous natural behavior central to many visual tasks (Wolfe, 2020). The properties and characteristics of visual search have been studied extensively, and include many cognitive (e.g., goals) and sensory (e.g., stimulus salience) factors (Wolfe & Horowitz, 2017). Color provides important information for visual search (D'Zmura, 1991; D'Zmura & Mangalick, 1994), and this may have been one of the forces driving the evolution of color vision in many species. For example, a prominent theory of primate trichromacy is that it evolved to help find ripe fruit among foliage (Osorio & Vorobyev, 1996; Regan et al., 2001).

To measure color search, we used a "color foraging" paradigm developed by McDermott et al. (McDermott et al., 2010) that simulates search for colored "fruit" on densely colored backgrounds, and examined how performance on the task was affected by the changes in chromatic contrast introduced by a notch filter. The specific filter we tested was the Enchroma SuperX® glasses (EnChroma Inc., Berkeley, California, USA), which were designed for normal trichromats, and which amplify chromatic contrasts or colorimetric purity along a roughly magenta-lime axis. As illustrated below, the filtering can selectively increase the color contrast of some broadband spectra by up to 50%. Previous studies of color search have shown that search improves as the difference between the target color and background color (or color axis)

increases (Bauer et al., 1996; D'Zmura, 1991; Manalansan et al., 2025; McDermott et al., 2010; Nagy & Sanchez, 1990). We therefore expected that the effects of the filters on the search task would be consistent with the chromatic differences they introduced in the stimuli, and with how these depend on the specific chromatic properties of the targets and backgrounds.

Method

Participants

Ten color-normal observers (mean age ± SD = 27.5 ± 4.4) were tested in the Visual Perception Laboratory at the University of Nevada, Reno (UNR). Most were undergraduate or graduate students at the university, who received course credit for participation. All had normal or corrected-to-normal vision and normal color vision as assessed by the Ishihara Pseudoisochromatic Plates (Clark, 1924). Participation required written informed consent, and all procedures were in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki) and approved by the Institutional Review board of the University of Nevada, Reno. A single session of the experiment lasted 1-1.5 hours, depending on the participant.

Stimuli

All stimuli were presented on a 32'' Cambridge Research Systems Display++ controlled by a computer running experimental code written in Matlab and PsychToolBox. The monitor primaries were calibrated with a spectroradiometer (Photo Research PR 655). The monitor was driven at a refresh rate of 120Hz and had a resolution of 1920 x 1080 pixels.

Chromaticities were defined within a version of Derrington-Krauskopf-Lennie cone-opponent color space, which represents the chromatic plane in terms of axes corresponding to variations in the LvsM cone and the SvsLM cone-antagonistic signals at constant luminance (L+M) (Derrington et al., 1984). The cone spectra were based on the Stockman-Sharpe fundamentals (Stockman & Sharpe, 2000) and the space was defined so that the gray origin corresponded to the chromaticity of illuminant D65 and a luminance of 20 cd/m^2. Units along the two axes were scaled so that a distance of 1 unit corresponded roughly to the threshold for detecting a chromatic change away from white along each axis, based on measurements from prior studies, and so that positive values along the SvsLM axis corresponded to +S increments. The equiluminant plane in the resulting space is related to the MacLeod-Boynton chromaticity diagram by the following equations:

$$\text{LvsM} = 2500 \times (l_{mb} - 0.6981) \tag{1}$$

$$\text{SvsLM} = 5000 \times (s_{mb} - 0.02065)$$

where 0.6981 and 0.02065 represent the MacLeod-Boynton L/L+M) and S/(L+M) values of the D65 gray point based on the Stockman-Sharpe fundamentals (Stockman, 2019).

The use of a monitor was necessary to allow systematic variation and control of the stimulus and experiment. However, for our study we were interested in the impact of the notch filters on naturalistic spectra, and not on the spectra generated by a typical RGB display. Monitor spectra are relatively narrowband and in particular already have lower emission at wavelengths that the notch filters are designed to block. Thus, the filters have more limited effects on colors viewed on a monitor (Somers and Bosten, 2024). To overcome this limitation, testing was performed without wearing the filters, and instead we simulated the filter effects by modeling their impact on naturalistic reflectance spectra and then presenting the monitor stimuli with the same

chromaticity. The natural spectra were based on the first 3 principal components of variation in the Munsell reflectance spectra (Cohen, 1964) when shown under D65 illumination. This transform allowed us to first specify the DKL coordinates of our stimuli and then construct the corresponding Munsell spectra (Ilic et al., 2022; Webster & Mollon, 1995). These simulated spectra were then adjusted by the transmittance of the notch filter and weighted by the LMS cones to yield the altered chromaticities. Finally, to control for changes in the mean luminance and chromaticity after filtering, the cone responses were adjusted assuming von Kries adaptation to the mean stimulus.

The panels of Figure 1 illustrate the DKL chromatic plane (left) and the transmittance of the SuperX filter (right). The effect the filtering is to increase chromatic contrast roughly along the magenta-green (45-225°) axis of the space (Figure 2).

1a.   1b.

Figure 1. (a) 90-270° and 135-325° (shown and referred to as 90° and -45°) background chromatic axes in the DKL color space. (b) Enchroma SuperX® glasses spectra transmittance (400 – 700 nm).

For the experimental stimuli, the background colors were varied along two different axes in the plane. For one, the colors varied along angles of 135-315° , a roughly bluish to yellowish-orange direction. For the second, they were confined to variations along the 90-270° SvsLM axis, which varies from a purple to yellow-green. For shorthand, we refer to these two conditions as the -45° and 90° axes respectively. The two axes were chosen to span the range of variation in the principal axes of natural daytime scenes, with the blue-yellow typical of arid and panoramic scenes while the SvsLM more typical of scenes dominated by lush foliage (Ruderman et al., 1998; Skelton et al., 2024; Webster et al., 2007; Webster & Mollon, 1997). As discussed below, the different axes also allowed us to assess how the impact of the filter on color salience depended on the background colors. For each background, the chromaticities of individual elements were selected randomly from contrasts spanning +/-80 units. The luminance of each background element was also randomly varied by +/-30% relative to the 20 cd/m^2 mean of the background, in order to mask luminance differences as a potential cue to the target (Regan et al., 1994). Target colors were instead chosen to sample 8 different color directions or hue angles, (at steps of 45° around the cone-opponent plane, Figure 1a) each shown at 4 different contrast levels ranging from 10 to 80 units in 0.3 log steps, and all at the mean luminance before the filtering. The filtering also changed the relative luminances of the target and background colors, and this effect was included in the rendered stimuli. However, these

luminance changes were less than +/-6% and thus small relative to the chromatic changes and to the random luminance variation added to the background.

Figure 2 shows the chromatic coordinates of the targets and the two backgrounds. Note that the same targets were used for both backgrounds. Note also that the targets include colors at different angles relative to each background and include colors along the background axis. The angles and contrasts of both the targets and backgrounds are altered by the filters. In the analyses below, we represent the effective contrast of the target by the distance of the target chromaticity to the background (TBD) axis. This varies from 0 for all targets along the background axis to the full nominal contrast for the targets at angles perpendicular to the axis.

2a. 2b. 2c. 2d.

Figure 2. Top : targets and background coordinates in the LvsM and SvsLM plane without (a) and with (b) the filter for the -45° background. Bottom: targets and background coordinates without (c) and with (d) the filter in the 90° background. Note that the same set of targets were used for both backgrounds, so that only the chromatic axis of the background differs between the upper and lower panels..

To display the stimuli, a variegated color background was displayed composed of 10,000 superimposed ellipses whose minor axis randomly varied from 0.2° to 0.39° of visual angle and major axis from 0.62° to 0.8°, according to a uniform distribution. The ellipses' positions and orientations were also randomly varied. The ellipses occluded those below them and the large number was to ensure the screen was fully tiled. The circular target had a diameter of 0.5° and was randomly positioned on top of the background, so that it was always fully visible. The circular shape cue meant that the target could be detected even when the color was along the background axis, which allowed observers to identify the target on all trials. Examples of the backgrounds and targets are illustrated in Figure 3.

.

3a.

3b.

3c.

3d.

Figure 3. Examples of the stimuli for -45° background without (a) and with (b) the filter simulation, and for 90° background without (c) and with (d) the filter simulation. Arrows show the location of the targets, which vary in visibility based on the color difference from the background.

Procedure

Observers viewed the display binocularly in an otherwise dark room and were seated at 120cm from the monitor, at which the full-screen background subtended 20° x 11.6°. On each trial the target was displayed at a random location on the left or right side of the display, and the participant had to identify the side by using a handheld keypad. There was no time limit to respond, and a response was required in order to advance to the next stimulus.

Each participant completed 20 repetitions for each target/background condition (one experimental block). Each session was composed of four experimental blocks (two with the filters and two without the filters) with the order of the filter condition counterbalanced and unknown to participants. For each background two sessions were completed on separate days; the whole experiment, thus, lasted 4 days for the 4 sessions, with the time of a single session

lasting between 1 and 1.5 hrs. The order of the two backgrounds was also counterbalanced between the participants. Results reported are for the aggregate data across observers: 2560 total observations per observer were recorded (1280 for each background).

Statistical Analyses

To formally assess the results, the data were analyzed in the R environment for statistical computing (R Core Team, 2024). For both backgrounds, we conducted sign tests of the median reaction times to compare the filter vs. no-filter condition results, for session 1, for session 2 and for the two sessions together, both for between-observer and within-observer comparisons. We then fitted nonlinear mixed-effects models (NLME) to account for the effects of the Target-Background Difference (TBD). We initially also analyzed the Target Contrast, which showed a linear dependence on the dependent variable after it had been transformed logarithmically and similar filter effects to the TBD. A principal component analysis of the explanatory variables, however, indicated that the TBD and Target Contrast were highly correlated. The first principal component depended on both variables with approximately equal weighting. Therefore, we opted to analyze just the TBD here.

Histograms of reaction times (RT) are typically highly positively skewed, which leads to overestimated and biased confidence intervals using standard mean and variance estimators, and thereby leading to reduced power in hypothesis testing (Zandt, 2002). We evaluated the residuals of a linear model including all variables with Box-Cox power transformations (Box & Cox, 1964) using the *boxcox* function in the **MASS** package (Venables & Ripley, 2002). The analysis suggested that the reciprocal of the RT or speed would stabilize the variance and approximately normalize the distribution of responses, considered desirable properties of a good measurement scale (McCullagh, 2019). Simulations have indicated that the inverse transform minimizes the effects of outliers (Ratcliff, 1993), An example of the extreme skewness in the RT distribution for one observer for the 90° background condition, both filter conditions confounded, is shown in Fig 4a. The same data are shown in Fig 4b with the histogram now plotted as a function of the speed or reciprocal reaction time. As the results were similar for all subjects for both background conditions, the speed was analyzed as the dependent variable in subsequent analyses. The exception is the initial analysis of medians, which is a robust (but less powerful) statistic that depends much less on the distribution shape.

Figure 4. (a) Histogram of frequencies of reaction times in seconds for observer S7 for the 90° background condition. (b) Histogram of frequencies of reciprocal reaction times in inverse seconds for the same data in part a).

Nonlinear mixed-effects models were fit to the transformed data using the nlme function in the **nlme** package (Pinheiro et al., 2025; Pinheiro & Bates, 2000) to model the effect of the filter as a function of TBD and Target Angle. Initial evaluations indicated that the results did not depend on Target Angle and it was excluded from the main analyses. A Michaelis-Menten function was found to provide a good description of the dependence of the speed on TBD. In preliminary analyses a power law was found to provide a similar description. However, as the best fits with the Michaelis-Menten function yielded a smaller AIC, we used this model for subsequent analyses. Random effects and then fixed effects were evaluated with χ^2 statistics obtained from nested likelihood ratio tests. The full NLME initially fit to the data is formalized as

$$S=\left(R_m+R_f+r_o\right)\frac{|\mathrm{TBD}|}{|\mathrm{TBD}|+\left(\varsigma+\varsigma_f+s_o\right)}+\epsilon,\qquad(2)$$

where

$$\epsilon \sim N\left(0,\sigma^2\right)$$

$$r_o \sim N\left(0,\sigma_{r_o}^2\right)$$

$$s_o \sim N\left(0,\sigma_{s_o}^2\right)$$

and where S is the speed (in reciprocal seconds), TBD the target-background distance (as defined below), the R and ς parameters indicating the maximum speed and TBD at 50% maximum response (semi-saturation constant) for the no filter condition, respectively. R_f and ς_f are the respective fixed effects of the filter on the two preceding parameters and r_o and s_o are

random observer predictions of each of the preceding parameters, each assumed to be normally distributed with its own variance. The parameter ϵ is the residual random variation that is assumed to be normally distributed with mean zero and variance σ^2. Thus, the full model has 3 variance terms and the correlation between the two parameter random effects to estimate in addition to the fixed effects of the no filter and change due to the filter on each of the parameters. Nested likelihood tests were performed initially to evaluate the significance of the random effect terms. Subsequently, the fixed effects of adding the filter were evaluated for each parameter, R_f and ς_f. We set the level of significance at α = 0.05 and report statistical analyses only for correct responses, which were overall very high across conditions and observers (99.1%). No outliers were removed.

Results

Figure 5 shows the reaction times for detecting the targets under the 4 conditions tested: on the two different backgrounds, and with or without the filter. As illustrated in Figure 2, the RTs are plotted as a function of the TBD, which is defined by

$$\text{TBD} = T_c \sin(T_a - B_a) \tag{3}$$

where T_c equals the target contrast and T_a and B_a are the target and background chromatic angles, respectively. As in many previous studies, RTs decrease asymptotically as the separation between the target and background or distractor colors increases (Bauer et al., 1996; Manalansan et al., 2025; McDermott et al., 2010; McKeefry et al., 2003; Nagy & Sanchez, 1990; Vanston et al., 2021; Webster & Mollon, 1994). In our case, these effects are bounded by roughly a parallel or "pop out" search when the target is very different from the background, to a serial search based instead on the spatial cue alone as the target color becomes indistinguishable from the background (McDermott et al., 2010).

5a.   5b.

5a.   5b.

Figure 5.

Reaction times plotted as a function of the target-to-background chromatic difference for the -45° (a) and 90° (b) backgrounds. Blue asterisks represent the average reaction times in the no-filter condition, while red circles denote the corresponding values for the filter condition. Specifically, each data point reflects the mean across observers of the median reaction time calculated within each run, where the median is derived from five repeated reaction time measurements per observer. This approach ensures a robust

central tendency measure by first summarizing individual trial variability within runs before aggregating across observers.

The questions of interest were whether the search times differed for the filter vs. no-filter condition, and whether this depended on the chromatic properties of the backgrounds and targets. To visualize these differences, in Figure 6 the scatterplots directly compare the average reaction times for each target under the filtered or unfiltered condition. For the -45° background, search times appeared consistently faster with the simulated filters. Alternatively, there is little evident difference for the 90° background. These patterns are confirmed statistically below. Note also that overall, the search times were substantially faster on the blue-yellow background, even though the two backgrounds were matched for nominal chromatic contrast. Below we consider the implications of this difference and how it impacted the observed effects of the filter.

6a.   6b.

Figure 6. Average reaction times for the same targets compared for the filtered vs. unfiltered conditions. (a) -45° background; (b) 90° background. Symbols plot the RTs for all responses (blue diamonds) or correct responses only (red circles).

-45° Background

Within-subjects comparisons for the -45° background revealed a global trend of the filter effect, without any distinction between the first and the second session: speed was always higher in the filter condition.   Sign tests for the between-subjects comparison showed statistically significant results across all the data: the filter had a greater effect both in the individual sessions (2- tailed: Session 1 - p = 0.002, Session 2 – p = 0.02) and in the two sessions together (2- tailed: p = 0.002). The assessment of the individual sessions was done to confirm that the differences were stable across the sessions. Participants were, thus, always faster in the experimental blocks with the filters. The average filter vs. no filter difference for correct trials in session 1 was -207 ms, -134 ms in session 2, and -171 ms in the two sessions together. The general accuracy was 99.4%, indicating that global performance was highly accurate.

Fitting the data to the NLME of Equation 2 proceeded in two steps. First, the random effects were evaluated to simplify them, if necessary, and then the fixed effects. Tests of the random effects indicated that the correlation between the two random effects was not significant (LRT: $\chi^2(1)$= 0.001, p = 0.97) permitting subsequent analyses to be performed with a diagonal covariance matrix. Random variances associated with the maximum value, $\sigma^2_{r_o}$ and the semi-saturation constant, $\sigma^2_{s_o}$, however, were required as leaving either of them out yielded a significant difference from the model containing both ($\sigma^2_{r_o}$, LRT: $\chi^2(1)$= 334.4, p << 0.001; $\sigma^2_{s_o}$, $\chi^2(1)$= 8.9, p = 0.003).

With the random structure of the model set, the fixed effects were evaluated. A filter effect on the maximum value parameter was not significant (LRT: $\chi^2(1)$ = 0.339, p = 0.56), but the filter effect on the semi-saturation constant was (LRT: $\chi^2(1)$ = 250.0, p << 0.001). A summary of the fixed and random effects parameter estimates are given in column 2 of Table 1 with their 95% confidence intervals in column 3.

Table 1. Estimates of fitted parameters and standard errors (SE) from best fitting NLME models for the -45° and 90° deg data sets.

	-45		90	
	Estimate	95% Conf. Int.	Estimate	95% Conf. Int.
R_m	2.212	(2.072, 2.351)	2.336	(2.183, 2.488)
ς	8.996	(8.480, 9.512)	15.303	(14.346, 16.259)
ς_f	-3.003	(-3.305, -2.700)	-0.710	(-1.220, -0.201)
$\sigma^2_{r_o}$	0.049	(0.020, 0.119)	0.057	(0.023, 0.141)
$\sigma^2_{s_o}$	0.329	(0.089, 1.216)	1.184	(0.814, 1.723)
σ^2	0.168	(0.163, 0.173)	0.181	(0.177, 0.187)

Figure 7a shows the average speed as a function of TBD for no filter (blue) and filter (orange) conditions. The error bars indicate 95% confidence intervals. The curves are based on the estimated fixed effects parameters from the best fit model for the no filter (solid) and filter (dashed) conditions based on the fixed effect parameters in Table 1, and indicate the generally faster responses to the filter condition. Note the similarity in the size of the confidence intervals across TBD, supporting the transformation of RT to speed to stabilize the variance. (Compare with the larger variation in RTs as a function of TBD in Figures 5). Concurrently, the filter had an approximately constant effect on all targets, independently of their angle, showing faster detection on experimental blocks with the filter on average (Figure 7b).

7a.   7b.

Figure 7. (a) Average speed (1/RT) with and without the filter as a function of the TBD. The curves indicate the fixed-effect model fit using the Nonlinear Mixed-Effect Model. Reaction times are indicated on the right ordinate. The Michaelis-Menten equation for the curves is indicated as an inset. Error bars are 95% confidence intervals. (b) Average speed as a function of Target Angle with the corresponding Reaction Times (in s) indicated on the right ordinate. The curves are local regression curves fit to the data. Error bars are 95% confidence intervals.

90° Background

Sign tests for the comparison between the filter and no-filter runs were not statistically significant for either individual session or the two sessions together (2-tailed: p=0.75, in all 3 cases). The average difference in session 1 was 81.4 ms, -33.2 ms in session 2, and 24.1 ms in the two sessions together. The global accuracy was 98.8%, similar to the -45° condition.

As with the -45° background, speed again varied systematically as a function of TBD with speed increasing with the magnitude of TBD (Fig 8a) and again was relatively independent of Target Angle (Fig. 8b). The differences between the no filter and filter conditions appear less obvious on this background.

8a.   8b.

Figure 8. (a) Speed (1/RT) with and without the filter as a function of the TBD. Reaction times are indicated on the right ordinate. The points indicate the means with 95% confidence intervals computed across subjects and color coded the same as in Figure 7. The curves indicate a Michaelis-Menten function (inset equation) fit with a nonlinear mixed-effects model for the filter (dashed) and no filter (solid) conditions. Error bars correspond to 95% confidence intervals. (b) Speed as a function of the Target

Angle (with reaction times indicated on the right ordinate). The points indicate the means and 95% confidence intervals computed across subjects. The curves are local regression fit (loess) to indicate the general trends through points.

We again fit the data with the NLME using Eq 2 to describe the dependence of the speed on TBD. As before we began by testing the significance of the random effect variance terms. As at -45°, the correlation between the two random effects for the model parameters was not significant (LRT: $\chi^2(1) = 0.898$, p = 0.34). Also, similarly to the -45° condition, both random effect variances were required to model the individual variability of the parameter estimates (LRT: $\sigma^2_{r_o}$ $\chi^2(1) = 159.3$, $p << 0.001$; $\sigma^2_{s_o}$, $\chi^2(1) = 5.52$, $p = 0.02$). With the random effects structure established, we tested the fixed effects parameters. As for the -45° background, the estimate of the difference in values for maximum response, R_f, did not differ significantly from 0 (LRT: $\chi^2(1) = 1.862$, p = 0.17) but the difference in values for the semi-saturation constant, ς_f did (0 (LRT: $\chi^2(1) = 7.53$, p = 0.006). The parameter estimates for the fixed and random effects are indicated in column 4 Table 1 with their 95% confidence intervals in column 5. The fixed effect predictions of the two models are shown as solid and dashed curves in Fig 8a and suggest that while the filter effect on the semi-saturation constant is significant, it is very slight.

Analyses of the search differences on the two backgrounds

Why was the filter enhancement on color search on the bluish-yellowish background more effective than on the SvsLM background? The basis for this difference is not readily evident in the context of the changes the filters induced in the cone opponent contrasts. We assumed that the salience of the targets was largely captured by the chromatic distance of the target from the background axis. As Figure 2 showed, filtering expanded chromatic contrasts along the magenta-lime direction. This left the -45° background axis largely unaffected, while rotating the 90° background by roughly 20°. However, the TBD remained similar across the two backgrounds, and in particular, the increase in distances with the filtering was also similar for the two conditions, as illustrated in Figure 9. Moreover, while the 90° axis was rotated by filtering, the overall contrast remained similar, so that the chromatic variance along each background remained largely the same. Thus, based on the net cone-opponent contrasts, the effects of the filters on color search might be expected to be similar, in contrast to the differences we observed.

Figure 9. Chromatic difference between the targets and background axes within the cone opponent plane, for the -45° background (red circles) or the 90° background (blue circles).

However, a clue to the basis for this difference is seen in average search times in Figures 5 and 6, which again show that locating targets on the -45° background was substantially faster than on the 90° background. This effect is consistent with previous results of McDermott et al. and Manalansan et al, who found that chromatic targets are easier to find on bluish-yellowish (or warm-cool) backgrounds than on orthogonal (magenta-greenish) backgrounds (Manalansan et al.,2025; McDermott & Webster, 2012). In turn, these effects are consistent with a large number of findings demonstrating weaker sensitivity to the bluish-yellowish axis of color space (Webster, 2020).

To explore this for our results, we converted the coordinates of the stimuli from the cone-opponent axes to the corresponding values in CIELAB, a "perceptually uniform" space designed to approximate perceptual rather than chromatic contrast differences. A circle in cone-opponent space, representing equal contrasts at different hue angles, warps into an elongated contour in LAB (McDermott & Webster, 2012). Manalansan et al. showed that the LAB contour minima correspond closely to the stimuli associated with warm and cool (orange and teal), while the maxima of the contour align with the categorical boundaries between warm and cool (Manalansan, Whitehead and Webster, 2025). In CIELAB, distances from gray approximate the chroma or perceptual strength of the color. The distortions thus imply that for the same cone-opponent contrasts, the perceptual strength is weakest along the warm-cool dimension and strongest along roughly orthogonal directions.

Figure 10a shows a similar analysis for our stimulus set and represents the LAB values for the target and background colors depicted in Figure 2 within the cone-opponent space. (Note that in addition to the warping there is an inversion of the blue-yellow axis between the spaces.) Within this perceptual metric, the 90° axis is strongly expanded relative to the -45° axis. Moreover, the distances between the targets and background are markedly larger for the -45° background (Fig. 10b). These asymmetries provide a plausible basis for the large asymmetries in search times on the two backgrounds.

Figure 10. (a) coordinates of the targets (circles) and backgrounds (-45° - solid lines; 90° – dashed lines) replotted in CIELAB. Black circles and lines show values without the filter. Red circles and lines show values with the filter. Labels show the original angles of the background axes in the cone-opponent space. (b) Chromatic difference between the targets and background axes within the perceptually uniform (CIELAB) space; -45° background (red circles), 90° background (blue circles).

When the filter is applied, there is again little effect on the -45° background colors, while other stimuli are biased away from this axis. Thus, the faster search times with the filter are consistent with the increased TBD for the stimuli. In contrast, the filtering rotated the 90° background toward an axis of even greater perceptual sensitivity; and while the TBD was also amplified, the enhancements relative to the background were weaker in the perceptually-based units. Thus, in this case the advantages of the filter for highlighting the targets are partially offset by the enhancement of the background. Finally, this analysis only considers the TBD of the stimuli, and not the variance of the background. However, the contrast or range of the background colors also influences the salience of the targets, which for example are all much easier to detect on an achromatic background (McDermott et al., 2010). As noted, this variance is much higher for the 90° background, and modestly increased further by the filtering, and this is potentially an additional factor accounting for why filtering did not facilitate color search for this background for the specific conditions we tested.

Discussion

To summarize, we found that the increases in chroma afforded by notch filtering led to predictable effects on color salience as measured by a color search task, and that the effects also depended in predictable ways on how the filter altered the chromatic properties of both the background and targets. In particular, for the conditions we tested, search times were significantly faster when the targets were shown on the bluish-yellowish background (-45°). In this case, the filter had only a marginal effect on the background chromaticities, while boosting other chromatic directions relative to the background axis, thus increasing the colorimetric distance from the background. On the other hand, for the purple to yellow-green background (90°), the targets were not easier to find even though the chromatic differences to the

background were again increased. Notably, the basis for the lack of improvement on this background was only apparent when the stimuli were analyzed within a perceptual rather than cone-opponent space. While the backgrounds were equated in terms of cone-opponent contrasts, by the perceptual metric (CIELAB), the 90° deg background had a higher perceptual gamut than the blue-yellow background (Manalansan, Whitehead, et al., 2025; McDermott et al., 2010), and the filtering was less effective at selectively boosting the perceptual distances of the targets relative to this background.

Search times on the -45° background were also substantially faster than the 90° background, both with and without the filtering. This difference is again consistent with previous studies of color search using similar paradigms (Manalansan, Simoncelli, et al., 2025; McDermott et al., 2010). The weaker sensitivity and salience for the bluish-yellowish dimension has also been documented with a variety of other techniques e.g. (Bosten et al., 2015; Boynton et al., 1986; Goddard et al., 2010; Juricevic et al., 2010; Skelton et al., 2023). A common account of this bias is that the world varies more along the bluish-yellowish dimension, so that vision is more strongly adapted or desensitized to this axis (Juricevic & Webster, 2009; Skelton et al., 2024; Webster, 2020; Webster & Mollon, 1997). In this regard, adaptation also alters the relative salience of colors, but in a complementary way to the filtering we examined. Specifically, adaptation changes salience by reducing sensitivity to the background while "sparing" sensitivity to novel stimuli (McDermott et al., 2010; Wissig et al., 2013). In contrast, filtering is predicted to increase salience by enhancing the color of the target relative to the background (Somers & Bosten, 2024; Werner et al., 2020). In both cases, the changes in target salience should largely depend on how selective the effects are for the targets vs. the backgrounds. When the targets fall within the background distribution, salience may be relatively unaffected because both stimuli are biased in similar ways. Such predictions can also be readily extended to other background directions. For example, for the specific filter we examined, filtering the spectra should reduce color search on magenta-lime (e.g., 45-225°) backgrounds, where the effect would instead be to enhance the background colors relative to the targets. However, such color backgrounds are atypical, at least for natural environments.

As noted in the Introduction, the two backgrounds we examined were chosen to roughly span the range of dominant chromatic axes of many natural scenes – and in particular to capture the color signatures of lush or arid environments (Nascimento et al., 2021; Ruderman et al., 1998; Skelton et al., 2024; Webster & Mollon, 1997). However, in other ways these backgrounds were also highly unnatural: the colors were confined to a single hue angle and sampled the axis uniformly, and had high variance. In contrast, natural color distributions are more irregular and clustered (e.g., between regions of sky and earth), and often have a lower variance with higher proportion of low-contrast levels (Laughlin, 1981; Nascimento et al., 2002; von der Twer & MacLeod, 2001). Moreover, in real scenes colors are usually associated with objects and surfaces and thus have very different spatial characteristics from the uniform textures we used. A further limitation is again that we simulated the filters rather than directly wearing them, and did not evaluate the effects the filters have on the mean light level (Somers & Bosten, 2024). The simulations did, however, capture the pattern of cone excitations predicted if observers in each case were adapted to the mean level of the display. This light and chromatic adaptation should largely offset the effects of absolute intensity differences but does not fully address potential changes in chromatic discrimination at different light levels.

The highly constrained conditions we tested are useful for controlling the stimuli and revealing the principles affecting the search performance, but are unlikely to fully generalize to realistic viewing environments. As one example, we used the target to background distance (TBD) as a simple predictor for salience. In real scenes, however, the background distribution will not be confined to a single axis, so that the target salience is more likely to be a function of the signal to noise ratio. By this metric, even targets that lie along the dominant axis can (and do) become salient as they fall outside the background gamut (McDermott et al., 2010; Manalansan, Simoncelli, et al., 2025). In such cases, an important question for understanding the effect of the filters is how amplifying both background and target (while for example maintaining the contrast ratio) might impact the color search. Similarly, in natural scenes, search is strongly affected by the specific task demands and expectations about the potential presence, location, or nature of specific targets, and thus factors such as expectations and familiarity, both with objects and their colors, will also impact the salience (Võ, 2021; Wolfe & Horowitz, 2017). In future work we plan to evaluate the contrast enhancements under these more naturalistic conditions.

A further factor determining salience is how the color signals are represented in the visual system (at least at the level limiting performance). For example, as our results revealed, the different effects of the filtering could not be explained by the cone-opponent contrasts in the stimuli, and instead depended on sensitivity biases that were better captured by a color space that approximates perceptual uniformity. This bias itself argues against stages of color coding where the LvsM and SvsLM cardinal axes are independent, because the blue-yellow bias (i.e. at -45°) occurs relative to stimuli that are matched for the cardinal axis responses (i.e. for axes at +45°, where the LvsM and SvsLM signals are equal but combined in opposite phase) (Goddard et al., 2010). Finally, predicting the salience also depends on the underlying structure of the color coding. Our TBD metric assumed that backgrounds had no effect on orthogonal target directions. However, at cortical stages where color is encoded by multiple hue selective mechanisms (Gegenfurtner, 2003), these orthogonal signals can interact. For example, the broad color tuning at early cortical levels means that while cells tuned to 0° or 90° do not "see" both axes, cells tuned to intermediate axes (e.g. 45°) respond and contribute to both (Webster & Mollon, 1994). Thus, it is an oversimplification to assume that there is no effect of the backgrounds on the orthogonal targets. Finally, we have considered only the changes in chromatic contrast and perceived hue and chroma. However, increases in saturation are also associated with changes in perceived brightness, which could serve as a further potential cue for salience (Corney et al., 2009). A more formal model of color search would need to incorporate these different factors.

Our results have a number of implications for the potential to enhance color vision through spectral filtering. First, these results are specific to the specific filter design we evaluated, which again was developed for normal trichromatic vision. In other designs the notch filtering has been tailored to enhance chromatic contrasts for different types of anomalous trichromacy (i.e., deuteranomalous or protanomalous) (Somers & Bosten, 2024). The current study tested only normal color vision, but the color search task could be readily adopted for assessing the impact of the filters for color deficiencies. An advantage of the paradigm is that it can evaluate a wide range of both threshold and suprathreshold colors while exploiting a common and naturalistic visual task. The present analyses and measurements are also relevant to understanding consequences of varying the illumination rather than filtering. As noted, wide gamut lighting with narrowband primaries applies similar principles to enhance chromatic contrasts, and thus

predicts similar consequences for color salience (Ilic et al., 2022), for both normal and color-anomalous vision (Tamura, Okamoto et al., 2017ì5; Tamura, Okamoto et al., 2017; Joyce et al., 2021). Finally, filters (or lighting) could also be engineered for different environments or goals rather than observers, depending on the availability of appropriate dyes. For example, a filter for normal color vision could in principle be designed to minimize the changes specifically along the 90° axis rather than the bluish-yellow axis, as in the model we tested. This could selectively expand chromatic contrasts relative to the SvsLM opponent axis and thus better isolate the signals in the LvsM axis. These LvsM signals are carried by the "modern subsystem" of primate trichromacy and may have evolved specifically to make ripening fruit more distinguishable from the foliage background, by restricting the chromatic noise from the background of foliage to primarily SvsLM variations (Osorio & Vorobyev, 1996; Regan et al., 2001; Sumner & Mollon, 2000). As the two backgrounds we tested again illustrate, natural (as well as carpentered) environments do vary substantially across both location and time (e.g., seasons (Webster et al., 2007)). Adaptation is thought to help tune vision for specific environmental contexts (Webster, 2015). In a similar way, notch filtering could potentially be optimized for specific environments or tasks.

Acknowledgments

Supported by DOD W911NF-24-1-0328 (DM, MW, KK), NIH EY010834 and NIH P30 GM145646 (MW) and LABEX CORTEX (ANR- 11-LABX-0042) of the Université de Lyon (ANR-11-IDEX-0007) from the French National Research Agency (KK).

References

Bauer, B., Jolicoeur, P., & Cowan, W. B. (1996). Visual search for colour targets that are or are not linearly separable from distractors. *Vision Res*, *36*(10), 1439-1465. https://doi.org/10.1016/0042-6989(95)00207-3

Bosten, J. M., Beer, R. D., & MacLeod, D. I. (2015). What is white? *J Vis*, *15*(16), 5. https://doi.org/10.1167/15.16.5

Box, G. E., & Cox, D. R. (1964). An analysis of transformations. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, *262*(2), 211-243.

Boynton, R. M., Nagy, A. L., & Eskew, R. T. (1986). Similarity of normalized discrimination ellipses in the constant-luminance chromaticity plane. *Perception*, *15*(6), 755-763. https://doi.org/10.1068/p150755

Clark, J. H. (1924). The Ishihara Test for Color Blindness. *American Journal of Physiological Optics*, 5, 269–276.

Cohen, J. (1964). Dependency of the spectral reflectance curves of the Munsell color chips. *Psychonomic Science*, *1*(1), 369-370.

Corney, D., Haynes, J. D., Rees, G., & Lotto, R. B. (2009). The brightness of colour. *PLoS One*, *4*(3), e5091. https://doi.org/10.1371/journal.pone.0005091

D'Zmura, M. (1991). Color in visual search. *Vision Res*, *31*(6), 951-966. https://doi.org/10.1016/0042-6989(91)90203-h

D'Zmura, M. and Mangalick, A., 1994. Detection of contrary chromatic change. *Journal of the Optical Society of America A*, *11*(2), pp.543-546.

Derrington, A. M., Krauskopf, J., & Lennie, P. (1984). Chromatic mechanisms in lateral geniculate nucleus of macaque. *J Physiol*, *357*, 241-265. https://doi.org/10.1113/jphysiol.1984.sp015499

Feng, X., Xu, W., Han, Q., & Zhang, S. (2016). LED light with enhanced color saturation and improved white light perception. *Optics express*, *24*(1), 573-585.

Foster, D. H. (2011). Color constancy. *Vision Res*, *51*(7), 674-700. https://doi.org/10.1016/j.visres.2010.09.006

Gegenfurtner, K. R. (2003). Cortical mechanisms of colour vision. *Nat Rev Neurosci*, *4*(7), 563-572. https://doi.org/10.1038/nrn1138

Goddard, E., Mannion, D. J., McDonald, J. S., Solomon, S. G., & Clifford, C. W. (2010). Combination of subcortical color channels in human visual cortex. *J Vis*, *10*(5), 25. https://doi.org/10.1167/10.5.25

Gómez-Robledo, L., Valero, E. M., Huertas, R., Martínez-Domingo, M. A., & Hernández-Andrés, J. (2018). Do EnChroma glasses improve color vision for colorblind subjects? *Opt Express*, *26*(22), 28693-28703. https://doi.org/10.1364/OE.26.028693

Ilic, I., Lee, K. R., Mizokami, Y., Whitehead, L., & Webster, M. A. (2022). Adapting to an enhanced color gamut - implications for color vision and color deficiencies. *Opt Express*, *30*(12), 20999-21015. https://doi.org/10.1364/OE.456067

Joyce, D. S., Whitehead, L. A., Webster, M. A. (4 August 2021). Wide gamut lighting and color contrast in anomalous trichromacy. Proc. SPIE 11814, *Current Developments in Lens Design and Optical Engineering XXII, 118140D*. https://doi.org/10.1117/12.2594334

Juricevic, I., Land, L., Wilkins, A., & Webster, M. A. (2010). Visual discomfort and natural image statistics. *Perception*, *39*(7), 884-899. https://doi.org/10.1068/p6656

Juricevic, I., & Webster, M. A. (2009). Variations in normal color vision. V. Simulations of adaptation to natural color environments. *Vis Neurosci*, *26*(1), 133-145. https://doi.org/10.1017/S0952523808080942

Laughlin, S. (1981). A simple coding procedure enhances a neuron's information capacity. *Zeitschrift für Naturforschung c ,*, *36*(9-10).

Lynch, H., & Ng, J. S. (2024).. Hue Discrimination in Patients with Red/Green Deficiency Using Multi-Notch Filters. O*ptometry & Visual Performance* , 1*2* (4). *Optometry and Visual Performance*, *12*(4).

Manalansan, J., Simoncelli, C., & Webster, M. A. (2025). Visual search for warm and cool colors. *J Opt Soc Am A Opt Image Sci Vis*, *42*(5), B148-B154. https://doi.org/10.1364/JOSAA.545307

Manalansan, J., Whitehead, L. A., & Webster, M. A. (2025). Warm versus cool colors and their relation to color perception. *J Vis*, *25*(4), 13. https://doi.org/10.1167/jov.25.4.13

Marques, D. N., Gomes, A. E., Linhares, J. M. M., & Nascimento, S. M. C. (2023). Discrimination of natural colors in anomalous trichromacy and the effects of EnChroma and Vino filters. *Opt Express*, *31*(11), 18075-18087. https://doi.org/10.1364/OE.451326

McCullagh, P. Nelder J. A. (1989). Generalized Linear Models.Chapman & Hall/CRC, Boca Raton.

McDermott, K. C., Malkoc, G., Mulligan, J. B., & Webster, M. A. (2010). Adaptation and visual salience. *J Vis*, *10*(13), 17. https://doi.org/10.1167/10.13.17

McDermott, K. C., & Webster, M. A. (2012). Uniform color spaces and natural image statistics. *J Opt Soc Am A Opt Image Sci Vis*, *29*(2), A182-187. https://doi.org/10.1364/JOSAA.29.00A182

McKeefry, D. J., Parry, N. R., & Murray, I. J. (2003). Simple reaction times in color space: the influence of chromaticity, contrast, and cone opponency. *Invest Ophthalmol Vis Sci*, *44*(5), 2267-2276. https://doi.org/10.1167/iovs.02-0772

Nagy, A. L., & Sanchez, R. R. (1990). Critical color differences determined with a visual search task. *J Opt Soc Am A*, *7*(7), 1209-1217. https://doi.org/10.1364/josaa.7.001209

Moreland, J. D., Dain, S. J., Cheung, V., & Westland, S. (2022). A model for assessing the efficacy of colour vision aids. *Optics Express*, *30*(15), 27903-27911.

Nascimento, S. M., Ferreira, F. P., & Foster, D. H. (2002). Statistics of spatial cone-excitation ratios in natural scenes. *J Opt Soc Am A Opt Image Sci Vis*, *19*(8), 1484-1490. https://doi.org/10.1364/josaa.19.001484

Nascimento, S. M. C., & Foster, D. H. (2022). Information gains from commercial spectral filters in anomalous trichromacy. *Opt Express*, *30*(10), 16883-16895. https://doi.org/10.1364/OE.451407

Nascimento, S. M. C., Marit Albers, A., & Gegenfurtner, K. R. (2021). Naturalness and aesthetics of colors - Preference for color compositions perceived as natural. *Vision Res*, *185*, 98-110. https://doi.org/10.1016/j.visres.2021.03.010

Osorio, D., & Vorobyev, M. (1996). Colour vision as an adaptation to frugivory in primates. *Proc Biol Sci*, *263*(1370), 593-599. https://doi.org/10.1098/rspb.1996.0089

Patterson, E. J., Mastey, R. R., Kuchenbecker, J. A., Rowlan, J., Neitz, J., Neitz, M., & Carroll, J. (2022). Effects of color-enhancing glasses on color vision in congenital red-green color deficiencies. *Opt Express*, *30*(17), 31182-31194. https://doi.org/10.1364/OE.451295

Pattie, C., Aston, S., & Jordan, G. (2022). Do EnChroma glasses improve performance on clinical tests for red-green color deficiencies? *Opt Express*, *30*(18), 31872-31888. https://doi.org/10.1364/OE.456426

Pinheiro, C., Bates, D., & R Core Team. (2025). nlme: Linear and Nonlinear Mixed   Effects Models,   R package version 3.1-168. doi:10.32614/CRAN.package.nlme <https://doi.org/10.32614/CRAN.package.nlme>

Pinheiro, J. C., & Bates, D. M. (2000). *Mixed-Effects Models in S and S-PLUS*. Springer New York.

R Core Team (2025). _R: A Language and Environment for Statistical Computing_. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Rabin, J., Silva, F., Trevino, N., Gillentine, H., Li, L., Inclan, L.,…Vo, H. (2022). Performance enhancement in color deficiency with color-correcting lenses. *Eye (Lond)*, *36*(7), 1502-1503. https://doi.org/10.1038/s41433-021-01924-0

Ratcliff, R. (1993). Methods for dealing with reaction time outliers. *Psychol Bull*, *114*(3), 510-532. https://doi.org/10.1037/0033-2909.114.3.510

Regan, B. C., Julliot, C., Simmen, B., Viénot, F., Charles-Dominique, P., & Mollon, J. D. (2001). Fruits, foliage and the evolution of primate colour vision. *Philos Trans R Soc Lond B Biol Sci*, *356*(1407), 229-283. https://doi.org/10.1098/rstb.2000.0773

Regan, B. C., Reffin, J. P., & Mollon, J. D. (1994). Luminance noise and the rapid determination of discrimination ellipses in colour deficiency. *Vision Res*, *34*(10), 1279-1299. https://doi.org/10.1016/0042-6989(94)90203-8

Ruderman, D. L., Cronin, T. W., & Chiao, C. C. (1998). Statistics of cone responses to natural images: implications for visual coding. *Journal of trhe Optical Society of America A*, *15*(8), 2036-2045.

Skelton, A. E., Franklin, A., & Bosten, J. M. (2023). Colour vision is aligned with natural scene statistics at 4 months of age. *Dev Sci*, *26*(6), e13402. https://doi.org/10.1111/desc.13402

Skelton, A. E., Maule, J., Floyd, S., Wozniak, B., Majid, A., Bosten, J. M., & Franklin, A. (2024). Effects of visual diet on colour discrimination and preference. *Proc Biol Sci*, *291*(2031), 20240909. https://doi.org/10.1098/rspb.2024.0909

Somers, L. P., & Bosten, J. M. (2024). Predicted effectiveness of EnChroma multi-notch filters for enhancing color perception in anomalous trichromats. *Vision Res*, *218*, 108381. https://doi.org/10.1016/j.visres.2024.108381

Somers, L. P., Franklin, A., & Bosten, J. M. (2024). Empirical tests of the effectiveness of EnChroma multi-notch filters for enhancing color vision in deuteranomaly. *Vision Res*, *218*, 108390. https://doi.org/10.1016/j.visres.2024.108390

Stockman, A. (2019). Cone fundamentals and CIE standards. *Current Opinion in Behavioral Sciences*, *30*, 87-93.

Stockman, A., & Sharpe, L. T. (2000). The spectral sensitivities of the middle- and long-wavelength-sensitive cones derived from measurements in observers of known genotype. *Vision Res*, *40*(13), 1711-1737. https://doi.org/10.1016/s0042-6989(00)00021-3

Sumner, P., & Mollon, J. D. (2000). Catarrhine photopigments are optimized for detecting targets against a foliage background. *J Exp Biol*, *203*(Pt 13), 1963-1986. https://doi.org/10.1242/jeb.203.13.1963

Tamura, S., Okamoto, Y., Nakagawa, S., Sakamoto, T., Ando, M. and Shigeri, Y. (2017). Light wavelengths of LEDs to improve the color discrimination in Ishihara test and Farnsworth Panel D-15 test for deutans. *Color Res. Appl., 42: 424-430*. https://doi.org/10.1002/col.22106.

Tamura, S., Okamoto, Y., Nakagawa, S., Sakamoto, T. and Shigeri, Y. (2015). Practical color barrier-free illumination for deuteranopia using LEDs. *Color Res. Appl., 40: 218-223.* https://doi.org/10.1002/col.21884

Vanston, J. E., Tregillus, K. E. M., Webster, M. A., & Crognale, M. A. (2021). Task-dependent contrast gain in anomalous trichromats. *Vision Res*, *184*, 14-22. https://doi.org/10.1016/j.visres.2021.02.003

Venables, W. N., & Ripley, B. D. (2002). Random and mixed effects. In *Modern applied statistics with S-PLUS* (pp. 297-322). Springer New York.

Venables, W. N. & Ripley, B. D. (2002). Modern Applied Statistics with S. Fourth Edition. Springer, New York. ISBN 0-387-95457-0

von der Twer, T., & MacLeod, D. I. (2001). Optimal nonlinear codes for the perception of natural colours. *Network*, *12*(3), 395-407.

Võ, M. L. (2021). The meaning and structure of scenes. *Vision Res*, *181*, 10-20. https://doi.org/10.1016/j.visres.2020.11.003

Webster, M. A. (2015). Visual Adaptation. *Annu Rev Vis Sci*, *1*, 547-567. https://doi.org/10.1146/annurev-vision-082114-035509

Webster, M. A. (2020). The Verriest Lecture: Adventures in blue and yellow. *J Opt Soc Am A Opt Image Sci Vis*, *37*(4), V1-V14. https://doi.org/10.1364/JOSAA.383625

Webster, M. A., Mizokami, Y., & Webster, S. M. (2007). Seasonal variations in the color statistics of natural images. *Network*, *18*(3), 213-233. https://doi.org/10.1080/09548980701654405

Webster, M. A., & Mollon, J. D. (1994). The influence of contrast adaptation on color appearance. *Vision Res*, *34*(15), 1993-2020. https://doi.org/10.1016/0042-6989(94)90028-0

Webster, M. A., & Mollon, J. D. (1995). Colour constancy influenced by contrast adaptation. *Nature*, *373*(6516), 694-698. https://doi.org/10.1038/373694a0

Webster, M. A., & Mollon, J. D. (1997). Adaptation and the color statistics of natural images. *Vision Res*, *37*(23), 3283-3298. https://doi.org/10.1016/s0042-6989(97)00125-9

Wen, Z., Zhou, Z., Liu, H., Wang, Z., Li, X., Fang, F., ... & Sun, X. W. (2021). Color revolution: Toward ultra-wide color gamut displays. *Journal of Physics D: Applied Physics*, *54*(21), 213002.

Werner, J. S., Marsh-Armstrong, B., & Knoblauch, K. (2020). Adaptive Changes in Color Vision from Long-Term Filter Usage in Anomalous but Not Normal Trichromacy. *Curr Biol*, *30*(15), 3011-3015.e3014. https://doi.org/10.1016/j.cub.2020.05.054

Wissig, S. C., Patterson, C. A., & Kohn, A. (2013). Adaptation improves performance on a visual search task. *J Vis*, *13*(2), 6. https://doi.org/10.1167/13.2.6

Wolfe, J. M. (2020). Visual Search: How Do We Find What We Are Looking For? *Annu Rev Vis Sci*, *6*, 539-562. https://doi.org/10.1146/annurev-vision-091718-015048

Wolfe, J. M., & Horowitz, T. S. (2017). Five Factors that Guide Attention in Visual Search. *Nat Hum Behav*, *1*(3). https://doi.org/10.1038/s41562-017-0058

Zandt, T. V. (2002). Analysis of response time distributions. In H. Pashler (Ed.), *Steven's Handbook of Experimental Psychology*.

Zhu, R., Luo, Z., Chen, H., Dong, Y., & Wu, S. T. (2015). Realizing Rec. 2020 color gamut with quantum dot displays. *Optics express*, *23*(18), 23680-23693.