Distinct D-box Motifs in SPD-2 Mediate APC/C^{FZR-1}-Dependent Degradation and Centrosomal Localization in *Caenorhabditis elegans* Embryos

Running title: SPD-2 as a substrate of APC/C^{FZR-1}

Rachel N. Yim[1], Joseph R. DiPanni[1], Paollette A. Rivera[1], and Mi Hye Song[1,2]

[1]Department of Biological Sciences, Oakland University, Rochester, MI 48309, USA.
[2]Coresponding Author.

Contact Information: Mi Hye Song (msong2@oakland.edu)

Keywords: Anaphase Promoting Complex (APC/C); FZR-1; *C. elegans*; Centrosome; E3 Ubiquitin Ligase; SPD-2; ZYG-1; D-box; Degron; Proteasomal degradation

Abstract

Centrosome duplication must be tightly regulated to maintain genomic stability. In *Caenorhabditis*

elegans, the APC/C and co-activator FZR-1 function as negative regulators of centrosome

duplication by targeting specific substrates for proteolytic degradation. While *C. elegans* SAS-5

and ZYG-1 have been identified as substrates of APC/C^{FZR-1}, the mechanism by which APC/C^{FZR-1}

^{1}-dependent degradation influences centrosome assembly remains unclear. Here, we identified

SPD-2, the conserved homolog of human CEP192, as a substrate of APC/C^{FZR-1}. We show that

loss of APC/C^{FZR-1} increases both cellular and centrosomal SPD-2 levels, and that SPD-2

physically associates with FZR-1 *in vivo*. Functional analyses of canonical D-box motifs reveal

that D-box1, D-box2, and D-box3 each contribute to SPD-2 degradation, each with different

functional consequences. Mutation of D-box3 alone partially rescued *zyg-1* mutant phenotypes

by restoring centrosome duplication and embryonic viability through increased centrosomal SPD-

2 and ZYG-1. In contrast, mutating D-box1 or D-box2 elevated cellular SPD-2 but did not rescue

zyg-1, with the D-box1 mutation further reducing centrosomal SPD-2 and exacerbating duplication

defects and lethality in *zyg-1* mutants. Our results reveal a conserved mechanism for APC/C^{FZR-1}

^{1}-dependent degradation of SPD-2 and show that its degron motifs have dual functions in

degradation and centrosomal localization, ensuring robust control of centrosome assembly during

C. elegans embryogenesis.

Introduction

Centrosomes are the primary microtubule-organizing centers (MTOCs) in animal cells and play crucial roles in spindle assembly, chromosome segregation, and cell fate determination (Nigg and Holland 2018). Their functional capacity depends on the dynamic recruitment of pericentriolar material (PCM) around a centriole core, ensuring that centrosomes are competent for duplication and microtubule nucleation during each cell cycle. Errors in centrosome regulation can lead to defective mitotic spindles, genomic instability, and developmental disorders, including microcephaly and cancer (Fernandes-Mariano et al. 2025). Understanding how centrosome assembly and activity are coordinated in both time and space is therefore crucial in cell and developmental biology.

A key step in centrosome biogenesis and maturation in *Caenorhabditis elegans* is the recruitment of centrosome components to the site of new centriole formation, a process initiated by the conserved scaffold protein SPD-2 (Kemp et al. 2004; Pelletier et al. 2004) and its upstream factor SAS-7 (Sugioka et al. 2017). SPD-2 localizes to both centrioles and PCM, where it acts upstream of the kinase ZYG-1 (Pelletier et al. 2006; Shimanovskaya et al. 2014) and promotes the recruitment of SPD-5 and γ-tubulin to link PCM assembly with centriole duplication (Kemp et al. 2004; Pelletier et al. 2004). Loss of SPD-2 function significantly impairs centrosome biogenesis, maturation, and microtubule dynamics, underscoring its vital role in centrosome assembly and function (Kemp et al. 2004; Pelletier et al. 2004). Recent structural analyses have further uncovered the modular and intrinsically disordered structure of SPD-2, highlighting its capacity to mediate protein–protein interactions that link centrosome duplication with PCM expansion (Murph et al. 2022). Although SPD-2 is known to be crucial for centrosome regulation, its abundance and activity during centrosome assembly in *C. elegans* embryos remain unclear.

Proteolytic regulation by the anaphase-promoting complex/cyclosome (APC/C), an E3 ubiquitin ligase, has emerged as a key mechanism for controlling centrosome activity and maintaining centrosome copy number (Hames et al. 2001; Strnad et al. 2007; Arquint and Nigg 2014; Meghini et al. 2016; Medley et al. 2017; Medley et al. 2021; Tischer et al. 2022; Meghini et al. 2023). In *Drosophila*, the APC/C, in complex with its co-activator Fizzy-related (Fzr), targets Spd2 for degradation (Meghini et al. 2016). This degradation maintains centrosome asymmetry by limiting Spd2 accumulation at the daughter centrosome, ensuring proper PCM recruitment and maintaining spindle orientation. Without APC/C^{Fzr}-dependent proteolysis, excess Spd2 disrupts centrosome behavior, impairs microtubule nucleation, and randomizes the division axis (Meghini

et al. 2023). These findings demonstrate that proteasome-mediated turnover of conserved
centrosome proteins is a crucial mechanism for maintaining centrosome function. Given the
conserved role of SPD-2 in centrosome assembly, a similar APC/C-dependent regulation may
occur in *C. elegans*. Indeed, SPD-2 functions upstream of ZYG-1 in the centrosome assembly
pathway, indicating that precise regulation of SPD-2 levels is critical for promoting centrosome
duplication and coordinating centrosome maturation (Kemp et al. 2004; Pelletier et al. 2004).
Dysregulation of SPD-2 levels could therefore have serious consequences for centrosome
assembly and function during the cell cycle.

In *C. elegans*, it has been shown that APC/C[FZR-1] regulates centrosome duplication by promoting
the degradation of the centrosome protein SAS-5 through its KEN-box motif (Medley et al. 2017)
and by modulating centrosomal ZYG-1 levels via D-box motifs (Medley et al. 2021). Loss of FZR-
1 or core APC/C subunits elevates SAS-5 and ZYG-1 levels at centrosomes, thereby restoring
bipolar spindle formation in hypomorphic *zyg-1* mutants, revealing that APC/C[FZR-1] functions at
multiple points in the centrosome assembly pathway (Medley et al. 2017; Medley et al. 2021).
Together, these genetic analyses indicate that APC/C[FZR-1] targets multiple substrates, including
ZYG-1 and SAS-5, as well as additional factors acting upstream of ZYG-1. Here, we investigate
the hypothesis that APC/C[FZR-1]-mediated degradation of SPD-2 provides a conserved mechanism
to regulate centrosome assembly in *C. elegans* embryos.

74   **Materials and Methods**

C. elegans **Culture and Genetic Analysis**

The *C. elegans* strains used in this study are listed in Supplementary **Table 1**. All strains were
derived from the wild-type Bristol N2 strain (Brenner 1974; Church et al. 1995) and maintained on
MYOB plates seeded with *Escherichia coli* OP50 at 16°C or 20°C. Some strains were provided
by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).
For genetic analysis, individual L4 hermaphrodites were transferred to new plates and allowed to
produce progeny for 0-24, 24-48, and 48-72 hours at the indicated temperatures. Progeny were
allowed to develop for 18-24 hours before counting the number of larvae and dead eggs.

Immunofluorescence and Cytological analysis

Immunofluorescence and confocal microscopy were performed as described (Stubenvoll et al.
2016). The following primary and secondary antibodies were used at 1:3000 dilutions: DM1a
(Millipore Sigma, #T9026), α-ZYG-1 (Stubenvoll et al. 2016), α-TBG-1 (Stubenvoll et al. 2016), α-

HA (Millipore Sigma, #11867423001, #11583816001; Thermo Fisher, #26183-A647), Alexa Fluor

488 and 568 secondary antibodies (Thermo Fisher, #A11001, A11004, A11006, A11034,

A11036). Confocal microscopy was performed using a Nikon Eclipse Ti-U microscope equipped

with a Plan Apo 60x1.4 NA lens, a Spinning Disk Confocal (CSU X1), and a Photometrics Evolve

512 camera. MetaMorph software (Molecular Devices, Sunnyvale, CA, USA) was employed for

image acquisition and fluorescence intensity quantification, and Adobe Photoshop/Illustrator 2025

were used for image processing. To quantify fluorescent signals at centrosomes, the average

intensity within an 8- or 9-pixel (1 pixel = 0.151 µm) diameter region was recorded from the

highest-intensity focal plane within an area centered on each centrosome. The average intensity

within a 25-pixel diameter region outside the embryo was used for background subtraction.

CRISPR/Cas9 Genome Editing

For genome editing, we used the co-CRISPR method described previously (Arribere et al. 2014;

Paix et al. 2015). crRNA was designed using the CRISPOR web server (crispor.tefor.net;

Concordet and Haeussler, 2018). Animals were microinjected with a mixture of commercially

available SpCas9 (IDT, Coralville, IA) and custom-designed oligonucleotides, including crRNAs

(Supplementary **Table 2**) at 0.4–0.8 mg/ml, tracrRNA at 12 mg/ml, and single-stranded DNA

oligonucleotides (Supplementary **Table 3**) at 25–100 ng/ml. The amount of crRNA was tripled for

low-efficiency crRNAs (SPD-2:D-box2). After injection, we screened for *dpy-10(cn64) II/+* rollers

in F1 progeny and genotyped F2. The genome editing was confirmed by Sanger sequencing

(GeneWiz, South Plainfield, NJ).

Immunoprecipitation (IP) and Immunoblot

IP experiments were performed as described previously (Medley et al. 2021). Embryos were

collected from young gravid worms using hypochlorite treatment (1:2:1 ratio of M9 buffer, 5.25%

sodium hypochlorite, and 5 M NaOH), frozen in liquid nitrogen, and stored at -80°C until use.

Embryos were suspended in lysis buffer [50 mM HEPES, pH 7.4, 2 mM EDTA, pH 8.0, 1 mM

$MgCl_2$, 150 mM KCl, 0.5 mM DTT, 0.5% NP-40 (v/v) and 10% glycerol (v/v)] with complete

protease inhibitor cocktail (Millipore Sigma) and MG132 (Tocris, Avonmouth, Bristol, UK), milled

for 8 minutes (repeat x2) at 30 Hz using a Retsch MM 400 mixer-mill (Verder Scientific, Newtown,

PA), then sonicated for 5 minutes in an ultrasonic bath (Thermo Fisher). Lysates were spun at

45,000 rpm for 45 minutes using a Sorvall RC M120EX ultracentrifuge (Thermo Fisher), and the

supernatant was recovered into clean tubes. An equal amount of total protein lysates was used

for each IP. Lysates and α-HA tag magnetic beads (MBL, #M181-11; Millipore Sigma, #SAE0197)

were incubated with rotation for 1 hour at 4°C and washed (5x 5 minutes) with PBST (PBS + 0.1%

Tween-20). IP with beads and input samples were resuspended in 2X Laemmli Sample Buffer

(Sigma) and boiled for 5 minutes before fractionation on a 4–12% NuPAGE Bis-Tris gel (Thermo

Fisher Scientific). Proteins on a gel were transferred to a nitrocellulose membrane and analyzed

using the antibodies at 1:3000- 10,000 dilutions: α-SPD-2 (Song et al., 2008), α-TBG-1

(Stubenvoll et al., 2016), DM1a (Sigma, #T9026), α-Ollas (Thermo Fisher, #MA5-16125), α-HA

(Millipore Sigma, #11867423001, #11583816001), IRDye secondary antibodies (LI-COR

Biosciences). Blots were imaged with the Odyssey M scanner (LI-COR Biosciences) and

analyzed using Image Studio software (LI-COR Biosciences).

Statistical analysis

Statistics were produced using R statistical software and presented as average ± standard

deviation (SD). Dot plots were generated using the R 'beeswarm' package. In the dotplots, the

box ranges from the first through the third quartile of the data. The thick bar indicates the median.

A solid grey line extends 1.5 times the interquartile range, or to the minimum and maximum data

points. All P-values were calculated using two-tailed t-tests: [ns] $p>0.05$, * $p<0.05$, ** $p<0.01$, ***

$p<0.001$.

140   **Results and Discussion**

Loss of FZR-1 stabilizes cellular SPD-2

In *C. elegans* embryos, APC/C^{FZR-1} negatively regulates centrosome duplication by promoting

proteasomal degradation of SAS-5 (Medley et al. 2017) and by modulating centrosomal ZYG-1

levels (Medley et al. 2021). Previous genetic analyses indicate that APC/C^{FZR-1} targets multiple

substrates, including ZYG-1 and SAS-5, as well as other factors acting upstream of ZYG-1

(Medley et al. 2021). Here, we investigate SPD-2, which functions upstream of ZYG-1 in the

centrosome assembly pathway, as a potential substrate of APC/C^{FZR-1} in *C. elegans* embryos.

If APC/C^{FZR-1} directly targets SPD-2 for proteasomal degradation, then inhibiting APC/C^{FZR-1}

should block SPD-2 degradation, thereby hyper-stabilizing SPD-2 and elevating its overall cellular

levels. We first tested whether loss of APC/C^{FZR-1} affected SPD-2 abundance by quantitative

Western blot analysis using anti-SPD-2 (Song et al. 2008) and protein lysates extracted from *fzr-*

1(bs31) mutant (Medley et al. 2017) and wild-type (WT) embryos (**Fig. 1A**). Our quantification

data showed that SPD-2 levels in *fzr-1(bs31)* mutant embryos are significantly higher (1.89±0.71-

fold, $p<0.001$, n=27) compared to WT controls (1.00±0.0-fold, n=24), while *zyg-1* mutant embryos

show no significant change (0.94±0.16-fold, n=10). In contrast, TBG-1 levels appear unaffected

in both mutants. We also observed a similar trend in *mat-3(or180)* for the core APC8/CDC23 subunit (Golden et al. 2000) and *emb-1(hc57)* mutants for the APC16 (Shakes et al. 2011) in the *C. elegans* APC/C complex (Supplementary **Fig. 1A**). These findings indicate that loss of APC/C^{FZR-1} results in elevated cellular SAS-5 levels, most likely due to increased protein stability caused by impaired proteasomal degradation.

Then, increased levels of cellular SPD-2 in the embryo can influence the amount of SPD-2 recruited to centrosomes. To assess SPD-2 levels at centrosomes, we stained embryos with anti-HA and measured the fluorescence intensity of the SPD-2::2xHA signal at centrosomes (Medley et al. 2023). As SPD-2 localizes to both centrioles and PCM, we measured total centrosomal (centriolar and PCM; **Fig. 1B**) and only centriolar SPD-2 signals (Supplementary **Fig. 1C**). Quantitative immunofluorescence (IF) using SPD-2 tagged with 2xHA (SPD-2::2xHA) reveal that during the first mitosis, *fzr-1(bs31)* mutant embryos exhibit increased levels of total centrosomal SPD-2::2xHA (1.50±0.44-fold, n=92 at metaphase; 1.42±0.37-fold, n=39 at anaphase) compared to WT controls (1.00±0.36-fold, n=47 at metaphase; 1.00±0.31-fold, n=33 at anaphase). A comparable increase was seen in *mat-3(or180)* for the core APC8/CDC23 subunit (Golden et al. 2000) (Supplementary **Fig. 1B**). Additionally, a similar pattern was observed for the centriolar SPD-2 levels (Supplementary **Fig. 1C**). The *fzr-1(bs31)* mutant embryo exhibits increased levels of centriolar SPD-2::2xHA (1.50±0.37-fold, n=98 at metaphase; 1.36±0.44-fold, n=39 at anaphase) compared to WT controls (1.00±0.19-fold, n=34 at metaphase; 1.00±0.30-fold, n=36 at anaphase). Since our prior study showed that *fzr-1(bs31)* mutant embryos display elevated centrosomal ZYG-1 levels (Medley et al. 2021), we also measured ZYG-1 levels by co-staining embryos with anti-HA and anti-ZYG-1 (Stubenvolle et al. 2016; **Fig. 1B**). Consistent with prior observations, *fzr-1(bs31)* mutant embryos exhibit increased levels of centrosomal ZYG-1 (1.35±0.35-fold, n=80 at metaphase; 1.55±0.49-fold, n=31 at anaphase) compared to WT controls (1.00±0.26-fold, n=31 at metaphase; 1.00±0.23-fold, n=32 at anaphase). Our data show that loss of APC/C^{FZR-1} leads to a significant increase in both cellular and centrosomal levels of SPD-2, supporting a model in which APC/C^{FZR-1}-dependent proteolysis regulates SPD-2 abundance during *C. elegans* embryogenesis.

Furthermore, our immunoprecipitation (IP) assays demonstrate a physical interaction between SPD-2 and FZR-1 *in vivo* (**Fig. 1C**). We detected SPD-2::2xHA and Ollas::FZR-1 in the pull-down using embryonic lysates that express SPD-2::2xHA (Medley et al. 2023) and Ollas::FZR-1 (Medley et al. 2021). These results collectively support our hypothesis that APC/C^{FZR-1} directly

190 targets SPD-2 for proteasomal degradation. These findings support our model in which the APC/C

191 and its co-activator FZR-1 (APC/C^{FZR-1}) control SPD-2 levels in *C. elegans* embryos.

SPD-2 contains five putative D-box motifs

The E3 ubiquitin ligase APC/C^{FZR-1} targets specific substrates through the coactivator FZR-1,

which directly recognizes conserved motifs such as the destruction (D)-box and KEN-box within

the substrates (Glotzer et al. 1991; Pfleger and Kirschner 2000). Our *in silico* analysis identified

five potential D-box motifs (RxxL), but no KEN-box, within the *C. elegans* SPD-2 (**Fig. 2A**). In

SPD-2, D-box1 is located within the coiled-coil (CC3: aa291-322) domain, D-box2 within the

Aspm-SPD-2-Hydin (ASH) domain, D-box3 between the Ig-like and PDZ domains, and both D-

box4 and D-box5 within the PDZ domain (Murph et al. 2022). The sequence alignments indicate

that D-box4 is the only fully conserved motif across closely related nematodes, while the other

motifs retain partial but detectable sequence conservation (**Fig. 2B**).

We first examined whether APC/C^{FZR-1} targets SPD-2 by directly recognizing the canonical D-box

degron. To determine if any of the five putative D-box motifs function as degrons *in vivo* by

mediating FZR-1 and SPD-2 interaction, we used CRISPR/Cas9 genome editing to generate

degron mutants by substituting two key residues within each D-box motif (RxxL) with alanine

(AxxA) at their endogenous loci. Viable degron mutants were successfully produced for D-box1,

D-box2, and D-box3, each with two-alanine substitutions, referred to as the SPD-2:DB1(2A),

SPD-2:DB2(2A), and SPD-2:DB3(2A) mutations (**Table 1**). However, homozygous 2A mutations

in D-box4 and D-box5 were lethal, indicating that the normal function of these motifs is essential

for viability. Notably, a genome-wide genetic screen identified the *spd-2(oj29: R710Q)* mutation,

which alters the first Arg (R) residue of D-box4 (710-RTRL-713) and causes centrosome

duplication failure and embryonic lethality (**Fig. 2A**, Kemp et al. 2004). As a result, we were unable

to evaluate the functional roles of the D-box4 and D-box5 degron motifs *in vivo*. Due to this

limitation, we focused our analysis on D-boxes 1, 2, and 3 (DB1, 2, and 3) using viable degron

mutants to assess their functional effects in this study.

Mutating the D-box leads to increased overall cellular SPD-2 levels

If APC/C^{FZR-1} recognizes SPD-2 through its D-box motif, then mutating the D-box degron should

inhibit APC/C^{FZR-1} binding to SPD-2, preventing its degradation and resulting in higher SPD-2

protein levels. To assess how the D-box motif affects overall SPD-2 levels, we conducted

quantitative Western blot analysis using protein lysates from each degron mutant embryo

expressing SPD-2::2xHA (Medley et al. 2023) (**Fig. 2C**). Our results show that SPD-2 levels in all

three D-box mutant embryos are significantly higher (DB1: 1.72±0.43-fold, n=24; DB2: 2.10±0.57-

fold, n=17; DB3: 2.09±0.43-fold, n=11, $p<0.001$) compared to wild-type embryos (1.00±0.0-fold,

n=17). In contrast, TBG-1 levels remain unaffected in degron mutants. These findings suggest

that mutating the D-box motifs renders SPD-2 resistant to degradation, supporting the idea that

the three D-box motifs (DB1, DB2, and DB3) in SPD-2 function as degrons directly recognized by

APC/C^{FZR-1}. Blocking this recognition prevents SPD-2 degradation, leading to its accumulation

and mimicking SPD-2 overexpression.

Only the SPD-2:DB3(2A) mutation results in elevated centrosomal SPD-2 levels

Quantitative Western blot shows that the SPD-2 degron mutations lead to increased total SPD-2

levels in the embryo, consistent with enhanced protein stabilization. To assess whether this

stabilization influences SPD-2 levels at centrosomes, we examined how each degron mutation

changes centrosomal SPD-2 levels. We performed quantitative immunofluorescence (IF) to

compare centrosome-associated SPD-2 signals during the first mitosis in embryos expressing

SPD-2::2xHA, stained with anti-HA tag and anti-tubulin (Medley et al. 2021; **Fig. 2D**).

The DB3(2A) mutation resulted in elevated centrosomal SPD-2 levels (1.35±0.43-fold, $p<0.01$,

n=45 at metaphase; 1.32±0.42-fold, $p<0.0001$, n=34 at anaphase) compared to WT controls

(1.00±0.33-fold, n=60 at metaphase; 1.00±0.27-fold, n=38 at anaphase), indicating that mutating

the D-box3 does not impair centrosomal localization. Instead, the increased centrosomal signal

likely reflects the elevated pool of cellular SPD-2 caused by protein stabilization. In contrast, the

DB1(2A) mutation caused a significant decrease in centrosomal SPD-2 (0.59±0.20-fold, $p<0.001$,

n=28 at metaphase; 0.66±0.26-fold, $p<0.001$, n=10 at anaphase) compared to controls. This

dramatic reduction of centrosomal SPD-2 in the DB1(2A) mutant indicates that the D-box1 motif

is critical for SPD-2 localization to centrosomes, implicating its dual role as both a degron and a

key site for subcellular localization. Meanwhile, the DB2(2A) mutation caused no significant

changes in centrosomal SPD-2 (1.06±0.40-fold, $p=0.55$, n=21 at metaphase; 1.12±0.68-fold,

$p=0.40$, n=6 at anaphase) although the DB2(2A) mutation increases overall SPD-2 levels,

indicating that the SPD-2:DB2(2A) mutation partially impairs SPD-2 localization to centrosomes.

These degron mutants also showed consistent trends in centriolar SPD-2 levels (Supplementary

Fig. 2A). The DB3(2A) mutant embryos exhibit higher levels (1.23±0.28-fold, n=31 at metaphase;

1.13±0.22-fold, n=35 at anaphase), while the DB1(2A) mutants have reduced levels (0.57±0.18-

fold, n=28 at metaphase; 0.63±0.17-fold, n=10 at anaphase) and the DB2(2A) mutants show no

significant change (1.07±0.46-fold, n=21 at metaphase; 0.98±0.2-fold, n=8 at anaphase),
compared to WT controls (1.00±0.33-fold, n=56 at metaphase; 1.00±0.23-fold, n=27 at
anaphase).

Furthermore, to assess the impact of SPD-2 degron mutations on ZYG-1 recruitment, we
quantified centrosomal ZYG-1 levels in the degron mutants stained with anti-ZYG-1 and anti-
tubulin (**Fig. 2E**). Consistent with their elevated SPD-2 levels, DB3(2A) mutants exhibited a
significant increase in centrosomal ZYG-1 (1.14±0.29-fold, $p<0.01$, n=57 at metaphase;
1.21±0.21-fold, $p<0.001$, n=31 at anaphase) compared to WT controls (1.00±0.28-fold, n=72 at
metaphase; 1.00±0.25-fold, n=45 at anaphase). In contrast, DB1(2A) mutants, which display
reduced SPD-2 at centrosomes, showed a marked decrease in centrosomal ZYG-1 (0.50±0.13-
fold, $p<0.001$, n=24 at metaphase; 0.45±0.13-fold, $p<0.001$, n=19 at anaphase). DB2(2A) mutants
exhibited a slight reduction with no significant difference (0.84±0.10-fold, $p=0.07$, n=10 at
metaphase; 0.88±0.09-fold, $p=0.09$, n=14 at anaphase). These results align with the observed
patterns of centrosomal SPD-2 levels across the degron mutants (**Fig. 2D**). These data reveal a
strong correlation between centrosomal SPD-2 levels and ZYG-1 recruitment: mutants with
elevated centrosomal SPD-2 recruit more ZYG-1 to centrosomes, while those with reduced
centrosomal SPD-2 show a proportional decrease in ZYG-1.

The SPD-2:DB3(2A) mutation leads to *zyg-1* suppression
Next, we investigated how specific degron mutations affected SPD-2 activity in regulating
centrosome assembly, using the hypomorphic *zyg-1(it25)* genetic background. The *zyg-1(it25)*
allele carries a temperature-sensitive (ts) mutation (P442L) in the cryptic polo-box (CPB) domain,
which is critical for centrosomal recruitment of ZYG-1 (Kemphues et al. 1988; Shimanovskaya et
al. 2014). At the restrictive temperature 24°C, *zyg-1(it25)* mutant embryos fail to duplicate
centrosomes during the first mitosis, leading to monopolar spindle formation in the second mitosis
and resulting in 100% embryonic lethality (O'Connell et al. 2001). In the molecular hierarchy, SPD-
2 acts upstream of ZYG-1 and is required for ZYG-1 loading to centrioles before procentriole
formation (Pelletier et al. 2006).
285
If APC/C[FZR-1] targets SPD-2 for degradation through its D-box degron, then mutating the D-box
motif should block APC/C[FZR-1] binding, thus stabilizing SPD-2. Consequently, hyper-stabilized
SPD-2, similar to overexpression, could compensate for the reduced ZYG-1 activity in *zyg-1(it25)*
mutants. To determine whether any SPD-2 D-box mutation could rescue *zyg-1(it25)* phenotypes,
we introduced each mutation into the *zyg-1(it25)* background by genetic crossing and examined

their effects on embryonic survival (**Fig. 3A**, **Table 1**) and centrosome duplication (**Fig. 3B**).
Consistent with our hypothesis, the DB3(2A) mutation restored embryonic viability (35±16% vs
18±14%, *p*<0.001) and bipolar spindle formation (61±15% vs 34±22%, *p*<0.01) in *zyg-1(it25)*
mutants, compared to *zyg-1(it25)* controls, at the semi-restrictive temperature of 22.5°C. This
rescue effect was more potent at 22.5°C (34.5±15.9%) than at a more restrictive temperature of
23°C (5.0±4.3%; **Table 1**), suggesting that the SPD-2:DB3(2A) mutation depends on residual
ZYG-1 activity for suppression. However, suppression of *zyg-1* by the SPD-2:DB3(2A) mutation
appeared significantly weaker than that observed with the *fzr-1* mutation (100% bipolar spindles
and embryonic viability; **Fig. 3B, Table 1**) (Medley et al. 2021).
300
In contrast, the DB1(2A) mutation markedly decreased embryonic viability (0.5±0.9% vs 18±14%,
p<0.001; **Fig. 3A, Table 1**) and bipolar spindles (1.2±2.2% vs 34±22%, *p*<0.001; **Fig. 3B**) in *zyg-*
1(it25) mutants, although the DB1(2A) mutant alone remained largely viable (99.2±1.1%; **Table**
1). These results indicate that *zyg-1(it25)* provides a sensitized genetic background where SPD-
2:DB1(2A) causes near-complete lethality, demonstrating a strong synthetic effect. Similarly,
although *spd-2* and *zyg-1* mutations are individually recessive, *spd-2/+; zyg-1/+* double
heterozygotes exhibit ~75% embryonic lethality (Song et al. 2011). DB1(2A) mutant embryos
show centrosomal SPD-2 levels at 59% of those in wild-type embryos, comparable to those in
spd-2 heterozygotes (*spd-2/+*), despite exhibiting elevated cellular SPD-2 levels. Thus, D-box1
appears to regulate centrosomal loading of SPD-2 independently of its degron function. Although
59% of centrosomal SPD-2 is sufficient to support ZYG-1 loading and normal centrosome
duplication in a wild-type background, combining the SPD-2:DB1(2A) mutation with homozygous
zyg-1(it25) severely disrupts centrosome duplication and mitosis, leading to near-complete
embryonic lethality at the semi-restrictive temperature (22.5 °C), where *zyg-1(it25)* alone causes
~80% lethality (**Table 1**). These findings suggest that the D-box1 motif promotes proper
localization of SPD-2 at centrosomes and may facilitate its interaction with ZYG-1. D-box1 resides
within the coiled-coil (CC3) domain (**Fig. 2A**), and the DB1(2A) mutation consists of two alanine
substitutions (R298A and L301A). Since coiled-coil domains are known to mediate protein–protein
interactions (Murph et al. 2022), these substitutions likely impair the ability of SPD-2 to interact
with binding partners necessary for centrosome function.
321
Meanwhile, the DB2(2A) mutation had a minimal negative impact on embryonic viability
(14.4±17.6% vs 18±14%, *p*=0.36; **Table 1, Fig. 3A**) and bipolar spindle formation (30.2±18.5%
vs 34±22%, *p*=0.55; **Fig. 3B**) in *zyg-1(it25)* mutants. Like DB1(2A), the DB2(2A) mutation

increased overall SPD-2 levels but did not enhance its centrosomal activity, suggesting a partial
defect in SPD-2 loading, though much milder than DB2(2A). Since it resides within the ASH
domain, which is known to target SPD-2 to centrosomes (Murph et al. 2022), this may partly
explain the modest yet negative impact of the DB2(2A) mutation on embryonic viability, unlike its
effect on SPD-2 stability.

Increased centrosomal ZYG-1 levels in SPD-2:DB3(2A) mutants

In *zyg-1(it25)* embryos, centrosomal ZYG-1 is reduced to ~40% of WT controls (Medley et al.
2017), impairing the recruitment of SAS-5 and SAS-6, which are required for subsequent SAS-4
loading to centrosomes (Pelletier et al. 2006). Since SPD-2 acts upstream of ZYG-1 and promotes
centrosomal recruitment of ZYG-1, the observed suppression of *zyg-1* mutant phenotypes likely
results from increased centrosomal ZYG-1 levels caused by SPD-2 stabilization. Consequently,
increased levels of centrosomal ZYG-1 may restore the recruitment of downstream factors,
rendering centrosomes competent to assemble new centrioles in *zyg-1(it25)* mutants.

To assess how the SPD-2 degron mutation influenced ZYG-1 recruitment in *zyg-1(it25)* mutants,
we measured centrosomal ZYG-1 levels in *zyg-1(it25)* embryos carrying each degron mutation
(**Fig. 3C**). As expected, the SPD-2:DB3(2A) mutation led to increased centrosomal ZYG-1 levels
(1.15±0.22-fold, $p<0.001$, n=52 at metaphase; 1.20±0.29-fold, $p<0.001$, n=43 at anaphase)
compared to *zyg-1(it25)* controls (1.00±0.14-fold, n=44 at metaphase; 1.00±0.15-fold, n=33 at
anaphase). In contrast, the DB1(2A) mutation significantly decreased centrosomal ZYG-1
(0.53±0.14-fold, $p<0.001$, n=24 at metaphase; 0.31±0.06-fold, $p<0.001$, n=6 at anaphase), while
DB2(2A) mutants exhibited a slight decrease (0.92±0.18-fold, $p=0.466$, n=12 at metaphase;
0.91±0.29-fold, $p=0.02$, n=10 at anaphase). These results align with the patterns of centrosomal
SPD-2 levels across each degron mutant combined with *zyg-1(it25)* (Supplementary **Fig. 3A)**.
The SPD-2:DB3(2A) mutation caused higher centrosomal SPD-2 levels (1.22±0.35-fold, $p<0.001$,
n=66 at metaphase; 1.25±0.28-fold, $p<0.001$, n=69 at anaphase), while DB1(2A) resulted in a
dramatic decrease (0.74±0.25-fold, $p<0.001$, n=22 at metaphase; 0.74±0.30-fold, $p<0.001$, n=33
at anaphase), and DB2(2A) showed a slight reduction (0.86±0.22-fold, $p<0.01$, n=42 at
metaphase; 1.03±0.24-fold, $p=0.67$, n=13 at anaphase), relative to *zyg-1(it25)* controls
(1.00±0.29-fold, n=76 at metaphase; 1.00±0.19-fold, n=30 at anaphase). These findings indicate
that the SPD-2:DB3(2A) mutation leads to increased centrosomal ZYG-1, which enhances the
recruitment of downstream factors and restores centrosome duplication and embryonic viability

in *zyg-1* mutants. This suggests a possible mechanism by which SPD-2:DB3(2A) suppresses the
zyg-1 phenotype.
360
Finally, we observed that the SPD-2:DB3(2A) mutation results in lower centrosomal SPD-2 levels
than the *fzr-1(bs31)* mutation in *zyg-1(it25)* mutant backgrounds (1.25±0.28-fold, n=69 vs
2.59±0.89-fold, n=14, *p*<0.001; Supplementary **Fig. 3B**), consistent with the weaker suppression
of the *zyg-1* phenotype by the SPD-2:DB3(2A) mutation relative to the *fzr-1* mutation. SPD-
2:DB3(2A) mutant embryos show increased cellular SPD-2 levels, resulting in higher centrosomal
levels of SPD-2 and ZYG-1. However, the SPD-2:DB3(2A) mutation alone produces a much
weaker rescue of embryonic viability (1.25±0.28% vs 100%, **Table 1**) and bipolar spindle
formation (60±15% vs 100±0%, **Fig. 2B**) in *zyg-1(it25)* mutants compared to the *fzr-1* mutation.
Similar results were observed in our earlier studies, where ZYG-1 or SAS-5 degron mutations led
to *zyg-1* suppression that was less effective than the *fzr-1* mutation (Medley et al. 2017; Medley
et al. 2021). Several factors may account for the relatively weak suppression of *zyg-1* by the
DB3(2A) single mutation. First, we were unable to test the *in vivo* functions of D-box4 and D-box5
because their 2A substitutions are homozygous lethal, suggesting that these degrons may play a
more dominant role in SPD-2 turnover than D-box3. Second, we could not assess the combined
effects of D-boxes 1 through 3 on *zyg-1* suppression because D-box1 and D-box2 appear to be
required for centrosomal localization of SPD-2 and fail to rescue *zyg-1* phenotypes, making triple
mutant analysis challenging. Thus, the DB3(2A) mutation alone is unlikely to entirely block
APC/C^{FZR-1}-dependent degradation of SPD-2, resulting in only a partial rescue of the *zyg-1(it25)*
phenotype, which is weaker than the *fzr-1* mutation. Another possibility is that APC/C^{FZR-1} targets
additional centrosome proteins upstream of SPD-2, such as SAS-7 (Sugioka et al. 2017). In this
scenario, loss of APC/C^{FZR-1} would enhance ZYG-1 recruitment both by stabilizing ZYG-1 directly
and by increasing the levels of SPD-2 and other upstream factors, such as SAS-7, required for
ZYG-1 loading. However, testing the contribution of SAS-7 is beyond the scope of the present
study and will require additional investigation.
385
SPD-2:DB1(2A) mutant embryos exhibit abnormal cell division patterns
During our genetic analysis at 25°C, we observed a low but significant rate of embryonic lethality
in the SPD-2:DB1(2A) mutants, and this lethality was further exacerbated in the SPD-2:DB1(2A)
mutant embryos tagged with 2xHA (1.2% vs 4.8% lethality, **Table 1**). To investigate the cause of
embryonic lethality, we examined SPD-2:DB1(2A):2xHA mutant embryos stained for
centrosomes, microtubules, and DNA (**Fig. 4**). At 25°C, DB1(2A) mutant embryos display

abnormal cell division phenotypes. Consistent with reduced levels of SPD-2 (**Fig. 2D**) and ZYG-
1 (**Fig. 2E**) at centrosomes, these embryos display tiny and often highly asymmetric centrosomes
during the first mitotic metaphase. Such small and asymmetric centrosomes have also been
observed in ZYG-1:4D mutant embryos, which exhibit unstable and reduced levels of centrosomal
ZYG-1, indicating a partial block in centrosome duplication (Medley et al. 2023). DB1(2A) mutants
also display abnormal patterns in DNA segregation, cell division timing, and mitotic division
patterns, compared to the WT control, although at a low frequency (<5%, n>100). These cell
division defects likely account for embryonic lethality and further confirm that the D-box1 motif
functions not only as a degron motif but also plays a critical role in SPD-2 localization and its
centrosomal activity. Therefore, our study identified new key sites essential for SPD-2 activity in
regulating centrosome assembly during early embryonic development.

In summary, this study reports that *C. elegans* SPD-2 is a direct substrate of APC/C^{FZR-1} and
illustrates a regulatory network that links protein degradation, centrosomal localization, and
interactions with other partners. We showed that SPD-2 physically interacts with FZR-1 *in vivo*,
and that loss of APC/C^{FZR-1} results in elevated levels of both cellular and centrosomal SPD-2.
These results expand the repertoire of APC/C^{FZR-1} substrates beyond SAS-5 and ZYG-1 (Medley
et al. 2017; Medley et al. 2021), highlighting the crucial role of APC/C^{FZR-1} in controlling
centrosome assembly during *C. elegans* embryogenesis. Importantly, our findings also suggest
that APC/C^{FZR-1} may target additional centrosome factors, with SAS-7 (Sugioka et al. 2017) as a
strong candidate for future investigation.

Our degron motif analysis reveals both canonical and noncanonical functions. While all three D-
box mutations (DB1, DB2, and DB3) stabilize SPD-2, only the DB3 mutation restores centrosome
duplication and embryonic viability in *zyg-1* mutants. Our results also reveal that D-box1 and D-
box2 have functions beyond their roles as degron motifs. The D-box1 is required for efficient
centrosomal localization of SPD-2 and may facilitate its interaction with ZYG-1. Consistently,
DB1(2A) mutants show reduced centrosomal SPD-2 and ZYG-1 levels, leading to monopolar
spindle formation and increased embryonic lethality in *zyg-1* mutants. This suggests that D-box1
is important for both stabilizing and centrosomal targeting of SPD-2. By contrast, DB2 mutation
increased total SPD-2 levels but did not alter centrosomal pools or rescue *zyg-1* phenotypes.
These findings demonstrate that degron motifs serve multiple roles, with D-box1 and D-box2
involved in both proteasomal degradation and centrosomal functions.

Finally, our findings have broader evolutionary implications. SPD-2/CEP192 is a highly conserved

key regulator of centrosome assembly in metazoans (Kemp et al. 2004; Pelletier et al. 2004; Zhu

et al. 2008). Because dysregulation of centrosome factors is closely associated with genomic

instability and tumorigenesis in humans (Nigg and Holland 2018), understanding the degron-

dependent regulation of SPD-2/DSpd-2 in *C. elegans* and *Drosophila* (Meghini et al. 2016;

Meghini et al. 2023) provides an evolutionary framework for investigating analogous mechanisms

in human cells. Future work should explore how APC/C^{FZR-1} influences centrosome assembly and

microtubule dynamics, with potential implications for human health and cancer biology.

Table 1. Genetic Analysis

Strain	°C	% Embryonic viability (ave±sd)	n (progeny)
wild-type (N2)	25	99.8±0.3	2214
SPD-2^{DB1(2A)}		98.8±1.0*	2298
SPD-2^{DB2(2A)}		100.0±0.0	1224
SPD-2^{DB3(2A)}		99.9±0.2	2313
SPD-2::2xHA	25	99.2±0.7	2322
SPD-2^{DB1(2A)}*::2xHA*		95.2±2.7**	2124
SPD-2^{DB2(2A)}*::2xHA*		99.4±0.5	2431
SPD-2^{DB3(2A)}*::2xHA*		100.0±0.0**	1648
zyg-1(it25)	22.5	18.2±14.4	6053
SPD-2^{DB1(2A)}*; zyg-1(it25)*		0.5±0.9***	6249
SPD-2^{DB2(2A)}*; zyg-1(it25)*		14.4±17.6	4913
SPD-2^{DB3(2A)}*; zyg-1(it25)*		34.5±15.9***	6285
zyg-1(it25) fzr-1(bs31)		100.0±0.0***	1848
zyg-1(it25)	23	0.0±1.0	1163
SPD-2^{DB1(2A)}*; zyg-1(it25)*		0.0±0.0	859
SPD-2^{DB2(2A)}*; zyg-1(it25)*		0.0±0.6	710
SPD-2^{DB3(2A)}*; zyg-1(it25)*		5.0±4.3***	1095

*** $p<0.001$, ** $p<0.01$, * $p<0.05$, compared to the corresponding controls.

Data and Reagent Availability

The authors confirm that all data necessary to support the conclusions of the article are presented
within the article, figures, tables, and supplementary material. Strains used in this study are
available upon request.

Acknowledgements

We thank former and current members of the Song lab for their technical support. We especially
thank WormBase and the *Caenorhabditis* Genetics Center (CGC). WormBase is supported by
grant U24HG002223 from the National Human Genome Research Institute at the US National
Institutes of Health, the UK Medical Research Council, and the UK Biotechnology and Biological
Sciences Research Council. The CGC (St. Paul, MN), is funded by the National Institutes of
Health Office of Research Infrastructure Programs (P40 OD010440).

Competing Interests

No competing interests declared.

Author Contributions

M.H.S. designed the experiments, wrote the manuscript, and performed quantifications of
confocal imaging and protein levels from western blots. R.N.Y., J.D., P.R., and M.H.S. performed
experiments and provided data.

Funding

This work was supported by a grant 1R15GM147857 to M.H.S. from the National Institute of
General Medical Sciences. The funders had no role in study design, data collection and analysis,
decision to publish, or manuscript preparation.

References

Arquint, C., and E. A. Nigg, 2014 STIL microcephaly mutations interfere with APC/C-mediated

degradation and cause centriole amplification. Curr Biol 24**:** 351-360.

Arribere, J. A., R. T. Bell, B. X. Fu, K. L. Artiles, P. S. Hartman *et al.*, 2014 Efficient marker-free

recovery of custom genetic modifications with CRISPR/Cas9 in Caenorhabditis elegans.

Genetics 198**:** 837-846.

Brenner, S., 1974 The genetics of Caenorhabditis elegans. Genetics 77**:** 71-94.

Church, D. L., K. L. Guan and E. J. Lambie, 1995 Three genes of the MAP kinase cascade,

mek-2, mpk-1/sur-1 and let-60 ras, are required for meiotic cell cycle progression in

Caenorhabditis elegans. Development 121**:** 2525-2535.

Fernandes-Mariano, C., J. N. Bugalhao, D. Santos and M. Bettencourt-Dias, 2025 Centrosome

biogenesis and maintenance in homeostasis and disease. Curr Opin Cell Biol 94**:**

102485.

Glotzer, M., A. W. Murray and M. W. Kirschner, 1991 Cyclin is degraded by the ubiquitin

pathway. Nature 349**:** 132-138.

Golden, A., P. L. Sadler, M. R. Wallenfang, J. M. Schumacher, D. R. Hamill *et al.*, 2000

Metaphase to anaphase (mat) transition-defective mutants in Caenorhabditis elegans. J

Cell Biol 151**:** 1469-1482.

Hames, R. S., S. L. Wattam, H. Yamano, R. Bacchieri and A. M. Fry, 2001 APC/C-mediated

destruction of the centrosomal kinase Nek2A occurs in early mitosis and depends upon

a cyclin A-type D-box. EMBO J 20**:** 7117-7127.

Kemp, C. A., K. R. Kopish, P. Zipperlen, J. Ahringer and K. F. O'Connell, 2004 Centrosome

Maturation and Duplication in C. elegans Require the Coiled-Coil Protein SPD-2. Dev

Cell 6**:** 511-523.

Kemphues, K. J., M. Kusch and N. Wolf, 1988 Maternal-effect lethal mutations on linkage group

II of Caenorhabditis elegans. Genetics 120**:** 977-986.

Medley, J. C., L. E. DeMeyer, M. M. Kabara and M. H. Song, 2017 APC/CFZR-1 Controls SAS-

5 Levels To Regulate Centrosome Duplication in Caenorhabditis elegans. G3

Genes|Genomes|Genetics 7**:** 3937-3946.

Medley, J. C., J. R. DiPanni, L. Schira, B. M. Shaffou, B. M. Sebou *et al.*, 2021 APC/CFZR-1

regulates centrosomal ZYG-1 to limit centrosome number. J Cell Sci 134.

Medley, J. C., R. N. Yim, J. DiPanni, B. Sebou, B. Shaffou *et al.*, 2023 Site-specific

phosphorylation of ZYG-1 regulates ZYG-1 stability and centrosome number. iScience

26**:** 108410.

Meghini, F., T. Martins, X. Tait, K. Fujimitsu, H. Yamano *et al.*, 2016 Targeting of Fzr/Cdh1 for

timely activation of the APC/C at the centrosome during mitotic exit. Nat Commun 7**:**

12607.

Meghini, F., T. Martins, Q. Zhang, N. Loyer, M. Trickey *et al.*, 2023 APC/C-dependent

degradation of Spd2 regulates centrosome asymmetry in Drosophila neural stem cells.

EMBO Rep 24**:** e55607.

Murph, M., S. Singh and M. Schvarzstein, 2022 A combined in silico and in vivo approach to the

structure-function annotation of SPD-2 provides mechanistic insight into its functional

diversity. Cell Cycle 21**:** 1958-1979.

Nigg, E. A., and A. J. Holland, 2018 Once and only once: mechanisms of centriole duplication

and their deregulation in disease. Nat Rev Mol Cell Biol 19**:** 297-312.

O'Connell, K. F., C. Caron, K. R. Kopish, D. D. Hurd, K. J. Kemphues *et al.*, 2001 The C.

elegans zyg-1 gene encodes a regulator of centrosome duplication with distinct maternal

and paternal roles in the embryo. Cell 105**:** 547-558.

Paix, A., A. Folkmann, D. Rasoloson and G. Seydoux, 2015 High Efficiency, Homology-Directed

Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein

Complexes. Genetics 201**:** 47-54.

Pelletier, L., E. O'Toole, A. Schwager, A. A. Hyman and T. Muller-Reichert, 2006 Centriole

assembly in Caenorhabditis elegans. Nature 444**:** 619-623.

Pelletier, L., N. Ozlu, E. Hannak, C. Cowan, B. Habermann *et al.*, 2004 The Caenorhabditis

elegans centrosomal protein SPD-2 is required for both pericentriolar material

recruitment and centriole duplication. Curr Biol 14**:** 863-873.

Pfleger, C. M., and M. W. Kirschner, 2000 The KEN box: an APC recognition signal distinct from

the D box targeted by Cdh1. Genes Dev 14**:** 655-665.

Shakes, D. C., A. K. Allen, K. M. Albert and A. Golden, 2011 emb-1 encodes the APC16 subunit

of the Caenorhabditis elegans anaphase-promoting complex. Genetics 189**:** 549-560.

Shimanovskaya, E., V. Viscardi, J. Lesigang, M. M. Lettman, R. Qiao *et al.*, 2014 Structure of

the C. elegans ZYG-1 cryptic polo box suggests a conserved mechanism for centriolar

docking of Plk4 kinases. Structure 22**:** 1090-1104.

Song, M. H., L. Aravind, T. Muller-Reichert and K. F. O'Connell, 2008 The conserved protein

SZY-20 opposes the Plk4-related kinase ZYG-1 to limit centrosome size. Dev Cell 15**:**

901-912.

Song, M. H., Y. Liu, D. E. Anderson, W. J. Jahng and K. F. O'Connell, 2011 Protein
phosphatase 2A-SUR-6/B55 regulates centriole duplication in C. elegans by controlling
the levels of centriole assembly factors. Dev Cell 20**:** 563-571.
Strnad, P., S. Leidel, T. Vinogradova, U. Euteneuer, A. Khodjakov *et al.*, 2007 Regulated
HsSAS-6 levels ensure formation of a single procentriole per centriole during the
centrosome duplication cycle. Dev Cell 13**:** 203-213.
Stubenvoll, M. D., J. C. Medley, M. Irwin and M. H. Song, 2016 ATX-2, the C. elegans Ortholog
of Human Ataxin-2, Regulates Centrosome Size and Microtubule Dynamics. PLoS
Genet 12**:** e1006370.
Sugioka, K., D. R. Hamill, J. B. Lowry, M. E. McNeely, M. Enrick *et al.*, 2017 Centriolar SAS-7
acts upstream of SPD-2 to regulate centriole assembly and pericentriolar material
formation. Elife 6.
Tischer, T., J. Yang and D. Barford, 2022 The APC/C targets the Cep152-Cep63 complex at the
centrosome to regulate mitotic spindle assembly. J Cell Sci 135.
Zhu, F., S. Lawo, A. Bird, D. Pinchev, A. Ralph *et al.*, 2008 The mammalian SPD-2 ortholog
Cep192 regulates centrosome biogenesis. Curr Biol 18**:** 136-141.
544
545

Figure 1. Loss of FZR-1 stabilizes SPD-2. (**A**) Representative western blot with anti-SPD-2 and

anti-TBG-1 using embryonic lysates extracted from *fzr-1(bs31)* mutants, wild-type(*N2*), and *zyg-*

1(it25) mutants (Left). Tubulin was used as a loading control. Quantitative Western blot analyses

show significantly increased levels of SPD-2 (Middle) in *fzr-1(bs31)* mutants (1.89±0.71-fold,

p<0.001, n=27), but no change in *zyg-1(it25)* mutants (0.94±0.16-fold, *p*=0.49, n=10), compared

to wild-type (WT) controls (1.00±0.0-fold, n=24). In contrast, there were no significant differences

in TBG-1 levels (Right) among *fzr-1(bs31)* mutants (1.01±0.12-fold, *p*=0.80, n=14), *zyg-1(it25)*

mutants (1.01±0.07-fold, *p*=0.53, n=15), and wild-type controls (1.00±0.0-fold, n=15). (**B**) (Top)

Quantification of total centrosomal SPD-2::2xHA and ZYG-1 levels during the first mitotic cycle.

Each dot represents a centrosome. (Bottom) Embryos stained for SPD-2::2xHA and ZYG-1 at the

first mitotic metaphase and anaphase, with magnified images of posterior centrosomes. Bars, 10

μm. In the plots, the box ranges from the first through the third quartile of the data. The thick bar

indicates the median. (**A-B**) The solid grey line extends 1.5 times the interquartile range or to the

minimum and maximum data points. [ns]*p*>0.05, ***p*<0.001 (two-tailed t-tests). (**C**) IP using anti-

HA suggests that SPD-2 physically interacts with FZR-1. Protein lysates extracted from embryos

expressing only Ollas::FZR-1 were used as a negative control; ~1% of the total lysates were

loaded in the input lanes. Tubulin was used as a loading control in the input lanes.

Figure 2. Mutating D-box motifs elevates overall cellular levels of SPD-2. (**A**) The SPD-2

protein structure illustrates the locations of D-boxes (DB:1 through 5), functional domains, and

mutant alleles. (**B**) Sequence alignments of D-box motifs in the *Caenorhabditis* species. (**C**) (Left)

Representative western blot with anti-HA and anti-TBG-1 using embryonic lysates from individual

degron mutants and wild-type (WT) controls expressing SPD-2::2xHA. Tubulin was used as a

loading control. (Middle) Quantitative western blot analysis shows significantly increased levels

of SPD-2::2xHA in all three degron mutants. (Right) In contrast, there were no significant

differences in TBG-1 levels among the degron mutants and wild-type controls. (**D-E**) (Left)

Quantification of total centrosomal SPD-2::2xHA levels during the first mitotic cycle. Each dot

represents a centrosome. (Right) Embryos stained for (**D**) SPD-2::2xHA and (**E**) ZYG-1 at the first

mitotic metaphase and anaphase, with magnified images of posterior centrosomes. Bars, 10 μm.

(**C-E**) In the plots, the box ranges from the first through the third quartile of the data. The thick bar

indicates the median. The solid grey line extends 1.5 times the interquartile range or to the

minimum and maximum data points. [ns]*p*>0.05, ***p*<0.01, ****p*<0.001 (two-tailed t-tests).

Figure 3. Mutating SPD-2 D-box3 leads to the suppression of *zyg-1* phenotypes and increased centrosomal ZYG-1 levels. (**A**) Embryonic viability at 22.5°C (See **Table 1**). Each dot represents a hermaphrodite. (**B**) Quantification of monopolar (dark grey) and bipolar (light grey) spindles in two-cell stage embryos at 23°C. Average and standard deviation (SD) are presented. n is the number of blastomeres. Embryos stained for centrosomes (TBG-1), microtubules, and DNA, illustrate mitotic spindles at the second mitosis. (**C**) (Left) Quantification of centrosome-associated ZYG-1 levels at the first mitotic metaphase and anaphase in *zyg-1(it25)* mutant backgrounds with individual degron mutations. Each dot represents a centrosome. (Right) Embryos stained for ZYG-1 and microtubules during the first mitotic cycle, with magnified images of posterior centrosomes. (**B-C**) Bars, 10 μm. (**A,C**) In the plots, the box ranges from the first through the third quartile of the data. The thick bar indicates the median. A solid grey line extends 1.5 times the interquartile range, or to the minimum and maximum data points. (**A-C**) $^{ns}p>0.05$, *$p<0.05$, **$p<0.01$, ***$p<0.001$ (two-tailed t-tests).

Figure 4. SPD-2:DB1(2A) mutants exhibit cell division defects during early embryogenesis. Embryos expressing SPD-2::2xHA stained for centrosomes (SAS-4), microtubules, and DNA. At 25°C, SPD-2:DB1(2A) mutant embryos exhibit abnormal cell division phenotypes, including small centrosomes and highly asymmetric centrosome sizes during the first mitotic metaphase, DNA missegregation (arrowhead), synchronous mitotic timing, and aberrant patterns of mitotic divisions, compared to the WT embryos (Top). Bar, 10 μm.

Figure 1.

Figure 2.

Figure 3.

Figures 1-4 Yim et al. 2025

Figure 4.