Ecological context shapes microbial contributions to

nutrition and development in *Drosophila melanogaster*

Short title: Ecological context shapes nutritional symbiosis in *Drosophila*

Elisabeth K. Riedel [1*], Laura Lender [1,2], Vienna Kowallik [3], Melanie M. Pollierer [4,5], Marko
Rohlfs [1*]

[1] Institute of Ecology, University of Bremen, Bremen, Germany

[2] Institute of Biology, Freie Universität Berlin, Berlin, Germany

[3] Chair of Forest Entomology and Protection, University of Freiburg, Freiburg, Germany

[4] Julius Kühn-Institute (JKI) - Federal Research Center for Cultivated Plants, Institute for
Forest Protection, Quedlinburg, Germany

[5] JFB Institute of Zoology and Anthropology, University of Göttingen, Göttingen, Germany

Corresponding authors:

Elisabeth K. Riedel: eriedel@uni-bremen.de, Marko Rohlfs: rohlfs1@uni-bremen.de

17 Abstract

Microbial symbionts play a critical role in shaping insect development by supplying
essential nutrients, influencing host phenotype, and mediating context-dependent
mutualistic outcomes. In the fruit fly *Drosophila melanogaster,* larvae develop in
ephemeral breeding substrates such as decaying fruits, where they acquire maternally
transmitted bacterial and fungal symbionts. These substrates vary in their structural and
chemical properties, potentially altering the composition and function of associated
microbial communities. Here, we combine semi-natural microcosm experiments with
compound-specific isotope analysis of essential amino acids to assess how variation in
fruit substrate and symbiont origin affect larval nutrition, development, and adult
phenotypes. We find that microbial community composition and environmental context
jointly shape amino acid flow, with larvae deriving variable proportions of their nutrition
from plant, bacterial, and fungal sources. These differences translate into distinct
developmental outcomes, including shifts in larval development time – either delayed or
accelerated, depending on the symbiont-substrate combination – and significant
variation in adult body weight. Our findings introduce symbiont-environment interactions
as a framework for understanding how ecological context shapes nutritional mutualisms,
showing that environmental heterogeneity can alter microbial composition, nutrient
sources, and larval performance. Together, these results reveal how ecological context
modulates mutualistic dependence and underscores the context sensitivity of host-
microbe associations.

Introduction

The evolutionary success of insects in occupying a wide variety of ecological niches is outstanding (1). Interactions with microorganisms, such as bacteria and fungi, which lead to the formation of temporarily stable or transient host-microbe symbioses contribute to this success (2,3). Symbiotic microbiota facilitate the realization of the host's ecological niche through nutritional supplementation, protection from pathogens, and detoxification or other environmental modifications (4–8). The composition of symbiotic microbiota is shaped by a diversity of both microbe-microbe interactions and host control mechanisms, which operate across open ecosystems to precise genetic shaping and vertical transmission (9).

In systems where both hosts and microbial symbionts can associate flexibly with multiple partners (10), ecological context becomes a key driver of symbiont community composition, with potentially significant consequences for host phenotypes and the stability of the mutualism (11). This is especially true for the vast diversity of insects that develop in ephemeral resource patches, such as dung, carrion, mushroom, or rotting plant tissue (12). The benefits of microbiota that vary across time and space are less evident than those of mutualistic partners with well-defined functions or identities (10,13–15). Rather than symbiont composition or environmental factors alone, it is their dynamic interplay that ultimately determines the expression of insect life-history traits and, by extension, the breadth of the host's ecological niche (10).

Cases where the impact of microbial symbionts on host development depends on the breeding environment, defined as symbiont-environment interactions, remain largely unexplored regarding the arising benefits and costs for the host (15). In this study, we address this gap by investigating how variation in the breeding environment influences symbiont community composition, nutrient acquisition during larval development, and the resulting host phenotype in the fruit fly *Drosophila melanogaster*.

Drosophila melanogaster, a cosmopolitan decomposer insect, is a representative of the vast diversity of insect species that are generalists for transient breeding sites, including dung, carrion, fruits, and plant litter. For example, *D. melanogaster* was observed to breed in 15 of 24 plant substrates, primarily fruits, examined by Atkinson and Shorrocks (16). During juvenile development the larval stages of *Drosophila* spp. engage in prolonged and intimate interactions with microbiota residing both inside their gut and in the decaying plant substrate (17,18). Symbionts serve as a vital source of amino acids within predominantly low-protein plant substrates (19,20) thus, playing a pivotal role in shaping the expression of adult life-history traits that are largely contingent upon preadult growth and development (21). Furthermore, dietary symbionts can also exhibit a protective role against pathogens like toxin-producing filamentous mold fungi in *Drosophila* larvae (22). The deposition of fecal maternal microbiota during egg-laying significantly influences the symbiont community composition in localized breeding patches (23–27). Additionally, the inherent characteristics of the breeding substrate and seasonal variation seem to have a profound influence on the symbiont composition of *Drosophila* (14,28–30). Beyond seasonal weather variation, a complex interplay of factors, including plant substrate properties such as pH, water availability, and nutrient content, along with microbial interactions, shapes these dynamics (31). To date, how variation in symbiotic microbes influence *D. melanogaster* development across different

environments and thus contribute to generalization in mutualism (10) remains an open
question. In particular, the consequences of fruit substrate variation, and its impact on
the bacterial and fungal symbionts that provide essential nutrition during larval
development, is unknown.

Carbon (^{13}C) and nitrogen (^{15}N) stable isotope analysis offers an effective method for
uncovering the specific plant substrates that serve as nutritional sources for *Drosophila*
larvae (32). However, bulk stable isotope analyses fall short in distinguishing between
plants and microorganisms as nutritional sources for fruit fly larvae. Compound specific
stable isotope analysis (CSIA) of essential amino acids (EAA) is an emerging tool to
investigate both basal resources and trophic positions of consumers (33–36). Unique
patterns, or 'fingerprints', of δ^{13}C in EAA produced by bacteria, fungi or plants allow to
quantitatively trace the biosynthetic origin of these EAA in consumers (36–38). We used
this approach to quantify, for the first time, how plant substrates, bacteria, and fungi
differentially contribute to larval nutrition in *Drosophila*, depending on breeding
environment and symbiont composition.

We analyzed EAA profiles of adult *Drosophila* and the respective ontogenetic
environment. These adult flies developed from larvae reared under distinct symbiont–
environment combinations, established by inoculating different plant substrates with
female-derived fecal microbiota from flies maintained in source microcosms. The
microcosms, based on field-exposed substrates, sustained fly-symbiont associations
and enabled successive microbial transmission across generations. Generally, we used
two types of microbiota-inoculated plant substrates: substrate with symbionts from
females that lived in the same substrate environment (autochthonous symbiont
inoculum, *autoch*SI) and substrate with symbionts from females that came from a
different substrate environment (allochthonous symbiont inoculum, *alloch*SI; Fig 1). We
employed a metabarcoding approach to, first, characterize the bacterial and fungal
symbionts transmitted through fecal matter deposition by females from different
substrate environments (39). Microdiversity of strain diverse bacterial phylotypes was
assessed to reveal the potential of environment-specific symbiont strain selection (40).
Second, we analyzed microbiota changes during *Drosophila* development in substrate
with *autoch*SI and *alloch*SI. Lastly, we recorded larval development time and biomass
gain through quantification of adult body weight to investigate how the consequences of
*autoch*SI versus *alloch*SI are connected to larval nutrient intake and variation in the
expression of insect life-history traits. In this context, it is assumed that ephemeral
resources, which can be subject to unpredictable catastrophic events (12) exert selective
pressure for rapid development while simultaneously selecting for maximal biomass gain
as key fitness components in holometabolous insects (21,41).

By emphasizing the often overlooked yet pervasive influence of environmental
heterogeneity on symbiont composition in a generalist host-microbe system, our study
advances the understanding of nutritional symbioses shaping host development and
mutualistic dependence.

Fig 1. Ecological context as determined by autochthonous versus allochthonous symbiont transmission may cause variation in the expression of insect life-history traits.
Many insects develop in short-lived, ephemeral resource patches. The subsequent generation of adult insects may encounter similar resource patches and reproduce there. However, due to the temporal and spatial variability in the occurrence of these habitats, dispersing individuals may instead encounter novel and dissimilar breeding sites that differ substantially in both structure and chemical composition. Microbial symbionts, which are essential for larval development and transmitted maternally, are thereby exposed to new environmental conditions. These shifts may alter symbiont growth dynamics and community composition, potentially affecting host developmental outcomes. Thus, the breeding substrate type determines whether symbionts supporting juvenile development encounter a familiar (autochthonous) or a novel (allochthonous) ecological context.

Thus, it provides new insights into the limitations that environmental heterogeneity imposes on niche width (42,43) and the evolution of mutualistic symbioses in animal hosts that have been selected to reproduce in ephemeral resource patches (12,44,45). More broadly, we propose symbiont-environment interaction as a conceptual framework for uncovering the ecological processes that generate variation in partner breadth and mutualistic dependence (10,11).

Material and methods

Drosophila stock and rearing

The *Drosophila melanogaster* laboratory stock originated from wild-caught individuals collected in Kiel, Germany in 2006 (46). Since then, the population has been maintained as an outbred stock of several hundred individuals per generation. Flies are reared on a standard laboratory diet (S1 Text), under controlled conditions of 20 °C and a 16:8 h light-dark cycle.

Field-derived microbes and substrate-specific *Drosophila* microcosms

Drosophila melanogaster reared in laboratory environments typically harbor a reduced and less diverse microbiota compared to their wild counterparts (18,28,30). To generate microbiota inocula more representative of natural conditions, we aimed to recolonize our laboratory *D. melanogaster* population with field-acquired microbial symbionts. In September 2019 we exposed cut apple and tomato substrates on enclosed pedestals coated with tangle-trap paste (TEMMEN Insektenleim, Temmen GmbH) allowing access to flying insects only. This exposure allowed the substrates to be colonized by environmental microbes and insects visiting the patches. After one week, we retrieved the baits and removed any remaining arthropods. The microbially enriched substrates were then provided as the sole food source to our laboratory stock of *D. melanogaster*, thereby introducing field-derived microbiota to the lab-reared flies. After 24 hours, these flies were transferred to fresh apple or tomato pieces to facilitate egg laying and the transmission of associated microbes. New fly populations were subsequently reared from these substrates, with the respective fruit type regularly replenished to maintain continuous development and microbial transfer across generations. *Drosophila*-microbe microcosms were maintained in 22 L (39 × 28 × 28 cm) plastic containers, where adult flies had continuous access to water and a mixture of sucrose and hydrolyzed brewer's yeast. Approximately once per month, microcosms were renewed by allowing flies to oviposit into fresh pieces of the respective fruit substrate, which were then transferred to a new container. The number of these transfer events was recorded for each microcosm. Female flies from the populations were used as symbiont donors and are referred to as "apple" and "tomato" flies. A second set of "apple" and "tomato" populations was created in a similar manner in 2020.

Collection of female fecal symbionts

A fundamental assumption of this study is that bacterial and fungal microbiota transmitted via female fecal deposition play a major role in establishing the larval microbiota and shaping offspring-microbe symbioses during development (23–27). To minimize the effects of strong inter-individual variation in microbial transmission (25), we standardized the deposition of female-derived symbionts across all microbiota community analyses and experiments described herein. To standardize symbiont inoculation, 30 female flies from each tomato- or apple-associated population were placed in sterile 15 mL screw-cap tubes and allowed to defecate overnight, depositing their microbiota as fecal droplets on the tube walls. For metabarcoding, flies were removed, and the tubes were immediately frozen at −80 °C until DNA extraction (see

below). For experiments assessing essential amino acid flow and larval development, freshly collected fecal material served as the inoculum. In these cases, after removing the flies, the fecal deposits were suspended in 5 mL phosphate-buffered saline and dislodged by shaking (180 rpm) for 60 minutes. Aliquots of 50 µL were pipetted into each experimental unit (see below).

Microbiota were collected from the 2019 and 2020 microcosms after varying numbers of transfer events. Details of the microcosms used in the larval development experiments are provided in S2 Table. For metabarcoding-based characterization of symbiont communities from "apple" and "tomato" flies, we sampled fecal material from both apple and tomato microcosms established in 2019 after twelve and seventeen transfer events, and from the 2020 microcosms after one, two, and five transfer events. Fecal material from apple-associated flies experimentally applied to apple substrate is termed an autochthonous symbiont inoculum (*autoch*SI), whereas the same material on tomato substrate is termed allochthonous (*alloch*SI). Conversely, fecal material from tomato-associated flies is *autoch*SI on tomato and *alloch*SI on apple (Fig 1).

Metabarcoding of *Drosophila melanogaster* bacterial and fungal microbiota

DNA-Extraction and amplicon sequencing

For the extraction of microbial DNA from fly fecal material, we adapted the protocol described by Fink et al. (47). Briefly, screw-cap tubes containing frozen samples were thawed, and fecal deposits adhering to the tube walls were collected using sterile swabs pre-moistened with sterile phosphate-buffered saline (PBS). The cotton tip of each swab was aseptically removed with sterilized scissors and placed directly into the first reaction tube of the Qiagen DNeasy PowerSoil Kit. To each sample, we added 60 µL of the kit's buffer C1 along with 20 µL of proteinase K, followed by incubation for 2 h at 50 °C in a thermomixer to facilitate cell lysis. Samples were then vortexed for 10 min using a vortex adapter before proceeding with the manufacturer's standard protocol. Two empty screw-cap tubes, processed alongside the fecal samples, served as negative controls for the extraction workflow. For the analysis of symbiont communities in the developmental substrate (see substrate preparation below in *Larval development experiment*), we used material collected after the development of single larvae in the same experimental setup described for the larval development assays (see below). Surface scrapings from substrates were randomly pooled in groups of 10 experimental units, and DNA was extracted using the Qiagen DNeasy PowerSoil Kit according to the manufacturer's protocol.

The DNA samples were sent to Advanced Identification Methods (AIM) GmbH, Leipzig, Germany, for dual-tag library preparation, amplicon sequencing, and demultiplexing. For bacterial community profiling, the V3–V4 hypervariable region of the 16S ribosomal RNA gene using primers 341F and 785R (48) was amplified. Fungal communities were profiled by amplifying the internal transcribed spacer 2 (ITS2) region with primers ITS3_KYO2 (forward) and ITS4_KYO3 (reverse) (49), which are optimized for broad taxonomic coverage of fungi. In total, eleven fecal samples, nine substrate-scrape samples, and two negative extraction controls were processed. Libraries were sequenced on an Illumina

MiSeq platform (v3 chemistry) using a paired-end strategy (2 × 300 bp), generating high-
resolution reads suitable for downstream community composition analyses.

Taxonomic identification
The demultiplexed FASTQ files were processed in QIIME 2 version 2022.2 (50) on a Linux-
based virtual machine (kernel 5.4.0-148-generic). Sequence quality control and
denoising were performed using the *denoise-paired* command, which implements the
DADA2 algorithm (51) as integrated within the QIIME 2 framework, using default settings
unless stated otherwise below.

For the 16S sequences, forward reads were truncated at 299 bp and reverse reads at 255
bp after visual inspection. Primer sequences were trimmed from the start of each read
(17 bp from forward and 21 bp from reverse), leaving an overlap of ~110 bp for merging.
Taxonomic assignment was performed in QIIME 2 using the q2-feature-classifier plugin
(52) with a naïve Bayes classifier. The classifier was trained against the SILVA database
release 132 April 2018 (53), reduced to 99% identity sequences with taxonomy resolved
to seven levels limited to sequences containing our specific primer set.

Due to length variability in the ITS2 region, which increases the likelihood of primer read-
through, primer sequences and their reverse complements were removed using
Cutadapt in QIIME 2 (54). Quality trimming involved removing the first 30 low-quality
bases from both forward and reverse reads, and truncating reverse reads at 215 bp after
visual quality inspection. Forward reads were processed with a quality threshold of 2.0,
while reverse reads were allowed an error rate of up to 5.0 due to lower overall quality.
We did not filter for chimeras since they are much less likely to occur when using ITS as a
marker gene (55). Taxonomic assignment was based on the UNITE database for fungal
sequences, adapted for the QIIME 2 environment. We used the "dynamic" taxonomy, in
which clustering thresholds vary between 1–3% depending on fungal lineage
characteristics (56). A naïve Bayes classifier was trained on the full taxonomy, following
QIIME 2 development team recommendations (57), and applied to our ITS reads for
taxonomic identification (52). A total of 136 unassigned amplicon sequence variants
(ASV) were manually blasted (BLAST+ 2.13.0, December 2022) against the NCBI
"Nucleotide collection (nr/nt)," including GenBank, EMBL, DDBJ, PDB, and RefSeq. Most
unassigned features were of non-fungal origin (plants, animals, etc.) and were excluded
from further analysis. Three additional fungal ASVs were identified and manually added
to the taxonomy table: one Saccharomycetaceae species (one read, NCBI accession:
7ZW0_LD, 100% coverage) and two features belonging to the *Penicillium chrysogenum*
complex (two reads each, NCBI accession: MW774585.1, 98.39% coverage).

Microbial community analysis
All analyses of microbial alpha and beta diversity, as well as taxonomic composition at
the genus, species, and ASV levels, were performed in R (R Core Team, 2022; RStudio
Team, 2022) using the *phyloseq* package (58). For fungal community analysis, ASVs
represented by a single read were removed as likely artifacts, and samples with fewer
than 1,000 reads were excluded. This resulted in the removal of one pooled tomato
substrate sample. Potential contaminants were identified using the *decontam* package
(59) with a threshold-based selection strategy. After filtering, the dataset contained 2,051

taxa represented by more than 1.1 million reads across 16 samples. Similarly, for bacterial communities we used the *decontam* package to identify and remove 27 putative contaminants based on prevalence. Blank samples, empty taxa, and sequences assigned to chloroplasts or mitochondria were excluded. ASVs represented by a single read were removed, and one apple substrate sample was excluded due to low read counts (<1,000). The final dataset comprised 614 taxa represented by ~300,000 reads across 16 samples. Rarefaction curves indicated sufficient sequencing depth for all samples, with asymptotic behavior reached in each case (Fig S1). Therefore, no rarefaction was applied, and analyses proceeded with the full dataset.

Alpha and *beta* diversity of fungal and bacterial communities were analyzed separately for substrate and fecal microbiota. Alpha diversity was quantified as Shannon diversity and observed richness (taxon count) using the *estimate_richness* function in the *phyloseq* package. Richness estimates were compared using Wilcoxon rank-sum tests (*stats* package). *Beta* diversity was calculated as Bray-Curtis dissimilarity (60) and visualized by principal coordinates analysis (PCoA) using multidimensional scaling (MDS). Differences in community composition were tested by permutational multivariate analysis of variance (PERMANOVA) with the *adonis* function in the *vegan* package (61).

To assess strain-level variation in the fecal microbiota and its potential ecological relevance, we rarefied samples to an even sequencing depth of 9,400 reads, matching the smallest sample size. We selected three phylotypes identified to species level that contained at least five unique ASVs within each sample type, *Klebsiella pneumoniae*, *Bacillus subtilis*, and *Lactobacillus buchneri*, as well as *Caulobacteraceae* sp., which reached high relative abundance in fecal microbiota from apple environments (up to 80% of bacterial reads; Fig 2). These taxa were analyzed for *alpha* and *beta* diversity at the strain (ASV) level, following the procedures described above, to explore whether substrate-dependent environments are associated with fine-scale genetic variation in dominant symbionts.

Compound specific isotope analysis of amino acids

Compound-specific isotope analysis (CSIA) of essential amino acids (EAAs) was performed using fly and substrate material from the October 2021 run of the larval development assay (see below). Within each treatment, fly samples were pooled by stratified random sampling (n = 8 per pool) to obtain at least 2 mg dry weight (following Larsen et al. 2013), resulting in five replicate pools per treatment. Developmental substrates from these assays were frozen, pre-dried for 24 h at 65 °C, lyophilized, and pooled accordingly. Fresh substrate samples were also dried (48 h at 65 °C) and lyophilized. For each substrate type (apple, tomato, and additionally, standard laboratory diet), five replicates (10-15 mg dry weight) were prepared for amino acid extraction. Flies and substrates were processed in separate extraction batches for synchronized handling, with samples randomized across treatments.

Amino acids were extracted and derivarized as described in Larsen et al. (62). Derivatized amino acids were analyzed in triplicate by gas chromatography–combustion isotope ratio mass spectrometry (GC-C-IRMS). They were injected into a Thermo Finnigan Trace 1310 gas chromatograph coupled to a Delta Plus IRMS (Finnigan, Bremen, Germany),

equipped with an Agilent J&W VF-35ms GC column (30 m × 0.32 mm × 1.00 µm). Carbon isotope values were corrected for derivatization (63) and expressed relative to Vienna Pee Dee Belemnite in δ-notation (‰). We obtained isotope values of 12 amino acids including alanine (*Ala)*, asparagine/aspartic acid (*Asx)*, glutamine/glutamic acid (*Glx)*, isoleucine (*Ile**), leucine (*Leu**), methionine (*Met**), phenylalanine (*Phe**), proline (Pro), serine (Ser), threonine (*Thr**), tyrosine (Tyr) and valine (*Val**); asterisks denote essential amino acids.

A linear discriminant analysis (LDA) was trained with $δ^{13}C$ values of essential amino acids from bacteria, fungi, and plants, published by Larsen et al. (38,62) and Pollierer et al. (64) to represent potential basal resources. Decision boundaries were constructed to project substrate, substrate post-eclosion, and fly samples into the LDA space, based on their essential amino acid profiles. To estimate the proportional contribution of basal resources used by flies we ran Bayesian mixing models implemented in R package MixSIAR (65) based on mean-centered $δ^{13}C$ values of essential amino acids in the potential basal resources as used in LDA and in the flies.

Larval development experiments

To ensure that variation in larval developmental success could be attributed solely to differences in substrate and symbiont inoculum conditions, all experiments were conducted with axenic larvae. *D. melanogaster* eggs were obtained overnight from the laboratory stock maintained on standard laboratory diet. The following morning, eggs were surface sterilized by dechorionation in 6% sodium hypochlorite for 10 min, followed by repeated rinses, first with ethanol and then with autoclaved tap water. Embryos were transferred to sterile glucose agar plates and incubated at 25 °C for 24 h to allow hatching. Newly emerged first-instar larvae were then used in the development assays. Smear-plating of freshly hatched larvae showed no visible microbial growth on either bacterial- or fungal-selective agar (S1 Text for media composition). Because all larvae originated from the same large, outbred laboratory population, there was no prior adaptation to either substrate or inoculum. Any observed interaction effects would therefore reflect differences in how microbial communities from different origins function within distinct substrate environments, rather than genetic or epigenetic differences among flies.

Across five independent experimental runs (March 2020, September 2020, March 2021, June 2021, and October 2021), we tested the effects of fecal microbial inocula from "apple" and "tomato" flies, transferred to either autochthonous or allochthonous fruit substrates (see above). Larval development was quantified in four of these runs, while the June 2021 run was dedicated exclusively to substrate microbiota analysis. The October 2021 run additionally included amino acid analysis (S1 Table for sample details).

Organic apples (*Holsteiner Cox*) and vine tomatoes were purchased from a local store (ALECO, Bremen). Apples were peeled and cored, while seeds were removed from tomatoes. Fruits were homogenized to produce a semi-liquid substrate, which was microwaved (800 W) three times until boiling commenced. This treatment standardized substrate texture and moisture content while substantially reducing the initial microbial load prior to inoculation. Plating of the prepared substrates showed no visible microbial growth on either bacterial- or fungal-selective agar.

For each experimental unit, 1 mL of prepared fruit substrate was pipetted into a sterile 2
mL collection tube, followed by 50 µL of fecal suspension originating from "apple" or
"tomato" flies. A single axenic larva was then transferred into each tube, and the tubes
were sealed with autoclaved dental cotton rolls. Experimental units were placed in racks
and incubated under the same environmental conditions as the laboratory stock and
microcosms (see above). The following day, all units were checked for the presence of
alive larva to ensure that only replicates unaffected by accidental mortality during
transfer were included in subsequent analyses. The total number of prepared replicates
for each run, the origin of the fecal microbiota, and the final number used in statistical
analyses are provided in S1 Table. Replicates that passed this quality check were
monitored daily to record the number of days from larval transfer to adult emergence.
Newly emerged flies were sexed, immediately frozen, and dried over silica gel before
weighing on a precision scale. The June 2021 run, designed to generate samples for
metabarcoding of the developmental substrate, used only female fecal microbiota from
"tomato" flies to establish *autoch*SI or *alloch*SI conditions; developmental parameters
from this assay were therefore excluded from analysis. In the October 2021 run, we
included an additional treatment with standard laboratory diet without microbial
inoculation, enabling comparison of essential amino acid profiles between flies reared
on fruit substrates and those reared on the laboratory diet. Larval development on the
standard laboratory diet was not further analyzed.

Statistical analysis

Our statistical approach was designed to test the hypothesis that host developmental
traits are shaped by symbiont–environment interactions, i.e., that the effect of microbial
inoculum origin (*autoch*SI vs. *alloch*SI) on *Drosophila* development depends on the
substrate in which larvae develop. Interaction terms between substrate type, inoculum
origin, and sex were therefore explicitly included in all models, as significant interaction
effects would indicate the presence of such context-dependent symbiont effects.

Model fit was first assessed using data from the initial experimental run for both adult
weight and development time. For both traits, a full model including all interaction terms
provided an adequate description of the data. We then analyzed data from all four
experimental runs, with "run" initially included as a random effect to account for
potential variation among runs. For adult weight, a linear mixed-effects model was fitted
first. The intraclass correlation coefficient for "run" was low (ICC = 0.03), indicating that
between-run variance was negligible. We therefore proceeded with a standard linear
model for the final analysis, using z-transformed weight data. The final model structure
was:

Fly weight ~ Substrate × SI condition × Sex

407 For development time, data were Box–Cox transformed to meet model assumptions and
408 analyzed using a linear mixed-effects model using *lmerTest* (66) with the same fixed-
409 effects structure, yet experimental run was included as a random effect (ICC = 0.45):

Development time ~ Substrate × SI condition × Sex + (1|Run)

411 Across all treatments, sample sizes were *n* = 817 for development time and *n* = 796 for
412 weight (exact numbers by treatment are given in S1 Table). Variation in replicate numbers
413 arose from larval mortality and occasional measurement failures. Ninety-five percent

confidence intervals for predicted means were calculated from fitted model values using the *predictInterval* function in the *merTools* package (67), based on 1,000 simulated datasets. All statistical analyses of the development assays were conducted in R version 4.2.2 (R Core Team, 2022) using the RStudio interface (version 2022.07.02+576; RStudio Team, 2022).

Figure design

Graphical outputs involving data from either metabarcoding, CSIA or larval development experiments were generated with the ggplot2 package (68) and adapted in Inkscape v1.1.0 using original designs. Fig 1 was designed in Inkscape. Accessibility of graphics was assessed using Color Oracle 1.3.0.

Results

Symbiotic microbiota in the feces of fly populations differ between breeding environments

To assess whether the symbiotic microbiota transmitted by female *D. melanogaster* to egg-laying sites differ by breeding environment, we analyzed fecal material from flies reared in separate populations maintained on distinct substrates (apple or tomato). Across both substrate environments, the fecal material from "apple" and "tomato" flies contained seven bacterial families that comprised at least 80% of the total bacterial microbiota. These families were Bacillaceae, Burkholderiaceae, Caulobacteraceae, Enterobacteriaceae, Lactobacillaceae, Rhizobiaceae, and Sphingomonadaceae. The predominant (at least 70% of all samples) fungal members of the microbiota were yeasts from the order Saccharomycetales (specifically *Pichia* and *Starmerella* species), as well as some *Geotrichum* and filamentous *Penicillium* species (Fig 2, Fig S2). The *alpha* diversity of bacteria showed only marginal difference between symbionts derived from apple and tomato flies (Fig 3A); however, this could be due to between-replicate variability in phylotype numbers (Fig S3). Notably higher diversity among fungal symbionts was observed in the apple flies compared to the tomato flies (Fig 3B, Fig S3).

Fig 2. Breeding environment determines *Drosophila melanogaster* symbiont community composition.

Heatmaps showing the nine most abundant bacterial and fungal taxa found in the fecal material of *Drosophila melanogaster* females from apple and tomato environments (n = 5). The ITS2 region and the 16S ribosomal subunit were used as marker genes for identification of fungal and bacterial communities. Amplicon sequence variants were agglomerated at species level and the nine most abundant taxa are displayed. All other taxa are grouped under "Other". The relative abundance of taxa in each sample is depicted as color intensity from blue to red as percentage of total reads.

454
Fig 3. Community composition of *Drosophila melanogaster* symbionts differs
between breeding environments.
Boxplots depicting observed and Shannon diversity of symbiont communities in
the fecal material of *Drosophila melanogaster* females from apple or tomato
environments (n = 5). The ITS2 region and the 16S ribosomal subunit were used as
marker genes to characterize fungal (A) and bacterial (B) communities,
respectively. Symbiont community differences between apple and tomato were
assessed using the Wilcoxon Rank-Sum Test. Principal Coordinate Analysis
(PCoA) based on Bray-Curtis dissimilarity showing the bacterial (C) and fungal (D)
symbiont community composition. Ellipses represent the 95% confidence
interval.

The composition of bacterial symbionts differed significantly between the substrate environments (PERMANOVA; $F_{1,8}$ = 4.03, p = 0.012, Fig 3C). Bacterial symbionts from "apple" flies were dominated by a species from the gram-negative Caulobacteraceae family (Fig 2). Also, the fungal symbiont composition exhibited distinctions between "apple" and "tomato" flies, although the differentiation was subtle (PERMANOVA; $F_{1,8}$ = 1.09, p = 0.156, Fig 3D). Notably, symbionts in the tomato environment were predominantly (70-80% of reads) composed of *Geotrichum silvicola* (Fig 2, Fig S2). Additionally, many phylotypes were found to consist of more than five strain variants (S2 Table). A deeper analysis of ASV microdiversity within the most divergent bacterial phylotypes revealed substrate-dependent differentiation in a Caulobacteraceae phylotype, *Klebsiella pneumoniae*, *Bacillus subtilis*, and the *Lactobacillus buchneri* phylotype (Fig 4). This indicates that environment-driven differentiation in *Drosophila* symbionts is evident not only at the phylotype level but also at the level of intraspecific variation.

Symbiont community composition changes during *Drosophila* development under *autoch*SI and *alloch*SI conditions

Next, we investigated the consequences of autochthonous (*autoch*SI) versus allochthonous symbiont inoculation (*alloch*SI) for microbiota composition in the larval environment. While *autoch*SI conditions in a tomato environment resulted in substrate bacterial and fungal microbiota similar to the initial fecal symbionts, *alloch*SI in an apple environment leads to a quite distinct microbial community (Fig 5). The shift in symbiont composition during larval development on the apple substrate was characterized by an increase in the relative abundance of *Pichia* yeasts (Fig 5) and the dominance of the Caulobacteraceae phylotype (Fig 5).

Source-specific contribution of essential amino acids (EAA) during *Drosophila* development depends on symbiont inoculum conditions

We performed compound specific isotope analysis of EAA in two contexts: 1) the breeding substrate after fly eclosion and 2) freshly emerged *Drosophila* females. This analysis aimed to characterize changes in EAA composition within the habitat during larval development and to identify the sources of these EAA – whether derived from bacteria, fungi, or plant substrates (apple or tomato) – utilized by or available to the insects.

During *Drosophila* larval development, the EAA profiles of both plant substrates shifted toward profiles resembling those of bacteria and fungi (Fig 6A, Fig S4). However, compared to apple, tomato substrates tended to be more dominated by bacteria-derived EAA, particularly under *autoch*SI conditions. Notably, EAA profiles on both substrates were influenced by symbiont inoculum conditions (Fig 6A). On apple, *alloch*SI conditions caused a pronounced shift of substrate EAA profile toward bacteria derived EAA.

Fig 4. Microdiversity in bacterial symbionts of *Drosophila melanogaster* differs between breeding environments.

Strain level diversity was analyzed in four selected bacterial phylotypes found in the fecal material of female *Drosophila melanogaster* reared in both apple and tomato environments. *Alpha*-diversity is shown as Observed and Shannon diversity, and environmental effects were analyzed by means of Wilcoxon rank sum test. Principal Coordinate Analysis (PCoA) based on Bray-Curtis dissimilarity was used to visualize differences in strain community composition (*beta*-diversity), ellipses represent 95% confidence. PERMANOVA were applied to identify influences of the insect breeding environment.

Fig 5. Allochthonous microbiota transmission alters larval *Drosophila melanogaster* symbiont composition.

Relative abundance of the 10 most abundant bacterial (A) and fungal (B) symbionts in the fecal inoculum (single stacked bars, left) and on the autochthonous (tomato) and allochthonous (apple) substrates following larval development. *alpha*-diversity (Observed and Shannon) of symbionts following autochthonous and allochthonous inoculum and larval development (C, D). Principal Coordinate Analysis (PCoA) based on Bray-Curtis dissimilarity was used to visualize differences in bacterial and fungal community composition (*beta*-diversity) following larval development on the autochthonous (tomato) and allochthonous (apple) substrates, ellipse represent 95% confidence. PERMANOVA was applied to identify influences of the breeding substrate (E, F).

Fig 6. Symbiont inoculum conditions affect essential amino acid (EAA) flow during *Drosophila melanogaster* development.
(A) Centroids calculated based on a linear discriminant analysis (LDA) of EAA-specific $\delta^{13}C$ values of fresh tomato and apple, and substrates following larval development under different symbiont inoculum conditions. (B) Fly EAA profiles following development under autochthonous and allochthonous symbiont inoculum condition. The ellipses represent the positioning of EAA profiles from different plant, bacterial, and fungal sources, respectively. They were used as classifiers in LDA (64). See Figure S4 for polygons depicting variation in EAA profiles of substrates and flies.

Interestingly, the EAA profiles of the flies showed only little resemblance to those of the substrates following fly eclosion; in fact, they remained similar to the EAA profiles of fresh substrate (Fig 6). Under *alloch*SI conditions on tomato, fly and substrate EAA profiles following fly eclosion clustered similarly relative to each other. In contrast, fly EAA profiles under *alloch*SI conditions on apple, resembled those of fresh tomato substrate. Notably, the EAA profiles of flies reared on a standard laboratory diet showed no overlap with those developed under semi-natural conditions on either apple or tomato substrates, irrespective of symbiont inoculum (Fig S4).

Independent of symbiont inoculum conditions, tomato substrate following fly eclosion and respective fly EAA profiles were lower on LD2 compared to fresh substrates, i.e. closer to microbial resources (Fig 6). The EAA profiles of both the tomato substrate and the eclosed flies tended toward a bacterial profile under *autoch*SI (LD1, negative) and toward a fungal profile under *alloch*SI conditions (LD1, positive). Apple substrate and the respective flies did not show these similarities in EAA profiles. Compared to fresh apple substrate EAA profiles of apple substrate after fly eclosion followed the same vector as the tomato substrates post eclosion. EAA profiles of flies that developed on apple did not follow these vectors.

Regardless of the plant substrate and symbiont inoculum conditions, both tomato and apple consistently served as the primary source of EAA for flies, contributing on average between over 50% to more than 75% (Fig 7). When developing in an apple environment, bacteria contributed slightly more than fungi to the EAA in flies that fed and grew as larvae under both *autoch*SI and *alloch*SI conditions. However, under *alloch*SI conditions, the relative contribution of bacterial and fungal EAA dropped to well below 20%, resulting in the contribution of the apple substrate increasing to more than 75%. In the tomato environment, symbiont inoculum conditions hardly affected EAA incorporation by the larvae.

Fig 7. Symbiont inoculum conditions influence essential amino acid (EAA) uptake in *Drosophila melanogaster* larvae.

Estimated relative contribution (%) of bacteria, fungi, and the plant substrates to EAA assimilated by larvae under autochthonous and allochthonous symbiont inoculum conditions. EAA were extracted from freshly eclosed female flies. Estimates are based on a Bayesian mixing model of mean-centered $\delta^{13}C$ values and displayed with 95% confidence interval.

Expression of *Drosophila* life-history traits depends on symbiont inoculum conditions

We investigated whether environment-dependent differences in fecal symbionts and observed shifts in microbiota composition in response to *alloch*SI impact *Drosophila* larval development. In a series of independent experiments (see Methods for details), we

found a consistent pattern of effects of *autoch*SI versus *alloch*SI conditions on adult fly body weight and development time, respectively. In the presence of symbionts from an apple environment, fly weight remained consistent regardless of sex or symbiont inoculum conditions (Fig 8). Interestingly, larval development was faster under *alloch*SI conditions compared to *autoch*SI conditions. However, in an apple environment, *alloch*SI conditions resulted in a significant decrease in body weight and simultaneously prolonged development times in both female and male flies (Fig 9). Given the significant effect of the statistical interaction between breeding substrate and symbiont origin on both body weight and development time (Table 1 and 2), these results indicate a symbiont-environment interaction.

Fig 8. Biomass gain during development of *Drosophila melanogaster* larvae is affected by symbiont inoculum conditions.

Body weight of freshly emerged male and female *Drosophila melanogaster* reared under different symbiont inoculum conditions. Larvae developed on either apple or tomato substrates inoculated with fly-derived fecal microbiota from an autochthonous (*autoch*SI) or allochthonous (*alloch*SI) environment. Raw data were z-transformed to account for baseline differences in average body weight across four experimental runs. Estimates (filled symbols) and corresponding 95% confidence intervals were generated using a general linear model. *n* indicates the number of individual flies measured. See Table 1 for Type III ANOVA.

Fig 9. Development time of *Drosophila melanogaster* larvae is affected by symbiont inoculum conditions.

Development time of male and female *Drosophila melanogaster* reared on apple and tomato substrates inoculated with fly fecal microbiota from either autochthonous (*autoch*SI) or allochthonous (*alloch*SI) environments. Development time was measured in days from first instar larva to adult eclosion. Estimates were generated for each combination of substrate, symbiont inoculum, and sex using a linear mixed-effects model that corrected for variation across experimental runs (*n* = 4 per condition, *n* = 2 sexes). Error bars represent 95% confidence intervals. *n* indicates the number of individual flies for which development could be recorded. See Table 2 for Type III ANOVA.

Table 1. Type III ANOVA result for a general linear model testing predictors of z-transformed adult *Drosophila melanogaster* weight (mg).

Explanatory variable	Sum of squares	DF	F-value	p-value
Substrate	5.82	1	9.08	0.003
***SI* condition**	73.89	1	115.29	<0.001
Sex	88.28	1	137.74	<0.001
Substrate x *SI* condition	50.06	1	78.11	<0.001
Substrate x Sex	2.89	1	4.52	0.034
***SI* condition x Sex**	13.25	1	20.67	<0.001
Substrate x *SI* condition x Sex	6.87	1	10.72	0.001
Residuals	505.03	788		

Substrate: apple or tomato. **SI**: symbiont inoculum condition, either autochthonous or allochthonous

Table 2: Type III ANOVA result for a linear mixed model testing predictors of Box-Cox transformed development time in *Drosophila melanogaster* (days). Experimental run was specified as random factor.

Explanatory variable	Sum of squares	DF (num)	DF (den)	F-value	p-value
Substrate	8.547e-05	1	806.083	411.179	<0.001
***SI* condition**	9.017e-06	1	806.025	43.380	<0.001
Sex	4.530e-06	1	806.022	21.795	<0.001
Substrate x *SI* condition	1.829e-05	1	806.086	87.994	<0.001
Substrate x Sex	4.901e-08	1	806.036	0.236	0.627
***SI* condition x Sex**	9.406e-07	1	806.013	4.525	0.034
Substrate x *SI* condition x Sex	9.983e-07	1	806.126	4.803	0.029

Substrate: apple or tomato. **SI**: symbiont inoculum condition, either autochthonous or allochthonous

The relatively long development time and low body weight of female flies reared as larvae under *alloch*SI conditions on apple correlate with the highest contribution of plant-derived EAA and the lowest contribution of fungus-derived EAA to their EAA profile (Fig 7, Fig S5). Variation in bacteria-derived EAA provides limited explanatory power for differences in insect life-history traits.

Discussion

In this study, we provide evidence of symbiont-environment interactions affecting the developmental success of *D. melanogaster*. We define symbiont-environment interactions as a situation where the influence of microbial symbiont composition on host development depends on the ecological context of the breeding environment. We explored this experimentally by simulating autochthonous and allochthonous symbiont scenarios within the lifecycle of *D. melanogaster* (Fig 1). To bridge the gap between laboratory and natural conditions, we established fly populations in semi-natural microcosms using fruit substrates (apple and tomato), where field-derived microbiota was transmitted and maintained across multiple generations. This approach allowed us to observe consistent environment-dependent variation in symbiont communities, both in adult females and in larval developmental substrates. By linking symbiont composition, essential amino acid uptake, and life-history trait expression, we present a novel approach to investigate environmental heterogeneity shaping host-microbe interactions in the *D. melanogaster* model system.

The breeding environment determines bacterial and fungal symbiont diversity of *D. melanogaster*

Our results experimentally verify the finding of observational studies (28,32,69) that the plant substrate defining the *Drosophila* breeding habitat is a fundamental driver of the microbiota forming a symbiotic relationship with both the adult flies and the larvae. The deposition of fecal material near oviposition sites is a primary route through which larvae acquire microbial symbionts (25,27). The fecal material deposited by female flies reared in a tomato environment tended to contain more diverse bacterial communities than those reared in an apple environment. Despite variation in bacterial *alpha*-diversity across years and transmission histories (Fig S3), substrate environment consistently explained differences in community composition (*beta*-diversity). First, this demonstrates a robust, environment-dependent association of bacterial symbionts across generations. Second, it challenges the common view of *Drosophila* microbiota as "simple" e.g., Ludington and Ja (70), revealing substantial variation shaped by ecological context.

While several Enterobacteriaceae phylotypes and a Caulobacteraceae phylotype dominated the fecal microbiota of flies from an apple environment, the community composition in the fecal material of flies from a tomato environment appeared more even with higher proportions of Burkholderiaceae and Lactobacillaceae, among others. Except for the Caulobacteraceae phylotype, the families and phylotypes identified in both populations generally align with those reported to be fly associated in nature by other studies (14,18,71–74). Surprisingly, we did not detect a dominance of Acetobacteraceae, often reported in other studies (18,71,72,74). Acetobacteraceae phylotypes have previously been shown to establish a stable association with *Drosophila*, supported by gut-specific conditions that promote their growth (24). Some of the mechanisms that enable such stable gut-bacterial associations in *Drosophila* have only recently been revealed (75). These mechanisms involve a complex interplay of host–microbe and microbe–microbe interactions that shape a specialized niche in the proventriculus of adult flies, ultimately favoring the persistence of microbial taxa capable of propagating and establishing stable associations with the host. Dodge et al (75) links the establishment of stable symbiosis to chemical alterations of the fly gut by microbial partners. As our study demonstrates, the pool from which microbes may be selected varies greatly across environments, leading to the establishment of distinct symbiont communities. Accordingly, we expect that the mechanisms identified by Dodge et al. (75)are sufficiently adaptable to respond to the substrate environment's influence on shaping the microbiota ingested by flies. The dominant bacterial taxa transmitted by flies in our study likely possess traits that enable their selection through these mechanisms, possibly facilitating their integration into stable, habitat-dependent mutualistic relationships (see also below).

In addition to environment-dependent differences in phylotype diversity, a more detailed analysis of amplicon sequence variants (ASVs) revealed microdiversity within several bacterial phylotypes, including *Klebsiella pneumoniae, Bacillus subtilis, Lactobacillus buchneri* and a Caulobacteraceae phylotype. Notably, both number of strains and strain composition analyses showed, that the composition of ASVs, within single phylotypes in the fecal material of female *Drosophila,* can vary depending on the substrate

environment in which the flies were kept. In a study on the honeybee gut microbiome,
Baud et al. (76) link microdiversity differences between nursing and foraging bees to
dietary shifts, specifically the cessation of pollen consumption in foragers. Similarly, a
cautious interpretation of our results could be that within-phylotype genetic variation in
a given environment undergoes selection, favoring variants that are best adapted to thrive
in and be transmitted by flies between substrate patches of a specific kind. This finding
thus highlights two important novel aspects: First, microdiversity in bacterial symbionts
must be considered in *Drosophila* microbiome analyses to uncover their potential impact
on host phenotypic trait expression. Second, the ecological context – specifically, the
breeding substrate environment – should be regarded as a potential selective filter for
ASV composition. The extent to which host control mechanisms (see above) additionally
contribute to the maintenance and differentiation of microdiversity remains to be
determined.

Fungal symbiont diversity in the feces of female *Drosophila* was also influenced by the
substrate environment. The *alpha*-diversity analysis revealed greater fungal diversity in
the fecal material of flies reared in an apple environment compared to those in a tomato
environment, while *beta*-diversity analysis showed only marginal differences. Fungal
symbionts in "apple" flies were dominated by frequently found species of *Starmerella*
and *Pichia* (e.g.,23,25,76) which were nearly absent in the "tomato" flies. Most strikingly,
Geotrichum silvicola overwhelmingly dominated the fungal community in the feces of
flies from the tomato environments. *G. silvicola* was originally isolated from insects,
including *Drosophila* (78). Furthermore, various *Geotrichum* species, including those
that are plant pathogens, are frequently found in association with *Drosophila*
(73,77,79,80). Since fly populations thrived in our tomato microcosms, *Geotrichum*,
which exhibits morphological traits intermediate between yeasts and filamentous mold
fungi (81), does not seem to be pathogenic for the insects.

Although fungi can persist in the flies' gut system for an extended period (25), there is
currently no evidence that they establish a long-term association without ongoing
replenishment. This suggests that fungal symbiont communities are likely to change
when the flies' breeding environment changes. Supporting evidence comes from Gurung
et al. (82), who found that fungi associated with *D. suzukii* were similarly associated to
their breeding substrate, unlike their bacterial symbionts. Since we observed effects of
the breeding substrate on both fly-associated bacteria and fungi, we cannot confirm that
bacteria are more persistently associated with *D. melanogaster*. A possible explanation
could be that apples and tomatoes differ more significantly from each other in their
properties (see below) than the different types of berries used by Gurung et al. (82). The
plant substrate can be a very strong filter for the symbionts that establish in the flies'
breeding site, which becomes apparent when comparing symbiont community
development under autochthonous inoculation (*autoch*SI) conditions to that under
allochthonous inoculation (*alloch*SI) conditions (Fig 1), as discussed below.

Under *alloch*SI conditions, we observed a significant shift in bacterial symbionts from
"tomato" flies that established on apple substrate. Unlike the *autoch*SI conditions, which
led to a community similar to the initial inoculate, bacterial diversity on apple was notably
reduced and characterized by a pronounced shift towards a Caulobacteraceae
phylotype. In contrast, fungal phylotypes slightly increased under *alloch*SI conditions,

however this was just a trend. This demonstrates how common environmental variation
in the *Drosophila* habitat can lead to substantial changes in symbiont composition.

In conclusion, the breeding substrate environment of *Drosophila* influences both
bacterial and fungal symbionts. This is evident for symbionts transmitted by adult flies
from different habitats, as well as for those that grow in the presence of larvae under
*alloch*SI versus *autoch*SI conditions. While we can currently only speculate on the
mechanisms underlying these differences, it likely involves significant variation in
substrate acidity, water availability, and nutrient content. For example, we found the
substrate pH in apple and tomato substrates to vary between 2.6 - 3.2, and 3.6 - 4.4,
respectively (S3 Table). Oakeshott (31) identified a correlation between substrate-
dependent differences in pH, the abundance of bacteria and different kinds of yeast
fungi, and the distribution of various *Drosophila* species. Although substrate properties
directly shape microbe–microbe interactions that influence host performance(22,83–
86), both adult and larval flies can further modulate symbiont composition (e.g. Stamps
et al. 2012). This has recently been shown in *Ceratitis capitata* (Tephritidae) harboring a
microbiota strongly influenced by host fruit but regulated through life-stages and host
functional needs (87). Such effects may arise through individual immune traits (88), the
gut's capacity to lyse or promote specific microbes (24,75,89), and cooperative
interactions within larval aggregations (44,90,91). It remains unknown if host traits and
system properties allow *D. melanogaster* to exert a density-dependent "leash" on its
microbiome (9).

Essential amino acid contribution to *D. melanogaster* tissue depends on plant substrate-symbiont combination

One of the most relevant "services" of symbionts is to promote host fitness through
improving dietary conditions, such as biopolymer degradation, provisioning of essential
vitamins and amino acids (2,92). A similar importance of symbiont microbes providing
essential amino acids (EAA) in an otherwise (artificial) low-protein environment has been
demonstrated for both larval *Drosophila* development, and adult lifespan and
reproduction (19,93). To the best of our knowledge, combining our semi-natural
experimental approach with stable isotope analysis of EAA in freshly emerged flies has
not been done so far. Thus, we were able to reveal for the first time the actual sources of
EAA incorporated during larval development. High proportion of EAA found in the flies
indeed came from the plant substrate. This proportion even increased under *alloch*SI
conditions, whereas the proportion of EAA contribution from the plant to the insect tissue
was always higher in the presence of symbionts from a tomato environment. Possibly,
the environmental context-dependent shifts in EAA contributions from the three sources,
plant, bacteria and fungi, is due to substrate-induced changes in the prevalence of
nutrient-rich, nutrient-poor, unpalatable, or even pathogenic symbionts.

Interestingly, EAA profiles of flies did not resemble the EAA profile of the substrate after
larval development (see Fig 6). Again, differences in symbiont communities may create
nutritional niches that constrain EAA uptake. Furthermore, larvae may directly influence
EAA uptake through selective feeding (94). While the EAA profile of the substrate reflects
the endpoint of microbial succession during larval development, the larval EAA profile
results from the dietary intake of EAA throughout ontogeny. Therefore, the substrate EAA

profiles cannot represent microbial dynamics during larval development. These, still
unknown, dynamics in symbiont community succession – relative abundances and
absolute population sizes – may additionally explain the mismatch between substrate
and insect EAA profiles.

Lastly, we note that the EAA profiles of flies reared on an artificial laboratory diet showed
no overlap with those of flies developing in natural breeding substrates, regardless of
their associated symbionts. Although we can, beyond variation in body size (see below),
only speculate on the consequences for *Drosophila* nutritional physiology (95), our
findings show that the quality of microbial partners depends strongly on the breeding
substrate. This makes explicitly including ecological context essential for understanding
the selective environments that shape the evolution of both *Drosophila* life-history (21)
and its mutualisms with microorganisms (see below).

D. melanogaster phenotypic trait expression is shaped by symbiont-environment interactions

Symbiont-environment interactions, shaped by variation in *D. melanogaster* microbiota
across habitats and substrate-dependent dynamics *within* substrate patches, manifest
in significant differences in two key life-history traits: larval development time until adult
eclosion and adult body weight. Fly body weight was not uniformly affected by *alloch*SI
conditions, instead, our experiment revealed that larval weight gain was relatively lower
when symbionts originated from a tomato environment. In contrast, the effect of
symbionts from "apple" flies did not vary significantly between *autoch*SI and *alloch*SI
conditions. Development time was prolonged under *alloch*SI conditions in the presence
of microbiota from "tomato" flies. Interestingly, when larvae were reared with symbionts
from "apple" flies under *alloch*SI conditions, their development time was even
accelerated. These findings suggest that when substrate availability varies and female
flies transmit allochthonous microbes, symbiont-environment interactions can lead to
both fitness gains – such as faster development without a reduction in body mass – and
fitness losses, as seen in slower development and reduced body mass under *alloch*SI
conditions. Therefore, symbiont-environment interactions offer a new conceptual
framework for better understanding microbiota-driven phenotypic variation in
Drosophila, which in consequence determines the width of the reproductive niche of the
entire symbiotum.

As discussed above, the most striking outcome of the symbiont-environment interaction
observed in this study is the massively reduced body size and prolonged development
time under *alloch*SI conditions on apple. Possibly, this could be attributed to the higher
intake of EAA from the plant substrate, relatively to the intake from the microbial
symbionts. Since yeast fungi alone have been shown to provide a sufficient nutrient
source that promotes larval growth (25,94), the stability in body weight observed with
symbionts from apples may be due to a largely unaltered EAA contribution from fungi,
even under *alloch*SI conditions. The striking dominance of plant-derived EAA in flies that
developed under *alloch*SI conditions on apple may negatively impact larval development
through several potential mechanisms. One possibility is that symbiont-derived EAA and
other essential nutrients, such as steroids, become scarce due to reduced microbial
(fungal?) growth in the new environment. This forces larvae to rely on plant EAA, which

may exist in lower concentrations and be harder to extract from the plant matrix.
Additionally, certain *alloch*SI conditions may alter the symbiont community composition,
promoting the proliferation of low-quality or even pathogenic microbes (22). This shift
could trigger secondary metabolite-mediated dysbiosis, potentially disrupting larval
development (96,97). The observed changes in larval weight gain over time could also
result from a complex interplay between changes in symbiont communities and host
behavior. For instance, microbe-induced changes in feeding behavior (50) and the varying
prevalence of symbionts with beneficial or detrimental traits might collectively influence
development outcomes, and thus the reproductive niche space of the host (98). To
deepen our understanding of these dynamics, future research should focus on isolating
key microbial taxa, assessing their individual and combined effects on host traits, and
applying synthetic communities under controlled yet ecologically relevant conditions
that mirror the environments in which host-microbe interactions naturally evolve (15).

Implications for understanding the evolution of *Drosophila*-microbe interactions

The substrate-dependent differences in microbial community composition across
breeding environments indicate that mutualistic dependence between *Drosophila* and
its microbiota is largely facultative. Although larval development and adult reproduction
remain viable across a range of microbial assemblages (99), not all symbiont
communities contribute equally to host development. Variation in fitness outcomes
under *alloch*SI conditions suggests that microbial partners differ in the relative benefit
depending on ecological context. This supports the view that *Drosophila* interacts with its
microbial partners along a continuum of mutualistic quality, with weak dependence on
individual phylotypes likely limiting the evolution of specialization (10).

Crucially, while *Drosophila* larvae require microbial input for successful development,
this dependency is functional rather than obligate in the evolutionary sense.
Development relies on microbial functions, such as essential amino acid provisioning,
which can be fulfilled by a wide range of bacteria and fungi. This positions *Drosophila* as
an obligate generalist that cannot develop without microbial partners, but partner
breadth is potentially broad (10,11). For microbial symbionts, the unpredictability of fly
breeding environments and thus the lack of consistent transmission between host
generations limit the evolution of stable associations and cooperative traits. Most
identified symbionts can associate with a variety of invertebrate hosts and persist
independently in the environment e.g., Carramachi et al. (100). Altogether, these findings
suggest that ecological context constrains the evolution of specialization in *Drosophila*-
microbe interactions and, through broad partner breadth, stabilizes a functionally
essential yet evolutionarily loose form of mutualism.

Despite limited specialization in *Drosophila*-microbe interactions, a deeper
understanding of symbiont-environment dynamics offers a framework for identifying
microbial taxa with the potential to form stable, long-term associations across diverse
ecological contexts. For example, a single variant of the *Lactobacillus buchneri* phylotype
was detected in the fecal material of flies from diverse microcosm environments beyond
apple and tomato, including plum, grape, lemon, and onion (Riedel & Rohlfs 2025
unpublished). The broad occurrence of this variant suggests it may have the potential to

evolve into an obligate specialist under specific ecological conditions, a hypothesis that could be tested experimentally by evaluating its functional contributions to larval development across diverse substrates. A similar case was reported for *Acetobacter thailandicus* in *D. melanogaster* (24), where an isolate of *A. thailandicus* colonized the gut of several *D. melanogaster* genotypes, proliferating and providing sufficient nutrients to sustain the host on otherwise sterile figs. However, this interaction was only tested within the fig environment. Alternatively, rather than relying on specialized microbial partnerships, *Drosophila* may benefit from actively shifting breeding substrates to replace symbionts with limited functionality (10,11). As outlined previously, while *Drosophila* can influence its microbiota through individual behavior, social interactions, and physiology (18), it remains unclear to what extent this control can override the strong filtering imposed by the breeding environment. To disentangle relative roles of host control and environmental filtering will be essential for linking natural ecological dynamics to the evolution of host-microbe associations in this model system.

Lastly, our findings raise the possibility that variation in symbiont community composition contributes to host adaptation in novel environments (43,98,101). The faster development of flies under *alloch*SI conditions on tomato, compared to those reared with autochthonous microbes, suggests that non-local microbial communities may sometimes confer unexpected nutritional or physiological advantages. These results point to a broader role for flexible generalist symbioses in promoting adaptive plasticity, potentially explaining how *Drosophila* populations maintain breeding niche breadth across diverse and ephemeral plant substrates and may even expand into previously unexploited resources (102).

Our study highlights how ecological context shapes nutritional interactions between *D. melanogaster* and its microbial associates. Symbiont-environment interactions reveal context-dependent costs and benefits of symbiont-dependent development, underscoring the contingencies of generalist associations. This framework offers new directions for exploring the ecological limits of evolutionary processes in host-microbe interactions.

Acknowledgments

We thank Stefan Scheu for sharing knowledge and lab resources. We thank Guido Humpert and Nico Binnemann for technical support, and Jens Dyckmans for compound-specific isotope measurements of amino acids. Data availability of this study was supported by the Data Science Center of the University of Bremen (DSC@UB) funded by the State of Bremen.

References

1.   Tihelka E, Cai C, Giacomelli M, Lozano-Fernandez J, Rota-Stabelli O, Huang D, et
al. The evolution of insect biodiversity. Curr Biol. 2021 Oct;31(19):R1299–311.

2.   Douglas AE. Insects and Their Beneficial Microbes. Vol. 1st Edition. Princeton
University Press; 2022.

3.   Feldhaar H. Bacterial symbionts as mediators of ecologically important traits of
insect hosts. Ecol Entomol. 2011;36(5):533–43.

4.   Kaufman MG, Walker ED, Odelson DA, Klug MJ. Microbial Community Ecology &
Insect Nutrition. Am Entomol. 2000 July 1;46(3):173–85.

5.   Douglas AE. Lessons from Studying Insect Symbioses. Cell Host Microbe. 2011
Oct;10(4):359–67.

6.   Douglas AE. Multiorganismal Insects: Diversity and Function of Resident
Microorganisms. Annu Rev Entomol. 2015 Jan 7;60(1):17–34.

7.   Oliver KM, Martinez AJ. How resident microbes modulate ecologically-important
traits of insects. Curr Opin Insect Sci. 2014 Oct;4:1–7.

8.   Flórez LV, Biedermann PHW, Engl T, Kaltenpoth M. Defensive symbioses of
animals with prokaryotic and eukaryotic microorganisms. Nat Prod Rep. 2015 June
26;32(7):904–36.

9.   Foster KR, Schluter J, Coyte KZ, Rakoff-Nahoum S. The evolution of the host
microbiome as an ecosystem on a leash. Nature. 2017 Aug;548(7665):43–51.

10.   Batstone RT, Carscadden KA, Afkhami ME, Frederickson ME. Using niche breadth
theory to explain generalization in mutualisms. Ecology. 2018 May;99(5):1039–50.

11.   Chomicki G, Kiers ET, Renner SS. The Evolution of Mutualistic Dependence. Annu
Rev Ecol Evol Syst. 2020 Nov 2;51(Volume 51, 2020):409–32.

12.   Butterworth NJ, Benbow ME, Barton PS. The ephemeral resource patch concept.
Biol Rev. 2023 June 1;98(3):697–726.

13.   Borges RM. Co-niche construction between hosts and symbionts: ideas and
evidence. J Genet. 2017 July 1;96(3):483–9.

14.   Lemoine MM, Wöhner T, Kaltenpoth M. Microbial Community Dynamics in
Natural Drosophila melanogaster Populations Across Seasons. Environ Microbiol.
2025;27(6):e70104.

15.   Holt JR, Cavichiolli de Oliveira N, Medina RF, Malacrinò A, Lindsey ARI. Insect–
microbe interactions and their influence on organisms and ecosystems. Ecol Evol.
2024;14(7):e11699.

16.   Atkinson W, Shorrocks B. Breeding Site Specificity in the Domestic Species of
Drosophila. Oecologia. 1977;29(3):223–32.

17.   Markow TA, O'Grady P. Reproductive ecology of Drosophila. Funct Ecol.
2008;22(5):747–59.

18.   Broderick NA, Lemaitre B. Gut-associated microbes of Drosophila melanogaster.
Gut Microbes. 2012 July 14;3(4):307–21.

19.   Yamada R, Deshpande SA, Bruce KD, Mak EM, Ja WW. Microbes Promote Amino
Acid Harvest to Rescue Undernutrition in Drosophila. Cell Rep. 2015 Feb
17;10(6):865–72.

20.   Steck K, Walker SJ, Itskov PM, Baltazar C, Moreira JM, Ribeiro C. Internal amino
acid state modulates yeast taste neurons to support protein homeostasis in
Drosophila. Ramaswami M, editor. eLife. 2018 Feb 2;7:e31625.

21.   Flatt T. Life-History Evolution and the Genetics of Fitness Components in
Drosophila melanogaster. Genetics. 2020 Jan 1;214(1):3–48.

22.   Caballero Ortiz S, Trienens M, Pfohl K, Karlovsky P, Holighaus G, Rohlfs M.
Phenotypic responses to microbial volatiles render a mold fungus more susceptible
to insect damage. Ecol Evol. 2018 Apr 1;8(8):4328–39.

23.   Stamps JA, Yang LH, Morales VM, Boundy-Mills KL. Drosophila Regulate Yeast
Density and Increase Yeast Community Similarity in a Natural Substrate. Kytöviita
MM, editor. PLoS ONE. 2012 July 31;7(7):e42238.

24.   Pais IS, Valente RS, Sporniak M, Teixeira L. Drosophila melanogaster establishes
a species-specific mutualistic interaction with stable gut-colonizing bacteria. Moran
N, editor. PLOS Biol. 2018 July 5;16(7):e2005710.

25.   Cho H, Rohlfs M. Transmission of beneficial yeasts accompanies offspring
production in Drosophila—An initial evolutionary stage of insect maternal care
through manipulation of microbial load? Ecol Evol. 2023 June 1;13(6):e10184.

26.   Mure A, Sugiura Y, Maeda R, Honda K, Sakurai N, Takahashi Y, et al. Identification
of Core Yeast Species and Microbe-Microbe Interactions Impacting Larval Growth of
Drosophila in the Wild. eLife. 2023 Sept 12;12.

27.   Guilhot R, Xuéreb A, Lagmairi A, Olazcuaga L, Fellous S. Microbiota acquisition
and transmission in Drosophila flies. iScience. 2023 Sept 15;26(9):107656.

28.   Chandler JA, Morgan Lang J, Bhatnagar S, Eisen JA, Kopp A. Bacterial
Communities of Diverse Drosophila Species: Ecological Context of a Host–Microbe
Model System. Malik HS, editor. PLoS Genet. 2011 Sept 22;7(9):e1002272.

29.   Chandler JA, Eisen JA, Kopp A. Yeast Communities of Diverse Drosophila
Species: Comparison of Two Symbiont Groups in the Same Hosts. Appl Environ
Microbiol. 2012 Oct 15;78(20):7327–36.

30.   Staubach F, Baines JF, Künzel S, Bik EM, Petrov DA. Host Species and
Environmental Effects on Bacterial Communities Associated with Drosophila in the
Laboratory and in the Natural Environment. Quince C, editor. PLoS ONE. 2013 Aug
13;8(8):e70749.

31.   Oakeshott JG, Vacek DC, Anderson PR. Effects of microbial floras on the
distributions of five domestic Drosophila species across fruit resources. Oecologia.
1989 Mar 1;78(4):533–41.

32.   Markow TA, Anwar S, Pfeiler E. Stable isotope ratios of carbon and nitrogen in
natural populations of Drosophila species and their hosts. Funct Ecol. 2000
Apr;14(2):261–6.

33.   Whiteman JP, Elliott Smith EA, Besser AC, Newsome SD. A Guide to Using
Compound-Specific Stable Isotope Analysis to Study the Fates of Molecules in
Organisms and Ecosystems. Diversity. 2019 Jan;11(1):8.

34.   Pollierer MM, Scheu S. Stable isotopes of amino acids indicate that soil
decomposer microarthropods predominantly feed on saprotrophic fungi. Ecosphere.
2021;12(3):e03425.

35.   Vesala R, Ghosh S, Murphy R, Poulsen M, Schmidt S, Mwangala L, et al.
Nutritional dynamics in fungus-growing termites: Integration of biological nitrogen
fixation and microbial amino acids. Funct Ecol. 2025 July 11;00:1–15.

36.   Pollierer MM, Larsen T, Potapov A, Brückner A, Heethoff M, Dyckmans J, et al.
Compound-specific isotope analysis of amino acids as a new tool to uncover trophic
chains in soil food webs. Ecol Monogr. 2019 Nov;89(4).

37.   Larsen T, Ventura M, Andersen N, O'Brien DM, Piatkowski U, McCarthy MD.
Tracing Carbon Sources through Aquatic and Terrestrial Food Webs Using Amino Acid
Stable Isotope Fingerprinting. PLOS ONE. 2013 Sept 17;8(9):e73441.

38.   Larsen T, Taylor DL, Leigh MB, O'Brien DM. Stable isotope fingerprinting: a novel
method for identifying plant, fungal, or bacterial origins of amino acids. Ecology. 2009
Dec 1;90(12):3526–35.

39.   Fink C, Staubach F, Kuenzel S, Baines JF, Roeder T. Noninvasive Analysis of
Microbiome Dynamics in the Fruit Fly Drosophila melanogaster. Appl Environ
Microbiol. 2013 Nov 15;79(22):6984–8.

40.   Kowallik Vienna, Mikheyev Alexander S. Honey Bee Larval and Adult Microbiome
Life Stages Are Effectively Decoupled with Vertical Transmission Overcoming Early
Life Perturbations. mBio. 2021 Dec 21;12(6):e02966-21.

41.   Manthey C, Metcalf CJE, Monaghan MT, Steiner UK, Rolff J. Rapid growth and the
evolution of complete metamorphosis in insects. Proc Natl Acad Sci. 2024 Sept
17;121(38):e2402980121.

42.   Joy JB. Symbiosis catalyses niche expansion and diversification. Proc R Soc B
Biol Sci. 2013 Apr 7;280(1756):20122820.

43.   Lemoine MM, Engl T, Kaltenpoth M. Microbial symbionts expanding or
constraining abiotic niche space in insects. Curr Opin Insect Sci. 2020 June 1;39:14–
20.

44.   Biedermann PH, Rohlfs M. Evolutionary feedbacks between insect sociality and
microbial management. Vectors Med Vet Entomol Soc Insects. 2017 Aug 1;22:92–
100.

45.   Henry LP, Bruijning M, Forsberg SKG, Ayroles JF. The microbiome extends host
evolutionary potential. Nat Commun. 2021 Aug 26;12(1):5141.

46.   Wölfle S, Trienens M, Rohlfs M. Experimental evolution of resistance against a
competing fungus in Drosophila melanogaster. Oecologia. 2009 Oct 1;161(4):781–90.

47.   Fink C, Von Frieling J, Knop M, Roeder T. Drosophila Fecal Sampling. BIO-Protoc.
2017;7(18).

48.   Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation
of general 16S ribosomal RNA gene PCR primers for classical and next-generation
sequencing-based diversity studies. Nucleic Acids Res. 2013 Jan 1;41(1):e1.

49.   Toju H, Tanabe AS, Yamamoto S, Sato H. High-Coverage ITS Primers for the DNA-
Based Identification of Ascomycetes and Basidiomycetes in Environmental Samples.
PLOS ONE. 2012 July 12;7(7):e40863.

50.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
Reproducible, interactive, scalable and extensible microbiome data science using
QIIME 2. Nat Biotechnol. 2019 Aug;37(8):852–7.

51.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
High-resolution sample inference from Illumina amplicon data. Nat Methods. 2016
July;13(7):581–3.

52.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al.
Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME
2's q2-feature-classifier plugin. Microbiome. 2018 May 17;6(1):90.

53.   Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, et al. The SILVA and
"All-species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Res.
2014 Jan 1;42(D1):D643–8.

54.   Martin M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.journal. 2011 May 2;17(1):10–2.

55.   Bjørnsgaard Aas A, Davey ML, Kauserud H. ITS all right mama: investigating the
formation of chimeric sequences in the ITS2 region by DNA metabarcoding analyses
of fungal mock communities of different complexities. Mol Ecol Resour.
2017;17(4):730–41.

56.   Abarenkov K, Nilsson RH, Larsson KH, Taylor AFS, May TW, Frøslev TG, et al. The
UNITE database for molecular identification and taxonomic communication of fungi
and other eukaryotes: sequences, taxa and classifications reconsidered. Nucleic
Acids Res. 2023 Nov 11;1–7.

57.   QIIME 2 Development Team. Training feature classifiers with q2-feature-classifier
[Internet]. 2022. Available from: https://docs.qiime2.org/2022.8/tutorials/training-
feature-classifiers/

58.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLOS ONE. 2013 Apr
22;8(4):e61217.

59.   Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical
identification and removal of contaminant sequences in marker-gene and
metagenomics data. Microbiome. 2018 Dec 17;6(1):226.

60.   Faith DP, Minchin PR, Belbin L. Compositional dissimilarity as a robust measure
of ecological distance. Vegetatio. 1987 Apr 1;69(1):57–68.

61.   Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, et al.
vegan: Community Ecology Package. 2025.

62.   Larsen T, Pollierer MM, Holmstrup M, D'Annibale A, Maraldo K, Andersen N, et al.
Substantial nutritional contribution of bacterial amino acids to earthworms and
enchytraeids: A case study from organic grasslands. Soil Biol Biochem. 2016 Aug
1;99:21–7.

63.   O'Brien DM, Fogel ML, Boggs CL. Renewable and nonrenewable resources:
Amino acid turnover and allocation to reproduction in Lepidoptera. Proc Natl Acad
Sci. 2002 Apr 2;99(7):4413–8.

64.   Pollierer MM, Scheu S, Tiunov AV. Isotope analyses of amino acids in fungi and
fungal feeding Diptera larvae allow differentiating ectomycorrhizal and saprotrophic
fungi-based food chains. Briones M, editor. Funct Ecol. 2020 Nov;34(11):2375–88.

65.   Stock BC, Jackson AL, Ward EJ, Parnell AC, Phillips DL, Semmens BX. Analyzing
mixing systems using a new generation of Bayesian tracer mixing models. PeerJ. 2018
June 21;6:e5096.

66.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear
Mixed Effects Models. J Stat Softw. 2017 Dec 6;82:1–26.

67.   Knowles JE, Frederick C. merTools: Tools for Analyzing Mixed Effect Regression
Models. 2015. p. 0.6.2.

68.   Wickham H. A Layered Grammar of Graphics. J Comput Graph Stat. 2010 Jan
1;19(1):3–28.

69.   Keebaugh ES, Yamada R, Ja WW. The Nutritional Environment Influences the
Impact of Microbes on Drosophila melanogaster Life Span. Garsin DA, editor. mBio.
2019 Aug 27;10(4):e00885-19.

70.   Ludington WB, Ja WW. Drosophila as a model for the gut microbiome. PLOS
Pathog. 2020 Apr 23;16(4):e1008398.

71.   Adair KL, Wilson M, Bost A, Douglas AE. Microbial community assembly in wild
populations of the fruit fly Drosophila melanogaster. ISME J. 2018 Apr;12(4):959–72.

72.   Wang Y, Staubach F. Individual variation of natural D.melanogaster-associated
bacterial communities. FEMS Microbiol Lett. 2018 Mar 1;365(6):fny017.

73.   Ramírez-Camejo LA, Maldonado-Morales G, Bayman P. Differential Microbial
Diversity in Drosophila melanogaster: Are Fruit Flies Potential Vectors of
Opportunistic Pathogens? Int J Microbiol. 2017;2017(1):8526385.

74.   Corby-Harris V, Pontaroli AC, Shimkets LJ, Bennetzen JL, Habel KE, Promislow
DEL. Geographical Distribution and Diversity of Bacteria Associated with Natural
Populations of Drosophila melanogaster. Appl Environ Microbiol. 2007
June;73(11):3470–9.

75.   Dodge R, Jones EW, Zhu H, Obadia B, Martinez DJ, Wang C, et al. A symbiotic
physical niche in Drosophila melanogaster regulates stable association of a multi-
species gut microbiota. Nat Commun. 2023 Mar 21;14(1):1557.

76.   Baud GLC, Prasad A, Ellegaard KM, Engel P. Turnover of strain-level diversity
modulates functional traits in the honeybee gut microbiome between nurses and
foragers. Genome Biol. 2023 Dec 8;24(1):283.

77.   Jones R, Fountain MT, Andreani NA, Günther CS, Goddard MR. The relative
abundances of yeasts attractive to Drosophila suzukii differ between fruit types and
are greatest on raspberries. Sci Rep. 2022 June 20;12(1):10382.

78.   Pimenta RS, Alves PDD, Corrêa A, Lachance MA, Prasad GS, Rajaram, et al.
Geotrichum silvicola sp. nov., a novel asexual arthroconidial yeast species related to
the genus Galactomyces. Int J Syst Evol Microbiol. 2005;55(1):497–501.

79.   Yaghmour MA, Bostock RM, Morgan DP, Michailides TJ. Biology and Sources of
Inoculum of Geotrichum candidum Causing Sour Rot of Peach and Nectarine Fruit in
California. Plant Dis. 2012 Feb;96(2):204–10.

80.   Morais PB, Hagler AN, Rosa CA, Mendonca-Hagler LC, Klaczko LB. Yeasts
associated with Drosophila in tropical forests of Rio de Janeiro, Brazil. Can J
Microbiol. 1992 Nov;38(11):1150–5.

81.   Perkins V, Vignola S, Lessard MH, Plante PL, Corbeil J, Dugat-Bony E, et al.
Phenotypic and Genetic Characterization of the Cheese Ripening Yeast Geotrichum
candidum. Front Microbiol. 2020 May 7;11.

82.   Gurung K, Vink SN, Salles JF, Wertheim B. More persistent bacterial than fungal
associations in the microbiota of a pest insect. J Pest Sci. 2023 Mar 1;96(2):785–96.

83.   Consuegra J, Grenier T, Akherraz H, Rahioui I, Gervais H, da Silva P, et al.
Metabolic Cooperation among Commensal Bacteria Supports Drosophila Juvenile
Growth under Nutritional Stress. iScience. 2020 June;23(6):101232.

84.   Corby-Harris V, Promislow DEL. Host ecology shapes geographical variation for
resistance to bacterial infection in Drosophila melanogaster. J Anim Ecol. 2008 July
1;77(4):768–76.

85.   Gould AL, Zhang V, Lamberti L, Jones EW, Obadia B, Korasidis N, et al.
Microbiome interactions shape host fitness. Proc Natl Acad Sci. 2018 Dec
18;115(51).

86.   Chakraborty A, Mori B, Rehermann G, Hernández Garcia A, Lemmen-Lechelt J,
Hagman A, et al. Yeast and fruit fly mutual niche construction and antagonism
against mould. Funct Ecol. 2022;36(7):1639–54.

87.   Jose PA, Yuval B, Jurkevitch E. Maternal and host effects mediate the adaptive
expansion and contraction of the microbiome during ontogeny in a holometabolous,
polyphagous insect. Funct Ecol. 2023 Apr 1;37(4):929–46.

88.   Broderick NA. Friend, foe or food? Recognition and the role of antimicrobial
peptides in gut immunity and Drosophila–microbe interactions. Philos Trans R Soc B
Biol Sci. 2016 May 26;371(1695):20150295.

89.   Inamine H, Ellner SP, Newell PD, Luo Y, Buchon N, Douglas AE. Spatiotemporally
Heterogeneous Population Dynamics of Gut Bacteria Inferred from Fecal Time Series
Data. mBio. 2018 Jan 9;9(1):10.1128/mbio.01453-17.

90.   Trienens M, Rohlfs M. A Potential Collective Defense of Drosophila Larvae
Against the Invasion of a Harmful Fungus. Front Ecol Evol. 2020 May 14;8:1–10.

91.   Bronstein JL, Sridhar H. Connecting and integrating cooperation within and
between species. Philos Trans R Soc B Biol Sci. 2024 July 22;379(1909):20230203.

92.   Moriyama M, Fukatsu T. Host's demand for essential amino acids is
compensated by an extracellular bacterial symbiont in a hemipteran insect model.
Front Physiol. 2022 Sept 30;13:1028409.

93. Keebaugh ES, Yamada R, Obadia B, Ludington WB, Ja WW. Microbial Quantity
Impacts Drosophila Nutrition, Development, and Lifespan. iScience. 2018 June
29;4:247–59.

94. Anagnostou C, Dorsch M, Rohlfs M. Influence of dietary yeasts on Drosophila
melanogaster life-history traits: Influence of yeast on Drosophila melanogaster.
Entomol Exp Appl. 2010 May 17;136(1):1–11.

95. Piper MD. Using artificial diets to understand the nutritional physiology of
Drosophila melanogaster. Glob Change Biol Mol Physiol. 2017 Oct 1;23:104–11.

96. Trienens M, Keller NP, Rohlfs M. Fruit, flies and filamentous fungi – experimental
analysis of animal–microbe competition using Drosophila melanogaster and
Aspergillus mould as a model system. Oikos. 2010 Nov 1;119(11):1765–75.

97. Caballero Ortiz S, Trienens M, Rohlfs M. Induced Fungal Resistance to Insect
Grazing: Reciprocal Fitness Consequences and Fungal Gene Expression in the
Drosophila-Aspergillus Model System. PLOS ONE. 2013 Aug 30;8(8):e74951.

98. Lange C, Boyer S, Bezemer TM, Lefort MC, Dhami MK, Biggs E, et al. Impact of
intraspecific variation in insect microbiomes on host phenotype and evolution. ISME
J. 2023 Nov;17(11):1798–807.

99. Koerte S, Keesey IW, Easson MLAE, Gershenzon J, Hansson BS, Knaden M.
Variable dependency on associated yeast communities influences host range in
Drosophila species. Oikos. 2020 July;129(7):964–82.

100. Carramaschi I, Lery LMS, Chagas TPG, Zahner V. Klebsiella spp. carried by
insects as a reservoir of virulence and resistance to antimicrobials. One Health. 2025
June 1;20:101064.

101. Zhang S, Song F, Wang J, Li X, Zhang Y, Zhou W, et al. Gut microbiota facilitate
adaptation of invasive moths to new host plants. ISME J. 2024 Jan 1;18(1):wrae031.

102. Lu M, Hulcr J, Sun J. The Role of Symbiotic Microbes in Insect Invasions. Annu Rev
Ecol Evol Syst. 2016 Nov 1;47(Volume 47, 2016):487–505.

1175