PyMSQ: a python package for fast Mendelian sampling (co)variance and haplotype-based similarity in genomic selection

Abdulraheem Arome Musa and Norbert Reinsch

Research Institute for Farm Animal Biology (FBN), Wilhelm-Stahl-Allee 2, 18196 Dummerstorf, Germany.

Corresponding author: reinsch@fbn-dummerstorf.de

Abstract

Background: While genomic selection (GS) boosts rapid genetic gains by leveraging dense marker data for genomic estimated breeding values (GEBVs), prolonged application can reduce haplotype diversity and increase inbreeding. To address these risks, recent research has emphasized Mendelian sampling variance (MSV) and covariance (MSC), which capture within-family (co)variation not fully reflected in GEBVs. In parallel, theoretical advances have introduced a haplotype-based similarity measure that targets shared heterozygous segments, enabling more direct control over haplotype diversity—either standalone or in combination with conventional coancestry-based and genomic relationship matrices.

Results: We present PyMSQ, an open-source Python package that implements two key developments: (1) a matrix-based approach for computing MSV and MSC in single-trait, multi-trait, and zygotic contexts, and (2) a haplotype-based similarity metric. By combining this matrix-based framework with optimized scientific libraries, PyMSQ achieves up to 332-fold faster computations than gamevar—a publicly available alternative—while preserving numerical accuracy. Using a Holstein-Friesian dataset, we demonstrate PyMSQ's effectiveness in deriving MSV and MSC, as well as its novel similarity measure, which complements standard genomic relationship matrices by explicitly quantifying shared heterozygous segments rather than overall allele sharing, thereby providing additional insights for balancing immediate gains with long-term diversity.

Conclusion: By facilitating the practical use of MSV, MSC, and a haplotype-based similarity metric, PyMSQ enables breeders and quantitative geneticists to adopt haplotype diversity constraints—whether as a standalone criterion or in synergy with optimal contribution selection. This framework opens new possibilities for preserving key haplotypic segments, ultimately supporting more sustainable genomic selection strategies. PyMSQ is freely available under an MIT License at https://github.com/aromemusa/PyMSQ.

Keywords: Mendelian sampling variance, gametic variance, haplotype similarity matrices, optimal mating decisions, genetic diversity management.

Background

Genomic selection (GS) relies on genomic prediction, which uses dense marker data and predictive models to estimate breeding values more accurately than traditional pedigree-based methods [1, 2]. Although genomic estimated breeding values (GEBVs) have expedited short-term genetic progress, relying solely on GEBVs—without integrating additional criteria such as inbreeding—may exacerbate inbreeding and erode haplotype diversity over time [3–6]. Preserving genetic variation at the haplotype level is especially important for traits influenced by multiple linked loci, where the loss of crucial segments could irreversibly limit future breeding opportunities.

A key step toward more sustainable breeding decisions involves quantifying within-family genetic variation by way of Mendelian sampling variance (MSV) and covariance (MSC). These metrics require phased genotypes, a marker map, and estimated marker effects. Unlike GEBVs—an individual's average genetic merit—MSV and MSC capture the range of possible gametic outcomes a parent (or mating pair) could produce [7–15]. Identifying parents whose gametes exhibit high variability for one or multiple traits can help breeders "hedge" against narrowing genetic diversity [14, 16–21]. MSV therefore fulfils two complementary roles. First, it supports short-term gain: parents with larger MSV increase the chance that at least a few offspring will outrank their own GEBV, thereby raising the expected within-family response (the rationale behind strategies such as "usefulness-criterion" selection) [12, 13]. Second, it provides the variance component exploited by optimization frameworks—haplotype-similarity constraints[14, 20], coancestry-based optimal contributions [14], genomic-mating [22] and optimal cross-selection schemes [18]—to balance short-term gain with the long-term preservation of diverse haplotypes. However, computing MSV and MSC at scale has historically been computationally challenging, especially for large populations with high-density marker panels. Early simulation-based methods that enumerate or randomly sample parental haplotypes [7–10] become impractical for tens or hundreds of thousands of markers.

Subsequent analytical methods express breeding values in terms of marker effects [11, 23, 24], yet many still require constructing a parent-specific (co)variance matrix with phase indicators for heterozygous loci—an intensive step in large populations with dense marker sets. In contrast, Musa and Reinsch [14] recently proposed an efficient matrix-based representation that sets up parent-specific marker effects (including phase indicators for heterozygous loci) while building only a single population covariance matrix for each chromosome. This approach avoids creating a

unique covariance matrix per parent, thereby enabling rapid computation of MSV and MSC under

single-trait, multi-trait, and zygotic models. Beyond speed, the same framework yields a

haplotype-based similarity measure that complements coancestry or genomic relationship matrices

(GRMs) by targeting shared heterozygous segments (see "Haplotype-Based Similarity" below).

This measure quantifies the extent to which MSVs of potential parents arise from identical

chromosomal segments. By optimizing mate selection to minimize parental similarities, breeders

can preserve broader allelic diversity while maintaining expected genetic gain [14, 20].

Despite this theoretical progress, accessible software implementing both the matrix-based

MSV/MSC framework and haplotype-based similarity has been limited. Many breeding programs

still rely on simulation-heavy or inefficient analytical methods (e.g., gamevar [24]) that do not

provide haplotype-level diversity metrics, or that repeat computationally demanding steps for each

parent. To bridge this gap, we developed PyMSQ—an open-source Python package that merges

the efficient MSV/MSC derivations of Musa and Reinsch [14] with haplotype-based similarity

into a single platform. By leveraging optimized scientific libraries, PyMSQ can handle large

datasets, multi-trait analyses, and zygotic variance calculations more feasibly than older methods.

In what follows, we detail PyMSQ's design and implementation, validate its performance

on a Holstein-Friesian dataset, and demonstrate how breeders can integrate MSV, MSC, and

haplotype-based similarity to maintain diversity while achieving robust genetic gains in modern

genomic selection programs.

Implementation

Overview

PyMSQ is a Python 3.8+ package designed to compute MSV and MSC, alongside a haplotype-

based similarity metric, using the matrix-based framework presented by Musa and Reinsch [14].

By uniting parent-specific phase indicators with a single population covariance matrix $\boldsymbol{R}^c$ per

chromosome, PyMSQ eliminates the need for parent-specific covariance matrices—thus

significantly shortening runtime. Built primarily with NumPy and pandas for data manipulation, it

also uses Numba to just-in-time (JIT) compile numeric routines, enabling faster processing of large

genomic datasets.

Although written in Python, PyMSQ is compatible with R via the reticulate package,

ensuring straightforward integration with R-centric breeding pipelines. The following subsections

describe PyMSQ's internal data structures, matrix computations, and core algorithms.

Comprehensive package documentation, including examples and user guides, is available at

https://github.com/aromemusa/PyMSQ.

Data inputs and preparation

All genotype data must be fully phased and free of missing calls; PyMSQ neither performs phasing

nor imputation. Users should provide genotype arrays (or DataFrames) in which each marker is

assigned one of the two parental haplotypes. A genetic map is also required, specifying a

chromosome identifier (e.g., 1, 2) for each marker, along with either a genetic distance (d_{kl}^c) or

recombination rate (r_{kl}^c) between markers k and l on a chromosome c. These values populate the

population-level covariance matrix $\boldsymbol{R}^c$ for each chromosome c, reflecting linkage disequilibrium

patterns [14]. Markers on different chromosomes are considered independent.

Finally, marker-effect estimates must be supplied (commonly from a genomic prediction

model like genomic best linear unbiased prediction), ensuring their order and dimension exactly

match the SNP order of the phased genotype data. In a single-trait setting, PyMSQ expects a vector

$\boldsymbol{m}^c \in \mathbb{R}^K$ for each chromosome c. For multiple traits, it requires a matrix $\boldsymbol{M}^c \in \mathbb{R}^{T \times K}$, where T

is the number of traits. PyMSQ terminates if it detects inconsistencies in marker order or shape

among these datasets, prompting users to correct input files.

Construction of the population covariance matrix

Musa and Reinsch [14] emphasize constructing a single population covariance matrix $\boldsymbol{R}^c \in \mathbb{R}^{K \times K}$

per chromosome c, thereby avoiding parent-specific covariance overhead. The off-diagonal

entries ρ_{kl}^c encode the expected recombination-adjusted linkage between markers k and l.

PyMSQ supports two mapping functions:

1. Haldane's function (assuming no interference) [11, 25]:

128
$$\rho_{kl}^c = \frac{\exp(-2d_{kl}^c)}{4} \text{ or } \frac{(1-2r_{kl}^c)}{4}.$$

129   2. Linear approximation (with $\rho_{kl}^c = 0$ if $r_{kl}^c \geq 0.5$ (or $d_{kl}^c \geq 0.5$ Morgans) [23, 24]:

130
$$\rho_{kl}^c = \left(-\frac{d_{kl}^c}{2} + 0.25 \right) \text{ or equivalently } \left(-\frac{r_{kl}^c}{2} + 0.25 \right).$$

In both models, the diagonal elements of $\boldsymbol{R}^c$ are 0.25 [11, 14] , reflecting the variance contribution

of a fully heterozygous locus. Internally, PyMSQ stores each $\boldsymbol{R}^c$ in a dictionary or list for quick

access.

Parent-specific marker effects

Where earlier methods often built a phase-adjusted matrix $\boldsymbol{R}_i^c$ for each parent i [11, 23, 24],

PyMSQ encodes phase indicators directly into marker-effect vectors (or matrices), following Musa

and Reinsch [14]. Let $\boldsymbol{m}^c \in \mathbb{R}^K$ be the additive marker-effect vector for chromosome c . For parent

i , PyMSQ constructs a vector $\boldsymbol{m}_i^c$ by multiplying each entry in $\boldsymbol{m}^c$ by a phase indicator δ_{ik} :

$m_{ik}^c = \delta_{ik} m_k^c$, with $\delta_{ik} = \begin{cases} +1 & \text{if parent } i \text{ has the reference allele on the first haplotype at marker } k \\ -1 & \text{if parent } i \text{ has that allele on the second haplotype at marker } k \\ 0 & \text{if parent } i \text{ is homozygous at marker } k. \end{cases}$

This marker-effect adjustment eliminates the need for parent-specific covariance matrices and

smoothly extends to multi-trait contexts by applying δ_{ik} to each rows of $\boldsymbol{M}^c \in \mathbb{R}^{T \times K}$.

Computing MSV and MSC for gametes

Once $\boldsymbol{R}^c$ and $\boldsymbol{m}_i^c$ are established, the MSV of parent i (single trait) becomes:

$$\sigma_{b_i}^2 = \sum_c (\boldsymbol{m}_i^{\,c})' \boldsymbol{R}^c \boldsymbol{m}_i^{\,c} , \qquad (1)$$

while in multi-trait settings, each chromosome's marker-effect matrix $\boldsymbol{M}_i^c$ yields:

$$\boldsymbol{V}_i = \sum_c \boldsymbol{M}_i^c \boldsymbol{R}^c \left(\boldsymbol{M}_i^c \right)' , \qquad (2)$$

where $\boldsymbol{V}_i$ contains MSVs on the diagonal and MSCs off-diagonal [14]. An aggregate genotype is

computed via $\boldsymbol{a}'\boldsymbol{V}_i\boldsymbol{a}$, given a vector of index weight $\boldsymbol{a} \in \mathbb{R}^{T \times 1}$.

Computing MSV and MSC for zygotes (parent-pair)

When considering a parent pair ij , PyMSQ sums each parent's gametic variance to derive the

(co)variance of their potential offspring [11]:

155
$$var\left(\boldsymbol{b}_{ij}\right) = \sum_{c}\left[\boldsymbol{M}_{i}^{c}\boldsymbol{R}^{c}\left(\boldsymbol{M}_{i}^{c}\right)' + \boldsymbol{M}_{j}^{c}\boldsymbol{R}^{c}\left(\boldsymbol{M}_{j}^{c}\right)'\right] = \boldsymbol{V}_{ij}.$$
(3)

In single-trait mode, this reduces to $\sigma_{b_{ij}}^{2}$. This enables direct comparisons of within-family

(co)variation for multiple mating pairs under single- or multi-trait models. The aggregated

genotype variance for pair ij is $\boldsymbol{a}'\boldsymbol{V}_{ij}\boldsymbol{a}$, and PyMSQ outputs each trait's MSV, the aggregate MSV,

and any inter-trait covariances.

Haplotype-Based Similarity

PyMSQ implements the haplotype-based similarity measure from Musa and Reinsch [14],

explicitly targeting shared heterozygous segments among parents. Specifically, the similarity

measure quantifies the expected covariance between the additive genetic values of gametes

(Mendelian sampling terms) produced by pairs of parents. In single-trait scenarios, the similarity

s_{ij} between parents i and j is:

167
$$s_{i,j} = \sum_{c}\left|\left(\boldsymbol{m}_{i}^{c}\right)'\boldsymbol{R}^{c}\boldsymbol{m}_{j}^{c}\right|,$$
(4)

where the absolute value maintains invariance to haplotype order. A similarity matrix $\boldsymbol{S} \in \mathbb{R}^{N \times N}$

for N parents can be set up such that a parent's self-similarity equals its MSV, i.e., $s_{i,i} = \sigma_{b_{i}}^{2}$.

Marker-effect values in $\boldsymbol{m}_{i}^{c}$ and $\boldsymbol{m}_{j}^{c}$ are set to zero for loci homozygous in both parents, as these

loci do not generate MSV and hence do not contribute to similarity. Consequently, high similarity

values identify parent pairs whose MSVs originate from the same chromosomal segments, whereas

low similarity values indicate parents with MSVs arising from distinct chromosomal regions.

Breeding schemes can thus optimize mate selection by minimizing average similarity s_{ij},

effectively preserving allelic diversity and moderating long-term inbreeding without

compromising immediate genetic gain.

For multi-trait contexts, a user-specified index-weight vector $\boldsymbol{a}$ yields:

178
$$s_{ij} = \sum_{c}\left|\boldsymbol{a}' \cdot \left(\boldsymbol{M}_{i}^{c}\boldsymbol{R}^{c}\left(\boldsymbol{M}_{j}^{c}\right)'\right) \cdot \boldsymbol{a}\right|.$$

179 PyMSQ also calculates similarities $s_{ij,uv}$ between zygotes from parent pairs ij and uv:

180
$$s_{ij,uv} = \sum_c \left[\left| \boldsymbol{a}' \cdot \left(\boldsymbol{M}_i^c \boldsymbol{R}^c \left(\boldsymbol{M}_u^c \right)' \right) \cdot \boldsymbol{a} \right| + \left| \boldsymbol{a}' \cdot \boldsymbol{M}_j^c \boldsymbol{R}^c \left(\boldsymbol{M}_v^c \right)' \cdot \boldsymbol{a} \right| \right], \tag{5}$$

181 placing the respective MSVs on the diagonal [14]. Here, the row vectors in $\boldsymbol{M}_u^c$ and $\boldsymbol{M}_v^c$

182 correspond to the trait-specific marker effects of parents u and v for chromosome c.

184 ***Standardized similarity***

185 Because similarity values can range widely, PyMSQ offers a standardized matrix $\boldsymbol{K}$ derived as:

186
$$\boldsymbol{K} = \boldsymbol{D}^{-1} \cdot \boldsymbol{S} \cdot \boldsymbol{D}^{-1}, \tag{6}$$

where $\boldsymbol{D}$ is diagonal, holding each parent's Mendelian standard deviation $\sqrt{\sigma_{b_i}^2}$. This transforms

off-diagonal entries to $[0,1]$ and can help breeders impose constraints on haplotype similarity [14,

20].

Selection criteria

PyMSQ calculates GEBVs for parent i as $b_i = \boldsymbol{c}_i' \boldsymbol{m}$, with $\boldsymbol{c}_i^c \in \mathbb{R}^K$ denoting the marker genotype

and $\boldsymbol{m} \in \mathbb{R}^K$ the marker-effect vector derived from genomic prediction models [1]. It then derives

an index $I_i = b_i + \lambda \sigma_{b_i}$, combining GEBV and MSV, where σ_{b_i} represents the Mendelian standard

deviation. The constant λ may reflect selection intensity [13] or optimize the probability of

producing top-ranking offspring (e.g., $\sqrt{2}x$, where x is the standardized normal truncation point

of the selected proportion [12]). Under a zygotic approach, GEBVs and indices for a parent pair

ij are averaged to explore the joint potential.

Algorithmic and software optimization

Most core operations in PyMSQ—such as setting up $\boldsymbol{R}^c$, $\boldsymbol{m}_i^c$ or $\boldsymbol{M}_i^c$—are vectorized or just-in-

time (JIT) compiled using Numba, reducing Python's overhead in loops and arithmetic on large

arrays. For instance, repetitive tasks like $(\boldsymbol{m}_i^c)' \boldsymbol{R}^c \boldsymbol{m}_i^c$ benefit significantly from JIT speedups. The

software caches each parent's phase-adjusted marker effects to avoid recomputing $\boldsymbol{m}_i^c$ or $\boldsymbol{M}_i^c$

multiple times. It also parallelizes matrix multiplications, leveraging NumPy's underlying

BLAS/LAPACK routines.

Outputs and integration

PyMSQ returns results primarily as NumPy arrays or pandas DataFrames, facilitating export to

.csv or .npy. Because many breeding analyses are R-based, PyMSQ is fully callable from R via

reticulate, supporting seamless integration with other R-based genomic pipelines (e.g., rrBLUP,

BGLR). By accommodating both Python and R environments, PyMSQ aims to reach a broad

community of breeders and geneticists.

We envision that MSV, MSC, and haplotype-based similarity matrices will typically be

computed centrally at evaluation centers or large-scale breeding units where phased genotypes and

genomic predictions are available. These pre-computed values can then be securely and efficiently

distributed (e.g., via routine evaluation reports or secure APIs) to breeding advisors or farms. Thus,

direct access to raw genotype data at the farm level is unnecessary. Furthermore, PyMSQ readily

accommodates complex multi-trait indices involving numerous traits or sub-indexes, since multi-

trait indices are easily represented by single weighted marker-effect vectors, ensuring

computational efficiency.

Data and analysis

Using PyMSQ and a publicly available Holstein-Friesian cattle dataset [14], we derived MSCs and

haplotype similarity matrices for the aggregate genotype of gametes, assigning equal index weights

of 1. The dataset includes 265 cows from five paternal half-sib families (sizes 32–106), along with

a physical map, genotypic data, pedigree information, and phenotypes for three milk traits—pH,

fat yield (FY), and protein yield (PY) [26, 27]. Marker effects for this population had been

estimated previously [26], and the genotypes were phased into haplotypes (yielding 10,304

markers). Approximate genetic map positions were also established [14]. These data files (map,

phased genotypes, marker effects, and pedigree details) are bundled with PyMSQ to ensure full

reproducibility.

Following the computation of MSVs and MSCs in PyMSQ, we visualized the results with

ggplot2 (v 3.3.3) [28], generating density plots of MSVs and MSCs. We then constructed and

examined haplotype similarity matrices using corrplot. We then generated and examined haplotype

similarity matrices with corrplot (v 0.84) [29] to assess interrelationships among parental

genotypes. Supporting scripts and their outputs are included in Additional File 1.

To compare PyMSQ's performance with gamevar [24], we ran simulation studies with two
data groups, each focused on a single chromosome and ten traits. In the Individuals scenario, we
fixed the number of markers and varied the number of individuals to test computational scalability
with increasing population size. We generated 20 datasets each containing 1,000 markers but
varied the number of individuals. In the Markers scenario, we fixed the number of individuals at
500 while varying marker numbers. Each dataset underwent ten replicates, measuring the
computation time (minutes) and peak memory usage (GB) needed to compute MSCs, as well as
the time and memory required for haplotype similarity matrices. All computations were conducted
on a multi-user Linux server (Intel Xeon Gold 6130, 128 CPU cores at 2.10 GHz, 1536 GB
RAM)—a high-memory environment typical of large-scale genomic breeding analyses. Detailed
scripts and outputs are provided in Additional File 1.
249
Results
Mendelian variances and trait correlations
Applying PyMSQ to the Holstein-Friesian dataset revealed a wide range of MSVs for the three
milk traits: pH, FY, and PY. As illustrated in Figure 1A, MSV distributions for each trait showed
coefficients of variation exceeding 90%, underscoring the substantial within-family genetic
diversity stemming from heterozygous loci and linkage phase [11, 14]. The FY trait in particular
exhibited a bimodal distribution—possibly indicating DGAT1 as a major locus—previously noted
to affect milk-fat segregation [8, 11, 23, 30]. Moving beyond single-trait analyses, we derived
multi-trait MSCs for each cow, thus quantifying how alleles for pH, FY, and PY co-segregate. The
resulting correlations (Figure 1B) revealed that while most cows had moderately positive inter-
trait correlations, a subset displayed mildly negative correlations—a reflection of individual
differences in linkage phases or pleiotropic effects [11, 23, 31]. These findings highlight
opportunities for breeders to strategically select or mate parents to enhance beneficial multi-trait
associations or reduce antagonistic linkages.
264
Haplotype-based similarity and its standardization
We next constructed a haplotype-based similarity matrix for an aggregate genotype combining pH,
FY, and PY with equal weights. Diagonal elements in this matrix correspond to each cow's MSV,
and off-diagonals measure the overlap in high-impact heterozygous segments [14]. Figure 2A

illustrates that some pairs of cows share extensive haplotype regions, while others remain
relatively dissimilar. We also generated a standardized matrix K (Figure 2B) by dividing off-
diagonal entries by the product of each parent's Mendelian standard deviation, which rescales
similarities to a [0,1] interval and accentuates smaller overlaps. This haplotype-level focus
complements the GRM by highlighting heterozygous loci relevant to within-family variance,
rather than just overall allele sharing [14, 20]. In practical breeding settings, imposing limits on
pairwise similarity can retain critical haplotypes, thus promoting the transmission of diverse
haplotypes, thereby moderating rather than accelerating long-term inbreeding.
277
Comparative benchmarking with gamevar
In the Individuals scenario, the number of markers was fixed at 1,000 while individual counts
ranged from 5,000 to 100,000. PyMSQ's runtime (Figure 3A) increased from about 0.07 min to
1.16 min, whereas gamevar required 1.69 min to 30.80 min—yielding a time ratio
(gamevar/PyMSQ) of 23 to 27. Memory usage (Figure 3C) showed the opposite trend: PyMSQ
scaled from ~0.55 GB to ~4.97 GB, while gamevar remained near 0.015 GB throughout, reflecting
gamevar's genotype line-streaming approach versus PyMSQ's in-memory design.
In the Markers scenario (Figure 3B, D), 500 individuals were held constant while marker
counts rose from 2,500 to 50,000. PyMSQ's runtime climbed from 0.03 min to 3.25 min, whereas
gamevar soared from 2.44 min to ~1,078.89 min, producing time ratios of up to 332. Gamevar's
memory usage grew from ~0.07 GB to ~27.95 GB, reflecting the quadratic expansion of marker-
level arrays. Although PyMSQ's memory usage also increased (from ~1.10 GB to ~36.35 GB), its
matrix-based approach remained markedly faster, particularly for higher marker densities and
multi-trait analyses. Overall, PyMSQ provides substantial speed advantages over gamevar,
provided sufficient RAM is available.
293
Benchmarking of haplotype similarity matrix
We further evaluated PyMSQ's construction of the haplotype-based similarity matrix, an $O(N^2)$
procedure for N parents (Additional File 1, Table 3). In the Individuals scenario (population sizes
of 5,000 to 100,000, with 1,000 markers), PyMSQ's runtime for building this matrix rose from
about 0.08 min to 7.93 min, and peak memory from 0.76 GB to nearly 70 GB. Conversely, in the
Markers scenario (500 individuals, 2,500–50,000 markers), similarity matrix runtime grew from

0.04 min to 2.8 min, and memory usage climbed from 0.37 GB to ~34.28 GB. Although large

servers can handle these demands, breeding programs with constrained resources may choose to

subset parents (e.g., based on GEBVs) rather than compute a full $N \times N$ matrix. Nonetheless,

PyMSQ's ability to handle large-scale haplotype similarity computations—whether driven by

large N or substantial marker densities—enables refined selection and mating strategies that

directly target critical heterozygous segments.

Limitations and future directions

Though PyMSQ is both fast and versatile, its in-memory design can prove demanding for those

with restricted hardware. For MSV/MSC calculations, one can process chromosomes sequentially,

merging results to reduce memory usage. However, generating a full haplotype similarity matrix

for very large N may be infeasible unless breeders subset parents based on GEBVs or other

criteria. PyMSQ currently focuses on additive genetic models; adding dominance or epistasis

would broaden its scope, while further chunk-based or parallel-streaming optimizations could

shrink memory overhead. We emphasize that PyMSQ does not replace genomic prediction

methods but explicitly serves as a post-processing step applied directly to marker effects derived

from genomic prediction methods (Single-Step or Two-Step). PyMSQ explicitly calculates MSV,

MSC, and haplotype similarity based exclusively on genomic marker-effect estimates. It does not

directly incorporate the additional polygenic (residual additive genetic) component that is

explicitly modeled using pedigree or genomic relationships in single-step genomic evaluations.

Finally, our tests focused on one cattle population; applying PyMSQ to other species (swine,

poultry, crops) should help confirm its broader applicability for sustaining within-family genetic

diversity.

Conclusions

PyMSQ unifies advanced Mendelian (co)variance computation with haplotype-based similarity in

a single, open-source platform, enabling faster multi-trait analyses than existing methods. It affords

breeders a potent tool to harness within-family variance and safeguard genetic diversity—

potentially working in concert with coancestry-based or optimal contribution approaches. Future

PyMSQ releases will refine data handling, incorporate non-additive genetic models, and explore

multi-species applications, ensuring it remains a practical, high-performance resource for modern

genomic selection.

Availability and requirements

Project name: PyMSQ

Project home page: https://github.com/aromemusa/PyMSQ

Operating systems: Platform independent

Programming language: Python 3.8+

Other requirements: pandas, NumPy, SciPy, and Numba Python libraries

License: MIT license

Any restrictions to use by non-academics: None

All code, documentation, and the Holstein-Friesian dataset used in the examples are available

through the GitHub repository. Users may run PyMSQ directly in Python or call it from R via the

reticulate package. Contributions—including issue reports, feature requests, or new

functionalities—are welcomed via GitHub's issue tracker.

Figures

Fig. 1 Density plots of Mendelian sampling variances and trait correlations. Panel A displays the

variance in fat yield (kg), protein yield (kg), and pH (mol/L). Panel B shows correlations between

these traits.

Fig. 2 Unstandardized (A) and standardized (B) similarity matrices for the aggregate genotype of

265 Holstein-Friesian cows from five half-sib families (delineated by red lines). In (A), diagonal

values represent individual Mendelian sampling variances, and off-diagonal values represent

similarity between parent pairs in terms of shared gametic variance. Panel (B) shows the

standardized similarity matrix, where values range from 0 (no similarity) to 1 (maximum

similarity).

Fig. 3 Benchmark plots comparing computation time (Panels A, B) and memory usage (Panels C,

D) for PyMSQ and gamevar.

Supplementary information

Supplementary information accompanies this paper at

https://github.com/aromemusa/PyMSQ/blob/main/paper/Additional_file_1.pdf.

Additional file 1: Analysis of Holstein-Friesian cattle data and benchmark test

Abbreviations

FY – fat yield (kg); GEBV – genomic estimated breeding value; GRM – genomic relationship matrix; GS – genomic selection; JIT – just-in-time (compilation); MSC – Mendelian sampling covariance; MSV – Mendelian sampling variance; pH – acidity measure (hydrogen-ion concentration); PY – protein yield (kg); RAM – random access memory

Acknowledgments

We thank Nina Melzer and Dörte Wittenburg of FBN Dummerstorf for providing the empirical Holstein-Friesian dataset used in this study.

Author's contributions

AAM developed the Python package, analyzed the data, and wrote the manuscript. NR conceptualized and supervised the entire work and revised the manuscript. All of the authors read and approved the final manuscript.

Funding

Financial support for AAM from Bundesanstalt für Landwirtschaft und Ernährung (BLE) under Grant 281B101516 is gratefully acknowledged. The publication of this article was funded by the Open Access Fund of the Leibniz Association.

Availability of data and materials

The code, documentation, illustration, and Holstein-Friesian cattle data are available at https://github.com/aromemusa/PyMSQ.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Meuwissen THE, Hayes BJ, Goddard ME. Prediction of total genetic value using genome-wide dense marker maps. Genetics. 2001;157:1819–29.

2. Hickey JM, Chiurugwi T, Mackay I, Powell W. Genomic prediction unifies animal and plant breeding programs to form platforms for biological discovery. Nat Genet. 2017;49:1297–303.

3. Schaeffer LR. Strategy for applying genome-wide selection in dairy cattle. J Anim Breed Genet. 2006;123:218–23.

4. Liu H, Meuwissen TH, Sørensen AC, Berg P. Upweighting rare favourable alleles increases long-term genetic gain in genomic selection programs. Genet Sel Evol. 2015;47:19.

5. Lin Z, Cogan NOI, Pembleton LW, Spangenberg GC, Forster JW, Hayes BJ, et al. Genetic Gain and Inbreeding from Genomic Selection in a Simulated Commercial Breeding Program for Perennial Ryegrass. Plant Genome. 2016;9:1–12.

6. Rutkoski J, Singh RP, Huerta-Espino J, Bhavani S, Poland J, Jannink JL, et al. Genetic gain from phenotypic and genomic selection for quantitative resistance to stem rust of wheat. Plant Genome. 2015;8:1–10.

7. Mohammadi M, Tiede T, Smith KP. Popvar: A genome-wide procedure for predicting genetic variance and correlated response in biparental breeding populations. Crop Sci. 2015;55:2068–77.

8. Segelke D, Reinhardt F, Liu Z, Thaller G. Prediction of expected genetic variation within groups of offspring for innovative mating schemes. Genet Sel Evol. 2014;46:42.

9. Cole JB, VanRaden PM. Use of haplotypes to estimate Mendelian sampling effects and selection limits. J Anim Breed Genet. 2011;128:446–55.

10. Bernardo R. Genomewide selection of parental inbreds: Classes of loci and virtual biparental populations. Crop Sci. 2014;54:2586–95.

11. Bonk S, Reichelt M, Teuscher F, Segelke D, Reinsch N. Mendelian sampling covariability of marker effects and genetic values. Genet Sel Evol. 2016;48:1–11.

12. Bijma P, Wientjes YCJ, Calus MPL. Breeding Top Genotypes and Accelerating Response to Gametic Variance. Genetics. 2020;214:91–107.

13. Lehermeier C, Teyssèdre S, Schön CC. Genetic gain increases by applying the usefulness criterion with improved variance prediction in selection of crosses. Genetics. 2017;207:1651–61.

14. Musa AA, Reinsch N. A similarity matrix for hedging haplotype diversity among parents in genomic selection. J Anim Breed Genet. 2025. https://doi.org/https://doi.org/10.1111/jbg.12930.

15. Wellmann R, Bennewitz J. Key genetic parameters for population management. Front Genet. 2019;10 JUN:1–20.

16. Wolfe MD, Chan AW, Kulakow P, Rabbi I, Jannink JL. Genomic mating in outbred species: Predicting cross usefulness with additive and total genetic covariance matrices. Genetics. 2021;219.

17. Neyhart JL, Lorenz AJ, Smith KP. Multi-trait Improvement by Predicting Genetic Correlations in Breeding Crosses. G3 Genes, Genomes, Genet. 2019;9:3153–65.

18. Allier A, Lehermeier C, Charcosset A, Moreau L, Teyssèdre S. Improving short-and long-term genetic gain by accounting for within-family variance in optimal cross-selection. Front Genet. 2019;10 OCT:1–15.

19. Niehoff TAM, ten Napel J, Bijma P, Pook T, Wientjes YCJ, Hegedűs B, et al. Improving selection decisions with mating information by accounting for Mendelian sampling variances looking two generations ahead. Genet Sel Evol. 2024;56.

20. Musa AA, Reinsch N. Hedging genetic diversity using a similarity matrix based on Mendelian
sampling. In: Proceedings of 12th World Congress on Genetics Applied to Livestock Production
(WCGALP). 2022. p. 936–9.
21. Reinsch N, Musa AA. Neue Kriterien für die genomische Selektion. Züchtungskunde.
2021;93:201–9.
22. Akdemir D, Sánchez JI. Efficient breeding by genomic mating. Front Genet. 2016;7 NOV:1–
12.
23. Santos DJA, Cole JB, Lawlor TJ, VanRaden PM, Tonhati H, Ma L. Variance of gametic
diversity and its application in selection programs. J Dairy Sci. 2019;102:5279–94.
24. De Abreu Santos DJ, Cole JB, Liu GE, Vanraden PM, Ma L. Gamevar.f90: A software package
for calculating individual gametic diversity. BMC Bioinformatics. 2020;21:3–7.
25. Haldane JBS. The combination of linkage values and the calculation of distances between the
loci of linked factors. J Genet. 1919;8:299–309.
26. Melzer N, Wittenburg D, Repsilber D. Integrating Milk Metabolite Profile Information for the
Prediction of Traditional Milk Traits Based on SNP Information for Holstein Cows. PLoS One.
2013;8:e70256.
27. Hampel A, Teuscher F, Gomez-Raya L, Doschoris M, Wittenburg D. Estimation of
Recombination Rate and Maternal Linkage Disequilibrium in Half-Sibs. Front Genet. 2018;9
JUN:186.
28. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York; 2016.
29. Wei T, Viliam S. R package "corrplot": Visualization of a Correlation Matrix. R package
version 0.92; 2021. https://github.com/taiyun/corrplot. 2021.
30. Thaller G, Krämer W, Winter A, Kaupe B, Erhardt G, Fries R. Effects of DGAT1 variants on
milk production traits in German cattle breeds. J Anim Sci. 2003;81:1911–8.
31. Musa AA, Reinsch N. Unravelling the Potential of Mendelian Covariability for Multi-Trait
Genomic Selection. In: Proceedings of the 35th Annual Conference of the Society for Animal
Sciences (GfT) and the German Society for Animal Production (DGfZ) held on 13-14 September
in Halle, Germany. 2023.
479
480