Main Manuscript for Biophysics and Computational Biology

Structural basis for gating inhibition by the cytoplasmic domain in HCN1 channels

Valentina Zorzini[1*], Jessica M. Kleiz-Ferreira[1], Marijke Brams[1], Rodrigo Gallardo[2*], Steven De Gieter[3,4], Jana Kusch[5], Chris Ulens[1]

[1]Laboratory of Structural Neurobiology, Department of Cellular and Molecular Medicine, Faculty of Medicine, KU Leuven, 3000 Leuven, Belgium.

[2]School of Molecular and Cellular Biology, Faculty of Biological Sciences and Astbury Centre for Structural Molecular Biology, University of Leeds, Leeds, LS2 9JT, United Kingdom.

*Current affiliation: Switch Laboratory, Department of Cellular and Molecular Medicine, KU Leuven, 3000 Leuven, Belgium.

[3]Center for Structural Biology, Vlaams Instituut voor Biotechnologie, 1050 Brussels, Belgium.

[4]Structural Biology Brussels, Department of Bioengineering Sciences, Vrije Universiteit Brussel, 1050 Brussels, Belgium.

[5]Institut für Physiologie II, Universitätsklinikum Jena, Jena, Germany.

Email correspondence to: chris.ulens@kuleuven.be

Author Contributions: VZ performed protein expression, purification, sample and grid preparation, collected and processed Cryo-EM data. MB performed cloning and helped with purification. RG helped with grid preparation, collection and processing of cryo-EM data. VZ, JMKF, SDG, RGE and CU analyzed the data and prepared figures. CU designed and guided the project. VZ and CU wrote the manuscript with input from other authors. All authors have read the manuscript.

Competing Interest Statement: the authors declare no competing interests.

Classification: Biophysics and Computational Biology

Keywords: voltage-activated ion channel, cryo-electron microscopy, electrophysiology

This PDF file includes:

    Main Text
    Figures 1 to 5
    Table 1

Abstract

Hyperpolarization-activated, cyclic nucleotide-gated (HCN) channels generate rhythmic electrical activity in cardiac and neuronal tissues, with isoform-specific cAMP sensitivity remaining poorly understood. While HCN2 exhibits strong cAMP regulation, HCN1 shows minimal response. To investigate the structural basis of this divergence, we analyzed two engineered HCN1 variants using cryo-electron microscopy. One variant (HCN112) incorporates the C-linker and CNBD from HCN2 into the HCN1 backbone and exhibited enhanced cAMP sensitivity, with structural analysis revealing a compressed cytoplasmic domain arrangement that may facilitate regulatory interactions. In contrast, the truncated HCN1ΔC variant (lacking the cytoplasmic domain) displayed an intermediately open pore conformation, supporting auto-inhibitory regulation by the CNBD in wild-type channels. These structural insights elucidate how domain-specific interactions modulate cAMP-dependent gating and intrinsic auto-inhibition, resolving long-standing questions about mechanistic divergence among HCN isoforms. Our findings not only shed new light on the structural mechanisms underlying isoform-specific cAMP sensitivity but also have implications for the development of therapeutic strategies targeting HCN channels in neurological and cardiac disorders.

Significance Statement

Hyperpolarization-activated cyclic nucleotide-gated (HCN) channels are essential regulators of rhythmic electrical activity in the heart and brain, and their dysfunction is linked to disorders such as epilepsy, depression, and cardiac arrhythmias. Although HCN channel isoforms display highly divergent responses to cAMP modulation, the structural basis for these differences has remained elusive. This study reveals, through cryo-EM structures, how domain-specific interactions within the cytoplasmic regions of HCN1 channels underlie their unique gating and auto-inhibition properties, as well as their muted cAMP sensitivity compared to HCN2. These insights resolve long-standing mechanistic questions about HCN channel regulation and pave the way for rational design of targeted therapies that selectively modulate HCN isoform activity in neurological and cardiac disease.

Main Text

Introduction

Ion channels are integral membrane proteins that act as molecular switches of electrochemical signaling in the central nervous system. More specifically, the Hyperpolarization-activated Cyclic Nucleotide-sensitive (HCN) channels belong to a family of cation channels that is activated by hyperpolarizing potentials and regulated by cyclic AMP (cAMP). HCN channels serve as nonselective voltage-gated cation channels that contribute to spontaneous rhythmic activity in the plasma membranes of heart and brain cells. This type of channel is also concentrated in cortical and hippocampal pyramidal cell dendrites, where it plays an important role in determining synaptic input integration and thus neuronal output. Therefore, it has been suggested to be involved in physiological processes such as cognition as well as pathophysiological states such as epilepsy. Recent evidence suggests that these channels may also be therapeutic targets for treatment of depressive disorders (1).

Wang et al., 2001 have identified structural domains in HCN1 that are important for the basal voltage dependence of gating and the extent to which cAMP modulates the gating (2). Specifically, channels containing either the core transmembrane domain or C-terminus of HCN2 channels activate at more negative voltages compared with channels that contain the corresponding regions of HCN1. Differences in cAMP modulation are largely localized to the C-terminus, with channels containing the HCN2 C-terminus showing a larger shift in response to cAMP compared to channels containing the HCN1 C-terminus (2). Thus, removal of the entire C-terminus fully eliminates the

52 response to cAMP. Yet the precise nature and molecular mechanism of this structural modulation
53 is unknown.
54
Although the HCN1 structures in its apo-state and also cAMP-bound have already been solved by
cryo-EM (3), together with other HCN1 variants (4), none of these structures showed an open gate
and therefore they do not fully explain the HCN1 mechanism of gating. More recently, another
group reports a full panorama of structures on HCN4, in its apo-state and cAMP-bound, in the
presence of different detergents. The group also presents a HCN4 structure with the gate open
shedding light onto gating movement and ion permeation in HCN4 pacemaker channels (5).
Multiple other studies support the HCN role also in the function of sensory neurons and pain
sensation, particularly pain associated with nerve or tissue injury (6). Therefore, understanding the
structural basis of the HCN channel regulation could offer unique insights into the molecular basis
of its modulation and its druggability.
65
66 However, although lots of structures have already been solved, the regulation of HCN gating activity
67 is not fully understood. How the different domains change conformation, communicate between
68 each other's, and also inhibit gating remains unknown.
69
This study aims to structurally unveil how the HCN gating mechanisms are modulated by the C-
terminus in its apo-state and by cAMP ligand binding. To understand the role of the C-terminus in
the modulation of gating induced by cAMP, we have expressed and purified two HCN variants:
HCN112 and HCN1ΔC (2, 7). HCN112 is a chimera between the cytoplasmic NH2 terminus and
transmembrane domain of HCN1 and the cytoplasmic C-terminus of HCN2. On the other hand,
HCN1ΔC has a deletion corresponding to the full cytoplasmic C-terminus. These constructs allowed
us to understand a long-standing question whether the C-terminal domain acts in an auto-inhibitory
manner on the gate of the channel.
78
79 **Results and Discussion**
80
C-terminal region of HCN2 shows a compressed conformation in an HCN1 channel
background
In recent years, several research groups have published three-dimensional structures for three of
the four HCN isoforms, namely HCN1, HCN3 and HCN4 channels (3–5, 8, 9). For HCN2, however,
only the three-dimensional structure of the isolated CL-CNBD has been elucidated so far (10, 11).
To study the C-terminal region of the HCN2 channel not only as an isolated protein but in a full
channel environment, we took advantage of the HCN112 chimera (Figure 1A). This chimera is
composed of the N-terminal domain and transmembrane domain of HCN1 and the C-terminal
domain of HCN2 channels (2).
To show that the CL-CNBD region in the HCN112 chimera is still functional, we tested the effect of
cAMP on the current amplitude in inside-out macropatches of *Xenopus laevis* oocytes. Data
presented in Figure 1B (left panel) show that the ability of cAMP to increase the maximum current
amplitude, I_{max}, evoked by a hyperpolarizing voltage jump is still intact after binding to an HCN2
CNBD in an HCN1 background. From this data we conclude that the HCN112 chimera is a valid
tool to compare HCN1 and HCN2 CNBD structures in a functional channel system.
We expressed and purified HCN112 and HCN1ΔC according to Lee and MacKinnon (3). Cryo-EM
structures of HCN112 in the ligand-free and cAMP-bound state were determined to an overall
resolution of 3.5 Å and 3.26 Å (Figure 1C-D). This is comparable to the resolution of HCN1
structures (3.5 Å) by Lee and MacKinnon (3). The overall channel structure of HCN112 in the
ligand-free state is comparable to HCN1 ligand-free (pdb code 5u6o) with an RMSD of 1.4250 Å.
Similar to HCN1, HCN112 channels consist of four identical subunits forming a 4-fold symmetric
channel which is surrounded by the membrane. It has a membrane-embedded set of six alpha-
helices (S1-S6, transmembrane domain = TMD) constituting the pore (S5-S6) and the voltage
sensor (S1-S4). The HCN112 C-terminal cytoplasmic domain is formed by the C-linker and the
cyclic nucleotide binding domain (CNBD). Unique to HCN channels is the N-terminal HCN domain

spanning the first 45 residues preceding S1 and forming a 3-alpha-helical domain. This domain is
wedged between the voltage sensor and the cytoplasmic domain. The HCN domain contacts the
S4 helix from the same subunit and the C-linker and CNBD from an adjacent subunit. This
constrains the channel at the level of the C-linker disk (3).
Besides these above-mentioned similarities in the overall channel structure, comparison of
HCN112 and HCN1 in the ligand-free states reveals an important change in the quaternary
conformation (Figure 2). In HCN112, the cytoplasmic domain and transmembrane domain are
closer to each other in the direction perpendicular to the plane of the membrane, resulting in a
'compressed' mode of the TMD and CNBD (supplementary movie S1) in HCN112 compared to a
'relaxed' mode in HCN1. This compression amounts to ~3.5 Å when measuring the distance from
the top of the S5 helix (residue Q323) to the distal end of the CNBD (residue G547) (Figure 2A-C).
Importantly, as a result of the compression movement in HCN112 we observe a drastic increase in
the inter-chain interactions through hydrogen bonds and salt bridges (Figure 2D-F, Table 1). These
interactions (indicated in purple, Figure 2E-F) are unique for HCN112 and not present in HCN1.
For example, in the pore helix and middle of S6 there are unique H-bond interactions between
chain A residues L357, Y386 and chain B residue C358. Additionally, at the top of the pore helix
there is a unique H-bond between chain A residue A363 and chain B residue Q364. However, most
of the unique interactions in HCN112 between chain A and B are localized in the C-linker, a region
that has been shown repeatedly to be important for coupling ligand binding to channel opening (2,
5, 10, 12–31). In a recent study, the C-linker has also been implicated in the gating polarity of a
CNBD channel chimera composed of the EAG pore domain in a HCN1 background (32) . The
unique inter-chain salt bridges in the C-linker of HCN112 are D401-R297, R404-E283 and K430-
E460. The unique inter-chain H-bonds are R405-Q398, Q408-S293, K412-S399/L400, Q416-K442,
Y435-I449, Y439-E452 in the C-linker (Figure 2F). Lastly, there are unique inter-chain H-bonds
involving the CNBD, namely residues H517-F114 and R560-R112. As shown in Table 1 most of
the H-bonds present in HCN112 but not in HCN1 are localized in the C-linker, a region important
for coupling ligand binding to channel opening.
In summary, in HCN112 we observe a more compressed conformation of the structure in the plane
perpendicular to the membrane compared to HCN1. As a result, there are more inter-chain
interactions, especially in the C-linker which couples the CNBD to the TMD. We suggest that this
compressed conformation is one of the reasons why cAMP works more efficiently in HCN2 than in
HCN1 channels.
138
139 *cAMP binding breaks interactions in the compressed HCN112 conformation*
To study the compressed mode and the unique inter-chain interactions in HCN112 CL-CNBD in
more detail we next analyzed HCN112 in its cAMP-bound form.
The cryo-EM structure of cAMP-bound HCN112 was obtained from data at 3.26 Å resolution
(Figure 3A-C). The structure of cAMP-bound HCN112 superimposes onto the ligand-free HCN112
structure with an RMSD of 0.542 Å. The largest conformational change upon cAMP-binding can be
observed in the CNBD, which rotates and tilts away from the C-linker, and toward the C-terminal
helices D and E (Supplementary movie S2). These helices rigidify and become more structured
upon cAMP binding.
As a result of this conformational change there is a large decrease in the number of inter-chain
interactions under the form of hydrogen bonds and salt bridges. Specifically, there are 17 inter-
chain H-bonds and salt bridges in the ligand-free form of HCN112 (see Table 1). Most of these
interactions are broken in the cAMP-bound state except for 3 remaining H-bonds (K412-L400,
Y434-E452, Y439-E452) and 3 H-bonds that are newly formed (H421-N465, V495-N454, R548-

E575) (Figure 3F-H). For HCN112 in the ligand-free conformation we will only discuss those
interactions that likely contribute to gating of the channel.
155
156 **Table 1: Unique H-bonds and salt bridges observed in HCN112 but not in HCN1.**

Salt bridges	cAMP-free	cAMP-bound
CL	D401-R297	
	R404-E283	
	K430-E460	
H-bonds		
TMD	L357-C358	
	A363-Q364	
	Y386-C358	
CL	R405-Q39	
	Q408-S293	
	K412-S399/L400	K412-L400
	Q416-K442	
		H421-N465
		Y434-E452
	Y435-I449	
	Y439-E452	Y439-E452
		V495-N454
		R548-E575
CL-CNBD	H517-F114	
	R560-R112	

157
First, a key role appears to be played by R112 in the HCND domain. Three residues project
interactions onto R112, namely H437, Q440 (in the C-linker) and R560 (in the CNBD). Second,
there are 4 residues at the bottom of S4 and the S4-S5 linker that interact with the C-linker. These
include M287 (S4), D290 and S293 (S4-S5 linker), and R297 (S5). Lastly, two residues at the
bottom of S6, namely S399 and L400 interact with K412 in the C-linker.
In summary, our interpretation is that there are tight inter-chain interactions in the CNBD and C-
linker, also involving the HCND domain and S4-S5 linker, and that these interactions are broken
upon cAMP-binding. The breaking of these interactions could then contribute to the facilitation of
channel opening.
167
168
169 ***Deletion of the C-linker and CNBD causes partial channel opening in the HCN1ΔC channel***
170
It has been shown that the intracellular channel region is not only responsible for mediating cAMP
regulation, but also for the integrity of the channel gate, which is formed by the intracellular helix
bundle of the S6 segments. Previous studies proposed a gating mechanism in which the C-linker
disk plays an important role in keeping the gate closed at depolarized voltages: (1) by forming
intense contacts with the unusual S4 helix (3), and (2) by contacting the HCN domain of opposite
subunits (30). These interactions help to stabilize the gate in a closed right-handed helix bundle.
For channel opening, the CL-CNBD has to perform a leftward rotation to unwind the right-handed
S6 helix bundle gate. Such a leftward rotation is suggested to be induced by the movement of the
voltage sensor upon a hyperpolarizing voltage jump. Cyclic nucleotide binding can cause a CL-
CNBD leftward rotation, too. However, this cAMP-triggered rotation is not sufficient to open the

channel but is supportive for voltage-induced rotation. In this sense, cAMP causes the CL-CNBD
to adopt a conformation, which can be interpreted as relief of autoinhibition. To test this proposed
mechanism, we took advantage of the HCN1ΔC construct, in which the C-terminal cytoplasmic
domain was deleted (7). The deletion spans two domains, the C-linker and the CNBD, starting from
D401 till the last residue of HCN1 (Figure 1A-E). From the mechanism described above we expect
that the gate and the pore move towards their open configurations when the CL-CNBD is missing.
Gating of HCN channel is activated by membrane hyperpolarization and is also modulated by
cAMP. It has been shown that the cyclic nucleotide–binding domain has an inhibitory effect on
channel gating that is relieved by cAMP binding (7). Such a relieving effect is reflected by an
increase in current amplitude at a given voltage, by a shift of the voltage of half-maximum activation,
$V_{1/2}$, to more depolarized values, and by an acceleration of activation kinetics and a deceleration
of deactivation kinetics (33).
193

Here, we determined the cryo-EM structure of HCN1ΔC, which reached an overall resolution of
3.96Å (Figure 4). HCN1ΔC shows an RMSD value of 1.114Å and 1.096Å when aligned to the TMD
domain of ligand-free HCN112 and cAMP-bound HCN112, respectively. Our HCN1ΔC structure
shows a movement of pore helices S6 and S5, as shown in the inset in Figure 4 and supplementary
movie S3. Overlay of cryo-EM densities of ligand-free HCN112 and HCN1ΔC (bottom left inset)
clearly shows this redistribution of helices and inner sidechains along the pore.
200

Determinant aspects confirming the widening of the pore in HCN1ΔC are sidechains of V390, T394
and Q398. Interestingly, these residues have also been shown before to be key elements in the
inner channel gate (3). The long sidechain of Q398 at the intracellular gate of the pore tilts away
from the inner pore cavity, widening the diameter of the intracellular gate of HCN1ΔC (colored in
green mesh - bottom right inlet). The HCN1ΔC pore diameter is close to double in size of the
HCN112 pore, which is depicted in yellow colour and overlayed in the bottom-up view in Figure 4A
and B. These small differences in conformation of S6-S5 helices in HCN1ΔC led the overall pore
diameter to be halfway toward being fully open (~3Å vs ~5Å) and thus resembles the open HCN1
conformation as well as the rabbit HCN4 open pore conformation (Figure 4B-C).
210

In summary, we hypothesized that the S6-S5 and S4 helices movements cause a loss of contacts
between these pore helices and the C-linker, as well as between the HCND and the CNBD. We
therefore suggest that these rearrangements reflect the transition state towards an open pore in
HCN1, triggering a release of constraints and allowing the pore to open. We also propose that our
HCN1ΔC structure, with its deletion reflecting an absence of auto-inhibition, mimics a transitioning-
hyperpolarized state, that most likely precedes the hyperpolarized state described in Lee and
MacKinnon (4).
218

220
221   **Materials and Methods**
222

Constructs design
All HCN1 constructs were engineered in the background of the human HCN1$_{EM}$ plasmid made by
Lee and MacKinnon (3) and was cloned into a pEG-BacMam vector (34) for expression in
mammalian cells and which tags the protein of interest with both N-His10, eGFP and 3C protease
site. The following constructs were generated on top of the HCN1$_{EM}$ constructs: 1. Chimera
HCN112 contains region Met1-Leu400 from human HCN1 and Asp401-Leu658 from human HCN2,
with the region between Ala705-Ser866 of HCN2 which was deleted, while the HCN2 extreme C-
terminal region including the SNL tripeptide sequence was preserved Ala867-Leu889 (residue
numbering according to full-length human HCN2); 2. HCN1ΔC corresponds to the HCN1 gene
covering the sequence between Met1-Asp401, with a deletion that spans the entire C-terminal
region.

Electrophysiological recordings

For expression in *Xenopus* oocytes HCN112 and HCN1ΔC were subcloned into pGEM-HE using *Xba*I and *Hind*III restriction sites. Plasmids were linearized with *Nhe*I and transcribed with T7 RNA polymerase using the mMessage mMachine T7 transcription kit. The RNA quality was verified by loading RNA samples on 1% agarose gel.

Oocytes were surgically removed from adult female South African claw frogs *Xenopus laevis* under anesthesia with 0.3% tricaine methanesulfonate (MS-222) (Pharmaq Ltd. Fordingbridge, United Kingdom). After removal, the oocytes were treated with collagenase A (3 mg/ml; Roche, Grenzach-Wyhlen, Germany) for 105 min in Ca^{2+}-free Barth's solution containing (in mM) 82.5 NaCl, 2 KCl, 1 $MgCl_2$, and 5 Hepes, pH 7.5. Oocytes of stages IV and V were manually dissected and injected with cRNA encoding either HCN112 or HCN1ΔC channels. After injection with cRNA, the oocytes were incubated at 18°C for 2–6 days in Barth's solution containing (in mM) 84 NaCl, 1 KCl, 2.4 $NaHCO_3$, 0.82 MgSO4, 0.41 $CaCl_2$, 0.33 Ca(NO3)$_2$, 7.5 TRIS, pH 7.4. Oocytes harvested in our lab were complemented with ready-to-use oocytes purchased from Ecocyte Bioscience (Dortmund, Germany). The surgery procedures were carried out in accordance with the German Animal Welfare Act with the approval of the Thuringian State Office for Consumer Protection on 30.08.2013 and 09.05.2018.

Macroscopic currents were recorded using the inside-out configuration of the patch-clamp technique. All measurements were started after a delay of 3.5 min to minimize run-down phenomena. Patch pipettes were pulled from quartz tubings with outer and inner diameters of 1.0 and 0.7 mm (VITROCOM, New Jersey, United States), respectively, using a laser puller (P-2000, Sutter Instrument, Novato, United States). The pipette resistance was 1.2–2.1 MOhm. The bath solution contained (in mM) 100 KCl, 10 EGTA, and 10 Hepes, pH 7.2, and the pipette solution contained (in mM) 120 KCl, 10 Hepes, and 1.0 $CaCl_2$, pH 7.2. For parts of the experiments, a saturating concentration of 10 µM cAMP (BIOLOG LSI GmbH & Co. KG, Bremen, Germany) was applied with the bath solution. A HEKA EPC 10 USB amplifier (Harvard Apparatus, Holliston, United States) was used for current recording. Pulsing and data recording were controlled by the Patchmaster software (Harvard Apparatus, Holliston, United States). The sampling rate was 5 kHz. The holding potential was generally −30 mV. Maximally two membrane patches were excised from one individual oocyte.

Protein expression, purification and sample preparation for Cryo-EM

The different HCN constructs were expressed in HEK293S GnTI⁻ cells (Goehring et al., 2014) at 37ºC and to a density of ~$3x10^6$ cells/ml for ~72h, after addition of baculoviruses carrying HCN1 genes (10% v/v). After 15h, 10 mM sodium butyrate was added to cells and the culture temperature was shifted to 30ºC for ~72 h, after which cells were harvested and pellets were flash frozen and stored at -80ºC.

Cell pellets were resuspended in 30% glycerol for 10 min and lysed by gentle homogenization in 10 mM Tris pH 8.0, 20 mM KCl, 2 mM DTT, 0.1 mg/ml DNAse, 0.5 mM $MgCl_2$, 2 mg/ml leupeptin, 10 mg/ml aprotinin, 1 mg/ml pepstatin, 50 mg/ml benzamidine HCl, 1 mM AEBSF, 30% glycerol for 25 min at 4ºC. The lysate was spun at 39,800g for 35 min at 4ºC to sediment crude membranes. The membrane pellet was homogenized and solubilized in the extraction buffer (20 mM Tris pH 8.0, 300 mM KCl, 2 mM DTT, 2 mg/ml leupetin, 10 mg/ml aprotinin, 1 mg/ml pepstatin, 50 mg/ml benzamidine HCl, 1 mM AEBSF, 10 mM LMNG + 2 mM CHS) for 2 h. Solubilized membranes were clarified by centrifugation at 39,800g for 35 min, 4ºC. The supernatant was incubated with the GFP nanobody-coupled sepharose resin prepared in-house (35) for 1.5 h at 4ºC and subsequently washed for 10 column volumes with wash buffer (20 mM Tris pH 8.0, 300 mM KCl, 2 mM DTT, 2 mg/ml leupetin, 10 mg/ml aprotinin, 1 mg/ml pepstatin, 50 mg/ml benzamidine HCl, 1 mM AEBSF, 0.05% digitonin). An extra 5 column volumes of buffer with 1 mM EDTA was used to incubate the proteins of interest overnight at 4ºC with PreScission protease to cleave the GFP tag. HCN variants were eluted straight on a Glutathione Sepharose 4B column and washed out with wash buffer. The eluted proteins were concentrated and purified further through a Superose 6 10/300 column using gel filtration buffer (20 mM Tris pH 8.0, 150 mM KCl, 2 mM DTT, 2 mg/ml leupeptin, 10 mg/ml aprotinin, 1 mg/ml pepstatin, 50 mg/ml benzamidine HCl, 1 mM AEBSF, 0.05% digitonin). Peak

fractions of interest were pooled, concentrated to 3-4 mg/ml, flash frozen in liquid nitrogen and
stored at -80C.
Once thawed, the purified proteins were spun at 13,000 rpm for 5 min at 4ºC and the supernatant
was used for grid preparation. 0.4 mM cAMP was added to 3.5 mg/ml HCN112 in order to perform
single particle analysis on the HCN112 cAMP-bound sample. 3 ml of the purified HCN variants
were applied on glow discharged AuFoil grid R2/2 200 mesh within a Vitrobot IV (FEI) operated a
4ºC, 100% humidity (blot time 3 s, waiting time 3 s) and plunged into liquid ethane.
295
296 **Cryo-EM data acquisition**
297
298 Two data acquisitions were acquired using two Titan Krios transmission electron microscopes (FEI)
299 operated at 300keV with energy filter, equipped with Falcon 4 electron detector (ABSL, Leeds, UK)
300 at a nominal magnification of 130k, with a pixel size of 0.91Å (HCN112+cAMP and HCN1ΔC). Two
301 other datasets, later merged as one single dataset, were acquired at a nominal magnification of
302 96k, with a pixel size of 0.81Å (ligand-free HCN112).
303
The nominal defocus range chosen was between the following values and changing in increments
of 0.2: for apoHCN112 this was 1.6 - to 3.2 mm, for HCN112+cAMP this was 1.2 to 3 mm, for
HCN1ΔC this was 1.6 to 3.4 mm. Respectively, the total doses were: 41.42 e$^-$/Å^2 (merged ligand-
free HCN112), 32.44 e$^-$/Å^2 (HCN112+cAMP) and 39.81 e$^-$/Å^2 (HCN1ΔC). The exposure time for
each image was 6.26 s (ligand-free HCN112), 3.98s (HCN112+cAMP), 6.49s (HCN1ΔC),
fractionated over 54, 41, 40 EER fractions, respectively. Acquisition parameters are summarized
in Table S1.
311
312 **Image processing and single-particle analysis**
313
Movies were corrected for beam-induced motion using MotionCor (relion3.1.1) (36) and contrast
transfer function parameters were estimated from motion-corrected images using CTFFIND 4.1.
For all datasets, particle picking was performed in crYOLO v1.61_GPU (37) using an integrated
general model and cutoff 0.1 and JANNI denoised. Particles were extracted in Relion with a box
size of 350px. The particle.star file was then imported in cryoSPARC v.3.2.0 (38), where 2D
classification, ab-initio reconstruction and refinements were performed (Suppl Fig. 2-7 for Cryo-EM
flowchart). During refinement, C4 symmetry was imposed in all datasets. The same single-particle
processing was applied for all datasets, leading to a local resolution of 2.9 Å ligand-free HCN112,
2.7Å HCN112+cAMP, 3.45Å HCN1ΔC.
323
Model building, refinement and validation
Manual building and real-space refinement was performed using HCN1 (PDB 5u6o) as starting
model for the transmembrane region (TM) and HCN2 crystal structure (PDB 3u10) for the
cytoplasmic domain (CD) in Coot v0.8.9.2 (39) and then in Phenix v1.19.2 (40). These homology
models were docked into the density map of the ligand-free HCN112 using UCSF chimera (41) in
order to build a starting model for one protomer of the chimeric HCN112. A tetramer model of the
channel was obtained by fitting copies of the same model in the C4 symmetric map of the tetramer.
The starting model for the cAMP-bound HCN112 was the refined ligand-free HCN112. The
placement of the cAMP ligand in the holoHCN112 was based on the crystal structure of the HCN2
in complex with cAMP (PDB 3u10). As a model for HCN1ΔC TM region, the cAMP-bound HCN112
was used but manual rebuilding was necessary for amino acid fragments that undergo
conformational changes i.e. S6, S5 and HCN domain.
The final ligand-free structure includes residues 97–240, and 252–579, whereas the cAMP-bound
structure includes additional residues 580–605 and part of the E-helices built as a poly-Ala helix.
For the HCN1ΔC structure building was more challenging due to the lower quality of the cryo-EM
map, but residues of the pore region (361-398) could be built using the HCN112+cAMP structure
as a reference thus allowing analysis of the pore radius profile. The quality of the final models was
evaluated by MolProbity (42) and EMRinger (43) (Table S1). Figures were prepared using Chimera

(41), ChimeraX (44)and PyMOL Molecular Graphics System (v1.8, Schrodinger, LLC). Pore analysis was performed in HOLE v2.2.005 (45) after superposition of all protein structures of interest. Interactions between residues were calculated using Contact/Act in the CCP4 suite (46) using a maximum distance cut-off of 3.8 Å. H-bonds and salt bridges were considered between 2.6-3.4 Å and van der Waals interactions were considered between 3.2-3.8 Å.

Acknowledgments

The research by CU was supported by KU Leuven grant C14/23/128 and grants G0C1319N, G087921N from FWO-Vlaanderen. We thank Dr. Rebecca Thompson and all the Cryo-EM team at Astbury Center for Molecular and Structural Biology, Leeds, UK for access to and technical support for Krios Titans. The access was granted by iNext-Discovery. VZ was funded by Fund for Scientific Research Flanders Postdoctoral Fellowship (FWO 12W4618N).

Data resources
Cryo-EM density maps of HCN112 apo, HCN112+cAMP, and HCN1ΔC have been deposited in the electron microscopy data bank under accession code EMD-53516, EMD-53514 and EMD-53515, respectively. Atomic coordinates of HCN112 apo, HCN112+cAMP and HCN1ΔC have been deposited in the protein data bank under accession code 9R1V, 9R1T and 9R1U, respectively.

References

1. B. Santoro, M. M. Shah, Hyperpolarization-Activated Cyclic Nucleotide-Gated Channels as Drug Targets for Neurological Disorders. *Annu. Rev. Pharmacol. Toxicol.* **60**, 109–131 (2020).
2. J. Wang, S. Chen, S. A. Siegelbaum, Regulation of hyperpolarization-activated HCN channel gating and cAMP modulation due to interactions of COOH terminus and core transmembrane regions. *The Journal of General Physiology* **118**, 237–250 (2001).
3. C.-H. Lee, R. Mackinnon, Structures of the Human HCN1 Hyperpolarization- Activated Channel. *Cell* **168**, 111-120.e11 (2017).
4. C.-H. Lee, R. Mackinnon, Voltage Sensor Movements during Hyperpolarization in the HCN Channel. *Cell* (2019). https://doi.org/10.1016/j.cell.2019.11.006.
5. A. Saponaro, *et al.*, Gating movements and ion permeation in HCN4 pacemaker channels. *Mol Cell* (2021). https://doi.org/10.1016/j.molcel.2021.05.033.
6. A. Wickenden, *et al.*, HCN Channels as Targets for Drug Discovery. *Comb. Chem. High Throughput Screen.* **12**, 64–72 (2009).
7. B. J. Wainger, M. DeGennaro, B. Santoro, S. A. Siegelbaum, G. R. Tibbs, Molecular mechanism of cAMP modulation of HCN pacemaker channels. *Nature* **411**, 805–810 (2001).
8. B. Yu, *et al.*, Cryo-EM structure of human HCN3 channel and its regulation by cAMP. *J. Biol. Chem.* **300**, 107288 (2024).
9. V. Burtscher, *et al.*, Structural basis for hyperpolarization-dependent opening of human HCN1 channel. *Nat. Commun.* **15**, 5216 (2024).
10. W. N. Zagotta, *et al.*, Structural basis for modulation and agonist specificity of HCN pacemaker channels. *Nature* **425**, 200–205 (2003).
11. A. Saponaro, *et al.*, Structural basis for the mutual antagonism of cAMP and TRIP8b in regulating HCN channel function. *Proceedings of the National Academy of Sciences* **111**, 14577–14582 (2014).
12. N. Decher, J. Chen, M. C. Sanguinetti, Voltage-dependent Gating of Hyperpolarization-activated, Cyclic Nucleotide-gated Pacemaker Channels MOLECULAR COUPLING BETWEEN THE S4–S5 AND C-LINKERS*. *J. Biol. Chem.* **279**, 13859–13865 (2004).

13. L. Zhou, N. B. Olivier, H. Yao, E. C. Young, S. A. Siegelbaum, A Conserved Tripeptide in CNG
and HCN Channels Regulates Ligand Gating by Controlling C-Terminal Oligomerization. *Neuron*
44, 823–834 (2004).
14. K. B. Craven, W. N. Zagotta, Salt bridges and gating in the COOH-terminal region of HCN2 and
CNGA1 channels. *The Journal of General Physiology* **124**, 663–677 (2004).
15. J. P. Johnson, W. N. Zagotta, The carboxyl-terminal region of cyclic nucleotide-modulated
channels is a gating ring, not a permeation path. *Proc. Natl. Acad. Sci.* **102**, 2742–2747 (2005).
16. D. L. Prole, G. Yellen, Reversal of HCN channel voltage dependence via bridging of the S4-S5
linker and Post-S6. *The Journal of General Physiology* **128**, 273–282 (2006).
17. L. Zhou, S. A. Siegelbaum, Gating of HCN Channels by Cyclic Nucleotides: Residue Contacts
that Underlie Ligand Binding, Selectivity, and Efficacy. *Structure* **15**, 655–670 (2007).
18. X. Xu, Z. V. Vysotskaya, Q. Liu, L. Zhou, Structural Basis for the cAMP-dependent Gating in
the Human HCN4 Channel. *Journal of Biological Chemistry* **285**, 37082–37091 (2010).
19. M. Lolicato, *et al.*, Tetramerization dynamics of C-terminal domain underlies isoform-specific
cAMP gating in hyperpolarization-activated cyclic nucleotide-gated channels. *Journal of Biological*
Chemistry **286**, 44811–44820 (2011).
20. D. C. H. Kwan, D. L. Prole, G. Yellen, Structural changes during HCN channel gating defined
by high affinity metal bridges. *The Journal of General Physiology* **140**, 279–291 (2012).
21. M. Lolicato, *et al.*, Cyclic dinucleotides bind the C-linker of HCN4 to control channel cAMP
responsiveness. *Nature chemical biology* (2014). https://doi.org/10.1038/nchembio.1521.
22. M. Akimoto, *et al.*, A Mechanism for the Auto-Inhibition of Hyperpolarization-Activated Cyclic
Nucleotide-Gated (HCN) Channel Opening and its Relief by cAMP. *Journal of Biological Chemistry*
(2014). https://doi.org/10.1074/jbc.m114.572164.
23. M. Nieves-Cordones, I. Gaillard, Involvement of the S4-S5 linker and the C-linker domain
regions to voltage-gating in plant Shaker channels: Comparison with animal HCN and Kv channels.
Plant Signal. Behav. **9**, e972892 (2014).
24. H. A. DeBerg, P. S. Brzovic, G. E. Flynn, W. N. Zagotta, S. Stoll, Structure and Energetics of
Allosteric Regulation of HCN2 Ion Channels by Cyclic Nucleotides*. *J. Biol. Chem.* **291**, 371–381
(2016).
25. S. Weißgraeber, A. Saponaro, G. Thiel, K. Hamacher, A reduced mechanical model for cAMP-
modulated gating in HCN channels. *Sci. Rep.* **7**, 40168 (2017).
26. C. Gross, *et al.*, Mechanical transduction of cytoplasmic-to-transmembrane-domain
movements in a hyperpolarization-activated cyclic nucleotide–gated cation channel. *J. Biol. Chem.*
293, 12908–12918 (2018).
27. C. P. Alvarez-Baron, V. A. Klenchin, B. Chanda, Minimal molecular determinants of isoform-
specific differences in efficacy in the HCN channel family. *J. Gen. Physiol.* **150**, 1203–1213 (2018).
28. C. Pfleger, *et al.*, Allosteric signaling in C-linker and cyclic nucleotide-binding domain of HCN2
channels. *Biophys. J.* **120**, 950–963 (2021).
29. G. Dai, T. K. Aman, F. DiMaio, W. N. Zagotta, Electromechanical coupling mechanism for
activation and inactivation of an HCN channel. *Nat. Commun.* **12**, 2802 (2021).
30. M. Kondapuram, *et al.*, Functional and structural characterization of interactions between
opposite subunits in HCN pacemaker channels. *Commun. Biol.* **5**, 430 (2022).
31. S. Yüksel, *et al.*, Uncoupling of Voltage- and Ligand-Induced Activation in HCN2 Channels by
Glycine Inserts. *Front. Physiol.* **13**, 895324 (2022).
32. J. L. Lin, Y. Chang, D. Tewari, J. Cowgill, B. Chanda, Mapping the contribution of the C-linker
domain to gating polarity in CNBD channels. *Biophys. J.* **123**, 2176–2184 (2024).
33. D. DiFrancesco, P. Tortora, Direct activation of cardiac pacemaker channels by intracellular
cyclic AMP. *Nature* **351**, 145–147 (1991).
34. A. Goehring, *et al.*, Screening and large-scale expression of membrane proteins in mammalian
cells for structural studies. *Nature protocols* **9**, 2574–2582 (2014).
35. A. Kirchhofer, *et al.*, Modulation of protein properties in living cells using nanobodies. *Nature*
structural & molecular biology **17**, 133–138 (2010).
36. S. H. W. Scheres, RELION: Implementation of a Bayesian approach to cryo-EM structure
determination. *J. Struct. Biol.* **180**, 519–530 (2012).

37. T. Wagner, *et al.*, SPHIRE-crYOLO is a fast and accurate fully automated particle picker for
cryo-EM. *Commun. Biol.* **2**, 218 (2019).
38. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).
39. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta*
crystallographica Section D, Biological crystallography **66**, 486–501 (2010).
40. P. D. Adams, *et al.*, PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta crystallographica Section D, Biological crystallography* **66**, 213–221 (2010).
41. E. F. Pettersen, *et al.*, UCSF Chimera--a visualization system for exploratory research and
analysis. *Journal of Computational Chemistry* **25**, 1605–1612 (2004).
42. V. B. Chen, *et al.*, MolProbity: All-atom structure validation for macromolecular crystallography.
Acta crystallographica Section D, Biological crystallography **66**, 12–21 (2010).
43. B. A. Barad, *et al.*, EMRinger: side chain–directed model and map validation for 3D cryo-
electron microscopy. *Nat. Methods* **12**, 943–946 (2015).
44. E. F. Pettersen, *et al.*, UCSF ChimeraX : Structure visualization for researchers, educators, and
developers. *Protein Sci.* **30**, 70–82 (2020).
45. O. S. Smart, J. G. Neduvelil, X. Wang, B. A. Wallace, M. S. P. Sansom, HOLE: A program for
the analysis of the pore dimensions of ion channel structural models. *Journal of molecular graphics*
14, 354–360 (1996).
46. M. D. Winn, *et al.*, Overview of the CCP4 suite and current developments. *Acta*
crystallographica Section D, Biological crystallography **67**, 235–242 (2011).

469
470
Figure legends

Figure 1. Structure and function of HCN112 and HCN1DC channels. a. Transmembrane
topology of the HCN112 and HCN1ΔC channel constructs. The dashed line in HCN112 indicates
the C-linker and CNBD, which have been replaced with the corresponding sequence from HCN2
channels. **b**. Representative current traces from HCN112 and HCN1ΔC channels expressed in
Xenopus oocytes and evoked by a voltage step to -100 mV in the absence (black) and presence
(red) of 10 mM cAMP. The graphs on the right show the summary of peak current amplitudes from
4 individual experiments. **c-e**. Cryo-EM maps of HCN112 ligand-free (yellow), HCN112 cAMP-
bound (cyan) and HCN1ΔC (green). The 3D-structures are docked in the map and shown in cartoon
representation. The top panels are side-views, the bottom panels are a bottom-up views.

Figure 2. HCN112 channels show a compressed conformation and unique inter-chain
interactions. a-c. Cartoon representations of HCN1 ligand-free (**b**, red, pdb code 5u6o), HCN112
ligand-free (**c**, yellow) and overlay (**a**). Only 2 opposing subunits are shown for clarity. The green
spheres and the dashed line show the distance between the top of S5, residue Q323 and the distal
end of the CNBD, residue G547. **d-e**. Inter-chain interactions between protomor A (yellow ribbon)
and protomor B (blue ribbon) for HCN1 ligand-free (**d**, pdb code 5u6o) and HCN112 ligand-free (**e**).
Interacting residues are shown in sphere representation. Hydrogen bonds are shown between
yellow (protomor A) and dark blue (protomor B) residues. Salt bridges are shown between pale
yellow (promotor A) and pale blue (protomor B) residues. Hydrogen bonds and salt bridges that are
unique in HCN112 (**e**) are shown as magenta and pink residues. **f**. Detailed view of the uniquely
interacting residues in the C-linker region. Residues of protomor A are shown in magenta ball and
stick representation, residues of protomor B are shown in pink. Oxygen atoms are shown in red,
nitrogen atoms are shown in blue.

Figure 3. cAMP binding causes a conformational change of the C-linker and CNBD, and
breaks inter-chain interactions in HCN112. a-c. Cartoon representation of HCN112 in the ligand-

free conformation (**a**, yellow), cAMP-bound conformation (**b**, cyan) and overlay (**c**). Only 2 opposing
subunits are shown for clarity. **d**. The inset shows a detailed view of conformational changes in the
C-linker and CNBD. **e-h.** Interchain-interactions between promotor A (yellow) and promotor B (blue)
in the HCN112 ligand-free conformation and promotor A (cyan) and protomer B (magenta) in the
HCN112 cAMP-bound conformation. **g-h**. The insets show a detailed view of inter-chain
interactions. Interacting residues are shown in ball and stick representation. Dashed lines show
hydrogen bonds or salt bridges. Oxygen atoms are shown in red, nitrogen atoms are shown in blue.

Figure 4. Relief from auto-inhibition in HCN1ΔC reveals a pore in an intermediately open
conformation. a. Pore radius calculations for HCN112 ligand-free (yellow), HCN112 cAMP-bound
(cyan), and HCN1ΔC (green) are shown on the graph (left). The corresponding structures of the
pore region (A and C chains, residues 358 to 401) are shown as cartoons (right), with the sidechains
of key pore-lining amino acids shown as licorice sticks. The selectivity filter, cavity, and gate regions
are indicated on both the graph (left) and structures (right) with dotted lines. Grey spheres within
the structures represent the calculated pore. **b.** Overlay of two subunits of HCN112 ligand-free
(yellow) and HCN1ΔC (green), is shown as cartoons on the top left, with a zoomed bottom-up view
on the top right. The C-linker and CNBD domain from the HCN112 ligand-free structure are colored
grey. A zoomed-in view of the pore region in the overlapped structures is displayed in the bottom
panels. The bottom left panel shows the overlapped structures with their corresponding cryo-EM
maps, while the bottom right panel focuses on displaying the calculated pore radius within the
structures (yellow sphere and green mesh) and highlighting the residue Q398 for all S6 helices. A
side view of the calculated pore for both structures is included in the bottom-right corner,
emphasizing the position of Q398 in the gate region. **c.** Pore radius calculations of the HCN1ΔC
structure, compared to published HCN1 structures in different conformational states, are plotted.
The approximate locations of the selectivity filter, cavity, and gate regions are indicated for
reference. Comparison of the pore radius in the closed ligand-free HCN1 conformation (pdb code
5u6o), the closed cAMP-bound HCN1 conformation (pdb code 5u6p), the closed HCN1 F186C
S264C bound to cAMP (pdb code 8t4m), the open HCN1 F186C S264C bound to cAMP (pdb code
8T50) and the intermediately open HCN1ΔC.

Figure 5. Models of domain compression in HCN112 and relief from auto-inhibition in
HCN1ΔC. The models illustrate the domain compression in HCN112 (yellow) upon comparison
with HCN1 in the closed ligand-free conformation (red). The green model shows relief from auto-
inhibition in HCN1ΔC resulting in a partially open pore conformation. The figure was adapted from
Lee and MacKinnon (2017) with permission from Elsevier.

Figure 1

a

HCN112

HCN1ΔC

b

-30 mV
-100 mV

50 pA
500 ms

HCN112

20 pA
500 ms

HCN1ΔC

HCN112

HCN1ΔC

I_{max} (pA)

control   10 μM cAMP

control   10 μM cAMP

c   **d**   **e**

HCN112 ligand-free   HCN112 cAMP-bound   HCN1ΔC

TMD

90°   90°   90°

Bottom-up view   Bottom-up view   Bottom-up view

Figure 2

Figure 3

a

Figure 4

HCN112 ligand-free

HCN112 cAMP-bound

HCN1ΔC

b

TMD

Pore
S6

Bottom-up view

Bottom-up views

Q398

Pore

Selectivity Filter

Cavity

Gate

c

HCN1 pore analysis

— HCN1ΔC
— 8T50 cAMP_opened
— 8T4M cAMP_closed
— 5U6O ligand-free
— 5U6P cAMP

Figure 5

HCN1 channel

relaxed

HCN112 channel

compressed

HCN1ΔC channel

relieved from
auto-inhibition