A mathematical kinetic model of memory in *Bacillus subtilis* spore germination

Chris G. de Koster[a] [□], Xiuping Lin[c], Yong-qing Li[c], Stanley Brul[b*], Leo J. de Koning[a] and Peter Setlow[d]

a) *Mass Spectrometry of Biomacromolecules and (b) Molecular Biology & Microbial Food Safety, Swammerdam Institute for Life Sciences, University of Amsterdam, Science Park 904, 1098 XH, Amsterdam, The Netherlands.*

c) *Department of Physics, East Carolina University, Greenville, North Carolina, USA[c].*

d) *Department of Molecular Biology and Biophysics, UConn Health, Farmington, Connecticut, USA[d].*

□ In memory of prof. Chris de Koster who passed away 5th August 2019 * Author to whom correspondence should be addressed s.brul@uva.nl

Keywords: *Bacillus subtilis*, germination, memory, germinosome, model

Abstract

Dormant *Bacillus subtilis* spores germinate through interaction of germinants with germinant receptors (GRs). Subsequently, the activated GR signal is transduced to SpoVA proteins that constitute transport channels. The opening of the SpoVA channel then leads to calcium dipicolinic acid (CaDPA) release and then completion of spore germination. Spores are known to exhibit memory in germination, as spores given an initial short germinant pulse respond more readily to a second pulse. We have developed a mathematical model to identify a minimal reconstructed molecular network that is crucial for germination kinetics leading to memory of germinant exposure; the model reproduces experimental double germinant pulse germination curves. Analysis of the reconstructed

network indicates that a minimal set of inactive and active GRs and a SpoVA channel in three states -

closed inactive, closed active and open - is needed to reproduce memory. Spore germination memory

is introduced in the network by the activation and deactivation rates of GRs, and by the interplay

between activation of closed SpoVA channels and their rate of opening and closing.

Author summary

Dormant bacterial spores return to life in germination when exposed to germinants which are

recognized by germinant receptors (GRs). Spores have memory of germinant exposure, such that

more spores germinate after a 2^{nd} germinant pulse than after the 1^{st} one, if intervals between pulses are

short. In this work, a mathematical model has been developed that reproduces experimental memory

behavior. Computer simulations of the model reveal the crucial factors in spore memory of germinant

exposure as: i) active and inactive GR states and ii) a membrane channel for a major spore core

molecule, Ca-dipicolinic acid (CaDPA), with the channel existing in closed inactive, closed active and

open states, all of which are indispensable for spore memory. The model will be part of an iterative

cycle of modelling of network dynamics and generating hypotheses for further experimentation.

Introduction

Spores of bacteria of *Bacillus subtilis* are formed in sporulation, and are dormant and with an overall structure quite different from that of a growing cell (1). From the outside in, spores contain an exosporium in some species (but not *B. subtilis* spores!), then a layer of protein termed the coat, then an outer membrane above the spore's peptidoglycan cortex that contains muramic acid lactam, next a layer of germ cell peptidoglycan with no muramic acid lactam, then an inner membrane (IM) encircling the spore core that contains all nucleic acids, most spore enzymes and a high level of the 1:1 chelate of Ca^{2+} with pyridine-2,6-dicarboxlic acid (CaDPA, ~ 10% of spores core dry wt), as well as low levels of core water, as low as 20% of core wet weight (1). Spores are extremely resistant to all manner of treatments, have no ATP, but can survive harsh environmental conditions for many years (1-3). However, when nutrients become available and conditions are favorable for growth, spores can germinate and outgrow into vegetative cells via a sequence of molecular processes and signaling through pathways and networks (4-7). In the 1[st] stage of germination the spore probes its environment for the availability of nutrient germinants. Upon interaction of germinants with germinant receptors (GRs) in the IM, spores germinate. *Bacillus* GRs generally have 3 subunits, A, B and C and these are gated ion channels with other ion channels participating (8-12). The binding of germinants, in particular to GR's B subunits, leads to release of the spore's huge depot of (CaDPA). This release then triggers the activation of two cortex-lytic enzymes (CLEs), CwlJ and SleB. In the 2[nd] stage of germination, the CLEs hydrolyze the large spore cortex peptidoglycan (PG) layer, as these enzymes recognize muramic acid lactam which is only present in cortex PG (1-4). Cortex degradation then allows core expansion with full hydration of the spore core which allows resumption of normal metabolism, macromolecular synthesis, and finally outgrowth into a vegetative growing cell.

The GR protein assemblies GerA, GerB and GerK in *B. subtilis* are each encoded by homologous tricistronic operons *gerA*, *gerB* and *gerK* (1-4). GR subunits A and B are integral membrane proteins that are embedded in the spores' IM through multiple predicted α-helices. Subunit C is a peripheral membrane protein, attached to the IM by a diacylglycerol anchor (13). The GerA GR responds to L-alanine or L-valine, the GerB and GerK GRs cooperatively respond to a mixture of L-asparagine, D-

glucose, D-fructose and K^+ (AGFK). A GerB variant, GerB*, germinates in response to L-asparagine

alone. Average wild-type GR spore copy numbers have been determined by Western blotting and are

~2,500 total GRs/spore, with GerA at ~1,100 molecules/spore; and GerB and GerK each at ~700

molecules/spore (14). As expected, GR levels in spores affect the germination rate, and GR

overexpression increases rates of spore germination by these GRs (15). All *B subtilis* GRs co-localize

in the spores' IM in a cluster termed the germinosome (16), and germinosomes have also been found

in other *Bacillus* species (17). The GerD protein co-localizes with GRs in the germinosome and is

essential for germinosome formation, presumably acting as a scaffold, although the germinosome

structure has not been determined. GerD is also essential for normal GR-dependent spore germination,

as *gerD* spores germinate extremely poorly with L-valine via GerA or AFGK via GerB plus GerK

(18). Like GR C subunits, GerD is also likely attached to the IM through an N-terminal diacylglycerol

anchor attached to an N-terminal cysteine residue (18).

The signal generated upon GR activation, likely involving cations released from the spore core, is

transduced to SpoVA proteins that make up a channel for CaDPA in the spore IM. However, the

signaling mechanism is not fully understood, and no proteins that act in between the GRs and the

SpoVA channel protein complex have been identified. The *Bacillus subtilis spoVA* operon encodes

SpoVAA, SpoVAB, SpoVAC, SpoVAD, SpoVAEb, SpoVAEa and SpoVAF proteins. SpoVAA,

SpoVAB, SpoVAC, SpoVAEb and SpoVAF are predicted to be integral membrane proteins based on

their primary sequences, and SpoVAD and SpoVAEa are hydrophilic, soluble proteins that are on the

outer IM surface (19,20). The copy numbers of SpoVAD and SpoVAEa have been determined by

western blotting a ~6,000 - 6,500 and 750 molecules/spore, respectively (19,20). SpoVAC alone

functions as a mechanosensitive channel for small molecules such as CaDPA, with several treatments

including membrane tension and the non-GR-dependent germinant dodecylamine opening this

channel in vivo and in vitro (4,21,22). While the arrangement of SpoVA proteins in a supramolecular

structure is not known, it seems most likely that the SpoVA proteins are subunits of a protein

assembly in the spore's IM that constitutes a channel for CaDPA movement into the developing spore

during sporulation and out of the germinating spore (23). In contrast to GRs which are in the

multiprotein germinosome, SpoVA proteins are more evenly distributed over the spore IM (16).

Opening of the CaDPA channel is triggered not only by GR activation, but also by dodecylamine or

hydrostatic pressures of ≥ 4000 atmospheres (4). Notably, while there is slow leakage of CaDPA from

spores after germination is begun (24), almost all CaDPA is released in a very short time and in

parallel with water uptake into the core (4).

In the 2^{nd} germination stage, CaDPA activates the CLEs which hydrolyze the cortex PG. CwlJ is

activated first by CaDPA released from the spore core and SleB is activated only after substantial

CaDPA release, perhaps by changes in cortex PG structure accompanying CaDPA release and water

uptake. CwlJ can also be activated by exogenous CaDPA. The structure of the dormant spore cortex

PG and physical effects due to elastic and contractile properties of the cortical PG sacculus

presumably affect the susceptibility of opening of SpoVA channels , as does crosslinking in the spore

coat which continues even after spores are released from sporangia (25,26). Cortex hydrolysis

degrades the cortex PG macromolecular network restricting expansion of the spore core, thus enabling

further uptake of water. As core water content increases, eventually full core hydration allows onset of

metabolic reactions including biosynthesis of molecules needed for cell growth.

Spore germination is a heterogeneous process, as in populations some spores germinate rapidly and

some extremely slowly (27-31). This heterogeneity is due primarily to a variable time period between

the application of a germinant stimulus and the onset of rapid CaDPA release from the spore core.

This time interval is denoted as the lag time T_{lag}; all CaDPA is released by $T_{release}$, and cortex PG

hydrolysis and core swelling are complete by T_{lysis}. Spores that are exposed to germinant molecules

for a short time interval can be committed to germination at time T_c even when the germinants are

removed and further binding to GRs is blocked (28-31); the slow leakage of 15-25% of CaDPA

begins at or close to T_c and this slow release ends at T_{lag}, when rapid CaDPA release begins (24). T_{lag}

values are decreased by spore heat activation, higher GR levels and higher levels of GR-dependent

germinants up to saturation (19). Spores with ~4 fold higher SpoVA copy numbers have lower T_{lag}

values in GR germination than those for wild type spores (15). The times needed for rapid release of

all CaDPA remaining at T_{lag} ($\Delta T_{release} = T_{release} - T_{lag}$) (0.5-3 min), and for cortex hydrolysis and core

swelling ($\Delta T_{\text{lysis}} = T_{\text{lysis}} - T_{\text{release}}$) (10-20 min) are generally shorter than T_{lag}. GR levels and nutrient

germinant concentrations do not appreciably affect $\Delta T_{\text{release}}$ or ΔT_{lysis} values. However, loss of the CwlJ

CLE, but not SleB, markedly decreases rates of CaDPA release during spore germination.

Notably, spores were found to exhibit memory of prior germinant exposures (30,32,33). Thus spores

given a brief exposure to a germinant that activates the GerA or GerB* GRs responded more readily

to a 2$^{\text{nd}}$ short germinant pulse. However, the response to the 1$^{\text{st}}$ germinant pulse diminished when the

time interval in between pulses was increased (30, 32), indicating that there is memory in the

germination system. Furthermore, nonnutrient germinants such as GR-independent CaDPA or

dodecylamine, which do not require GRs, exhibit memory either alone or in combination with nutrient

germinants (32), suggesting that the SpoVA channel proteins are the most likely candidates for

memory storage while GRs are the candidates for memory generation. It has been proposed that spore

memory is primarily stored in metastable states of SpoVA proteins (32), and/or that the spore's

electrochemical potential plays a role in enabling this memory (34). However, the involvement of

electrochemical potential in the memory of germinant exposure appears unlikely as it was found that

Thioflavin-T cannot report on electrochemical potential of germinating spores (33), but mechanisms

underlying spore germination memory are not clear.

The pioneering work on mathematically modeling germination kinetics and heterogeneity was

conducted by Woese et al. in 1968 (35). Spore germination was assumed to result from the

accumulation of an unknown substance P, with its production rate being proportional to the number n

of activated germination "enzymes". By assuming that a threshold amount of P was required for

initiating germination, it was predicted that time-to-germination of a spore with n active enzymes is

proportional to 1/n. Furthermore, by assuming that the fraction of spores with n active enzymes in a

population follows a Poisson distribution, this simple model explained spore heterogeneity and

germination kinetics. Recently, Woese's work was extended to a comprehensive mathematical model

to describe heterogeneity, commitment, memory, and kinetic CaDPA release of spore germination

(36), in which each spore in a population possesses m functional GRs following a statistical

distribution, the binding of nutrient germinant activates a number of GRs and produces a time-

dependent germination trigger signal. It was assumed that when the trigger signal is above a critical

value, the spore is committed to germinate. This commitment results in the opening of the SpoVA

channel, leading to the release of CaDPA and full spore germination (36). This model's predictions

fitted very well with the experimental data of heterogeneity, commitment, memory, and CaDPA

release at single spore level. However, the biochemical detail for the trigger signal or germination

substance P remains unclear.

In the current work, we have attempted to develop a mathematical model to identify a minimal

reconstructed molecular network that is crucial for germination kinetics and memory. We consider a

minimal set of GRs in two states – inactive and active, and a SpoVA channel in three states - closed

inactive, closed active and open. When the number of SpoVA proteins in the open active state reaches

a threshold level (e.g. 50 copies/spore used in this paper), the channel is largely opened for rapid

CaDPA release. Numerical simulation reveals that memory can reside with GRs and/or be stored in

the CaDPA channel complex, which is consistent with the previous proposal that spore memory is

stored in metastable states of SpoVA proteins (32). Obviously, this model must be able to predict both

the heterogeneity of T_{lag} and the percentages of spore germination in 1st and 2nd germinant pulses. In

addition, with computer simulations we can identify network topologies that exhibit minimal

deviation between predicted and experimental data, and reveal reaction dynamics of molecular

processes between GR activation and rapid CaDPA release. Finally, analysis of the optimized model

could shed light on the question of how the network's molecules and their interactions determine its

biological function.

Results

Spore germination model.

In our modelling strategy, we follow a top-down approach. First we develop an overall mathematical

model and then we dissect the mathematics into a set of mass action kinetics based differential

equations based on the components of the germinosome and the SpoVA channel which must

somehow interact. As a boundary condition we opted for the most parsimonious number of building

blocks, and that the model must be able to recreate published experimental spore memory data (32).

One common premise for commitment in molecular networks is a bistable switch (37). In such

bistable systems two stable states coexist, and thus spores are dormant or are committed to germinate

(38). Upon activation, the system is irreversibly committed to one state. A condition for bistability is

non-linear dynamics involving positive feedback in which a molecular species upregulates its

production or autoactivates its function, *e.g.* channel opening, a mechanism to filter out noise and a

mechanism to prevent numeric explosion of the upregulated or activated species (38,39). A simple

deterministic molecular kinetic model for the autoactivation and deactivation rates of the state

variable *X* commonly used to model bistability is given in Eq (1) (40,41).

$$\frac{dX}{dt} = P + k_s \frac{X^n}{X^n + \theta^n} - k_d X \tag{1}$$

In this equation, *X* denotes the concentration [molecules/spore] of a crucial state of a germination

protein, k_s is its maximum production rate, *n* represents cooperativity of variable *X*, θ^n sets the value

at which the rate of production of *X* is half its maximum value, k_d is the decay rate of protein *X* and *P*

is a basal production rate (40-42). Positive feedback is introduced by cooperative action of the system

variable *X* for *n* >1. The multistabilty of Eq (1) depends on the settings of the various parameters.

There are some settings in which *X* gradually increases upon an inductive stimulus and there is a

parameter subspace where *X* exhibits a bistable response. In the latter case, there are two stable states,

one at low *X* and one at high *X*, and an unstable state in between. Fluctuations of *X* mediated by *P*

around the lower stable state (state 1) but not progressing to the metastable state fall back to the lower

stable state. Fluctuations of *X* through *P* surpassing the unstable state are auto-activated and increase

to the stable state 2 at high *X*. Fluctuations in *X* above the high stable state fall back to stable state 2,

thereby preventing a runaway of variable *X* to infinity. Eq (1) is thus an autocatalytic off-on switch

with parameter subspace for bistability and is the core of our model.

Starting from this equation we designed an input section, signal transduction and finally we describe

SpoVA channel opening kinetics to estimate T_{lag} times in germination. The molecular wiring of the

reconstructed molecular network of the model is given in Fig 1. In our model, we do not differentiate

between GerA, GerB and GerK GRs, and thus do not differentiate between germinants. GRs can be in

an inactive R_{i} state or an active R_{a} state, and germinant S binding converts an R_i GR to R_a with a rate

constant k_1. Low copy protein numbers in individual cells in bacterial populations usually vary

according to a gamma distribution (43). Measurements of levels of GFP-labelled GRs in spores have

shown that GR copy numbers also vary according to a gamma distribution function (43,44). Thus, the

distribution of total levels of R_{i} in spores in the absence of germinant is given by the probability

density (gamma) function

$$f(R_i; a, b) = R_i^{a-1} \frac{e^{-R_i/b}}{b^a \Gamma(a)} \tag{2}$$

208   in which $\Gamma(a) = (a-1)!$, a is the shape parameter and b is an inverse scale parameter. The

209   distribution mean $E(R_{\text{i}}) = ab$ and the variance $Var(R_{\text{i}})$ of the distributions is ab^2. The skewness of the

210   distribution is equal to $2/a^{1/2}$. The skewness depends only on the shape parameter a and approaches a

211   normal distribution when approximately $a > 10$.

213   Fig 1   Architecture of a *Bacillus subtilis* germinosome that exhibits spore germination

214   memory. The molecular network is composed of GR proteins that are activated by a

215   germinant, and channel proteins are in a closed, activated closed and an open state

216   and undergo mutual interactions.

Germinant binding in our model is reversible and dissociation takes place with rate constant k_{-1}. Eq

(3) and Eq (4) describe rates of formation of active GR molecules R_a and rates of depletion of the

inactive GR population R_i. In contrast to GRs which we propose exist in only two states, R_i and R_a, we

need a SpoVA channel that can adopt three states. While there is no experimental proof for the latter

assumption, a model with two SpoVA states does not reproduce experimental germination curves

(data not shown). The SpoVA channel C_c is closed and inactive, the channel C_{ca} is closed and in an

activated state, or the channel C_o is open and in an activated state. Interaction of activated GRs R_a with

closed SpoVA channels C_c activates the latter (Eq (5)). We assume a direct interaction, although there

is no conclusive experimental data about signaling from activated GRs to the SpoVA channel.

However, yeast two hybrid experiments suggest that there is interaction between *B. subtilis* GerA

proteins and several SpoVA proteins (45). The function of GerD is ignored in the model. A premise

for commitment in molecular networks is a bistable switch with delay. Bistability is introduced here

through intersection of a production function comprising a Hill function with $n > 1$ and a linear decay

function in Eq (7). The production function describes the rate of opening of SpoVA channels.

Cooperativity of open SpoVA channels also contributes to the opening rate. This means that open

SpoVA channels (C_o) have multiple mutual interactions which enhance the opening rate of closed

channels. Cooperativity of SpoVA channel gating is introduced through a Hill function with apparent

dissociation constant θ and coefficient n. The decay function describes closure of channels with

reaction rate $-k_5 C_o$. Channel closure is first order in the concentration of open channels. Upon closure

the closed channel falls back to the inactive ground state C_c.

$$\frac{dR_i}{dt} = -k_1 S R_i + k_{-1} R_a \tag{3}$$

$$\frac{dR_a}{dt} = k_1 S R_i - k_{-1} R_a \tag{4}$$

$$\frac{dC_c}{dt} = -k_2 R_a C_c - \frac{k_4 C_o^n}{C_o^n + \theta^n} + k_5 C_o \tag{5}$$

$$\frac{dC_{ca}}{dt} = k_2 R_a C_c - k_3 C_{ca} \tag{6}$$

$$\frac{dC_o}{dt} = k_3 C_{ca} + \frac{k_4 C_o^n}{C_o^n + \theta^n} - k_5 C_o \qquad (7)$$

In these equations, S is germinant concentration, R_i inactive GR, R_a active GR, C_c closed SpoVA

channel, C_{ca} activated closed SpoVA channel, and C_o open SpoVA channel. The units of the GR and

SpoVA variables are numbers of the various complexes/spore. Constants k_1, k_{-1}, k_2, k_3, k_4 and k_5 are

rate constants, n is a Hill coefficient, and θ^n sets the value at which the production rate in the second

term is half its maximum value. The final stage of the model is SpoVA channel opening. Whether

core CaDPA is exported by concentration differences and/or an electrochemical gradient of ions

inside and outside the inner membrane, CaDPA efflux will be proportional to the number of open

SpoVA channels, and we assume that T_{lag} is determined by a minimal number of open SpoVA

channels needed for rapid CaDPA release. It is known that CaDPA activates the CLE CwlJ and this

enzyme's action stimulates CaDPA efflux. However, CwlJ activation and its effect on the SpoVA

channel opening rate is not yet a part of the model, and we expect that this effect will become

significant only after T_{lag}. If there is a contribution of CwlJ activation to SpoVA opening than this will

lead to a minor shift of T_{lag}. In our computer model, we use a cycle of 500 numeric simulations to

solve the set of coupled ordinary differential equations. In each simulation, we predict the number of

active open SpoVA channels as a function of time and we determine for each simulation the time of

opening, T_{open}, of SpoVA channels. This time point corresponds with the time point at the beginning

of fast CaDPA release (28), and is related to T_{lag} through $T_{lag} = T_{open} - T_{block}$, where T_{block} is the time of

the end of a germinant pulse. We define T_{open} as the time point in a simulation where C_o exceeds 50

copies/spore. This number is arbitrarily selected as experimental data are lacking.

Modeling of spore memory of germinant stimuli

First, we reproduce with our mathematical model an earlier published percentage germination curve

showing spore memory for stimulation of wild-type *B. subtilis* spores with L-valine (32). Next, we

study the relationship between the mean $E(R_i)$ of the GR distribution and the percentage germination

upon the 1[st] and the 2[nd] germinant pulses. We then compare the predictions with experimental results

of the percentage germination of wild-type spores and mutant spores with elevated GR levels. Wild-

type spores can acquire memory upon short pulses of L-valine to induce germination as follows. After

the 1[st] pulse a fraction of the spores is committed to germinate and does so. From the subpopulation

that does not germinate after the 1[st] pulse, a fraction germinates upon a 2[nd] pulse and finally a fraction

does not germinate at all. Interestingly, spore populations can exhibit 2- to 3-fold more germination in

the 2[nd] germinant pulse following the 1[st] one. Apparently, there is memory in the molecular system for

external perturbations such as a germinant pulse that occurred before the 2[nd] pulse (32). Consequently,

we optimize model parameters to reproduce the spore germination curve for an experiment with two

short germinant pulses (32). In our simulation, we select two pulses of germinant exposure from $0 < t$

< 5 min and $30 < t < 35$ min. We set the germinant concentration to 3.5 mM. A percentage

germination curve in our definition is a cumulative distribution function of T_{open} to give rapid CaDPA

release. Spores that do not germinate have an infinite T_{open} and do not contribute to the distribution in

the time frame of the simulations. Heterogeneity of T_{open} and as result T_{lag} determines the shape of the

curve. Here, we introduce heterogeneity through a gamma distribution of GR copy numbers (Eq (2)).

For each simulation of a cycle of 500 we randomly select a GR copy number from a gamma

distribution, but keep SpoVA copy numbers constant. This means that $C_c + C_{ca} + C_o$ is constant in

each cycle of a set of simulations. This approach is justified by the experimental observation that

overexpression or deletion of GR's has no effect on levels of SpoVAD and by inference levels of all

proteins encoded by the heptacistronic *spoVA* operon (15). From Western blotting, it is known that on

average ~700 to ~1,100 of each GR are present in a spores' germinosome. The starting value of at

least SpoVAD is 6,500 as determined earlier by Western Blotting (20). By tuning of k_4, the SpoVA

copy number of activated and open channels in our simulations asymptotically approaches 6,500

copies/spore (*vide infra*). This copy number is also an arbitrary selection as we do not know by

experimentation the number of open SpoVA channels at the end of rapid CaDPA release. At the end

of the simulations we construct a predicted percentage germination curve. The experimental

germination curve was determined by monitoring germinating spores with DIC microscopy. However,

most of the intensity change seen in DIC microscopy is caused by rapid CaDPA release, and the

295 kinetic parameters for which averages were determined are T_{lag}, $T_{release}$ and $\Delta T_{release} = T_{release}\text{-}T_{lag}$

296 (28,32,46).

297 We start with varying the mean $E(R_i)$ of the GR distribution and keep the standard deviation $Var^{\frac{1}{2}}(R)$

298 fixed. The parameters of the randomly accessed gamma distribution are estimated at the end of a set

299 of 500 cycles from the fitted gamma distribution. A typical GR protein distribution is depicted in Fig

300 2A. The average of this GR copy number distribution is set to 1100 copies/spore and we selected

301 combinations of distribution parameters a and b with the condition that $Var^{\frac{1}{2}}(x) = a^{\frac{1}{2}}b = 220$. With

302 this setting the GR distribution tails to higher copy numbers, and a fraction of the spores germinate

303 after the 1st pulse and a fraction germinates after the 2nd germinant pulse (Fig 2A, B). Notably, with

304 the 1st pulse 15% of the spore population germinates, while after the 2nd pulse 68% of the remaining

305 spores germinate. Although the germinant concentration and pulse lengths are equal for both pulses,

306 germination after the 2nd pulse is approximately four-fold higher than after the 1st pulse. Thus, the

307 model reproduces published experimental data (32). Simulations for other $E(R_i)$ are compiled in S1

308 Figures.

310 Fig 2    Simulation of spore germination memory

(A)   Gamma distribution of GR copy numbers used for simulation of the percentage germination efficiency curve (Fig.2b) with $E(R_i) = 1100$ at t is 0 min. The red line is the fitted distribution. The GR estimated mean is 1105 molecules/spore, and the estimated standard deviation is 219 molecules/spore; the parameters of the fitted gamma distribution are: a is 25.4491 [22.5002, 28.7844] and b is 43.4004 [38.3247, 49.1484]; Right panel: Percentage germination of spores in a two pulse germination experiment.

(B)   Percentage germination in a two pulse germination experiment. After the 1st pulse the curve levels off to an asymptote at 15%. After the 2nd pulse the curve approaches 68%. Two pulses of 3.5 mM germinant for receptor R_i activation are administered from $0 < t < 5$ and $30 < t < 35$ min. At zero time $R_a = 0$ copies/spore; $C_c = 6500$ copies/spore. $C_a = C_o = 0$ copies/spore; $k_1 = 14e\text{-}3$; $k_{-1} = 100e\text{-}3$; $k_2 = 25e\text{-}7$; $k_3 = 60e\text{-}3$; $k_4 = 195e2$; $k_5 = 305e\text{-}2$; $n = 3$; $\theta = 20$.

We repeated a double pulse germination experiment with *B. subtilis* wild-type spores and spores of an isogenic strain that overexpresses the GerA GR ~8-fold as determined by Western Blotting, with an experimental error of 25% (7). The experimental percentage germination curves are depicted in Fig 3A, B. We simulated the curves with our model (Fig 3C, D). Compared to germination of wild-type spores we had to adjust reaction rate constants for a 2 min pulse for wild-type spores that overexpress GerA with 10 mM L-valine as germinant. A broad GR copy number distribution, with $E(R_i)$ at 1100, a is 1 and b is 1100, reproduces approximately 4% germination upon the first pulse and approximately 6% germination at the second pulse (Fig 3C). Next, we overexpress the GerA GR by increasing $E(R_i)$ and keep the reaction rate constants similar. In this approach, we put forward the constraint that overexpression of GR GerA from a more active promotor does not affect the rate constants for

binding and dissociation of the germinant to the receptor. A second assumption is that expression of

GerA from another promotor affects the parameters of the gamma distribution. It is known from

modelling of burst like stochastic gene expression that the rate at which the gene transitions from the

inactive state to the active state and vice versa determines the momenta of the mRNA copy number

distribution and consequently the protein copy number distribution (47). Our modelling results are in

accord with an earlier prediction using a stochastic gene expression model that a high burst frequency

produces a narrow mRNA copy number distribution and a low burst frequency a broad mRNA

distribution (47). However, experimental data for the distribution of copy numbers of an

overexpressed GFP-labelled GerA in individual spores are lacking. With $E(R_i)$ at 3350 a as 67 and b

as 50 we reproduce 3% germination at the first pulse and 64% germination at the second pulse (Fig

3D). In this case, we had to adjust the variance of the GR distribution to reproduce the percentage

germination curve. For $E(R_i)$ at 3350 the overexpression is approximately 3.3-fold. For further

increase of $E(R_i)$ we have to diminish k_1 to keep the second pulse at 63% without surpassing the 3%

germination at the first pulse. These results validate simulations of the relation between GR protein

distributions and percentage germination curves in a double pulsed memory experiment. Interestingly,

the model predicts constant $\Delta T_{release}$ times and that $\Delta T_{release}$ is independent of T_{lag}, corroborating

observations from earlier experimental work (28). Next, we probed for the effect of varying the

concentration of germinants used during the pulses, from 2 mM to 5 mM (S2 Figures). At 2 mM

germinant no germination is predicted with the 1[st] pulse and approximately 1% after the 2[nd] pulse.

This latter percentage increases to 12% at 2.5 mM germinant, but still with no germination after the 1[st]

pulse. This implies that the 1[st] pulse primes 12% of the spore population for germination after the 2[nd]

pulse. For higher germinant concentrations, significant germination with the 1[st] pulse as well as

substantial memory effects are predicted, *e.g.* S2 Fig 2C. The influence of the standard deviation of

the distribution is further highlighted in S3 Figures. There is an obvious relationship between the

variance of the distribution and the fraction of spores germinating after the 1[st] and 2[nd] pulses (S3

Figures). As the variance of the distribution increases and tails to higher GR copy numbers, more

spores germinate with the 1[st] germinant pulse.

Fig 3   A double pulse germination experiment with *B. subtilis* wild-type spores and spores

366   of an isogenic strain that overexpresses the GerA GR.

367   (A)   Wild-type *B. subtilis* spores (PS533) were given two 2-min 10 mM L-valine

368   (L-val) germinant pulses separated by 30 min.

369   (B)   Spores of *B. subtilis* PS3476 (GerA GR over-expressed ~ 8-fold) were given

370   two 2-min 10 mM L-valine germinant pulses separated by 30 min. Gray bars

371   above the horizontal axis indicate pulse durations.

372   (C)   Simulated percentage germination of WT spores in a two pulse germination

373   experiment. The estimated distribution mean is 1118 and the standard

374   deviation is 1064 with *a* is 1.10476 [0.989585, 1.23334] and *b* is 1012.19

375   [881.615, 1162.1]. At zero time $R_a = 0$ copies/spore; $C_c = 6500$ copies/spore.

376                $C_a = C_o = 0$ copies/spore; $k_1 = 350e\text{-}5$; $k_{-1} = 100e\text{-}3$; $k_2 = 25e\text{-}7$; $k_3 = 60e\text{-}3$; k_4

377                $= 195e2$; $k_5 = 305e\text{-}2$; $n = 3$; $\theta = 20$.

(D)   Simulated percentage germination of overexpressed GR spores in a two pulse

germination experiment. The estimated distribution mean is 3367 and the

standard deviation is 399 with a is 71.2149 [62.9306, 80.5898] and b is

47.2855 [41.7667, 53.5336]. At zero time $R_a = 0$ copies/spore; $C_c = 6500$

copies/spore. $C_a = C_o = 0$ copies/spore; $k_1 = 350e\text{-}5$; $k_{-1} = 100e\text{-}3$; $k_2 = 25e\text{-}7$;

$k_3 = 60e\text{-}3$; $k_4 = 195e2$; $k_5 = 305e\text{-}2$; $n = 3$; $\theta = 20$.

Dynamics of SpoVA channel opening

The kinetics of SpoVA channel opening are studied in more detail for GR copy numbers ranging from

600 to 1,800 molecules/spore. The increment of GRs is 50 copies per cycle of numeric evaluation of

the set of ordinary differential equations (ODE's) in a simulation. The number of GRs is defined by R_i

and at time point t is 0 seconds. The results C_o and the copy number time profiles of the state variables

R_i, R_a, C_c, and C_{ca} are given in Fig 4. There are three subpopulations in the uniform GR distribution.

With 600 to 950 GRs/spore, there is no germination, with 1000 to 1350 GRs/spore there is

germination after the 2nd pulse and with 1400 to 1800 GRs/spore there is germination after the 1st and

2nd pulses. Fig 4 shows that only in a fraction of the spore population are GRs sufficiently activated by

germinant binding to lead to commitment to germinate. We can tune this fraction by variation of the

rate constants k_1 and k_{-1} for binding of S to GR and dissociation thereof. This tuning also affects the

number of spores opening SpoVA channels during the 1st and 2nd pulses. With the parameter settings

of our simulation only a few SpoVA channels are activated by interaction with GRs. Because of

different copy numbers between GRs and channels there cannot be a one to one stoichiometry

between GR activation and channel opening. Approximately 1,100 GerA GRs are in the

germinosome, while ~6,500 SpoVA channels are dispersed over the IM. The surface area of the spore

IM can be approximated by an ellipsoid. The surface area of an ellipsoid (A_{im}) is calculated as $A_{im} =$

$2\pi(s^2+ls)$, where l is the length of the long axis and s of the short axis (48). The long axis l and short

axis s of the IM are estimated by electron microscopy and l is 837 nm and s is 554 nm for a typical

spore (49). These are obtained by subtraction of coat and cortex layers' thickness from the spore long

axis and short axis (49). The surface area of the IM is $A_{im} = \sim 4{,}835{,}800$ nm^2, and on average there

will be one SpoVA molecule per 744 nm^2 of IM. Next, we calculate the area of the germinosome. As

input we use the normalized average distribution of the fluorescence signal along the long axis of a

spores. The full width at half max intensity for GerKB-GFP signal is 300 nm (9). Assuming a planar

circle on the surface of the IM for the contour of the germinosome its surface $A_{ger} = \pi r^2 = 502{,}655$

nm^2. Assuming a homogeneous distribution of 744 nm^2 per SpoVAE then the germinosome contains

at most 676 SpoVA molecules. In a stoichiometry where one GR activates one channel maximally,

1,100 channels can be activated. Thus, there are $\sim 5{,}800$ SpoVA channels that cannot directly interact

with a GR. Thus, SpoVA channel cooperativity is introduced in the model to relay the signal from

GRs over the entire IM surface to open more than approximately 676 channels. In our model, closed

SpoVA channels are directly opened by activated and open SpoVA channels. Immediately after the 1st

germinant pulse the number of open SpoVA channels gradually increases. This slow increase is the

gross difference between opening by activated SpoVA channels through $k_4 C_a$ and closing of SpoVA

channels through decay $-k_5 C_o$ (Eq (7)). Beyond 10 min sufficient channel opening takes place to force

cooperative opening of the dark blue subpopulation through the Hill function term in Eq (5). Channel

opening is autocatalytically driven, all SpoVA channels open, and hence the C_o curve approaches

6,500 molecules. The SpoVA channels from spores that germinate after the 2nd pulse are partially

activated by the 1st pulse but remain below a critical copy number needed to force an autocatalytic

opening mediated by the Hill function in Equation 6. However, the extent of channel opening does not

fully return to the baseline. A 2nd pulse at $30 < t < 35$ min contributes additionally to channel opening

and drives through cooperative action channel opening to completion at 6,500 copies/spore. As a

result, channels open that are not opened by the 1st germinant pulse. Thus, the system exhibits

memory for an earlier germination stimulus. Finally, the curve levels off to a plateau at 6500 open

SpoVA channels. The level of the plateau of open SpoVA channels is mainly determined by k_4 and k_5.

A third fraction does not open at all, as the 2nd germinant pulse is insufficient to trigger autocatalytic

SpoVA opening and the number of open channels decays to the baseline.

Fig 4   The copy numbers of R_i, R_a, C_c, C_a, C_o and an extension of C_o as a function of time.

433   Each trace represents a state variable profile for a variable GR R_i copy number. The

434   starting GR values are $600 \leq R_i \leq 1800$ copies/spore and the increment is $\Delta R_i = 50$

435   copies/spore. The yellow spore population ($600 \leq R_i \leq 950$) does not germinate, the

436   red one ($1000 \leq R_i \leq 1350$) germinates at the 2[nd] pulse and the blue ones ($1400 \leq R_i \leq$

437   1600) germinate at the first pulse. Two pulses of 3.5 mM germinant for R_i activation

438   are administered from $0 < t < 5$ and $30 < t < 35$ min. At time zero, $R_a = 0$

439   copies/spore; $C_c = 6500$ copies/spore. $C_a = C_o = 0$ copies/spore; $k_1 = 14e\text{-}3$; $k_{-1} =$

440              $100e\text{-}3$; $k_2 = 25e\text{-}7$; $k_3 = 60e\text{-}3$; $k_4 = 195e2$; $k_5 = 305e\text{-}2$; $n = 3$; $\theta = 20$.

442   **Discussion**

443 Spore germination experiments with sequential germinant stimuli separated in time are very suitable

444 to construct and optimize a kinetic germination model. The shape of the percentage germination

445 curves with multiple inflection points and asymptotes enables a more robust fitting of model

446 parameters than a single stimulus sigma-shaped germination response experiment. The germination

447 model based on the minimal molecular network as presented in Fig 1 and described by differential Eq

448 (3-7) reproduces percentage germination curves in double pulse memory experiments (32) and newly

449 acquired data (Fig 3A, B). Upon germinant exposure in two 5 min pulses separated by thirty min, two

450 subpopulations of the whole population germinate. The second one exhibits a memory effect, in that

451 the amounts of spores that germinate after the 2nd pulse are larger than those germinating after the first

452 pulse. Memory has been attributed to a conformational change in some germination protein, that

453 activates this protein for germination, although the activated protein can also decay to the inactivate

454 ground state. When the activated protein is not fully decayed, then a 2nd germinant pulse leads to a

455 larger population of activated proteins and a higher percentage of germination after the 2nd pulse (32).

456 The reconstructed network shows that a minimal set of an inactive and active GR and a transport

457 channel in three states - closed inactive, closed active and open is needed to reproduce memory. Thus,

458 memory can reside with GRs and/or be stored in the CaDPA channel complex. This is consistent with

459 generation of memory in spores that lack all GRs when germination is triggered by direct or indirect

460 stimulation of CaDPA release by exogenous CaDPA itself that activates CwlJ, or by dodecylamine

461 that may cause the SpoVA channels to open (32).

462 Memory is stored in an activated GR R_a, although this can decay to the inactive state R_i. As is seen in

463 Fig 4, GR inactivation does not fully return to the baseline before the 2nd germinant pulse and thus GR

464 inactivation characteristics contribute to memory. The fraction of activated GRs and its relaxation to

465 an inactive state is determined in our model by the ratio k_1/k_{-1}, although we do not know this ratio

466 from experimental data. However, memory in our model is not uniquely confined to GR activation

467 and inactivation kinetics, as there is a model parameter setting with $k_{-1} = 225.10^{-3}$ and $k_2 = 45.10^{-7}$

468 where R_a returns to baseline before the 2nd germinant pulse and the network still shows memory for

469 germination without adjusting the other rate constants. In this case memory is introduced in the

network by the interplay between activation of closed channels and their rate of opening to C_o. The

spore channel activation rate is proportional to $k_2 R_a$ and the number of closed inactive SpoVA

channels C_c (Equation 6), and the rate of opening is proportional to $k_3 C_a$. We assume that the number

of activated GRs at 3.5 mM is approximately 17% and we have optimized k_2 accordingly. This needs

further experimentation to dissect $k_2 R_a$ into k_2 and R_a. Rate constant k_3 has an effect on T_{lag} because it

feeds the rate of open channel formation (Eq (7)). The Hill coefficient n, θ together with k_4 and k_5

determine the bistability of the model needed to irreversibly open CaDPA channels. In this simulation

n is 3 pointing to cooperativity effects of open activated SpoVA channels (Eq (7)). Furthermore, rate

constants k_4 and k_5 in combination with θ determine the number of open SpoVA channels at the end-

point of simulations. This number is not known by experimentation and is arbitrarily set.

The time to the onset of rapid CaDPA release T_{lag} is heterogeneous and shows a probability

distribution. We introduce germination time heterogeneity through the GR distribution in spore

population. In a cycle of computer simulations, we keep the rate constants fixed. Rate constants k_1 and

k_{-1} are on-rate and off-rate constants and determined by the 3-D structure of the GR proteins. As the

primary sequence of the GR proteins is not variable within an isogenic strain, we expect that a k_1 and

k_{-1} probability distribution does not contribute to heterogeneity of T_{lag} at the population level. At

present, our model is a minimal one, and we do not differentiate between actions of GerA, GerB and

GerK GRs. Consequently, we do not integrate the signal from different GRs and germinants. We also

do not know the fraction of SpoVA channels in an inactive state that is activated by a population of

activated GRs. The network in our model is capable of signal amplification, as with current parameter

settings approximately 1-27 closed SpoVA channels are activated (Fig 4). This is sufficient to

irreversibly open the entire channel population through cooperative opening which is needed to open

channels outside the perimeter of the germinosome. Another benefit from the network wherein only

fractions of molecular populations are active is that it dampens fluctuations in germinant exposure.

This ensures that there is a threshold for GR activation that will prevent unwanted early germination.

The model generated in this work has been optimized to reproduce memory of germinant exposures.

Other combinations of reaction rate constants will likely also give kinetics that show memory, but the

molecular wiring and the kinetic differential equations will likely remain identical. However, further,

experimentation is needed to narrow down the parameter space. In particular, the contributions of GR

activation and deactivation, and SpoVA channel activation, deactivation and opening should be

studied. The endpoint of germination in our model is the rapid opening of SpoVA channels. In a

future extension of the model, the effects of the CLEs SleB and CwlJ should also be included. In

addition, as noted above GR-independent germination by CaDPA or dodecylamine also show

memory, presumably through effects on the SpoVA channel alone, and this can also be introduced

into the model. The model in its current state will be used in future research to reduce the number of

experimental variables in an iterative cycle of prediction of the dynamics of a hypothetical molecular

network, the design of an experiment, comparison with the outcome, and further tuning of the model

until agreement between theoretical predictions and experimental data is reached.

Materials and methods

Simulations were carried out using the MATLAB package of software (the Mathworks, Natick,

USA). M-files with source code for simulation of percentage germination curves and copy number

time profile plots are given in S Matlab m-file one and S Matlab m-file two.

Phase contrast microscopy.

Spores (1 µl; ~10^8 spores/ml in water) were spread on the surface of a microscope coverslip that was

then dried in a vacuum desiccator for 5 min. Coverslips were suspended in water and then mounted on

and sealed to a microscope sample holder kept at a constant temperature of 37°C. A phase contrast

microscope was used to record the images of ~500 individual spores at a rate of 15 s per frame.

Spores that contain DPA appear as bright and spores that do not contain DPA appear as dark.

Percentage of spores containing DPA was defined as the number of bright spores divided by the total

number of spores.

Spore germination with germinant pulses.

B. subtilis spores of strains PS533 (wild-type) and PS3476 (overexpresses the GerA GR ~ 8-fold)

were prepared, purified and stored as described previously (14). The purified spores were free (>98%)

of sporulating or growing cells or germinated spores as observed by phase contrast microscopy. *B.*

subtilis spores were heat activated prior to nutrient germination by incubation in water at 70°C for 30

min and then by cooling on ice for at least 15 min. *B. subtilis* spores attached on the surface of a glass

coverslip were germinated in 25 mM K-HEPES buffer (pH 7.4) with a pulse of a germinant (10 mM

L-valine) for 2 or 4 min before the germinant was removed and spores were rinsed 5 times with 25

mM HEPES buffer (pH 7.4) using vacuum pump suction (32). The rinsed spores were then incubated

at 37°C in 25 mM K-HEPES buffer with a pH of 7.4 unless noted otherwise. After various incubation

times, the spores were given a 2nd exposure to germinants for various times at 37°C in 25 mM K-

HEPES buffer (pH 7.4) followed by germinant removal, rinsing by vacuum pump suction, and further

incubation at 37°C in 25 mM K-HEPES buffer (pH 7.4).

Acknowledgements

CdK, LdK and SB acknowledge the University of Amsterdam for funding of their research.

Author contributions

CdK developed the model and performed computational analyses with input from LdK and SB. XL

and YQL conducted experiments. PS provided spores of *Bacillus subtilis* strains. CdK wrote the

manuscript with input from LdK, SB, YQL and PS. YQL, JL and PS designed the germination

experiments.

Conflict of interest

The authors declare that they have no conflict of interest.

References

(1) Setlow P and Johnson EA. (2019) Spores and their significance. In *Food Microbiology,*
Fundamentals and Frontiers, 5th edition (M.P. Doyle and R. Buchanan, ed.), ASM Press,
Washington, DC, p. 23-64.

(2) Setlow P, Christie G. If ATP and macromolecular synthesis are needed for dormant spores of
Bacillota species to trigger spore germinatuon, where do the energy and precursors come from. J Appl
Microbiol 2025 Jul: 136(7); lxaf177.

(3) Christie G, Setlow P. 2021. What's new and notable in bacterial spore killing. World J. Microbiol.
Biotechnol. **37**:144.

(4). Setlow P. Germination of spores of *Bacillus species*: what we know and do not know. J Bacteriol
2014 Apr;196(7):1297-1305.

(5) Christie G, Setlow P, Bacillus spore germination: Knowns, unknowns and what we need to learn.
Cell Signalling 2014 Oct; 74: 109729

(6) Setlow P. Spore germination. Curr Opinion Microbiol 2003 Dec;6(6):550-556. (7) Ross C, Abel-
Santos E. The Ger receptor family from sporulating bacteria. Curr Issues Mol Biol 2010;12(3):147-
158.

(8) Amon JD, Artzi L, Rudner DZ. Genetic evidence for signal transduction within the *Bacillus*
subtilis GerA germinant receptor.J. Bacteriol 2022 Feb 15:204:e0047021

(9) Artzi L, Alon A, Brock KP, Green AG et al. Dormant spores sense anino acids through the B
subunits of their germinant receptors. Nat Commun 2021:12:6842.

(10) Gao Y, Amon JD, Ramirez-Guadiana FH, Brock KP *et al*. Bacterial spores germinant receptors
are nutrient gated ion channels. Science 2023:380 (6643):3870391.

(11) Eichenberger P Spore germination. Two ion channels better than one. Genes Dev 2024; 38:1-3.

(12) Gao Y, Amon JD, Brogan AP, Artzi L et al. SpoFV and Fig P assembe into oligomeric ion
channels that enhance spore germination. Genes Dev 2014: 38:31-45.

(13) Igarashi T, Setlow B, Paidhungat M, Setlow P. Effects of a *gerF* (*lgt*) mutation on the
germination of spores of *Bacillus subtilis*. J Bacteriol 2004 May;186(10):2984-2991.

14) Stewart KA, Setlow P. Numbers of individual nutrient germinant receptors and other germination
proteins in spores of *Bacillus subtilis*. J Bacteriol 2013 Aug;195(16):3575-3582.

(15) Stewart KA, Yi X, Ghosh S, Setlow P. Germination protein levels and rates of germination of
spores of *Bacillus subtilis* with overexpressed or deleted genes encoding germination proteins. J
Bacteriol 2012 Jun;194(12):3156-3164.

(16) Griffiths KK, Zhang J, Cowan AE, Yu J, Setlow P. Germination proteins in the inner membrane
of dormant *Bacillus subtilis* spores colocalize in a discrete cluster. Mol Microbiol 2011
Aug;81(4):1061-1077.

(17) Wang Y, de Boer R, Vischer N, Haastrecht N *et al*. Visualization of germination proteins in
putative *Bacillus cereus* germinosomes. Int J Mol Sci 2020; 21(15):5198.

(18) Li Y, Jin K, Ghosh S, Devarakonda P, Carlson K *et al.* Structural and functional analysis of the
GerD spore germination protein of *Bacillus* species. J Mol Biol 2014 May 1;426(9):1995-2008.

(19) Perez-Valdespino A, Li Y, Setlow B, Ghosh S, Pan D, Korza G, et al. Function of the SpoVAEa
and SpoVAF proteins of *Bacillus subtilis* spores. J Bacteriol 2014 Jun;196(11):2077-2088.

(20) Vepachedu VR, Setlow P. Localization of SpoVAD to the inner membrane of spores of *Bacillus*
subtilis. J Bacteriol 2005 Aug;187(16):5677-5682.

(21) Velásquez J, Schuurman-Wolters G, Birkner JP, Abee T, Poolman B. *Bacillus subtilis* spore
protein SpoVAC functions as a mechanosensitive channel. Mol Microbiol 2014 May;92(4):813-823.

(22) Setlow B, Cowan A, Setlow P. Germination of spores of *Bacillus subtilis* with dodecylamine. J
Appl Microbiol 2003;95(3):637-648.

(23) Gao Y, Barajas-Ornelas DRC, Amon JD, Ramirez-Guadiana FH *et al*. The SpoVA membrane
complex is required for dipicolinic acid import during sporulation and export during germination.
Genes Dev 2022; 36:634-646.

(24)Wang S, Setlow P, Li Y-Q. Slow leakage of Ca-dipicolinic acid from individual *Bacillus* spores
during initiation of spore germination. J Bacteriol 2015; 197:1095-1103.

(25) Sanchez-Salas J-L, Setlow B, Zhang P, Li YQ *et al*. Maturation of released spores is necessary
for acquisition of full spore heat resistance during *Bacillus subtilis* sporulation. *Appl. Environ.*
Microbiol. **2011; 77:**6746-6754.

(26). Abhyankar W, Pandey R, Ter Beek A, Brul S *et al*. Reinforcement of *Bacillus subtilis* spores by
cross-linking of outer coat proteins during maturation. *Food Microbiol.* 2015; **45(**Pt A):54-62. doi:
10.1016/j.fm.2014.03.007.

(27) Zhang P, Thomas S, Li YQ, Setlow P. Effects of cortex peptidoglycan structure and cortex
hydrolysis on the kinetics of Ca(2+)-dipicolinic acid release during *Bacillus subtilis* spore
germination. J Bacteriol 2012 Feb;194(3):646-652.

(28) Zhang P, Garner W, Yi X, Yu J, Li YQ, Setlow P. Factors affecting variability in time between
addition of nutrient germinants and rapid dipicolinic acid release during germination of spores of
Bacillus species. J Bacteriol 2010 Jul;192(14):3608-3619.

(29) Yi X, Setlow P. Studies of the commitment step in the germination of spores of *Bacillus* species.
J Bacteriol 2010 Jul;192(13):3424-3433.

(30) Zhang P, Liang J, Yi X, Setlow P, Li YQ. Monitoring of commitment, blocking, and continuation
of nutrient germination of individual *Bacillus subtilis* spores. J Bacteriol 2014 Jul;196(13):2443-2454.

(31) Chen D, Huang SS, Li YQ. Real-time detection of kinetic germination and heterogeneity of
single *Bacillus* spores by laser tweezers Raman spectroscopy. Anal Chem 2006 Oct 1;78(19):6936-
6941.

(32) Wang S, Faeder JR, Setlow P, Li YQ. Memory of germinant stimuli in bacterial spores. MBio
2015 Nov 24;6(6):e01859-15.

(33) Li Y-Q, He L, Aryal M, Wicander J, Korza G, Setlow P. Thioflavin-T does not report on
electrochemical potential and memory of dormant or germinating bacterial spores. MBio 2023 Oct 13;
14(5):e02220-23.

(34) Kikuchi K, Galera-Laporta L, Weatherwax C, Lam JY, Moon EC, Theodorakis EA, Garcia-
Ojalvo J, Süel GM. Electrochemical potential enables dormant spores to integrate environmental
signals. Science 2022; 378:43–49.

(35) Woese CR, Vary JC, Halvorson HO. A kinetic model for bacterial spore germination. Proc Natl
Acad Sci U S A 1968; 59:869–875.

(36) Li W, Mednick W, Setlow P, Li Y-Q, Modeling heterogeneity, commitment, and memory of
bacterial spore germination, MBio 2025 Apr 2; 16(5):e00596-25.

(37) Xiong W, Ferrell JE,Jr. A positive-feedback-based bistable 'memory module' that governs a cell
fate decision. Nature 2003 Nov 27;426(6965):460-465.

(38) Smits WK, Kuipers OP, Veening JW. Phenotypic variation in bacteria: the role of feedback
regulation. Nat Rev Microbiol 2006 Apr;4(4):259-271.

(39) Tiwari A, Ray JC, Narula J, Igoshin OA. Bistable responses in bacterial genetic networks:
designs and dynamical consequences. Math Biosci 2011 May;231(1):76-89.

(40) Frigola D, Casanellas L, Sancho JM, Ibanes M. Asymmetric stochastic switching driven by
intrinsic molecular noise. PLoS One 2012;7(2):e31407.

(41) Tyson JJ, Chen KC, Novak B. Sniffers, buzzers, toggles and blinkers: dynamics of regulatory
and signaling pathways in the cell. Curr Opin Cell Biol 2003 Apr;15(2):221-231.

(42) Becskei A, Seraphin B, Serrano L. Positive feedback in eukaryotic gene networks: cell
differentiation by graded to binary response conversion. EMBO J 2001 May 15;20(10):2528-2535.

(43) Friedman N, Cai L, Xie XS. Linking stochastic dynamics to population distribution: an analytical
framework of gene expression. Phys Rev Lett 2006 Oct 20;97(16):168302.

(44) Zhang JQ, Griffiths KK, Cowan A, Setlow P, Yu J. Expression level of *Bacillus subtilis*
germinant receptors determines the average rate but not the heterogeneity of spore germination. J
Bacteriol 2013 Apr;195(8):1735-1740.

(45) Vepachedu VR, Setlow P. Analysis of interactions between nutrient germinant receptors and
SpoVA proteins of Bacillus subtilis spores. FEMS Microbiol Lett 2007 SEP;274(1):42-47.

(46) Zhang P, Kong L, Wang G, Setlow P, Li YQ. Combination of Raman tweezers and quantitative
di42erential interference contrast microscopy for measurement of dynamics and heterogeneity during
the germination of individual bacterial spores. J Biomed Opt 2010 Sep-Oct;15(5):056010.

(47) Raj A, van Oudenaarden A. Single-molecule approaches to stochastic gene expression. Ann Rev
Biophys 2009;38:255-270.

(48) Chada VG, Sanstad EA, Wang R, Driks A. Morphogenesis of *Bacillus* spore surfaces. J Bacteriol
2003 Nov;185(21):6255-6261.

(49) Driks A. *Bacillus subtilis* spore coat. Microbiol Mol Biol Rev 1999 Mar;63(1):1-20.