POI-associated LMD-3 mutation impairs B12-regulated lysosomal function and reproductive capacity in *C. elegans*

Yile Zhai[1]*, Tiantian Wang[2]*, Meixi Gong[2], Yuhan Liang[2], Wenfei Li[2#], Xin Wang[3#],

Zhe Zhang[1,2#]

[1]Institute of Medical Sciences, the Second Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong, China.

[2]Department of Histology and Embryology, School of Basic Medical Sciences, Cheeloo College of Medicine, Shandong University, Jinan, Shandong, China.

[3]State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, Yunnan University, Kunming, Yunnan, China.

*These authors contributed equally: Yile Zhai, Tiantian Wang.

[#]Correspondence: wenfeili@sdu.edu.cn (W. Li), xinwang2@ynu.edu.cn (X. Wang), zhezhang@sdu.edu.cn (Z. Zhang)

1   **Abstract**

Reproductive longevity decline is a key feature of aging and premature ovarian

insufficiency (POI). While proteostasis collapse is implicated in reproductive aging, the

molecular link remains elusive. Here, we engineered the clinically relevant W690C

mutation into *C. elegans lmd-3* gene, the human *NCOA7* ortholog, to establish a

valuable in vivo POI model. The mutation severely compromises LMD-3 protein

stability and expression, leading to a profound collapse in reproductive capacity driven

by germline apoptosis. We demonstrate that this defect drives catastrophic

autophagic-lysosomal dysfunction, blocking degradation and causing proteotoxic

accumulation of misfolded proteins. Crucially, we define this pathology as a functional

B12 deficiency and show that mecobalamin (meCbl) supplementation successfully

restores proteostasis. Together, these findings delineate a conserved genetic pathway

from LMD-3 destabilization to reproductive failure and propose vitamin B12 as a

readily translatable therapeutic intervention for age-related reproductive decline.

Introduction

Reproductive longevity is fundamental for female fertility and healthy aging, yet its decline is one of the first biological processes to emerge in many organisms, from *C. elegans* to humans[1,2]. In humans, the ovary is uniquely susceptible, with its function diminishing decades before other somatic systems[3,4]. This early and pronounced decline contributes significantly to age-related infertility and premature ovarian insufficiency (POI), posing a major global health challenge[1,5]. A growing body of evidence links this decline to impaired proteostasis, the cellular network that maintains protein quality control[6]. While cellular stress responses and protein quality control systems are evolutionarily shared across species[7-9], the specific molecular pathways that connect these processes to reproductive failure remain elusive.

The nematode *C. elegans* provides a powerful and evolutionarily conserved model for investigating reproductive aging[10,11]. Its distinct and early decline in fertility mirrors the human condition, making it an ideal platform to dissect the molecular basis of reproductive health[2,12]. This decline is largely driven by a loss of germline integrity and function, which is highly dependent on the maintenance of a healthy proteome, a process regulated by conserved cellular systems, including the lysosomal and autophagic pathways[13-15].

The oxidation resistance 1 (OXR1) protein family is a key player in this process, comprised of OXR1 and nuclear receptor coactivator 7 (NCOA7) in mammals[16]. This evolutionarily conserved family acts as a critical protective mechanism against cellular

stress[17,18]. NCOA7 has recently been identified as a key regulator of ovarian longevity in humans and mice, and its orthologs have been shown to protect against oxidative damage in other species like *C. elegans* and zebrafish[19-21]. The TLDc domain of these proteins is known to directly interact with the V-type ATPase, a proton pump essential for lysosomal acidification and degradative function[22,23]. This shared function and evolutionary conservation highlight the potential of using *C. elegans* model to gain insights into human disease.

In this study, we engineered a clinically relevant human mutation in *C. elegans* to investigate the pathogenic mechanism of POI. We created the W690C single-point mutation in the conserved TLDc domain of *C. elegans* LMD-3, which perfectly corresponds to a pathogenic NOCA7[W804C] mutation found in human patients with POI[19]. We demonstrate that this single-point mutation significantly impairs reproductive capacity, causing reduced germline cell numbers and increased apoptosis. Mechanistically, our findings reveal that the mutation severely compromises LMD-3 protein stability and gene expression, leading to profound cellular autophagic-lysosomal dysfunction. This defect impairs both lysosomal acidification and its degradative capacity, causing the widespread accumulation of misfolded proteins and aggregates in the germline.

Critically, we show that this defect results in a functional B12 deficiency that can be corrected by mecobalamin (meCbl) supplementation, thereby restoring proteostasis. These findings provide critical insights into the molecular basis of female reproductive longevity, establishing a valuable genetic model for studying POI and highlighting a

potential therapeutic target for mitigating age-related reproductive decline.

Results

Genetic and structural evidence for a shared POI-associated W690C mutation in

LMD-3

The *C. elegans* LMD-3 protein, which serves as the sole OXR1 family member in the

nematode, possesses a domain architecture highly conserved with its human

orthologs, OXR1 and NCOA7 (Fig. 1A). This architecture is defined by three highly

conserved LysM, GRAM, and TLDc domains. Despite the low 26.46% overall amino

acid sequence identity between LMD-3 and human NCOA7[24], the key functional

domains, including the TLDc domain and other conserved motifs, are remarkably

homologous (Supplementary Fig. 1).

The conservation of these domains is further highlighted by the POI-associated

tryptophan residue (W804) in the TLDc domain of human NCOA7[19]. This residue is

perfectly conserved across multiple species, including the corresponding W690

residue in *C. elegans* LMD-3 (Fig. 1B and Supplementary Fig. 1)[25]. The remarkable

conservation of this clinically relevant residue across species provides a strong genetic

basis for a shared pathogenic mechanism and validates the use of *C. elegans* W690C

mutant as a model for human POI.

Following strong genetic evidence for a shared mechanism, we next sought to

determine if this functional similarity was underpinned by structural conservation. A

comparative structural analysis using AlphaFold3 revealed a remarkable similarity

between the predicted *C. elegans* LMD-3 structure and its human NCOA7 ortholog

(Fig. 1C). Despite differences in unique α-helices and intrinsic disordered regions

(IDRs), the core LysM, GRAM, and TLDc domains were highly conserved. This

structural homology was particularly striking in the V-ATPase-interacting TLDc domain,

which showed an almost identical conformation and the conserved W804/W690 site

(Fig. 1C and Supplementary Movie 1). These results provide compelling structural

evidence that the TLDc-V-ATPase interaction is a central, conserved mechanism

governing lysosomal activity and proteostasis across species.

Based on this strong evidence for a shared mechanism, we created an in vivo

model of human POI by introducing the orthologous W690C mutation into *C. elegans*

lmd-3 using CRISPR/Cas9 (Fig. 1D and supplementary Table 1). A preliminary analysis

of the *lmd-3*[W690C] homozygous mutant animals revealed no obvious developmental

defects, as confirmed by a lack of significant difference in body length compared to

wild-type animals (Fig. 1F, G). The lack of significant developmental defects, consistent

with the primary reproductive phenotype observed in human POI[19], establishes the

lmd-3[W690C] strain as a high-fidelity in vivo model for investigating the molecular

mechanisms of this disease.

The POI-associated W690C mutation impairs fecundity and germline integrity in

C. elegans

The deleterious W804C mutation in human NCOA7 has been shown to contribute to

POI by accelerating oxidative stress-induced cellular senescence in ovarian granulosa

cells[19]. Given the perfect conservation of this residue in *C. elegans* LMD-3 and the

established role of this protein in oxidative stress resistance[20,26], we investigated

whether the orthologous *lmd-3*[W690C] mutation would impair reproductive capacity in a

manner consistent with the human disease. Our results showed a pronounced defect

in fecundity, with the mutant strain exhibiting a 32.1% reduction in total progeny over

their lifespan compared to wild-type animals (Fig. 2A and Supplementary Fig. 2A, B).

Mirroring the rapid decline in fertility observed in human POI, this reduction was

particularly striking during the peak reproductive period, suggesting that the mutation

causes an accelerated and severe decline in fertility.

To understand the cellular basis for this reduced fecundity, we next assessed

germline integrity, hypothesizing that the pronounced reduction in fecundity was a

direct result of germ cell deficiency[27]. The *C. elegans* germline is a dynamic, spatially

organized tissue that produces all progeny. It is divided into distinct zones of

proliferation, meiosis, and differentiation, all of which are essential for producing viable

offspring (Fig. 2B). Germ cells are maintained by a distal stem cell region, regulated

by the distal tip cell (DTC), which allows for clear visualization and analysis of germ

cell progression throughout the gonad[28].

Quantitative analysis of DAPI-stained gonads confirmed a significant reduction in

the total number of germ cells in Day 1 adult *lmd-3*[W690C] mutants (Fig. 2C, D). This

defect was accompanied by a specific decline in the number of proliferative germline

stem cells (Fig. 2E), indicating that the *lmd-3*[W690C] mutation fundamentally

compromises the maintenance of the germline progenitor pool, which is essential for

ongoing germ cell production and homeostasis[29].

To investigate the mechanism driving germ cell loss, we assessed germline

apoptosis, a process frequently linked to oxidative stress[30]. Using two complementary

methods, the nucleic acid dye SYTO12 and a *lim-7*p::*ced-1*::GFP reporter[31,32], we

observed a significant increase in apoptotic cell corpses within the germline of W690C

mutants at Day1 and Day2 of adulthood (Fig. 2F, G and Supplementary Fig. 2C, D).

The finding directly links increased germ cell death to reduced fecundity in this POI

model. Our results collectively demonstrate the *lmd-3*[W690C] mutation compromises

germline integrity and reproductive capacity by promoting apoptosis, validating the

mutant as a valuable in vivo model for human POI.

The LMD-3 W690C mutation induces widespread and chronic cellular stress

To gain a deeper understanding of the physiological role of LMD-3, we first

characterized its expression and subcellular localization. An integrated (Is)

transcriptional reporter revealed that *lmd-3* is broadly expressed, including the nervous

system, intestine, and vulva (Fig. 3A)[26]. Subcellular fractionation of the N-terminal

mCherry-tagged LMD-3 protein and its TLDc domain confirmed their predominant

localization to the cytoplasm and membrane fractions (Fig. 3B), consistent with its

known association with lysosomes and the V-ATPase complex[22,23]. A minor nuclear

fraction was also observed, which aligns with the transcriptional and antioxidant

functions of its orthologs NCOA7 and OXR1 (Fig. 3B)[33,34].

Based on the established link between oxidative stress-induced cellular

senescence and POI[19], and given that the LMD-3 orthologs in yeast and humans are

known to be stress-inducible[26,35], we hypothesized that LMD-3 expression would

increase with age as a protective response. Indeed, we found a gradual, age-

dependent increase in *lmd-3* transcription (Fig. 3C, D), a pattern suggesting that LMD-

3 is actively involved in preserving cellular health and longevity as the organism ages.

This finding further supports a conserved role for LMD-3 as a protective factor against

age-related stress, thereby promoting survival.

To determine if the W690C mutation compromises cellular stress resilience, we

monitored cytoplasmic and endoplasmic reticulum (ER) stress throughout the late larval

and early adult stages. We found that the cytosolic unfolded protein response (UPR^{cyto}),

monitored by the *hsp-16.2*p::GFP reporter[36], showed a gradual but dramatic induction

in the mutants (Fig. 3E, F and Supplementary Table 2), indicating an increase in

oxidative stress. In parallel, the ER unfolded protein response (UPR^{ER}), as indicated

by the increased expression of the *hsp-4*p::GFP reporter[37], also showed a progressive

but apparent activation (Fig. 3G, H and Supplementary Table 2).

These results show that the POI-associated W690C mutation induces widespread

but relatively mild and chronic cellular stress, as evidenced by the activation of both

cytosolic and ER stress pathways. However, this general stress response is not

sufficient to fully account for the early and severe reproductive decline observed in the

mutant. This discrepancy suggests that the reproductive failure is not a secondary

consequence of this general chronic stress but rather stems from a direct and severe

breakdown in essential germline protein quality control, specifically through impaired

lysosomal function and protein degradation[6,19,38].

The W690C-induced lysosomal dysfunction drives proteostasis collapse

Building on the established interaction between the TLDc domain and V-ATPases[22,23],

and the central role of lysosomes in aging, we investigated the impact of the W690C

mutation on lysosomal function. Our transcriptional analysis revealed a significant

downregulation of a wide range of lysosome-related genes in the mutant (Fig. 4A and

Supplementary Fig. 3A). This included key components for acidification (V-ATPases),

lysosomal structural integrity (LMP-2, NCR-1), and degradation (cathepsin proteases

ASP-3, ASP-4). The initial finding suggests that the W690C mutation severely impairs

the biogenesis and overall function of the lysosomal system.

To directly confirm that the transcriptional defects led to a loss of lysosomal

function, we measured lysosomal acidification, a process critical for protein

degradation[39]. The use of a heat-shock-inducible pH-sensitive reporter, NUC-

1::pHTomato[40], revealed a significantly higher fluorescence intensity in the mutant,

indicating a less acidic, more neutralized lysosomal pH (Fig. 4B, C and Supplementary

Fig. 3B, C). This defect was independently corroborated by co-staining with the

acidotropic dye LysoSensor Green (LSG) and the lysosome-specific dye LysoTracker

Red (LTR)[41]. The resulting reduced LSG/LTR fluorescence ratio in the mutant

definitively confirmed that the transcriptional defects resulted in a severe impairment

of lysosomal acidity in *C. elegans* (Fig. 4D, E and Supplementary Fig. 3D, E).

The consequence of the acidification defect was a severe breakdown in

degradative capacity, which we assessed by monitoring the clearance of secreted

ssGFP[42]. The reporter showed a significant accumulation of fluorescence in the

coelomocytes of the mutant (Fig. 4F, G), demonstrating a failure of lysosomal

degradation. This defective degradation led to a broad collapse in proteostasis, as

evidenced by extensive aggregation of the age-related protein NMY-2::GFP in

oocytes[43] (Fig. 4H) and the polyglutamine reporter Q40::YFP in body wall muscles[44]

(Fig. 4I). The vital role of LMD-3 in protein homeostasis was further highlighted by the

accumulation of vitellogenin, a major egg yolk protein, in the mutant's intestine,

pseudocoelom, and germline (Supplementary Fig. 3F), confirming that LMD-3

deficiency impairs the processing of this crucial reproductive protein.

To establish that the lysosomal defect is a direct cause of the reproductive failure,

we used the *vha-12(ok821)* mutant to genetically neutralize V-ATPase activity. This

loss-of-function allele not only recapitulated the reproductive defects observed in the

W690C mutant but also exacerbated the decline (Supplementary Fig. 3G). These

findings cement a causal link between the W690C-induced V-ATPase dysfunction and

reproductive decline, underscoring the vital role of LMD-3 in maintaining lysosomal

function for reproductive longevity.

Vitamin B12 restores proteostasis by repairing LMD-3(W690C)-mediated autophagic-lysosomal defects

Following our discovery of profound lysosomal dysfunction and proteostasis collapse in the LMD-3(W690C) mutant, which mirrors the defective autophagy and senescence linked to NCOA7 dysfunction in human POI[19], we investigated the integrity of the autophagic pathway. We first utilized a transcriptional reporter for the stress-inducible autophagy gene, *tts-1*[45]. In contrast to wild-type animals, the W690C mutants showed a significant reduction in *tts-1*p::GFP fluorescence (Fig. 5A). This was further supported by the notable downregulation of key autophagy genes, including the master transcription factor *hlh-30*[46], along with genes essential for autophagosome formation, cargo recognition, fusion, and lysosomal degradation (Fig. 5B, Supplementary Fig. 5A). These findings suggested that the W690C mutation severely compromises the biogenesis and overall function of the autophagic system.

To confirm that this transcriptional defect leads to a functional impairment, we proceeded to examine the accumulation of SQST-1, the *C. elegans* ortholog of p62. Efficient clearance of SQST-1 is indicative of a healthy autophagic flux, and we observed a massive accumulation of SQST-1::GFP aggregates in the adult W690C mutants (Fig. 5C and Supplementary Fig. 4B), providing strong evidence that the overall autophagic degradation pathway is severely compromised.

To precisely characterize autophagic failure, we first utilized tandem fluorescent

reporter mCherry::GFP::LGG-1 to distinguish between autophagosomes formation

(yellow puncta) and maturation into autolysosomes (red puncta), where acid-sensitive

GFP is quenched in the acidic lysosomal environment[47]. Confocal analysis revealed a

dramatic increase in yellow puncta and a parallel decrease in red puncta in the W690C

mutant (Fig. 5D, E and Supplementary Fig. 4C, D). This block in autophagic flux at the

fusion/degradation stage directly reinforces the severe lysosomal acidification defect

observed previously (Fig. 4B-E). Furthermore, GFP::LGG-1 Western blot analysis

revealed a simultaneous defect in autophagosome biogenesis[48], shown by the

significant drop in the ratio of membrane-associated LGG-1-II (lipidated) to cytosolic

LGG-1-I (unlipidated) (Fig. 5F). These findings confirm the W690C mutation induces a

comprehensive, multi-stage failure of the autophagic-lysosomal pathway.

Given that LMD-3[W690C] dysfunction impairs lysosomal function and reproductive

health, we hypothesized that enhancing cellular resilience could restore proper

lysosomal function. Since lysosomes are critical for the uptake and processing of

dietary vitamin B12, the catastrophic lysosomal failure in *lmd-3* mutants would result

in a state of functional B12 deficiency[49]. We therefore tested the capacity of the active

form, meCbl, to mitigate this lysosomal dysfunction. meCbl supplementation

significantly rescued the accumulation of the aggregation-prone protein NMY-2::GFP

in the germline (Fig. 5G, H) and improved protein clearance in coelomocytes, as

evidenced by reduced ssGFP fluorescence (Supplementary Fig. 4E, F). Collectively,

these results demonstrate that B12 supplementation mitigates cellular stresses and

directly improves the defective lysosomal function, thereby restoring germline

proteostasis and protein clearance in *lmd-3*[W690C] mutants.

The W690C mutation compromises LMD-3 structure and expression

Having demonstrated the essential role of the W690C mutation in disrupting lysosomal

function and cellular homeostasis in *C. elegans*, we set out to determine the precise

molecular basis of this effect. We first investigated whether the mutation affected LMD-

3 expression at the transcriptional level. Using primers designed to target three distinct

locations along the gene (Fig. 6A and supplementary Table 3), we detected a

significant and consistent reduction in the overall *lmd-3* transcript level in the mutant

strain compared to the wild type (Fig. 6B and Supplementary Fig. 5A). This reduction

in transcript levels exacerbates the loss-of-function phenotype.

With the established link between human POI-associated NCOA7 mutations and

protein stability[19], we reasoned that the W690C mutation would compromise the

structure of LMD-3. To confirm this hypothesis, three-dimensional structural models of

the *C. elegans* wild-type and W690C-mutated LMD-3 protein were developed[50]. Our

model showed that the LMD-3 protein, like its human ortholog NCOA7, forms a

conserved trimer-like region comprised of its LysM, GRAM, and TLDc domains (Fig.

6C). Although the overall core domains maintained their general conformation, we

found that the W690C mutation significantly altered the inter-subunit distances within

the predicted trimer-like region (Fig. 6D). This structural change suggests a

destabilization of a key binding interface. Critically, six machine learning-based

methods confirmed that the W690C mutation leads to a compromise in overall protein

stability (supplementary Table 4)[51-55], consistent with findings from a human study[19].

To directly confirm that the W690C mutation results in reduced protein stability, we

performed a cycloheximide (CHX) chase assay to evaluate the turnover of LMD-3

protein[56]. As a result, while wild-type LMD-3 protein remained relatively stable, the

mutant protein exhibited a significant decrease in levels, showing a marked

degradation after 12 hours of CHX treatment (Fig. 6E, F and Supplementary Fig. 5B).

This result provides compelling evidence that the W690C mutation compromises the

stability of LMD-3 protein, mechanistically linking the structural defect to its rapid

degradation. Collectively, these findings establish that the POI-associated W690C

mutation compromises both *lmd-3* gene expression and protein stability, thereby

leading to the lysosomal defects and reproductive decline.

Discussion

Reproductive longevity is a fundamental aspect of healthy aging, yet its molecular

mechanisms remain incompletely understood. Our study establishes a novel POI-

associated *C. elegans* model to dissect the pathophysiological basis underlying fertility

decline. By introducing an orthologous W690C mutation into LMD-3, the single-gene

family member of human NCOA7 and OXR1, we found compelling evidence that this

genetic lesion impairs reproductive capacity by compromising protein stability and

inducing a profound collapse in cellular proteostasis. These findings define a specific

pathogenic mechanism in which a single amino acid substitution leads to widespread

cellular dysfunction, thereby directly linking a conserved protein to the maintenance of

reproductive health.

Our study reveals that the W690C mutation, located within the conserved TLDc

domain, severely compromises LMD-3 protein stability. This finding provides a crucial

mechanistic link between the genetic mutation and the observed loss-of-function

phenotypes. A key insight from our data is that this mutation also severely reduces the

overall *lmd-3* mRNA level (Fig. 6B), which directly contributes to the loss-of-function

phenotype. This observation contrasts sharply with the human NCOA7[W804C] mutation,

where mRNA levels are unaffected[19]. This observed discrepancy in mRNA levels is a

critical point that highlights the mechanistic advantage of studying a non-redundant

system. In mammals, the presence of the ortholog OXR1 likely provides cellular

buffering or compensation that masks any regulatory feedback related to NCOA7 loss.

As *lmd-3* is the sole member of this gene family in *C. elegans*, its dysfunction has a

more direct, non-redundant, and severe impact, making our model an ideal system to

study the core pathogenic mechanism.

The destabilization of LMD-3[W690C] protein leads to profound lysosomal dysfunction,

characterized by defective acidification and impaired degradative capacity. These data

align with emerging evidence that lysosomes are not merely "waste disposal plants"

but critical signaling hubs whose dysfunction is linked to aging and neurodegenerative

diseases[57-59]. Our study now unequivocally links organelle failure to a major defect in

reproductive capacity, thereby establishing a direct causal link. This primary lysosomal

defect triggers a downstream cascade of cellular damage, as the accumulation of

misfolded proteins and aggregates creates a proteotoxic environment that is

detrimental to sensitive, non-dividing cells like oocytes[6,60]. Such chronic cellular stress

eventually manifests as increased germ cell apoptosis and a reduction in total cell

number, fully accounting for the decline in reproductive capacity in the mutant.

Ultimately, our findings reveal an elegant pathway linking impaired proteostasis, a well-

known hallmark of aging[61], to the specific context of ovarian aging.

Beyond identifying the pathogenic mechanism, our study provides pioneering

evidence for a therapeutic strategy targeting this conserved pathway. Given that

lysosomal failure compromises the cellular uptake and processing of vitamin B12, the

LMD-3^{W690C} mutation creates a state of functional B12 deficiency that subsequently

disrupts propionate metabolism. This is critical because B12 is an essential cofactor

for the enzyme methylmalonyl-CoA mutase, which catalyzes a required step in

propionate metabolism[62]. A functional deficiency in meCbl impairs this pathway, which

is known to lead to toxic metabolite accumulation and cellular stress, thereby

exacerbating the lysosomal dysfunction[63]. Supplementation with the active form,

meCbl, successfully rescued the LMD-3^{W690C}-induced proteostasis collapse, directly

restoring germline protein homeostasis and enhancing clearance. This highly

significant finding suggests a simple nutritional supplement could mitigate the effects

of this severe genetic dysfunction, highlighting the immediate clinical relevance of our

C. elegans model.

Our human disease-driven *C. elegans* model represents a significant

advancement in the study of reproductive aging. By establishing a direct causal link

between a specific genetic lesion, lysosomal dysfunction, proteotoxicity, and

reproductive failure, our model moves beyond correlational studies. Furthermore, the

model is highly relevant to human POI, as it recapitulates key features of the human

condition, including germ cell depletion and premature reproductive decline[64].

Beyond the specific W690C mutation, our work suggests that LMD-3/NCOA7 and

its genetic interactors are promising candidates for genetic screening in patients with

idiopathic POI. Furthermore, our findings raise broader questions about the role of wild-

type LMD-3/NCOA7 in age-related fertility decline. Recent evidence has shown that

the expression and function of this protein naturally wane with age, leading to a gradual

decline in the lysosomal and proteostasis health observed in aged oocytes[19].

Consequently, our mutant model serves as an accelerated version of the natural aging

process, providing a powerful system for dissecting these fundamental mechanisms.

The *C. elegans* model we established provides an invaluable high-throughput in

vivo platform for screening potential drug interventions that can enhance lysosomal

activity through mechanisms like TFEB activation or other protein homeostasis-

restoring pathways like autophagy and proteostasis[47,65,66]. Furthermore, while our

study establishes LMD-3 as a crucial regulator of lysosomal activity, its precise

biochemical function within this organelle remains to be fully elucidated. Therefore,

further studies are required to identify its binding partners and determine whether it

acts as a scaffold, a regulator, or serves another function in maintaining lysosomal pH

and enzymatic activity.

In summary our study establishes that the POI-associated W690C mutation in

LMD-3, supported by genetic and structural evidence, compromises protein stability

and expression, leading to a profound collapse in lysosomal function. This collapse

triggers proteotoxic stress, germ cell death, and reproductive failure. Crucially, we

define the resultant phenotype as a functional B12 deficiency, rationalizing

mecobalamin supplementation as a viable therapeutic strategy. This discovery

provides a valuable in vivo platform for screening potential therapeutic compounds.

Furthermore, it establishes lysosomal activity as a crucial determinant of reproductive

longevity and a promising target for extending reproductive capacity.

Methods

C. elegans **strains and maintenance**

We maintained all *C. elegans* strains (Supplementary Table 1), including the wild-type

N2 Bristol strain and all other strains, at 20°C on standard nematode growth medium

(NGM) plates. These plates were uniformly seeded with *Escherichia coli* OP50 as a

food source. To ensure experimental synchrony, we obtained age-matched

populations by treating gravid adults with a bleaching solution to isolate and hatch eggs.

CRISPR-Cas9 mediated mutagenesis

The *lmd-3*^{W690C} point mutation was generated by SunyBiotech using the established

CRISPR-Cas9 gene editing system. The presence of the desired point mutation was then definitively confirmed by Sanger sequencing (Supplementary Table 3). To remove any potential off-target mutations and ensure a clean genetic background, the verified mutant was subsequently outcrossed to the wild-type N2 strain three times before all phenotypic and functional analyses commenced.

Plasmid construction

To facilitate biochemical analyses in mammalian cells, we generated expression vectors for 1xFLAG-tagged LMD-3 proteins. The full-length *lmd-3* cDNA sequence were amplified by PCR from both the wild-type and *lmd-3*^{W690C} mutant strains. These amplified fragments were subsequently subcloned into the pcDNA3.1 vector. Primers used for the amplification are listed (Supplementary Table 3).

Quantitative real-time PCR (qRT-PCR)

To quantify target gene transcripts, 200 synchronized Day 1 adult animals were collected into M9 buffer, lysed using Trizol Reagent (Invitrogen, 15596026CN), and immediately frozen in liquid nitrogen. Total RNA was subsequently isolated via standard chloroform extraction and isopropanol precipitation. A total of 1.2 μg isolated RNA was then reverse-transcribed into complementary DNA (cDNA) using the HiScript III 1st Stand cDNA Synthesis Kit (+ gDNA wiper) (Vazyme, R312-01). Quantitative real-time PCR (qRT-PCR) was performed on an ABI QuantStudio 1 machine using ChamQ

SYBR Color qPCR Master Mix (Vazyme, Q411-02). Transcript levels for the target

genes were calculated and normalized to the endogenous control gene, *act−1*.

Reproductive span analysis

As previously described, a minimum of 10 synchronized hermaphrodite larvae at L4

stage for each strain were individually transferred to fresh NGM plates at every 12

hours[67]. This transfer process continued until the cessation of reproduction, allowing

for the isolation of all laid eggs. The number of progenies was counted at each transfer,

and the total number of progenies per animal was calculated. The experiment was

performed in three biological replicates.

Measurement of body length

Synchronized *C. elegans* at designated developmental stages were first anesthetized

using levamisole and mounted onto agarose pads. Images were captured using a

Nexcope 910 microscopy system. The linear distance from the head to the tail of each

worm was then measured using the Image View software to determine body length. A

minimum of 22 worms were analyzed for each group.

Microscopy and imaging analysis

For all standard fluorescence and bright-field microscopy experiments, synchronized

animals were first anesthetized using levamisole and mounted onto agarose pads for

imaging. Images were captured using either a Nexcope 910 fluorescence microscope

or a high-resolution laser scanning confocal microscope (Carl Zeiss, LSM 980),

depending on the required resolution and image type. Subsequently, fluorescence

intensity and other quantitative metrics, such as puncta number or size, were

measured and analyzed using the ImageJ software package.

Germline staining and quantification

To analyze the integrity and morphology of the germline, synchronized Day 1 adult

animals were first fixed by immersion in ice-cold methanol for 5 minutes. Following the

removal of methanol, the samples were washed three times with M9 buffer. Germ cell

nuclei were then stained by incubating the samples in a DAPI solution (Macklin,

D807022) for 30 minutes in the dark, followed by three additional washes. The stained

worms were then mounted, and their gonads were examined using a laser scanning

confocal microscope. The total number of germ cells and the specific count of

proliferative germline stem cells in each gonad were subsequently determined

quantitatively using ImageJ software.

SYTO12 staining for germline apoptotic cells

Synchronized larvae from control and experimental groups were collected and allowed

to develop into Day 1 and Day 2 adults. These adult animals were then stained by

incubation in SYTO 12 dye (Thermo Fisher Scientific, S7575) in the dark for 3 hours. Following staining, worms were washed three times with buffer via centrifugation (4,000 rpm, 1 minute) and transferred to fresh NGM plates for a 1-hour recovery period to clear dye-labeled bacteria from the intestines. Finally, the worms were anesthetized with levamisole, mounted on agarose pads, and imaged using an objective under green fluorescence microscopy to count the apoptotic cell corpses.

Dual lysosomal staining and quantification

Synchronized Day 1 adult animals were co-stained for 1 hour at 20°C in the dark with 10 mM LysoSensor Green DND-189 (Yeasen) and LysoTracker Red DND-99 (Yeasen), as previously described[68]. The animals were transferred to fresh NGM plates seeded with OP50 for a 1-hour recovery period to clear residual dye from the gut bacteria. Lysosomal acidification was then quantitatively evaluated by measuring the relative fluorescence intensity ratio of LysoSensor Green to LysoTracker Red using ImageJ software, where a lower ratio indicates reduced lysosomal acidity.

Cell culture and Transient transfection

HEK-293T cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM) (Gibco, C119955-00BT) supplemented with 10% fetal bovine serum (FBS) (Excell, FSP-500) in a humidified atmosphere at 37°C with 5% CO_2. For transient transfection experiments, cells were transfected with the appropriate plasmid DNA using

polyethyleneimine (PEI) (Sigma-Aldrich, 408727). The transfection mixture was

prepared using a DNA-to-PEI mass ratio of 1:4.5, with a total of 1 µg DNA per 10^6

cells used for each condition.

Cycloheximide pulse chase assay

The compromised stability of the 1xFLAG-tagged protein was evaluated in cells using

a Cycloheximide (CHX) chase assay, which monitors protein turnover[69]. Thirty-six

hours after transfection, cells were subjected to 50 µg/mL CHX to block protein

synthesis. At designated time points, cells were collected, and the degradation of wild-

type and mutant were determined by quantifying protein levels via Western blotting.

Autophagy flux and aggregation analysis

To assess autophagy flux, synchronized worms expressing the tandem fluorescent

reporter mCherry::GFP::LGG-1 were used. Puncta were counted in the anterior

intestinal cells, where the number of autophagosomes (AP) was determined by

counting the puncta positive for both the GFP and mCherry (yellow/orange), and the

number of autolysosomes (AL) was determined by counting mCherry-only positive

puncta (red, due to quenching in the acidic lysosome). Additionally, to evaluate the

accumulation of aggregated cargo, the number of SQST-1::GFP positive puncta was

counted specifically within the head neurons of synchronized worms. Both assays

were conducted with a minimum of 11 worms examined per condition across three

independent experiments.

Mecobalamin supplementation

A 150 µM stock solution of meCbl (Macklin, M812748) was initially prepared in H_2O

(which also served as the control solvent). This stock solution was then diluted to a

final working concentration of 0.15 µM directly into the NGM agar prior to pouring the

plates, ensuring the worms were exposed to the compound throughout their life cycle.

Quantification of *myo-3*p::ssGFP intensity

Fluorescence images were captured from synchronized adult *C. elegans* at Day 1 of

adulthood expressing the secreted reporter *myo-3*p::ssGFP using Nexcope

microscopy. The average fluorescence intensity of the ssGFP signal was quantified

using ImageJ software. At least 18 animals were analyzed for each strain and condition.

Quantification of NUC-1::pHTomato intensity

Synchronized Day 1 adult *C. elegans* expressing P_{hs}NUC-1::pHTomato were first

subjected to a 33°C heat shock for 30 minutes to induce expression. Following

induction, animals were allowed to recover at 20°C for 24 hours before imaging[40,43].

The average intensity of pHTomato was then quantified per area in the hypodermis

using ImageJ software. A minimum of 20 worms were analyzed for each strain.

Western blotting

A total of 200 synchronized day 1 adult worms were collected, lysed directly in 1×

sample buffer, and boiled at 95°C for 10 minutes. The resulting protein extracts were

separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred

onto a PVDF membrane. The membrane was blocked for 1 hour at room temperature

using 5% non-fat milk in TBST before being incubated overnight at 4°C with the

following primary antibodies: anti-GFP rabbit polyclonal antibody (1:1000, abclonal,

AE011), anti-α-tubulin mouse monoclonal antibody (1:5000, Sigma-Aldrich, T5168),

anti- flag DDDDK-tag mAb antibody (1:10000, MBL, M185) and anti-GAPDH antibody

(1:500, Absin, abs830030ss). After washing, the membrane was incubated with HRP-

conjugated secondary antibodies for 1 hour at room temperature. Finally, protein bands

were visualized using an enhanced chemiluminescence (ECL) detection system.

Protein Sequence and Structure Prediction

The protein sequences for human NCOA7 (NCBI accession: NP_001186548.1) and

its *C. elegans* ortholog LMD-3 (NCBI accession: NP_505173.2) were retrieved from

the National Center for Biotechnology Information (NCBI) database. Comparative

three-dimensional (3D) structural modeling for wild-type and the W690C mutant was

performed using. The predicted structures were visualized and analyzed using the

ChimeraX 1.10.1 software package.

Inter-domain distance measurement

525   To quantify the structural impact of the mutation on the predicted LMD-3 trimer-like

526   region, the distances between the conserved, and domains were measured. The

527   centroid of each respective domain was defined as a vertex of a triangle. The distances

528   between these centroids were then calculated using the "Distance" function within the

529   "Annotation" module of ChimeraX 1.10.1.

Protein stability prediction

To assess the effect of the point mutation on protein stability, we employed six distinct

machine learning-based prediction tools[51-55]. The LMD-3 protein sequence and 3D

structural model were the same as those used for the structural analysis above. For

each web-based tool, the W690C substitution was specified to calculate the change in

free energy ($\Delta\Delta G$). The definition of a "Destabilizing" prediction varied among the

different websites, corresponding to specific $\Delta\Delta G$ thresholds.

Statistical analysis

All data were statistically analyzed using GraphPad Prism 9. Data are presented as

the mean, with error bars representing the standard error of the mean (SEM). Student's

t-tests were utilized for comparisons between two distinct groups, while one-way

ANOVA tests were performed for multiple group comparisons. Statistical significance

was established at $p < 0.05$ (indicated by *), $p < 0.01$ (indicated by **), and $p <$

0.001(indicated by ***).

Data availability

All data supporting the findings of this study, presented within the main manuscript and

the Supplementary Information, are available in the accompanying Source data file

provided with this paper. All unique reagents and strains generated throughout this

study are available from the corresponding author upon reasonable request and the

completion of a standard Material Transfer Agreement (MTA).

Acknowledgements

We thank the Caenorhabditis Genetics Center (CGC), Dr. Xiaochen Wang (Southern

University of Science and Technology), Dr. Dengke Ma (UCSF), and Dr. Chenggang

Zou (Yunnan University) for strains. Dr. Yingying Qin, Dr. Ting Dong, and Dr. Xue Jiao

for their valuable suggestions (Shandong University).

Funding

This study was supported by a grant (no. 2022KF006 to Z.Z.) from YNCUB, the

National Natural Science Foundation of China (32170781 to Z.Z.), and the Natural

566 Author contributions

Conceptualization: W.L, X.W, Z.Z; Data Curation: Y.Z, T.W, Z.Z; Formal Analysis: Y.Z,

T.W; Funding Acquisition: Z.Z; Investigation: Y.Z, T.W, M.G, W.L, X.W, Z.Z;

Methodology: Y.Z, T.W, M.G, W.L, Z.Z; Project Administration: Y.Z, T.W, W.L, X.W, Z.Z;

Resources: T.W, M.G, Z.Z; Supervision: W.L, X.W, Z.Z; Validation: Y.Z, T.W, M.G, W.L;

Visualization: Y.Z, T.W, M.W, Z.Z; Writing – Original Draft: Y.Z, T.W, M.W; Writing –

Review & Editing: W.L, X.W, Z.Z.

574 Competing interests

575 The authors declare no competing interests.

References

1 Broekmans, F., Soules, M. & Fauser, B. Ovarian aging: mechanisms and clinical
consequences. *Endocrine reviews* **30**, 465-493 (2009).

2 Scharf, A., Pohl, F., Egan, B. M., Kocsisova, Z. & Kornfeld, K. Reproductive aging
in Caenorhabditis elegans: from molecules to ecology. *Frontiers in cell and*
developmental biology **9**, 718522 (2021).

3 Benayoun, B. A., Kochersberger, A. & Garrison, J. L. Studying ovarian aging and
its health impacts: modern tools and approaches. *Genes & Development* **39**, 975-
990 (2025).

4 Te Velde, E. R. & Pearson, P. L. The variability of female reproductive ageing.
Human reproduction update **8**, 141-154 (2002).

5   Park, S. U., Walsh, L. & Berkowitz, K. M. Mechanisms of ovarian aging.
Reproduction **162**, R19-R33 (2021).

6   Sala, A. J. & Morimoto, R. I. Protecting the future: balancing proteostasis for
reproduction. *Trends in cell biology* **32**, 202-215 (2022).

7   Agarwal, A., Gupta, S. & Sharma, R. K. Role of oxidative stress in female
reproduction. *Reproductive biology and endocrinology* **3**, 28 (2005).

8   Agarwal, A., Aponte-Mellado, A., Premkumar, B. J., Shaman, A. & Gupta, S. The
effects of oxidative stress on female reproduction: a review. *Reproductive biology*
and endocrinology **10**, 49 (2012).

9   Fredriksson, Å. *et al.* Effects of aging and reproduction on protein quality control
in soma and gametes of Drosophila melanogaster. *Aging cell* **11**, 634-643 (2012).

10   Ye, A. L. & Bhalla, N. Reproductive aging: insights from model organisms.
Biochemical Society Transactions **39**, 1770-1774 (2011).

11   Hughes, S. E., Evason, K., Xiong, C. & Kornfeld, K. Genetic and pharmacological
factors that influence reproductive aging in nematodes. *PLoS genetics* **3**, e25
(2007).

12   Zimmerman, S. M., Hinkson, I. V., Elias, J. E. & Kim, S. K. Reproductive aging
drives protein accumulation in the uterus and limits lifespan in C. elegans. *PLoS*
genetics **11**, e1005725 (2015).

13   Bohnert, K. A. & Kenyon, C. A lysosomal switch triggers proteostasis renewal in
the immortal C. elegans germ lineage. *Nature* **551**, 629-633 (2017).

14   Song, Z. *et al.* Germ cell-specific Atg7 knockout results in primary ovarian
insufficiency in female mice. *Cell death & disease* **6**, e1589-e1589 (2015).

15   Labbadia, J. & Morimoto, R. I. The biology of proteostasis in aging and disease.
Annual review of biochemistry **84**, 435-464 (2015).

16   Durand, M. *et al.* The OXR domain defines a conserved family of eukaryotic
oxidation resistance proteins. *BMC cell biology* **8**, 13 (2007).

17   Finelli, M. J. & Oliver, P. L. TLDc proteins: new players in the oxidative stress
response and neurological disease. *Mammalian Genome* **28**, 395-406 (2017).

18   Finelli, M. J., Sanchez-Pulido, L., Liu, K. X., Davies, K. E. & Oliver, P. L. The
evolutionarily conserved Tre2/Bub2/Cdc16 (TBC), lysin motif (LysM), domain
catalytic (TLDc) domain is neuroprotective against oxidative stress. *Journal of*
Biological Chemistry **291**, 2751-2763 (2016).

19   Dong, T. *et al.* Stress granule clearance mediated by V-ATPase-interacting protein
NCOA7 mitigates ovarian aging. *Nature Aging*, 1-20 (2025).

20   Sanada, Y. *et al.* Oxidation resistance 1 is essential for protection against oxidative
stress and participates in the regulation of aging in Caenorhabditis elegans. *Free*

radical research **48**, 919-928 (2014).

21 Xu, H. *et al.* Zebrafish Oxr1a knockout reveals its role in regulating antioxidant
defenses and aging. *Genes* **11**, 1118 (2020).

22 Eaton, A. F., Brown, D. & Merkulova, M. The evolutionary conserved TLDc domain
defines a new class of (H+) V-ATPase interacting proteins. *Scientific reports* **11**,
22654 (2021).

23 Oot, R. A. & Wilkens, S. Human V-ATPase function is positively and negatively
regulated by TLDc proteins. *Structure* **32**, 989-1000. e1006 (2024).

24 Madeira, F. *et al.* The EMBL-EBI Job Dispatcher sequence analysis tools
framework in 2024. *Nucleic acids research* **52**, W521-W525 (2024).

25 Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M. & Barton, G. J. Jalview
Version 2—a multiple sequence alignment editor and analysis workbench.
Bioinformatics **25**, 1189-1191 (2009).

26 Moriwaki, T., Kato, S., Kato, Y., Hosoki, A. & Zhang-Akiyama, Q.-M. Extension of
lifespan and protection against oxidative stress by an antioxidant herb mixture
complex (KPG-7) in Caenorhabditis elegans. *Journal of Clinical Biochemistry and*
Nutrition **53**, 81-88 (2013).

27 Hayashi, Y. *et al.* Nutritional and metabolic control of germ cell fate through O-
GlcNAc regulation. *EMBO reports* **24**, e56845 (2023).

28 Albert Hubbard, E. J. & Schedl, T. Biology of the Caenorhabditis elegans germline
stem cell system. *Genetics* **213**, 1145-1188 (2019).

29 Kimble, J. & Seidel, H. C. elegans germline stem cells and their niche. *StemBook*
[Internet] (2013).

30 Gartner, A., Boag, P. R. & Blackwell, T. K. Germline survival and apoptosis.
WormBook: The Online Review of C. elegans Biology [Internet] (2018).

31 Wu, Y.-C., Wang, X. & Xue, D. Methods for studying programmed cell death in C.
elegans. *Methods in Cell Biology* **107**, 295-320 (2012).

32 Li, X., Singh, N., Miller, C., Sosseh, B. & Gordon, K. L. The C. elegans gonadal
sheath Sh1 cells extend asymmetrically over a differentiating germ cell population
in the proliferative zone. *Elife* **11**, e75497 (2022).

33 Halachmi, S. *et al.* Estrogen receptor-associated proteins: possible mediators of
hormone-induced transcription. *Science* **264**, 1455-1458 (1994).

34 Oliver, P. L. *et al.* Oxr1 is essential for protection against oxidative stress-induced
neurodegeneration. *PLoS genetics* **7**, e1002338 (2011).

35 Elliott, N. A. & Volkert, M. R. Stress induction and mitochondrial localization of
Oxr1 proteins in yeast and humans. *Molecular and cellular biology* **24**, 3180-3187
(2004).

36 Rong, B. *et al.* Ribosome 18S m6A methyltransferase METTL5 promotes
translation initiation and breast cancer cell growth. *Cell reports* **33** (2020).

37 Higuchi-Sanabria, R. *et al.* Divergent nodes of non-autonomous UPRER signaling
through serotonergic and dopaminergic neurons. *Cell reports* **33** (2020).

38 Kaushik, S. & Cuervo, A. M. Proteostasis and aging. *Nature medicine* **21**, 1406-
1415 (2015).

39 Lo, C. H. & Zeng, J. Defective lysosomal acidification: a new prognostic marker
and therapeutic target for neurodegenerative diseases. *Translational*
neurodegeneration **12**, 29 (2023).

40 Wang, Z. *et al.* The RBG-1–RBG-2 complex modulates autophagy activity by
regulating lysosomal biogenesis and function in C. elegans. *Journal of Cell*
Science **132**, jcs234195 (2019).

41 Barral, D. C. *et al.* Current methods to analyze lysosome morphology, positioning,
motility and function. *Traffic* **23**, 238-269 (2022).

42 Egami, Y., Taguchi, T., Maekawa, M., Arai, H. & Araki, N. Small GTPases and
phosphoinositides in the regulatory mechanisms of macropinosome formation and
maturation. *Frontiers in physiology* **5**, 374 (2014).

43 Sun, Y. *et al.* Lysosome activity is modulated by multiple longevity pathways and
is important for lifespan extension in C. elegans. *elife* **9**, e55745 (2020).

44 Jia, K., Hart, A. C. & Levine, B. Autophagy genes protect against disease caused
by polyglutamine expansion proteins in Caenorhabditis elegans. *Autophagy* **3**, 21-
25 (2007).

45 Nakamura, S. *et al.* Mondo complexes regulate TFEB via TOR inhibition to
promote longevity in response to gonadal signals. *Nature communications* **7**,
10944 (2016).

46 Chen, H.-D. *et al.* HLH-30/TFEB-mediated autophagy functions in a cell-
autonomous manner for epithelium intrinsic cellular defense against bacterial
pore-forming toxin in C. elegans. *Autophagy* **13**, 371-385 (2017).

47 Chang, J. T., Kumsta, C., Hellman, A. B., Adams, L. M. & Hansen, M.
Spatiotemporal regulation of autophagy during Caenorhabditis elegans aging.
Elife **6**, e18459 (2017).

48 Wu, F. *et al.* Structural basis of the differential function of the two C. elegans Atg8
homologs, LGG-1 and LGG-2, in autophagy. *Molecular cell* **60**, 914-929 (2015).

49 Wei, W. & Ruvkun, G. Lysosomal activity regulates Caenorhabditis elegans
mitochondrial dynamics through vitamin B12 metabolism. *Proceedings of the*
National Academy of Sciences **117**, 19970-19981 (2020).

50 Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with

698   AlphaFold 3. *Nature* **630**, 493-500 (2024).

699   51   Pires, D. E., Ascher, D. B. & Blundell, T. L. mCSM: predicting the effects of
700   mutations in proteins using graph-based signatures. *Bioinformatics* **30**, 335-342
701   (2014).

702   52   Pandurangan, A. P., Ochoa-Montano, B., Ascher, D. B. & Blundell, T. L. SDM: a
703   server for predicting effects of mutations on protein stability. *Nucleic acids*
704   *research* **45**, W229-W235 (2017).

705   53   Pires, D. E., Ascher, D. B. & Blundell, T. L. DUET: a server for predicting effects of
706   mutations on protein stability using an integrated computational approach. *Nucleic*
707   *acids research* **42**, W314-W319 (2014).

708   54   Savojardo, C., Fariselli, P., Martelli, P. L. & Casadio, R. INPS-MD: a web server to
709   predict stability of protein variants from sequence and structure. *Bioinformatics* **32**,
710   2542-2544 (2016).

711   55   Capriotti, E., Calabrese, R. & Casadio, R. Predicting the insurgence of human
712   genetic diseases associated to single point protein mutations with support vector
713   machines and evolutionary information. *Bioinformatics* **22**, 2729-2734 (2006).

714   56   Kao, S.-H. *et al.* Analysis of protein stability by the cycloheximide chase assay.
715   *Bio-protocol* **5**, e1374-e1374 (2015).

716   57   Zhao, Q., Gao, S. M. & Wang, M. C. Molecular mechanisms of lysosome and
717   nucleus communication. *Trends in biochemical sciences* **45**, 978-991 (2020).

718   58   Settembre, C. *et al.* A lysosome-to-nucleus signalling mechanism senses and
719   regulates the lysosome via mTOR and TFEB. *The EMBO journal* **31**, 1095-1108
720   (2012).

721   59   Lamming, D. W. & Bar-Peled, L. Lysosome: The metabolic signaling hub. *Traffic*
722   **20**, 27-38 (2019).

723   60   Jentoft, I. M. A. & Schuh, M. Protein Storage in Oocytes: Implications for Oocyte
724   Quality, Embryonic Development, and Female Fertility. *Annual Review of Cell and*
725   *Developmental Biology* **41** (2025).

726   61   López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks
727   of aging: An expanding universe. *Cell* **186**, 243-278 (2023).

728   62   Takahashi-Iñiguez, T., García-Hernandez, E., Arreguín-Espinosa, R. & Flores, M.
729   E. Role of vitamin B12 on methylmalonyl-CoA mutase activity. *Journal of Zhejiang*
730   *University Science B* **13**, 423-437 (2012).

731   63   Costanzo, M. *et al.* Methylmalonic acidemia triggers lysosomal-autophagy
732   dysfunctions. *Cell & Bioscience* **14**, 63 (2024).

733   64   Touraine, P. *et al.* Premature ovarian insufficiency. *Nature Reviews Disease*
734   *Primers* **10**, 63 (2024).

65 Bretou, M. *et al.* Lysosome signaling controls the migration of dendritic cells.
Science immunology **2**, eaak9573 (2017).

66 Powers, E. T., Morimoto, R. I., Dillin, A., Kelly, J. W. & Balch, W. E. Biological and
chemical approaches to diseases of proteostasis deficiency. *Annual review of*
biochemistry **78**, 959-991 (2009).

67 Wang, M. C., Oakley, H. D., Carr, C. E., Sowa, J. N. & Ruvkun, G. Gene pathways
that delay Caenorhabditis elegans reproductive senescence. *PLoS genetics* **10**,
e1004752 (2014).

68 Pu, X. & Qi, B. Lysosomal dysfunction by inactivation of V-ATPase drives innate
immune response in C. elegans. *Cell Reports* **43** (2024).

69 Liu, Y. *et al.* ALKBH1 promotes HIF-1 α-mediated glycolysis by inhibiting N-
glycosylation of LAMP2A. *Cellular and Molecular Life Sciences* **81**, 130 (2024).

2 Fig. 1

3 Fig 1. Characterization of the POI-associated W690C Mutation in the conserved

4 TLDc domain of *C. elegans* LMD-3. A Schematic representation of conserved

5 domains in *C. elegans* LMD-3 and its human orthologs, OXR1 and NCOA7. The

6 schematic diagrams illustrate the conserved LysM, GRAM, and TLDc domains. The

total number of amino acids (aa) is indicated for each protein. **B** The conserved

tryptophan residue (W804) associated with premature ovarian insufficiency (POI) is

highly conserved across species. A multiple sequence alignment of NCOA7 orthologs

from five different species demonstrates the conservation of POI-associated W804

residue in human NCOA7. The residue, corresponding to W690 in *C. elegans* LMD-3,

is highlighted with a red arrowhead. **C** Comparative structural analysis of the TLDc

domain from human NCOA7 and *C. elegans* LMD-3. The TLDc domains are shown in

detail to highlight their conserved structure similarity. The intrinsically disordered

region (IDR) is colored gray. The conserved POI-associated tryptophan residues

(human W804 and *C. elegans* W690) are indicated by black arrows and are colored

red in the detailed view. **D** Sequencing confirmation of the *C. elegans lmd-3*^{W690C}

knock-in allele. A representative DNA sequencing trace shows the successful creation

of the POI-associated W690C mutation. A red arrow marks the target base change.

The target knock-in site is indicated by a red arrow. The resulting amino acid change

(W to C) is highlighted in red, while a silent mutation introduced for screening is

highlighted in light blue. **F**, **G** Development defects detection between wild-type and

lmd-3^{W690C} animals at D1 and D2 adult stages with bright-field image (**F**) and

quantification (**G**) of animal body lengths (n ≥ 22 for each group, unpaired t-tests: ns,

no significant difference). Scale bar, 50 µm.

 Fig. 2

Fig 2. The POI-associated W690C mutation in *C. elegans* LMD-3 impairs

fecundity and germline integrity. A Total brood sizes indicate the overall

reproductive ability of wild-type and *lmd-3*^{W690C} mutants. Data are presented as mean

± SEM (n = 10 per group, **p < 0.01). **B** Spatial organization of the C. elegans germline

stem cell and meiotic regions. The schematic diagram illustrates one of the two distinct

gonad arms, showing the spatial organization of the stem cell region, transition zone,

and the meiotic phase along the distal-proximal axis. The distal tip cell (DTC) is also

depicted, highlighting its role in enclosing the distal arm. **C, D** Representative

fluorescence images (**C**) and quantification (**D**) of DAPI-stained total germline cells in

wild-type and *lmd-3*^W690C mutants (n = 14 for each group, ***p < 0.001). Scale bars, 50

µm. **D** Representative fluorescence images and quantification of DAPI staining for

germline stem cells and meiotic cells in wild-type and *lmd-3*^W690C mutants (n = 14 for

each group, **p < 0.01, ***p < 0.001). Scale bars, 20 µm. **F, G** Germline apoptosis in

wild-type and *lmd-3*^W690C mutants with SYTO12 staining (**F**) and a *lim-7p::ced-1::GFP*

reporter (**G**). White arrowheads indicate apoptotic cell corpses. n ≥ 13 per group;

unpaired t-tests, ***p < 0.001. D1, Day 1 of adulthood; D2, Day 2 of adulthood. Scale

bars, 50 µm.

Fig. 3

Fig 3. The expression pattern of LMD-3 and the role of W690C mutation in stress

response. A Expression pattern of *lmd-3* at the L4 larval stage. Representative images

show the expression of the *lmd-3p*::GFP transcriptional reporter). Is, integrated

transgene. Scale bars, 100 μm. **B** Subcellular localization of full-length LMD-3 and its

TLDc domain. The distribution of mCherry-tagged full-length LMD-3 (upper) and

mCherry-tagged TLDc domain (lower) was analyzed. The fractions analyzed were

cytoplasmic (Cyto), membrane (Memb), soluble nuclear (SolNu), chromatin-bound

(ChroB), and cytoskeletal (CytoSkel). **C, D** Transcriptional level of *lmd-3* during

different developmental stages with representative fluorescence images (**C**) and

quantification (**D**) using the *lmd-3*p::GFP reporters (n ≥ 6 for each group, ***p < 0.001).

Is, integrated transgene. Scale bars, 100 μm. a.u., arbitrary units. **E, F** Representative

fluorescence images (**E**) and quantification (**F**) of the UPR^{cyto} reporter *hsp-16.2*p::GFP

in wild type and *lmd-3*^{W690C} animals at indicated stages (n ≥ 20 for each group, ns, no

significant difference). Scale bars represent 100 μm. a.u., arbitrary units. L4, the fourth

larval stage. D1, Day 1 of adulthood. D2, Day 2 of adulthood. **G, H** Representative

fluorescence images (**G**) and quantification (**H**) of the UPR^{ER} reporter *hsp-*

*4*p::GFP in wild-type and *lmd-3*^{W690C} animals at indicated stages (n ≥ 10 for each

group, unpaired t-tests: ns, no significant difference, *p < 0.05). Scale bars

represent 100 μm. a.u., arbitrary units.

  Fig. 4

Fig 4. The W690C mutation affects proteostasis and lysosomal function in _C._

elegans. A Transcriptional analysis of lysosome-related genes in wild type and _lmd-_

_3_^W690C animals at the D1 adult stage (n = 3 for each group, unpaired t-tests: ***p* < 0.01,

****p* < 0.001). **B, C** Representative fluorescence images (**B**) and quantification (**C**) of

NUC-1::pHTomato expression in the hypodermis driven by a heat-shock (hs) promoter

(n ≥ 20 per group, ***p < 0.001). a.u., arbitrary units. Scale bar, 10 μm. **D, E**

Representative confocal fluorescence images (**D**) and quantification (**E**) of intestines

co-stained with Lysosome-Specific Green (LSG DND-189) and Lysosome-Targeting

Red (LTR DND-99) in wild type and *lmd-3*^{W690C} animals at the D1 adult stage. White

arrowheads indicate vesicular lysosomes stained by both dyes. Relative intensity of

LSG/LTR were quantified (n ≥ 13 per group, ***p < 0.001). Scale bars, 10 μm. **F, G**

Representative fluorescence images (**F**) and quantification (**G**) of secreted protein

accumulation (*myo-3*p::ssGFP) in the coelomocytes of wild type and *lmd-3*^{W690C}

animals at the D1and D2 stages (n ≥ 18 per group, unpaired t-tests: **p < 0.01, ***p <

0.001). Scale bars, 50 μm. a.u., arbitrary units. **H** Confocal fluorescence images and

quantification of NMY-2::GFP accumulation in wild type and *lmd-3*^{W690C} animals at D1

stage (mean ± SEM, n = 3 biological replicates, 10 animals per condition per replicate,

unpaired t-tests: ***p < 0.001). **I** Fluorescence images and quantification of Q40::YFP

accumulation in the body wall muscles in wild type and *lmd-3*^{W690C} animals at the L4

stage (n ≥ 16 per group, unpaired t-tests: ***p < 0.001). Scale bars, 50 μm.

 Fig. 5

Fig 5. meCbl restores proteostasis in LMD-3(W690C) mutants by repairing

autophagic-lysosomal defects. A Representative fluorescence images and

quantification of the autophagy reporter *tts-1*p::GFP in wild type and *lmd-3*^{W690C} animals

at the indicated stages (n ≥ 10 per group, ***p < 0.001). a.u., arbitrary units. Scale bars

represent 100 μm. **B** Transcriptional analysis of key autophagy genes was performed

in wild type and *lmd-3*^{W690C} animals, and gene expression was quantified by RT-qPCR

(n = 3 per group, unpaired t-tests: *p < 0.05, **p < 0.01, ***p < 0.001, ns, no significant

difference). **C** Representative fluorescence images and quantification of SQST-1::GFP

accumulation in wild type and *lmd-3*^{W690C} animals at D1 adult stage (n ≥ 14 per group,

***p < 0.001). White arrowheads denote fluorescent protein aggregates. Scale bars,

50 μm. **D, E** Representative confocal images (**D**) and quantification (**E**) of the

mCherry::GFP::LGG-1 reporter in wild type and *lmd-3*^{W690C} animals at D1 and D2 adult

stages. The ratio of autophagosomes (yellow puncta, yellow arrow) to total puncta and

autolysosomes (red puncta, red arrow) to total puncta was quantified (n ≥ 11 per group,

***p < 0.001). Scale bar, 10 μm. **F** Representative Western blot and corresponding

quantification of the ratio of lipidated LGG-1-II to unlipidated LGG-1-I protein in wild-

type and *lmd-3*^{W690C} mutant animals (n = 3 biological replicates, ***p < 0.001). a.u.,

arbitrary units. IB, immunoblotting. **G, H** Confocal fluorescence images (**G**) and

corresponding quantification (**H**) of *nmy-2*p::*nmy-2*::GFP accumulation in the oocytes

of wild-type and *lmd-3*^{W690C} mutants at D1 of adulthood, with mecobalamin (meCbl)

supplementation (mean ± SEM, n=3 biological replicates, animals per condition per

replicate = 10, ***p < 0.001, ns, no significant differences). Scale bars, 10 μm.

Fig. 6

Fig 6. The W690C mutation affects LMD-3 protein structure and stability. **A** The

schematic diagram shows the genomic structure of *lmd-3*, including the positions of its

conserved LysM, GRAM, and TLDc domains. The locations of the qRT-PCR primers

(P1, P2, P3) designed to quantify lmd-3 mRNA are also shown. **B** qRT-PCR

measurements for endogenous *lmd-3* mRNA levels in wild type (wt) and *lmd-3*[W690C] animals (n = 3 per group, unpaired t-tests: **$p < 0.01$, ***$p < 0.001$). **C, D**

Structural analysis of the LMD-3[W690C] protein. Predicted three-dimensional structures

of wild-type (**C**) and W690C (**D**) proteins are presented to illustrate the effects of the

W690C mutation on protein folding. Key domains are color-coded: LysM (cornflower

blue), GRAM (light salmon), and TLDc (pale violet red). A magnified view shows the

trimer-like region, where the mutation leads to altered inter-subunit distances in the

p.W690C protein (49.97 Å, 53.43 Å, 36.33 Å) compared to the wild-type (47.58 Å,

52.70 Å, 39.58 Å). **E, F** The W690C mutation reduces LMD-3 protein stability. Western

blot images (**E**) and semi-quantitative analysis (**F**) show the turnover of 1xFLAG-

tagged LMD-3 in H293T cells treated with cycloheximide (CHX) for 0, 6, and 12 h (n =

3 biological replicates, *$p < 0.5$, ***$p < 0.001$). a.u., arbitrary units.