Title

Investigating the potential for mirror self-recognition in honeybees (*Apis mellifera*): a

preliminary study using the mark test

Gentaro Shishimi[1]*, Rena Akaishi[1]†, Mana Matsukami[1]†

[1] Mita International School of Science, Tokyo, Japan

† These authors contributed equally to this work.

* E-mail: g.shishimi@gmail.com (GS)

Abstract

Mirror self-recognition (MSR) is a key indicator of self-awareness, primarily studied in

vertebrates. Investigating MSR in invertebrates such as the western honeybee (*Apis*

mellifera), an insect with advanced cognition, is essential to understanding the

evolutionary origins and potential convergent evolution of self-recognition. We

conducted two experiments using a classic mark test, applying a paint mark to the

clypeus, a body part invisible to the bee without a mirror, and compared head-directed

grooming between bees with and without mirror access. In Experiment 1 (n=12), without

prior mirror exposure, bees did not increase their total grooming activity, although they

did groom for a significantly higher proportion of time while facing the mirror. In

Experiment 2 (n=9), bees with prior mirror exposure showed significantly higher total

grooming activity than controls, an effect driven by an increase in grooming while facing

the mirror. However, the critical test for MSR, a direct comparison of grooming while

facing the mirror versus facing away from it, was not significant in either experiment's

mirror group. Our findings suggest the observed behaviors are not indicative of MSR.

Instead, they are better explained by simpler mechanisms, such as an orienting

response to a novel stimulus (Exp 1), generalized agitation from a perceived social

stimulus (Exp 2), or tactile irritation from the mark itself, an interpretation consistent with

findings in other invertebrates. This exploratory research, limited by small sample sizes,

provides preliminary yet inconclusive evidence. We discuss our results in the context of

key methodological prerequisites for the mark test, such as mark salience, and the

broader possibility that MSR-like behaviors reflect a learned perceptual-motor skill

rather than an abstract self-concept. Thus, this study contributes not by providing a

definitive answer, but by highlighting these critical challenges and establishing a

rigorous framework for future investigations designed to disentangle these alternative

hypotheses.

Introduction

39 The capacity for self-recognition, the ability to identify oneself as a distinct entity

40 separate from the environment and others, has long been a central topic in cognitive

science. Mirror self-recognition (MSR), typically assessed using the mark test, is a

primary experimental benchmark for this capacity in non-human animals and is widely

regarded as a key indicator of self-awareness and potentially other higher-order

cognitive functions [1, 2]. Since Gallup's seminal study in 1970, which first provided

experimental evidence of MSR in chimpanzees [3], this phenomenon has attracted

substantial interest across multiple disciplines, including primatology, comparative

psychology, and neuroscience. MSR is primarily assessed using the mark test. This

classic method involves applying a mark to a body part invisible to the subject without a

mirror and observing whether self-directed behaviors (e.g., touching, inspecting) of the

marked area increase in the presence of a mirror compared to its absence [1, 2].

Although MSR was initially considered a hallmark of great apes and humans, a large-

scale study on chimpanzees revealed that MSR is a developmental phenomenon, not

universally present in all individuals [4]. Subsequent research has reported MSR or

MSR-like behaviors in a select group of other mammals and birds, including bottlenose

dolphins [5], Asian elephants [6], and Eurasian magpies [7]. These species often,

though not exclusively, possess relatively large brain capacities and exhibit complex

social structures [8, 9]. However, the interpretation of these findings, particularly in non-

primate species, remains a subject of considerable debate and methodological scrutiny

[10, 11]. Recent research has further explored the phylogenetic breadth of MSR,

questioning 'how simple a nervous system can support MSR?'. In this context, the

success reported in the cleaner wrasse (*Labroides dimidiatus*), a fish species [12],

received significant attention by suggesting that MSR capability is not necessarily

limited to large brains or specific lineages, although this interpretation itself remains

highly contested and subject to alternative explanations [13, 14]. Furthermore, studies in

mice have shown that the activation of hippocampal neuronal ensembles by visuotactile

integration promotes mirror-induced self-directed behavior, suggesting a role for the

hippocampus in MSR in mammals [15].

Evidence for MSR in invertebrates is particularly limited and often debated. For

instance, a study on three ant species (*Myrmica sabuleti*, *Myrmica rubra*, and *Myrmica*

ruginodis) reported mark-directed behavior in adults but not in very young subjects,

requiring cautious interpretation [16]. Research on paper wasps (*Polistes metricus*)

found increased exploratory behaviors towards mirrors but did not yield conclusive

evidence of MSR [17].

Among invertebrates, arguably the most extensively studied group for complex

cognition is the cephalopods (octopus, squid, and cuttlefish). However, a recent

comprehensive review concluded that evidence for mirror self-recognition (MSR) in

cephalopods remains weak and contested [18]. While some species like the oval squid

show strong exploratory behaviors such as touching their mirror image [19], a

preliminary attempt to apply the mark test to the common octopus found that the

observed mark-directed grooming was likely triggered by the tactile sensation of the

mark itself, rather than visual self-recognition [20].

Currently, the ghost crab (*Ocypode quadrata*) is reported as potentially one of the

phylogenetically lowest organisms showing evidence of MSR [21], though again,

interpretation requires careful consideration of alternative explanations.

This approach, focusing on an invertebrate, directly addresses a foundational critique

within comparative psychology. For decades, influential voices have called for the

investigation of fundamental cognitive processes at widely separated points in the

phylogenetic scale to truly test the generality of psychological principles, arguing against

the field's historical over-reliance on a few mammalian models [22]. Against this

complex and evolving background, the present study investigates MSR in the western

honeybee (*Apis mellifera*). The honeybee is an insect phylogenetically situated between

crustaceans and vertebrates and is renowned for its remarkable cognitive abilities

despite possessing a brain containing only about one million neurons [23]. Honeybees

exhibit a surprisingly diverse suite of complex cognitive functions. Indeed, a long

tradition of research has systematically analyzed honeybee learning by drawing direct

comparisons to vertebrate performance, establishing the honeybee as a powerful model

for the comparative analysis of cognition [24, 25]. This comparative work has been

extensively reviewed in the context of visual cognition and perception [26, 27]. Their

capacity for concept learning is particularly notable [28]. For instance, their ability to

learn the abstract concepts of 'sameness' and 'difference' has been reported [29, 30],

and the topic has been recently reviewed [31]. A key factor underpinning this ability is a

robust visual working memory. While an early study reported this memory to last for 6.5

seconds [32], subsequent research has demonstrated that it can, under certain

conditions, persist for approximately 2.5 minutes [30]. They can also learn to apply two

abstract concepts simultaneously [33], learn concepts of oddity/nonoddity [34, 35],

relative size [36], and above/below spatial relationships [37]. Their cognitive suite further

includes maze navigation [38], numerical cognition including understanding zero and

parity [39, 40], perception of visual illusions [41], and flexible behavioral strategies in

learning tasks [42]. This suite of abilities has also inspired research in robotics and

artificial intelligence, such as the development of neuromorphic vision systems [43].

They also exhibit emotion-like states interpreted as "pessimistic decision-making" [44].

In social contexts, evidence suggests sophisticated social cognitive capabilities, such as

learning from conspecifics using visual cues [45], as well as nestmate recognition using

olfactory and tactile cues [46], and remarkably, they have even been reported to

discriminate human faces [47]. Such complex visual recognition is likely important for

recognizing conspecifics [48] and adjusting behavior accordingly, particularly during

foraging [49].

Particularly relevant to MSR is the finding by Stach & Giurfa (2001) that honeybees can

generalize learned visual patterns to their mirror images [50]. This suggests a basic

ability to process mirror image information, a potential prerequisite for MSR. However,

their study focused on generalization in pattern recognition, a different cognitive process

from recognizing one's own image in a mirror (MSR). The present study aims to test

whether this mirror image processing ability extends to the recognition of the self-image

using the mark test.

Investigating MSR in honeybees holds significant implications. Firstly, if MSR is

confirmed in an insect with a relatively simple nervous system, it would strongly suggest

that higher cognitive functions like self-recognition do not necessarily require large

brains or specific structures (e.g., neocortex), thereby deepening our understanding of

the evolutionary origins and neural basis of self-recognition [8]. It might eventually open

possibilities for identifying the neural circuits involved in MSR within the relatively

tractable honeybee nervous system [23, 24]. Secondly, findings on honeybee MSR,

highlighting complex cognition within a relatively simple nervous system [24], could

contribute not only to psychology and biology but also inform robotics and artificial

intelligence research [51]. For example, computational approaches have already

demonstrated the feasibility of functionally imitating honeybee concept learning

processes with simple neural networks inspired by the insect brain [52], as well as with

other computational frameworks such as Vector Symbolic Architectures [53], potentially

shedding light on the foundations required for capabilities like self-recognition.

We chose honeybees over paper wasps for several reasons. First, honeybees exhibit a

higher degree of social complexity [54]. This is a key consideration, as a prominent

theory for the evolution of self-recognition, the social intelligence hypothesis, posits that

the cognitive demands of complex social life are a primary driver for the evolution of

self-awareness [9, 55, 56]. Indeed, strong evidence from paper wasps suggests that

selection for individual recognition can lead to the evolution of specialized cognitive

abilities, such as dedicated face learning [57]. This proposed link between sociality and

mirror-directed behavior has also been observed in cephalopods. For example, highly

social species (squid) have been shown to have strong interest in their mirror image,

whereas asocial species (octopus) tend to show little or no reaction, suggesting that the

cognitive pressures of social life may be a key factor in the evolution of such behaviors

[58]. Therefore, highly social insects like honeybees represent a critical test case for the

taxonomic limits of this hypothesis. Honeybee colonies comprise tens of thousands of

individuals (compared to tens or hundreds in independent-founding paper wasps) and

feature distinct morphological and behavioral caste differentiation (queen, workers,

drones) [46, 54]. They have developed sophisticated social systems, including

advanced communication via the waggle dance [46] and collective defense behaviors

like heat-balling [59]. Such a complex social environment likely imposes strong selection

pressure on the ability to accurately discriminate self from others (nestmates, non-

nestmates, predators, etc.) and adjust behavior accordingly, potentially favoring self-

related cognitive abilities. Efficient functioning of a system where tens of thousands live

closely, dividing and coordinating diverse tasks (e.g., foraging, brood care, nest

construction/maintenance/defense, cleaning, etc.), arguably requires individuals to

differentiate, to some extent, their own state and actions from those of others. Mistaking

reflections (e.g., from water surfaces or dew) during activities outside the hive could be

maladaptive, wasting energy and time [60]; thus, the ability to distinguish their own

reflection from other individuals could be adaptive. Second, as mentioned earlier,

various cognitive abilities potentially related to MSR expression, such as the ability to

recognize and learn from conspecifics using visual cues [45], complex visual pattern

recognition [47], mirror-image generalization [50], and concept learning [29-31, 33-37,

42, 43, 49], have already been reported in honeybees, suggesting a higher potential for

MSR presence compared to species where these prerequisite abilities are less

documented.

It is noteworthy that previous studies have used mirrors (on the floor of experimental

tunnels or during flight over water) to investigate how honeybees use optic flow from the

ground for altitude control [60, 61]. These studies examined flight behavior changes

when visual cues were manipulated using horizontal mirrors and differ significantly in

purpose (investigating visually guided flight, not MSR via self-directed behavior facing a

vertical mirror) and experimental setup from the present study. This study aimed to

determine whether honeybees exhibit mirror self-recognition (MSR) by applying the

classic mark test. Furthermore, by exploring a species with a miniature nervous system,

this research also addresses the important call to investigate the cognitive limits of

animals, which is as crucial as demonstrating their capabilities for understanding the

diversity and evolution of cognition [62]. We hypothesized that if honeybees possess

MSR capabilities, subjects exposed to a mirror would show increased self-directed

grooming towards a mark placed on their head, compared to control subjects without a

mirror. To investigate this hypothesis, we compared the frequency and duration of head-

directed grooming directed at a marked area on the head between bees presented with

a mirror (mirror group) and those without (no-mirror/sham mirror group). By applying the

same classic paradigm to a phylogenetically distant invertebrate, this study also

provides an opportunity to explore the potential for the convergent evolution of complex

cognitive traits [22, 25, 63] such as self-recognition. Furthermore, to examine the effect

of prior mirror experience, we conducted Experiment 1 (without pre-exposure) and

Experiment 2 (with pre-exposure).

Materials and methods

Animals

196 Foraging worker bees (*Apis mellifera*) were collected as needed from hives on the

197 rooftop of Mita International School of Science (Tokyo, Japan) as they departed from

198 the hive entrance. Collected bees were therefore foragers, though precise age was not

199 controlled. The sample sizes were smaller than initially planned due to colony collapses

200 during the experimental periods, which is a notable limitation of this study.

Experiment 1: Mark test (no prior mirror exposure)

Subjects

203 Thirteen worker bees were subjected to the experimental procedure. They were

204 randomly assigned to the Mirror group (n = 7; Ss 1, 2, 5, 6, 9, 10, 13) or the Control

205 group (n = 6; Ss 3, 4, 7, 8, 11, 12). (As noted later and in the Results section, the final

206 number of subjects analyzed was reduced due to marking failures, poor recovery from

207 anesthesia, or abnormal behavior during the experiment.)

Apparatus

Transparent acrylic boxes served as individual test boxes for observation. For subjects

up to S8, the box size was 13.5 cm L × 9.7 cm W × 7.1 cm H; for S9 onwards, a slightly

smaller box (10.0 cm L × 10.2 cm W × 4.6 cm H) was used. This change was

implemented to improve visibility for subsequent video analysis. A horizontal platform

served as the base for the test box, and the platform was enclosed by white paper

(overall enclosure size: approx. 52 cm L × 55 cm W × 38 cm H) to block external visual

disturbances and avoid reflections on the test box (Fig 1).

216

217 Fig 1. Experimental setup showing a test box on a horizontal platform, enclosed

218 by white paper to block external visual disturbances and avoid reflections on the

219 test box.

221 Marking procedure

Approximately 10 bees were captured near the hive entrance in a clear plastic bag. To

induce a temporary hypothermic state, the bag containing the bees was placed in a

freezer (approx. -18°C) until they became immobile (approx. 4 min 30 s). The

anesthetized bees were then removed from the bag and each was individually marked

on the clypeus. A small amount of white correction fluid (Pentel Co., Ltd. XEZL1-W) was

applied to the tip of a toothpick and used to place a small mark on the bee's head,

specifically on the clypeus below the compound eyes and above the mandibles (Fig 2),

adopting a similar mark placement to that employed in tests with paper wasps [17]. This

marking procedure presented technical challenges due to the small size of the bees and

their rapid arousal. A number of failures occurred, such as incorrect mark size or

placement, or application of fluid to unintended areas (e.g., antennae, mouthparts).

Subjects judged to have been improperly marked were excluded before proceeding.

This cooling procedure, a common method for temporarily immobilizing honeybees for

manipulation, was used to minimize invasiveness while allowing for accurate marking.

However, it is important to acknowledge that the procedure itself, particularly the

potential for physical irritation or olfactory cues from the marking fluid, could influence

subsequent behavior and thus requires careful interpretation of the results.

239

240   **Fig 2. Honeybee (*Apis mellifera*) with a white mark applied to the clypeus (below**

241   **the compound eyes and above the mandibles) for the mark test.**

Experimental procedure

After the marking procedure, bees were allowed to recover. Once a subject began

moving, it was fed a small amount of 50% (w/w) sucrose solution via pipette to provide

energy. Each bee was then placed individually into a transparent test box enclosed by

white paper on all sides except the top. For each trial, the test box containing the

individual that appeared most physically intact and exhibited the most normal recovery

behavior was selected. This selected box was placed onto a horizontal platform, and a

5-minute acclimatization period commenced when the bee inside started walking

spontaneously. After acclimatization, one of two conditions was applied. For the Mirror

group, an acrylic mirror (commercially available, cut to fit the side) was inserted

vertically along one of the side walls of the test box; specifically, a short side wall for the

larger rectangular box (subjects up to S8) and the slightly longer side wall for the

smaller box (subjects from S9 onwards). For the Control group, subjects up to S8

received no mirror. For subjects S9 onwards, a 'sham mirror' (the back of a mirror

covered with non-reflective white cardstock) was inserted to control for the potential

effects of the insertion procedure and to provide a non-reflective visual stimulus,

ensuring similar physical manipulation between groups. Behavior was video-recorded

from directly above the test box using an iPad (Apple Inc.) for 30 minutes following the

acclimatization period. Note that during recovery from anesthesia, a few subjects died

or exhibited prolonged immobility, and these were excluded from the experiment.

262 Experiment 2: Mark test (prior mirror exposure)

263 Purpose

264 Given that Experiment 1 yielded no clear MSR evidence, and considering the

265 importance of prior mirror experience often highlighted in MSR research [2], Experiment

266 2 aimed to test the effect of pre-exposure under more controlled conditions (consistent

267 sham mirror control, extended observation time, uniform box size).

268 Subjects

269 Nine new worker bees were subjected to the procedure. They were randomly assigned

270 to the Mirror group (n = 5; Ss 1, 2, 4, 6, 9) or the Sham Mirror group (n = 4; Ss 3, 5, 7,

271 8). (As in Experiment 1, the number analyzed reflects exclusions due to the criteria

272 mentioned above).

Apparatus

To maintain consistency and improve visibility during video analysis, the same smaller

transparent acrylic boxes (10.0 cm L × 10.2 cm W × 4.6 cm H) were used for initial

capture, the pre-exposure session, and as test boxes for observation. The indoor setup

for Experiment 2 consisted of a cardboard box (32 cm L × 40 cm W × 26 cm H) with an

open top and lined entirely with white paper on the interior walls to reduce reflections

and provide a uniform background. The cardboard box was placed on a horizontal

platform, and a portion of the box top was covered with white paper to further minimize

reflections. Additionally, one or two white paper enclosures, each used in Experiment 1

to cover a test box, were placed inside the cardboard box.

Pre-exposure

Pre-exposure was conducted in two stages: at the hive and subsequently in the

laboratory.

(1) **At the hive.** A few days before starting to collect experimental subjects, a large

mirror (reflective surface: 28.5 cm × 38.5 cm) was placed in front of the entrance of the

target colony's hive (Fig 3). This provided bees entering and leaving the hive with

natural exposure opportunities to their moving reflections. This procedure was intended

to allow for potential associative learning between the bees' own movements and the

movements observed in the mirror reflection, which might facilitate recognition of the

mirror image as their own in a more natural context, but its effectiveness remains

speculative. Additionally, the exact duration and degree of attention paid by each

individual were not recorded or controlled.

(2) **In the laboratory.** The second stage of pre-exposure was conducted in the

laboratory to habituate the bees to the presence of a mirror within the testing context,

thus minimizing responses to the mirror as a novel object during the subsequent

experimental phase. Bees were individually captured in the experimental boxes in front

of the hive with the large mirror. To eliminate the potential impact of the mirror insertion

process on the bees' behavior, these experimental boxes, each containing a bee, were

then directly placed inside the white paper enclosures with a mirror pre-installed on one

side, within the indoor experimental setup for an additional pre-exposure session in the

lab. Bees in both the Mirror and Sham Mirror groups had a mirror placed along one side

wall (within the experimental box). We aimed for counterbalancing; however, the small

sample sizes limited its full implementation. The mirror position remained consistent for

each individual between the pre-exposure and testing periods. They were allowed to

explore freely for 20 minutes.

Fig 3. A large mirror (reflective surface: 28.5 cm × 38.5 cm) for Experiment 2 pre-exposure, placed in front of the colony hive.

Experimental procedure

The overall procedure for Experiment 2 generally followed that of Experiment 1, but the observation time was extended to 40 minutes to gather more behavioral data. The following steps were performed after the pre-exposure phase:

1. After the 20-minute pre-exposure session, bees were removed and anesthetized by cooling as in Experiment 1 (approx. 4 min 30 s).
2. Marking was performed as in Experiment 1. Exclusions due to marking failures occurred here as well.
3. Upon recovery, bees were individually fed 50% (w/w) sucrose solution and placed in transparent, clean test boxes. To avoid any behavioral changes potentially induced by inserting either the mirror or the sham mirror during the experiment, the experimental setup already included individual white paper enclosures, each with either a mirror (Mirror group) or a sham mirror (Sham

Mirror group) pre-installed along one longer side. Then, one or two transparent

test boxes, each containing a bee, were placed inside the corresponding white

paper enclosures for each trial.

4. The bee in its test box was observed, and its behavior was video-recorded for 40

minutes starting from when it began walking spontaneously. In Experiment 2, in

order to observe the effects of prior mirror experience in more detail, unlike

Experiment 1, video recording was started immediately after the bee began

walking spontaneously. The aim was to capture a wider range of behavioral

changes by including the behavior immediately after recovery from anesthesia.

To improve efficiency, two subjects were sometimes tested concurrently in

separate boxes and their respective white paper enclosures, but they remained

visually and physically isolated from each other. Subjects showing poor recovery

from anesthesia were excluded from analysis.

337 Data collection and behavioral analysis

After the experiment, recorded videos were observed by two independent observers

(RA and MM, the second and third authors, who were also the experimenters). To

ensure consistency and assess inter-observer reliability, all video analysis was

conducted by the two observers (RA and MM). This process included obtaining

measurements from videos played back at slow speed as needed to reliably capture

fast movements. The following behavioral metrics were measured.

344 Grooming

Grooming was defined as any action in which the forelegs touched or scrubbed any part

of the head region, including the clypeus, compound eyes, and antennae. During

analysis, it was not possible for observers to reliably distinguish between actions

specifically directed at the paint mark and more general grooming actions directed at

nearby areas, such as the antennae. Therefore, all recorded grooming data represent

general head-directed grooming rather than exclusively mark-directed actions. Total (T)

frequency and duration (in seconds, measured with 0.01 second precision) were

recorded. Due to the fast movements of some grooming behaviors, which made

accurate measurement difficult at normal playback speed, videos were slowed down as

needed for reliable measurement.

Body Orientation during Grooming

The bee's body orientation relative to the mirror (or sham mirror/hypothetical mirror

location for some controls) was recorded at the moment grooming occurred.

Orientations were coded as Facing (F) if the bee's body axis was within approximately

±90° of perpendicular to the mirror surface; Away (A), with the body axis outside the

±90° range; or Not facing or away (N), for orientations that were roughly sideways or

ambiguous (Fig 4). For Control group subjects up to S8 in Exp 1, a mirror position was

randomly assumed to determine orientation. For all subsequent control and sham mirror

subjects, the inserted sham mirror served as the reference.

Fig 4. Classification of honeybee body orientation relative to the mirror during

grooming.*

* F: Facing the mirror, A: Away from the mirror, N: Not facing or away.

The "N" category was included to avoid forcing ambiguous observations into the "F" or

"A" classifications, thereby ensuring a more rigorous application of these criteria.

However, data for the "N" category proved to have low inter-observer reliability

(Pearson's r: -0.232 to 0.968), likely due to its infrequent occurrence. Therefore, these

data were excluded from the main behavioral analyses and are presented only as

supplementary information. The main statistical analyses focused on the more reliable

Total (T), Facing (F), Away (A), and Facing/Total ratio (F/T) metrics.

Statistical analysis

For each individual, the values measured by the two observers were averaged for

analysis. Inter-observer reliability was assessed using Pearson's correlation coefficient

(Pearson's r) for each metric (T, F, A, N frequency and duration). Consistency between

duration and frequency metrics was also assessed using Pearson's r. Between-group

comparisons (Mirror vs. Control/Sham Mirror) for T, F, A, and N grooming frequency

and duration were performed using Welch's t-test, selected to accommodate potentially

unequal variances and small sample sizes. The proportion of mirror-facing grooming

(F/T, where T = F+A+N) was calculated as an index of directional bias and compared

between groups (Welch's t-test). Within-group comparisons (F vs. A frequency and

duration) in the Mirror group were performed using paired t-tests. The statistical

significance level was set at $\alpha = .05$. Given the small sample sizes, exact p-values are

reported, and effect sizes (Cohen's d) are provided for context, but all statistical results

should be interpreted with caution due to limited power.

Individual analysis (binomial test)

To examine individual variation in addition to group-level trends, data from each

observer for each individual were analyzed separately. The number of grooming bouts

Facing (F) versus Away (A) from the mirror/sham was compared using a two-tailed

binomial test ($\alpha = .05$) to assess significant directional bias. Raw count data from each

observer were used as the test requires integer values.

Ethical considerations

All experimental procedures were conducted with the highest regard for animal welfare.

We aimed to minimize the number of bees used, acknowledging challenges in handling

foraging bees from outdoor hives, including recovery from cooling anesthesia, marking

precision, or behavior. In Experiment 1, approximately 10 bees were captured per

session in a plastic bag, an approach that sometimes led to necessary exclusions. In

Experiment 2, bees were introduced individually into experimental boxes, minimizing

unused individuals. Cooling anesthesia, when applied, was limited to the minimum

necessary (approx. 4 min 30 s) for safe handling and rapid recovery. Marking was

performed swiftly to minimize stress. Bees showing prolonged immobility or abnormal

behavior were immediately excluded.

Results

Inter-observer reliability and correlation between duration and frequency

In both experiments, high inter-observer reliability was found for all grooming behavior

measurements, including total and orientation-specific frequency and duration (T, F, A)

and the facing/total (F/T) ratio. In Experiment 1, Pearson's r was ≥ .960 for duration

metrics and ≥ .913 for frequency metrics. Similarly, in Experiment 2, r was ≥ .898 for

duration and ≥ .918 for frequency. Strong positive correlations were also observed

between duration and frequency for each metric (T, F, A, and F/T) within each

experiment (Experiment 1: r ≥ .984; Experiment 2: r ≥ .890). This indicates that duration

and frequency were closely related measures of grooming behavior, reflecting similar

underlying tendencies. Therefore, we report statistical results for both metrics, as the

trends were consistent across both.

Experiment 1: Mark test (no prior mirror exposure, 30-min observation)

To visually represent the results of Experiment 1, the following box plots are provided:

Fig 5 shows the grooming duration for T, F, A, and N; Fig 6 shows the F/T ratio for

grooming duration; Fig 7 shows the grooming frequency for T, F, A, and N; and Fig 8

shows the F/T ratio for grooming frequency. Of the 13 bees subjected to the procedure,

S13 was excluded from analysis due to abnormal behavior (e.g., prolonged immobility)

during the video recording. Data from the remaining 12 bees (Mirror group n = 6,

Control group n = 6) were analyzed. Between-group comparisons were performed using

Welch's t-test.

Fig 5. Box plots showing grooming duration (seconds) for T, F, A, and N for the

Mirror group and Control group in Experiment 1. *

435   * T: Total, F: Facing the mirror, A: Away from the mirror, N: Not facing or away.

438 **Fig 6. Box plots showing the F/T ratio of grooming duration for the Mirror group**

439 **and Control group in Experiment 1. ***

440 * F: Facing the mirror, T: Total.

Fig 7. Box plots showing grooming frequency (counts) for T, F, A, and N for the

Mirror group and Control group in Experiment 1. *

* T: Total, F: Facing the mirror, A: Away from the mirror, N: Not facing or away.

Fig 8. Box plots showing the F/T ratio of grooming frequency for the Mirror group and Control group in Experiment 1. *

* F: Facing the mirror, T: Total.

Grooming duration

Within-group (Mirror, n = 6): No significant difference between mean duration Facing (M = 5.997, SD = 8.219) and Away (M = 4.941, SD = 8.069) from the mirror (Paired $t(5)$ = 1.864, p = .121, d = 0.761, 95% CI for mean difference [-0.400, 2.512], mean difference = 1.056 seconds). Within-group (Control, n = 6): No significant difference between mean duration Facing (M = 10.724, SD = 16.722) and Away (M = 10.189, SD = 14.815) from

the hypothetical/sham mirror (Paired t(5) = 0.558, p = .601, d = 0.228, 95% CI for mean

difference [-1.930, 3.000], mean difference = 0.535 seconds). Between-group (Total):

No significant difference between Mirror (M = 11.218, SD = 16.276) and Control (M =

22.518, SD = 31.848) groups (Welch's t(7.445) = -0.774, p = .463, d = -0.447, 95% CI

for mean difference [-45.413, 22.813], mean difference = -11.300 seconds). Between-

group (Facing): No significant difference between Mirror (M = 5.997, SD = 8.219) and

Control (M = 10.724, SD = 16.722) groups (Welch's t(7.283) = -0.621, p = .553, d = -

0.359, 95% CI for mean difference [-22.574, 13.119], mean difference = -4.728

seconds). Between-group (Away): No significant difference between Mirror (M = 4.941,

SD = 8.069) and Control (M = 10.189, SD = 14.815) groups (Welch's t(7.727) = -0.762,

p = .469, d = -0.440, 95% CI for mean difference [-21.228, 10.732], mean difference = -

5.248 seconds). Between-group (Facing/Total Ratio): The Mirror group (M = .649, SD =

.243) showed a significantly higher proportion of grooming time facing the mirror

compared to the Control group (M = .371, SD = .158) (Welch's t(8.576) = 2.354, p =

.044, d = 1.359, 95% CI for mean difference [.009, .549], mean difference = .279).

Grooming frequency

Within-group (Mirror, n = 6): A non-significant trend for more frequent grooming Facing

(M = 33.167, SD = 40.163) than Away (M = 26.333, SD = 40.595) from the mirror

(Paired t(5) = 2.165, p = .083, d = 0.884, 95% CI for mean difference [-1.280, 14.946],

mean difference = 6.833). Within-group (Control, n = 6): No significant difference

between frequency Facing (M = 66.417, SD = 95.439) and Away (M = 58.083, SD =

73.755) from the hypothetical/sham mirror (Paired t(5) = 0.865, p = .427, d = 0.353, 95%

CI for mean difference [-16.427, 33.093], mean difference = 8.333). Between-group

(Total): No significant difference between Mirror (M = 61.750, SD = 82.049) and Control

(M =134.167, SD = 170.019) groups (Welch's t(7.209) = -0.940, p = .378, d = -0.542,

95% CI for mean difference [-253.592, 108.759], mean difference = -72.417). Between-

group (Facing): No significant difference between Mirror (M = 33.167, SD = 40.163) and

Control (M = 66.417, SD = 95.439) groups (Welch's t(6.717) = -0.787, p = .458, d = -

0.454, 95% CI for mean difference [-134.068, 67.568], mean difference = -33.250).

Between-group (Away): No significant difference between Mirror (M = 26.333, SD =

40.595) and Control (M = 58.083, SD = 73.755) groups (Welch's t(7.775) = -0.924, p =

.383, d = -0.533, 95% CI for mean difference [-111.409, 47.909], mean difference = -

31.750). Between-group (Facing/Total Ratio): The proportion of grooming frequency

Facing the mirror was significantly higher in the Mirror group (M = .660, SD = .232) than

the Control group (M = .378, SD = .158) (Welch's t(8.837) = 2.462, p = .036, d = 1.422,

95% CI for mean difference [.022, .542], mean difference = .282). Note on procedural

variations in Exp 1 Control: Additional analyses (Welch's t-tests) confirmed that changes

in experimental box size (used for Ss up to S8 vs. S9 onwards) and the

presence/absence of a sham mirror in the Control group (no mirror up to S8 vs. sham

mirror S9 onwards) did not significantly affect the measured grooming metrics (ps >

.05), suggesting these procedural variations did not introduce significant confounds

within the limits of this small sample analysis. However, the definition of F and A for

each bee in the Control group up to S8 was based on a hypothetical mirror location,

which was randomly determined for each individual by a separate coin toss, whereas

from S9 onwards it was based on the sham mirror.

Summary for Experiment 1

Overall, Exp 1 showed no significant difference in the absolute amount of grooming between groups, but the Mirror group showed a significantly higher proportion of grooming frequency facing the mirror. However, since both the total grooming duration and frequency did not increase and there was no significant difference between facing and away grooming within the mirror group, the observed proportional shift provides limited support for MSR.

Experiment 2: Mark test (prior mirror exposure, 40-min observation)

To visually represent the results of Experiment 2, the following box plots are provided: Fig 9 shows the grooming duration for T, F, A, and N; Fig 10 shows the F/T ratio for grooming duration; Fig 11 shows the grooming frequency for T, F, A, and N; and Fig 12 shows the F/T ratio for grooming frequency. Of the 9 bees subjected to the procedure, data from all 9 bees (Mirror group n = 5, Sham Mirror group n = 4) were analyzed after exclusions prior to the test phase due to marking failure, poor recovery, etc. Note that the very small sample sizes (n = 4-5 per group) significantly limit statistical power and necessitate caution in interpreting these results. Between-group comparisons were performed using Welch's t-test.

Fig 9. Box plots showing grooming duration (seconds) for T, F, A, and N for the

Mirror group and Sham Mirror group in Experiment 2. *

526   * T: Total, F: Facing the mirror, A: Away from the mirror, N: Not facing or away.

Fig 10. Box plots showing the F/T ratio of grooming duration for the Mirror group

and Sham Mirror group in Experiment 2. *

* F: Facing the mirror, T: Total.

Fig 11. Box plots showing grooming frequency (counts) for T, F, A, and N for the

Mirror group and Sham Mirror group in Experiment 2. *

* T: Total, F: Facing the mirror, A: Away from the mirror, N: Not facing or away.

Fig 12. Box plots showing the F/T ratio of grooming frequency for the Mirror

group and Sham Mirror group in Experiment 2. *

541   * F: Facing the mirror, T: Total.

Grooming duration

Within-group (Mirror, n = 5): No significant difference between mean duration Facing (M

= 16.951, SD = 8.140) and Away (M = 11.399, SD = 9.017) from the mirror (Paired t(4)

= 1.110, p = .329, d = 0.496, 95% CI for mean difference [-8.334, 19.438], mean

difference = 5.552 seconds). Within-group (Sham Mirror, n = 4): No significant

difference between mean duration Facing (M = 5.741, SD = 2.852) and Away (M =

5.888, SD = 3.613) from the sham mirror (Paired t(3) = -0.050, p = .963, d = -0.025,

95% CI for mean difference [-9.369, 9.077], mean difference = -0.146 seconds).

Between-group (Total): Significantly longer duration in the Mirror group (M=28.549, SD

= 12.839) than the Sham Mirror group (M = 12.748, SD = 2.806) (Welch's t(4.471) =

2.673, p = .050, d = 1.600, 95% CI for mean difference [0.051, 31.553], mean difference

= 15.802 seconds). Between-group (Facing): Significantly longer duration Facing the

mirror in the Mirror group (M = 16.951, SD = 8.140) than the Sham Mirror group (M =

5.741, SD = 2.852) (Welch's t(5.160) = 2.867, p = .034, d = 1.743, 95% CI for mean

difference [1.253, 21.167], mean difference = 11.210 seconds). Between-group (Away):

No significant difference between Mirror (M = 11.399, SD = 9.017) and Sham Mirror (M

= 5.888, SD = 3.613) groups (Welch's t(5.473) = 1.247, p = .263, d = 0.764, 95% CI for

mean difference [-5.558, 16.581], mean difference = 5.512 seconds). Between-group

(Facing/Total Ratio): No significant difference between Mirror (M = .598, SD = 0.177)

and Sham Mirror (M = .456, SD = 0.206) groups (Welch's t(6.008) = 1.100, p = .314, d =

0.752, 95% CI for mean difference [-0.175, 0.460], mean difference = .143).

Grooming frequency

Within-group (Mirror, n = 5): A non-significant trend for more frequent grooming Facing

(M = 85.600, SD = 34.976) than Away (M = 57.500, SD = 33.227) from the mirror

(Paired t(4) = 1.413, p = .231, d = 0.632, 95% CI for mean difference [-27.125, 83.325],

mean difference = 28.100). Within-group (Sham Mirror, n = 4): No significant difference

between frequency Facing (M = 31.750, SD = 16.454) and Away (M = 33.625, SD =

15.892) from the sham mirror (Paired t(3) = -0.129, p = .905, d = -0.065, 95% CI for

mean difference [-47.967, 44.217], mean difference = -1.875). Between-group (Total):

Significantly higher frequency in the Mirror group (M = 145.000, SD = 49.793) than the

Sham Mirror group (M = 72.625, SD = 13.066) (Welch's t(4.672) = 3.119, p = .029, d =

1.875, 95% CI for mean difference [11.438, 133.312], mean difference = 72.375).

Between-group (Facing): Significantly higher frequency Facing the mirror in the Mirror

group (M = 85.600, SD = 34.976) than the Sham Mirror group (M = 31.750, SD =

16.454) (Welch's t(5.916) = 3.047, p = .023, d = 1.886, 95% CI for mean difference

[10.455, 97.245], mean difference = 53.850). Between-group (Away): No significant

difference between Mirror (M = 57.500, SD = 33.223) and Sham Mirror (M = 33.625, SD

= 15.892) groups (Welch's t(5.965) = 1.417, p = .207, d = 0.878, 95% CI for mean

difference [-17.412, 65.162], mean difference = 23.875). Between-group (Facing/Total

Ratio): No significant difference between Mirror (M = .589, SD = 0.139) and Sham

Mirror (M = .430, SD = 0.193) groups (Welch's t(5.318) = 1.383, p = .222), d = 0.966,

95% CI for mean difference [-0.131, 0.449], mean difference = .159.

Summary for Experiment 2

In Experiment 2, bees with prior mirror exposure (Mirror group) exhibited significantly

higher total grooming duration and frequency, and significantly more grooming bouts

facing the mirror, compared to the Sham Mirror group. Specifically, the Mirror group

showed a significant increase in grooming bouts facing the mirror (Facing), but no

significant difference in grooming bouts away from the mirror (Away). However, there

was no significant difference in the Facing/Total ratio between the groups. Importantly,

however, this increase in grooming did not result in a statistically significant bias facing

the mirror when compared to grooming away from it within the Mirror group, nor did the

proportion of mirror-facing grooming differ significantly between groups.

595   Individual analysis (binomial test)

596   Directional bias in grooming frequency at the individual level was assessed by

597   comparing F versus A frequencies for each observer per bee (two-tailed binomial test, α

598   = .05).

599   Experiment 1: Mirror group

600   For the Mirror group (n = 6), only S5 showed a significant Facing > Away bias according

601   to both observers (ps < .05). S1 showed this bias only for observer MM. No significant

602   bias was found for the other 4 bees by either observer.

603   Experiment 1: Control group

604   For the Control group (n =6), while some subjects showed significant biases (S3 (RA),

605   S11 (MM) showed Facing > Away), no individual showed a consistent directional bias

606   according to both observers.

607   Experiment 2: Mirror group

608   For the Mirror group (n = 5), S1 and S6 showed a significant Facing > Away bias

609   according to both observers (ps < .05). No significant bias was found for the other 3

610   bees by either observer.

611   Experiment 2: Sham Mirror group

612   For the Sham Mirror group (n = 4), S3 showed a significant Facing > Away bias

613   according to both observers (ps < .05). S8 showed a significant Away > Facing bias

614   according to both observers (ps < .05). S7 showed an Away > Facing bias only for

615   observer MM. The remaining bee (S5) showed no significant bias according to either

616   observer.

Summary of individual results

These individual-level analyses reveal that while group averages showed some

significant effects (especially in Exp 2), only a minority of subjects in the mirror groups

consistently directed significantly more grooming facing the mirror than away (1/6 or

~17% in Exp 1; 2/5 or 40% in Exp 2, based on agreement between observers).

Furthermore, the presence of significant biases (both facing and Away) in some

control/sham mirror subjects complicates simple interpretation based solely on

directional preference. These findings highlight pronounced individual variation and

suggest factors beyond MSR could influence the results, highlighting the need to look

beyond simple group averages when evaluating MSR potential.

Discussion

This study investigated the potential for mirror self-recognition (MSR) in the western

honeybee using the classic mark test paradigm. The core logic relied on comparing

head-directed grooming directed at a mark (invisible without a mirror) between subjects

with and without mirror access, thereby controlling for the baseline effects of the mark

itself. Any significant increase in mark-directed grooming specifically in the mirror

condition would suggest recognition of the marked self-image.

Interpretation of findings

Preliminary insights requiring cautious interpretation

Our findings provide preliminary insights into how honeybees respond to mirrors after

being marked, but the assessment regarding MSR potential requires considerable

caution. The results of Experiment 1 particularly highlight this need for caution. In this

experiment, a key indicator of MSR was absent, as the mirror group showed no

significant increase in the total amount of grooming compared to the control group. And

yet, the proportion of grooming directed towards the mirror (F/T ratio) was significantly

higher for both duration and frequency. This shift in proportion, without an increase in

the total quantity of grooming, can be explained by a simpler mechanism than MSR.

Specifically, it is possible that the bees exhibited an orienting response to the mirror as

a novel visual stimulus. This well-documented behavior, in which an organism's

attention is captured by a new object, would explain why they spent more time oriented

in its direction. Consequently, the baseline grooming behavior—likely prompted by the

physical irritation of the mark—was more likely to be observed coincidentally while the

bee was facing the mirror. This explanation, which favors a simpler, non-cognitive

explanation, is consistent with findings in other complex invertebrates. For instance, in a

preliminary mark test on the common octopus (*Octopus vulgaris*), subjects were found

to frequently explore the mark with their arms even in control trials without a mirror. The

study concluded that these mark-directed behaviors were likely driven by proprioceptive

stimuli from the mark itself, rather than by visual feedback from the mirror [20].

This alternative hypothesis is strongly supported by the finding that even within the

mirror group, there was no clear and significant preference for grooming while facing the

mirror over grooming while facing away. Therefore, the elevated F/T ratio cannot be

conclusively construed as a cognitive leap of self-recognition and may instead be an

artifact of the spatial overlap between attentional orientation and a pre-existing

grooming pattern.

In contrast, Experiment 2 revealed a different behavioral pattern after mirror pre-

exposure. Here, bees in the mirror group exhibited significantly higher total grooming

activity than the sham-mirror controls. While this increase might initially seem more

consistent with MSR, other key details suggest a different conclusion, as emphasized in

critical reviews of MSR research [13]. Crucially, the significant bias in the F/T ratio

observed in Experiment 1 disappeared in Experiment 2. Furthermore, as in Experiment

1, bees in the mirror group still did not groom significantly more when facing the mirror

compared to when facing away. This implies that while pre-exposure caused the mirror

to become a trigger for increased grooming, this behavior was not visually guided by the

reflection. Instead, the mirror image may have been perceived as a social stimulus—

that is, a conspecific. From this perspective, the increased grooming is not an act of

self-recognition, but rather a sign of the resulting heightened arousal, stress, or a social

response to this stimulus [e.g., 2, 9, 11, 58].

Taken together, the results from both experiments do not contradict each other; rather,

they form a consistent narrative that argues against MSR in honeybees. The initial

reaction to the mirror (Experiment 1) appears to be an attentional artifact, while the

subsequent reaction after mirror pre-exposure (Experiment 2) appears to be one of

generalized agitation. In neither case did the bees display the key criterion for MSR: the

specific, tool-like use of the mirror to guide actions toward an otherwise invisible part of

the body [e.g., 1-3]. Ultimately, determining whether an animal perceives its reflection

as self, another individual, or simply a novel visual stimulus remains a central challenge

in MSR research [10, 11]. Our results reinforce the interpretation that the observed

behaviors can be explained by simpler, non-cognitive mechanisms. This supports a

foundational debate in comparative psychology, where behaviors often attributed to

higher-order cognition, such as a 'Theory of Mind,' have been compellingly reinterpreted

as the products of associative learning [64], and is a recurring theme in insect cognition

where seemingly abstract abilities have been alternatively explained by lower-level

processes [30, 40, 42, 50, 52, 53, 62, 65-67].

Lack of clear within-group difference (facing vs. away)

A key observation requiring careful consideration is the general lack of a statistically

significant preference for grooming while facing the mirror versus grooming while facing

away from it within the mirror group in both experiments, particularly in Experiment 2

where overall grooming increased. Classic MSR, as demonstrated in chimpanzees [3],

involves the animal specifically using the mirror to direct actions towards the mark. The

absence of a clear bias towards mirror-facing grooming for the mark itself in honeybees

weakens the MSR interpretation. This is consistent with findings suggesting that some

seemingly complex cognitive tasks in bees, such as the learning of 'above/below' spatial

relationships [37], may be solved by employing stereotyped flight movements and

sequential inspection that transform the problem into a simpler discrimination task [42].

Similarly, numerosity estimation has been explained by a simple neural network coupled

with a sequential inspection strategy [67]. These precedents suggest the observed

grooming could be a byproduct of a simpler behavioral mechanism rather than genuine

self-recognition.

Substantial individual variation

The pronounced individual variation revealed by both group-level statistics and binomial

tests further complicates interpretation. In both experiments, only a minority of subjects

in the mirror groups consistently directed significantly more grooming towards the

mirror. Moreover, the occurrence of directional biases in some Control/Sham mirror

subjects suggests that factors unrelated to MSR could influence the results. This

heterogeneity highlights the need to look beyond simple group averages when

assessing the possibility of MSR and reinforces the importance of analyzing individual

response patterns [4, 13].

Alternative interpretations

Conspecific or novel stimulus recognition

A plausible alternative interpretation, as often discussed in MSR literature [11], is that

bees might have responded to the reflection as if it were a novel bee, a familiar bee with

an unusual mark, or simply an unfamiliar visual stimulus. This type of social responding

to a mirror image has been clearly demonstrated in other invertebrates. For example,

female cuttlefish (*Sepia officinalis*) display a specific body pattern called 'Splotch' almost

exclusively towards other females and their own mirror image, suggesting they perceive

their reflection as a same-sex conspecific [68]. This principle of seeking simpler, more

parsimonious explanations is a foundational tenet of comparative psychology, famously

expressed as Morgan's Canon [69], and is echoed in the rigorous analytical framework

required for comparing diverse species, which cautions against inferring complex

cognitive processes when simpler mechanisms cannot be ruled out [63, 64]. Indeed, the

challenge of disentangling abstract conceptual understanding from reliance on lower-

level mechanisms, such as graded perceptual similarity, has been a central debate in

the field ever since the pioneering work on relational matching in primates [70]. A prior

investigation into same-different learning argued that bees' performance in visual

matching tasks could instead be explained by reliance on lower-level mechanisms such

as graded perceptual similarity or other simple visual cues [30]. Applying this same

critical lens to the current study, it is imperative to consider that the observed mirror-

induced grooming may not reflect the cognitive leap of self-recognition, but rather a

manifestation of these more fundamental behavioral or perceptual responses. This

misinterpretation could trigger various behaviors such as increased antennation and

touching, potentially alongside social, defensive, or exploratory responses. Notably, a

similar study on paper wasps [17] found increased exploratory behaviors (antennation

and touching) but a decrease in self-directed grooming in response to mirrors,

suggesting these behaviors may serve different functions across insect species and not

necessarily all point towards MSR. The honeybee's grooming could be a displacement

activity triggered by the ambiguous stimulus of its reflection [16], rather than a mark-

directed action guided by self-recognition. The phenomenon of social responding to

mirrors, even after extensive habituation, is well-documented in primates such as

rhesus monkeys, where a simple change in the mirror's location can reinstate social

responses [71], and notably, self-recognition is absent in social pairings like mother-

infant or infant-infant groups [72]. The current study design, lacking a crucial control

condition where bees are presented with a live, marked conspecific, makes it difficult to

definitively distinguish this social/novelty-response hypothesis from the MSR

hypothesis.

Methodological prerequisites and the mark test

Beyond cognitive capacity, recent critiques emphasize that an animal's performance on

the mark test is contingent on fulfilling several methodological prerequisites [73]. First,

the animal must understand the properties of a mirror. Our pre-exposure in Experiment

2 was designed to facilitate this, and the resulting increase in grooming suggests the

mirror had an effect, but it is uncertain if this constitutes true mirror understanding.

Second, the mark itself must be salient enough to motivate investigation. While the

white paint was visually conspicuous to humans, its motivational salience for a

honeybee, which relies heavily on tactile and olfactory cues for grooming, is

questionable. Third, the animal must possess a repertoire of self-exploratory behavior.

While bees groom, it is unknown if this extends to visually-guided exploration of a

specific, marked body part. The failure to meet one or more of these criteria, as outlined

by Kakrada & Colombo (2022) [73], could explain our inconclusive results, independent

of whether honeybees possess a self-concept.

The perceptual-motor skill hypothesis

An even more parsimonious explanation is that MSR-like behavior may not reflect an

abstract self-concept at all, but rather a learned perceptual-motor skill [74]. Landmark

studies demonstrated that pigeons, through explicit operant conditioning that links

seeing a mark in the mirror with pecking their own body, can be trained to pass the

mark test [75, 76]. From this perspective, the increased grooming we observed in

Experiment 2 could be a low-level associative response to the novel visuotactile

contingency created by the mark and the mirror, rather than an act of self-recognition.

This critical stance is warranted even when bees demonstrate robust cognitive abilities

under rigorous conditions. For instance, even when honeybees showed evidence of

concept learning using more rigorous methods, such as a trial-unique procedure

designed to minimize simpler associative explanations, the potential for alternative,

lower-level mechanisms still required careful consideration [30]. This aligns with our

conclusion that seemingly complex behaviors can often be explained by simpler

mechanisms [22, 24, 25, 28, 30, 34, 35, 39, 40, 42, 43, 51, 52, 62, 64-67, 74-76].

Mark/anesthesia effects

While the comparison design aimed to control for the effects of the marking and

anesthesia procedures, their potential influence as confounding variables cannot be

entirely dismissed. For instance, the physical irritation or olfactory cues from the

marking fluid itself may have increased baseline grooming across all conditions. Future

research should incorporate dedicated no-mark control groups to fully isolate these

potential effects [1, 2, 10]. Consequently, the observed grooming behavior—even when

performed while facing the mirror—may not be a visually-guided attempt to touch the

mark, but rather a generalized response to the physical irritation or olfactory cues from

the marking fluid itself. The antennae are vital sensory organs, and any irritation nearby

could easily trigger cleaning behaviors directed at them, independent of any visual

information from the mirror.

Non-specific mirror effects

793 The vertical mirror might induce spatial confusion or general arousal unrelated to self-

794 recognition [cf. 60, 61]. Honeybees are known to be highly sensitive to visual

795 environmental changes [45].

Methodological limitations

Several methodological limitations must be considered when interpreting our findings.

Primarily, the small sample sizes (n=4−6 per group in Experiment 1 after exclusions,

and n=4−5 per group in Experiment 2) result in low statistical power, which significantly

affects the ability to draw firm conclusions. Additionally, the observers were not blind to

the experimental conditions, which introduces a potential for unconscious observer bias.

Perhaps the most significant methodological limitation, central to the interpretation of all
our findings, was the ambiguity in our primary dependent variable: the observers'
inability to reliably distinguish between grooming actions specifically targeting the mark
and more generalized head-grooming (e.g., of the antennae). This means we cannot
definitively conclude that the observed grooming, even when performed while facing the
mirror, was a contingent response to *seeing* the mark. Instead, it is highly plausible that
the underlying motivation for grooming stems from non-visual stimuli (e.g., tactile
irritation or olfactory cues) from the marking fluid itself. In this view, the mirror does not
serve as a tool for guided self-inspection, but rather as an arousing visual stimulus that
triggers an increase in this pre-existing, generalized grooming behavior. Because our
methodology could not falsify this simpler, non-cognitive hypothesis, the data do not
allow us to confirm that the bees were using the mirror as a tool for targeted self-
inspection. Given all these limitations, the statistical findings should be interpreted with
extreme caution, and we stress the necessity of future replications with larger sample
sizes and refined methodologies.

However, as has been argued for cephalopods, a failure to pass the classic mark test
does not necessarily confirm the absence of self-awareness [18]. The interpretation of
mirror-induced behaviors must consider the specific sensory world of the animal in
question. Honeybees [23, 24, 26], much like cuttlefish, are sensitive to polarized light.
Research on cuttlefish has shown that they react differently to their mirror image
depending on whether the reflection's polarization patterns are intact, suggesting they
are perceiving subtle cues in their reflection that go beyond simple shape and color [77].
This highlights the possibility that insects may also perceive their reflections in ways not

immediately obvious to human observers, adding another layer of complexity to the

interpretation of the mark test.

Comparison with other invertebrate studies

Our preliminary findings, particularly the increased grooming in Experiment 2, contrast

somewhat with studies on paper wasps [17] that found no clear MSR evidence,

although they reported increased exploratory behaviors like antennation and touching

the mirror. The wasp study employed crucial control conditions absent here (e.g., sham-

marked wasps with a mirror). The findings highlight the sensitivity of wasps to mirror

presence and marking, suggesting they perceive these alterations but respond

differently than classic MSR would predict (e.g., with more exploration rather than self-

directed grooming). Their results emphasize the difficulty in demonstrating conclusive

MSR in insects using this paradigm. Our results add to the sparse literature on potential

MSR in invertebrates like ants [16] and crabs [21], all of which present interpretational

challenges. While evidence for MSR in cephalopods is also contested, recent work

using a different paradigm—the rubber arm illusion—suggests that octopuses possess

a sophisticated sense of body ownership by integrating visual, tactile, and

proprioceptive information [78]. This finding indicates that complex forms of self-

perception may exist in invertebrates, which the classic mark test might not be suited to

detect [78]. Further supporting this notion, recent studies on bumblebees have shown

that they perceive the affordance of narrow gaps in relation to their own body size,

turning sideways to avoid collisions in a manner strikingly similar to humans [79]. This

capacity to gauge the relationship between the environment and one's own body

dimensions suggests a form of bodily self-awareness that operates independently of

mirror recognition [80].

Broader implications and future directions

If honeybees possess even a rudimentary form of MSR—a possibility that remains to be

rigorously tested—it would significantly challenge assumptions about the cognitive and

neural requirements for this ability, contributing to the broader debate on whether larger

brains are necessarily "better" [8], and aligns with the emerging perspective that the

neural substrates for subjective experience, a foundational aspect of self-awareness,

may not be exclusive to vertebrates. Indeed, it has been compellingly argued that the

insect brain possesses the necessary functional organization to support a capacity for

basic consciousness [81]. It would support a "gradualist perspective" on self-awareness,

suggesting that this capacity is not an all-or-nothing phenomenon but rather exists on a

continuum across the animal kingdom [14, 17, 21], ranging from fundamental bodily

self-perception [21, 74, 78, 81] to more abstract self-concepts [9, 13, 74, 81]. This

finding would reinforce the idea that complex behaviors can arise from relatively small

nervous systems [8, 28, 30, 31, 65], a notion corroborated by computational modeling

studies demonstrating that abstract concept learning can be achieved with simple

neural networks inspired by the insect brain [52] and other computational frameworks

[53], and even that numerosity estimation can be enabled by an artificial network of just

four neurons [67]. The prerequisite ability for honeybees to generalize visual patterns to

their mirror images [50] provides a fascinating cognitive foundation, suggesting that

while bees can process mirror-image information, this does not automatically translate

to recognizing their own image.

However, given the significant limitations and strong critiques leveled against non-

primate MSR claims [13], firm conclusions are premature. The history of MSR research

includes numerous claims that did not stand up to rigorous scrutiny, such as initial

failures to find MSR in great apes like gorillas, or early studies on monkeys that lacked

adequate controls [1, 2, 10]. Indeed, a central critique within the primate literature itself

is that even when chimpanzees pass the test, their behavior can be interpreted as a

sophisticated understanding of the mirror's properties for guiding the body, rather than

as evidence for a human-like, introspective self-concept [11]. This history provides a

critical lesson for invertebrate research. As Heyes argued in her review of the primate

literature, the burden of proof for complex cognitive claims lies in experimentally ruling

out simpler, non-mentalistic alternative explanations [64]. Therefore, future research on

honeybees should not only aim for replication but must incorporate the rigorous control

conditions necessary to falsify these leaner hypotheses. Specifically, this requires

studies designed with: (1) increased statistical power through larger sample sizes, (2)

the implementation of essential control groups (e.g., no-mark, conspecific controls), (3)

refined behavioral analysis, and (4) further investigation of individual differences.

Conclusion

This exploratory study employed a mark test to investigate MSR potential in honeybees.

Experiment 2 (with mirror pre-exposure) showed increased overall grooming, more

likely attributable to generalized arousal than specific self-recognition. Our findings must

be considered preliminary due to the lack of a significant preference for mirror-facing

grooming within the mirror group, substantial individual variability, plausible alternative

explanations, and critical methodological limitations, including low statistical power. The

results do not provide conclusive evidence for MSR capability in honeybees at this

stage. However, a failure to pass the classic mark test does not necessarily confirm the

absence of self-awareness, particularly when considering the specific sensory world of

the animal in question [10, 14, 18, 20, 21, 55, 74, 77].

Ultimately, our preliminary findings underscore a critical principle in comparative

psychology, echoing the tenet of Morgan's Canon [69] and the modern intellectual

caution that demands the rigorous exclusion of simpler explanations before accepting

complex cognitive claims [25, 62-65]. Therefore, while our results do not provide

conclusive evidence for MSR, their primary value lies in highlighting specific and critical

avenues for future investigation. Building upon the framework established here, future

research should not only prioritize increasing statistical power but also incorporate

refined control conditions to directly address the alternative hypotheses discussed, such

as testing for mark salience, distinguishing MSR from social responses, and designing

paradigms that can disentangle associative learning from a potential self-concept.

Furthermore, exploring alternative paradigms that may reveal different facets of self-

perception, such as the bodily self-awareness demonstrated in other invertebrates [18,

20, 21, 78-80], will be essential. By systematically exploring these issues, future work

can clarify the cognitive limits of the honeybee mind and its place in the broader debate

on the convergent evolution of cognition [25] and self-awareness.

Acknowledgments
914

915 The authors acknowledge the use of Gemini 2.5 pro for assistance with language

916 editing and improving the clarity of the manuscript. The authors reviewed and take full

917 responsibility for the final content.

Funding
919

920 The authors received no specific funding for this work. All research materials and costs

921 were supported by institutional funds from Mita International School of Science. The

922 funders had no role in study design, data collection and analysis, decision to publish, or

923 preparation of the manuscript.

Author contributions
925

Conceptualization: Gentaro Shishimi

Data curation: Gentaro Shishimi, Rena Akaishi, Mana Matsukami

Formal analysis: Gentaro Shishimi

Investigation: Rena Akaishi, Mana Matsukami

Methodology: Gentaro Shishimi

Project administration: Gentaro Shishimi

Supervision: Gentaro Shishimi

Visualization: Gentaro Shishimi, Rena Akaishi, Mana Matsukami

Writing – original draft: Gentaro Shishimi, Rena Akaishi, Mana Matsukami

Writing – review & editing: Gentaro Shishimi, Rena Akaishi, Mana Matsukami

Data availability statement

938   The raw data for Experiment 1 and Experiment 2 are provided as Supporting

939   Information files S1 and S2, respectively. A video illustrating head-directed grooming

940   behavior of S6 in the mirror group of Experiment 2 is available as Supporting

941   Information file S3 (Video S1).

References

1. Gallup GG Jr, Anderson JR, Shillito DJ. The mirror test. In: Bekoff M, Allen C,

Burghardt GM, editors. *The Cognitive Animal: Empirical and Theoretical*

Perspectives on Animal Cognition. Cambridge (MA): MIT Press; 2002. pp. 325–

333. doi: 10.7551/mitpress/1885.003.0046

2. Anderson JR, Gallup GG Jr. Mirror self-recognition: a review and critique of

attempts to promote and engineer self-recognition in primates. *Primates.*

2015;56(4):317–26. doi: 10.1007/s10329-015-0488-9.

3. Gallup GG Jr. Chimpanzees: self-recognition. *Science.* 1970;167(3914):86–7.

doi: 10.1126/science.167.3914.86.

4. Povinelli DJ, Rulf AB, Landau KR, Bierschwale DT. Self-recognition in

chimpanzees (*Pan troglodytes*): distribution, ontogeny, and patterns of

emergence. *J Comp Psychol.* 1993;107(4):347–72. doi: 10.1037/0735-

7036.107.4.347.

5. Reiss D, Marino L. Mirror self-recognition in the bottlenose dolphin: a case of

cognitive convergence. *Proc Natl Acad Sci U S A.* 2001;98(10):5937–42. doi:

10.1073/pnas.101086398.

6. Plotnik JM, de Waal FBM, Reiss D. Self-recognition in an Asian elephant. *Proc*

Natl Acad Sci U S A. 2006;103(45):17053–7. doi: 10.1073/pnas.0608062103.

7. Prior H, Schwarz A, Güntürkün O. Mirror-induced behavior in the magpie (*Pica*

pica): evidence of self-recognition. *PLoS Biol.* 2008;6(8):e202. doi:

10.1371/journal.pbio.0060202.

8. Chittka L, Niven J. Are bigger brains better? *Curr Biol.* 2009;19(21):R995–1008.

doi: 10.1016/j.cub.2009.08.023.

9. Gallup GG Jr. Self-awareness and the evolution of social intelligence. *Behav*
Processes. 1998;42:239–47. doi: 10.1016/s0376-6357(97)00072-9.

10. De Veer MW, Van den Bos R. A critical review of methodology and interpretation
of mirror self-recognition research in nonhuman primates. *Anim Behav*.
1999;58(3):459–68. doi: 10.1006/anbe.1999.1132.

11. Heyes CM. Reflections on self-recognition in primates. *Anim Behav*.
1994;47(4):909–19. doi: 10.1006/anbe.1994.1126.

12. Kohda M, Hotta T, Takeyama T, Awata S, Tanaka H, Asai J-y, et al. If a fish can
pass the mark test, what are the implications for consciousness and self-
awareness testing in animals? *PLoS Biol.* 2019;17(2):e3000021. doi:
10.1371/journal.pbio.3000021.

13. Gallup GG Jr, Anderson JR. Self-recognition in animals: where do we stand 50
years later? Lessons from cleaner wrasse and other species. *Psychol Conscious*
(Wash D C). 2020;7(1):46–58. doi: 10.1037/cns0000181.

14. de Waal FBM. Fish, mirrors, and a gradualist perspective on self-awareness.
PLoS Biol. 2019;17(2):e3000112. doi: 10.1371/journal.pbio.3000112.

15. Yokose J, Marks WD, Kitamura T. Visuotactile integration facilitates mirror-
induced self-directed behavior through activation of hippocampal neuronal
ensembles in mice. *Neuron*. 2024;112(2):306–318.e7. doi:
10.1016/j.neuron.2023.10.015.

16. Cammaerts MC, Cammaerts R. Are ants (*Hymenoptera, Formicidae*) capable of
self recognition? *J Sci*. 2015;5(7):521–32.

17. Riojas-Schnier JL, Toth AL. Insights into insect cognition through mirror-induced

behaviour in paper wasps. *Behaviour*. 2022;159(8-9):735–58. doi:

10.1163/1568539X-bja10175.

18. Schnell AK, Farndale Wright NR, Clayton NS. The inner lives of cephalopods.

Integr Comp Biol. 2023;63(6):1298–1306. doi: 10.1093/icb/icad122.

19. Ikeda Y, Matsumoto G. Mirror image reactions in the oval squid *Sepioteuthis*

lessoniana. *Fish Sci*. 2007;73(6):1401–1403. doi: 10.1111/j.1444-

2906.2007.01485.x.

20. Amodio P, Fiorito G. A preliminary attempt to investigate mirror self-recognition in

Octopus vulgaris. *Front Physiol*. 2022;13:951808. doi:

10.3389/fphys.2022.951808.

21. Robinson T. Mirror self-recognition in the ghost crab (*Ocypode quadrata*). *Anim*

Cogn. 2023;26(5):1539–49. doi: 10.1007/s10071-023-01777-w.

22. Bitterman ME. Toward a comparative psychology of learning. *Am Psychol*.

1960;15(11):704–712. doi: 10.1037/h0048359.

23. Menzel R, Giurfa M. Cognitive architecture of a mini-brain: the honeybee. *Trends*

Cogn Sci. 2001;5(2):62–71. doi: 10.1016/s1364-6613(00)01601-6.

24. Menzel R. The honeybee as a model for understanding the basis of cognition.

Nat Rev Neurosci. 2012;13(11):758–70. doi: 10.1038/nrn3357.

25. Bitterman ME. Comparative analysis of learning in honeybees. *Anim Learn*

Behav. 1996;24(2):123–141. doi: 10.3758/BF03198961.

26. Srinivasan MV. Honey bees as a model for vision, perception, and cognition.

Annu Rev Entomol. 2010;55:267–84. doi: 10.1146/annurev-ento-112408-085435.

27. Dyer AG. The mysterious cognitive abilities of bees: why models of visual

processing need to consider experience and individual differences in animal

performance. *J Exp Biol.* 2012;215(Pt 3):387–95. doi: 10.1242/jeb.059405.

28. Avarguès-Weber A, Giurfa M. Conceptual learning by miniature brains. *Proc R*

Soc B Biol Sci. 2013;280(1772):20131907. doi: 10.1098/rspb.2013.1907.

29. Giurfa M, Zhang S, Jenett A, Menzel R, Srinivasan MV. The concepts of

'sameness' and 'difference' in an insect. *Nature.* 2001;410(6831):930–3. doi:

10.1038/35073582.

30. Shishimi G. *Working memory and concept learning in honeybees* (Doctoral

dissertation). Honolulu (HI): University of Hawai'i at Manoa; 2013.

31. Giurfa M. Learning of sameness/difference relationships by honey bees:

performance, strategies and ecological context. *Curr Opin Behav Sci.* 2021;37:1–

6. doi: 10.1016/j.cobeha.2020.08.003.

32. Zhang S, Bock F, Si A, Tautz J, Srinivasan MV. Visual working memory in

decision making by honey bees. *Proc Natl Acad Sci U S A.* 2005;102(14):5250–

5. doi: 10.1073/pnas.0501440102.

33. Avarguès-Weber A, Dyer AG, Combe M, Giurfa M. Simultaneous mastering of

two abstract concepts by the miniature brain of bees. *Proc Natl Acad Sci U S A.*

2012;109(19):7481–6. doi: 10.1073/pnas.1202576109

34. Muszynski NM, Couvillon PA. Relational learning in honeybees (*Apis mellifera*):

oddity and nonoddity discrimination. *Behav Processes.* 2015;115:81–93. doi:

10.1016/j.beproc.2015.03.011.

35. Muszynski NM, Couvillon PA. Category difference facilitates oddity learning in

honeybees (*Apis mellifera*). *J Comp Psychol*. 2020;134(3):266–79. doi:

10.1037/com0000216.

36. Avarguès-Weber A, d'Amaro D, Metzler M, Dyer AG. Conceptualization of

relative size by honeybees. *Front Behav Neurosci*. 2014;8:80. doi:

10.3389/fnbeh.2014.00080.

37. Avarguès-Weber A, Dyer AG, Giurfa M. Conceptualization of above and below

relationships by an insect. *Proc R Soc B Biol Sci*. 2011;278(1707):898–905. doi:

10.1098/rspb.2010.1596.

38. Zhang S, Mizutani A, Srinivasan MV. Maze navigation by honeybees: learning

path regularity. *Learn Mem*. 2000;7(6):363–74. doi: 10.1101/lm.32300.

39. Howard SR, Avarguès-Weber A, Garcia JE, Greentree AD, Dyer AG. Numerical

ordering of zero in honey bees. *Science*. 2018;360(6393):1124–6. doi:

10.1126/science.aar4975.

40. Howard SR, Greentree J, Avarguès-Weber A, Garcia JE, Greentree AD, Dyer

AG. Numerosity categorization by parity in an insect and simple neural network.

Front Ecol Evol. 2022;10:805385. doi: 10.3389/fevo.2022.805385.

41. Howard SR, Avarguès-Weber A, Garcia JE, Stuart-Fox D, Dyer AG. Perception

of contextual size illusions by honeybees in restricted and unrestricted viewing

conditions. *Proc R Soc B Biol Sci*. 2017;284(1866):20172278. doi:

10.1098/rspb.2017.2278.

42. Guiraud M, Roper M, Chittka L. High-speed videography reveals how honeybees

can turn a spatial concept learning task into a simple discrimination task by

stereotyped flight movements and sequential inspection of pattern elements. *Front Psychol.* 2018;9:1347. doi: 10.3389/fpsyg.2018.01347.

43. Sandin F, Khan AI, Dyer AG, Amin AHM, Indiveri G, Chicca E, et al. Concept learning in neuromorphic vision systems: What can we learn from insects? *J Softw Eng Appl.* 2014;7(5):387–95. doi: 10.4236/jsea.2014.75037.

44. Bateson M, Desire S, Gartside SE, Wright GA. Agitated honeybees exhibit pessimistic cognitive biases. *Curr Biol.* 2011;21(12):1070–3. doi: 10.1016/j.cub.2011.05.017.

45. Romero-González JE, Solvi C, Chittka L. Honey bees adjust colour preferences in response to concurrent social information from conspecifics and heterospecifics. *Anim Behav.* 2020;170:155–65. doi: 10.1016/j.anbehav.2020.10.012.

46. Seeley TD. *Honeybee Ecology: A Study of Adaptation in Social Life.* Princeton (NJ): Princeton University Press; 1985.

47. Dyer AG, Neumeyer C, Chittka L. Honeybee (*Apis mellifera*) vision can discriminate between and recognise images of human faces. *J Exp Biol.* 2005;208(Pt 24):4709–14. doi: 10.1242/jeb.01956.

48. Avarguès-Weber A, d'Amaro D, Metzler M, Finke V, Baracchi D, Dyer AG. Does holistic processing require a large brain? Insights from honeybees and wasps in fine visual recognition tasks. *Front Psychol.* 2018;9:1313. doi: 10.3389/fpsyg.2018.01313.

49. Howard SR, Avarguès-Weber A, Garcia JE, Dyer AG. Free-flying honeybees extrapolate relational size rules to sort successively visited artificial flowers in a

realistic foraging situation. *Anim Cogn.* 2017;20(4):627–38. doi: 10.1007/s10071-017-1087-3.

50. Stach S, Giurfa M. How honeybees generalize visual patterns to their mirror image and left-right transformation. *Anim Behav.* 2001;62(5):981–92. doi: 10.1006/anbe.2001.1833.

51. Webb B. Can robots make good models of biological behaviour? *Behav Brain Sci.* 2001;24(6):1033-50; discussion 1050–98. doi: 10.1017/s0140525x01000127.

52. Cope AJ, Vasilaki E, Minors D, Sabo C, Marshall JAR, Barron AB. Abstract concept learning in a simple neural network inspired by the insect brain. *PLoS Comput Biol.* 2018;14(9):e1006435. doi: 10.1371/journal.pcbi.1006435.

53. Kleyko D, Osipov E, Gayler RW, Khan AI, Dyer AG. Imitation of honey bees' concept learning processes using Vector Symbolic Architectures. *Biol Inspir Cognit Archit.* 2015;14:57–72. doi: 10.1016/j.bica.2015.09.001.

54. Bourke AFG. Colony size, social complexity and reproductive conflict in social insects. *J Evol Biol.* 1999;12(2):245–57. doi: 10.1046/j.1420-9101.1999.00028.x.

55. Lei Y. Sociality and self-awareness in animals. *Front Psychol.* 2023;13:1065638. doi: 10.3389/fpsyg.2022.1065638.

56. Humphrey NK. *The social function of intellect.* In: Bateson PPG, Hinde RA, editors. Growing Points in Ethology. Cambridge: Cambridge University Press; 1976. pp. 303–317.

57. Sheehan MJ, Tibbetts EA. Specialized face learning is associated with individual

recognition in paper wasps. *Science*. 2011;334(6060):1272–1275. doi:

10.1126/science.1211334.

58. Ikeda Y. A perspective on the study of cognition and sociality of cephalopod

mollusks, a group of intelligent marine invertebrates. *Jpn Psychol Res*.

2009;51(3):146–153. doi: 10.1111/j.1468-5884.2009.00401.x.

59. Ono M, Igarashi T, Ohno E, Sasaki M. Unusual thermal defence by a honeybee

against mass attack by hornets. *Nature*. 1995;377(6547):334–6. doi:

10.1038/377334a0.

60. Heran H, Lindauer M. Windkompensation und seitenwindkorrektur der bienen

beim flug über wasser. *Z Vergl Physiol*. 1963;47(1):39–55. doi:

10.1007/BF00340578.

61. Serres JR, Morice AHP, Blary C, Miot R, Montagne G, Ruffier F. Floor and ceiling

mirror configurations to study altitude control in honeybees. *Biol Lett*.

2022;18(3):20210534. doi: 10.1098/rsbl.2021.0534.

62. Howard SR, Barron AB. Understanding the limits to animal cognition. *Curr Biol*.

2024;34(7):R294–R300. doi: 10.1016/j.cub.2024.02.043.

63. Bitterman ME. The comparative analysis of learning. *Science*.

1975;188(4189):699–709. doi: 10.1126/science.188.4189.699.

64. Heyes CM. Theory of mind in nonhuman primates. *Behav Brain Sci*.

1998;21(1):101–148. doi: 10.1017/s0140525x98000039.

65. Brown MF, Austin BP. Bees and abstract concepts. *Curr Opin Behav Sci*.

2021;37:140–5. doi: 10.1016/j.cobeha.2021.01.006.

66. Chittka L, Jensen K. Animal cognition: Concepts from apes to bees. *Curr Biol*.

2011;21(3):R116–9. doi: 10.1016/j.cub.2010.12.023.

67. Vasas V, Chittka L. Insect-inspired sequential inspection strategy enables an

artificial network of four neurons to estimate numerosity. *iScience*. 2019;11:85–

92. doi: 10.1016/j.isci.2018.12.009.

68. Palmer ME, Calvé MR, Adamo SA. Response of female cuttlefish *Sepia*

officinalis (Cephalopoda) to mirrors and conspecifics: evidence for signaling in

female cuttlefish. *Anim Cogn*. 2006;9(2):151–155. doi: 10.1007/s10071-005-

0009-0.

69. Morgan CL. *An introduction to comparative psychology*. London: Walter Scott;

1894.

70. Premack D. The codes of man and beasts. *Behav Brain Sci*. 1983;6(1):125–67.

doi: 10.1017/s0140525x00015330.

71. Gallup GG Jr, Suarez SD. Social responding to mirrors in rhesus monkeys

(*Macaca mulatta*): effects of temporary mirror removal. *J Comp Psychol*.

1991;105(4):376–9. doi: 10.1037/0735-7036.105.4.376.

72. Gallup GG Jr, Wallnau LB, Suarez SD. Failure to find self-recognition in mother-

infant and infant-infant rhesus monkey pairs. *Folia Primatol* (Basel).

1980;33(3):210–9. doi: 10.1159/000155928.

73. Kakrada E, Colombo M. Mirror mirror on the wall, it's not the mark I care about at

all. *Learn Motiv*. 2022;77:101785. doi: 10.1016/j.lmot.2022.101785.

74. Adolph KE, Tamis-LeMonda CS. Self-recognition: From touching the body to knowing the self. *Curr Biol.* 2024;34(6):R239–R241. doi: 10.1016/j.cub.2024.02.007.

75. Epstein R, Lanza RP, Skinner BF. "Self-awareness" in the pigeon. *Science.* 1981;212(4495):695–6. doi: 10.1126/science.212.4495.695.

76. Uchino E, Watanabe S. Self-recognition in pigeons revisited. *J Exp Anal Behav.* 2014;102(3):327–34. doi: 10.1002/jeab.112.

77. Shashar N, Rutledge PS, Cronin TW. Polarization vision in cuttlefish–a concealed communication channel? *J Exp Biol.* 1996;199(Pt 9):2077–2084. doi: 10.1242/jeb.199.9.2077.

78. Kawashima S, Ikeda Y. Rubber arm illusion in octopus. *Curr Biol.* 2025;35:R702-R703. doi: 10.1016/j.cub.2025.05.017.

79. Ravi S, Siesenop T, Bertrand O, Li L, Doussot C, Warren WH, et al. Bumblebees perceive the spatial layout of their environment in relation to their body size and form to minimize inflight collisions. *Proc Natl Acad Sci U S A.* 2020;117(49):31494–31499. doi: 10.1073/pnas.2016872117.

80. Brebner J, Chittka L. Animal Cognition: The Self-Image of a Bumblebee. *Curr Biol.* 2021;31(4):R207–R209. doi: 10.1016/j.cub.2020.12.027.

81. Barron AB, Klein C. What insects can tell us about the origins of consciousness. *Proc Natl Acad Sci U S A.* 2016;113(18):4900–4908. doi: 10.1073/pnas.1520084113.