The *Stenotrophomonas maltophilia* MntR miniregulon includes novel extracytoplasmic components and affects replication in *Acanthamoeba castellanii* phagosomes

Fulvia-Stefany Argueta-Zepeda[1,2], Javier Rivera[1], Julio César Valerdi-Negreros[1,3] https://orcid.org/0000-0001-7536-5483, Christopher Rensing[4] https://orcid.org/0000-0002-5012-7953, and Pablo Vinuesa[1] https://orcid.org/0000-0001-6119-2956

[1]Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México (UNAM), Cuernavaca, Morelos, México
[2]Programa de Maestría y Doctorado en Ciencias Bioquímicas, UNAM, Mexico City, Mexico
[3]Programa de Doctorado en Ciencias Biomédicas, UNAM, Mexico City, Mexico
[4]Fujian Agriculture and Forestry University, Fuzhou, China.

Address correspondence to Pablo Vinuesa, vinuesa@ccg.unam.mx

Keywords: metal-homeostasis, transcriptomics, host-microbe interactions, intracellular replication, nutritional immunity, free-living amoebae, oxidative stress

ABSTRACT

Manganese homeostasis is crucial for opportunistic pathogens such as *Stenotrophomonas*
maltophilia, which switch lifestyles between contrasting environments. This study elucidated
the manganese homeostasis system in *S. maltophilia* strain Sm18, a Mexican environmental
isolate, revealing a sophisticated network that includes conserved and novel components. We
focused on the MntR miniregulon, which comprises the well-known inner membrane importer
MntH and exporter MntP, both regulated by the Mn^{2+}-responsive MntR metalloregulator.
However, it also contains a TonB-dependent receptor (TBDR) with a distinctive CQNC motif
likely involved in Mn^{2+} coordination and a periplasmic thioredoxin-fold protein (pTFP). Both
are co-expressed from a bicistronic operon under severe manganese and iron limitations and
belong to novel families with a limited phyletic distribution. Transcriptomic profiling under
varying Mn and Fe concentrations revealed a tight interplay between these metals,
emphasizing the ferrophilic nature of *S. maltophilia* and the differential expression of MntR
regulon core members. Additional transporters of the RND and CDF families were modulated
under dual metal limitation. Mutant analysis demonstrated that MntP is critical for survival,
even at Mn levels that do not affect growth kinetics. MntH is required for optimal growth under
oxidative stress and intracellular replication within *Acanthamoeba castellanii* phagosomes,
highlighting its role in overcoming nutritional immunity. Our findings provide new insights into
the remarkable adaptability of *S. maltophilia* to diverse environments, contributing to its
success as an opportunistic pathogen causing a broad range of infections in humans.

1. Introduction

Stenotrophomonas maltophilia is a gram-negative, non-fermentative bacterium recognized
for its ecological versatility and opportunistic pathogenicity (Brooke, 2021). It thrives in diverse
environments, including soil, water, plant rhizospheres, free-living amoebae, and diverse
animal organs and cells, demonstrating remarkable adaptability to contrasting ecological
niches (Ochoa-Sánchez and Vinuesa, 2017; Rivera et al., 2024; Ryan et al., 2009). These bacteria
can survive harsh conditions, such as oligotrophic or metal-rich environments, and are highly
resistant to disinfectants and multiple drugs due to robust biofilm formation and multiple efflux
systems. Recent research has highlighted these adaptive mechanisms, showing how they
enhance the resilience and persistence of *S. maltophilia* in nosocomial environments (Brooke,
2021, 2012), making it one of the ten most prevalent microorganisms in hospital care units

(Rello et al., 2019). This pathogen primarily affects immunocompromised individuals, leading

to respiratory tract, bloodstream, and urinary tract infections (Brooke, 2021), and poses a

significant challenge in clinical settings because of its intrinsic resistance to multiple

antibiotics, including carbapenems (Chang et al., 2015; Ochoa-Sánchez and Vinuesa, 2017).

However, little is known about the genes and mechanisms involved in the adaptation of

Stenotrophomonas to host-associated and intracellular lifestyles in diverse host tissues and

phagocytic cells, including free-living amoebae (Denet et al., 2018; Rivera et al., 2024) and

mammalian macrophages (Takahashi et al., 2020). In recent years, mounting evidence has

revealed that professional phagocytes subject ingested bacteria to limiting or toxic

concentrations of transition metals, inhibiting their proliferation in phagosomes, a process

known as nutritional immunity (Murdoch and Skaar, 2022).

Homeostasis of essential transition metals, such as manganese, iron, copper, and zinc, is

crucial for various physiological processes (Palmer and Skaar, 2016; Waldron and Robinson,

2009). Metallostasis is maintained by a network of allosteric metalloregulators, metal

transporters, metallochaperones, and efflux pumps (Porcheron et al., 2013). The concerted

action of these components regulates the intracellular metal concentrations, ensuring that

they are available for essential functions without reaching toxic levels (Chandrangsu et al.,

2017). The ability of bacteria to rapidly adapt to different environments is tightly coupled to

their metal homeostasis capabilities, particularly during host infection, as metal concentrations

vary widely across organs, tissues, and cellular compartments (Kehl-Fie et al., 2013; Pompilio

et al., 2014).

Studies on *S. maltophilia* metallostasis have focused on iron homeostasis, highlighting its

importance in bacterial virulence (Mikhailovich et al., 2024). In contrast, our knowledge of the

molecular mechanisms underlying manganese homeostasis in *Stenotrophomonas* is limited

to a single transcriptomic study performed on *Stenotrophomonas* sp. MNB17, an organism

isolated from a deep-sea mud sample in the western Pacific, removes Mn^{2+} by efficiently

forming Mn-precipitates ($MnCO_3$ and Mn oxides) at high Mn concentrations (Song et al.,

2022). However, manganese homeostasis has been well studied in some Gram-positive and

Gram-negative bacteria (Bosma et al., 2021; Waters, 2020). A crucial component is the

manganese transporter protein MntH, which facilitates manganese uptake into the cytoplasm

under limiting conditions (Kehres et al., 2000). Conversely, the conserved MntP exporter

removes excess manganese, thereby maintaining intracellular levels within the optimal range

(Waters et al., 2011). This process is regulated by various regulatory proteins and signaling

pathways that sense intracellular manganese levels and adjust the expression or activity of

manganese transporters (Bosma et al., 2021; Martin and Waters, 2022; Sengupta et al.,
2018; Wright et al., 2024), with MntR being the master regulator (Que and Helmann, 2000;
Waters et al., 2011). Furthermore, under manganese limitation and oxidative stress, a Mn^{2+}-
scavenging pathway has been identified in *Burkholderia thailandensis*, comprising a newly
discovered TonB-dependent outer membrane (OM) manganese transporter, MnoT, and a
Mn^{2+}-binding protein, TseM, secreted by a type VI secretion pathway (T6SS) which is
apparently unique to Burkholderia species (Si et al., 2017). The requirement for Mn^{2+} and Fe^{2+}
varies greatly among bacterial species (Bosma et al., 2021). *S. maltophilia* features an iron-
centric metabolism that requires iron for optimal growth and virulence (Kalidasan et al., 2018;
Mikhailovich et al., 2024). However, Mn^{2+} becomes essential under oxidative stress or low
iron availability (Anjem et al., 2009), conditions that mirror the environments that *S.*
maltophilia likely encounters as an intracellular opportunistic pathogen in phagosomes
subjected to nutritional immunity (Čapek and Večerek, 2023; Murdoch and Skaar, 2022). This
bacterium has been reported to survive in human epidermoid epithelial cells (Zhang et al.,
2024), human lung epithelial cells (DuMont et al., 2015; Nas et al., 2019), free-living amoebae
(Denet et al., 2018; Rivera et al., 2024), and mammalian macrophages (Nas et al., 2019;
Takahashi et al., 2020). The divalent Mn^{2+}/Fe^{2+} transporter Nramp1 protein, located in the
phagosomal membrane and ubiquitous in eukaryotes, plays a crucial role in nutritional
immunity by depleting phagocytosed bacteria of these micronutrients. Interestingly, its
bacterial homolog, MntH, is also essential for countering this stress by importing Mn^{2+} into
bacterial cells (Cellier, 2001; Juttukonda and Skaar, 2015).
Here, we present a detailed phylogenomic and functional analysis of *Stenotrophomonas*
manganese homeostasis systems, demonstrating their contribution to bacterial fitness in
Acanthamoeba castellanii phagosomes for the first time. We identified the key components of
the MntR miniregulon in *S. maltophilia* Sm18, a Mexican environmental isolate with complete
genome sequence (Ochoa-Sánchez and Vinuesa, 2017; Vinuesa et al., 2026), and subjected them
to genetic and functional analyses. We demonstrated that the Sm18Δ*mntP* deletion strain
was susceptible to Mn^{2+} *in vitro*. We identified two novel genes as members of the
miniregulon. They are organized in an operon, encoding a TonB-dependent receptor (TBDR)
featuring a conserved CQNC motif, and a periplasmic thioredoxin-fold protein (pTFP), and
have not been previously reported to be manganese-regulated. Phylogenomic and
comparative genomic analyses revealed that both proteins belong to novel families. RNA-
Seq-based transcriptome analyses performed under limiting iron and manganese
concentrations revealed that the TBDR-pTFP locus and *mntH* gene are derepressed,

underscoring their importance in maintaining metal homeostasis in metal-restricted
environments such as the phagosomes of diverse phagocytes. We recently reported that
Sm18 can survive and replicate in the acidified Rab7A-positive vacuoles of *Acanthamoeba*
castellanii trophozoites (Rivera et al., 2024). Here, we show that the *mntH* gene is expressed
and required for optimal intracellular replication in the phagosomes of this free-living
professional phagocyte.
128

2. Material and methods

2.1 Bacterial strains and culture conditions

Stenotrophomonas maltophilia ESTM1D_MKCAZ16_6a (Sm18) was isolated from the
sediments of a clean river in Morelos, Mexico (Ochoa-Sánchez and Vinuesa, 2017), and has its
complete genome sequenced (Vinuesa et al., 2026). Sm18 was routinely cultivated
aerobically in Lysogeny Broth (LB) at 30°C or in MOPS, previously proven effective for
cultivating *Pseudomonas* strains, using 0.4 % (w/v) glucose as the carbon source (LaBauve
and Wargo, 2012). We optimized minimal medium MOPS (MM) for *S. maltophilia* growth by
adding thiamine and biotin at 0.5% and 0.1% (w/v), respectively, along with 0.2% (w/v)
Bacto™ casamino acids (CAA). Our MM lacks manganese in the micronutrient stock. For the
growth kinetics, LB was supplemented with 10mM of MOPS buffer (pH 6.8) and antibiotics, as
required.
142

2.2 Generation of markerless deletion mutants

Two genomic DNA fragments flanking the *mntP* gene were PCR-amplified using total DNA
from Sm18 as the template with primer pairs Sm18DmntP_PCR1.FOR/
Sm18DmntP_PCR1.REV (570 bp) and Sm18DmntP_PCR2.FOR / Sm18DmntP_PCR2.REV
(512 pb). The amplicons were cloned in pEX18Tc (Hoang et al., 1998) linearized with EcoRI
and HindIII using NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, NEB),
following the manufacturer's instructions, and transformed into chemically competent *E. coli*
DH5α. The resulting suicide vector pEX18Tc::DmntP was transferred into Sm18 by
conjugation from *E. coli* S17-1, selecting for Sm18 merodiploids on LB plates supplemented
with 20 µg/mL tetracycline and 30 µg/mL gentamicin. Sm18 merodiploids were grown in LB
without NaCl and supplemented with 10% sucrose directly from glycerol stocks to select for
double recombinants. Markerless deletion mutants were confirmed by tetracycline sensitivity,

colony PCR, and amplicon sequencing (primers Sm18DmntP_VER.F/Sm18DmntP_VER.R;
Supplementary Tables A4, A5, and A6). The same procedure was used to generate
markerless deletions of *orf02357*, *orf02358*, and *mntH*.

2.3 Generation of Vector Integration Mutants (VIM)

An internal (401 bp) DNA fragment from the coding region of the *mntP* gene in Sm18 was
PCR amplified using primers Sm18-VIM-*mntP*.F1/Sm18-VIM-*mntP*.R1 (Supplementary
Tables A4, A5, and A6), directionally cloned into the *Ba*mHI and *Hin*dIII sites of pEX18TcVIM-
GFP, and transformed into chemically competent *E. coli* DH5α following standard procedures.
pEX18TcVIM-GFP_D*mntP* was transferred to Sm18 by conjugation from *E. coli* S17-1. The
same procedure was used to generate VIM derivatives of *orf02357* and *mntH*.
165

2.4 Construction of Sm18 expression plasmids

We modified standard SEVA plasmids (Martínez-García et al., 2023) for the construction of
pSEVA327 (oriVRK2, Cm^R) and pSEVA337R (oriVpBBR1, Cm^R) bacterial expression
plasmids, following the standardized SEVA modular architecture and nomenclature.
Intergenic regions of *orf02357*, *mntH,* and *mntP* were amplified by PCR from genomic DNA
using Phusion polymerase (NEB) and primers listed in Supplementary Tables A4, A5 and A6.
Selected vectors and amplicons were digested with proper restriction enzymes and
directionally cloned upstream of the promoterless reporter genes (GFP or mCherry) to
generate the Sm18 expression plasmids 327-*02357_02358*, 327-*mntH*, 327-*mntP*, 337R-
02357_02358, 337R-*mntH,* and 337R-*mntP* (Supplementary Tables A4, A5 and A6).

2.5 Fluorescence kinetics and microbial growth

The Fluorescence and growth kinetics of strains carrying transcriptional fusions were
assessed across three or more biological replicates in 96-well microplates. The bacterial
optical density (OD_{600}) was adjusted to 0.05 in LB and 0.01 in MM at a final volume of 200 μL
of medium supplemented with increasing $MnCl_2$ concentrations. For growth kinetics, cultures
were grown at 30°C and 205 rpm in a Biotek Epoch 2 microplate spectrophotometer, with
OD_{600} readings taken every 30 or 45 min. The same setup was used for fluorescence kinetics
analysis, with fluorescence (GFP: 479/ 520 nm) and growth (600 nm) measurements
recorded in parallel using a BioTek Synergy H1 microplate reader.

2.6 Statistical analysis of fluorescence and growth kinetics

186   Uninoculated wells with LB or MM served as blanks for microbial growth curves, while 187 strains carrying empty plasmids served as controls. Fluorescence measurements were 188 normalized with OD_{600} readings. The non-parametric Kruskal-Wallis test was used to compare 189 the medians across independent groups based on the Area Under the Curve (AUC) 190 calculated using the Growthcurver R package (v. 0.3.1). When significant differences were 191 detected, Dunn's *post hoc* pairwise comparisons were performed, applying the Bonferroni or 192 Benjamini-Hochberg (BH) correction for multiple testing, as appropriate. Bootstrapping of 193 AUC data was used to compute 95% confidence intervals (CI), represented as error bars in 194 the corresponding figures. All statistical analyses and graphs were performed in R (v 4.3.2), 195 using the ggplot2 (v 3.5.1), ggpubr (v 4.3.2) and rstatix (v0.7.2) packages. Error bars on the 196 growth kinetic graphs represent the standard error of the mean for the technical and biological 197 replicates.

2.7 Cell sample preparation for RNA-sequencing

199   Three biological replicates were obtained from glycerol stocks stored at -80°C and cultured 200 on LB agar at 30°C for 34 h to isolate colonies with a one-day interval between each replicate. 201 A single isolated colony was subcultured in 3 mL of MM and incubated at 30°C with agitation 202 at 205 rpm for 24 h. Precultures (1:50 dilution) were established in 25 mL of MM in a 50 ml 203 Erlenmeyer flask and incubated at 30°C/205 rpm for 14 h. 10 mL of the pre-cultures were 204 collected by centrifugation (5000 × g for 5 min), washed three times with 5 mL of MM without 205 Fe (MM-Fe0), resuspended in 5 mL of the same culture medium, and used to inoculate four 206 25 ml flasks with metal-supplemented MM media at OD_{600} ~ 0.005, labeled as MM-Fe0Mn0, 207 MM-Fe0Mn8, MM-Fe10Mn0, and MM-Fe10Mn8. For all conditions, Sm18 cultures were 208 incubated at 30 °C/205 rpm until reaching the early log phase (OD_{600} 0.1-0.2), after which the 209 cultures were split into two technical replicates, and the cells harvested by centrifugation. 210 Immediately, the pellets were covered with 900µL of RNA later. One technical replicate was 211 stored at 4°C until shipped to GENEWIZ from Azenta Life Science for library preparation and 212 sequencing. The other was stored at -80°C for subsequent experiments. Standard RNA-seq 213 (150 bp paired-end reads) was performed on an Illumina HiSeq 2500 platform.

2.8 RNA-seq analysis data and gene expression

Read quality assessment was performed using Fastqc v 0.12.1 (Andrews, S., 2010), using
Trimmomatic v 0.39 (Bolger et al., 2014) to trim the index adapter and the initial 15 bp from
each read, resulting in 135 bp reads and > 23 million paired-end reads per sample. The
complete genome sequence of S. maltophilia strain ESTM1D_MKCAZ16_6a ((Vinuesa et al.,
2026); CP146374) served as the reference for aligning the reads using Bowtie2 v 2.4.4
(Langmead and Salzberg, 2012). SAM files generated by Bowtie2 (overall alignment rate per
sample >96%) served as input for featuresCounts from the R-Bioconductor software package
Rsubreads v 2.14.2 (Liao et al., 2019). FeaturesCount generated an expression matrix
containing the expression values of each sample for every gene using the GFF genome
annotation file (successfully assigned alignments per sample >92%). Differential gene
expression analysis between treatments was performed using the DESeq2 v1.42.0 (Love et
al., 2014) package, which normalizes the raw counts through the median of ratios method and
conducts differential expression analysis based on the negative binomial distribution of the
data. DESeq2 adjusted p-values using Benjamini and Hochberg's method were used to
reduce the false discovery rate. Differentially expressed genes (DEGs) were identified using
the adjusted p-value (Padj) < 0.01 and log2 fold change ≥ 1 or ≤ -1 as cutoffs. The protein
sequences of DEGs were analyzed using the UniProt database (The UniProt Consortium,
2017), InterProScan (Blum et al., 2025), and MOTIF search
(https://www.genome.jp/tools/motif/) to assign functions and Gene Ontology (GO) terms.
235

2.9 Identification of transcription factor binding sites and protein homologs

We used the motif discovery algorithm MEME (v 5.5.5) from the MEME suite (Bailey et al.,
2015) to identify 16 to 25 nt long sequence motifs in the 5'-UTR regions of
Stenotrophomonas mntH (V8P27_002360) orthologs identified with GET_HOMOLOGUES
(Contreras-Moreira and Vinuesa, 2013). The resulting position-specific probability matrices
(PSPMs) were compared with entries in the CollecTF (Erill, 2015) and PRODORIC (Dudek
and Jahn, 2022) databases of experimentally validated transcription factor-binding sites in
bacteria, using TOMTOM (v 5.5.5) to determine the identity of the discovered motifs. The
collection of 5'-UTR regions containing the MntR-box motif was extended by adding new hits
discovered by MAST (v 5.5.5) scans of complete genome sequences representative of the
currently described taxonomic diversity of *Stenotrophomonas* and related genera in the
Lysobacteriaceae (Chauviat et al., 2025; Vinuesa et al., 2018). Homologs of the *S. maltophilia*
MntR miniregulon proteins were identified by hmmsearch (v3.4) from the HMMER3 suite (44)

using the gathering threshold (--cut_ga) option for calibrated profile hidden Markov models (HMMs) retrieved from Pfam (PF01566 for MntH, PF02659 for MntP), or built locally with hmmbuild (v3.4) from clustal omega (Sievers and Higgins, 2018) alignments (with –iter 2 and –use-kimura options) of our curated datasets of MntR, V8P27_002357, and V8P27_002358 homologs. The latter three HMMs and the PSPM for the *Lisobacteraceae* MntR-box are available for download from https://github.com/vinuesa/supplementary_materials/.

2.10 Live-cell imaging (LCI) using widefield fluorescence microscopy of Sm18/337R-mntH in co-culture with A. castellanii

The assay was reported in detail by (Rivera et al., 2024). Briefly, co-cultures were established by seeding 20-mm diameter glass-bottom cell culture dishes with 5×10^5 *A. castellanii* strain Neff trophozoites, which were incubated for 1 h at 30°C in MMsalts-MOPS-Glc medium. *S. maltophilia* Sm18/337R-*mntH* was grown in LB, washed, and adjusted to an OD_{600} of ~1.0 (~4.8×10^8 CFU/mL). Diluted bacterial suspensions were added to the culture dishes to achieve a multiplicity of infection (MOI) of 25, followed by centrifugation at 500 × g for 5 min to synchronize phagocytosis. LCI was performed using an AxioVertA1 inverted microscope (Karl Zeiss) at three and 18 h ppc.

2.11 Quantitative evaluation of the intracellular replication of mutant strains Sm18VIMmntH and Sm18ΔmntR in Acanthamoeba trophozoites

Experiments were performed using the microtiter plate assay described by (Rivera et al., 2024). Briefly, each well contained 1×10^5 trophozoites in MMsalts-MOPS-Glc and was incubated for 30 min at 25°C. After replacing the medium with the tagged bacteria (Sm18VIM*mntH*::mTn7_Pc_mScarlet-I and Sm18Δ*mntR*::mTn7_Pc_mScarlet-I mutants) at MOI10, phagocytosis was synchronized by centrifugation at 500 × g for 5 min, and the plates placed in a plate reader Synergy 2 and incubated statically at 25°C. Red fluorescence was measured every 30 min for 48 h at 568 nm/594 nm (excitation/emission). The controls included wells with medium, trophozoites in the medium, or bacteria in the medium.

3 RESULTS

3.1 Manganese tolerance of S. maltophilia Sm18 in complex and defined media

To assess the strain's manganese tolerance, we ran 48-hour growth kinetics using complex lysogeny broth (LB) and minimal MOPS (MM) media with increasing $MnCl_2$ concentrations. Fig. 1A shows that Sm18 grew in LB and MM up to 16 mM and 8 mM of $MnCl_2$, respectively. In both media, the maximal growth rate and total yield of the culture remained unaffected, except at the highest Mn concentration tested. However, statistical analysis of the area under the curve (AUC) for population density (OD_{600}) data (Fig. 1B) revealed a significant decrease in growth in both media starting at 8 mM $MnCl_2$, due to the extended lag phase observed with increasing Mn concentrations. These results indicate that Sm18 exhibits notably higher tolerance to manganese than other gram-negative bacteria, such as *Escherichia coli (Zeinert et al., 2018)*.

3.2 Genome-wide identification of manganese-responsive genes in S. maltophilia Sm18 by RNA-seq analysis under varying manganese and iron concentrations

To identify Sm18 genes involved in Mn^{2+} hemostasis under contrasting Fe^{2+} regimes, we analyzed the transcriptional profile of Sm18 cultures (three biological replicates per treatment) using RNA-sequencing (RNA-Seq). Cultures were established in defined MM medium supplemented with casamino acids, under four combinations of Fe^{2+} and Mn^{2+} in micromolar (μM) concentrations: (i) Fe0 Mn0, (ii) Fe0 Mn8, (iii) Fe10 Mn0, and (iv) Fe10 Mn8. Principal component analysis (PCA) shown in Fig. 2A demonstrated that replicates within each group were clustered and separated according to metal concentration. PC1 revealed that 64% of the variance was related to varying iron conditions, while PC2 explained 10% of the variance due to manganese concentrations, supporting the classification of Sm18 as a bacterium with an iron-centric metabolism. To validate this hypothesis, we conducted differential gene expression analysis using DESeq2. We identified a total of 124 DEGs when comparing Fe^{2+} 10 μM vs. Fe^{2+} 0 μM in MM-Mn0, and 89 DEG in MM-Mn8, based on an adjusted *p*-value (padj) < 0.01 and a log_2 fold change $|(log_2FC)| \geq 1.0$ (see Methods). In contrast, only five DEGs were detected when comparing Mn^{2+} 0 μM vs. Mn^{2+} 8 μM in MM-Fe10, and 16 DEG in MM-Fe0. The number of DEGs for each condition is summarized as a Venn diagram (Fig. 2B), revealing that 88 DEGs were regulated solely by iron levels, irrespective of manganese concentrations in the medium. A detailed analysis of these genes will be presented elsewhere (Argueta-Zepeda and Vinuesa, *in preparation*). Among the elements of the MntR miniregulon

in Sm18, *mntP* is the only gene exclusively regulated by manganese levels in the medium,
independent of iron availability (Fig. 2B). In contrast, the expression of *mntH* was affected by
any change in Mn or Fe concentration (Fig. 2B). The latter result aligns with previous reports
showing that *mntH* transcription is regulated by MntR and Fur (Bosma et al., 2021). Nine
genes displayed differential expression only when cultures were subjected to simultaneous
Fe^{2+} and Mn^{2+} limitation (Fig. 2B). Two were down-regulated (blue-colored locus tags) under
these conditions, whereas the remaining genes were up-regulated, as detailed below. Figure
2C highlights the DEGs identified when comparing Mn^{2+} 0 µM vs. Mn^{2+} 8 µM in both MM-Fe10
and MM-Fe0 media (see Supplementary Data, Tables A1 and A2 for more details). These
DEGs are represented as colored dots on the log_2FC plots, and their corresponding genomic
loci are indicated by green vertical bars mapped to the Sm18 chromosome. Their gene
ontologies (GOs) are highlighted in different colors with the transmembrane transport
category (yellow points) being the most prevalent. Fig. 2C also shows the log_2FC values in a
bar plot for a better quantitative representation, including the protein annotations of each
DEG. It shows that V8P27_002357, V8P27_002358, and V8P27_004008, encoding a TonB-
dependent receptor (TBDR), a periplasmic thioredoxin-fold protein (pTFP), and the well-
known cytoplasmic membrane Mn exporter MntP, respectively, display the most significant
log_2FC changes, exceeding 2.8 units in absolute values. TBDR and pTFP are circled and,
with a few others mentioned below, are likely involved in the Sm18 transition metal
homeostasis system under simultaneous manganese and iron limitation. These proteins were
subjected to additional bioinformatic, genetic, and functional characterization, as detailed in
the following sections. Finally, in MM supplemented with 8 µM Mn^{2+}, *mntH* expression was
repressed, whereas *mntP* showed a >3 Log_2FC increase, presumably to export excess
cytoplasmic manganese. Furthermore, in the absence of Fe supplementation, other genes
were up-regulated along with *mntP*, including four predicted transporter proteins (Fig. 2C). In
contrast, five genes primarily associated with transmembrane transport were down-regulated
together with *mntH*, V8P27_002357, and V8P27_002358 (Fig. 2C).
337

3.3 Genome-wide scanning for potential MntR binding sites identified the core components of
the MntR mini regulon in S. maltophilia Sm18

Several Mn^{2+}-responsive genes are known to be regulated by MntR, the master allosteric
manganese transport metalloregulator of the DtxR family (Patzer and Hantke, 2001), which
recognizes a short inverted palindromic sequence in the 5' UTR region of target loci known as

the MntR-box. We performed iterative MEME-MAST (Bailey et al., 2015) searches using 5'-
UTR regions of *Stenotrophomonas mntH* (V8P27_002360) orthologues (see Methods) to
identify potential MntR-boxes in a collection of 103 Lysobacteraceae genomes described in
the following section, resulting in the MEME profile shown in Fig. 3A. MAST-based (Bailey et
al., 2015) genome-wide scanning of the Sm18 assembly with the MntR-box profile revealed
the presence of two additional MntR boxes, defining two additional genomic loci potentially
under MntR regulation. The first site mapped 75 bp upstream of the putative V8P27_002357
(*orf2357*) start codon, which was shown by RNA-seq to be the DEG with the highest log_2FC
value of those displaying differential regulation specifically under both Mn^{2+} and Fe^{2+} limitation
(Figs. 2B and 2C), and closely linked to the *mntR* (V8P27_002359) and *mntH* genes. The
latter two share a bidirectional promoter, with the MntR-box positioned 10 bp upstream of the
mntH putative start codon (Fig. 3B). The third site maps 247 bp upstream of the putative *mntP*
start codon which is differentially expressed in media with high Mn^{2+} concentrations,
irrespective of Fe^{2+} concentration. Similar to *E. coli*, *mntP* in Sm18 is preceded by a putative
yybP-ykoY-like riboswitch (Fig. 3C). Based on this evidence, we postulated that *orf2357,*
orf2358, mntH and *mntP* (Fig. 3B and 3C) form the core of a putative MntR miniregulon in *S.*
maltophilia Sm18, similar to that defined by (Waters et al., 2011) for *Escherichia coli*.
Interestingly, *orf2357* and *orf2358*, are not found in MntR regulons reported in other bacteria
(Martin and Waters, 2022).
362

3.4 Phylogenomic analysis revealed a lineage-specific distribution of MntR miniregulon core
members across Lysobacteraceae
We used hmmsearch from the HMMER3 suite (Eddy, 2011) with calibrated profile hidden
Markov models (see Methods) to identify MntR*,* MntH, MntP, ORF2357 (TBDR), and
ORF2358 (pTFP) homologs across a collection of 103 RefSeq genomes representative of the
currently described taxonomic diversity of *Stenotrophomonas* and related Lysobacteraceae
(Chauviat et al., 2025). Genome-wide MAST scans using the MntR-box profile described in
the previous section (Fig. 3A) identified 0 to 4 MntR-boxes per genome at a cutoff P-value ≤
$5e^{-8}$, suggesting that not all genomes contain the MntR miniregulon. To study the phylogenetic
distribution of its members, we estimated a maximum likelihood species tree for the 103
genomes based on the concatenated supermatrix of 155 top-scoring core genome proteins
selected with GET_PHYLOMARKERS (Vinuesa et al., 2018) out of 601 input core gene
families computed with GET_HOMOLOGUES (Contreras-Moreira and Vinuesa, 2013). The

resulting tree was used to map taxonomic information and integrate it with the presence-

absence patterns of hmmsearch hits and the count data of MAST hits for the MntR motif (Fig.

4). Only *Stenotrophomonas* species, including *S. maltophilia*, encode these five proteins,

which correlates with the presence of three MntR boxes. However, *S. pavanii*, *S. beteli*, and

S. hibiscicola lack TBDR and pTFP proteins and contain only two MntR boxes each. TBDR

and pTFP homologs were not detected in the *Allostenotrophomonas*,

Pseudostenotrophomonas, and *Parastenotrophomonas* proteomes. Only the MntP exporter

was conserved across the four genera analyzed (Fig. 4). In conclusion, within the analyzed

Lyosobacteraceae*,* the five-member MntR miniregulon was found to be restricted to species

in the *S. maltophilia* complex (Chauviat et al., 2025; Vinuesa et al., 2018).

3.5 The expression of S. maltophilia Sm18 MntR regulon core genes is modulated by iron and

manganese concentrations

To validate the bioinformatic predictions supporting the existence of an MntR miniregulon

in *S. maltophilia* complex genomes (Figs. 3 and 4), we constructed transcriptional fusions of

the 5'-UTRs of each of the three core loci of Sm18 to GFP using the broad-host-range

plasmid pSEVA327 (Sm18/327-2357_2358, Sm18/327-*mntH*, and Sm18/327-*mntP;*

Supplementary Data, Table A4). Fig. 5A presents the corresponding promoter activities

measured over 40 h as the AUC values of GFP relative fluorescence units (RFU), normalized

to the population density (OD_{600}), in response to varying Mn and Fe concentrations.

Expression data for the Sm18/327-2357_2358 fusion (Fig. 5A, first panel) revealed that the

operon encoding the novel TBDR and pTFP protein was expressed solely under Mn and Fe

limitation (Mn^{2+} 0 µM and Fe^{2+} 0 µM), consistent with the RNA-seq data (Fig. 2). However, no

GFP signal was detected in MM containing 80 nM Mn^{2+}, indicating that low Mn^{2+}

concentrations abolish their expression. Activity of the Sm18/327-*mntH* fusion (Fig. 5A,

second panel) was negatively correlated with increasing Mn^{2+} concentrations in the culture

medium. However, low activity was still measurable at 8 µM Mn^{2+}. A significantly higher

activity in the Sm18/327-*mntH* expression profile was observed at Fe^{2+} 10 µM vs. 0 µM,

except when the medium contained 8 µM Mn^{2+}. These results confirm a strong relationship

between Mn and Fe homeostasis in Sm18, specifically in manganese uptake systems, as

revealed by our RNA-seq data, and is consistent with observations made in *E. coli* (Taudte et

al., 2016; Wang et al., 2023). In line with our RNA-seq results, the expression profile of

Sm18/327-*mntP* increased significantly at 8 µM Mn^{2+}, compared to Mn^{2+} 0µM, irrespective of

Fe availability in the medium (Fig. 5A, third panel). As shown in Fig. 1, although the upper

Mn^{2+} concentration (8 µM) used for these experiments is orders of magnitude below the

threshold tolerated by Sm18, it is high enough to diminish or abolish the expression of

importers and activate the MntP exporter.

3.6 The MntR metalloregulator controls the expression of the core components of the Sm18

Mn homeostasis system

To confirm that the promoter activity observed in the transcriptional fusions reported in Fig.

5A is regulated by MntR, the three fusions were transferred into a Sm18Δ*mntR* strain carrying

a markerless *mntR* deletion (Supplementary Data, Table A3). Promoter activity in iron-

deficient MM with 0 µM and 8 µM manganese was measured as described in the previous

section (Fig. 5A). The Sm18Δ*mntR*/327-2357_2358 fusion (Fig. 5B, first panel) revealed a

significant increase in expression in the absence of supplemented manganese, suggesting

derepression in the absence of the MntR transcription factor. However, adding manganese

led to reduced expression, following a similar trend to that observed in the Sm18 wild-type

background, suggesting that manganese may also negatively regulate this operon via an

alternative MntR-independent mechanism. Similarly, derepression of the 327-*mntH* fusion

was observed in the Sm18Δ*mntR* genetic background compared to the wild-type growing in

MM without supplemented manganese (Fig. 5B, second panel). However, no statistically

significant activity was detected in the Sm18ΔmntR/327-*mntH* fusion upon addition of 8 µM

Mn^{2+}. This suggests that manganese-dependent regulation of *mntH* promoter activity is

mediated exclusively by MntR. Finally, the *mntP* fusion exhibited manganese-dependent

regulation exclusively through MntR, as no expression was detected in the absence of MntR

(Fig. 5B), independent of manganese concentration, in line with the RNA-seq data (Fig. 2).

3.7 The V8P27_002357 TonB-dependent receptor belongs to a novel family with a restricted

phylogenetic distribution featuring a conserved CQNC motif in its plug domain

After confirming that V8P27_002357 (TBDR) and V8P27_002358 (pTFP) are members of

the MntR miniregulon restricted within Lysobacteraceae to some Stenotrophomonas species

(Fig. 4), and are subjected to Mn^{2+} regulation (Figs. 2 and 5), we performed a domain, motif,

and phylogenomic characterization of these novel proteins. Fig. 6 summarizes the findings for

V8P27_002357. It encodes a 719 aa TonB-dependent receptor with a 28 bp-long N-terminal

signal peptide (Fig. 6A), resulting in a mature protein of 76.6 kDa (691 aa) containing an N-

terminal plug or core domain (141 aa, 15 kDa) featuring a centrally located CQNC motif,

followed by a TBDR β-barrel domain (550 aa, 61.6 kDa). AlphaFold3 (AF3) (Abramson et al.,

2024) modeling of V8P27_002357 in the apo and holo conformations strongly suggests (ipTM

= 0.84; pTM =0.90; mean pLDDT = 87.08) that a Mn^{2+} cation could be coordinated by two

sulfur, one nitrogen, and one oxygen atom arranged in a tetrahedral geometry, contributed by

residues C63, C66 from the CQNC motif, and residues E247, and H446 from the β-barrel

domain, respectively (Supplementary Figure S1).

We used BlastP to search for V8P27_002357 homologs (excluding *Stenotrophomonas*

hits) in the PDB, Swiss-Prot, and RefSeq protein databases. This analysis revealed a

restricted and sparse phylogenetic distribution of the closest hits, which belonged to a few

genera in the classes Bacteroidia, Flavobacteriia, Holophagae, Opitutia, Alphaproteobacteria,

and Gammaproteobacteria. These sequences were subjected to structure-guided multiple-

sequence alignment and maximum likelihood phylogeny estimation (see Methods section).

Fig. 6B depicts a small summary tree, revealing that the closest V8P27_002357 homologs

form a perfectly supported (100% bootstrap support) and tightly clustered lineage of

sequences, all of which share the distinct CQNC motif in their plug domain (Fig. 6C: Fig S1).

Based on this evidence, we propose that V8P27_002357-like proteins comprise a new family

(CQNC) of TBDRs (Fig. 6B).

3.8 V8P27_002358 belongs to a novel family of periplasmic thioredoxin-fold proteins (pTFPs)

linked to CQNC-type TBDRs by synteny and co-expression, featuring a WC[AG]PC motif

The presence of a periplasmic thioredoxin-fold protein (pTFP; matching IPR013766) co-

expressed with a preceding CQNC-type TBDR (V8P27_002357 in Sm18) has not been

previously reported and suggests that both proteins might be functionally related. To

challenge this hypothesis, we performed phylogenomic and comparative genomic analyses of

this protein. BLAST searching for V8P27_002358 homologs was done as described in the

previous section for the TBDR protein, extending the collection of homologues with the

recently reported metagenomic dataset of thioredoxin superfamily proteins compiled by

Nilewski et al. (Nilewski et al., 2021), and hmmsearch hits found by screening the Reference

Proteomes 2025_01 database through the HMMER portal with our pTFP.hmm profile HMM

as the query (see Methods), resulting in a collection of 278 proteins. After alignment with

clustal-omega, the resulting matrix was manually curated, generating a final set of 78 closely-
related homologs, including plant sequences automatically annotated (TrEMBL) as protein-
disulfide reductases. A maximum likelihood phylogeny was estimated from this set, revealing
that V8P27_002358 clustered in a highly supported (99% bootstrap support) and distinct
clade (Supplementary Fig. S2) with uncharacterized sequences from a few bacterial genera
belonging to the orders Lysobacterales (Gammaproteobacteria), Sphingomonadales
(Alphaproteobacteria), and Opitutales (Verrucomicrobiota; Opitutia). Synteny analysis of
representative bacterial sequences revealed that the V8P27_002358 clade members were
located immediately downstream of the CQNC-type TBDR (Fig. 6D), strongly reinforcing the
notion that they are functionally related. However, the genes flanking the TBDR-pTFP locus
were variable (Fig. 6D).

V8P27_002358 is a 173 aa long protein and features an N-terminal signal peptide of 23 aa.
The mature protein contained 159 aa, with a computed mass of 17.64 kDa and a periplasmic
location, as predicted by Phobius (p = 0.94) and Psortb (score = 9.84). All pTFPs share a
minimal thioredoxin fold (TRX-fold) characterized by an N-terminal βαβ fold separated by an
α-helix from the C-terminal ββα fold (Martin, 1995). The highly diversified and ubiquitous
superfamily of pTFPs (Anjou et al., 2024; Atkinson and Babbitt, 2009; Nilewski et al., 2021)
encodes thiol:disulfide oxidoreductases (TDORs) that contain additional secondary structural
elements and loops inserted between the conserved domains of the TRX fold. Figure 6E
displays the highly conserved CXXC (C39, C42) motif of thioredoxins and thioredoxin-like
proteins found at the beginning of the α1 helix of the TRX fold, as confirmed by our AF3
modeling (Supplementary Figure S3). V8P27_002358 contains an unusual CAPC dipeptide in
the presumed catalytic CXXC redox center. Motif analysis around the β1-α1 region of the
TRX-fold, including well-characterized pTFPs (SwissProt codes) with contrasting reduction
potentials and pK_a values previously analyzed by Mavridou et al. (Mavridou et al., 2014)
revealed a considerable diversity in and around the CXXC motif. The pTFPs associated with
CQNC-type TBDRs are characterized by a distinctive and lineage-specific
Y[FYC]GA[DS]WC[AG]PCH[AE]F[VL]P profile overlapping the CXXC motif (Fig. 6E).
501
3.9 Mutants of MntR miniregulon members result in hydrogen peroxide and manganese
sensitivity

To investigate whether the Sm18 *mntH* and *mntP* homologs encode functional manganese
importer and exporter proteins, we constructed the vector integration mutant (VIM)

Sm18VIM*mntH* and the markerless deletion mutant Sm18Δ*mntP* (see Methods). The growth

kinetics of the Sm18VIM*mntH* mutant indicated that the *mntH* gene is not essential under

Fe^{2+}- or Mn^{2+}-limiting conditions in axenic cultures. However, it is required for growth under

oxidative stress induced by hydrogen peroxide (H_2O_2), particularly at neutral pH (Fig. 7A).

Statistical analysis revealed that the difference in growth was significant only at 500 μM H_2O_2

compered with the wild type (Fig. 7B). These results are consistent with other studies, which

demonstrated that *mntH* mutants are sensitive to H_2O_2, a phenotype largely attributed to their

impaired ability to import manganese into the intracellular space (Anjem et al., 2009).

Conversely, it has been well established in diverse bacterial models that manganese exporter

proteins are required to counter manganese toxicity (Waters, 2020; Zeinert et al., 2018).

Sm18Δ*mntP* was challenged with increasing Mn^{2+} concentrations (5-160 μM; Fig. 7C), with

statistically significant sensitivity detectable at 40 μM, and complete growth inhibition

observed at 160 μM (Fig. 7D). The wild-type phenotypes of the Sm18VIM-*mntH* and

Sm18Δ*mntP* strains were restored by complementing plasmids Sm18VIM-*mntH*/332::*mntH*

and Sm18Δ*mntP*/332::*mntP* provided in *trans*, and carrying the *mntH* and *mntP* genes under

the control of their native promoters (Fig. 7). These results demonstrate that *mntH* and *mntP*

expression is essential for H_2O_2 resistance and manganese detoxification, respectively, and

confirm that the uncovered MntR miniregulon is functional. Notably, no growth phenotypes

were detected for the markerless V8P27_002357 and V8P27_002358 mutants under all

conditions tested.

3.10 The mntH gene of S. maltophilia is expressed in A. castellanii phagosomes and is

required for optimal intracellular replication

We recently found that *S. maltophilia* Sm18 resists digestion by *A. castellanii* strain Neff

trophozoites and can replicate in acidified, Rab7A-positive *Stenotrophomonas*-containing

vacuoles (SCV) of this free-living professional phagocyte (Rivera et al., 2024). Given that the

phagosomes of *Dictyostelium discoideum* and human macrophages subject bacteria to iron

and manganese limitation via the NRAMP exporter (Morey et al., 2015; Peracino et al., 2013),

we postulated that the Sm18 MntH Mn importer is required to survive in the SCV. To test this

hypothesis, we first evaluated the transcriptional fusion of the *mntH* promoter region to

mCherry2 by live-cell imaging (LCI) of *A. castellanii* strain Neff co-cultured with

Sm18/pSEVA337R_mntH set at MOI25, using wide-field fluorescence microscopy. As shown

in Fig. 8A, the fusion was expressed at 3 and 18 h post-primary contact (ppc). This result

strongly suggests that phagocytosed Sm18 cells experience Mn and possibly Fe limitations in
the SCV. We used our *A. castellanii-Stenotrophomonas* co-culture assay in 96-well microtiter
plates coupled to a fluorescence reader (Rivera et al., 2024) to evaluate the replication
capacity of strains Sm18VIM*mntH* and Sm18Δ*mntR* in trophozoites. We tagged
Sm18VIM*mntH* and Sm18Δ*mntR* with our pUC18T_mTn7TC1_Pc_mScarlet-I
minitransposon, which constitutively expresses the mScarlet-I fluorescent protein from the
strong Pc promoter (Rivera et al., 2024), resulting in strains
Sm18VIM*mntH*::mTn7_Pc_mScarlet-I and Sm18Δ*mntR*::mTn7_Pc_mScarlet-I, respectively
(Supplementary Data, Table A3). Their replication capacity was compared to that of Sm18
tagged with the same transposon (Sm18:: mTn7_Pc_mScarlet-I). As shown in Fig. 8B, the
Sm18VIM*mntH* mutant had a significantly reduced replication capacity in Neff trophozoites
compared to Sm18, whereas that of Sm18Δ*mntR* was indistinguishable from that of the wild
type strain (Figs. 8B and 8C). These findings support the hypothesis that *Acanthamoeba*
phagosomes restrict manganese availability to ingested bacteria and demonstrate that MntH
is indispensable for optimal intracellular replication.

555 4. Discussion

This study provides the first insights into the MntR miniregulon and manganese homeostasis
of *Stenotrophomonas maltophilia* and related species, revealing similarities to known systems
in other bacteria and novel cell envelope components. As expected, the core elements of the
S. maltophilia MntR system include the inner-membrane importer and exporter proteins MntH
and MntP under the control of the MntR allosteric metalloregulator, which are highly
conserved throughout evolution (Martin and Waters, 2022; Waters et al., 2011). However, *S.*
maltophilia and other *Stenotrophomonas* species also feature a novel bicistronic operon
encoding an outer membrane (OM) TonB-dependent receptor (TBDR, V8P17_002357) and a
periplasmic thioredoxin-fold protein (pTFP, V8P17_002358). Gram-negative bacteria typically
encode multiple TBDRs, many of which have unknown functions (Blanvillain et al., 2007;
Schauer et al., 2008). It is well established that in *S. maltophilia,* iron limitation enhances
virulence and biofilm formation (García et al., 2015; Kalidasan et al., 2018), and the expression of
TBDRs involved in heme (Shin et al., 2024), hemin (Liao et al., 2024; Shih et al., 2022), and
ferri-stenobactin (Wu et al., 2022; Yeh et al., 2025) uptake, as well as others involved in
competition for secreted pyochelin siderophores in co-cultures with *Pseudomonas aeruginosa*
(Pan et al., 2022). In addition to iron acquisition, *S. maltophilia* TBDRs have been implicated

in β-lactam (Calvopiña et al., 2020) and keratin monomer uptake (Xue et al., 2024).

Xanthomonas species, which are close relatives of *Stenotrophomonas*, feature around 70

TBDRs, many of which are required for scavenging plant carbohydrates (Blanvillain et al.,

2007). At least 14 TonB-dependent iron acquisition and virulence pathways have been

extensively studied in *P. aeruginosa (Hubert et al., 2023)*. Other well-known TBDRs, such as

BtuB, OprC, FhuA, and FecA, translocate vitamin B12 (cyanocobalamin), copper, ferric

hydroxamate, and Fe^{3+} dicitrate, respectively. However, MnoT (Q2SY62_BURTA) from

Burkholderia is the only TBDR currently known to import Mn (Si et al., 2017). Analysis of the

S. maltophilia Sm18 genome (Vinuesa et al., 2026) revealed that it encodes 65 putative

TBDRs, representing one of the most expanded paralog gene families in this organism (data

not shown). Our MEME-MAST analyses (Fig. 3B) revealed that V8P27_002357 is the only

TBDR associated with an MntR-binding box in Sm18. This box is located 75 bp upstream of

the putative start codon, between the -10 and -35 regions (Fig. 3B), supporting the notion that

MntR is a transcriptional repressor of this locus. Transcriptional fusion and transcriptome

profiling data revealed that the *orf2357-orf2358* operon was derepressed only under severe

manganese and iron restriction, with its expression virtually abolished upon the addition of 80

nM manganese. This correlates with observations of other TBDRs, typically expressed only

under severely limiting nutritional conditions (Si et al., 2017; Silale and Van Den Berg, 2023),

and highlights the interplay of manganese and iron in the unknown regulatory mechanism of

the *orf2357-orf2358* locus. However, the Sm18Δ*orf2357* (TBDR) and Sm18Δ*orf2358* deletion

mutants did not exhibit a growth phenotype under these metal-restricted conditions, and the

Sm18/337R-pr_*2357-2358* transcriptional fusion was not expressed in co-cultures with *A.*

castellanii trophozoites (data not shown). Phylogenomic analysis of Sm18 TBDRs and a

broad collection of well-classified reference proteins from this family revealed that it belongs

to a novel clade possibly involved in Mn uptake, which is unrelated to the *Burkholderia* MnoT

clade or typical iron-importing TBDRs mentioned above (data not shown). This conclusion is

supported by detailed structrual and motif analyses of V8P27_002357 (Fig S1). Its most

distinguishing feature is the presence of a CQNC motif located in the center of the plug

domain. Based on the bioinformatic and transcriptional evidence gathered, we hypothesized

that V8P27_002357 might function as an OM Mn importer, with the CQNC motif likely

involved in Mn^{2+} coordination, as strongly suggested by our AlphaFold3 modeling (Fig. S1).

The discovery of a periplasmic thioredoxin-fold protein (pTFP; V8P27_002358) as part of the

MntR regulon expands the known components of bacterial manganese homeostasis and

thiol-disulfide exchange systems. Furthermore, this finding adds a new layer of complexity to

our understanding of how bacteria manage metal homeostasis and oxidative stress in the cell
envelopes. Synteny, phylogenetic co-occurrence (Fig. 6D), and co-transcription in a
bicistronic operon with the preceding CQNC-type TBDR (V8P27_002357) suggest a
functional coupling between both proteins. It is well established that most secreted proteins
enter the periplasm via the Sec pathway in an unfolded state (Smets et al., 2019). The
periplasm of Gram-negative bacteria is an oxidizing compartment in which sulfur-containing
amino acids, such as cysteine and methionine, are particularly vulnerable to oxidation by
ROS (Collet et al., 2020). Disulfide bond formation in the periplasm of didermic
Gammaproteobacteria is catalyzed by DsbA from the canonical DsbAB pathway as the
unfolded intermediate transits through the SecYEG channel (Kadokura and Beckwith, 2009;
Landeta et al., 2018). DsbA is a highly oxidizing soluble periplasmic oxidoreductase with a
canonical thioredoxin fold and CXXC motif, which is also present in *S. maltophilia* Sm18
(V8P27_002985, BsbAsm). Therefore, it is likely that the two consecutive cysteines found in
CQNC motif othe CQNC-TBDR are oxidized, forming a disulfide bridge by the action of
BsbAsm as it transits into the periplasm through the SecYEG channel during export. In this
model, the cognate and specialized periplasmic thioredoxin-fold protein V8P27_002358 is
required to reduce the disulfide bond in the CQNC motif as the protein folds and integrates
into the outer membrane to fulfill its predicted role as a Mn-importing TBDR. Therefore, we
postulate that V8P27_002358 might be a specialized disulfide reductase for the CQNC motif
in V8P27_002357 featuring a distinct WC[AG]PC motif (Fig. 6 and Fig. S3). This working
model for the novel TBDR-pTFP locus opens new avenues for research into bacterial metal
acquisition strategies and thiol-disulfide homeostasis in the periplasm.
Regarding the conserved and broadly distributed MntR miniregulon components, this study
reports the first genetic analysis of *mntH* in *S. maltophilia*. We demonstrate that it encodes its
primary manganese importer protein, and is relevant for replication in amoebal phagosomes.
There are several lines of evidence to support this conclusion. The *mntH* gene maps
downstream of the *mntR* gene. It is regulated by both manganese and iron availability, as
shown by our RNA-seq and transcriptional fusion data, and has been observed in *E. coli* and
other bacteria (Martin and Waters, 2022). Furthermore, transcriptome analysis suggests that
no other manganese inner-membrane transporter is solely regulated by manganese
availability in Sm18. Perhaps most significantly, *mntH* is reported herein for the first time to be
required for optimal replication of *S. maltophilia* within *A. castellanii*, a professional phagocyte.
This finding has broader implications for understanding the pathogenicity of *S. maltophilia*,
suggesting that manganese acquisition contributes significantly to bacterial fitness within host

cells. We observed active expression of the *mntH* promoter probe *in vivo* in co-cultures with

A. castellanii trophozoites. This indicates that Sm18 most likely faces manganese and iron

restrictions within amoebal phagosomes. *S. maltophilia*'s ability to overcome these limitations

through MntH-mediated manganese import may be a key factor in its success as an

intracellular pathogen, as found in other bacteria (Anderson et al., 2009; Boyer et al., 2002;

Champion et al., 2011; Zaharik et al., 2004), effectively overcoming the host's imposed

nutritional immunity (Chandrangsu et al., 2017; Murdoch and Skaar, 2022), an ancient trait of

eukaryotic phagocytes conserved from amoebae to macrophages (Yu et al., 2024). Given that

S. maltophilia, like most Gammaproteobacteria, is a ferrophilic bacterium (see further

discussion below), it is not surprising that the Sm18VIM*mntH* mutant did not display

significant impairment when grown in defined MOPS medium without Mn (Fig. 7A). This

observation suggests that the attenuated replication rate of this mutant in amoebal

phagosomes might be due to its limited capacity to contend with reactive oxygen species

(ROS), such as superoxide anions and peroxides (Anjem et al., 2009). *S. maltophilia*

genomes, including that of Sm18, encode a periplasmic Mn-dependent superoxide dismutase

(SOD) and Mn catalase (Crossman et al., 2008). In addition, some enzymes, such as

cambialistic SODs (Sendra et al., 2023) or ribonucleotide reductases (Makhlynets et al., 2014;

Martin and Imlay, 2011; Wang et al., 2023), can use either Fe or Mn as a co-factor, whereas

mismetallation or oxidation via Fenton reactions of iron-containing enzymes in metal-

restricted environments in the presence of ROS may be alleviated by using Mn as an

alternative co-factor, as described for diverse mononuclear iron enzymes, most notably the

key enzyme ribulose-5-phosphate 3-epimerase (Rpe) from the pentose-phosphate pathway

(PPP) (Anjem et al., 2009; Imlay, 2014; Sobota and Imlay, 2011). The metallation of Rpe with

oxidative stress-resistant Mn^{2+} enables carbon flow through the PPP, producing the NADPH-

reducing power required to maintain ROS-countering enzymes, such as glutathione or

thioredoxin reductase (Song et al., 2016; Uppalapati and Vazquez-Torres, 2022) in the

cytoplasm and diverse periplasmic disulfide isomerases and reductases, probably including

the novel pTFP-encoding gene discovered herein.

In this study, we analyzed the links between manganese and iron homeostasis systems,

focusing on the genes specifically induced by the former under the control of MntR and iron

limitation. However, in addition to the MntR miniregulon members, a few other genes were

differentially expressed under iron-replete and manganese excess conditions. Of particular

relevance was the identification of efflux pumps of the CDF and RND families, as they are

know to support intracellular detoxification of zinc and copper during nutritional immunity

(Hood and Skaar, 2012), when iron and manganese concentrations are low, as shown in our
integrative model (Fig. 9). Emerging evidence indicates that an imbalanced ratio of transition
metals can function as either a direct or indirect regulatory mechanism to mitigate metal
toxicity through efflux pumps. The well-studied YiiP (CDF) has been identified as a secondary
transporter for manganese export in Salmonella Typhimurium. Notably, its activity is not
regulated by Fe or Mn concentrations in the medium, indicating that YiiP may play a role in
balancing the levels of other metals (Ouyang et al., 2009). In *Streptococcus pneumoniae*,
expression of *czcD*, a zinc CDF efflux pump, is upregulated under Mn(II)-limiting conditions
(Ogunniyi et al., 2010). This is consistent with other studies showing that high zinc
concentrations can compete with manganese-binding sites on enzymes essential for cellular
physiology (Sullivan et al., 2024). In *Bacillus subtilis*, reduced intracellular iron levels heighten
cell sensitivity to elevated Co(II) or Mn(II) levels, particularly in strains deficient for export of
these metals (Pi et al., 2020). Furthermore, disruption of the Mg(II) efflux system alleviates
residual manganese toxicity in an *mntR mntH* mutant, likely by preventing mismetallation
through Mg(II) accumulation (Hohle and O'Brian, 2014; Pi et al., 2020). In summary, we
hypothesize that under manganese and iron limitation, in addition to uptake systems, Sm18
deploys additional survival mechanisms mediated by transporters that re-establish the
balance of transition metal pools by effluxing the most reactive ones, with more favorable free
energies (Irving and Williams, 1953; Osman et al., 2019), thereby minimizing mismetallation
(Fig. 9). However, the role of the differentially expressed CDF and RND transporters in metal
homeostasis needs to be experimentally established in Sm18.
695
696
5. Conclusion

This is the first comprehensive analysis of manganese homeostasis in *S. maltophilia.* It
revealed a sophisticated network of importers, exporters, and regulatory elements that work
coordinately to maintain appropriate intracellular manganese levels. The *S. maltophilia* MntR
miniregulon displays intricate connections with Fe and oxidative stress homeostasis systems
(Fig. 9). The identification of novel extracytoplasmic components (CQNC-TBDR and pTFP)
expands our understanding of bacterial metal homeostasis systems, opening new avenues
for future research. The *mntH* gene is required for optimal replication in *Acanthamoeba*
trophozoites and in minimal media depleted of Fe and Mn under oxidative stress, although
dispensable in complex LB medium. These findings highlight the ferrophilic nature of *S.*
maltophilia, advance our knowledge of *S. maltophilia* physiology and adaptation to

intracellular life in phagosomes, contributing to the broader field of bacterial metal

homeostasis, potentially opening new strategies to combat infections by this opportunistic,

multidrug- and amoeba-resistant pathogen.

712 CRediT authorship contribution statement

Fulvia Stefany Argueta Zepeda: Writing – original draft, Writing – review & editing, Validation,
Methodology, Investigation, Formal analysis, Data curation. **Javier Rivera Campos:** Writing – review
& editing, Validation, Methodology. **Julio Valerdi Negreros:** Writing – review & editing, Validation,
Methodology. **Christopher Rensing:** Writing – review & editing, Supervision. **Pablo Vinuesa:**
Writing – review & editing, Writing – original draft, Supervision, Resources, Project administration,
Methodology, Investigation, Funding acquisition, Formal analysis, Conceptualization

721 Conflict of interest

722 The authors declare no conflict of interest.

723
724

725 ACKNOWLEDGEMENTS

We thank Alfredo Hernández and Víctor del Moral for their expert technical support and

server maintenance at the Centro de Ciencias Genómicas, UNAM. We acknowledge the

excellent service provided by personnel at the Unidad de Síntesis y Secuenciación de DNA

from Instituto de Biotecnología, UNAM. We thank Dr. Ayari Fuentes from CCG-UNAM for

providing access to the multimodal plate reader. We thank Dr. Luis Lozano from the

Bioinformatics Unit at CCG-UNAM for his advice and support during the early phase of

transcriptome analysis. We thank M.Sc. Laura Cervantes for her laboratory assistance and

valuable feedback. We gratefully acknowledge the financial support received from Consejo

Nacional de Humanidades Ciencias y Tecnologías (CONAHCyT México, A1-S-11242),

Secretaría de Ciencia, Humanidades, Tecnología e Innovación (SECIHTI México CBF-2025-

I-1106), and Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica

(PAPIIT), Universidad Nacional Autónoma de México (DGAPA-PAPIIT, UNAM: IN209321 and

IN216424) to PV. F. S. A. Z. and J.C.V.N. were recipients of CONAHCyT scholarships

659308 and 745705, respectively.

741 Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at

doi:10.1016/j.micres.2026 ….

Data availability: The complete genome sequence of *S. maltophilia* Sm18 was deposited in

GenBank under accession number CP146374 and the associated genome and transcriptome

sequencing reads were deposited under BioProject PRJNA1081934

https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1081934. The RefSeq genome

assembly for strain Sm18 is publicly available from NCBI's genome datasets portal at

https://www.ncbi.nlm.nih.gov/datasets/genome/GCA_053078355.1/.

The sequence reads generated from the 18 RNAseq samples analyzed in this work were

deposited at NCBI's Sequence Read Archive https://www.ncbi.nlm.nih.gov/sra/ under

accession numbers SRR36898890 to SRR36898907, as part of BioProject PRJNA1081934

https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1081934.

The hidden Markov models built in this work are freely available from GitHub at:

https://github.com/vinuesa/supplementary_materials/tree/main/docs/Argueta-

Zepeda_Stenotrophomonas_MntR_miniregulon.

Conflict of interest disclosure statement: None declared.

Ethics statement: None required.

REFERENCES

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung, C.-C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain, R., Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Žídek, A., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., Jumper, J.M., 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature 630, 493–500. https://doi.org/10.1038/s41586-024-07487-w

Anderson, E.S., Paulley, J.T., Gaines, J.M., Valderas, M.W., Martin, D.W., Menscher, E., Brown, T.D., Burns, C.S., Roop, R.M., 2009. The Manganese Transporter MntH Is a Critical Virulence Determinant for *Brucella abortus* 2308 in Experimentally Infected Mice. Infect Immun 77, 3466–3474. https://doi.org/10.1128/IAI.00444-09

Andrews, S., 2010. Fastqc: A quality control tool for high throughput sequence data.

Anjem, A., Varghese, S., Imlay, J.A., 2009. Manganese import is a key element of the OxyR response

to hydrogen peroxide in *Escherichia coli*. Molecular Microbiology 72, 844–858. https://doi.org/10.1111/j.1365-2958.2009.06699.x

Anjou, C., Lotoux, A., Morvan, C., Martin-Verstraete, I., 2024. From ubiquity to specificity: The diverse functions of bacterial thioredoxin systems. Environmental Microbiology 26, e16668. https://doi.org/10.1111/1462-2920.16668

Atkinson, H.J., Babbitt, P.C., 2009. An Atlas of the Thioredoxin Fold Class Reveals the Complexity of Function-Enabling Adaptations. PLoS Comput Biol 5, e1000541. https://doi.org/10.1371/journal.pcbi.1000541

Bailey, T.L., Johnson, J., Grant, C.E., Noble, W.S., 2015. The MEME Suite. Nucleic Acids Res 43, W39–W49. https://doi.org/10.1093/nar/gkv416

Blanvillain, S., Meyer, D., Boulanger, A., Lautier, M., Guynet, C., Denancé, N., Vasse, J., Lauber, E., Arlat, M., 2007. Plant Carbohydrate Scavenging through TonB-Dependent Receptors: A Feature Shared by Phytopathogenic and Aquatic Bacteria. PLoS ONE 2, e224. https://doi.org/10.1371/journal.pone.0000224

Blum, M., Andreeva, A., Florentino, L.C., Chuguransky, S.R., Grego, T., Hobbs, E., Pinto, B.L., Orr, A., Paysan-Lafosse, T., Ponamareva, I., Salazar, G.A., Bordin, N., Bork, P., Bridge, A., Colwell, L., Gough, J., Haft, D.H., Letunic, I., Llinares-López, F., Marchler-Bauer, A., Meng-Papaxanthos, L., Mi, H., Natale, D.A., Orengo, C.A., Pandurangan, A.P., Piovesan, D., Rivoire, C., Sigrist, C.J.A., Thanki, N., Thibaud-Nissen, F., Thomas, P.D., Tosatto, S.C.E., Wu, C.H., Bateman, A., 2025. InterPro: the protein sequence classification resource in 2025. Nucleic Acids Research 53, D444–D456. https://doi.org/10.1093/nar/gkae1082

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bosma, E.F., Rau, M.H., van Gijtenbeek, L.A., Siedler, S., 2021. Regulation and distinct physiological roles of manganese in bacteria. FEMS Microbiology Reviews 45, fuab028. https://doi.org/10.1093/femsre/fuab028

Boyer, E., Bergevin, I., Malo, D., Gros, P., Cellier, M.F.M., 2002. Acquisition of Mn(II) in Addition to Fe(II) Is Required for Full Virulence of *Salmonella enterica* Serovar Typhimurium. Infect Immun 70, 6032–6042. https://doi.org/10.1128/IAI.70.11.6032-6042.2002

Brooke, J.S., 2021. Advances in the Microbiology of Stenotrophomonas maltophilia. Clin Microbiol Rev 34, e00030-19. https://doi.org/10.1128/CMR.00030-19

Brooke, J.S., 2012. Stenotrophomonas maltophilia: an Emerging Global Opportunistic Pathogen. Clin Microbiol Rev 25, 2–41. https://doi.org/10.1128/CMR.00019-11

Calvopiña, K., Dulyayangkul, P., Heesom, K.J., Avison, M.B., 2020. TonB-dependent uptake of β-lactam antibiotics in the opportunistic human pathogen Stenotrophomonas maltophilia. Mol Microbiol 113, 492–503. https://doi.org/10.1111/mmi.14434

Čapek, J., Večerek, B., 2023. Why is manganese so valuable to bacterial pathogens? Front. Cell. Infect. Microbiol. 13, 943390. https://doi.org/10.3389/fcimb.2023.943390

Cellier, M., 2001. Polyphyletic origins of bacterial Nramp transporters. Trends in Genetics 17, 365–370. https://doi.org/10.1016/S0168-9525(01)02364-2

Champion, O.L., Karlyshev, A., Cooper, I.A.M., Ford, D.C., Wren, B.W., Duffield, M., Oyston, P.C.F., Titball, R.W., 2011. Yersinia pseudotuberculosis mntH functions in intracellular manganese accumulation, which is essential for virulence and survival in cells expressing functional Nramp1. Microbiology 157, 1115–1122. https://doi.org/10.1099/mic.0.045807-0

Chandrangsu, P., Rensing, C., Helmann, J.D., 2017. Metal homeostasis and resistance in bacteria. Nat Rev Microbiol 15, 338–350. https://doi.org/10.1038/nrmicro.2017.15

Chang, Y.-T., Lin, C.-Y., Chen, Y.-H., Hsueh, P.-R., 2015. Update on infections caused by Stenotrophomonas maltophilia with particular attention to resistance mechanisms and therapeutic options. Front. Microbiol. 6. https://doi.org/10.3389/fmicb.2015.00893

Chauviat, A., Abrouk, D., Brothier, E., Muller, D., Meyer, T., Favre-Bonté, S., 2025. Genomic and phylogenetic re-assessment of the genus Stenotrophomonas: Description of Stenotrophomonas

thermophila sp. nov., and the proposal of Parastenotrophomonas gen. Nov., Pseudostenotrophomonas gen. Nov., Pedostenotrophomonas gen. Nov., and Allostenotrophomonas gen. Nov. Syst Appl Microbiol 48, 126630. https://doi.org/10.1016/j.syapm.2025.126630

Collet, J.-F., Cho, S.-H., Iorga, B.I., Goemans, C.V., 2020. How the assembly and protection of the bacterial cell envelope depend on cysteine residues. Journal of Biological Chemistry 295, 11984–11994. https://doi.org/10.1074/jbc.REV120.011201

Contreras-Moreira, B., Vinuesa, P., 2013. GET_HOMOLOGUES, a Versatile Software Package for Scalable and Robust Microbial Pangenome Analysis. Appl Environ Microbiol 79, 7696–7701. https://doi.org/10.1128/AEM.02411-13

Crossman, L.C., Gould, V.C., Dow, J.M., Vernikos, G.S., Okazaki, A., Sebaihia, M., Saunders, D., Arrowsmith, C., Carver, T., Peters, N., Adlem, E., Kerhornou, A., Lord, A., Murphy, L., Seeger, K., Squares, R., Rutter, S., Quail, M.A., Rajandream, M.-A., Harris, D., Churcher, C., Bentley, S.D., Parkhill, J., Thomson, N.R., Avison, M.B., 2008. The complete genome, comparative and functional analysis of Stenotrophomonas maltophiliareveals an organism heavily shielded by drug resistance determinants. Genome Biol 9, R74. https://doi.org/10.1186/gb-2008-9-4-r74

Denet, E., Vasselon, V., Burdin, B., Nazaret, S., Favre-Bonté, S., 2018. Survival and growth of Stenotrophomonas maltophilia in free-living amoebae (FLA) and bacterial virulence properties. PLoS ONE 13, e0192308. https://doi.org/10.1371/journal.pone.0192308

Dudek, C.-A., Jahn, D., 2022. PRODORIC: state-of-the-art database of prokaryotic gene regulation. Nucleic Acids Research 50, D295–D302. https://doi.org/10.1093/nar/gkab1110

DuMont, A.L., Karaba, S.M., Cianciotto, N.P., 2015. Type II Secretion-Dependent Degradative and Cytotoxic Activities Mediated by Stenotrophomonas maltophilia Serine Proteases StmPr1 and StmPr2. Infect Immun 83, 3825–3837. https://doi.org/10.1128/IAI.00672-15

Eddy, S.R., 2011. Accelerated Profile HMM Searches. PLoS Comput Biol 7, e1002195. https://doi.org/10.1371/journal.pcbi.1002195

Erill, I., 2015. Every Site Counts: Submitting Transcription Factor-Binding Site Information through the CollecTF Portal. J Bacteriol 197, 2454–2457. https://doi.org/10.1128/JB.00031-15

García, C.A., Alcaraz, E.S., Franco, M.A., Passerini De Rossi, B.N., 2015. Iron is a signal for Stenotrophomonas maltophilia biofilm formation, oxidative stress response, OMPs expression, and virulence. Front. Microbiol. 6. https://doi.org/10.3389/fmicb.2015.00926

Hoang, T.T., Karkhoff-Schweizer, R.R., Kutchma, A.J., Schweizer, H.P., 1998. A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked Pseudomonas aeruginosa mutants. Gene 212, 77–86. https://doi.org/10.1016/S0378-1119(98)00130-9

Hohle, T.H., O'Brian, M.R., 2014. Magnesium-dependent processes are targets of bacterial manganese toxicity. Molecular Microbiology 93, 736–747. https://doi.org/10.1111/mmi.12687

Hood, M.I., Skaar, E.P., 2012. Nutritional immunity: transition metals at the pathogen–host interface. Nat Rev Microbiol 10, 525–537. https://doi.org/10.1038/nrmicro2836

Hubert, T., Madec, M., Schalk, I.J., 2023. Experimental and computational methods to highlight behavioural variations in TonB-dependent transporter expression in Pseudomonas aeruginosa versus siderophore concentration. Sci Rep 13, 20015. https://doi.org/10.1038/s41598-023-46585-z

Imlay, J.A., 2014. The Mismetallation of Enzymes during Oxidative Stress. Journal of Biological Chemistry 289, 28121–28128. https://doi.org/10.1074/jbc.R114.588814

Irving, H., Williams, R.J.P., 1953. 637. The stability of transition-metal complexes. J. Chem. Soc. 3192–3210. https://doi.org/10.1039/JR9530003192

Juttukonda, L.J., Skaar, E.P., 2015. Manganese homeostasis and utilization in pathogenic bacteria. Molecular Microbiology 97, 216–228. https://doi.org/10.1111/mmi.13034

Kadokura, H., Beckwith, J., 2009. Detecting Folding Intermediates of a Protein as It Passes through the

Bacterial Translocation Channel. Cell 138, 1164–1173.
https://doi.org/10.1016/j.cell.2009.07.030

Kalidasan, V., Joseph, N., Kumar, S., Awang Hamat, R., Neela, V.K., 2018. Iron and Virulence in Stenotrophomonas Maltophilia: All We Know So Far. Front. Cell. Infect. Microbiol. 8, 401. https://doi.org/10.3389/fcimb.2018.00401

Kehl-Fie, T.E., Zhang, Y., Moore, J.L., Farrand, A.J., Hood, M.I., Rathi, S., Chazin, W.J., Caprioli, R.M., Skaar, E.P., 2013. MntABC and MntH Contribute to Systemic Staphylococcus aureus Infection by Competing with Calprotectin for Nutrient Manganese. Infect Immun 81, 3395–3405. https://doi.org/10.1128/IAI.00420-13

Kehres, D.G., Zaharik, M.L., Finlay, B.B., Maguire, M.E., 2000. The NRAMP proteins of *Salmonella typhimurium* and *Escherichia coli* are selective manganese transporters involved in the response to reactive oxygen. Molecular Microbiology 36, 1085–1100. https://doi.org/10.1046/j.1365-2958.2000.01922.x

LaBauve, A.E., Wargo, M.J., 2012. Growth and Laboratory Maintenance of *Pseudomonas aeruginosa*. CP Microbiology 25. https://doi.org/10.1002/9780471729259.mc06e01s25

Landeta, C., Boyd, D., Beckwith, J., 2018. Disulfide bond formation in prokaryotes. Nat Microbiol 3, 270–280. https://doi.org/10.1038/s41564-017-0106-2

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9, 357–359. https://doi.org/10.1038/nmeth.1923

Liao, C.-H., Lu, H.-F., Yang, C.-W., Yeh, T.-Y., Lin, Y.-T., Yang, T.-C., 2024. HemU and TonB1 contribute to hemin acquisition in Stenotrophomonas maltophilia. Front Cell Infect Microbiol 14, 1380976. https://doi.org/10.3389/fcimb.2024.1380976

Liao, Y., Smyth, G.K., Shi, W., 2019. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. Nucleic Acids Research 47, e47–e47. https://doi.org/10.1093/nar/gkz114

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15, 550. https://doi.org/10.1186/s13059-014-0550-8

Makhlynets, O., Boal, A.K., Rhodes, D.V., Kitten, T., Rosenzweig, A.C., Stubbe, J., 2014. Streptococcus sanguinis class Ib ribonucleotide reductase: high activity with both iron and manganese cofactors and structural insights. J Biol Chem 289, 6259–6272. https://doi.org/10.1074/jbc.M113.533554

Martin, J.E., Imlay, J.A., 2011. The alternative aerobic ribonucleotide reductase of Escherichia coli, NrdEF, is a manganese-dependent enzyme that enables cell replication during periods of iron starvation. Mol Microbiol 80, 319–334. https://doi.org/10.1111/j.1365-2958.2011.07593.x

Martin, J.E., Waters, L.S., 2022. Regulation of Bacterial Manganese Homeostasis and Usage During Stress Responses and Pathogenesis. Front. Mol. Biosci. 9, 945724. https://doi.org/10.3389/fmolb.2022.945724

Martin, J.L., 1995. Thioredoxin —a fold for all reasons. Structure 3, 245–250. https://doi.org/10.1016/S0969-2126(01)00154-X

Martínez-García, E., Fraile, S., Algar, E., Aparicio, T., Velázquez, E., Calles, B., Tas, H., Blázquez, B., Martín, B., Prieto, C., Sánchez-Sampedro, L., Nørholm, M.H.H., Volke, D.C., Wirth, N.T., Dvořák, P., Alejaldre, L., Grozinger, L., Crowther, M., Goñi-Moreno, A., Nikel, P.I., Nogales, J., de Lorenzo, V., 2023. SEVA 4.0: an update of the Standard European Vector Architecture database for advanced analysis and programming of bacterial phenotypes. Nucleic Acids Research 51, D1558–D1567. https://doi.org/10.1093/nar/gkac1059

Mavridou, D.A.I., Saridakis, E., Kritsiligkou, P., Mozley, E.C., Ferguson, S.J., Redfield, C., 2014. An Extended Active-site Motif Controls the Reactivity of the Thioredoxin Fold. Journal of Biological Chemistry 289, 8681–8696. https://doi.org/10.1074/jbc.M113.513457

Mikhailovich, V., Heydarov, R., Zimenkov, D., Chebotar, I., 2024. Stenotrophomonas maltophilia virulence: a current view. Front. Microbiol. 15, 1385631. https://doi.org/10.3389/fmicb.2024.1385631

Morey, J.R., McDevitt, C.A., Kehl-Fie, T.E., 2015. Host-imposed manganese starvation of invading pathogens: two routes to the same destination. Biometals 28, 509–519. https://doi.org/10.1007/s10534-015-9850-z

Murdoch, C.C., Skaar, E.P., 2022. Nutritional immunity: the battle for nutrient metals at the host-pathogen interface. Nat Rev Microbiol 20, 657–670. https://doi.org/10.1038/s41579-022-00745-6

Nas, M.Y., White, R.C., DuMont, A.L., Lopez, A.E., Cianciotto, N.P., 2019. Stenotrophomonas maltophilia Encodes a VirB/VirD4 Type IV Secretion System That Modulates Apoptosis in Human Cells and Promotes Competition against Heterologous Bacteria, Including Pseudomonas aeruginosa. Infect Immun 87, e00457-19. https://doi.org/10.1128/IAI.00457-19

Nilewski, S., Varatnitskaya, M., Masuch, T., Kusnezowa, A., Gellert, M., Baumann, A.F., Lupilov, N., Kusnezow, W., Koch, M.-H., Eisenacher, M., Berkmen, M., Lillig, C.H., Leichert, L.I., 2021. Functional metagenomics of the thioredoxin superfamily. J Biol Chem 296, 100247. https://doi.org/10.1074/jbc.RA120.016350

Ochoa-Sánchez, L.E., Vinuesa, P., 2017. Evolutionary Genetic Analysis Uncovers Multiple Species with Distinct Habitat Preferences and Antibiotic Resistance Phenotypes in the Stenotrophomonas maltophilia Complex. Front. Microbiol. 8, 1548. https://doi.org/10.3389/fmicb.2017.01548

Ogunniyi, A.D., Mahdi, L.K., Jennings, M.P., McEwan, A.G., McDevitt, C.A., Van Der Hoek, M.B., Bagley, C.J., Hoffmann, P., Gould, K.A., Paton, J.C., 2010. Central Role of Manganese in Regulation of Stress Responses, Physiology, and Metabolism in *Streptococcus pneumoniae*. J Bacteriol 192, 4489–4497. https://doi.org/10.1128/JB.00064-10

Osman, D., Martini, M.A., Foster, A.W., Chen, J., Scott, A.J.P., Morton, R.J., Steed, J.W., Lurie-Luke, E., Huggins, T.G., Lawrence, A.D., Deery, E., Warren, M.J., Chivers, P.T., Robinson, N.J., 2019. Bacterial sensors define intracellular free energies for correct enzyme metalation. Nat Chem Biol 15, 241–249. https://doi.org/10.1038/s41589-018-0211-4

Ouyang, Z., He, M., Oman, T., Yang, X.F., Norgard, M.V., 2009. A manganese transporter, BB0219 (BmtA), is required for virulence by the Lyme disease spirochete, *Borrelia burgdorferi*. Proc. Natl. Acad. Sci. U.S.A. 106, 3449–3454. https://doi.org/10.1073/pnas.0812999106

Palmer, L.D., Skaar, E.P., 2016. Transition Metals and Virulence in Bacteria. Annu. Rev. Genet. 50, 67–91. https://doi.org/10.1146/annurev-genet-120215-035146

Pan, S.-Y., Shih, Y.-L., Huang, H.-H., Li, L.-H., Lin, Y.-T., Yang, T.-C., 2022. The involvement of PacIRA system of Stenotrophomonas maltophilia in the uptake of Pseudomonas aeruginosa pyochelin and intraspecies competition for iron acquisition. Journal of Microbiology, Immunology and Infection 55, 273–281. https://doi.org/10.1016/j.jmii.2021.03.001

Patzer, S.I., Hantke, K., 2001. Dual Repression by Fe^{2+}-Fur and Mn^{2+}-MntR of the *mntH* Gene, Encoding an NRAMP-Like Mn^{2+} Transporter in *Escherichia coli*. J Bacteriol 183, 4806–4813. https://doi.org/10.1128/JB.183.16.4806-4813.2001

Peracino, B., Buracco, S., Bozzaro, S., 2013. The Nramp (Slc11) proteins regulate development, resistance to pathogenic bacteria and iron homeostasis in Dictyostelium discoideum. J Cell Sci 126, 301–311. https://doi.org/10.1242/jcs.116210

Pi, H., Wendel, B.M., Helmann, J.D., 2020. Dysregulation of Magnesium Transport Protects Bacillus subtilis against Manganese and Cobalt Intoxication. J Bacteriol 202. https://doi.org/10.1128/JB.00711-19

Pompilio, A., Ciavardelli, D., Crocetta, V., Consalvo, A., Zappacosta, R., Di Ilio, C., Di Bonaventura, G., 2014. Stenotrophomonas maltophilia Virulence and Specific Variations in Trace Elements during Acute Lung Infection: Implications in Cystic Fibrosis. PLoS ONE 9, e88769. https://doi.org/10.1371/journal.pone.0088769

Porcheron, G., Garénaux, A., Proulx, J., Sabri, M., Dozois, C.M., 2013. Iron, copper, zinc, and manganese transport and regulation in pathogenic Enterobacteria: correlations between strains, site of infection and the relative importance of the different metal transport systems for

virulence. Front. Cell. Infect. Microbiol. 3. https://doi.org/10.3389/fcimb.2013.00090

Que, Q., Helmann, J.D., 2000. Manganese homeostasis in *Bacillus subtilis* is regulated by MntR, a bifunctional regulator related to the diphtheria toxin repressor family of proteins. Molecular Microbiology 35, 1454–1468. https://doi.org/10.1046/j.1365-2958.2000.01811.x

Rello, J., Kalwaje Eshwara, V., Lagunes, L., Alves, J., Wunderink, R.G., Conway-Morris, A., Rojas, J.N., Alp, E., Zhang, Z., 2019. A global priority list of the TOp TEn resistant Microorganisms (TOTEM) study at intensive care: a prioritization exercise based on multi-criteria decision analysis. Eur J Clin Microbiol Infect Dis 38, 319–323. https://doi.org/10.1007/s10096-018-3428-y

Rivera, J., Valerdi-Negreros, J.C., Vázquez-Enciso, D.M., Argueta-Zepeda, F.-S., Vinuesa, P., 2024. Phylogenomic, structural, and cell biological analyses reveal that *Stenotrophomonas maltophilia* replicates in acidified Rab7A-positive vacuoles of *Acanthamoeba castellanii*. Microbiol Spectr 12, e02988-23. https://doi.org/10.1128/spectrum.02988-23

Ryan, R.P., Monchy, S., Cardinale, M., Taghavi, S., Crossman, L., Avison, M.B., Berg, G., Van Der Lelie, D., Dow, J.M., 2009. The versatility and adaptation of bacteria from the genus Stenotrophomonas. Nat Rev Microbiol 7, 514–525. https://doi.org/10.1038/nrmicro2163

Schauer, K., Rodionov, D.A., De Reuse, H., 2008. New substrates for TonB-dependent transport: do we only see the 'tip of the iceberg'? Trends in Biochemical Sciences 33, 330–338. https://doi.org/10.1016/j.tibs.2008.04.012

Sendra, K.M., Barwinska-Sendra, A., Mackenzie, E.S., Baslé, A., Kehl-Fie, T.E., Waldron, K.J., 2023. An ancient metalloenzyme evolves through metal preference modulation. Nat Ecol Evol 7, 732–744. https://doi.org/10.1038/s41559-023-02012-0

Sengupta, S., Mondal, A., Dutta, D., Parrack, P., 2018. HflX protein protects Escherichia coli from manganese stress. J Biosci 43, 1001–1013. https://doi.org/10.1007/s12038-018-9807-9

Shih, Y.-L., Wu, C.-M., Lu, H.-F., Li, L.-H., Lin, Y.-T., Yang, T.-C., 2022. Involvement of the *hemP-hemA-smlt0796-smlt0797* Operon in Hemin Acquisition by Stenotrophomonas maltophilia. Microbiol Spectr 10, e00321-22. https://doi.org/10.1128/spectrum.00321-22

Shin, H.E., Pan, C., Curran, D.M., Bateman, T.J., Chong, D.H.Y., Ng, D., Shah, M., Moraes, T.F., 2024. Prevalence of Slam-dependent hemophilins in Gram-negative bacteria. J Bacteriol 206, e0002724. https://doi.org/10.1128/jb.00027-24

Si, M., Zhao, C., Burkinshaw, B., Zhang, B., Wei, D., Wang, Y., Dong, T.G., Shen, X., 2017. Manganese scavenging and oxidative stress response mediated by type VI secretion system in *Burkholderia thailandensis*. Proc. Natl. Acad. Sci. U.S.A. 114. https://doi.org/10.1073/pnas.1614902114

Sievers, F., Higgins, D.G., 2018. Clustal Omega for making accurate alignments of many protein sequences. Protein Science 27, 135–145. https://doi.org/10.1002/pro.3290

Silale, A., Van Den Berg, B., 2023. TonB-Dependent Transport Across the Bacterial Outer Membrane. Annu. Rev. Microbiol. 77, 67–88. https://doi.org/10.1146/annurev-micro-032421-111116

Smets, D., Loos, M.S., Karamanou, S., Economou, A., 2019. Protein Transport Across the Bacterial Plasma Membrane by the Sec Pathway. Protein J 38, 262–273. https://doi.org/10.1007/s10930-019-09841-8

Sobota, J.M., Imlay, J.A., 2011. Iron enzyme ribulose-5-phosphate 3-epimerase in Escherichia coli is rapidly damaged by hydrogen peroxide but can be protected by manganese. Proc Natl Acad Sci U S A 108, 5402–5407. https://doi.org/10.1073/pnas.1100410108

Song, F., Zhang, G., Li, H., Ma, L., Yang, N., 2022. Comparative transcriptomic analysis of Stenotrophomonas sp. MNB17 revealed mechanisms of manganese tolerance at different concentrations and the role of histidine biosynthesis in manganese removal. Ecotoxicology and Environmental Safety 244, 114056. https://doi.org/10.1016/j.ecoenv.2022.114056

Song, M., Kim, J.-S., Liu, L., Husain, M., Vázquez-Torres, A., 2016. Antioxidant Defense by Thioredoxin Can Occur Independently of Canonical Thiol-Disulfide Oxidoreductase Enzymatic Activity. Cell Rep 14, 2901–2911. https://doi.org/10.1016/j.celrep.2016.02.066

Sullivan, M.J., Terán, I., Goh, K.G.K., Ulett, G.C., 2024. Resisting death by metal: metabolism and Cu/Zn homeostasis in bacteria. Emerging Topics in Life Sciences 8, 45–56. https://doi.org/10.1042/ETLS20230115

Takahashi, I., Hosomi, K., Nagatake, T., Toubou, H., Yamamoto, D., Hayashi, I., Kurashima, Y., Sato, S., Shibata, N., Goto, Y., Maruyama, F., Nakagawa, I., Kuwae, A., Abe, A., Kunisawa, J., Kiyono, H., 2020. Persistent colonization of non-lymphoid tissue-resident macrophages by *Stenotrophomonas maltophilia*. International Immunology 32, 133–141. https://doi.org/10.1093/intimm/dxz071

Taudte, N., German, N., Zhu, Y.-G., Grass, G., Rensing, C., 2016. Restoration of growth by manganese in a mutant strain of Escherichia coli lacking most known iron and manganese uptake systems. Biometals 29, 433–450. https://doi.org/10.1007/s10534-016-9927-3

The UniProt Consortium, 2017. UniProt: the universal protein knowledgebase. Nucleic Acids Res 45, D158–D169. https://doi.org/10.1093/nar/gkw1099

Uppalapati, S.R., Vazquez-Torres, A., 2022. Manganese Utilization in Salmonella Pathogenesis: Beyond the Canonical Antioxidant Response. Front Cell Dev Biol 10, 924925. https://doi.org/10.3389/fcell.2022.924925

Vinuesa, P., Argueta-Zepeda, F.-S., Valerdi-Negreros, J.C., Vázquez-Enciso, D.M., Osorio-Herrera, M.D., Gutiérrez-Hernández, D., Mendoza-Gómez, K., Rivera, J., 2026. Complete genome sequence of the environmental Stenotrophomonas maltophilia ESTM1D_MKCAZ16_6a (Sm18) strain recovered from sediments of a clean river in Morelos. In revision.

Vinuesa, P., Ochoa-Sánchez, L.E., Contreras-Moreira, B., 2018. GET_PHYLOMARKERS, a Software Package to Select Optimal Orthologous Clusters for Phylogenomics and Inferring Pan-Genome Phylogenies, Used for a Critical Geno-Taxonomic Revision of the Genus Stenotrophomonas. Front. Microbiol. 9, 771. https://doi.org/10.3389/fmicb.2018.00771

Waldron, K.J., Robinson, N.J., 2009. How do bacterial cells ensure that metalloproteins get the correct metal? Nat Rev Microbiol 7, 25–35. https://doi.org/10.1038/nrmicro2057

Wang, N.E., Courcelle, E.J., Coltman, S.M., Spolek, R.L., Courcelle, J., Courcelle, C.T., 2023. Manganese transporters regulate the resumption of replication in hydrogen peroxide-stressed Escherichia coli. Biometals 36, 1361–1376. https://doi.org/10.1007/s10534-023-00523-8

Waters, L.S., 2020. Bacterial manganese sensing and homeostasis. Current Opinion in Chemical Biology 55, 96–102. https://doi.org/10.1016/j.cbpa.2020.01.003

Waters, L.S., Sandoval, M., Storz, G., 2011. The Escherichia coli MntR Miniregulon Includes Genes Encoding a Small Protein and an Efflux Pump Required for Manganese Homeostasis. J Bacteriol 193, 5887–5897. https://doi.org/10.1128/JB.05872-11

Wright, Z., Seymour, M., Paszczak, K., Truttmann, T., Senn, K., Stilp, S., Jansen, N., Gosz, M., Goeden, L., Anantharaman, V., Aravind, L., Waters, L.S., 2024. The small protein MNTS evolved from a signal peptide and acquired a novel function regulating manganese homeostasis in *Escherichia coli*. Molecular Microbiology 121, 152–166. https://doi.org/10.1111/mmi.15206

Wu, C.-M., Li, L.-H., Lin, Y.-L., Wu, C.-J., Lin, Y.-T., Yang, T.-C., 2022. The *sbiTRS* Operon Contributes to Stenobactin-Mediated Iron Utilization in Stenotrophomonas maltophilia. Microbiol Spectr 10, e02673-22. https://doi.org/10.1128/spectrum.02673-22

Xue, K., Song, X., Zhang, W., Zhang, Y., Chen, T., Cao, Z., Hong, F., Zhang, X., 2024. Omics-Based Investigation on Mechanisms Controlling Cellular Internalization of Keratin Monomers during Biodegradation by Stenotrophomonas maltophilia DHHJ. Microb Physiol 34, 170–181. https://doi.org/10.1159/000540072

Yeh, T.-Y., Lu, H.-F., Li, L.-H., Lin, Y.-T., Yang, T.-C., 2025. Contribution of fepAsm, fciABC, sbaA, sbaBCDEF, and feoB to ferri-stenobactin acquisition in Stenotrophomonas maltophilia KJ. BMC Microbiol 25, 91. https://doi.org/10.1186/s12866-025-03792-0

Yu, Y., Li, Y.-P., Ren, K., Hao, X., Fru, E.C., Rønn, R., Rivera, W.L., Becker, K., Feng, R., Yang, J., Rensing, C., 2024. A brief history of metal recruitment in protozoan predation. Trends in Microbiology 32, 465–476. https://doi.org/10.1016/j.tim.2023.11.008

Zaharik, M.L., Cullen, V.L., Fung, A.M., Libby, S.J., Kujat Choy, S.L., Coburn, B., Kehres, D.G., Maguire, M.E., Fang, F.C., Finlay, B.B., 2004. The *Salmonella enterica* Serovar Typhimurium Divalent Cation Transport Systems MntH and SitABCD Are Essential for Virulence in an *Nramp1*[G169] Murine Typhoid Model. Infect Immun 72, 5522–5525. https://doi.org/10.1128/IAI.72.9.5522-5525.2004

Zeinert, R., Martinez, E., Schmitz, J., Senn, K., Usman, B., Anantharaman, V., Aravind, L., Waters, L.S., 2018. Structure–function analysis of manganese exporter proteins across bacteria. J Biol Chem 293, 5715–5730. https://doi.org/10.1074/jbc.M117.790717

Zhang, T., Wu, H., Ma, C., Yang, Y., Li, H., Yang, Z., Zhou, S., Shi, D., Chen, T., Yang, D., Li, J., Jin, M., 2024. Emergence of colistin-resistant Stenotrophomonas maltophilia with high virulence in natural aquatic environments. Science of The Total Environment 933, 173221. https://doi.org/10.1016/j.scitotenv.2024.173221

Evaluated contrasts

group1	group2	p.adj	p.adj.signif
0	8	1.101202e−02	*
0	16	1.470991e−04	***
0	32	4.379815e−07	****
4	16	1.101202e−02	*
4	32	1.470991e−04	***
8	32	1.101202e−02	*

group1	group2	p.adj	p.adj.signif
0	8	1.293498e−03	**
0	16	1.864407e−06	****
4	16	1.293498e−03	**

pwc: **Dunn test**; p.adjust: **Benjamini-Hochberg**

A

PC2: 10% variance

PC1: 64% variance

MM
Metal concentration
▲ Fe²⁺ 10 μM
● Fe²⁺ 0 μM

Mn²⁺ 8 μM
Mn²⁺ 0 μM

B

DEG Overlap Analysis

Mn²⁺ 0 μM vs Mn²⁺ 8μM   Fe²⁺ 10 μM vs Fe²⁺ 0μM

Fe10   Fe0   Mn8   Mn0

5   3

mntP   84

0   0

0   1

mntH   9

3   29

0   0

0   0

V8P27_001285
V8P27_001286
V8P27_001249
V8P27_001267
V8P27_001268
V8P27_002357   ◄
V8P27_002358   ◄
V8P27_003990
V8P27_003991

C

DEG: Mn²⁺ 0 vs. Mn²⁺ 8μM

MM-Fe10

Log₂FC

mntP

mntH

Chr Sm18

MM-Fe0

Log₂FC

mntP

mntH

TBDR-Trx fold operon

Chr Sm18

1Mbp   2Mbp   3Mbp   4Mbp

Molecular Funtion
- (GO: 0022857) Transmembrane transporter
- (GO: 0016531) Copper chaperone
- (GO: 0016491) Oxidoreductase
- Others

MM-Fe10   **MM-Fe0**

Log₂FC

Mn2+/Fe2+ uptake protein, MntH (2360)
Copper chaperone SCO1/SenC (3797)
DUF6436 domain-containing protein (3804)
HTH-type transcription Factor (3805)
Manganese efflux pump, MntP (4008)
propionate–CoA ligase, PreE (0822)
Methionyl aminopeptidase, map (1248)
Uncharacterized protein (1249)
CDF inner membrane transporter (1267)
Endonuclease V, nfi (1268)
TonB-dependent receptor (1285)
MerC domain-containing protein (1286)
TonB-dependent receptor (2357)
Thioredoxin fold protein (2358)
Mn2+/Fe2+ uptake protein, MntH (2360)
SMR inner membrane transporter (2373)
OMF of RND efflux pump (3889)
MFP of RND efflux pump (3890)
RND flux pump (3991)
MFS Inner membrane transporter (4004)
DcaP outer membrane transporter (4005)
Manganese efflux pump, MntP (4008)

A

B

MntR-box

signal peptide

signal peptide

MntR-box

2357

2358   mntR

mntH   2361

50 bp

```
tatttatatagccgttgctatatttataa        ccaaggacgacgccttcatgccc        tatagcaggagctatattcggccccatgaac
ataaatatatcggcaacgatataaatatt        ggttcctgctgcggaagtacggg        atatcgtcctcgatataagccggggtacttg
```

-30   MntR-box   -10

RBS

1
Met Pro
2357
SP

MntR-box

1
Met Asn
mntH

C

1 kb

4006   4007   mntP   4009

MntR-box
Riboswitch

54 bp

```
tatagccatcgctata        tgctaagcttcgcgtcccttggggagtagccggattccttcgcaggaatcgtccacgtcaacatactcggccggtgcgccgtggcgtggacaa
atatcggtagcgatat        acgattcgaagcgcaggggaacccctcatcggcctaaggaagcgtccttagcaggtgcagttgtatgagccggccacgcggcaccgcacctgtt
```

MntR-box   yybP-ykoY leader

```
ccatgatggttggcgagaccagcggcccgcgcgcgcaacgtcaggttgggcgcgagcgcaggccgtgcgtccgtcctggcccgacctcagccgctgtccatgtcc
ggtactaccaaccgctctggtcgccgggcgcgcgcgttgcagtccaacccgcgctcgcgtccggcacgcaggcaggaccgggctggagtcggcgacaggtacagg
```

yybP-ykoY leader

1
Met Ser
mntP

A

Kruskal-Wallis, $X^2 = 24.9$, $p = 1.5e^{-4}$, $n = 54$ $X^2 = 47.1$, $p = 5.4e^{-9}$, $n = 54$ $X^2 = 36.9$, $p = 6.2e^{-7}$, $n = 54$

Sm18/327−2357_2358

group1	group2	p.adj	p.adj.signif
Fe0_Mn0	Fe0_Mn0.08	0.0013072582	**
Fe0_Mn0	Fe0_Mn8	0.0013072582	**
Fe0_Mn0	Fe10_Mn0	0.0001318014	***
Fe0_Mn0	Fe10_Mn0.08	0.0015753157	**
Fe0_Mn0	Fe10_Mn8	0.0013072582	**

Sm18/327−mntH

group1	group2	p.adj	p.adj.signif
Fe0_Mn0	Fe0_Mn8	1.688007e−06	****
Fe0_Mn0	Fe10_Mn0	1.790375e−02	*
Fe0_Mn0	Fe10_Mn0.08	1.025144e−03	**
Fe0_Mn0	Fe10_Mn8	5.960094e−07	****
Fe0_Mn0.08	Fe0_Mn8	2.948261e−04	***
Fe0_Mn0.08	Fe10_Mn0.08	3.106638e−02	*
Fe0_Mn8	Fe10_Mn0	1.007357e−04	***
Fe10_Mn0	Fe10_Mn8	2.078660e−02	*
Fe10_Mn0	Fe10_Mn8	1.077084e−02	*

Sm18/327−mntP

group1	group2	p.adj	p.adj.signif
Fe0_Mn0	Fe0_Mn8	0.0016648267	**
Fe0_Mn0	Fe10_Mn8	0.0001353533	***
Fe0_Mn0.08	Fe0_Mn8	0.0061074939	**
Fe0_Mn0.08	Fe10_Mn0	0.0005486729	***
Fe0_Mn8	Fe10_Mn0	0.0010184605	**
Fe0_Mn8	Fe10_Mn0.08	0.0086319308	**
Fe10_Mn0	Fe10_Mn8	0.0001186073	***
Fe10_Mn0.08	Fe10_Mn8	0.0007528110	***

Fe concentration (μM)
- 0
- 10

B

$X^2 = 29.9$, $p = 1.5e^{-6}$, $n = 36$ $X^2 = 23.8$, $p = 2.8e^{-5}$, $n = 36$ $X^2 = 21.6$, $p = 8.0e^{-5}$, $n = 36$

group1	group2	p.adj	p.adj.signif
Sm18_0	Sm18_8	1.257391e−02	*
Sm18_0	Sm18ΔmnR_0	7.548757e−03	**
Sm18_8	Sm18ΔmnR_0	2.920875e−07	****
Sm18_8	Sm18ΔmnR_8	7.548757e−03	**
Sm18ΔmnR_0	Sm18ΔmnR_8	1.257391e−02	*

group1	group2	p.adj	p.adj.signif
Sm18_0	Sm18_8	1.771485e−02	*
Sm18_8	Sm18ΔmnR_0	2.183668e−05	****
Sm18_8	Sm18ΔmnR_8	8.694281e−04	***

group1	group2	p.adj	p.adj.signif
Sm18_0	Sm18_8	0.0077857539	**
Sm18_8	Sm18ΔmnR_0	0.0005108788	***
Sm18_8	Sm18ΔmnR_8	0.0001406068	***

Genetic background
- Sm18
- Sm18ΔmnR

AUC of normalized RFU (y-axis)

Mn concentration (μM) (x-axis)

pwc: **Dunn test**; p.adjust: **Benjamini-Hochberg**

A

| 0 | 0.2 | 0.5 |

pH 6.0

pH 7.0

Population density (OD$_{600nm}$)

Time (h)

B

$X^2 = 15.8$, $p = 7.3e^{-3}$, $n = 18$ $X^2 = 15.9$, $p = 7.1e^{-3}$, $n = 18$

| pH 6.0 | pH 7.0 |

AUC of OD$_{600nm}$

H$_2$O$_2$ concentration (μM)

Strain
- Sm18/332
- Sm18VIMmntH/332
- Sm18VIMmntH/332::mntH

Evaluated contrasts

pH 6.0

group1	group2	p.adj	p.adj.signif
Sm18/332_0	Sm18VIMmntH/332_500	0.01978453	*
Sm18VIMmntH/332_0	Sm18VIMmntH/332_500	0.02144954	*

pH 7.0

group1	group2	p.adj	p.adj.signif
Sm18/332_0	Sm18VIMmntH/332_500	0.03433498	*
Sm18VIMmntH/332_0	Sm18VIMmntH/332_200	0.03658535	*
Sm18VIMmntH/332_0	Sm18VIMmntH/332_500	0.03262763	*

C

| 0 | 10 | 20 |
| 40 | 80 | 160 |

Population density (OD$_{600nm}$)

Time (h)

D

Kruskal-Wallis, $X^2 = 28.0$, $p = 3.2e^{-3}$, $n = 36$

AUC of OD$_{600nm}$

Mn concentration (μM)

Strain
- Sm18/332
- Sm18ΔmntP/332
- Sm18ΔmntP/332::mntP

Evaluated contrasts

group1	group2	p.adj	p.adj.signif
Sm18ΔmntP/332_0	Sm18ΔmntP/332_160	0.008247813	**
Sm18ΔmntP/332_0	Sm18ΔmntP/332_40	0.037335551	*
Sm18ΔmntP/332_0	Sm18ΔmntP/332_80	0.016093707	*

pwc: **Dunn** test; p.adjust: **Benjamini-Hochberg**