# Manganese Homeostasis Drives *Stenotrophomonas maltophilia* Oxidative Stress Defense and Replication in *Acanthamoeba castellanii* phagosomes

Fulvia-Stefany Argueta-Zepeda[1,2], Javier Rivera[1], Julio César Valerdi-Negreros[1,3] https://orcid.org/0000-0001-7536-5483, Christopher Rensing[4] https://orcid.org/0000-0002-5012-7953, and Pablo Vinuesa[1] https://orcid.org/0000-0001-6119-2956

[1]Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México (UNAM), Cuernavaca, Morelos, México

[2]Programa de Maestría y Doctorado en Ciencias Bioquímicas, UNAM, Mexico City, Mexico

[3]Programa de Doctorado en Ciencias Biomédicas, UNAM, Mexico City, Mexico

[4]Fujian Agriculture and Forestry University, Fuzhou, China.

Address correspondence to Pablo Vinuesa, vinuesa@ccg.unam.mx

Keywords: metal-homeostasis, transcriptomics, host-microbe interactions, intracellular replication, nutritional immunity, free-living amoebae

## ABSTRACT

Manganese homeostasis is essential for the environmental adaptability and pathogenic potential of *Stenotrophomonas maltophilia*, a bacterium that thrives across diverse and fluctuating niches. Here, we characterize the manganese homeostasis network of *S. maltophilia* strain Sm18, revealing a coordinated system that integrates conserved transporters with previously unrecognized components. Central to this system is an MntR-controlled miniregulon that includes the canonical $Mn^{2+}$ importer MntH and exporter MntP, together with a TonB-dependent receptor (TBDR) and a periplasmic thioredoxin-fold protein (pTFP), both defining novel protein families with restricted phylogenetic distribution. Transcriptomic analyses under varying $Mn^{2+}$ and $Fe^{2+}$ conditions uncovered a tight interplay between these metals, highlighting the ferrophilic nature of *S. maltophilia* and the differential regulation of miniregulon components. Notably, the TBDR-pTFP module is strongly induced under combined $Mn^{2+}$ and $Fe^{2+}$ limitation, suggesting a specialized role in metal acquisition under nutrient-restricted conditions. Functional analyses demonstrated that MntP is required to prevent Mn toxicity even at sub-inhibitory concentrations, whereas MntH supports growth under oxidative stress and promotes intracellular replication within *Acanthamoeba castellanii* phagosomes. Together, these findings define a previously unrecognized Mn-responsive module that expands the MntR regulatory network and provides new insight into the mechanisms that enable *S. maltophilia* to adapt to metal-limited and host-associated environments.

## 1. Introduction

*S. maltophilia* is a gram-negative, non-fermentative bacterium recognized for its ecological versatility and opportunistic pathogenicity (Brooke, 2021). It thrives in diverse environments, including soil, water, plant rhizospheres, free-living amoebae, and diverse animal organs and cells, demonstrating remarkable adaptability to contrasting ecological niches (Ochoa-Sánchez and Vinuesa, 2017; Rivera et al., 2024; Ryan et al., 2009). These bacteria can survive harsh conditions, such as oligotrophic or metal-rich environments, and are highly resistant to disinfectants and multiple drugs due to robust biofilm formation and multiple efflux systems. Recent research has highlighted these adaptive mechanisms, showing how they enhance the resilience and persistence of *S. maltophilia* in nosocomial environments (Brooke, 2021, 2012), making it one of the ten most prevalent microorganisms in hospital care units (Rello et

al., 2019). This pathogen primarily affects immunocompromised individuals, leading to respiratory tract, bloodstream, and urinary tract infections (Brooke, 2021), and poses a significant challenge in clinical settings because of its intrinsic resistance to multiple antibiotics, including carbapenems (Chang et al., 2015; Ochoa-Sánchez and Vinuesa, 2017). However, little is known about the genes and mechanisms involved in the adaptation of *Stenotrophomonas* to host-associated and intracellular lifestyles in diverse host tissues and phagocytic cells, including free-living amoebae (Denet et al., 2018; Rivera et al., 2024) and mammalian macrophages (Takahashi et al., 2020). In recent years, mounting evidence has revealed that professional phagocytes subject ingested bacteria to limiting or toxic concentrations of transition metals, inhibiting their proliferation in phagosomes, a process known as nutritional immunity (Murdoch and Skaar, 2022).

Homeostasis of essential transition metals, such as manganese, iron, copper, and zinc, is crucial for various physiological processes (Palmer and Skaar, 2016; Waldron and Robinson, 2009). Metallostasis is maintained by a network of allosteric metalloregulators, metal transporters, metallochaperones, and efflux pumps (Porcheron et al., 2013). The concerted action of these components regulates the intracellular metal concentrations, ensuring that they are available for essential functions without reaching toxic levels (Chandrangsu et al., 2017). The ability of bacteria to rapidly adapt to different environments is tightly coupled to their metal homeostasis capabilities, particularly during host infection, as metal concentrations vary widely across organs, tissues, and cellular compartments (Kehl-Fie et al., 2013; Pompilio et al., 2014).

Studies on *S. maltophilia* metallostasis have focused on iron homeostasis, highlighting its importance in bacterial virulence (Mikhailovich et al., 2024). In contrast, our knowledge of the molecular mechanisms underlying manganese homeostasis in *Stenotrophomonas* is limited to a single transcriptomic study performed on *Stenotrophomonas* sp. MNB17, an organism isolated from a deep-sea mud sample in the western Pacific that removes $Mn^{2+}$ by efficiently forming Mn-precipitates ($MnCO_3$ and Mn oxides) at high Mn concentrations (Song et al., 2022). However, manganese homeostasis has been well studied in some Gram-positive and Gram-negative bacteria (Bosma et al., 2021; Waters, 2020). A crucial component is the manganese transporter protein MntH, which facilitates manganese uptake into the cytoplasm under limiting conditions (Kehres et al., 2000). Conversely, the conserved MntP exporter removes excess manganese, thereby maintaining intracellular levels within the optimal range (Waters et al., 2011). This process is regulated by various regulatory proteins and signaling pathways that sense intracellular manganese levels and adjust the expression or activity of

manganese transporters (Bosma et al., 2021; Martin and Waters, 2022; Sengupta et al.,
2018; Wright et al., 2024), with MntR being the master regulator (Que and Helmann, 2000;
Waters et al., 2011). Furthermore, under manganese limitation and oxidative stress, a $Mn^{2+}$-
scavenging pathway has been identified in *Burkholderia thailandensis*, comprising a newly
discovered TonB-dependent outer membrane (OM) manganese transporter, MnoT, and a
$Mn^{2+}$-binding protein, TseM, secreted by a type VI secretion pathway (T6SS) which is
apparently unique to *Burkholderia* species (Si et al., 2017). The requirement for $Mn^{2+}$ and $Fe^{2+}$
varies greatly among bacterial species (Bosma et al., 2021). *S. maltophilia* features an iron-
centric metabolism that requires iron for optimal growth and virulence (Kalidasan et al., 2018;
Mikhailovich et al., 2024). However, $Mn^{2+}$ becomes essential under oxidative stress or low
iron availability (Anjem et al., 2009), conditions that mirror the environments that *S.*
*maltophilia* likely encounters as an intracellular opportunistic pathogen in phagosomes
subjected to nutritional immunity (Čapek and Večerek, 2023; Murdoch and Skaar, 2022). This
bacterium has been reported to survive in human epidermoid epithelial cells (Zhang et al.,
2024), human lung epithelial cells (DuMont et al., 2015; Nas et al., 2019), free-living amoebae
(Denet et al., 2018; Rivera et al., 2024), and mammalian macrophages (Nas et al., 2019;
Takahashi et al., 2020). The divalent $Mn^{2+}/Fe^{2+}$ transporter Nramp1 protein, located in the
phagosomal membrane and ubiquitous in eukaryotes, plays a crucial role in nutritional
immunity by depleting phagocytosed bacteria of these micronutrients. Interestingly, its
bacterial homolog, MntH, is also essential for countering this stress by importing $Mn^{2+}$ into
bacterial cells (Cellier, 2001; Juttukonda and Skaar, 2015).
Here, we investigated the manganese homeostasis network of *S. Maltophilia* Sm18 to address
the limited understanding of this process in the genus. Through an integrated transcriptomic,
phylogenomic and functional approach, we identified the core components of an MntR-
controlled miniregulon, including the canonical transporters *mntH* and *mntP*, as well as a
previously uncharacterized bicistronic locus encoding a TonB-dependent receptor and a
thioredoxin-fold protein. We further assessed the contribution of these components to
bacterial fitness under metal stress conditions and within a biologically relevant intracellular
context. In particular, we demonstrate that *mntH* is expressed during interaction with *A.*
*castellani*i and is required for optimal intracellular replication, linking manganese acquisition to
adaptation to host-associated environments.
120
121

## 2. Material and methods

### 2.1 Bacterial strains and culture conditions

*S. maltophilia* ESTM1D_MKCAZ16_6a (Sm18) is an environmental isolate from river sediments in Morelos, Mexico (Ochoa-Sánchez and Vinuesa, 2017) with a fully sequenced genome (Vinuesa et al., 2026c). Sm18 was routinely cultivated aerobically in Lysogeny Broth (LB) at 30°C or in modified MOPS minimal medium (MM) supplemented with.4 % (w/v) glucose (LaBauve and Wargo, 2012). The MM was optimized for *S. maltophilia* growth by adding thiamine 0.5%, biotin 0.1% and 0.2% (w/v) casamino acids (CAA). Manganese was not included in the micronutrient stock of MM; however, trace amounts of metals, including iron and potentially manganese, may be present due to the addition of casamino acids, as indicated by the manufacturer. The addition of casamino acids was required to support robust growth of Sm18 strains carrying transcriptional fusions, which did not grow reproducibly in MOPS minimal medium supplemented only with methionine. For the growth kinetics, LB was supplemented with 10mM of MOPS buffer (pH 6.8) and antibiotics, as required.

### 2.2 Generation of deletion and vector integration mutants

Markless deletion mutants were generated using a two-step allelic exchange strategy (Huang and Wilks, 2017). Briefly, ~ 500-600 bp DNA fragments flanking the target genes were PCR-amplified from *S. maltophilia* Sm18 genomic DNA and assembled into the suicide vector pEX18Tc (Hoang et al., 1998) using NEBuilder HiFi DNA Assembly (NEB), following the manufacturer's instructions. The resulting constructs were transferred into Sm18 by conjugation from *E. coli* S17-1, as previously described (Simon et al., 1983). Transconjugants (merodiploids) were selected on LB agar supplemented with tetracycline (20 µg/mL) and gentamicin (30 µg/mL). Double recombinants were obtained by counter-selection on LB agar without NaCl and supplemented with 10% sucrose. Markerless deletions were confirmed by antibiotic sensitivity, colony PCR, and amplicon sequencing. This approach was used to generate deletion mutants of *orf02357*, *orf02358*, *mntP* and *mntR*. For *mntH*, repeated attempts to obtain a deletion mutant were unsuccessful. Therefore, a vector integration mutant (VIM) strategy was used to disrupt gene function. Briefly, an internal fragment of the target gene was PCR-amplified and cloned into pEX18TcVIM-GFP, followed by conjugation into Sm18 as described above. This approach was also applied to selected genes (*mntP* and *orf02357*) to validate that VIM and deletion mutants produced comparable

phenotypes (Supplementary Fig. S1). All primers and constructs used in this study are listed
in Supplementary Tables A4-A6.

*2.3 Construction of Sm18 expression plasmids*

Expression plasmids were constructed based on the standardized SEVA modular architecture
(Martínez-García et al., 2023). Derivatives of pSEVA327 (oriVRK2, CmR) and pSEVA337R
(oriVpBBR1, CmR) were used to generate transcriptional reporter fusions. Intergenic regions
upstream of *orf02357*, *mntH,* and *mntP* were PCR-amplified from Sm18 genomic DNA using
Phusion polymerase (NEB). The 5'-end UTR found upstream of *orf2357* also controls the
transcription of *orf2358*, which is co-expressed with the preceding gene and separated by a
short intergenic distance. Amplicons were directionally cloned upstream of promoterless GFP
or mCherry reporter genes using standard restriction-ligation procedures. The resulting
constructs (327-02357_02358, 327-*mntH,* 327-*mntP,* 337R-02357_02358, 337R-*mntH,* and
337R-*mntP*) are listed in Supplementary Tables A4-A6.

*2.4 Fluorescence kinetics and microbial growth*

Fluorescence and growth kinetics of strains were assessed in at least five and eight
independent biological replicates, respectively, using 96-well microplates. Cultures were
adjusted to an initial $OD_{600}$ of 0.05 in LB and 0.01 in minimal MOPS (MM) medium, in a final
volume of 200 μL supplemented with varying Mn and Fe concentrations. Plates were
incubated at 30 °C with orbital shaking (205 rpm), and $OD_{600}$ measurements were recorded
every 30-45 min using a BioTek Epoch 2 microplate spectrophotometer. Fluorescence (GFP;
excitation/emission: 479/520 nm) and $OD_{600}$ were measured in parallel under identical
conditions using a BioTek Synergy H1 microplate reader. Uninoculated media were used as
blanks, and strains carrying empty plasmids served as controls. Relative fluorescence units
(RFU) were normalized to $OD_{600}$ previous statistical analysis. Growth and fluorescence
kinetics were analyzed using the area under the curve (AUC), calculated with the
Growthcurver R package (v0.3.1). Statistical comparisons between conditions were
performed using the non-parametric Kruskal-Wallis test followed by Dunn's post hoc test with
Bonferroni or Benjamini-Hochberg correction for multiple testing. Bootstrapping was used to
estimate 95% confidence intervals. All analyses were conducted in R (v4.3.2) using standard
packages (ggplot2, ggpubr, rstatix). Error bars represent the standard error of the mean from
biological and technical replicates.

## 2.5 Cell sample preparation for RNA-sequencing

Three independent biological replicates were generated from glycerol stocks by streaking on LB agar and incubating at 30 °C. A single colony from each replicate was inoculated into 3 mL of MM and grown at 30 °C with shaking (205 rpm) for 24 h. Precultures (1:50 dilution) were then established in 25 mL MM and incubated under the same conditions for 14 h. Cells were harvested by centrifugation, washed three times with MM without added iron (MM-Fe0), and used to inoculate fresh MM at an initial $OD_{600}$ of ~0.005. Cultures were grown in media supplemented with defined concentrations of $Mn^{2+}$ and $Fe^{2+}$, denoted as Fe0Mn0, Fe0Mn8, Fe10Mn0, and Fe10Mn8, where the numbers indicate the micromolar concentration (µM) of each metal added (e.g., MM-Fe0Mn8 indicates 0 µM $Fe^{2+}$ and 8 µM $Mn^{2+}$). Fe0 and Mn0 refer to conditions without added metals; however, trace amounts may still be present due to the composition of the basal medium (see Section 2.1). All experimental conditions were prepared using the same basal medium, such that relative differences in gene expression and phenotypes reflect the defined supplementation of $Fe^{2+}$ and $Mn^{2+}$. Cultures were grown to early exponential phase ($OD_{600}$ 0.1–0.2), then split into two technical replicates and harvested by centrifugation. Cell pellets were immediately stabilized in RNA later for RNA sequencing (GENEWIZ, Azenta Life Sciences; Illumina HiSeq 2500, 150 bp paired-end).

## 2.6 RNA-seq analysis data and gene expression

Raw reads were quality-checked using FastQC (v0.12.1) (Andrews, S., 2010) and trimmed with Trimmomatic (v0.39) (Bolger et al., 2014). Clean reads (~135 bp; >23 million paired-end reads per sample) were aligned to the *S. maltophilia* Sm18 reference genome (Vinuesa et al., 2026c) using Bowtie2 (v2.4.4) (Langmead and Salzberg, 2012), achieving alignment rates above 96%. Gene-level counts were obtained using featureCounts from the Rsubread package (v2.14.2) (Liao et al., 2019), with >92% of reads successfully assigned to annotated features. Differential gene expression analysis was performed using DESeq2 (v1.42.0) (Love et al., 2014), applying median-of-ratios normalization and a negative binomial model. Adjusted p-values were calculated using the Benjamini-Hochberg method, and genes with Padj < 0.01 and $|\log_2$ fold change$| \geq 1$ were considered differentially expressed. Functional annotation of differentially expressed genes was performed using UniProt (The UniProt Consortium, 2017), InterProScan (Blum et al., 2025), and MOTIF search tools.

## 2.7 Identification of transcription factor binding sites and protein homologs

Putative MntR-binding motifs were identified using MEME (v 5.5.5) (Bailey et al., 2015) on
the 5'-UTR regions of *Stenotrophomonas mntH* (V8P27_002360) orthologs identified with
GET_HOMOLOGUES (Contreras-Moreira and Vinuesa, 2013). Motifs ranging from 16 to 25
nt were searched, and the resulting position-specific probability matrices (PSPMs) were
compared against experimentally validated transcription factor-binding sites, using TOMTOM
with the CollecTF (Erill, 2015) and PRODORIC (Dudek and Jahn, 2022) database. Motif
occurrences were further identified using MAST across representative genomes of
*Stenotrophomonas* and related genera in the *Lysobacteriaceae* (Chauviat et al., 2025;
Vinuesa et al., 2018). Homologs of MntR miniregulon components were identified by
hmmsearch (v3.4) (44) using the gathering threshold (--cut_ga) option for calibrated profile
hidden Markov models (HMMs) retrieved from Pfam (PF01566 for MntH, PF02659 for MntP),
or built locally with hmmbuild (v3.4) from clustal omega (Sievers and Higgins, 2018)
alignments (with –iter 2 and –use-kimura options) of our curated datasets of MntR,
V8P27_002357, and V8P27_002358 homologs. The latter three HMMs and the PSPM for the
*Lisobacteraceae* MntR-box are available for download from
https://github.com/vinuesa/supplementary_materials/.

*2.8 Phylogenomic and motive distribution analysis*

A dataset of 103 RefSeq genomes representing the taxonomic diversity of *Stenotrophomonas*
and related Lysobacteraceae was compiled. Homologs of MntR, MntH, MntP, ORF2357, and
ORF2358 were identified using hmmsearch (HMMER3) with curated profile HMMs. Motif
searches for the MntR-binding site were performed using MAST (MEME suite) with the
consensus motif defined in this study with a $p \leq 5\times10^{-8}$ cutoff. Phylogenetic relationships were
inferred using a maximum likelihood approach based on a concatenated alignment of 155
core proteins identified with GET_PHYLOMARKERS (Vinuesa et al., 2018) from a set of 601
core gene families generated with GET_HOMOLOGUES. Presence-absence patterns of
genes and motif counts were mapped onto the resulting phylogeny.

*2.9 Structural and phylogenomic characterization of miniregulon-associated proteins*

Protein domain architecture and signal peptides were predicted using standard bioinformatic
tools. Structural models were generated using AlphaFold3 (Abramson et al., 2024) and
visualized to uncover potential ligand-binding and other functional features. Homologs of

ORF2357 and ORF2357 were identified using BLASTP and HMMER searches against public
databases, including RefSeq and Reference Proteomes. Retrieved sequences were aligned
using Clustal Omega, and phylogenetic relationships were inferred using maximum likelihood
methods as described above. Synteny analysis was performed by examining the genomic
context of representative homologs across selected taxa. Conserved sequence motifs were
identified from multiple sequence alignments and visualized accordingly.

*2.10 Live-cell imaging (LCI) of of Sm18 expressing transcriptional fusions in co-culture with A.*
*castellanii*

Live-cell imaging assays were performed as previously described (Rivera et al., 2024). Briefly,
5 x $10^5$ trophozoites of *A. castellanii* strain Neff were seeded in 20-mm glass-bottom dishes
and incubated for 1 h at 30 °C in MMsalts-MOPS-Glc medium. *S. maltophilia* Sm18/337R-
*02357_02358*, Sm18/337R-*mntH* and Sm18/337R-*mntP* were grown in LB, washed, and
adjusted to an $OD_{600}$ of ~1.0 (~4.8 × $10^8$ CFU/mL). Bacterial suspensions were added to the
amoeba cultures at a multiplicity of infection (MOI) of 25, followed by centrifugation at 500 × g
for 5 min to synchronize phagocytosis. Live-cell imaging was performed using an AxioVert A1
inverted microscope (Zeiss) at 3 and 18 h post co-culture (ppc).

*2.11 Quantitative evaluation of the intracellular replication capacity of Sm18 mutants*

Intracellular replication assays were performed using a microtiter plate format as previously
described (Rivera et al., 2024). Briefly, 1 x $10^5$ trophozoites of *A. castellanii* (strain Neff) were
seeded per well in MMsalts-MOPS-Glc medium and incubated for 30 min at 25° C. The
medium was then replaced with Sm18VIM*mntH* and Sm18Δ*mntR* bacterial suspensions
tagged with the minitransposon pUC18T_mTn7TC1_Pc_mScarlet-I, which constitutively
expresses the mScarlet-I fluorescent protein from strong Pc promoter, at a MOI of 10.
Phagocytosis was synchronized by centrifugation and plates were incubated statically at 25
°C in a Synergy 2 plate reader. Red fluorescence (excitation/emission: 568/594 nm) was
measured every 30 min for 48 h. Control wells included medium only, trophozoites without
bacteria, and bacteria without trophozoites. All experiments were performed with five
independent biological replicates per strain, each including five technical replicates.
276

# 3 RESULTS

## 3.1 Manganese tolerance of S. maltophilia Sm18 in complex and defined media

To assess the strain's manganese tolerance, we ran 48-hour-long growth kinetics using LB and MM media with increasing $MnCl_2$ concentrations. Strain Sm18 grew in LB and MM up to 16 mM and 8 mM of $MnCl_2$, respectively (Fig. 1A). In both media, the maximal growth rate and total yield of the culture remained unaffected, except at the highest Mn concentration tested. However, statistical analysis of the area under the curve (AUC) for population density ($OD_{600}$) data revealed a significant decrease in growth in both media starting at 8 mM $MnCl_2$, due to the extended lag phase observed with increasing Mn concentrations (Fig. 1B). Consistent with these observations, complete growth inhibition was observed at higher $Mn^{2+}$ concentrations, occurring at 32 mM in LB and at 16 mM in MM medium. Although minimum inhibitory concentrations (MICs) were not determined using standardized assays, these results define the concentration range at which $Mn^{2+}$ becomes inhibitory under the tested conditions. The lower inhibitory thresholds were observed consistently across replicates.

## 3.2 Genome-wide identification of manganese-responsive genes in S. maltophilia Sm18

To identify Sm18 genes involved in $Mn^{2+}$ hemostasis, we designed experimental conditions that consider the well-established interplay between iron and manganese in bacterial metal homeostasis (Čapek and Večerek, 2023). These metals are known to be functionally interconnected, often sharing regulatory networks and compensatory roles under conditions of metal limitation (Helmann, 2014; Puri et al., 2010; Steingard et al., 2023; Taudte et al., 2016). Therefore, varying iron availability alongside manganese allows a more accurate assessment of manganese-dependent regulatory responses. Cultures were established in defined MM medium supplemented with casamino acids under four metal conditions: Fe0Mn0, Fe0Mn8, Fe10Mn0, and Fe10Mn8 (see methods). Principal component analysis (PCA) shown in Fig. 2A demonstrated that replicates within each group were clustered and separated according to metal concentration. PC1 revealed that 64% of the variance was related to varying iron conditions, while PC2 explained 10% of the variance due to manganese concentrations. We identified a total of 124 differentially expressed genes (DEGs) when comparing $Fe^{2+}$ 10 μM vs. $Fe^{2+}$ 0 μM in MM-Mn0, and 89 DEG in MM-Mn8, based on an adjusted *p*-value (padj) < 0.01 and a $\log_2$ fold change $|(\log_2 FC)| \geq 1.0$. In contrast, only five DEGs were detected when comparing $Mn^{2+}$ 0 μM vs. $Mn^{2+}$ 8 μM in MM-Fe10, and 16 DEG in

MM-Fe0. The number of DEGs for each condition is summarized as a Venn diagram (Fig.

2B), revealing that 88 DEGs were regulated solely by iron levels, irrespective of manganese

concentrations in the medium. Although this study focuses on manganese-related responses

modulated by manganese and iron availability, we identified a broader transcriptional

response reflecting the interplay between these two metals. A complementary analysis

specifically addressing iron homeostasis will be presented in a separate study. The *mntP*

gene was the only one found to be exclusively regulated by manganese levels in the medium,

independent of iron availability (Fig. 2B). In contrast, the expression of *mntH* was affected by

any change in Mn or Fe concentration (Fig. 2B). Nine genes displayed differential expression

only when cultures were subjected to simultaneous $Fe^{2+}$ and $Mn^{2+}$ limitation (Fig. 2B). Two

were down-regulated (blue-colored locus tags) under these conditions, whereas the

remaining genes were up-regulated, as detailed below. Figure 2C highlights the DEGs

identified when comparing $Mn^{2+}$ 0 µM vs. $Mn^{2+}$ 8 µM in both MM-Fe10 and MM-Fe0 media

(see Supplementary Data, Tables A1 and A2 for more details). These DEGs are represented

as colored dots on the $log_2FC$ plots, and their corresponding genomic loci are indicated by

green vertical bars mapped to the Sm18 chromosome. Their gene ontologies (GOs) are

highlighted in different colors with the transmembrane transport category (yellow points) being

the most prevalent. Fig. 2C also shows the $log_2FC$ values in a bar plot for a better quantitative

representation, including the protein annotations of each DEG. Loci displaying the most

significant $log_2FC$ changes were selected for further characterization.

*3.3 Genome-wide scanning for potential MntR binding sites*

MntR is a manganese-responsive metalloregulator that controls gene expression by binding

to conserved short inverted palindromic sequences (MntR-boxes) located in the 5′-UTR

regions of target genes (Patzer and Hantke, 2001). Using a motif derived from *mntH*

upstream regions, we identified three MntR-binding sites in the Sm18 genome (Fig. 3A-C). In

addition to the canonical target *mntH,* two additional loci were identified as putative members

of the MntR miniregulon. One binding site was located upstream of the *orf2357-orf2358* locus,

which lies in close proximity to *mntR* and *mntH*, which share a bidirectional promoter (Fig.

3B). Additionally, *orf2357* and *orf2358* are separated by a minimal intergenic region,

consistent with their organization as a putative operon (Fig. 3B). A third MntR-binding site was

identified upstream of *mntP*, where a putative yybP–ykoY riboswitch was also detected (Fig.

3C). Together, these findings identify a set of three loci associated with predicted MntR-

binding sites in the Sm18 genome which also correspond to the genes displaying the highest
differential expression in the transcriptomic analysis. We next examined the phylogenetic
distribution of these loci, considered here as candidate components of a MntR miniregulon of
manganese homeostasis, across the genus *Stenotrophomonas* and related Lysobacteraceae.
346

347 *3.4 Lineage-specific distribution of MntR miniregulon members*
348

To assess the evolutionary conservation of the proposed MntR miniregulon, we analyzed the
distribution of its components across 103 RefSeq genomes representing the diversity of
*Stenotrophomonas* and related *Lysobacteraceae* (Fig. 4). The presence-absence patterns of
MntR, MntH, MntP, ORF2357, and ORF2358, together with the occurrence of predicted
MntR-binding motifs, revealed a non-uniform distribution across the analyzed taxa. Notably,
the complete five-gene set, along with three associated MntR-binding sites, was exclusively
detected in *Stenotrophomonas* species, including *S. maltophilia*. In contrast, closely related
species such as *S. pavanii*,*S. beteli*, and *S. hibiscicola* lacked the TBDR and pTFP homologs
and contained only two predicted MntR-binding motifs (Fig. 4). Furthermore, these genes
were absent from the genomes of *Allostenotrophomonas*, *Pseudostenotrophomonas*, and
*Parastenotrophomonas* (Chauviat et al., 2025). Among the components analyzed, only the
MntP exporter was conserved across all four genera. Mapping these patterns onto the
species phylogeny revealed that the complete (three-loci) MntR miniregulon is restricted to
the *S. maltophilia* complex (Fig. 4). Importantly, the consistent co-occurrence of these genes
with conserved MntR-binding motifs supports the existence of a lineage-specific, coordinately
regulated gene module. These findings prompted us to experimentally validate the metal-
dependent regulation of these loci.
366

*3.5 The MntR miniregulon expression is modulated by iron and manganese*

To validate the metal-dependent regulation of MntR miniregulon elements, we constructed
transcriptional fusions of the 5'-UTRs of *orf2357-orf2358*, *mntH,* and *mntP* to evaluate their
expression in the Sm18 background. Expression data for the Sm18/327-2357_2358 fusion
revealed that  it was exclusively expressed under simultaneous Mn and Fe limitation, with
reporter activity completely abolished by a low $Mn^{2+}$ concentration (80 nM) (Fig. 5A, first
panel). Activity of the Sm18/327-*mntH* fusion was negatively correlated with increasing $Mn^{2+}$
concentrations in the culture medium. However, low activity was still measurable at 8 µM

$Mn^{2+}$, and a significantly higher expression was observed at $Fe^{2+}$ 10 µM compared with $Fe^{2+}$ 0

µM, except when the medium contained 8 µM $Mn^{2+}$ (Fig. 5A, second panel). Finally,

Sm18/327-*mntP* increased significantly at 8 µM $Mn^{2+}$ compared to $Mn^{2+}$ 0µM, irrespective of

Fe availability in the medium (Fig. 5A, third panel). Overall, these results show that the

expression of *orf2357-orf2358*, *mntH,* and *mntP* is differentially modulated by $Mn^{2+}$ and $Fe^{2+}$

availability, with distinct effects on each locus. Notably, the transcriptional responses

observed in these reporter assays are consistent with the trends identified in the RNA-seq

analysis, validating the described regulatory effects of both transition metals on the MntR

miniregulon members.

*3.6* MntR controls core manganese homeostasis genes in Sm18

To confirm that the promoter activity observed in the transcriptional fusions reported in Fig.

5A is regulated by MntR, and to further validate the core MntR miniregulon, the three fusions

were transferred into a Sm18Δ*mntR* strain, and promoter activity was measured in MM-

Fe0Mn0 and MM-Fe0Mn8 medium. The Sm18Δ*mntR*/327-2357_2358 fusion revealed a

significant increase in expression compared with Sm18/327-2357_2358. However, adding

manganese led to reduced expression, following a similar trend to that observed in the Sm18

background, although not to the same extent  (Fig. 5B, first panel). Similarly, derepression of

the 327-*mntH* fusion was observed in the Sm18Δ*mntR* genetic background compared to the

wild-type. However, no statistically significant activity was detected upon addition of 8 µM

$Mn^{2+}$ (Fig. 5B, second panel). Finally, no expression was detected for Sm18Δ*mntR*/327-*mntP*

independent of manganese concentration (Fig. 5B, third panel). Overall, these results indicate

that the expression of *orf2357–orf2358, mntH,* and *mntP* is dependent on MntR, with distinct

expression patterns observed for each locus in the Δ*mntR* background. Notably, *orf2357–*

*orf2358* and *mntH* show increased expression in the absence of MntR, whereas *mntP*

expression is not detected under these conditions.

*3.7 The orf2357-orf2358 locus encodes a novel protein module associated with manganese*

*homeostasis*

Following the identification of *orf2357* and *orf2358* as members of the MntR miniregulon, we

performed a structural and phylogenomic characterization of their encoded proteins.  *Orf2357*

encodes a TonB-dependent receptor (TBDR) with a predicted signal peptide and a canonical

β-barrel architecture containing an N-terminal plug domain (Fig 6A). Notably, this domain

harbors a conserved CQNC motif (Fig. 6B y C). Homology searches revealed a restricted
phylogenetic distribution of closely related sequences, which cluster into a well-supported
lineage characterized by the presence of the CQNC motif. These proteins are sparsely
distributed across diverse bacterial taxa. Structural modeling strongly suggests that this motif,
together with residues from the β-barrel domain, may coordinate a $Mn^{2+}$ ion in a tetrahedral
geometry (Fig. S2). The downstream gene, *orf2358,* encodes a predicted periplasmic
thioredoxin-fold protein (pTFP) (Nilewski et al., 2021). Phylogenomic analysis revealed that
pTFP homologs form a distinct and well-supported clade with a limited taxonomic distribution
(Fig. S3). Notably, these homologs are consistently found in synteny with CQNC-type TBDRs,
reinforcing their functional linkage (Fig 6D). Sequence analysis of pTFP identified a
conserved thioredoxin fold with a non-canonical CXXC motif (CAPC), embedded within a
distinctive lineage-specific sequence signature (Fig. 6E, S3).Together, these findings identify
a novel, MntR-regulated protein module comprising a CQNC-type TBDR and an associated
thioredoxin-fold protein, which are consistently found in synteny, with a restricted phylogenetic
distribution.

*3.8 MntR miniregulon mutants display sensitivity to hydrogen peroxide and manganese*

To determine whether the *mntH* and *mntP* homologs in Sm18 encode functional manganese
importer and exporter proteins, respectively, we constructed mutants of both genes and
evaluated their growth under different stress conditions in MM-Fe0Mn0. The growth kinetics
of the Sm18VIM*mntH* mutant indicated that the *mntH* gene is not essential under $Fe^{2+}$- or
$Mn^{2+}$-limiting conditions in axenic cultures. However, it is required for growth under oxidative
stress induced by hydrogen peroxide ($H_2O_2$), particularly at neutral pH (Fig. 7A). Statistical
analysis revealed that the difference in growth was significant only at 500 µM $H_2O_2$ compared
to the wild type (Fig. 7B).  Strain Sm18Δ*mntP* was challenged with increasing $Mn^{2+}$
concentrations (5-160 µM; Fig. 7C), with statistically significant sensitivity detectable at 40
µM, and complete growth inhibition observed at 160 µM (Fig. 7D). The wild-type phenotypes
of the Sm18VIM-*mntH* and Sm18Δ*mntP* strains were restored by plasmids Sm18VIM-
*mntH*/332::*mntH* and Sm18Δ*mntP*/332::*mntP* provided in *trans*, and carrying the *mntH* and
*mntP* genes under the control of their native promoters (Fig. 7). These results show that *mntH*
and *mntP* play key roles in resistance to $H_2O_2$ and manganese toxicity, respectively, as
supported by the stress-sensitive phenotypes of the mutants and their restoration upon
complementation. No growth phenotypes were detected for the markerless deletion *orf2357*

and *orf2358* mutants under the conditions tested.

*3.9 The S. maltophilia mntH gene is expressed and supports intracellular replication in A.*
*castellanii*

We recently found that *S. maltophilia* Sm18 resists digestion by *A. castellanii* strain Neff
trophozoites and can replicate in acidified, Rab7A-positive *Stenotrophomonas*-containing
vacuoles (SCV) of this free-living professional phagocyte (Rivera et al., 2024). Given that the
phagosomes of *Dictyostelium discoideum* and human macrophages subject bacteria to iron
and manganese limitation via the NRAMP exporter (Morey et al., 2015; Peracino et al., 2013),
we postulated that the Sm18 MntH Mn importer is required to survive in the SCV. To test this
hypothesis, we first evaluated the activity of the Sm18/337-*mntH* fusion in *A. castellanii* strain
Neff trophozoites using wide-field fluorescence microscopy. As shown in Fig. 8A, the fusion
was expressed at 3 and 18 h post-primary contact (ppc). In contrast, equivalent assays
performed with the Sm18/337R-*2357_2358* and Sm18/337R-*mntP* fusions did not reveal
detectable expression under the same conditions. Moreover, the replication capacity of Sm18
in trophozoites was compared with that of the Sm18VIM*mntH* (*VIMmntH*) and Sm18Δ*mntR*
(Δ*mntR*) mutants. As shown in Fig. 8B, the *VIMmntH* mutant had a significantly reduced
replication capacity in Neff trophozoites, whereas that of Δ*mntR* was indistinguishable from
Sm18 (Figs. 8B and 8C). Importantly, no differences in growth were observed among these
strains under *in vitro* conditions in MM or LB (Fig. S5). Together, these observations indicate
that the replication defect of the *VIMmntH* mutant is specific to the intracellular environment.

**4. Discussion**

The ability to maintain manganese homeostasis is critical for bacterial survival under
fluctuating environmental and host-imposed conditions. In this study, we identified and
characterized a regulatory and functional network that supports manganese homeostasis in
*S. maltophilia* Sm18, centered on the metalloregulator MntR and a set of core target genes.
Our results demonstrate that MntR plays a central role in modulating the expression of key
manganese homeostasis genes, including *mntH*, *mntP*, and the novel *orf2357-orf2358* locus.
Transcriptional analyses in the Δ*mntR* background revealed a clear derepression of *mntH* and
the *orf2357-orf2358* putative operon in the absence of manganese, supporting a model in
which MntR acts predominantly as a repressor under Mn-replete conditions. However, the

persistence of manganese-responsive expression patterns in the *ΔmntR* strain suggests that
additional regulatory layers contribute to fine-tuning this system. Thus, while our data
establish MntR as a major regulator, they also indicate that manganese homeostasis in Sm18
is controlled by a more complex regulatory network that remains to be fully elucidated.

Beyond regulation, our functional analyses provide direct evidence for the physiological roles
of MntR-regulated genes. The *VIMmntH* mutant displayed increased sensitivity to oxidative
stress, consistent with the established role of manganese importers in protecting against
reactive oxygen species (ROS). In contrast, deletion of *mntP* resulted in marked sensitivity
even at low manganese concentrations, confirming its role in manganese detoxification. The
restoration of wild-type phenotypes upon genetic complementation further supports the
specific involvement of these genes in manganese homeostasis. Together, these findings
validate the functional relevance of the MntR-controlled gene set and support the existence of
a coherent regulatory module governing manganese uptake and efflux.

In addition to these canonical components, we identified the *orf2357-orf2358* locus as a
previously uncharacterized element associated with manganese homeostasis. Structural and
phylogenomic analyses suggest that *orf2357* encodes a TonB-dependent receptor with a
conserved CQNC motif, while *orf2358* encodes a thioredoxin-fold protein that is consistently
found in synteny with CQNC-type receptors, and might be function as a specialized disulfide
reductase of the CQNC motif. The restricted phylogenetic distribution and conserved genomic
association of these proteins support the existence of a distinct protein module. Although
structural modeling suggests a potential role in manganese coordination, the precise function
of this system remains to be experimentally validated. Therefore, our findings provide a
foundation for future studies aimed at elucidating the mechanistic contribution of this module
to manganese homeostasis.

Importantly, our results extend the relevance of manganese homeostasis to a biologically
meaningful context. We show that *mntH* is expressed during interaction with *A. castellanii* and
is required for optimal intracellular replication. The impaired replication of the *VIMmntH*
mutant in amoebal trophozoites, together with its normal growth under *in vitro* conditions,
indicates that this phenotype is specifically linked to the intracellular environment. These
observations are consistent with previous reports showing that phagosomal compartments
impose metal limitation as an antimicrobial strategy.

In parallel, the increased sensitivity of the *VIMmntH* mutant to $H_2O_2$ observed *in vitro* indicates
that manganese uptake via MntH contributes to oxidative stress resistance. Although

oxidative stress was not directly measured during amoeba infection, these findings support a
model in which MntH-mediated manganese import enhances the bacterial capacity to cope
with host-derived ROS within the phagosomal environment. In this context, manganese likely
acts as a cofactor for antioxidant systems, including catalases and other ROS-detoxifying
enzymes, thereby sustaining bacterial survival under oxidative stress conditions.
Consequently, impaired manganese acquisition in the *VIMmntH* mutant is expected to
compromise these protective mechanisms, leading to reduced intracellular replication. This
integrated model linking manganese homeostasis, oxidative stress defense, and intracellular
survival is summarized in Fig. 9.

Taken together, this study defines a manganese homeostasis network in *S. maltophilia* Sm18
that integrates transcriptional regulation by MntR, functional roles of manganese transport
systems, and a previously unrecognized protein module. While our findings establish the
importance of this system under both environmental and intracellular conditions, they also
highlight the existence of additional regulatory and functional components that warrant further
investigation.

520
## 521 5. Conclusion

In this study, we defined a manganese homeostasis network in *S. maltophilia* Sm18 centered
on the metalloregulator MntR and its target genes. Our results demonstrate that *mntH* and
*mntP* play key roles in oxidative stress resistance and manganese detoxification, respectively,
while also identifying a previously uncharacterized TBDR-pTFP module associated with this
regulatory system. Importantly, we show that manganese acquisition via MntH is required for
optimal intracellular replication in *A. castellanii*, linking metal homeostasis with bacterial
adaptation to host-associated environments. Overall, this work established a functional
connection between manganese homeostasis, oxidative stress defense, and intracellular
survival, providing a framework for understanding how *S. maltophilia* adapts to the metal-
restricted and oxidative environment of phagosomes.

**533 CRediT authorship contribution statement**
534
**Fulvia Stefany Argueta Zepeda:** Writing – original draft,  Writing – review & editing, Validation,
Methodology, Investigation, Formal analysis, Data curation. **Javier Rivera Campos:** Writing – review
& editing, Validation, Methodology. **Julio Valerdi Negreros:** Writing – review & editing, Validation,
Methodology. **Christopher Rensing:** Writing – review & editing, Supervision. **Pablo Vinuesa:**

Writing – review & editing, Writing – original draft, Supervision, Resources, Project administration,
Methodology, Investigation, Funding acquisition, Formal analysis, Conceptualization
541

542 **Conflict of interest**

543     The authors declare no conflict of interest.

544
545

**ACKNOWLEDGEMENTS**

We thank Alfredo Hernández and Víctor del Moral for their expert technical support and
server maintenance at the Centro de Ciencias Genómicas, UNAM. We acknowledge the
excellent service provided by personnel at the Unidad de Síntesis y Secuenciación de DNA
from Instituto de Biotecnología, UNAM. We thank Dr. Ayari Fuentes from CCG-UNAM for
providing access to the multimodal plate reader. We thank Dr. Luis Lozano from the
Bioinformatics Unit at CCG-UNAM for his advice and support during the early phase of
transcriptome analysis. We thank M.Sc. Laura Cervantes for her laboratory assistance and
valuable feedback. We gratefully acknowledge the financial support received from Consejo
Nacional de Humanidades Ciencias y Tecnologías (CONAHCyT México, A1-S-11242),
Secretaría de Ciencia, Humanidades, Tecnología e Innovación (SECIHTI México CBF-2025-
I-1106), and Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica
(PAPIIT), Universidad Nacional Autónoma de México (DGAPA-PAPIIT, UNAM: IN209321 and
IN216424) to PV. F. S. A. Z. and J.C.V.N. were recipients of CONAHCyT scholarships
659308 and 745705, respectively.

**Appendix A. Supporting information**

Supplementary data associated with this article can be found in the online version at
[doi:10.1016/j.micres.2026](doi:10.1016/j.micres.2026) ….

**Data availability:** The complete genome sequence of *S. maltophilia* Sm18 was deposited in
GenBank under accession number CP146374 and the associated genome and transcriptome
sequencing reads were deposited under BioProject PRJNA1081934
[https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1081934](https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1081934). The RefSeq genome
assembly for strain Sm18 is publicly available from NCBI's genome datasets portal at
[https://www.ncbi.nlm.nih.gov/datasets/genome/GCA_053078355.1/](https://www.ncbi.nlm.nih.gov/datasets/genome/GCA_053078355.1/).

The sequence reads generated from the 18 RNAseq samples analyzed in this work were
deposited at NCBI's Sequence Read Archive [https://www.ncbi.nlm.nih.gov/sra/](https://www.ncbi.nlm.nih.gov/sra/) under

accession numbers SRR36898890 to SRR36898907, as part of BioProject PRJNA1081934

https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1081934.

The hidden Markov models built in this work are freely available from GitHub at:

https://github.com/vinuesa/supplementary_materials/tree/main/docs/Argueta-

Zepeda_Stenotrophomonas_MntR_miniregulon.

**Conflict of interest disclosure statement:** None declared.

**Ethics statement:** None required.

# 586 REFERENCES

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung, C.-C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain, R., Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Žídek, A., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., Jumper, J.M., 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature 630, 493–500. https://doi.org/10.1038/s41586-024-07487-w

Anderson, E.S., Paulley, J.T., Gaines, J.M., Valderas, M.W., Martin, D.W., Menscher, E., Brown, T.D., Burns, C.S., Roop, R.M., 2009. The Manganese Transporter MntH Is a Critical Virulence Determinant for *Brucella abortus* 2308 in Experimentally Infected Mice. Infect Immun 77, 3466–3474. https://doi.org/10.1128/IAI.00444-09

Andrews, S., 2010. Fastqc: A quality control tool for high throughput sequence data.

Anjem, A., Varghese, S., Imlay, J.A., 2009. Manganese import is a key element of the OxyR response to hydrogen peroxide in *Escherichia coli*. Molecular Microbiology 72, 844–858. https://doi.org/10.1111/j.1365-2958.2009.06699.x

Bailey, T.L., Johnson, J., Grant, C.E., Noble, W.S., 2015. The MEME Suite. Nucleic Acids Res 43, W39–W49. https://doi.org/10.1093/nar/gkv416

Blanvillain, S., Meyer, D., Boulanger, A., Lautier, M., Guynet, C., Denancé, N., Vasse, J., Lauber, E., Arlat, M., 2007. Plant Carbohydrate Scavenging through TonB-Dependent Receptors: A Feature Shared by Phytopathogenic and Aquatic Bacteria. PLoS ONE 2, e224. https://doi.org/10.1371/journal.pone.0000224

Blum, M., Andreeva, A., Florentino, L.C., Chuguransky, S.R., Grego, T., Hobbs, E., Pinto, B.L., Orr, A., Paysan-Lafosse, T., Ponamareva, I., Salazar, G.A., Bordin, N., Bork, P., Bridge, A., Colwell, L., Gough, J., Haft, D.H., Letunic, I., Llinares-López, F., Marchler-Bauer, A., Meng-Papaxanthos, L., Mi, H., Natale, D.A., Orengo, C.A., Pandurangan, A.P., Piovesan, D., Rivoire, C., Sigrist, C.J.A., Thanki, N., Thibaud-Nissen, F., Thomas, P.D., Tosatto, S.C.E., Wu, C.H., Bateman, A., 2025. InterPro: the protein sequence classification resource in 2025. Nucleic Acids Research 53, D444–D456. https://doi.org/10.1093/nar/gkae1082

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bosma, E.F., Rau, M.H., van Gijtenbeek, L.A., Siedler, S., 2021. Regulation and distinct physiological roles of manganese in bacteria. FEMS Microbiology Reviews 45, fuab028. https://doi.org/10.1093/femsre/fuab028

Boyer, E., Bergevin, I., Malo, D., Gros, P., Cellier, M.F.M., 2002. Acquisition of Mn(II) in Addition to Fe(II) Is Required for Full Virulence of *Salmonella enterica* Serovar Typhimurium. Infect Immun 70, 6032–6042. https://doi.org/10.1128/IAI.70.11.6032-6042.2002

Brooke, J.S., 2021. Advances in the Microbiology of Stenotrophomonas maltophilia. Clin Microbiol Rev 34, e00030-19. https://doi.org/10.1128/CMR.00030-19

Brooke, J.S., 2012. Stenotrophomonas maltophilia: an Emerging Global Opportunistic Pathogen. Clin Microbiol Rev 25, 2–41. https://doi.org/10.1128/CMR.00019-11

Calvopiña, K., Dulyayangkul, P., Heesom, K.J., Avison, M.B., 2020. TonB-dependent uptake of β-lactam antibiotics in the opportunistic human pathogen Stenotrophomonas maltophilia. Mol Microbiol 113, 492–503. https://doi.org/10.1111/mmi.14434

Čapek, J., Večerek, B., 2023. Why is manganese so valuable to bacterial pathogens? Front. Cell. Infect. Microbiol. 13, 943390. https://doi.org/10.3389/fcimb.2023.943390

Cellier, M., 2001. Polyphyletic origins of bacterial Nramp transporters. Trends in Genetics 17, 365–370. https://doi.org/10.1016/S0168-9525(01)02364-2

Champion, O.L., Karlyshev, A., Cooper, I.A.M., Ford, D.C., Wren, B.W., Duffield, M., Oyston, P.C.F., Titball, R.W., 2011. Yersinia pseudotuberculosis mntH functions in intracellular manganese accumulation, which is essential for virulence and survival in cells expressing functional Nramp1. Microbiology 157, 1115–1122. https://doi.org/10.1099/mic.0.045807-0

Chandrangsu, P., Rensing, C., Helmann, J.D., 2017. Metal homeostasis and resistance in bacteria. Nat Rev Microbiol 15, 338–350. https://doi.org/10.1038/nrmicro.2017.15

Chang, Y.-T., Lin, C.-Y., Chen, Y.-H., Hsueh, P.-R., 2015. Update on infections caused by Stenotrophomonas maltophilia with particular attention to resistance mechanisms and therapeutic options. Front. Microbiol. 6. https://doi.org/10.3389/fmicb.2015.00893

Chauviat, A., Abrouk, D., Brothier, E., Muller, D., Meyer, T., Favre-Bonté, S., 2025. Genomic and phylogenetic re-assessment of the genus Stenotrophomonas: Description of Stenotrophomonas thermophila sp. nov., and the proposal of Parastenotrophomonas gen. Nov., Pseudostenotrophomonas gen. Nov., Pedostenotrophomonas gen. Nov., and Allostenotrophomonas gen. Nov. Syst Appl Microbiol 48, 126630. https://doi.org/10.1016/j.syapm.2025.126630

Collet, J.-F., Cho, S.-H., Iorga, B.I., Goemans, C.V., 2020. How the assembly and protection of the bacterial cell envelope depend on cysteine residues. Journal of Biological Chemistry 295, 11984–11994. https://doi.org/10.1074/jbc.REV120.011201

Contreras-Moreira, B., Vinuesa, P., 2013. GET_HOMOLOGUES, a Versatile Software Package for Scalable and Robust Microbial Pangenome Analysis. Appl Environ Microbiol 79, 7696–7701. https://doi.org/10.1128/AEM.02411-13

Crossman, L.C., Gould, V.C., Dow, J.M., Vernikos, G.S., Okazaki, A., Sebaihia, M., Saunders, D., Arrowsmith, C., Carver, T., Peters, N., Adlem, E., Kerhornou, A., Lord, A., Murphy, L., Seeger, K., Squares, R., Rutter, S., Quail, M.A., Rajandream, M.-A., Harris, D., Churcher, C., Bentley, S.D., Parkhill, J., Thomson, N.R., Avison, M.B., 2008. The complete genome, comparative and functional analysis of Stenotrophomonas maltophiliareveals an organism heavily shielded by drug resistance determinants. Genome Biol 9, R74. https://doi.org/10.1186/gb-2008-9-4-r74

Denet, E., Vasselon, V., Burdin, B., Nazaret, S., Favre-Bonté, S., 2018. Survival and growth of Stenotrophomonas maltophilia in free-living amoebae (FLA) and bacterial virulence properties. PLoS ONE 13, e0192308. https://doi.org/10.1371/journal.pone.0192308

Dudek, C.-A., Jahn, D., 2022. PRODORIC: state-of-the-art database of prokaryotic gene regulation. Nucleic Acids Research 50, D295–D302. https://doi.org/10.1093/nar/gkab1110

DuMont, A.L., Karaba, S.M., Cianciotto, N.P., 2015. Type II Secretion-Dependent Degradative and

Cytotoxic Activities Mediated by Stenotrophomonas maltophilia Serine Proteases StmPr1 and StmPr2. Infect Immun 83, 3825–3837. https://doi.org/10.1128/IAI.00672-15

Erill, I., 2015. Every Site Counts: Submitting Transcription Factor-Binding Site Information through the CollecTF Portal. J Bacteriol 197, 2454–2457. https://doi.org/10.1128/JB.00031-15

García, C.A., Alcaraz, E.S., Franco, M.A., Passerini De Rossi, B.N., 2015. Iron is a signal for Stenotrophomonas maltophilia biofilm formation, oxidative stress response, OMPs expression, and virulence. Front. Microbiol. 6. https://doi.org/10.3389/fmicb.2015.00926

Helmann, J.D., 2014. Specificity of Metal Sensing: Iron and Manganese Homeostasis in Bacillus subtilis. Journal of Biological Chemistry 289, 28112–28120. https://doi.org/10.1074/jbc.R114.587071

Hoang, T.T., Karkhoff-Schweizer, R.R., Kutchma, A.J., Schweizer, H.P., 1998. A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked Pseudomonas aeruginosa mutants. Gene 212, 77–86. https://doi.org/10.1016/S0378-1119(98)00130-9

Hohle, T.H., O'Brian, M.R., 2014. Magnesium-dependent processes are targets of bacterial manganese toxicity. Molecular Microbiology 93, 736–747. https://doi.org/10.1111/mmi.12687

Hood, M.I., Skaar, E.P., 2012. Nutritional immunity: transition metals at the pathogen–host interface. Nat Rev Microbiol 10, 525–537. https://doi.org/10.1038/nrmicro2836

Huang, W., Wilks, A., 2017. A rapid seamless method for gene knockout in Pseudomonas aeruginosa. BMC Microbiol 17, 199. https://doi.org/10.1186/s12866-017-1112-5

Hubert, T., Madec, M., Schalk, I.J., 2023. Experimental and computational methods to highlight behavioural variations in TonB-dependent transporter expression in Pseudomonas aeruginosa versus siderophore concentration. Sci Rep 13, 20015. https://doi.org/10.1038/s41598-023-46585-z

Imlay, J.A., 2014. The Mismetallation of Enzymes during Oxidative Stress. Journal of Biological Chemistry 289, 28121–28128. https://doi.org/10.1074/jbc.R114.588814

Irving, H., Williams, R.J.P., 1953. 637. The stability of transition-metal complexes. J. Chem. Soc. 3192–3210. https://doi.org/10.1039/JR9530003192

Juttukonda, L.J., Skaar, E.P., 2015. Manganese homeostasis and utilization in pathogenic bacteria. Molecular Microbiology 97, 216–228. https://doi.org/10.1111/mmi.13034

Kadokura, H., Beckwith, J., 2009. Detecting Folding Intermediates of a Protein as It Passes through the Bacterial Translocation Channel. Cell 138, 1164–1173. https://doi.org/10.1016/j.cell.2009.07.030

Kalidasan, V., Joseph, N., Kumar, S., Awang Hamat, R., Neela, V.K., 2018. Iron and Virulence in Stenotrophomonas Maltophilia: All We Know So Far. Front. Cell. Infect. Microbiol. 8, 401. https://doi.org/10.3389/fcimb.2018.00401

Kehl-Fie, T.E., Zhang, Y., Moore, J.L., Farrand, A.J., Hood, M.I., Rathi, S., Chazin, W.J., Caprioli, R.M., Skaar, E.P., 2013. MntABC and MntH Contribute to Systemic Staphylococcus aureus Infection by Competing with Calprotectin for Nutrient Manganese. Infect Immun 81, 3395–3405. https://doi.org/10.1128/IAI.00420-13

Kehres, D.G., Zaharik, M.L., Finlay, B.B., Maguire, M.E., 2000. The NRAMP proteins of *Salmonella typhimurium* and *Escherichia coli* are selective manganese transporters involved in the response to reactive oxygen. Molecular Microbiology 36, 1085–1100. https://doi.org/10.1046/j.1365-2958.2000.01922.x

LaBauve, A.E., Wargo, M.J., 2012. Growth and Laboratory Maintenance of *Pseudomonas aeruginosa*. CP Microbiology 25. https://doi.org/10.1002/9780471729259.mc06e01s25

Landeta, C., Boyd, D., Beckwith, J., 2018. Disulfide bond formation in prokaryotes. Nat Microbiol 3, 270–280. https://doi.org/10.1038/s41564-017-0106-2

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9, 357–359. https://doi.org/10.1038/nmeth.1923

Liao, C.-H., Lu, H.-F., Yang, C.-W., Yeh, T.-Y., Lin, Y.-T., Yang, T.-C., 2024. HemU and TonB1

contribute to hemin acquisition in Stenotrophomonas maltophilia. Front Cell Infect Microbiol 14, 1380976. https://doi.org/10.3389/fcimb.2024.1380976

Liao, Y., Smyth, G.K., Shi, W., 2019. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. Nucleic Acids Research 47, e47–e47. https://doi.org/10.1093/nar/gkz114

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15, 550. https://doi.org/10.1186/s13059-014-0550-8

Makhlynets, O., Boal, A.K., Rhodes, D.V., Kitten, T., Rosenzweig, A.C., Stubbe, J., 2014. Streptococcus sanguinis class Ib ribonucleotide reductase: high activity with both iron and manganese cofactors and structural insights. J Biol Chem 289, 6259–6272. https://doi.org/10.1074/jbc.M113.533554

Martin, J.E., Imlay, J.A., 2011. The alternative aerobic ribonucleotide reductase of Escherichia coli, NrdEF, is a manganese-dependent enzyme that enables cell replication during periods of iron starvation. Mol Microbiol 80, 319–334. https://doi.org/10.1111/j.1365-2958.2011.07593.x

Martin, J.E., Waters, L.S., 2022. Regulation of Bacterial Manganese Homeostasis and Usage During Stress Responses and Pathogenesis. Front. Mol. Biosci. 9, 945724. https://doi.org/10.3389/fmolb.2022.945724

Martínez-García, E., Fraile, S., Algar, E., Aparicio, T., Velázquez, E., Calles, B., Tas, H., Blázquez, B., Martín, B., Prieto, C., Sánchez-Sampedro, L., Nørholm, M.H.H., Volke, D.C., Wirth, N.T., Dvořák, P., Alejaldre, L., Grozinger, L., Crowther, M., Goñi-Moreno, A., Nikel, P.I., Nogales, J., de Lorenzo, V., 2023. SEVA 4.0: an update of the Standard European Vector Architecture database for advanced analysis and programming of bacterial phenotypes. Nucleic Acids Research 51, D1558–D1567. https://doi.org/10.1093/nar/gkac1059

Mikhailovich, V., Heydarov, R., Zimenkov, D., Chebotar, I., 2024. Stenotrophomonas maltophilia virulence: a current view. Front. Microbiol. 15, 1385631. https://doi.org/10.3389/fmicb.2024.1385631

Morey, J.R., McDevitt, C.A., Kehl-Fie, T.E., 2015. Host-imposed manganese starvation of invading pathogens: two routes to the same destination. Biometals 28, 509–519. https://doi.org/10.1007/s10534-015-9850-z

Murdoch, C.C., Skaar, E.P., 2022. Nutritional immunity: the battle for nutrient metals at the host-pathogen interface. Nat Rev Microbiol 20, 657–670. https://doi.org/10.1038/s41579-022-00745-6

Nas, M.Y., White, R.C., DuMont, A.L., Lopez, A.E., Cianciotto, N.P., 2019. Stenotrophomonas maltophilia Encodes a VirB/VirD4 Type IV Secretion System That Modulates Apoptosis in Human Cells and Promotes Competition against Heterologous Bacteria, Including Pseudomonas aeruginosa. Infect Immun 87, e00457-19. https://doi.org/10.1128/IAI.00457-19

Nilewski, S., Varatnitskaya, M., Masuch, T., Kusnezowa, A., Gellert, M., Baumann, A.F., Lupilov, N., Kusnezow, W., Koch, M.-H., Eisenacher, M., Berkmen, M., Lillig, C.H., Leichert, L.I., 2021. Functional metagenomics of the thioredoxin superfamily. J Biol Chem 296, 100247. https://doi.org/10.1074/jbc.RA120.016350

Ochoa-Sánchez, L.E., Vinuesa, P., 2017. Evolutionary Genetic Analysis Uncovers Multiple Species with Distinct Habitat Preferences and Antibiotic Resistance Phenotypes in the Stenotrophomonas maltophilia Complex. Front. Microbiol. 8, 1548. https://doi.org/10.3389/fmicb.2017.01548

Ogunniyi, A.D., Mahdi, L.K., Jennings, M.P., McEwan, A.G., McDevitt, C.A., Van Der Hoek, M.B., Bagley, C.J., Hoffmann, P., Gould, K.A., Paton, J.C., 2010. Central Role of Manganese in Regulation of Stress Responses, Physiology, and Metabolism in *Streptococcus pneumoniae*. J Bacteriol 192, 4489–4497. https://doi.org/10.1128/JB.00064-10

Osman, D., Martini, M.A., Foster, A.W., Chen, J., Scott, A.J.P., Morton, R.J., Steed, J.W., Lurie-Luke, E., Huggins, T.G., Lawrence, A.D., Deery, E., Warren, M.J., Chivers, P.T., Robinson, N.J., 2019. Bacterial sensors define intracellular free energies for correct enzyme metalation. Nat

Chem Biol 15, 241–249. https://doi.org/10.1038/s41589-018-0211-4

Ouyang, Z., He, M., Oman, T., Yang, X.F., Norgard, M.V., 2009. A manganese transporter, BB0219 (BmtA), is required for virulence by the Lyme disease spirochete, *Borrelia burgdorferi*. Proc. Natl. Acad. Sci. U.S.A. 106, 3449–3454. https://doi.org/10.1073/pnas.0812999106

Palmer, L.D., Skaar, E.P., 2016. Transition Metals and Virulence in Bacteria. Annu. Rev. Genet. 50, 67–91. https://doi.org/10.1146/annurev-genet-120215-035146

Pan, S.-Y., Shih, Y.-L., Huang, H.-H., Li, L.-H., Lin, Y.-T., Yang, T.-C., 2022. The involvement of PacIRA system of Stenotrophomonas maltophilia in the uptake of Pseudomonas aeruginosa pyochelin and intraspecies competition for iron acquisition. Journal of Microbiology, Immunology and Infection 55, 273–281. https://doi.org/10.1016/j.jmii.2021.03.001

Patzer, S.I., Hantke, K., 2001. Dual Repression by $Fe^{2+}$ -Fur and $Mn^{2+}$ -MntR of the *mntH* Gene, Encoding an NRAMP-Like $Mn^{2+}$ Transporter in *Escherichia coli*. J Bacteriol 183, 4806–4813. https://doi.org/10.1128/JB.183.16.4806-4813.2001

Peracino, B., Buracco, S., Bozzaro, S., 2013. The Nramp (Slc11) proteins regulate development, resistance to pathogenic bacteria and iron homeostasis in Dictyostelium discoideum. J Cell Sci 126, 301–311. https://doi.org/10.1242/jcs.116210

Pi, H., Wendel, B.M., Helmann, J.D., 2020. Dysregulation of Magnesium Transport Protects Bacillus subtilis against Manganese and Cobalt Intoxication. J Bacteriol 202. https://doi.org/10.1128/JB.00711-19

Pompilio, A., Ciavardelli, D., Crocetta, V., Consalvo, A., Zappacosta, R., Di Ilio, C., Di Bonaventura, G., 2014. Stenotrophomonas maltophilia Virulence and Specific Variations in Trace Elements during Acute Lung Infection: Implications in Cystic Fibrosis. PLoS ONE 9, e88769. https://doi.org/10.1371/journal.pone.0088769

Porcheron, G., Garénaux, A., Proulx, J., Sabri, M., Dozois, C.M., 2013. Iron, copper, zinc, and manganese transport and regulation in pathogenic Enterobacteria: correlations between strains, site of infection and the relative importance of the different metal transport systems for virulence. Front. Cell. Infect. Microbiol. 3. https://doi.org/10.3389/fcimb.2013.00090

Puri, S., Hohle, T.H., O'Brian, M.R., 2010. Control of bacterial iron homeostasis by manganese. Proc. Natl. Acad. Sci. U.S.A. 107, 10691–10695. https://doi.org/10.1073/pnas.1002342107

Que, Q., Helmann, J.D., 2000. Manganese homeostasis in *Bacillus subtilis* is regulated by MntR, a bifunctional regulator related to the diphtheria toxin repressor family of proteins. Molecular Microbiology 35, 1454–1468. https://doi.org/10.1046/j.1365-2958.2000.01811.x

Rello, J., Kalwaje Eshwara, V., Lagunes, L., Alves, J., Wunderink, R.G., Conway-Morris, A., Rojas, J.N., Alp, E., Zhang, Z., 2019. A global priority list of the TOp TEn resistant Microorganisms (TOTEM) study at intensive care: a prioritization exercise based on multi-criteria decision analysis. Eur J Clin Microbiol Infect Dis 38, 319–323. https://doi.org/10.1007/s10096-018-3428-y

Rivera, J., Valerdi-Negreros, J.C., Vázquez-Enciso, D.M., Argueta-Zepeda, F.-S., Vinuesa, P., 2024. Phylogenomic, structural, and cell biological analyses reveal that *Stenotrophomonas maltophilia* replicates in acidified Rab7A-positive vacuoles of *Acanthamoeba castellanii*. Microbiol Spectr 12, e02988-23. https://doi.org/10.1128/spectrum.02988-23

Ryan, R.P., Monchy, S., Cardinale, M., Taghavi, S., Crossman, L., Avison, M.B., Berg, G., Van Der Lelie, D., Dow, J.M., 2009. The versatility and adaptation of bacteria from the genus Stenotrophomonas. Nat Rev Microbiol 7, 514–525. https://doi.org/10.1038/nrmicro2163

Schauer, K., Rodionov, D.A., De Reuse, H., 2008. New substrates for TonB-dependent transport: do we only see the 'tip of the iceberg'? Trends in Biochemical Sciences 33, 330–338. https://doi.org/10.1016/j.tibs.2008.04.012

Sendra, K.M., Barwinska-Sendra, A., Mackenzie, E.S., Baslé, A., Kehl-Fie, T.E., Waldron, K.J., 2023. An ancient metalloenzyme evolves through metal preference modulation. Nat Ecol Evol 7, 732–744. https://doi.org/10.1038/s41559-023-02012-0

Sengupta, S., Mondal, A., Dutta, D., Parrack, P., 2018. HflX protein protects Escherichia coli from

manganese stress. J Biosci 43, 1001–1013. https://doi.org/10.1007/s12038-018-9807-9

Shih, Y.-L., Wu, C.-M., Lu, H.-F., Li, L.-H., Lin, Y.-T., Yang, T.-C., 2022. Involvement of the *hemP-hemA-smlt0796-smlt0797* Operon in Hemin Acquisition by Stenotrophomonas maltophilia. Microbiol Spectr 10, e00321-22. https://doi.org/10.1128/spectrum.00321-22

Shin, H.E., Pan, C., Curran, D.M., Bateman, T.J., Chong, D.H.Y., Ng, D., Shah, M., Moraes, T.F., 2024. Prevalence of Slam-dependent hemophilins in Gram-negative bacteria. J Bacteriol 206, e0002724. https://doi.org/10.1128/jb.00027-24

Si, M., Zhao, C., Burkinshaw, B., Zhang, B., Wei, D., Wang, Y., Dong, T.G., Shen, X., 2017. Manganese scavenging and oxidative stress response mediated by type VI secretion system in *Burkholderia thailandensis*. Proc. Natl. Acad. Sci. U.S.A. 114. https://doi.org/10.1073/pnas.1614902114

Sievers, F., Higgins, D.G., 2018. Clustal Omega for making accurate alignments of many protein sequences. Protein Science 27, 135–145. https://doi.org/10.1002/pro.3290

Silale, A., Van Den Berg, B., 2023. TonB-Dependent Transport Across the Bacterial Outer Membrane. Annu. Rev. Microbiol. 77, 67–88. https://doi.org/10.1146/annurev-micro-032421-111116

Simon, R., Priefer, U., Pühler, A., 1983. A Broad Host Range Mobilization System for In Vivo Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. Nat Biotechnol 1, 784–791. https://doi.org/10.1038/nbt1183-784

Smets, D., Loos, M.S., Karamanou, S., Economou, A., 2019. Protein Transport Across the Bacterial Plasma Membrane by the Sec Pathway. Protein J 38, 262–273. https://doi.org/10.1007/s10930-019-09841-8

Sobota, J.M., Imlay, J.A., 2011. Iron enzyme ribulose-5-phosphate 3-epimerase in Escherichia coli is rapidly damaged by hydrogen peroxide but can be protected by manganese. Proc Natl Acad Sci U S A 108, 5402–5407. https://doi.org/10.1073/pnas.1100410108

Song, F., Zhang, G., Li, H., Ma, L., Yang, N., 2022. Comparative transcriptomic analysis of Stenotrophomonas sp. MNB17 revealed mechanisms of manganese tolerance at different concentrations and the role of histidine biosynthesis in manganese removal. Ecotoxicology and Environmental Safety 244, 114056. https://doi.org/10.1016/j.ecoenv.2022.114056

Song, M., Kim, J.-S., Liu, L., Husain, M., Vázquez-Torres, A., 2016. Antioxidant Defense by Thioredoxin Can Occur Independently of Canonical Thiol-Disulfide Oxidoreductase Enzymatic Activity. Cell Rep 14, 2901–2911. https://doi.org/10.1016/j.celrep.2016.02.066

Steingard, C.H., Pinochet-Barros, A., Wendel, B.M., Helmann, J.D., 2023. Iron homeostasis in Bacillus subtilis relies on three differentially expressed efflux systems. Microbiology 169. https://doi.org/10.1099/mic.0.001289

Sullivan, M.J., Terán, I., Goh, K.G.K., Ulett, G.C., 2024. Resisting death by metal: metabolism and Cu/Zn homeostasis in bacteria. Emerging Topics in Life Sciences 8, 45–56. https://doi.org/10.1042/ETLS20230115

Takahashi, I., Hosomi, K., Nagatake, T., Toubou, H., Yamamoto, D., Hayashi, I., Kurashima, Y., Sato, S., Shibata, N., Goto, Y., Maruyama, F., Nakagawa, I., Kuwae, A., Abe, A., Kunisawa, J., Kiyono, H., 2020. Persistent colonization of non-lymphoid tissue-resident macrophages by *Stenotrophomonas maltophilia*. International Immunology 32, 133–141. https://doi.org/10.1093/intimm/dxz071

Taudte, N., German, N., Zhu, Y.-G., Grass, G., Rensing, C., 2016. Restoration of growth by manganese in a mutant strain of Escherichia coli lacking most known iron and manganese uptake systems. Biometals 29, 433–450. https://doi.org/10.1007/s10534-016-9927-3

The UniProt Consortium, 2017. UniProt: the universal protein knowledgebase. Nucleic Acids Res 45, D158–D169. https://doi.org/10.1093/nar/gkw1099

Uppalapati, S.R., Vazquez-Torres, A., 2022. Manganese Utilization in Salmonella Pathogenesis: Beyond the Canonical Antioxidant Response. Front Cell Dev Biol 10, 924925. https://doi.org/10.3389/fcell.2022.924925

Vinuesa, P., Argueta-Zepeda, F.-S., Valerdi-Negreros, J.C., Vázquez-Enciso, D.M., Osorio-Herrera,

M.D., Gutiérrez-Hernández, D., Mendoza-Gómez, K., Rivera, J., 2026a. Complete genome sequence of the environmental Stenotrophomonas maltophilia ESTM1D_MKCAZ16_6a (Sm18) strain recovered from sediments of a clean river in Morelos. In revision.

Vinuesa, P., Argueta-Zepeda, F.-S., Valerdi-Negreros, J.C., Vázquez-Enciso, D.M., Osorio-Herrera, M.D., Gutiérrez-Hernández, D., Mendoza-Gómez, K., Rivera, J., 2026b. Complete genome sequence of the environmental Stenotrophomonas maltophilia ESTM1D_MKCAZ16_6a (Sm18) strain recovered from sediments of a clean river in Morelos. In revision.

Vinuesa, P., Argueta-Zepeda, F.-S., Valerdi-Negreros, J.C., Vázquez-Enciso, D.M., Osorio-Herrera, M.D., Gutiérrez-Hernández, D., Mendoza-Gómez, K., Rivera, J., Ochoa-Sánchez, L.E., 2026c. Complete genome sequence of *Stenotrophomonas maltophilia* ESTM1D_MKCAZ16_6a (Sm18), a Mexican environmental multidrug- and amoeba-resistant strain. Microbiol Resour Announc e01304-25. https://doi.org/10.1128/mra.01304-25

Vinuesa, P., Ochoa-Sánchez, L.E., Contreras-Moreira, B., 2018. GET_PHYLOMARKERS, a Software Package to Select Optimal Orthologous Clusters for Phylogenomics and Inferring Pan-Genome Phylogenies, Used for a Critical Geno-Taxonomic Revision of the Genus Stenotrophomonas. Front. Microbiol. 9, 771. https://doi.org/10.3389/fmicb.2018.00771

Waldron, K.J., Robinson, N.J., 2009. How do bacterial cells ensure that metalloproteins get the correct metal? Nat Rev Microbiol 7, 25–35. https://doi.org/10.1038/nrmicro2057

Wang, N.E., Courcelle, E.J., Coltman, S.M., Spolek, R.L., Courcelle, J., Courcelle, C.T., 2023. Manganese transporters regulate the resumption of replication in hydrogen peroxide-stressed Escherichia coli. Biometals 36, 1361–1376. https://doi.org/10.1007/s10534-023-00523-8

Waters, L.S., 2020. Bacterial manganese sensing and homeostasis. Current Opinion in Chemical Biology 55, 96–102. https://doi.org/10.1016/j.cbpa.2020.01.003

Waters, L.S., Sandoval, M., Storz, G., 2011. The Escherichia coli MntR Miniregulon Includes Genes Encoding a Small Protein and an Efflux Pump Required for Manganese Homeostasis. J Bacteriol 193, 5887–5897. https://doi.org/10.1128/JB.05872-11

Wright, Z., Seymour, M., Paszczak, K., Truttmann, T., Senn, K., Stilp, S., Jansen, N., Gosz, M., Goeden, L., Anantharaman, V., Aravind, L., Waters, L.S., 2024. The small protein MNTS evolved from a signal peptide and acquired a novel function regulating manganese homeostasis in *Escherichia coli*. Molecular Microbiology 121, 152–166. https://doi.org/10.1111/mmi.15206

Wu, C.-M., Li, L.-H., Lin, Y.-L., Wu, C.-J., Lin, Y.-T., Yang, T.-C., 2022. The *sbiTRS* Operon Contributes to Stenobactin-Mediated Iron Utilization in Stenotrophomonas maltophilia. Microbiol Spectr 10, e02673-22. https://doi.org/10.1128/spectrum.02673-22

Xue, K., Song, X., Zhang, W., Zhang, Y., Chen, T., Cao, Z., Hong, F., Zhang, X., 2024. Omics-Based Investigation on Mechanisms Controlling Cellular Internalization of Keratin Monomers during Biodegradation by Stenotrophomonas maltophilia DHHJ. Microb Physiol 34, 170–181. https://doi.org/10.1159/000540072

Yeh, T.-Y., Lu, H.-F., Li, L.-H., Lin, Y.-T., Yang, T.-C., 2025. Contribution of fepAsm, fciABC, sbaA, sbaBCDEF, and feoB to ferri-stenobactin acquisition in Stenotrophomonas maltophilia KJ. BMC Microbiol 25, 91. https://doi.org/10.1186/s12866-025-03792-0

Yu, Y., Li, Y.-P., Ren, K., Hao, X., Fru, E.C., Rønn, R., Rivera, W.L., Becker, K., Feng, R., Yang, J., Rensing, C., 2024. A brief history of metal recruitment in protozoan predation. Trends in Microbiology 32, 465–476. https://doi.org/10.1016/j.tim.2023.11.008

Zaharik, M.L., Cullen, V.L., Fung, A.M., Libby, S.J., Kujat Choy, S.L., Coburn, B., Kehres, D.G., Maguire, M.E., Fang, F.C., Finlay, B.B., 2004. The *Salmonella enterica* Serovar Typhimurium Divalent Cation Transport Systems MntH and SitABCD Are Essential for Virulence in an *Nramp1*[G169] Murine Typhoid Model. Infect Immun 72, 5522–5525. https://doi.org/10.1128/IAI.72.9.5522-5525.2004

Zeinert, R., Martinez, E., Schmitz, J., Senn, K., Usman, B., Anantharaman, V., Aravind, L., Waters, L.S., 2018. Structure–function analysis of manganese exporter proteins across bacteria. J Biol Chem 293, 5715–5730. https://doi.org/10.1074/jbc.M117.790717

Zhang, T., Wu, H., Ma, C., Yang, Y., Li, H., Yang, Z., Zhou, S., Shi, D., Chen, T., Yang, D., Li, J., Jin, M., 2024. Emergence of colistin-resistant Stenotrophomonas maltophilia with high virulence in natural aquatic environments. Science of The Total Environment 933, 173221. https://doi.org/10.1016/j.scitotenv.2024.173221

**FIGURE LEGENDS**

**Fig. 1.** Growth kinetics of *S. maltophilia* Sm18 under increasing Mn concentrations. (A) Growth in Lysogeny Broth (LB) and minimal MOPS (MM) medium with increasing millimolar (mM) concentrations of manganese ($Mn^{2+}$). The means of eight biological replicates per concentration are represented by distinct shapes and colors, with error bars indicating the standard error. (B) Statistical analysis of Mn tolerance in Sm18. Differences among groups were assessed using the non-parametric Kruskal-Wallis test, followed by Dunn's *post hoc* test with Benjamini-Hochberg (BH) correction. Significance levels are shown as: ns, $p > 0.05$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ***, $p < 0.0001$. Black asterisks indicate comparisons with the manganese-free control, while red asterisks indicate comparisons between other groups.

**Fig. 2.** Transcriptomic analysis and identification of differentially expressed genes (DEG). Data were obtained from three biological replicates. (A) Principal component analysis (PCA) based on DESeq2-normalized read counts from the full transcriptome ($n$ = 4,059 genes). (B) Venn diagram generated using the VennDiagram package in R, illustrating the overlap of DEG at different metal concentrations. Genes were considered differentially expressed if they met the criteria of adjusted $p$-value (padj) < 0.01 and |$\log_2$ fold change| ≥ 1. (C) DEG from $Mn^{2+}$ 0 µM vs. $Mn^{2+}$ 8 µM comparisons in MM-Fe10 and MM-Fe0 mapped across the Sm18 chromosome (Chr Sm18). Green loci denote regions with significant differential expression. Colored dots represent genes classified by Gene Ontology (GO) functional categories. The adjacent bar plot displays the protein annotations of each DEG.

**Fig. 3.** Core components of the MntR miniregulon in *S. maltophilia* Sm18. (A) Binding motif (MntR-box) for the MntR transcription factor identified using MEME-MAST. The MntR-box profile corresponds to a 25-bp inverted palindromic sequence. (B) Putative manganese uptake operon encoding an outer membrane TonB-dependent receptor and a periplasmic thioredoxin-fold protein, identified by locus tags 2357 and 2358, respectively. The MntR-box is located 75 bp upstream of the putative start codon, between the -10 and -35 promoter

regions. These genes are adjacent to *mntR* (encoding the transcription factor MntR) and
*mntH* (encoding the cytoplasmic Mn importer), which share a 117 bp intergenic region. The
mntH MntR-box is located 10 bp upstream of the putative start codon. (C) The gene encoding
the inner membrane manganese exporter MntP is predicted to be regulated by both the *yybP-*
*ykoY* riboswitch and MntR. Its MntR-box is located 247 bp upstream from the putative start
codon.

**Fig. 4.** Distribution of the MntR miniregulon proteins across *Stenotrophomonas* species and
closely related genera in the Lysobacteraceae family. Maximum-likelihood core-genome
phylogeny estimated with IQTree based on the concatenation of 155 top-scoring protein
families selected with GET_PHYLOMARKERS from 103 Lysobacteraceae RefSeq genomes
(see color code). The tree integrates taxonomic information (four inner rings) with the
presence/absence patterns of the MntR miniregulon genes (*orf02357*, *orf02358, mntH*, *mntR*,
*and mntP*) identified by hmmsearch (five outer rings), and the number of MntR-boxes
detected by MAST scanning (bars). *S. maltophilia* strains are labeled in bold font, with Sm18
highlighted in yellow. Bipartition support values are given as UFBoot bootstrap percentages
only for those ≥ 70%. The scale bar represents the number of expected substitutions per site
under the best-fitting Q.insect+F+R4 model, as selected by BIC.

**Fig. 5** Statistical analysis of fluorescence intensity for transcriptional fusions of Sm18 genes
within the manganese MntR miniregulon. (A) Area under the curve (AUC) of normalized
relative fluorescence units (RFU), normalized using the RFU of strains carrying the empty
pSEVA-327 plasmid. Nine biological replicates were grown in MM supplemented with 30
µg/mL of chloramphenicol. AUC values of transcriptional fusions Sm18/327-2357_2358,
Sm18/327-*mntH*, and Sm18/327-*mntP* are shown as mean ± 95% CI (bootstrap) for 0 and 10
µM $Fe^{2+}$. Each Fe condition was tested with 0, 0.08, and 8 µM of $Mn^{2+}$. (B) AUC of normalized
RFU for five biological and four technical replicates of Sm18 and Sm18ΔmntR strains
harboring the same fusions grown in MM-Fe0. Differences were assessed using the Kruskal-
Wallis test followed by Dunn's *post hoc* test with BH correction. Significance levels are shown
as: ns, $p > 0.05$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ***, $p < 0.0001$. Black asterisks
denote comparisons across Mn concentrations, red asterisks denote Fe comparisons, and
gray asterisks indicate combined Fe/Mn variations.

**Fig. 6** Structural, phylogenomic and comparative analysis of the novel CQNC family of

TBDRs and associated periplasmic thioredoxin-fold protein (pTFP). (A) Domain organization

of V8P27_002357 (TBDR) showing the β-barrel and plug domains, the N-terminal signal

peptide and the location of the CQNC motif. (B) Maximum-likelihood phylogeny showing the

monophyly and restricted taxonomic distribution of the CQNC-TBDR. (C) ESPript alignment

showing the conserved CQNC motif located in the center of the plug domain. (D)

Comparative genomics of the TBDR-pTFP showing conserved synteny of TBDR-pTFP loci

across Lysobacterales, Sphingomonadales, and Opitutales. Gray shading indicates sequence

identity. Loci are ordered according to a neighbor-joining tree based on Mash distances. E)

Structure-guided multiple-sequence alignment and a close-up of the CXXC motif of the pTFP

(V8P27_002358), with logo and profile (in PROSITE format) of the presumed catalytic

cysteines (C39, C42) and neighboring residues.

**Fig. 7.** Phenotypic characterization of *S. maltophilia* Sm18 *mntH* and *mntP* mutants. (A-B)

Growth kinetics of the Sm18VIM-*mntH* mutant in MM-MES under oxidative stress induced by

hydrogen peroxide ($H_2O_2$) at pH 7.0 and 6.0. The *mntH* mutant (orange triangles) exhibited

increased sensitivity to hydrogen peroxide (0.2 and 0.5 mM), particularly at neutral pH.

Providing pSEVA332::*mntH* in *trans* restores growth to wild-type levels. (C-D) Growth kinetics

of the markerless Sm18Δ*mntP* mutant (orange triangles) exposed to increasing $Mn^{2+}$

concentrations (5-160 µM). Deletion of *mntP* results in significant manganese sensitivity

starting at 40 µM, with complete growth inhibition at higher concentrations. Plasmid

pSEVA332::*mntP* expressing *mntP* under its native promoters restored the wild type

phenotype (green cubes). Error bars represent SEM for growth kinetics and 95% CI

(bootstrap) for AUC analysis for three biological replicates. Statistical significance was

evaluated using the Kruskal-Wallis test with Dunn's *post hoc* BH correction: ns, $p > 0.05$; *, *p*

$< 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$. Blue asterisks denote comparisons

between wild type (blue circles) and mutants; orange asterisks indicate comparisons across

$H_2O_2$ or $Mn^{2+}$ concentrations.

**Fig. 8.** Live-cell imaging of Sm18/337-*mntH* promoter activity and intracellular replication of

the Sm18VIM-*mntH* mutant in co-culture with *A. castellanii* trophozoites. (A) Representative

images from six biological replicates of Sm18/337R-*mntH* and *A. castellanii* co-cultures at 3

and 18 h post primary contact. Promoter activity was visualized in the red channel

(mCherry2). Trophozoites were imaged using brightfield microscopy. (B) Fluorescence

kinetics of mScarlet-I-labeled strains: Sm18 (blue triangles), Sm18VIM*mntH* (orange cubes),

and Sm18Δ*mntR* (green lines). Background fluorescence from medium-bacteria controls was
subtracted for normalization. (C) Kruskal-Wallis test was used to compare medians between
groups. When significant differences were detected, pairwise comparisons were performed
using Dunn's test with a Bonferroni correction. Significance levels: ns, $p > 0.05$; *, $p < 0.05$; **,
$p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

**Fig. 9.** Model of *S. maltophilia* manganese homeostasis and its role during the interaction with
*A. castellanii* trophozoites. (A) Schematic representation of manganese transport and gene
regulation in strain Sm18 under defined *in vitro* conditions with varying $Mn^{2+}$ and $Fe^{2+}$
concentrations. MntH and MntP mediate manganese uptake and efflux, respectively. (B)
Genes associated with manganese homeostasis that are specifically induced under combined
Mn and Fe limitation. Gene expression is regulated by MntR, with evidence from this study
supporting the existence of an additional MntR-independent regulatory mechanism. (C) Model
of Sm18 interaction with *A. castellanii*. Within the phagosomal environment, manganese
uptake via MntH contributes to resistance against oxidative stress generated by host-derived
$H_2O_2$. Disruption of *mntH* results in reduced intracellular replication. TBDR is shown with
reduced opacity to indicate lack of detectable expression under these conditions. Created
with https://BioRender.com.

**Evaluated contrasts**

| group1 | group2 | p.adj | p.adj.signif |
|--------|--------|-------|--------------|
| 0 | 8 | 1.101202e−02 | * |
| 0 | 16 | 1.470991e−04 | *** |
| 0 | 32 | 4.379815e−07 | **** |
| 4 | 16 | 1.101202e−02 | * |
| 4 | 32 | 1.470991e−04 | *** |
| 8 | 32 | 1.101202e−02 | * |

| group1 | group2 | p.adj | p.adj.signif |
|--------|--------|-------|--------------|
| 0 | 8 | 1.293498e−03 | ** |
| 0 | 16 | 1.864407e−06 | **** |
| 4 | 16 | 1.293498e−03 | ** |

pwc: **Dunn test**; p.adjust: **Benjamini-Hochberg**

**A** Kruskal-Wallis, $X^2 = 24.9$, $p = 1.5e^{-4}$, $n = 54$   $X^2 = 47.1$, $p = 5.4e^{-9}$, $n = 54$   $X^2 = 36.9$, $p = 6.2e^{-7}$, $n = 54$

Sm18/327−2357_2358 · Sm18/327−mntH · Sm18/327−mntP

**B** $X^2 = 29.9$, $p = 1.5e^{-6}$, $n = 36$   $X^2 = 23.8$, $p = 2.8e^{-5}$, $n = 36$   $X^2 = 21.6$, $p = 8.0e^{-5}$, $n = 36$

AUC of normalized RFU (y-axis)

Mn concentration (μM) (x-axis)

Fe concentration (μM): 0, 10

Genetic background: Sm18, Sm18ΔmnR

pwc: **Dunn test**; p.adjust: **Benjamini-Hochberg**

**A** Extracellular space

8µM Mn²⁺ 10µM Fe²⁺

Outer membrane

Periplasm

MntR/+ ⊣ MntH    MntR → MntP

Cytoplasmic membrane

Cytoplasm

**B** 0 µM Mn²⁺ 0 µM Fe²⁺

TBDR

MntR/+ → pTFP

MntR/+ → MntH    MntR ⊣ MntP

**C**

Amoeba cytoplasm

Fagosomal membrane — Nramp1

ADP + Pi    ATP    NADP⁺ + H⁺    NADPH

NOX

Fagosomal Lumen

2H⁺    O₂⁻    2O₂⁻

H₂O₂    O₂

Bacterial outer membrane — TBDR

Bacterial periplasm

Bacterial cytoplasmic membrane — MntH

Mn-SOD    Mn-CAT    **Reduction of oxidative stress**

O₂⁻    H₂O₂    O₂

Bacterial cytoplasm

Interaction Sm18 - *A. castellanii*

Interaction Sm18ΔmntH - *A. castellanii*

MntH ✗ → **Increase of oxidative stress** → **decrease of intracellular replication capacity**

- - - ▶ Mn²⁺ transport direction
⟶ Transcription activation
⊣ Transcription inactivation
+ Unknown MntR-independent regulation

● Mn²⁺