An *OpIE2*-DsRed marker disrupts female blood-feeding and shortens lifespan in the

malaria vector *Anopheles gambiae*

Arad Sarig[1], Evyatar Sar-Shalom[1], Ebrima SM Kolley[1], Elad Shmuel Yonah[1], Lee Benjamin

Lamdan[1], Ariel Roselin Lewin[1], Tamir Partosh[2], Gur Pines[2], Jonathan D. Bohbot[1] and Philippos

Aris Papathanos[1,3]

1. Department of Entomology, Institute of Environment Sciences, Robert H. Smith Faculty of

Agriculture, Food and Environment, Hebrew University of Jerusalem, Rehovot, Israel

2. Institute of Plant Protection, Agricultural Research Organization - Volcani Institute, Rishon

LeZion, Israel

3. Corresponding author: p.papathanos@mail.huji.ac.il

Abstract

Anopheles gambiae is one of the principal vectors of human malaria. Over the past two

decades, transgenic mosquito strains have been essential tools for studying mosquito biology

and developing genetic control strategies such as gene drives. Mosquito transformants are

typically identified using fluorescent markers, which are assumed to be phenotypically neutral.

While generating CRISPR-based gene drive strains carrying an *OpIE2*-DsRed marker we

unexpectedly found that transgenic females were unable to blood-feed and were consequently

sterile, whereas males initially appeared normal and fertile. Given the potential utility of

dominant, female-specific sterility for mosquito control, we established additional strains

controlling for transgene content and integration site, confirming that the *OpIE2*-DsRed cassette

caused the defect. Behavioral assays showed that females exhibited normal attraction to a

membrane feeder but failed to initiate blood-feeding, performing repeated cycles of probing and

proboscis grooming in rapid succession before ultimately leaving the feeder unfed. Microscopy

showed that both sexes possessed a distally curved proboscis, providing a morphological

explanation for the blood-feeding defect of females and the reduced male lifespan. A second

promoter variant (*OpIE2b*), differing in flanking sequences at the IE-2 junction, drove strong

marker expression without impairing blood-feeding or longevity. These findings demonstrate

that minor differences in promoter architecture can produce major, unexpected phenotypic

effects. *OpIE2b* provides a robust, phenotypically neutral marker for *An. gambiae* research,

while *OpIE2a* highlights the need for rigorous validation of transgenic components intended for
research and applied releases.

Introduction

The African malaria mosquito *Anopheles gambiae* is among the most efficient vectors of
Plasmodium parasites and remains a major public-health challenge in sub-Saharan Africa.
Transmission occurs exclusively through the blood-feeding of adult females, which digest blood
to obtain nutrients required for oocyte development and egg production. This link between
blood-feeding and reproduction makes female feeding behavior a critical determinant of both the
vectorial capacity and mosquito population dynamics.

Since stable germline transformation of *Anopheles* mosquitoes was first demonstrated in
the early 2000s (Catteruccia et al. 2000; Grossman et al. 2001), transgenic strains have
become important tools for mosquito research. They now underpin a wide range of studies in
mosquito biology, from malaria transmission biology (Pascini et al. 2022) to olfaction and
sensory behavior (Georgiades et al. 2023; Giraldo et al. 2024; Maguire et al. 2022; Riabinina et
al. 2016) and insecticide resistance (Adolfi et al. 2019). Transgenic approaches are also integral
to the development of genetic control strategies, including gene drives designed for population
suppression and replacement (Carballar-Lejarazú et al. 2020; Haber et al. 2024; Kyrou et al.
2018; Li et al. 2025; Strampelli et al. 2025; Tolosana et al. 2025; Verkuijl et al. 2025).

These advances depend not only on the ability to introduce and stably integrate
transgenes into the mosquito genome, but also on reliably identifying transformed individuals. In
practice, this is achieved through fluorescent reporters driven by 5' and 3' regulatory sequences
(hereafter 5' regulatory sequences are referred to as promoters for simplicity) that ensure strong
and easily detectable expression, most often during larval stages when screening is performed
under a fluorescent microscope. An ideal reporter is bright throughout development, spectrally
distinct from other markers (e.g., eGFP vs DsRed), and compatible with multiple constructs, for
instance through tissue-specific expression patterns (e.g., eye- versus muscle-specific).
Differences in brightness between homozygotes and heterozygotes can sometimes aid
husbandry and genetics, though these are highly dependent on the genome integration site and
are more reliably distinguished using automated optical sorters (e.g., COPAS) than by eye
(Marois 2024). Above all, reporters should be phenotypically neutral, imposing no measurable
fitness costs that might confound experiments or limit their application.

Several promoters have been used to drive fluorescent marker expression in *Anopheles*
gambiae, including the synthetic *3xP3* promoter, which drives strong eye- and neuron-specific
fluorescence (Berghammer, Klingler, and Wimmer 1999) and the heterologous *Actin5C*
promoter from *Drosophila melanogaster*, which provides broad somatic expression (Volohonsky
et al. 2015). Endogenous promoters have also been developed, including the *polyubiquitin*
promoter (Adolfi et al. 2018) and the *α-tubulin-1b* promoter (Lycett, Amenya, and Lynd 2012)
which drive multi-tissue expression. Viral regulatory elements like *hr5-IE1* have also been used
to drive ubiquitous expression (Grossman et al. 2001; Meredith et al. 2013). The *OpIE2*
promoter, derived from the immediate early gene 2 (*IE-2*) of the *Orgyia pseudotsugata*
multicapsid nucleopolyhedrovirus (OpMNPV) (Theilmann and Stewart 1992) has proven highly
effective in diverse insect systems. The *OpIE2* promoter works well in lepidopteran and dipteran
cell line transfections (Pfeifer et al. 1997; Xu et al. 2013) and in transgenic strains of several
insect species, including in *Drosophila melanogaster* (Buchman et al. 2018; Kandul et al. 2019),
Drosophila suzuki (Liu et al. 2025; Witherbee and Gamez 2025), *Bombyx mori* (Li et al. 2015;
Tang et al. 2018), *Pectinophora gossypiella* (Simmons et al. 2011)*, Plutella xylostella* (Martins et
al. 2012)*, Ceratitis Capitata* (Davydova et al. 2023) and *Aedes aegypti* (Estevez-Castro et al.
2024; Li et al. 2017; Rouyar et al. 2024; Vainer et al. 2024) *Aedes albopictus* (Lutrat et al. 2022;
Zaada et al. 2025) and *Culex quinquefasciatus* (Feng et al. 2021). However, its performance in
An. gambiae has not been systematically examined.

Here, we report that expression of DsRed in *An. gambiae* under a specific variant of the
OpIE2 promoter severely impairs female blood-feeding and reduces male longevity. To
investigate this effect, we generated a number of strains controlling for genomic integration site
and construct elements, characterized female feeding behavior, and tested an alternative *OpIE2*
promoter sequence. The alternative promoter maintained marker visibility while restoring normal
feeding, underscoring that promoter choice can have profound effects on mosquito biology and
is critical for both transgenic tool design and genetic control strategies.

Results

Generation of *OpIE2*-DsRed transgenic strains

In previous work, we used an *OpIE2*-DsRed fluorescent marker in *D. melanogaster* strains
containing split Cas9 and gRNA sex ratio distorters (Haber et al. 2024). Seeking to develop
similar capacities in *An. gambiae*, we cloned this *OpIE2*-DsRed to label sgRNA transgenic
constructs, complementing available Cas9 strains marked by fluorescent proteins driven by the

neuron-specific *3xP3* promoter. Four transformation constructs were initially generated, each
containing *OpIE2*-DsRed and single *U6*-driven sgRNA targeting different sex chromosome
sequences (**Figure 1A**). The constructs included *piggyBac* repeats for transposase-mediated
integration and an *attB* site for *phiC31*-mediated site-specific recombination. Constructs were
injected either into wild type eggs with a *piggyBac* transposase helper or into eggs of the *attP*-E
strain, which carries a well-characterized, *3xP3*-CFP-labelled *attP* docking site on chromosome
3 (Meredith et al. 2011). In total, eight independent strains were generated, four with random
piggyBac insertion (construct #1) and three in the *attP*-E line (constructs #2-4; **Supplementary**
Table 1).

Larvae of all strains displayed bright DsRed fluorescence in thoracic and abdominal
segments, and weaker expression in the larval head (**Figure 1B**). In pupae, DsRed
fluorescence was bright throughout the entire body (**Figure 1B**). However, while establishing
the strains, we observed that adult transgenic females from all eight lines failed to blood-feed
(**Figure 1A**), and oviposition bowls provided 48 hours later contained no eggs - a phenotype
described in detail below. In contrast, transgenic males appeared physiologically normal and
fertile, producing offspring when crossed to wild-type females and enabling maintenance of the
strains as heterozygotes. Because the defect was consistent across both *piggyBac* and *phiC31*
integrations, we concluded that it likely resulted from the *OpIE2*-DsRed cassette itself rather
than the insertion site or sgRNA sequence, which differed between constructs.

To test this, we generated a new transformation construct in which DsRed was driven by
the *Actin5C* promoter instead of *OpIE2*. When integrated at the same *attP-E* site, this construct
restored female blood-feeding (construct #5; **Supplementary Table 1; Figure 1A**), suggesting
that the phenotype was specific to *OpIE2*-DsRed expression. We confirmed this by generating a
new construct containing only the *OpIE2*-DsRed cassette and integrating it into the *attP*-E site
(construct #6; **Supplementary Table 1; Figure 1A**). As this strain exhibited the same non-
blood-feeding phenotype, we concluded that *OpIE2*-DsRed expression alone was sufficient to
interrupt female blood-feeding.

When we analyzed the cloned *OpIE2* promoter sequence, we noticed that the promoter
derived from the *Drosophila* constructs differed from another *OpIE2* promoter that we recently
used in *Aedes albopictus* (Zaada *et al.*, 2025). To better understand these differences we
aligned both promoters, which we called *OpIE2a* (for *D. melanogaster* constructs) and *OpIE2b*
(for *A. albopictus* constructs) to the *OpMNPV IE-2* gene region. Both promoters shared a 461 bp
core sequence, but there were several differences in the cloned flanking sequences. (**Figure**

1C). The *OpIE2a* promoter extended 66 bp into the viral *IE-2* coding sequence, including the

start codon and fusing 22 amino acids of *IE-2* to the N-terminus of DsRed. The *OpIE2b*

promoter included an additional 90 bp upstream sequence, but terminated prior to the *IE-2* start

codon in its 3' end, where it connects to DsRed. The 3' end of the *OpIE2b* promoter also

contained *T7* and *EM7* expression elements and restriction sites retained from previous cloning

steps.

To test whether these sequence differences accounted for the phenotype, we generated

a new strain carrying the *OpIE2b*-DsRed cassette integrated into the same *attP-E* site (construct

#7; **Supplementary Table 1; Figure 1A**). DsRed expression driven by *OpIE2b* was slightly

weaker in larvae, pupae and adults compared to *OpIE2a* but followed a similar spatial pattern

(**Figure 1C**). Importantly, *OpIE2b*-DsRed females blood-fed normally (**Figure 1A**),

demonstrating that *OpIE2a*-DsRed caused the feeding defect.

Blood-feeding capacity is impaired in *OpIE2a-DsRed* females.

To characterize and quantify the blood-feeding defect, we established triplicate cages of 14-20

OpIE2a-DsRed or *OpIE2b*-DsRed females, previously mated to wild type males, alongside wild

type control females. Females were offered a membrane feeder containing bovine blood for 25

minutes, and a camera inside each cage recorded female behavior (**Figure 2A**). At the end of

the assay, all females were aspirated from their cages to record the number of blood-fed

females. *OpIE2b*-DsRed females fed normally, with feeding rates indistinguishable from wild

type females ($p = 0.254$, unpaired two-sided *t*-test; **Figure 2B**). In contrast, none of the *OpIE2a*-

DsRed females fed during the 25-minute window ($p = 0.0002$, unpaired two-sided *t*-test; **Figure**

2B). To test whether additional stimulation could elicit feeding, *OpIE2a*-DsRed females were

given a 15-minute rest and re-offered fresh blood for an additional 30 minutes, accompanied by

intermittent human breath stimulation to mimic host-associated cues. Eventually a small fraction

of *OpIE2a*-DsRed females managed to feed (median = 6.25%; four females across all

replicates; **Figure 2B**).

Recordings of female blood-feeding behavior were analyzed using a fine-tuned

YOLOv11l-based object detection model (Khanam and Hussain, 2024) adapted for mosquito

detection by Vainer et al. (2024). *OpIE2a*-DsRed females showed a significantly steeper decline

in feeder visits over time compared to wild-type controls (interaction $\beta = -0.158 \pm 0.038$ SE, $p <$

0.001), whereas *OpIE2b*-DsRed females did not differ significantly ($\beta = -0.047 \pm 0.038$ SE, $p =$

0.21). At the beginning of the assay (time = 0), *OpIE2a*-DsRed females already exhibited fewer

visits than wild type (–6.66 ± 2.23 SE, p = 0.0028), while *OpIE2b*-DsRed females were

comparable to controls (p = 0.08; **Figure 2C**).

Behavioral ethograms for the first five minutes of each assay were also generated,

annotating arrival, probing, blood-feeding and departure events for each female. Wild-type and

OpIE2b-DsRed females began probing shortly after arrival and remained stationary once

feeding commenced, spending significantly more time on the feeder overall (**Figure 2C;**

Supplementary Figure 1). In contrast, *OpIE2a*-DsRed females displayed normal attraction to

the feeder, probing the membrane at a similar frequency to wild-type females, but all failed to

initiate blood-feeding. Instead, they performed repeated cycles of unsuccessful probing followed

by proboscis grooming, ultimately leaving the feeder unfed after brief visits (**Figure 2C;**

Supplementary Figure 1). These observations indicate that the *OpIE2a*-DsRed defect impairs

feeding initiation rather than attraction.

***OpIE2a*-DsRed adults exhibit a curved proboscis**

To investigate whether the behavioral defect corresponded to morphological abnormalities, we

examined the anterior feeding and sensory appendages of adult mosquitoes by

stereofluorescence microscopy. All *OpIE2a*-DsRed females displayed a conspicuous distal

curvature of the proboscis and maxillary palps, in contrast to the straight, parallel alignment

characteristic of wild-type and *OpIE2b*-DsRed females (**Figure 3A-C; Supplementary Figure**

2). The curvature averaged 21.3° along the distal end of the proboscis forming an abrupt ventral

deflection. The paired maxillary palps also appeared bent at the same position as the proboscis.

Because the feeding defect initially appeared female-specific, we also examined males

as controls (**Figure 3D-F)**. Unexpectedly, *OpIE2a*-DsRed males exhibited the same deformity,

with curvature of both the proboscis and palps even more pronounced than in females (mean of

39.2°). The male proboscis often appeared twisted or bent laterally at the tip, and in severe

cases the palps crossed over one another, partially covering the proboscis. In contrast, *OpIE2b*-

DsRed males retained normal morphology, with straight, evenly aligned proboscis and palps

indistinguishable from wild type.

Fluorescence imaging revealed notable DsRed signals within the proboscis and

maxillary palps of *OpIE2a*-DsRed adults, but not in *OpIE2b*-DsRed mosquitoes imaged under

identical exposure settings. The co-occurrence of proboscis-localized DsRed expression and

curvature in *OpIE2a*-DsRed individuals suggests that promoter sequences influence expression

in anterior appendages, coinciding with the observed morphological abnormalities.

OpIE2a-DsRed males exhibit reduced adult longevity

To determine whether the *OpIE2a*-DsRed construct was associated with broader fitness costs,

we assessed reproductive output and survival in both sexes. Among the few *OpIE2a*-DsRed

females that successfully blood-fed after stimulation (n = 4), only two laid eggs. The number of

eggs produced by these females was markedly lower than in controls, with a median of 28 eggs

compared to 97.5 in wild-type and 74.5 in *OpIE2b*-DsRed females (unpaired two-sided *t*-test, *p*

= 0.042; **Figure 4A**). In contrast, egg hatching rates did not differ significantly among strains

(**Figure 4B**), indicating that fertilization and embryonic development were unaffected. When the

blood-feeding assay was repeated with daughters of these females, none successfully blood-

fed, confirming that the phenotype was heritable and fully penetrant (**Supplementary Table 2**).

To evaluate potential effects on male fitness, we crossed transgenic males from each

strain to wild-type females, with wild-type intercrosses as controls. Egg hatching, larval-to-pupal,

and pupal-to-adult survival rates were comparable across all groups (unpaired two-sided *t*-tests,

all *p* > 0.05; **Figure 4C–E**), and adult sex ratios did not differ significantly (**Figure 4F**). However,

adult longevity was severely reduced in *OpIE2a*-DsRed males, which had a median lifespan of

only 6 days compared to 30 days in wild-type males (log-rank test, *p* < 0.001; **Figure 4G**).

OpIE2b-DsRed males showed no difference in survival relative to controls (median 25 days vs

30 days, *p* = 0.355). Among females, *OpIE2a*-DsRed also exhibited a modest reduction in

survival (median 20 days vs 25 days in wild type, *p* = 0.035), whereas *OpIE2b*-DsRed females

survived comparably to controls (median 28 days vs 25 days, *p* = 0.165, **Figure 4H**). Together,

these results overturned our initial conclusion that the *OpIE2a*-DsRed marker negatively

affected fitness in a female-specific manner, by being detrimental to adult longevity, particularly

among males, whereas the *OpIE2b*-DsRed variant maintains normal fecundity, survival, and

developmental progression.

Discussion

In this study, we demonstrate that expression of the fluorescent marker DsRed under the control

of a specific variant of the *OpIE2* promoter (*OpIE2a*) causes profound and unexpected

phenotypic effects in *An. gambiae*. Females carrying the *OpIE2a*-DsRed marker were unable to

blood-feed and were therefore sterile, while males exhibited markedly reduced adult lifespan. In

contrast, an alternative promoter variant, which we called *OpIE2b*, drove robust marker

expression without measurable effects on feeding, development or longevity. These results

reveal that even minor differences in promoter architecture can produce major biological

consequences, emphasizing the need to validate commonly used regulatory elements when
deployed in new species or experimental contexts.

The female blood-feeding impairment associated with *OpIE2a* was fully penetrant and
heritable, persisting across generations - some lines are now over 30 generations old.
Behavioral analyses showed that females were normally attracted to the feeder and initiated
probing, but were unable to transition to sustained feeding. Microscopic examination identified a
fully penetrant curvature of the proboscis and maxillary palps in *OpIE2a* adults of both sexes,
suggesting a structural rather than sensory or motivational basis for the defect. This distal bend
likely prevents proper fascicle unsheathing and membrane penetration of the proboscis (Jones
1978; Kong and Wu 2010), explaining the repeated probing-grooming cycles and failure to feed.
In males, the same curvature may interfere with sugar uptake, consistent with their severely
shortened lifespan.

The difference between the *OpIE2a* and *OpIE2b* promoters lies in a 66-bp extension of
OpIE2a into the viral *IE-2* coding region, which adds 22 amino acids fused to the N-terminus of
DsRed. Although both variants contain the core regulatory motifs required for promoter activity
(Theilmann and Stewart 1992), this fusion likely alters the transcriptional or translation context,
potentially modifying tissue specificity or protein localization. The presence of DsRed
fluorescence within the proboscis and maxillary palps of *OpIE2a* mosquitoes supports this
hypothesis and suggests that local expression or toxicity in mouthparts underlies the
morphological phenotype. While we did not directly measure promoter activity in dissected
tissues or test other reporters under the same sequence, the data clearly demonstrate that even
seemingly small differences in cloned promoter sequences can markedly change expression
domains in transgenic insects, as we have previously observed a native germline promoter in
transgenic *An. gambiae* strains (Papathanos et al. 2009).

Beyond its mechanistic implications, this finding carries practical significance for
transgenic mosquito research and genetic control tool development. First, transgenic reporters
are typically assumed to be phenotypically neutral (Catteruccia, Godfray, and Crisanti 2003;
McArthur, Meredith, and Eggleston 2014); however as we have shown here, this assumption
needs to be confirmed independently each time, since even standard markers can impose
unexpected fitness costs when newly introduced into an engineered insect (Irvin et al. 2004;
Marrelli et al. 2006).

In the case of *OpIE2a*-DsRed, the apparent female-specific sterility initially seemed
attractive for engineering dominant female lethality, for example in self-limiting or split gene-

drive systems (Champer et al. 2022; Verkuijl et al. 2025). However, the severe reduction in
male longevity makes this configuration unsuitable for practical applications of population-
suppression gene drives in its current form. If expression could be restricted to females, for
example through the use of sex-specific splicing (Fu et al. 2010; Liu, Rayes, and Akbari 2024;
Scali et al. 2005; Spinner et al. 2022; Weng et al. 2023, 2024), the dominant appendages
phenotype could be repurposed as a female-specific lethal trait in future designs.

Finally, the identification of *OpIE2b*-DsRed as a functionally neutral marker with bright,
broad expression provides a valuable addition to the transgenesis toolkit available for *An.*
gambiae. Because its expression pattern is distinct from commonly used markers such as *3xP3*
or *Actin5C*, *OpIE2b* is well suited for multiplexed or combinatorial transgenic designs, including
split gene drives and multi-color labeling systems.

In conclusion, our work reveals that promoter sequences alone can profoundly influence
expression and phenotype in transgenic insects, transforming an otherwise inert fluorescent
reporter into a dominant morphological impairment of mosquito physiology. This work offers
both a cautionary example for construct design and a validated promoter variant (*OpIE2b*)
suitable for future *An. gambiae* genetic engineering and gene-drive applications.

Materials and methods:
Mosquito rearing
An. gambiae mosquitoes from the G3 (wild-type) strain, and all transgenic mosquito lines were
reared under standard conditions of 28°C +/- 2°C and 80% +/- 10% relative humidity. Larvae
were fed Essence fish food (Alltech Coppens), while adult mosquitoes were provided with a
10% (wt/vol) sucrose solution. For egg production, adult mosquitoes were allowed to mate for 5-
7 days and then fed bovine blood using an artificial feeder (Hemotek, Ltd). An overnight egg-
bowl was placed in the mosquito cages 48 hours post-blood-feeding for oviposition. Eggs were
allowed 48 hours to hatch, and first instar larvae (L1) were transferred from the egg-bowl to
plastic trays, each containing approximately 250 larvae. The larvae were provided with ground
fish food daily until they reached the pupal stage.

Plasmid construction
The *OpIE2*-DsRed transformation constructs were assembled using Golden-Gate assembly.
Both plasmids contained an *OpIE2* promoter driving DsRed expression, differing only in the
promoter variant used. The *OpIE2a* promoter fragment (541 bp) and the *OpIE2b* promoter

fragment (652 bp) were PCR-amplified with primers incorporating BsaI recognition sites and
compatible overhangs for directional assembly. Each promoter fragment was assembled
upstream of the DsRed coding sequence to generate an *OpIE2*-DsRed cassette. In both
constructs, the cassette was flanked by *piggyBac* inverted terminal repeats and a single *attB*
site, enabling either random *piggyBac*-mediated or *phiC31* site-specific integration. Final
plasmids were validated by Oxford Nanopore sequencing (Plasmidsaurus) to confirm promoter
sequence identity and integrity of the assembled cassette.

Embryo microinjections
Microinjections were performed into eggs of the *attP*-E strain, a homozygous transgenic line
carrying an *attP* docking site on chromosome 3 and a *3xP3*-CFP fluorescence marker. *An.*
gambiae attP-E eggs were gently aligned on a glass slide against a strip of thin nitrocellulose
membrane, with moisture maintained using UltraPure™ Distilled Water (Invitrogen) applied to
dampened filter paper. Using a fine brush, eggs were oriented with the posterior end facing
upwards to maximize the chances of germline integration. A mixture of 300 ng/µl of the
transformation construct and 100 ng/µl of a helper construct encoding vasa-driven *phiC31*
integrase was injected. Hatched larvae were screened for transient expression of the *OpIE2*-
DsRed marker, and all transient-positive and transgenic individuals were backcrossed to G3
wild-type mosquitoes. Progeny were maintained under standard rearing conditions.

Blood-feeding assay and recording of female behavior
Adult males and females were allowed to mate for 5–7 days prior to blood-feeding. For each
strain, three biological replicates of 14–20 females were transferred to Bugdorm minicages
(Model BD4S1515) for feeding. Females were offered blood for 25 min using the Hemotek Ltd
membrane feeding system, with reservoirs filled with thawed bovine blood and sealed with
stretched Parafilm. Feeding sessions were recorded using an Arducam camera system with
underlighting to document female behavior for subsequent analysis. Reservoirs were replaced
with freshly thawed blood and new Parafilm membranes between assays performed on different
strains. If no feeding was observed, a 15 min rest period was followed by a stimulation feeding
session of 25 min, using the same Hemotek reservoir. During stimulation, air was exhaled
intermittently onto the reservoir surface to increase female attraction. At the end of each assay,
females were anesthetized at 4°C, and those with visibly blood-filled abdomens were aspirated
and counted as successfully fed. Blood-feeding recordings were analyzed using a trained

behavioral detection and classification neural network built on the YOLOv11 object detector
(Khanam and Hussain 2024), which was fine-tuned for mosquito detection and adapted from the
behavioral analysis pipeline developed by Vainer et al. (2024). The pre-trained YOLOv11l model
was tuned using a curated dataset of annotated images. Detection parameters were set to a
confidence threshold of 0.2 and an intersection-over-union (IoU) threshold of 0.5 for bounding-
box predictions. Feeder occupancy was analyzed using a linear mixed model with the different
strains and time as fixed effects and replicate as a random intercept.

Blood-feeding videos recorded during the assay were additionally analyzed using the
open-source event-logging software BORIS - Behavioral Observation Research Interactive
Software (Friard and Gamba 2016). For each strain, one video with the highest number of
females was selected for analysis (G3: 20 females, replicate R1; OpIE2b: 20 females, replicate
R3; OpIE2a: 20 females, replicate R1; see **Supplementary Table 3**). Female behavior was
tracked for the first 5 min following placement of the Hemotek feeder in the cage (time = 0).
Because individual females could not be distinguished, behaviors were scored per visit to the
feeder area. A visit was defined as entry into the area of interest over the Hemotek feeder, with
multiple visits by the same female possible. For each visit, arrival and departure times, the
number of probing events and, where applicable, the start and end times of blood-feeding were
recorded.

Female fertility assays

Females that blood-fed during the assay were individually isolated for oviposition. The number
of females per strain was determined by the number of *OpIE2a*-DsRed females that
successfully fed after stimulation. Wild-type and *OpIE2b*-DsRed females were randomly
selected to match this number. 48 h post blood-feeding, an egg-bowl was placed in each cage
and removed the following day. Egg papers were photographed for counting using
Clickmaster2000 (https://github.com/EuracBiomedicalResearch/clickmaster2000). Larvae were
provided food 24 h after egg collection, and the number of hatched larvae was recorded 24 h
later to calculate hatching rates per female. Progeny from these assays were subsequently
reared to adulthood and tested for blood-feeding behavior as described above (filming not
performed due to technical limitations).

Fitness and survival assays

To evaluate potential fitness costs of the *OpIE2*-DsRed cassette, paternal crosses were
established between transgenic males and wild-type females, with wild-type intercrosses as
controls. Eggs were collected and counted to determine hatching rates. Hatchlings were
distributed into trays at a density of 75 larvae per tray and reared under standard conditions. At
pupation, male and female pupae were collected and counted. Pupae were allowed to emerge
in Bugdorm minicages (Model BD4S1515), and the number of adults was determined by
counting remaining pupal cases. These data were used to calculate larval-to-pupal and pupal-
to-adult survival rates for each sex and strain, as well as the sex ratio of emerging adults.
Adult longevity was assessed in Bugdorm minicages (Model BD4S1515) containing 30–35
adults (24–72 h post-emergence) per strain, maintained on 10% sucrose solution. Dead adults
were recorded and removed daily until all individuals had died. If sex or age at death could not
be determined, the individual was censored on the day of discovery. Sucrose solution was
replaced with a new, clean feeder every 14 days. Survival data were analyzed using Kaplan–
Meier survival curves with log-rank tests.
374

Mosquito imaging
Photos of larvae, pupae, and adults were captured under standardized conditions to ensure
consistent exposure across strains. Larvae at the fourth instar and pupae were imaged using an
Olympus SZX16 binocular equipped with a U-HGLGPS fluorescence illumination system. All
fluorescence images were acquired using identical exposure settings for each strain. Adult
mosquitoes were imaged using a Nikon SMZ25 stereo microscope. 2 to 3 day old adults were
first anesthetized with CO_2, after which legs and wings were removed to expose the body and
proboscis for imaging. Dissected adults were then transferred to 50 mL plastic cups sealed with
Parafilm containing silica gel desiccant to prevent condensation and placed at –20 °C for 30 min
to achieve full immobilization. Following freezing, cups were thawed briefly at room temperature
immediately before imaging. Fluorescence parameters and exposure settings were kept
identical between strains to allow direct comparison of DsRed signal patterns.
387

Statistical analysis
Statistical tests were performed in Python using standard packages (3.10.9; scipy 1.15.3,
statsmodels 0.14.4) and JMP Pro 18.0.2 software. All experiments were independently
replicated a minimum of three times unless stated otherwise, and each replicate represented an

independent biological group (e.g., cage or clutch). The number of individuals or replicates used

for each analysis is indicated in the figure panels and methods sections.

 Figure Legends

 Figure 1. Transgenic constructs and marker expression in *OpIE2a* and *OpIE2b* promoters

 driving DsRed. (A) Transgenic construct design carrying either *OpIE2a*, *OpIE2b*, or *Actin5C*

 promoters driving DsRed, with or without a *U6*-sgRNA cassette. All were flanked by *piggyBac*

 inverted repeats and *attB* recombination sites enabling *phiC31*-mediated site-specific

 integration. The genomic location and female blood-feeding phenotype are summarized on the

 right. **(B)** Brightfield and fluorescence of fourth-instar larvae and pupae expressing DsRed under

 the *OpIE2a* promoter. **(C)** Alignment of the *OpIE2a* and *OpIE2b* promoter sequences relative to

 the *Orgyia pseudotsugata* immediate early-2 (*IE-2*) gene. Both promoters share a common 461

 bp core region, but *OpIE2a* extends 66 bp into the *IE-2* coding sequence, while *OpIE2b* includes

 an additional 90 bp upstream region and terminates before the start codon. **(D)** Brightfield and

 fluorescence of fourth-instar larvae and pupae expressing DsRed under the *OpIE2b* promoter.

 Figure 2. Blood-feeding behavior of *OpIE2*-DsRed females. **(A)** Illustration of the

 experimental setup for the blood-feeding assay. Females were placed in minicages containing a

 camera and light source. A membrane-feeder filled with bovine blood was placed on top of the

 cage at the beginning of the experiment. **(B)** Feeding success of wild type, *OpIE2b* and *OpIE2a*

 females. *OpIE2b*-DsRed females fed normally, whereas *OpIE2a*-DsRed females failed to feed.

 Re-exposure of *OpIE2a*-DsRed females to a new feeder following 15min of resting with human-

 breath stimulation induced only occasional feeding. Boxplots show medians ± IQR; points

 represent independent replicate cages. N indicates the total number of females tested per

 treatment. Statistical significance was derived from unpaired two-sided *t*-tests, assuming equal

 or unequal variances, comparing the feeding rates between the different strains to the wild-type

 control ($\alpha = 0.05$). **(C)** Feeder occupancy over time quantified by a YOLO-based behavioral

 tracking network. Solid lines show Gaussian-smoothed mean visits per minute ± SEM across

 (shaded ribbon) replicate cages. *OpIE2a*-DsRed females spent significantly less time on the

 membrane-feeder and their visits declined rapidly compared to WT and *OpIE2b*-DsRed

 controls. **(D)** Manually annotated behavioral ethograms during the first five minutes of the blood-

 feeding assay: Bars represent sequential visits by individual females, color-coded as on-feeder

 (gray), probing (blue), and blood-feeding (red). *OpIE2a*-DsRed females displayed repeated

 short probing attempts without transition to sustained feeding.

Figure 3. Proboscis morphology and DsRed fluorescence in *Anopheles gambiae* adults
carrying *OpIE2*-driven markers. (A–C) Females and **(D–F)** males from wild-type (WT),
OpIE2b-DsRed, and *OpIE2a*-DsRed strains imaged under bright-field (left) and fluorescence
(middle, grayscale; right, red channel) illumination. **(A,D)** WT adults display a straight proboscis
with no fluorescence. **(B,E)** *OpIE2b*-DsRed adults show strong, widespread DsRed expression
but retain normal proboscis morphology. **(C,F)** *OpIE2a*-DsRed adults exhibit a pronounced
distal curvature of the proboscis and strong DsRed signal localized on the proboscis and
maxillary palps. Images were acquired under identical exposure settings to permit direct
comparison of fluorescence intensity and anatomical deformation across strains and sexes.

Figure 4. Reproductive output and fitness of *OpIE2*-DsRed strains. (A–B) Reproductive
performance of females that successfully blood-fed. **(A)** Number of eggs laid per female and **(B)**
corresponding hatching rate (% L1 per egg batch). Each point represents an individual female;
Boxplots show medians ± IQR; N indicates the total number of eggs and hatched larvae tested
per female. Statistical significance was derived from unpaired two-sided *t*-tests, assuming equal
or unequal variances, comparing the feeding rates between the different strains to the wild-type
control ($\alpha = 0.05$). *OpIE2a*-DsRed females produced significantly fewer eggs than wild-type
(WT) or *OpIE2b*-DsRed females, while hatching rates did not differ significantly. **(C–F)** Fitness
parameters from crosses of transgenic males to WT females. **(C)** Hatching rate (% L1 per egg
batch), **(D)** larval-to-pupal survival (%) and **(E)** pupal-to-adult survival (%) were comparable
across strains. Boxplots show medians ± IQR; points represent replicate cages. **(F)** Adult sex
ratios of progeny from the same crosses. Stacked bars show the mean proportion of males and
females. Statistical comparisons were calculated relative to WT control using unpaired two-
sided *t*-tests; no significant differences were detected (all *p* > 0.05). **(G–H)** Adult longevity of
males **(G)** and females **(H)**. *OpIE2a*-DsRed males exhibited a markedly reduced lifespan
compared to WT and *OpIE2b*-DsRed males, while females showed a moderate but significant
reduction in survival. Survival curves were analyzed using Kaplan-Meier estimates and
compared using log-rank statistics. Diamonds denote median survival per strain.

Author contributions

AS and PAP conceived and designed the study. AS, ESY, ESMK and LBL generated transgenic mosquito lines. AS performed behavioral, imaging, and fitness assays, and analyzed the data. ARL helped with figure illustrations. ES-S and JB developed the trained behavioral detection and classification neural network. ESMK, ESY, and LBL assisted with mosquito genetic engineering. TP and GP contributed to adult proboscis imaging. PAP supervised the project, secured funding, and wrote the manuscript together with AS with input from all authors. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Daniella An Haber and Yael Arien for their valuable discussions and critical feedback and Gleb Ens, Ruth Shacham and Albert Nazarov for technical support.

Funding

This work was supported by research grants from the Gates Foundation (INV-004363 and INV-075374 to PAP) and the Israel Science Foundation (ISF) (2388/19 to PAP).

Ethics

All animals were handled in accordance and under the supervision of the ARO Institutional Animal Care and Use Committee approval number 2307-118-2-VOL-IL. All insect work was performed in facilities maintaining Arthropod Containment Level 2. This work received Institutional Approval and relevant authorizations from the Israel Ministry of Environmental Protection and Ministry of Agriculture (#31/2019).

Competing interests

The authors declare no competing interests.

Supplementary Figure 1. Quantification of *OpIE2*-DsRed female feeding behavior. (A)
Time spent on the feeder (min) per female visit. The time that each female spent on the feeder
area differed significantly between strains compared to the WT control (Kruskal–Wallis, *H* =
37.6634, *p* < 0.001). *OpIE2b*-DsRed females spent slightly more time on the feeder compared
to WT females (Dunn's comparisons with Bonferroni correction vs WT, *p* = 0.033), whereas
OpIE2a-DsRed females spent significantly less time (*p* < 0.001). **(B)** Frequency of probing
events per female visit. Group differences were significant (Kruskal–Wallis, *H* = 9.93, *p* = 0.007).
OpIE2b-DsRed females showed a slight reduction in probing frequency compared to wild-type
females (Dunn's comparisons with Bonferroni correction vs WT, *p* = 0.0485), while *OpIE2a*-
DsRed females did not differ significantly (*p* = 0.85). **(C)** Frequency of proboscis grooming
events per female visit. Group differences were significant (Kruskal–Wallis, *H* = 11.24, *p* =
0.0036). Grooming frequency was significantly higher in *OpIE2a*-DsRed females relative to WT
(Dunn's comparisons with Bonferroni correction vs WT, *p* = 0.042), whereas *OpIE2b*-DsRed
females were not different from WT (*p* = 0.66). For WT and   *OpIE2b*-DsRed females, probing
and proboscis grooming events prior to the initiation of blood-feeding are considered, to enable
comparison with *OpIE2a*-DsRed females, which do not initiate feeding.

Supplementary Figure 2. Quantification of proboscis curvature in OpIE2-DsRed
mosquitoes. (A) Mean ventral curvature angle angle (°) of female proboscises. *OpIE2a*-DsRed
females exhibited a significantly greater ventral curvature relative to wild-type females (unpaired
two-sided *t*-test, *p* = 0.007), whereas OpIE2b-DsRed females did not differ significantly from
controls (*p* = 0.63). Each data point represents an individual mosquito. **(B)** Mean ventral
curvature angle (°) of male proboscises. *OpIE2a*-DsRed males exhibited a significantly greater
ventral curvature relative to wild-type males (unpaired two-sided *t*-test, *p* < 0.001. No significant
difference was observed between *OpIE2b*-DsRed and wild-type males (*p* = 0.43).

Supplementary Table 1. Summary of transgenic strains.

Strain name	Date created	Construct #	Integration	sgRNA	Larval phenotype	Blood-feeding
AS-1 V.1	Oct-22	1	*pBac*	TCACCATCTGAGCGTTGCAC	Mixed (Strong and weak)	Impaired
AS-1 V.3	Oct-22	1	*pBac*	TCACCATCTGAGCGTTGCAC	Mixed (Strong and weak)	Impaired
AS-1 F.3	Nov-22	1	*pBac*	TCACCATCTGAGCGTTGCAC	Strong	Impaired
AS-1 F.5	Nov-22	1	*pBac*	TCACCATCTGAGCGTTGCAC	Strong	Impaired
AS-1 M.1	Dec-22	1	*pBac*	TCACCATCTGAGCGTTGCAC	Strong	Impaired
AS-5	Mar-23	2	*phiC31*	GGTGGGCGAGAGTTCACGTT	Strong	Impaired
AS-6	Apr-23	3	*phiC31*	GCAGTGAATCCATTGCTGGG	Strong	Impaired
AS-7	Jun-23	4	*phiC31*	GAGCTGGGCAGTTCAATGGG	Strong	Impaired
Xshrd	Jul-23	5	*phiC31*	GGGTACAAGCTTGCGTACGT	Strong	Fine
OpIE2a	Sept-24	6	*phiC31*	-	Strong	Impaired
OpIE2b	Oct-24	7	*phiC31*	-	Strong	Fine

Each strain is identified by an internal strain name. Construct number refers to the numbering
system used to distinguish transformation plasmids. Integration method indicates whether
transformation occurred through random *piggyBac*-mediated transposition or *phiC31*-mediated
site-specific integration. The sgRNA linker sequence shows the 5′-3′ sgRNA space sequence
cloned for each construct containing a *U6*-sgRNA cassette. Larval marker phenotype denotes
the strength and consistency of DsRed fluorescence, while female blood-feeding behavior and
male fertility summarize adult phenotypic outcomes for each line.

Supplementary Table 2. Heritable disruption of blood-feeding by *Opie2a*-DsRed.

Strain	Experiment date	Repeat	N females	N females fed
OpIE2a females	05.01.2024	1	9	0
OpIE2a females	05.01.2024	2	9	0
OpIE2a females	05.01.2024	3	9	0

Daughters of the two *OpIE2a*-DsRed females that successfully fed after additional stimulation (Figure 2C) were tested for blood-feeding using the same experimental setup as their mothers. Three replicate cages were established, each containing nine 5–7-day-old females previously mated to wild-type males. In all replicates, no females were observed with visibly blood-filled abdomens at the end of the feeding period, confirming complete penetrance and transmission of the non-feeding phenotype.

Supplementary File 1. Sequences of the *OpIE2a* and *OpIE2b* promoter variants and constructs generated in this study in FASTA format. File numbering corresponds to the constructs listed in **Supplementary Table 1** and shown in **Figure 1A**.

Supplementary File 2. Manual annotations of behavioral metrics of membrane feeding assays. The table contains output data of BORIS manually recorded behavioral observations for individual *An. gambiae* females of the indicated genotypes. Each row represents a single female visit. Behavioral events were recorded from 5-min video segments and include the number of grooming and probing events, total time spent in the area of interest (AOI) near the feeder, and total duration of blood-feeding. "Time at AOI" represents cumulative occupancy on the feeder, while "ROI before feeding" and associated probing metrics quantify pre-feeding activity. Frequencies are expressed as events per second or minute as indicated.

Supplementary File 3. Statistical analysis summary. All statistical tests, *p*-values, and significance levels corresponding to figures and supplementary figures presented in this study.

References

Adolfi, Adriana, Emilie Pondeville, Amy Lynd, Catherine Bourgouin, and Gareth J. Lycett. 2018. "Multi-Tissue GAL4-Mediated Gene Expression in All Anopheles Gambiae Life Stages Using an Endogenous Polyubiquitin Promoter." *Insect Biochemistry and Molecular Biology* 96:1–9. doi:10.1016/j.ibmb.2018.03.005.

Adolfi, Adriana, Beth Poulton, Amalia Anthousi, Stephanie Macilwee, Hilary Ranson, and Gareth J. Lycett. 2019. "Functional Genetic Validation of Key Genes Conferring Insecticide Resistance in the Major African Malaria Vector, Anopheles Gambiae." *Proceedings of the National Academy of Sciences of the United States of America* 116(51):25764–72. doi:10.1073/pnas.1914633116.

Berghammer, A. J., M. Klingler, and E. A. Wimmer. 1999. "A Universal Marker for Transgenic Insects." *Nature* 402(6760):370–71. doi:10.1038/46463.

Buchman, Anna B., Tobin Ivy, John M. Marshall, Omar S. Akbari, and Bruce A. Hay. 2018. "Engineered Reciprocal Chromosome Translocations Drive High Threshold, Reversible Population Replacement in Drosophila." *ACS Synthetic Biology* 7(5):1359–70. doi:10.1021/acssynbio.7b00451.

Carballar-Lejarazú, Rebeca, Christian Ogaugwu, Taylor Tushar, Adam Kelsey, Thai Binh Pham, Jazmin Murphy, Hanno Schmidt, Yoosook Lee, Gregory C. Lanzaro, and Anthony A. James. 2020. "Next-Generation Gene Drive for Population Modification of the Malaria Vector Mosquito, Anopheles Gambiae." *Proceedings of the National Academy of Sciences* 117(37):22805–14. doi:10.1073/pnas.2010214117.

Catteruccia, Flaminia, H. Charles J. Godfray, and Andrea Crisanti. 2003. "Impact of Genetic Manipulation on the Fitness of Anopheles Stephensi Mosquitoes." *Science* 299(5610):1225–27. doi:10.1126/science.1081453.

Catteruccia, Flaminia, Tony Nolan, Thanasis G. Loukeris, Claudia Blass, Charalambos Savakis, Fotis C. Kafatos, and Andrea Crisanti. 2000. "Stable Germline Transformation of the Malaria Mosquito Anopheles Stephensi." *Nature* 405(6789):959–62. doi:10.1038/35016096.

Champer, Samuel E., Isabel K. Kim, Andrew G. Clark, Philipp W. Messer, and Jackson Champer. 2022. "Anopheles Homing Suppression Drive Candidates Exhibit Unexpected Performance Differences in Simulations with Spatial Structure" edited by G. H. Perry, S. A. Verkuijl, and J. Bull. *eLife* 11:e79121. doi:10.7554/eLife.79121.

Davydova, Serafima, Junru Liu, Nikolay P. Kandul, W. Evan Braswell, Omar S. Akbari, and Angela Meccariello. 2023. "Next-Generation Genetic Sexing Strain Establishment in the Agricultural Pest Ceratitis Capitata." *Scientific Reports* 13(1):19866. doi:10.1038/s41598-023-47276-5.

Estevez-Castro, Carlos F., Murillo F. Rodrigues, Antinéa Babarit, Flávia V. Ferreira, Elisa G. de Andrade, Eric Marois, Rodrigo Cogni, Eric R. G. R. Aguiar, João T. Marques, and Roenick P. Olmo. 2024. "Neofunctionalization Driven by Positive Selection Led to the

Retention of the Loqs2 Gene Encoding an Aedes Specific dsRNA Binding Protein." *BMC Biology* 22:14. doi:10.1186/s12915-024-01821-4.

Feng, Xuechun, Víctor López Del Amo, Enzo Mameli, Megan Lee, Alena L. Bishop, Norbert Perrimon, and Valentino M. Gantz. 2021. "Optimized CRISPR Tools and Site-Directed Transgenesis towards Gene Drive Development in Culex Quinquefasciatus Mosquitoes." *Nature Communications* 12:2960. doi:10.1038/s41467-021-23239-0.

Friard, Olivier, and Marco Gamba. 2016. "BORIS: A Free, Versatile Open-Source Event-Logging Software for Video/Audio Coding and Live Observations." *Methods in Ecology and Evolution* 7. doi:10.1111/2041-210X.12584.

Fu, Guoliang, Rosemary S. Lees, Derric Nimmo, Diane Aw, Li Jin, Pam Gray, Thomas U. Berendonk, Helen White-Cooper, Sarah Scaife, Hoang Kim Phuc, Osvaldo Marinotti, Nijole Jasinskiene, Anthony A. James, and Luke Alphey. 2010. "Female-Specific Flightless Phenotype for Mosquito Control." *Proceedings of the National Academy of Sciences* 107(10):4550–54. doi:10.1073/pnas.1000251107.

Georgiades, Marcos, Alexandros Alampounti, Jason Somers, Matthew P. Su, David A. Ellis, Judit Bagi, Daniela Terrazas-Duque, Scott Tytheridge, Watson Ntabaliba, Sarah Moore, Joerg T. Albert, and Marta Andrés. 2023. "Hearing of Malaria Mosquitoes Is Modulated by a Beta-Adrenergic-like Octopamine Receptor Which Serves as Insecticide Target." *Nature Communications* 14(1):4338. doi:10.1038/s41467-023-40029-y.

Giraldo, Diego, Andrew M. Hammond, Jinling Wu, Brandon Feole, Noor Al-Saloum, and Conor J. McMeniman. 2024. "An Expanded Neurogenetic Toolkit to Decode Olfaction in the African Malaria Mosquito Anopheles Gambiae." *Cell Reports Methods* 4(2):100714. doi:10.1016/j.crmeth.2024.100714.

Grossman, G. L., C. S. Rafferty, J. R. Clayton, T. K. Stevens, O. Mukabayire, and M. Q. Benedict. 2001. "Germline Transformation of the Malaria Vector,Anopheles Gambiae, with the piggyBac Transposable Element." *Insect Molecular Biology* 10(6):597–604. doi:10.1046/j.0962-1075.2001.00299.x.

Haber, Daniella An, Yael Arien, Lee Benjamin Lamdan, Yehonathan Alcalay, Chen Zecharia, Flavia Krsticevic, Elad Shmuel Yonah, Rotem Daniel Avraham, Elzbieta Krzywinska, Jaroslaw Krzywinski, Eric Marois, Nikolai Windbichler, and Philippos Aris Papathanos. 2024. "Targeting Mosquito X-Chromosomes Reveals Complex Transmission Dynamics of Sex Ratio Distorting Gene Drives." *Nature Communications* 15(1):4983. doi:10.1038/s41467-024-49387-7.

Irvin, Nic, Mark S. Hoddle, David A. O'Brochta, Bryan Carey, and Peter W. Atkinson. 2004. "Assessing Fitness Costs for Transgenic Aedes Aegypti Expressing the GFP Marker and Transposase Genes." *Proceedings of the National Academy of Sciences* 101(3):891–96. doi:10.1073/pnas.0305511101.

Jones, Jack Colvard. 1978. "The Feeding Behavior of Mosquitoes." *Scientific American* 238(6):138–50.

Kandul, Nikolay P., Junru Liu, Hector M. Sanchez C., Sean L. Wu, John M. Marshall, and Omar S. Akbari. 2019. "Transforming Insect Population Control with Precision Guided Sterile Males with Demonstration in Flies." *Nature Communications* 10(1):84. doi:10.1038/s41467-018-07964-7.

Khanam, Rahima, and Muhammad Hussain. 2024. "YOLOv11: An Overview of the Key Architectural Enhancements."

Kong, X. Q., and C. W. Wu. 2010. "Mosquito Proboscis: An Elegant Biomicroelectromechanical System." *Physical Review E* 82(1):011910. doi:10.1103/PhysRevE.82.011910.

Kyrou, Kyros, Andrew M. Hammond, Roberto Galizi, Nace Kranjc, Austin Burt, Andrea K. Beaghton, Tony Nolan, and Andrea Crisanti. 2018. "A CRISPR–Cas9 Gene Drive Targeting Doublesex Causes Complete Population Suppression in Caged Anopheles Gambiae Mosquitoes." *Nature Biotechnology* 36(11):1062–66. doi:10.1038/nbt.4245.

Li, Ming, Michelle Bui, Ting Yang, Christian S. Bowman, Bradley J. White, and Omar S. Akbari. 2017. "Germline Cas9 Expression Yields Highly Efficient Genome Engineering in a Major Worldwide Disease Vector, Aedes Aegypti." *Proceedings of the National Academy of Sciences* 114(49):E10540–49. doi:10.1073/pnas.1711538114.

Li, Zhiqian, Yuemei Dong, Lang You, Rodrigo M. Corder, Jemariz Arzobal, Audrey Yeun, Lei Yang, John M. Marshall, George Dimopoulos, and Ethan Bier. 2025. "Driving a Protective Allele of the Mosquito FREP1 Gene to Combat Malaria." *Nature* 1–9. doi:10.1038/s41586-025-09283-6.

Li, Zhiqian, Baosheng Zeng, Lin Ling, Jun Xu, Lang You, Abu F. M. Aslam, Anjiang Tan, and Yongping Huang. 2015. "Enhancement of Larval RNAi Efficiency by Over-Expressing Argonaute2 in Bombyx Mori." *International Journal of Biological Sciences* 11(2):176–85. doi:10.7150/ijbs.10235.

Liu, Junru, Danny Rayes, and Omar S. Akbari. 2024. "A Fluorescent Sex-Sorting Technique for Insects with the Demonstration in Drosophila Melanogaster." *GEN Biotechnology* 3(1):35–44. doi:10.1089/genbio.2023.0041.

Liu, Junru, Danny Rayes, Minzhe Yang, and Omar S. Akbari. 2025. "Fluorescent-Based Sex-Separation Technique in Major Invasive Crop Pest, Drosophila Suzukii." *GEN Biotechnology* 4(1):29–36. doi:10.1089/genbio.2024.0056.

Lutrat, Célia, Roenick P. Olmo, Thierry Baldet, Jérémy Bouyer, and Eric Marois. 2022. "Transgenic Expression of Nix Converts Genetic Females into Males and Allows Automated Sex Sorting in Aedes Albopictus." *Communications Biology* 5(1):210. doi:10.1038/s42003-022-03165-7.

Lycett, Gareth J., D. Amenya, and A. Lynd. 2012. "The Anopheles Gambiae Alpha-Tubulin-1b Promoter Directs Neuronal, Testes and Developing Imaginal Tissue Specific Expression and Is a Sensitive Enhancer Detector." *Insect Molecular Biology* 21(1):79–88. doi:10.1111/j.1365-2583.2011.01112.x.

Maguire, Sarah E., Ali Afify, Loyal A. Goff, and Christopher J. Potter. 2022. "Odorant-Receptor-Mediated Regulation of Chemosensory Gene Expression in the Malaria Mosquito *Anopheles Gambiae.*" *Cell Reports* 38(10):110494. doi:10.1016/j.celrep.2022.110494.

Marois, Eric. 2024. "Sorting Mosquito Larvae with a COPAS Machine." *Cold Spring Harbor Protocols* 2024(6):pdb.prot108307. doi:10.1101/pdb.prot108307.

Marrelli, Mauro T., Cristina K. Moreira, David Kelly, Luke Alphey, and Marcelo Jacobs-Lorena. 2006. "Mosquito Transgenesis: What Is the Fitness Cost?" *Trends in Parasitology* 22(5):197–202. doi:10.1016/j.pt.2006.03.004.

Martins, S., N. Naish, A. S. Walker, N. I. Morrison, S. Scaife, G. Fu, T. Dafa'alla, and L. Alphey. 2012. "Germline Transformation of the Diamondback Moth, Plutella Xylostella L., Using the piggyBac Transposable Element." *Insect Molecular Biology* 21(4):414–21. doi:10.1111/j.1365-2583.2012.01146.x.

McArthur, Clare C., Janet M. Meredith, and Paul Eggleston. 2014. "Transgenic Anopheles Gambiae Expressing an Antimalarial Peptide Suffer No Significant Fitness Cost." *PLOS ONE* 9(2):e88625. doi:10.1371/journal.pone.0088625.

Meredith, Janet M., Sanjay Basu, Derric D. Nimmo, Isabelle Larget-Thiery, Emma L. Warr, Ann Underhill, Clare C. McArthur, Victoria Carter, Hilary Hurd, Catherine Bourgouin, and Paul Eggleston. 2011. "Site-Specific Integration and Expression of an Anti-Malarial Gene in Transgenic Anopheles Gambiae Significantly Reduces Plasmodium Infections." *PLOS ONE* 6(1):e14587. doi:10.1371/journal.pone.0014587.

Meredith, Janet M., Ann Underhill, Clare C. McArthur, and Paul Eggleston. 2013. "Next-Generation Site-Directed Transgenesis in the Malaria Vector Mosquito Anopheles Gambiae: Self-Docking Strains Expressing Germline-Specific phiC31 Integrase." *PLoS ONE* 8(3):e59264. doi:10.1371/journal.pone.0059264.

Papathanos, Philippos A., Nikolai Windbichler, Miriam Menichelli, Austin Burt, and Andrea Crisanti. 2009. "The Vasa Regulatory Region Mediates Germline Expression and Maternal Transmission of Proteins in the Malaria Mosquito Anopheles Gambiae: A Versatile Tool for Genetic Control Strategies." *BMC Molecular Biology* 10(1):65. doi:10.1186/1471-2199-10-65.

Pascini, Tales V., Yeong Je Jeong, Wei Huang, Zarna R. Pala, Juliana M. Sá, Michael B. Wells, Christopher Kizito, Brendan Sweeney, Thiago L. Alves e Silva, Deborah J. Andrew, Marcelo Jacobs--Lorena, and Joel Vega-Rodríguez. 2022. "Transgenic Anopheles Mosquitoes Expressing Human PAI-1 Impair Malaria Transmission." *Nature Communications* 13(1):1–16. doi:10.1038/s41467-022-30606-y.

Pfeifer, Tom A., Dwayne D. Hegedus, Thomas A. Grigliatti, and David A. Theilmann. 1997. "Baculovirus Immediate-Early Promoter-Mediated Expression of the Zeocin™ Resistance Gene for Use as a Dominant Selectable Marker in Dipteran and Lepidopteran Insect Cell Lines." *Gene* 188(2):183–90. doi:10.1016/S0378-1119(96)00756-1.

Riabinina, Olena, Darya Task, Elizabeth Marr, Chun-Chieh Lin, Robert Alford, David A. O'Brochta, and Christopher J. Potter. 2016. "Organization of Olfactory Centres in the Malaria Mosquito Anopheles Gambiae." *Nature Communications* 7(1):13010. doi:10.1038/ncomms13010.

Rouyar, Angela, Anandrao A. Patil, Melissa Leon-Noreña, Ming Li, Iliano V. Coutinho-Abreu, Omar S. Akbari, and Jeff A. Riffell. 2024. "Transgenic Line for Characterizing GABA-Receptor Expression to Study the Neural Basis of Olfaction in the Yellow-Fever Mosquito." *Frontiers in Physiology* 15:1381164. doi:10.3389/fphys.2024.1381164.

Scali, Christina, Flaminia Catteruccia, Qiuxiang Li, and Andrea Crisanti. 2005. "Identification of Sex-Specific Transcripts of the Anopheles Gambiae Doublesex Gene." *Journal of Experimental Biology* 208(19):3701–9. doi:10.1242/jeb.01819.

Simmons, Gregory S., Andrew R. McKemey, Neil I. Morrison, Sinead O'Connell, Bruce E. Tabashnik, John Claus, Guoliang Fu, Guolei Tang, Mickey Sledge, Adam S. Walker, Caroline E. Phillips, Ernie D. Miller, Robert I. Rose, Robert T. Staten, Christl A. Donnelly, and Luke Alphey. 2011. "Field Performance of a Genetically Engineered Strain of Pink Bollworm." *PLoS ONE* 6(9):e24110. doi:10.1371/journal.pone.0024110.

Spinner, Siân A. M., Zoe H. Barnes, Alin Mirel Puinean, Pam Gray, Tarig Dafa'alla, Caroline E. Phillips, Camila Nascimento de Souza, Tamires Fonseca Frazon, Kyla Ercit, Amandine Collado, Neil Naish, Edward Sulston, Gwilym C. Ll. Phillips, Kelleigh K. Greene, Mattia Poletto, Benjamin D. Sperry, Simon A. Warner, Nathan R. Rose, Grey K. Frandsen, Natalia C. Verza, Kevin J. Gorman, and Kelly J. Matzen. 2022. "New Self-Sexing Aedes Aegypti Strain Eliminates Barriers to Scalable and Sustainable Vector Control for Governments and Communities in Dengue-Prone Environments." *Frontiers in Bioengineering and Biotechnology* 10. doi:10.3389/fbioe.2022.975786.

Strampelli, Anna, Katie Willis, Hannah Robyn Gulliford, Matt Gribble, Barbara Fasulo, Austin Burt, Andrea Crisanti, and Federica Bernardini. 2025. "A Male-Drive Female-Sterile System for the Self-Limited Control of the Malaria Mosquito Anopheles Gambiae." 2025.06.10.658817.

Tang, X. F., X. L. Zhou, Q. Zhang, P. Chen, C. Lu, and M. H. Pan. 2018. "Bombyx Mori Cyclin-Dependent Kinase Inhibitor Is Involved in Regulation of the Silkworm Cell Cycle." *Insect Molecular Biology* 27(3):404–13. doi:10.1111/imb.12381.

Theilmann, D. A., and S. Stewart. 1992. "Molecular Analysis of the Trans-Activating IE-2 Gene of Orgyia Pseudotsugata Multicapsid Nuclear Polyhedrosis Virus." *Virology* 187(1):84–96. doi:10.1016/0042-6822(92)90297-3.

Tolosana, Ignacio, Katie Willis, Matthew Gribble, Lee Phillimore, Austin Burt, Tony Nolan, Andrea Crisanti, and Federica Bernardini. 2025. "A Y Chromosome-Linked Genome Editor for Efficient Population Suppression in the Malaria Vector Anopheles Gambiae." *Nature Communications* 16(1):206. doi:10.1038/s41467-024-55391-8.

Vainer, Yuri, Yinliang Wang, Robert M. Huff, Dor Perets, Evyatar Sar-Shalom, Esther Yakir, Majid Ghaninia, Iliano V. Coutinho-Abreu Gomes, Carlos Ruiz, Dhivya Rajamanickam,

A. Warburg, Omar S. Akbari, Philippos A. Papathanos, R. Ignell, Jeffrey A. Riffell, R.
Jason Pitts, and Jonathan D. Bohbot. 2024. "A Conserved Odorant Receptor Underpins
Borneol-Mediated Repellency in Culicine Mosquitoes." *bioRxiv* 2023.08.01.548337.
doi:10.1101/2023.08.01.548337.

Verkuijl, Sebald A. N., Giuseppe Del Corsano, Paolo Capriotti, Pei-Shi Yen, Maria Grazia
Inghilterra, Prashanth Selvaraj, Astrid Hoermann, Aida Martinez-Sanchez, Chiamaka
Valerie Ukegbu, Temesgen M. Kebede, Dina Vlachou, George K. Christophides, and
Nikolai Windbichler. 2025. "A Suppression-Modification Gene Drive for Malaria Control
Targeting the Ultra-Conserved RNA Gene Mir-184." *Nature Communications* 16(1):3923.
doi:10.1038/s41467-025-58954-5.

Volohonsky, Gloria, Olivier Terenzi, Julien Soichot, Daniel A. Naujoks, Tony Nolan, Nikolai
Windbichler, Delphine Kapps, Andrea L. Smidler, Anaïs Vittu, Giulia Costa, Stefanie
Steinert, Elena A. Levashina, Stéphanie A. Blandin, and Eric Marois. 2015. "Tools for
Anopheles Gambiae Transgenesis." *G3: Genes|Genomes|Genetics* 5(6):1151–63.
doi:10.1534/g3.115.016808.

Weng, Shih-Che, Igor Antoshechkin, Eric Marois, and Omar S. Akbari. 2023. "Efficient Sex
Separation by Exploiting Differential Alternative Splicing of a Dominant Marker in Aedes
Aegypti." *PLoS Genetics* 19(11):e1011065. doi:10.1371/journal.pgen.1011065.

Weng, Shih-Che, Fangying Chen, Ming Li, Sammy Lee, Connor Gerry, Dylan Can Turksoy, and
Omar S. Akbari. 2024. "Establishing a Dominant Early Larval Sex-Selection Strain in the
Asian Malaria Vector Anopheles Stephensi." *Infectious Diseases of Poverty* 13(1):83.
doi:10.1186/s40249-024-01256-7.

Witherbee, Avery D., and Stephanie Gamez. 2025. "Dissecting the Sterility Phenotype in Gene
Edited Drosophila Suzukii pgSIT Males." *Scientific Reports* 15:3903.
doi:10.1038/s41598-025-88598-w.

Xu, Jian, Yudai Nagata, Hiroaki Mon, Zhiqing Li, Li Zhu, Kazuhiro Iiyama, Takahiro Kusakabe,
and Jae Man Lee. 2013. "Soaking RNAi-Mediated Modification of Sf9 Cells for
Baculovirus Expression System by Ectopic Expression of Caenorhabditis Elegans SID-
1." *Applied Microbiology and Biotechnology* 97(13):5921–31. doi:10.1007/s00253-013-
4785-1.

Zaada, Doron SY, Or Toren, Flavia Krsticevic, Daniella A. Haber, Denys Gildman, Noam
Galpaz, Irina Häcker, Marc F. Schetelig, Eric Marois, Yael Arien, and Philippos A.
Papathanos. 2025. "Mosquito Sex Separation Using Complementation of Selectable
Traits and Engineered Neo-Sex Chromosomes." 2025.03.30.646185.

Figure 1

A

	Construct	Insertion	Blood-feeding
OpIE2a *attB - pBacL - DsRed - sgRNA - pBacR* (U6)	1	autosomal	No
OpIE2a *attL - pBacL - DsRed - sgRNA - pBacR - attR* (U6)	2-4	attP-E	No
Actin5C *attL - pBacL - DsRed - sgRNA - pBacR - attR* (U6)	5	attP-E	Yes
OpIE2a *attL - pBacL - DsRed - pBacR - attR*	6	attP-E	No
OpIE2b *attL - pBacL - DsRed - pBacR - attR*	7	attP-E	Yes

Transgene

B

OpIE2a-DsRed

DsRed

C

bps 90   461   13 66   *IE-2* ORF

OpIE2a
OpIE2b

D

OpIE2b-DsRed

DsRed

Figure 2

Figure 3

A WT females

B *OpIE2b*-DsRed females

C *OpIE2a*-DsRed females

D WT males

E *OpIE2b*-DsRed males

F *OpIE2a*-DsRed males

Figure 4