Proactive adjustments to cued gait perturbations in people with and without chronic stroke

Tara Cornwell[1] and James M. Finley[1-3]

[1]Department of Biomedical Engineering, University of Southern California, USA
[2]Division of Biokinesiology and Physical Therapy, University of Southern California, USA
[3]Neuroscience Graduate Program, University of Southern California, USA

Correspondence: *Tara Cornwell (tcornwell@u.northwestern.edu)*

Abstract

Balance disturbances exist along a continuum from those that are fully unexpected to those that are predictable based on cues from the environment. When people experience predictable disturbances while walking, they may proactively adjust their gait to minimize losses of balance. However, one's ability to effectively implement these proactive control strategies may be impaired after a stroke due to a combination of motor and cognitive impairments that result from brain lesions. Here, we used explicit audiovisual cues to characterize the proactive and reactive control strategies implemented by people with and without stroke during unexpected versus expected gait perturbations. Following unexpected treadmill accelerations, both groups had smaller margins of stability on the recovery step than during unperturbed walking. When we provided audiovisual cues specifying the impending perturbation step, people without stroke performed less leg and joint work, especially at the ankle, during the cued perturbations and increased their subsequent margins of stability by approximately 3 cm on the recovery step. However, people post-stroke did not make these proactive adjustments. Instead, after any perturbation, they modified their stepping to maintain their center of mass position within their base of support, and this reactive strategy remained unchanged with audiovisual cues. Our findings suggest that people post-stroke rely on a general reactive control strategy rather than proactively modifying push-off work, even when given precise timing information about the impending gait perturbations.

New & Noteworthy

People maintain balance during walking with a combination of proactive and reactive adjustments, but stroke-related impairments may hinder the effectiveness of these control adjustments. We varied audiovisual cues before single-belt treadmill accelerations and found

that people without stroke used the cues to modify their lower-extremity work, thereby improving subsequent stability. In contrast, people post-stroke did not use the cues to update their control strategy, which may contribute to their fall risk during predictable balance challenges.

Keywords: balance; biomechanics; gait; perturbations; stroke

Introduction

People maintain balance while walking through a combination of proactive and reactive control strategies. Unexpected balance challenges, such as black ice, depend on reactive control adjustments informed by sensory feedback, whereas expected challenges, such as a visible change in terrain, allow for proactive control informed by awareness of an upcoming disturbance (1). Anticipatory postural adjustments, one example of proactive control, are characterized by changes in muscle activity based on knowledge of the timing, direction, or size of the expected movement or perturbation and have been examined during arm flexion tasks (2–4), gait initiation (5–7), and standing perturbations (8). Recent studies have used auditory cueing to test how knowledge of perturbation timing influences proactive control strategies during walking (9,10). When an auditory warning was provided three seconds before waist-pulls, treadmill-belt accelerations, or treadmill decelerations, young adults made no anticipatory changes to whole-body angular momentum or fore-aft margins of stability before or during the perturbation (10). It is possible that the young adults examined in this study did not generate proactive responses because they were confident in their ability to maintain balance using purely reactive strategies.

It stands to reason that populations with higher fall risk and impaired reactive control, such as older adults and people post-stroke (11–14), may increase their reliance on proactive balance control strategies. For example, compared to younger adults, older adults demonstrate more proactive adjustments when approaching an overground obstacle, progressively slowing and taking shorter steps (15), which increases the time available to take in visual information and prepare their control strategy. However, the ability to make proactive adjustments may be affected by stroke-related impairments, because compared to similarly-aged peers without stroke, people post-stroke produce less anticipatory muscle activity before experiencing an expected standing perturbation (16) and before initiating gait (7), which results in slower stepping (17), and before clearing a visible obstacle on a treadmill (18), which results in a higher

failure rate (18,19). Proactive adjustments may be delayed or limited post-stroke due to somatosensory processing deficits (20), lower-extremity muscle weakness (21–23), abnormal muscle activation timing (24–26), or abnormal coordination (27,22,23). People post-stroke also frequently experience cognitive deficits in visual memory, executive function, and processing speed (28), which may affect their ability to understand audiovisual cues, plan an effective balance control strategy, and respond to cues promptly. Identifying how people post-stroke use timing cues to inform their control strategies could reveal how neurologic injury impairs proactive control during walking, possibly leading to falls.

The proactive balance adjustments made by older adults and people post-stroke while walking were recently examined by varying the predictability of perturbations using random- and regular-interval treadmill accelerations (29). Participants were not informed that perturbations would occur at fixed intervals in the regular condition, and, therefore, they needed to learn the regular pattern to proactively prepare for the perturbations. During regular perturbations, the group without stroke produced proactive adjustments that involved slowing the COM, but this strategy was not observed in their peers post-stroke (29). Given the need for participants to learn that the perturbations occurred regularly, it is possible that cognitive impairments limited the ability of people post-stroke to generate precise, proactive adjustments to their gait. In addition, the perturbations were applied unilaterally to the paretic side, but the control strategies may vary between the paretic and non-paretic sides due to the asymmetrical effects of a lesion.

Here, we examined how people with and without stroke used audiovisual cues to adjust proactive and reactive balance control strategies during single-belt treadmill accelerations. We modulated the predictability of these perturbations to evaluate how control strategies changed as a function of the timing information. Predictability was varied by changing the available response time (i.e., the time between the cue and perturbation) and timing precision (i.e., whether the perturbation occurred at a precise number of steps after the cue) of the audiovisual cues. We quantified changes in fore-aft MOS, positive mechanical leg work, and positive joint

work between cue conditions. We hypothesized that people without stroke would use timing cues to proactively perform less positive leg and joint work during the expected perturbation step, resulting in a larger MOS during the recovery step. Because a stroke can cause a range of impairments in perception, sensorimotor processing, and control, we expected people post-stroke to make smaller proactive changes with audiovisual cues, especially on their paretic side, resulting in smaller increases in the MOS during the recovery step compared to their peers without stroke.

Methods

Participants

Based on an a priori power analysis using pilot data, we recruited fourteen chronic stroke survivors and 14 age-/sex-matched adults without stroke (Table 1) to complete this protocol. Several of the participants post-stroke reported using an ankle brace or orthotic (n=6), cane (n=2), or wheelchair (n=1) at least some of the time. We recruited participants from the University of Southern California Registry for Aging and Rehabilitation Evaluation (RARE) and the Healthy Minds Research Volunteer Registry. People post-stroke were included if they had a single stroke over six months prior and could walk unassisted for at least five minutes. Exclusion criteria for either group included musculoskeletal conditions that affect walking ability, uncontrolled hypertension, cognitive impairment (Montreal Cognitive Assessment score <19/30 (30)), or uncorrected visual impairment.

Table 1. Group means ± standard deviations of demographic characteristics, clinical scores, and the p-values from Mann-Whitney U-tests evaluating group differences (bolded if p<0.05). MoCA: Montreal Cognitive Assessment, Mini-BEST: Mini Balance Evaluation Systems Test, ABC Scale: Activities-specific Balance Confidence Scale.

	Non-stroke	Stroke	p-value
Age (years)	58 ± 13	58 ± 13	0.945
Gender (M / F)	7 / 7	7 / 7	

Height (m)	1.70 ± 0.10	1.66 ± 0.09	0.222
Mass (kg)	76 ± 14	75 ± 17	0.800
Treadmill speed (m/s)	1.05 ± 0.11	0.66 ± 0.17	**<0.001**
MoCA (0-30)	27 ± 2	26 ± 2	0.523
Mini-BEST (0-28)	26 ± 2	21 ± 4	**<0.001**
ABC Scale (%)	96 ± 5	78 ± 15	**<0.001**
Fugl-Meyer (0-34)		23 ± 5	
Months since stroke		109 ± 55	

Experimental protocol

We first conducted a set of clinical exams, including the Fugl-Meyer Assessment of lower-extremity impairment post-stroke, Montreal Cognitive Assessment (MoCA) to screen for cognitive impairment, Activities-specific Balance Confidence (ABC) scale to evaluate subjective balance confidence during daily tasks, Mini Balance Evaluations Systems Test (Mini-BEST) to evaluate dynamic balance, and 10-Meter Walk Test (10MWT).

After completing the clinical exams, participants completed six trials walking on an instrumented split-belt treadmill (Bertec, OH) at their self-selected walking speed (Figure 1A). They completed two baseline trials, during which they acclimated to the treadmill for one minute and then experienced perturbations (three accelerations per side). Finally, participants completed four trials with trip-like perturbations, during which a custom Python script accelerated one belt at 5 m/s^2 up to 150% self-selected walking speed before returning to self-selected speed after the estimated stance time. We estimated swing and stance times as the average swing and stance times of the first 18 unperturbed steps per trial. Perturbations were triggered 250 ms before the estimated perturbation step to account for the time delays between sending the signal and accelerating the treadmill. We scaled perturbation magnitudes with walking speed to better normalize the relative effects of the perturbations across participants. Each perturbation trial consisted of 28 accelerations (14/side) every 12-22 steps with varied timing cues. The order of the four trials was randomized for every participant. The No-cue trial served as a reference condition. During the three trials with cues, participants received

audiovisual cues (Figure 1B) and information about what the cues signified. All cues included an arrow denoting the side on which the upcoming perturbation would occur. A green light signified there was no upcoming perturbation. General cues (beep and yellow light) were triggered three to eight steps before the perturbation, Exact cues (beep and red light) were triggered two steps before, and Countdown cues (beeps with 3 in green, 2 in yellow, and 1 in red lights) were triggered six steps before. On average, the cues allowed participants post-stroke 3.1, 1.0, and 3.7 seconds and participants without stroke 2.6, 0.8, and 3.2 seconds to prepare before perturbations in the General-, Exact-, and Countdown-cue conditions. The cues varied in the time available to prepare before the perturbation and the precision of the information provided, allowing us to evaluate several aspects of proactive control. Participants wore a safety harness but did not have access to handrails or assistive devices throughout the trials.

Figure 1. (A) The experimental setup with a participant wearing a full-body reflective marker set and safety harness on a split-belt treadmill while viewing cues on a monitor. (B) The four perturbation trials included No, General, Exact, and Countdown cues before the upcoming perturbation step (denoted by the red footstep). A green light indicated there were no upcoming perturbations. General cues (beep and yellow light) were triggered three to eight steps before the acceleration, Exact cues (beep and red light) were triggered two steps before the acceleration, and Countdown cues (beeps with 3 in green, 2 in yellow, and 1 in red lights) were triggered six, four, and two steps before the acceleration. All cues included an arrow to indicate the side of the upcoming perturbation.

Participants wore 54 reflective markers on their feet, shanks, thighs, pelvis, torso, upper arms, forearms, and head. We used Qualisys Track Manager to track their positions using a 10-camera motion capture system at 100 Hz (Qualisys, Sweden). We also collected force data from the treadmill force plates at 1000 Hz.

We analyzed the data in Visual3D (HAS-Motion, Ontario, Canada) and MATLAB (MathWorks, Natick, MA). A custom Visual3D pipeline gap-filled (third-order polynomial with a maximum gap of 10 frames) and low-pass filtered (Butterworth with 6-Hz cutoff) the marker data. We then fit a 13-segment model to the processed marker data and used this model to estimate the position of the body's center of mass (COM). Ankle, knee, and hip joint moments were calculated via inverse dynamics, and joint powers were computed as the dot products of these joint moments and their respective angular velocities. We also used Visual3D to identify and visually confirm the following events: 1) heel strikes (HS) and toe offs as the maximum and minimum fore-aft positions of the heel and toe markers, respectively, 2) perturbation timing based on peak belt speeds, 3) the onset of visual cues recorded in an analog channel, and 4) "crossovers" when participants' feet were placed on opposite belts. We removed force-based calculations during crossover events, which occurred, on average, four times per trial.

Data analysis

We evaluated how people with and without a stroke modulated the fore-aft margin of stability (MOS) at HS, its components (COM speed and its distance from the edge of the base of support), positive mechanical leg work, and positive ankle work in response to treadmill perturbations and how these variables changed with audiovisual cues.

Fore-aft margin of stability

We evaluated the fore-aft margin of stability (MOS) to test whether people proactively modulated dynamic stability before the perturbation and effectively reduced the destabilizing effects following perturbations with versus without audiovisual cues. The fore-aft MOS (Equation 1) was calculated as the anterior-posterior distance between the leading edge of the base of support (BOS) and the extrapolated COM position (31).

$$\text{MOS} = \text{BOS} - \left(x_{COM} + \frac{v_{COM}}{\sqrt{g/l}} \right) \qquad \text{(Eq. 1)}$$

We defined the BOS as the fore-aft position of the toe marker. The extrapolated COM was computed as the fore-aft COM position (x_{COM}) adjusted by fore-aft COM speed (v_{COM}), gravity (g), and leg length (l). Negative values indicate that the extrapolated COM is anterior to the BOS, and a corrective step or adjustment is generally required to prevent a fall. We hypothesized that people, especially those without stroke, would use audiovisual cues to proactively increase their MOS at the HS of the perturbation step to reduce the destabilizing effects. Subsequently, if participants effectively updated their strategy with audiovisual cues and reduced the destabilizing effects of the perturbation, we hypothesized that people, especially those without stroke, would have a larger MOS at the recovery step HS.

Since the MOS is a function of foot placement relative to the COM and COM speed, changes in the MOS could result from changes in either or both variables. Therefore, we decomposed the MOS into its components and evaluate how each changed across cueing conditions. These components included the COM speed at HS of the recovery step and the distance between the edge of the BOS and COM position (BOS-COM). We calculated the fore-aft COM speed relative to the treadmill belt at HS of the recovery step and BOS-COM as the fore-aft distance between the front toe marker and COM at HS of the perturbation and recovery steps.

Positive leg work

People may also proactively reduce mechanical leg work to reduce the destabilizing effects of expected treadmill perturbations (29). When given cues, people may perform less positive leg work during the pre-perturbation or perturbation steps to slow the forward COM speed or reduce the forward displacement of the COM to increase the recovery step MOS (Figure 2). We computed mechanical leg power (P_{leg}) using the individual limbs method (32) as the sum of the power performed by the leg on the body and the treadmill (Equation 2).

$$P_{leg} = \left(\vec{F} \cdot \vec{v}_{COM} \right) + \left(-\vec{F} \cdot \vec{v}_{TM} \right) \qquad \text{(Eq. 2)}$$

$\vec{F}$ is the ground reaction force vector, $\vec{v}_{COM}$ is the COM velocity vector, and $\vec{v}_{TM}$ is the treadmill velocity vector relative to the lab coordinate system. We calculated positive work for each step by integrating the positive portions of the power curve and then normalized work by the subject's mass.

Joint work

We characterized changes in ankle, knee, and hip joint work during the pre-perturbation and perturbation steps as potential contributors to the proactive changes in leg work. We computed joint power in Visual3D as the dot product of joint moment and angular velocity. Then, we calculated mass-normalized positive work per step by integrating the positive portions of joint power to evaluate energy generation. Due to a technical error, joint-level kinetic data were not collected during the Countdown cue trial for one of the participants post-stroke.

Statistical analysis

First, we determined if the effects of the perturbations on COM dynamics during the No-cue trial differed between groups by fitting a linear mixed-effects model for the change in median fore-aft COM speed at HS of the recovery step (Equation 3). The change in COM speed (Δv)

was computed relative to unperturbed walking. The model included an intercept ($\beta_{NS,No}$), fixed

effect of Leg (NS: non-stroke, P: paretic, NP: non-paretic), and random intercepts (b_i) to account

for variability between subjects and sides within the group post-stroke. If the results of an

ANOVA yielded a significant fixed effect of Leg, we reported the coefficients and their p-values,

which were Bonferroni corrected for the two comparisons.

$$\Delta v_i = \beta_{NS,No} + \beta_{P,No} P_i + \beta_{NP,No} NP_i + b_i + \varepsilon_i \qquad \text{(Eq. 3)}$$

Here, i is the observation number (1:42) and ε is the error.

Then, we fit separate linear mixed-effects models (Equation 4) for the changes in the

median fore-aft MOS, fore-aft COM speed, fore-aft BOS-COM distance, positive leg work, and

positive joint work relative to baseline (Δy) with fixed effects of Leg (NS: non-stroke, P: paretic,

NP: non-paretic), Cue (No, Gen: General, Ex: Exact, and Count: Countdown), and the

interaction between Leg and Cue. We considered changes in positive leg and joint work during

the pre-perturbation step, all metrics during the perturbation step, and MOS, COM speed, and

BOS-COM during the recovery step. The models included random intercepts (b_j) to account for

variability between subjects and sides within the group post-stroke. If the results of an ANOVA

yielded significant fixed effects, we reported the coefficients and their p-values, which were

Bonferroni corrected based on the number of comparisons (two for Leg, three for Cue, and six

for Leg*Cue).

$$\begin{aligned}
\Delta y_{ij} = {} & \beta_{NS,No} + \beta_{P,No} P_i + \beta_{NP,No} NP_i + \beta_{NS,Gen} Gen_i + \beta_{NS,Ex} Ex_i + \beta_{NS,Count} Count_i \\
& + \beta_{P,Gen}(P_i * Gen_i) + \beta_{NP,Gen}(NP_i * Gen_i) + \beta_{P,Ex}(P_i * Ex_i) \\
& + \beta_{NP,Ex}(NP_i * Ex_i) + \beta_{P,Count}(P_i * Count_i) \\
& + \beta_{NP,Count}(NP_i * Count_i) + b_j + \varepsilon_{ij}
\end{aligned} \qquad \text{(Eq. 4)}$$

where i is the observation number (1:168), j is the leg (1:42), β is the coefficient per fixed effect,

and ε is the error.

Lastly, we evaluated the relationships between changes in joint work and subsequent

changes to two components of MOS at the recovery step HS: the BOS-COM distance or the

COM speed. Here, we considered the components of MOS independently to determine the role of COM position versus speed in any MOS changes observed with cues. The two linear mixed-effects models included four predictors (the changes in ankle, knee, and hip work from the No-cue trial and Leg) but no intercept.

All statistical analyses were performed using MATLAB 2022b with an alpha level of 0.05. For every model, we plotted standardized residuals versus fitted values to check for heteroscedasticity, nonlinearity, and outliers. Outliers were identified if the standardized residual was outside three standard deviations, and the model was refit, excluding the outliers.

Results

Treadmill perturbations resulted in faster COM speeds in people with and without stroke

The treadmill perturbations caused an increase in fore-aft COM speed at the HS of the first recovery step for participants in both groups (Figure 3A). The perturbations in the No-cue condition increased COM speeds by 0.12 ± 0.01 m/s at the recovery step HS (Figure 3B) relative to unperturbed walking (Equation 3; $\beta_{NS,No}$ p<0.001). There was no significant difference between groups (Leg p=0.254), indicating that the perturbation was similar in size for participants with and without stroke.

Figure 3. (A) The fore-aft center of mass (COM) speed at heel strike (HS) of unperturbed (white), perturbed (black), and the first recovery (gray) steps for representative participants with and without stroke during the No-cue trial. (B) The change in COM speed at the recovery step HS after perturbations in the No-cue trial. The change was calculated relative to the average COM speed during unperturbed steps. Boxplot colors represent perturbations applied to the three legs: Non-stroke (blue), Paretic (gray), and Non-paretic (black). For paretic perturbations, the change in COM speed is computed as the change in speed during the non-paretic recovery step relative to non-paretic steps during the unperturbed baseline trial. Similarly, for non-paretic perturbations, the change in COM speed is computed as the change in speed during the paretic recovery step relative to paretic steps during the unperturbed baseline trial. The change in COM speed at the recovery step HS did not statistically differ between legs (p=0.254). Three total outliers were omitted to maximize the visual size of the boxplots.

Precise cues led to more positive MOS values during the recovery step, especially in people without stroke

Next, we tested whether people made proactive changes in the fore-aft MOS at the HS of the perturbation step (Figure 4A). The increase in belt speed during perturbations resulted in a general reduction in the MOS during the perturbation step relative to unperturbed walking. The MOS at the perturbation HS was 15.4 ± 0.7 cm smaller for the non-stroke group ($\beta_{NS,No}$ p<0.001), 9.0 ± 1.1 cm smaller in the group post-stroke during paretic perturbations ($\beta_{P,No}$ p<0.001), and 9.1 ± 1.1 cm smaller in the group post-stroke during non-paretic perturbations ($\beta_{NP,No}$ p<0.001). However, there was no effect of cueing on the perturbation step MOS (Cue: p=0.095; Interaction between Leg and Cue: p=0.965). Thus, cues did not elicit proactive changes in MOS at the time of the perturbation.

Figure 4. Change in the fore-aft (FA) margin of stability (MOS) during (A) perturbation and (B) recovery steps with No, General, Exact, and Countdown cues. The changes were calculated

relative to unperturbed walking. Boxplot colors represent perturbations to the Non-stroke (blue), Paretic (gray), and Non-paretic (black) limbs. Diamonds in the title indicate significant main effects of Leg. The results of post hoc analyses evaluating the main effect of Cue or its interaction with Leg are represented by vertical gray shading, indicating significant differences from the No-cue trial, and asterisks, indicating significant within-condition differences between the non-stroke and stroke groups. Five total outliers were omitted to maximize the visual size of the boxplots.

Next, we evaluated changes in the fore-aft MOS at HS of the recovery step to test if cues helped people improve stability (Figure 4B). For people without stroke, these MOS values remained 9.1 ± 0.9 cm smaller during the recovery step relative to unperturbed steps ($\beta_{NS,No}$ p<0.001), but this difference between unperturbed and recovery step MOS was smaller in the group post-stroke after both paretic ($\beta_{P,No}$ p<0.001) and non-paretic ($\beta_{NP,No}$ p=0.002) perturbations. There was a significant effect of cueing on the difference between the recovery step MOS and unperturbed MOS, but it differed between groups (Cue: p<0.001; Interaction between Leg and Cue: p=0.015). Compared to the No-cue trial, people without stroke increased MOS at the recovery step HS by 2.9 ± 0.5 cm with Exact cues ($\beta_{NS,Ex}$ p<0.001) and by 2.5 ± 0.5 cm with Countdown cues ($\beta_{NS,Count}$ p<0.001) but did not change MOS with General cues ($\beta_{NS,Gen}$ p=0.959). However, following paretic-leg perturbations, people post-stroke made smaller changes in recovery step MOS than their peers without stroke by 2.1 ± 0.7 cm with Exact cues ($\beta_{P,Exact}$ p=0.032) and by 2.3 ± 0.7 cm with Countdown cues ($\beta_{P,Count}$ p=0.012). Overall, more precise timing cues allowed people, especially those without stroke, to increase their recovery step MOS values.

We also evaluated changes in BOS-COM distance at the HS of perturbation (Figure 5A) and recovery (Figure 5B) steps separately from the calculation of MOS to understand the role of COM position in cue-related changes in stability. The BOS-COM distance on the perturbation step did not differ from unperturbed walking ($\beta_{NS,No}$ p=0.785) or between groups (Leg: p=0.875). However, there was a significant effect of cueing on BOS-COM distance (Cue: p=0.024; Interaction between Leg and Cue: p=0.927) with people increasing BOS-COM distances on the

perturbation step by 1.0 ± 0.4 cm with Exact cues ($\beta_{NS,Ex}$ p=0.040) but not with General ($\beta_{NS,Gen}$ p=1.000) or Countdown ($\beta_{NS,Count}$ p=0.055) cues. These changes were small, suggesting that proactively adjusting their COM position within their BOS was not the main control strategy, which is consistent with the lack of changes in the MOS during the perturbation step.

After the perturbation in the No-cue trial, the BOS-COM distance on the first recovery step (Figure 5B) was 5.5 ± 0.8 cm smaller for people without stroke ($\beta_{NS,No}$ p<0.001), 0.2 ± 1.1 cm larger for people post-stroke following paretic perturbations ($\beta_{P,No}$ p<0.001), and 1.5 ± 1.1 cm smaller following for people post-stroke non-paretic perturbations ($\beta_{NP,No}$ p<0.001) compared to unperturbed walking. Cueing significantly affected the size of the BOS-COM distance at the recovery step HS, but this effect differed between the groups (Cue: p<0.001; Interaction between Leg and Cue: p=0.017). For people without stroke, their BOS-COM distances on the recovery step were 1.3 ± 0.5 cm, 2.2 ± 0.5 cm, and 2.4 ± 0.5 cm less negative with General ($\beta_{NS,Gen}$ p=0.025), Exact ($\beta_{NS,Ex}$ p<0.001), and Countdown ($\beta_{NS,Count}$ p<0.001) cues, respectively, than in the No-cue trial. Compared to their peers without stroke, people post-stroke experienced smaller cue-related changes in their BOS-COM distances, as their changes were 1.8 ± 0.6 cm smaller after paretic perturbations with Exact cues ($\beta_{P,Ex}$ p=0.039) and 2.4 ± 0.6 cm and 1.9 ± 0.6 cm smaller after paretic ($\beta_{P,Count}$ p=0.002) and non-paretic ($\beta_{NP,Count}$ p=0.021) perturbations, respectively, with Countdown cues. Overall, people without stroke used cues to bring their COM closer to the edge of the BOS at the recovery step HS, mirroring the MOS results, but people post-stroke maintained similar BOS-COM distances, regardless of the perturbation or cueing condition.

Figure 5. The change in the fore-aft distance between the edge of the base of support (BOS) and center of mass (COM) at heel strike of (A) perturbation and (B) recovery steps with No, General, Exact, and Countdown cues. The changes were calculated relative to unperturbed walking. Boxplot colors represent perturbations to the Non-stroke (blue), Paretic (gray), and Non-paretic (black) limbs. Diamonds in the title indicate significant main effects of Leg. The results of post hoc analyses evaluating the main effect of Cue or its interaction with Leg are represented by vertical gray shading, indicating significant differences from the No-cue trial, and asterisks, indicating significant within-trial differences between the non-stroke and stroke groups. Eleven total outliers were omitted to maximize the visual size of the boxplots.

Lastly, we investigated the role of COM speed in cue-related changes to MOS by evaluating changes in fore-aft COM speed at the HS of perturbation (Figure 6A) and recovery steps (Figure 6B). Compared to unperturbed steps, COM speeds during perturbation steps in the No-cue trial were 0.53 ± 0.02 m/s faster for people without stroke ($\beta_{NS,No}$ p<0.001), 0.32 ± 0.03 m/s faster for people post-stroke on the paretic side ($\beta_{P,No}$ p<0.001), and 0.33 ± 0.03 m/s faster for people post-stroke on the non-paretic side ($\beta_{NP,No}$ p<0.001). While there was a significant effect of cueing on the change in COM speed at perturbation step HS (Cue: p=0.041), the effect did not yield significant Bonferroni-corrected comparisons between the No-cue trial and trials with General ($\beta_{NS,Gen}$ p=0.156), Exact ($\beta_{NS,Ex}$ p=1.000), or Countdown ($\beta_{NS,Count}$ p=1.000) cues, and this did not differ between groups (Interaction between Leg and Cue: p=0.074). Although the fore-aft COM speed was faster than unperturbed values at the recovery step HS, there were no effects of group (Leg: p=0.672), cueing (Cue: p=0.158), or the interaction between the two (Interaction between Leg and Cue: p=0.636). Therefore, while treadmill accelerations increased COM speed, as expected, these changes were not mitigated by audiovisual cues.

Figure 6. The change in the fore-aft center of mass (COM) speed at heel strike of (A) perturbation and (B) recovery steps with No, General, Exact, and Countdown cues. The changes were calculated relative to unperturbed walking. Boxplot colors represent perturbations to the Non-stroke (blue), Paretic (gray), and Non-paretic (black) limbs. Diamonds in the title indicate significant main effects of Leg. The results of post hoc analyses evaluating the main effect of Cue or its interaction with Leg are represented by vertical gray shading, indicating significant differences from the No-cue trial, and asterisks, indicating significant within-trial differences between the non-stroke and stroke groups. Fourteen total outliers were omitted to maximize the visual size of the boxplots.

People without stroke performed less positive leg work during cued perturbations

We next evaluated if participants proactively modified positive leg work during the pre-perturbation (Figure 7A) and perturbation (Figure 7B) steps when given cues about impending perturbations. While there was a significant main effect of cueing on changes in positive leg work during the pre-perturbation step (Cue: p=0.015), this effect did not remain after Bonferroni-corrected post hoc comparisons as there were no significant differences in positive leg work between the No-cue trial and the trials with General ($\beta_{NS,Gen}$ p=1.000), Exact ($\beta_{NS,Ex}$ p=0.086), or Countdown ($\beta_{NS,Count}$ p=1.000) cues.

Figure 7. The change in positive leg work during (A) pre-perturbation and (B) perturbation steps with No, General, Exact, and Countdown cues. The changes were calculated relative to unperturbed walking. Boxplot colors represent perturbations to the Non-stroke (blue), Paretic (gray), and Non-paretic (black) limbs. Diamonds in the title indicate significant main effects of Leg. The results of post hoc analyses evaluating the main effect of Cue or its interaction with Leg are represented by vertical gray shading, indicating significant differences from the No-cue trial, and asterisks, indicating significant within-trial differences between the non-stroke and stroke groups. Sixteen total outliers were omitted to maximize the visual size of the boxplots.

During the perturbation step (Figure 7B), the group without stroke performed 0.22 ± 0.02 J/kg more positive leg work than they did on unperturbed steps ($\beta_{NS,No}$ p<0.001), which was a greater change than the group post-stroke who performed by 0.14 ± 0.03 J/kg during paretic perturbations ($\beta_{P,No}$ p<0.001) and 0.10 ± 0.03 J/kg during non-paretic perturbations ($\beta_{NP,No}$ p=0.002). Audiovisual cues also affected changes in positive leg work during the perturbation step, but this change differed between the groups (Cue: p<0.001; Interaction between Leg and Cue: p=0.005). People without stroke performed 0.06 ± 0.01 J/kg and 0.08 ± 0.01 J/kg less positive leg work during perturbations with Exact ($\beta_{NS,Ex}$ p<0.001) and Countdown ($\beta_{NS,Count}$ p<0.001) cues, respectively, than during the No-cue trial. Compared to the group without stroke, the group post-stroke made smaller reductions in positive leg work by 0.06 ± 0.02 J/kg and 0.05 ± 0.02 J/kg during paretic ($\beta_{P,Count}$ p=0.008) and non-paretic ($\beta_{NP,Count}$ p=0.041) perturbations, respectively, with Countdown cues. Overall, audiovisual cues allowed people, especially those without stroke, to perform less positive leg work during the perturbation steps.

People without stroke performed less positive joint work during cued perturbations

Along with changes in leg work, we evaluated the changes in ankle, knee, and hip work to probe the joint-level contributions to this balance control strategy. During the pre-perturbation step, there were no changes in positive ankle (Cue: p=0.140; Interaction between Leg and Cue: p=0.739), knee (Cue: p=0.088; Interaction between Leg and Cue: p=0.398), or hip (Cue: p=0.393; Interaction between Leg and Cue: p=0.941). In general, during perturbations in people without stroke, the ankle performs more positive work during push-off (Figure 8A), the knee

performs similar amounts of positive work (Figure 8B), and the hip performs more positive work (Figure 8C) compared to unperturbed walking.

Figure 8. Average (A) ankle, (B) knee, and (C) hip power generated by each leg during unperturbed baseline steps (black), perturbed steps in the No-cue trial (gray), and perturbed steps in the Countdown-cue trial (blue). Trajectories were averaged across strides and then across participants. The ankle performs some negative work during the first half of the stance phase but generates much more positive work during push-off. The knee alternates between performing positive and negative work throughout the step cycle. The hip performs positive work during early to midstance, as well as push-off, and minimal negative work. In general, the paretic limb (middle) performs less positive work relative to the non-paretic limb (right) and to people without stroke (left).

During the perturbation step, people without stroke used cues to perform less positive ankle work. Compared to unperturbed steps, people without stroke performed 0.23 ± 0.01 J/kg more positive ankle work during perturbed steps (Figure 9A; $\beta_{NS,No}$ p<0.001). This was a greater increase than the group post-stroke made during paretic ($\beta_{P,No} = -0.17 \pm 0.02$ J/kg, p<0.001) and non-paretic perturbations ($\beta_{NP,No} = -0.15 \pm 0.02$ J/kg, p<0.001). Cueing also significantly affected the changes in positive ankle work performed during the perturbation step, though this again differed between groups (Cue p<0.001; Interaction between Leg and Cue: p<0.001). Compared to the No-cue trial, people without stroke performed 0.04 ± 0.01 J/kg, 0.07 ± 0.01 J/kg, and 0.08 ± 0.01 J/kg less positive ankle work during perturbations with General ($\beta_{NS,Gen}$ p=0.001), Exact ($\beta_{NS,Ex}$ p<0.001), and Countdown ($\beta_{NS,Count}$ p<0.001) cues, respectively. However, these cue-related reductions were smaller in the group post-stroke on the paretic side with Exact ($\beta_{P,Ex} = 0.05 \pm 0.01$ J/kg, p<0.001) and Countdown cues ($\beta_{P,Count} = 0.06 \pm 0.01$ J/kg, p<0.001) and on the non-paretic side with General ($\beta_{NP,Gen} = 0.04 \pm 0.01$ J/kg, p=0.046), Exact ($\beta_{NP,Ex} = 0.06 \pm 0.01$ J/kg, p<0.001), and Countdown ($\beta_{NP,Count} = 0.06 \pm 0.01$ J/kg, p<0.001) cues.

Figure 9. The change in positive (A) ankle, (B) knee, and (C) hip work during perturbation steps with No, General, Exact, and Countdown cues. The changes were calculated relative to unperturbed walking. Boxplot colors represent perturbations to the Non-stroke (blue), Paretic (gray), and Non-paretic (black) limbs. Diamonds in the title represent significant main effects of Leg. The results of post hoc analyses evaluating the main effect of Cue or its interaction with Leg are represented by vertical gray shading, indicating significant differences from the No-cue trial, and asterisks, indicating significant within-trial differences between the non-stroke and stroke groups. Outliers are omitted to maximize the visual size of the boxplots.

We also evaluated if positive knee work changed during perturbation steps (Figure 9B). While people without stroke did not modify positive knee work during perturbations in the No-cue trial ($\beta_{NS,No}$ p=0.120), people post-stroke performed 0.04 ± 0.01 J/kg more positive knee work during paretic perturbations than the group without stroke relative to unperturbed values ($\beta_{P,No}$ p=0.009). There was also a significant effect of cueing on the change in positive knee work during perturbation steps (Cue: p=0.005; Interaction between Leg and Cue: p=0.166), with people performing 0.01 ± 0.00 J/kg less work with General ($\beta_{NS,Gen}$ p=0.006) and Countdown ($\beta_{NS,Count}$ p=0.004) cues, but not Exact cues ($\beta_{NS,Ex}$ p=0.097). While the effect sizes were small, people may have used cues to reduce the amount of positive knee work performed during perturbation steps.

Lastly, we tested for changes in positive hip work during perturbation steps (Figure 9C). People performed 0.04 ± 0.01 J/kg more positive hip work during perturbed steps in the No-cue trial compared to unperturbed steps ($\beta_{NS,No}$ p<0.001), and there was no significant difference between groups (Leg: p=0.309). Cueing had a significant effect on positive hip work (Cue: p=0.027; Interaction between Leg and Cue: p=0.264), as people performed 0.02 ± 0.01 J/kg less work with General cues ($\beta_{NS,Gen}$ p=0.009) but not Exact ($\beta_{NS,Ex}$ p=0.266) or Countdown ($\beta_{NS,Count}$ p=0.131) cues. Therefore, people may have used General cues to perform less positive work at the hip during perturbation steps, but the effect size of this change is small relative to the changes at the ankle.

Changes to ankle and hip work are associated with changes to the BOS-COM distance

Finally, we evaluated the relationship between changes in positive joint work during the perturbation steps and each of the components of the recovery step fore-aft MOS: the BOS-COM distance and COM speed at HS. The two models were fit to data collapsed across all cued trials, and fixed effects included the change in positive ankle, knee, and hip work and Leg

(Non-stroke, Paretic, Non-paretic). We removed four outliers from the first model and one from the second model before refitting for the final analyses. Changes in positive ankle ($p<0.001$) and hip ($p=0.045$) work were associated with changes in BOS-COM distance at the recovery step HS, whereas the change in positive knee work ($p=0.684$) and Leg ($p=0.506$) were not ($R^2=0.366$). With every 0.01 J/kg reduction in positive ankle work, the BOS-COM distance at the recovery step increased by 0.33 ± 0.04 cm. Additionally, with every 0.01 J/kg reduction in positive hip work, the BOS-COM distance increased by 0.15 ± 0.07 cm. The change in positive knee work ($p=0.002$), change in positive hip work ($p=0.041$), and Leg ($p=0.018$) were associated with the change in COM speed, whereas the change in positive ankle work ($p=0.451$) was not ($R^2=0.092$). However, this model had a poor fit, suggesting that these changes in knee and hip work during cued perturbations were not strongly associated with changes in COM speed.

Discussion

People post-stroke report frequent falls (12), suggesting their balance control is impaired. Many real-world balance challenges are predictable, allowing for both proactive and reactive control, but little is known about how a stroke influences the roles of each control strategy during predictable gait perturbations. This study investigated how people with and without stroke use audiovisual cues to adjust their balance control strategies during trip-like treadmill perturbations. We found that people without stroke used the Exact and Countdown cues to increase recovery step MOS, and this was associated with a reduction in positive leg and joint work during the perturbations. On the other hand, people post-stroke did not make these proactive adjustments, but instead maintained the same reactive strategies, regardless of cueing, suggesting that control strategies are less tuned to information about impending perturbations in people post-stroke.

People without stroke use timing cues to reduce the destabilizing effects of treadmill perturbations

Participants without stroke reduced the destabilizing effects of treadmill perturbations when provided audiovisual cues that began one stride (Exact) or three strides (Countdown) before the perturbation. In both conditions, perturbations caused smaller reductions in the subsequent MOS compared to the No-cue trial. In contrast, there were no changes in the MOS with General cues provided three to eight steps before the perturbation, underlining the need for precise information about perturbation timing for participants to proactively update their motor plans. These findings complement prior work, which demonstrated that people without stroke reduce perturbation-induced increases in COM speed during regular versus random accelerations (29). However, these results differ from other findings that young adults made no significant changes to whole-body angular momentum (WBAM) or MOS during the pre-perturbation and perturbation strides when provided a three-second countdown before treadmill accelerations, decelerations, and waist-pulls (10). It is possible that the treadmill accelerations used in that study were not sufficiently destabilizing or that the young adults felt confident in their reactive strategies, both of which may negate the need to generate proactive adjustments. Notably, the average age of our participants was 58 years old, and we may expect older adults to make more proactive changes if they are less confident in their reactive balance than young adults. Therefore, although the mechanisms by which young adults modulate their proactive control strategies remain unclear, we found that older adults utilize precise audiovisual warnings to improve their stability following treadmill accelerations.

Modulating mechanical work with cues may be a key proactive control strategy for people without neurologic injury

Next, we investigated the strategies participants used to reduce the destabilizing effects of perturbations when provided with audiovisual cues. We tested for proactive changes during the pre- and perturbation steps and found that people without stroke reduced positive leg, ankle, knee, and hip work during the perturbation step, perhaps to slow their forward COM speed or move their COM to a more posterior position during the perturbation. Positive ankle work is generally performed during the push-off phase of stance, so reducing push-off could have helped people improve subsequent stability. Consistent with this interpretation, we found that changes in positive ankle work during the perturbation step were negatively correlated with changes in COM position relative to the edge of the BOS at HS of the recovery step, suggesting people reduced push-off to maintain their COM position. While changes in positive hip work were also associated with these changes in BOS-COM distance, the effect sizes were larger for the changes in positive ankle work. Parallel changes in positive leg and ankle work are consistent with prior research that found the ankle joint to best reflect changes in leg work during unexpected treadmill accelerations (33). Therefore, performing less positive leg and ankle work during cued perturbations – especially when the cues were more precise – appears to be a key strategy for mitigating the destabilizing effects of predictable perturbations that would otherwise lead to forward losses of balance.

A stroke may shift dependence onto a general reactive strategy versus proactive adjustments

In general, we found that people post-stroke during not make proactive balance adjustments when provided with cues about the timing of impending perturbations. Unlike their peers without stroke, people post-stroke did not increase their MOS during the recovery step following paretic-side perturbations with Exact or Countdown cues. Considering the

perturbations induced similar changes in fore-aft COM speed across groups, proactive adjustments could still have been beneficial for people post-stroke. Similarly, people post-stroke did not use the cues to perform less positive leg or ankle work during the perturbation steps. This group difference is consistent with our previous findings that only people without stroke performed less positive leg work during regular- versus random-interval perturbations (29). It is unclear whether this lack of proactive changes is due to abnormal muscle coordination (27,34) preventing effective changes in muscle activity, spasticity (35) preventing joint extension, sensory dysfunction (20) reducing trust in feedback that could inform proactive updates, or other stroke-related changes to neuromotor control. Future studies should incorporate electromyography, which could characterize anticipatory muscle activity and stroke-related differences in lower-level proactive control strategies. Overall, our findings add to the existing evidence that a stroke impairs proactive balance control strategies during standing (16) and walking (19,18,29).

If proactive control strategies are less effective following a stroke, a larger reactive response may be necessary to prevent falling. However, a stroke may also impair reactive control during walking, as people post-stroke experience more forward rotation during random treadmill accelerations and require an additional recovery step before returning to baseline than their peers without stroke (14). One reactive strategy may involve adjusting foot placement and COM position to maintain the size of the BOS, whether done passively or actively. Previous research has found strong associations between changes in COM dynamics and subsequent foot placement during walking in both unperturbed (36) and perturbed conditions (37). Our participants post-stroke demonstrated this proposed reactive response following all perturbations by maintaining BOS-COM distances that were similar in magnitude to unperturbed distances. However, the group post-stroke did not use the cues to modulate this reactive strategy as those without stroke did. This unchanging reactive response suggests that people

post-stroke implement control strategies that are more general and less flexible than those of people without neurologic injury.

Improving balance outcomes is a key rehabilitation goal following a stroke, and the lack of proactive adjustments in our participants post-stroke suggests that future interventions could focus on proactive control. Perturbation training has emerged as a research tool to enable people post-stroke to practice balance control strategies, with some mixed but promising initial improvements in dynamic balance and balance confidence (38–41). However, the emphasis has been on random perturbations that require reactive responses, and it remains unknown whether proactive strategies during gait can be trained and transferred to real-world conditions. Future studies should evaluate if people post-stroke can practice modulating leg work, ankle work, or step lengths – perhaps with biofeedback – in perturbed conditions to improve balance.

Limitations

There are a few limitations to our current study that could be addressed in future research. We measured changes in MOS, positive leg work, and lower-extremity joint work, but other proactive strategies, such as modulating limb impedance (42–44), may have been implemented to improve stability during cued perturbations. Additionally, because our groups walked at different speeds and the perturbations elicited changes in recovery step MOS with varying magnitudes, it may be challenging to disentangle the effects of stroke versus speed on one's ability to utilize precise timing cues to improve stability. We scaled the perturbation magnitudes with walking speed to allow people to walk at their comfortable speeds and normalize the perturbations, but it is possible that those walking at slower speeds, often in the group post-stroke, did not experience destabilizations large enough to warrant changes in control. However, our perturbations elicited comparable relative changes in COM speed across groups and conditions, suggesting we perturbed groups similarly. Therefore, while we demonstrated that people post-stroke made limited proactive adjustments during perturbed

walking, future studies should continue to investigate the underlying mechanisms of proactive and reactive balance control using other perturbation conditions and tools.

Conclusions

We aimed to investigate how people with and without stroke modify their proactive control strategies during treadmill perturbations with audiovisual cues. Adults without a history of stroke used audiovisual cues that began one and three strides before treadmill perturbations to mitigate the destabilizing effects, as measured by smaller reductions in the fore-aft margin of stability compared to the margins of stability in the absence of cues. This improvement in stability may have been accomplished by a range of proactive control strategies, including performing less positive leg and joint work, especially at the ankle, during the expected perturbation steps and then better positioning the body's center of mass relative to the base of support. As hypothesized, people post-stroke made fewer adjustments during predictable perturbations than their peers without stroke. People post-stroke did not use audiovisual cues to perform less work, but instead, they reactively maintained COM positioning regardless of perturbation predictability. Overall, our findings suggest that people with chronic stroke rely on a general reactive strategy, even when the balance disturbance is explicitly expected, and thus may be at risk for falls that could be prevented with proactive adjustments.

Data availability

The processed data and code to create the figures can be found https://osf.io/t8ue4/.

Acknowledgments

We thank our study participants for their time, effort, and contributions to this work.

Grants

This project was supported by the American Heart Association Predoctoral Fellowship 23PRE1012432, the Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health under award number R01HD091184, and NSF award number 2319710.

Disclosures

The authors do not have any conflicts of interest to report.

Author contributions

T.C. conceptualized and designed the research, performed experiments, analyzed data, interpreted results of experiments, prepared figures, drafted the manuscript, edited and revised the manuscript, and approved the final version of the manuscript. J.F. conceptualized and designed the research, analyzed data, interpreted results of experiments, edited and revised the manuscript, and approved the final version of the manuscript.

References

1. Patla AE. Strategies for dynamic stability during adaptive human locomotion. IEEE Engineering in Medicine and Biology Magazine. 2003 Mar;22(2):48–52.

2. Dick JP, Rothwell JC, Berardelli A, Thompson PD, Gioux M, Benecke R, et al. Associated postural adjustments in Parkinson's disease. J Neurol Neurosurg Psychiatry. 1986 Dec;49(12):1378–85.

3. Aruin AS, Latash ML. Directional specificity of postural muscles in feed-forward postural reactions during fast voluntary arm movements. Exp Brain Res. 1995 Mar 1;103(2):323–32.

4. Aruin AS, Latash ML. The role of motor action in anticipatory postural adjustments studied with self-induced and externally triggered perturbations. Exp Brain Res [Internet]. 1995 [cited 2025 June 3];106(2). Available from: http://link.springer.com/10.1007/BF00241125

5. Crenna P, Frigo C. A motor programme for the initiation of forward-oriented movements in humans. The Journal of Physiology. 1991;437(1):635–53.

6. Brunt D, Vander Linden DW, Behrman AL. The relation between limb loading and control parameters of gait initiation in persons with stroke. Archives of Physical Medicine and Rehabilitation. 1995 July 1;76(7):627–34.

7. Sousa ASP, Silva A, Santos R. Ankle anticipatory postural adjustments during gait initiation in healthy and post-stroke subjects. Clinical Biomechanics. 2015 Nov 1;30(9):960–5.

8. Kanekar N, Aruin AS. Aging and balance control in response to external perturbations: role of anticipatory and compensatory postural mechanisms. AGE. 2014 June 1;36(3):1067–77.

9. Kreter N, Lybbert C, Gordon KE, Fino PC. The effects of physical and temporal certainty on human locomotion with discrete underfoot perturbations. Journal of Experimental Biology. 2022 Oct 13;225(19):jeb244509.

10. Eichenlaub EK, Urrego DD, Sapovadia S, Allen J, Mercer VS, Crenshaw JR, et al. Susceptibility to walking balance perturbations in young adults is largely unaffected by anticipation. Human Movement Science. 2023 June 1;89:103070.

11. Blake AJ, Morgan K, Bendall MJ, Dallosso H, Ebrahim SBJ, Arie THD, et al. Falls by elderly people at home: Prevalence and associated factors. Age and Ageing. 1988 Jan 1;17(6):365–72.

12. Schmid AA, Yaggi HK, Burrus N, McClain V, Austin C, Ferguson J, et al. Circumstances and consequences of falls among people with chronic stroke. 2013;50(9):1277–86.

13. McCrum C, Karamanidis K, Grevendonk L, Zijlstra W, Meijer K. Older adults demonstrate interlimb transfer of reactive gait adaptations to repeated unpredictable gait perturbations. GeroScience. 2020 Feb 1;42(1):39–49.

14. Liu C, McNitt-Gray JL, Finley JM. Impairments in the mechanical effectiveness of reactive balance control strategies during walking in people post-stroke. Frontiers in Neurology. 2022;13.

15. Muir BC, Haddad JM, Heijnen MJH, Rietdyk S. Proactive gait strategies to mitigate risk of obstacle contact are more prevalent with advancing age. Gait & Posture. 2015 Jan 1;41(1):233–9.

16. Curuk E, Lee Y, Aruin AS. Individuals With Stroke Use Asymmetrical Anticipatory Postural Adjustments When Counteracting External Perturbations. Motor Control. 2019 Oct;23(4):461–71.

17. Melzer I, Goldring M, Melzer Y, Green E, Tzedek I. Voluntary stepping behavior under single- and dual-task conditions in chronic stroke survivors: A comparison between the involved and uninvolved legs. Journal of Electromyography and Kinesiology. 2010 Dec 1;20(6):1082–7.

18. van Swigchem R, van Duijnhoven HJR, den Boer J, Geurts AC, Weerdesteyn V. Deficits in Motor Response to Avoid Sudden Obstacles During Gait in Functional Walkers Poststroke. Neurorehabil Neural Repair. 2013 Mar 1;27(3):230–9.

19. Den Otter AR, Geurts ACH, de Haart M, Mulder T, Duysens J. Step characteristics during obstacle avoidance in hemiplegic stroke. Exp Brain Res. 2005 Feb 1;161(2):180–92.

20. Sullivan JE, Hedman LD. Sensory Dysfunction Following Stroke: Incidence, Significance, Examination, and Intervention. Topics in Stroke Rehabilitation. 2008 May 1;15(3):200–17.

21. Olney SJ, Richards C. Hemiparetic gait following stroke. Part I: Characteristics. Gait & Posture. 1996 Apr 1;4(2):136–48.

22. Neckel N, Pelliccio M, Nichols D, Hidler J. Quantification of functional weakness and abnormal synergy patterns in the lower limb of individuals with chronic stroke. J NeuroEngineering Rehabil. 2006 July 20;3(1):17.

23. Sánchez N, Acosta AM, Lopez-Rosado R, Stienen AHA, Dewald JPA. Lower Extremity Motor Impairments in Ambulatory Chronic Hemiparetic Stroke: Evidence for Lower Extremity Weakness and Abnormal Muscle and Joint Torque Coupling Patterns. Neurorehabil Neural Repair. 2017 Sept 1;31(9):814–26.

24. Jones RD, Donaldson IMacG, Parkin PJ. Impairment and recovery of ipsilateral sensory-motor function following unilateral cerebral infarction. Brain. 1989 Feb 1;112(1):113–32.

25. Marigold DS, Eng JJ, Timothy Inglis J. Modulation of ankle muscle postural reflexes in stroke: influence of weight-bearing load. Clinical Neurophysiology. 2004 Dec 1;115(12):2789–97.

26. Marigold DS, Eng JJ. Altered timing of postural reflexes contributes to falling in persons with chronic stroke. Exp Brain Res. 2006 June 1;171(4):459–68.

27. Higginson JS, Zajac FE, Neptune RR, Kautz SA, Delp SL. Muscle contributions to support during gait in an individual with post-stroke hemiparesis. Journal of Biomechanics. 2006 Jan 1;39(10):1769–77.

28. Nakling AE, Aarsland D, Næss H, Wollschlaeger D, Fladby T, Hofstad H, et al. Cognitive Deficits in Chronic Stroke Patients: Neuropsychological Assessment, Depression, and Self-Reports. Dementia and Geriatric Cognitive Disorders Extra. 2017 Aug 29;7(2):283–96.

29. Cornwell T, Finley J. Stroke impairs the proactive control of dynamic balance during predictable treadmill accelerations. Journal of The Royal Society Interface. 2025 Oct;22(231):20250336.

30. Trzepacz PT, Hochstetler H, Wang S, Walker B, Saykin AJ, for the Alzheimer's Disease Neuroimaging Initiative. Relationship between the Montreal Cognitive Assessment and Mini-mental State Examination for assessment of mild cognitive impairment in older adults. BMC Geriatr. 2015 Sept 7;15(1):107.

31. Hof AL, Gazendam MGJ, Sinke WE. The condition for dynamic stability. J Biomech. 2005 Jan;38(1):1–8.

32. Selgrade BP, Thajchayapong M, Lee GE, Toney ME, Chang YH. Changes in mechanical work during neural adaptation to asymmetric locomotion. Journal of Experimental Biology. 2017 Aug 15;220(16):2993–3000.

33. Golyski PR, Sawicki GS. Which lower limb joints compensate for destabilizing energy during walking in humans? Journal of The Royal Society Interface. 2022 June;19(191):20220024.

34. Arene N, Hidler J. Understanding motor impairment in the paretic lower limb after a stroke: a review of the literature. Top Stroke Rehabil. 2009;16(5):346–56.

35. Li S. Spasticity, Motor Recovery, and Neural Plasticity after Stroke. Front Neurol. 2017 Apr 3;8.

36. Wang Y, Srinivasan M. Stepping in the direction of the fall: the next foot placement can be predicted from current upper body state in steady-state walking. Biology Letters. 2014 Sept;10(9):20140405.

37. Liu C, Valero-Cuevas FJ, Finley JM. Generalizability of foot placement control strategies during unperturbed and perturbed gait. Royal Society Open Science. 2024 May;11(5):231210.

38. Handelzalts S, Kenner-Furman M, Gray G, Soroker N, Shani G, Melzer I. Effects of Perturbation-Based Balance Training in Subacute Persons With Stroke: A Randomized Controlled Trial. Neurorehabil Neural Repair. 2019 Mar 1;33(3):213–24.

39. Punt M, Bruijn SM, van de Port IG, de Rooij IJM, Wittink H, van Dieën JH. Does a Perturbation-Based Gait Intervention Enhance Gait Stability in Fall-Prone Stroke Survivors? A Pilot Study. Journal of Applied Biomechanics. 2019 June;35(3):173–81.

40. Esmaeili V, Juneau A, Dyer JO, Lamontagne A, Kairy D, Bouyer L, et al. Intense and unpredictable perturbations during gait training improve dynamic balance abilities in chronic hemiparetic individuals: a randomized controlled pilot trial. Journal of NeuroEngineering and Rehabilitation. 2020 June 17;17(1):79.

41. Bhatt T, Dusane S, Gangwani R, Wang S, Kannan L. Motor adaptation and immediate retention to overground gait-slip perturbation training in people with chronic stroke: an experimental trial with a comparison group. Front Sports Act Living [Internet]. 2023 Sept 13 [cited 2025 Feb 6];5. Available from: https://www.frontiersin.org/journals/sports-and-active-living/articles/10.3389/fspor.2023.1195773/full

42. Krutky MA, Trumbower RD, Perreault EJ. Influence of environmental stability on the regulation of end-point impedance during the maintenance of arm posture. Journal of Neurophysiology. 2013 Feb 15;109(4):1045–54.

43. Rouse EJ, Hargrove LJ, Perreault EJ, Kuiken TA. Estimation of Human Ankle Impedance During the Stance Phase of Walking. IEEE Transactions on Neural Systems and Rehabilitation Engineering. 2014 July;22(4):870–8.

44. Jakubowski KL, Ludvig D, Bujnowski D, Lee SSM, Perreault EJ. Simultaneous Quantification of Ankle, Muscle, and Tendon Impedance in Humans. IEEE Transactions on Biomedical Engineering. 2022 Dec;69(12):3657–66.