# Natural evolution of intermale sexual behavior by multiple pheromone switches among *Drosophila* species

Youcef Ouadah [1], Thomas H. Naragon [2], Hayley Smihula [2], Emily L. Behrman [3,†],
Mohammed A. Khallaf [4,‡], Yun Ding [5], David L. Stern [3], Joseph Parker [2], and David J. Anderson [1,*]

[1] Division of Biology and Biological Engineering, Tianqiao and Chrissy Chen Institute for Neuroscience, and Howard Hughes Medical Institute, Caltech, Pasadena, CA, USA

[2] Division of Biology and Biological Engineering, Caltech, Pasadena, CA, USA

[3] Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, USA

[4] Department of Evolutionary Neuroethology, Max Planck Institute for Chemical Ecology, Jena, Germany

[5] Department of Biology, University of Pennsylvania, Philadelphia, PA, USA

[†] Present address: Department of Biological Sciences, Dartmouth College, Hanover, NH, USA

[‡] Present Address: Molecular Physiology of Somatic Sensation Laboratory, Max Delbrück Center for Molecular Medicine, Berlin, Germany and Department of Zoology and Entomology, Faculty of Science, Assiut University, Assiut, Egypt

[*] Correspondence: wuwei@caltech.edu

**Abstract**

We have identified a *Drosophila* species in which males exhibit spontaneous, elaborate, and

robust intermale sexual behavior. Males of *D. santomea*, a West African island endemic, distinguish

conspecific sexes but court males and females promiscuously and seldom attack. Elevated intermale

courtship derives from at least three changes in two separate pheromone systems. In males, the sexually

monomorphic cuticular pheromone 7-tricosene promotes rather than inhibits courtship and the courtship-

inhibiting olfactory pheromone cVA is reduced 84-92% compared to close relatives, including the sibling

species *D. yakuba*. The third change is surprisingly in *D. santomea* females, where cVA suppresses rather

than promotes sexual receptivity. The female cVA switch and male cVA reduction may have co-evolved

to maintain efficient intraspecific mating in *D. santomea* but prevent sympatric hybridization with *D.*

*yakuba*, or to reduce intraspecific aggression. We find that high intermale courtship and low cVA also co-

occur and appear selectively derived in a distant monomorphic species *D. persimilis*, implying

pheromonal and behavioral convergence in the two recently speciated taxa. The data suggest that

sequential changes in the behavioral valence and levels of pheromones explain the recent evolutionary

emergence of intermale sexual behavior in *Drosophila*.

**Introduction**

Social behaviors, including aggregation, aggression, courtship, mating, parenting, and cooperation (*1*), are remarkably diverse in the animal kingdom, and can vary between closely related sibling species (*2*) or even entire orders (*3*). The ultimate causes of social behaviors (in terms of natural and sexual selection) have received considerable theoretical attention (*4*, *5*) and proximate causes (including genetic loci, hormones, neural circuits, pheromones, and sensory releasers) identified (*6–15*). The ~2,000 species of the genus *Drosophila* (*16*) have historically excelled in this regard as models for intersexual social behaviors (*17–22*), providing notably an unparalleled genetic, neural, and pheromonal understanding of the iconic courtship dance of *D. melanogaster* males (*23–25*). Comparisons of male courtship elsewhere in the genus have revealed flexible substrates, in genomes (*26–29*) and nervous systems (*30–33*), of its rapid evolution and revealed general principles of reproductive function and isolation (*34*, *35*). However, systematic investigations of intrasexual social behaviors between *Drosophila* males or females have comparatively lagged.

Interactions between same-sex conspecifics can vary greatly across species, strains, individuals, and time (*36*, *37*), and can be as opposed as altruistic cooperation (*38*) versus lethal aggression (*39*). Such flexibility implies that intrasexual social behaviors can evolve quickly under varying selection pressures or neutral drift, but underlying changes in pheromone systems, neural circuits, and genomes are often difficult to pinpoint. We sought to identify phylogenetic patterns of variation in intermale social behavior in *Drosophila* and, by leveraging experimental strengths of the genus, reveal proximate and ultimate evolutionary causes. We have found a species, *D. santomea*, in which males show promiscuous sexual behavior toward other males that includes all species-typical motor and acoustic features of the female-directed courtship dance, and is accompanied only rarely by overt aggression. Three distinct changes in pheromone levels or behavioral valence compared to *D. melanogaster* have occurred during *D. santomea* evolution, both before and after divergence from the sibling species *D. yakuba*. Together these changes have highly and selectively amplified intermale courtship without eliminating sex discrimination or the latent potential for intense attack, which can be revealed by optogenetic activation of conserved aggression-promoting neural circuits. The results, together with information from natural history and phylogenetic inference, provide insight into the pheromonal mechanism and evolutionary logic of recent social evolution in *Drosophila* that may apply broadly to other taxa.

**Results**

**Evolutionary variation in male-male social behavior within the genus *Drosophila***

In the presence of food or a female, pairs of single-housed *D. melanogaster* males will fight (*40–*

*47*), employing a stereotyped set of contact- and non-contact-mediated motor actions including lunging,

tussling, and wing threat (*48–50*). We assessed aggression (and other intermale social behaviors)

comparatively in 15 *Drosophila* species, screening for qualitative and quantitative variation in male social

engagement, frequency, and motor pattern. We sampled species densely within the *melanogaster*

subgroup (7 tested of 9 assigned) (*51*) to monitor potential behavior variation at close phylogenetic

distance to *D. melanogaster* with high resolution. At larger evolutionary distances we sampled sparsely

(typically one species per subgenus group or subgroup) to broadly survey genus social evolution, apart

from including both members of the *D. pseudoobscura* / *D. persimilis* sibling species pair from the

*obscura* group (*52*). We standardized assay conditions but occasionally varied the food source over which

flies competed to account for host specializations (*53*). Fly behavior was measured in dyadic male-male

interactions, by combining manual scoring on a subset of representative recordings (Fig. 1A,B, 1-S1A)

with automated behavior classification for the remainder (Fig. 1-S2A,B) (*54–56*).

Aggression among *D. melanogaster* males was the most prominent and intense of any

*melanogaster* subgroup species tested (Fig. 1A-C, 1-S2C,E). Species in the neighboring *simulans* clade

(*D. simulans*, *D. mauritiana*, and *D. sechellia*) also fought, though less frequently, using a similar set of

actions (Fig. 1A-C, 1-S2C,E). Varying foods usually had little effect on the behavior of specialists, with

the notable exception of *D. sechellia* which fought significantly more over *Morinda* than the standard

apple juice (Fig. 1-S3A), consistent with its specialization (*53*). Male-male interactions in *D.*

*melanogaster* and *simulans* clade species all included very little recognizable courtship, and the few bouts

we observed never progressed to attempted copulations (Fig. 1A,B). Thus, intermale aggressive actions in

the subgroup branch that includes *D. melanogaster* are almost entirely distinct from the courtship that

males perform toward females (Fig. 1E,F, 1-S4).

The other branch of the *melanogaster* subgroup, which includes *D. erecta*, *D. yakuba*, and *D.*

*santomea*, exhibited dramatically different male-male dynamics. *D. erecta* males hardly interacted or

locomoted, even on the preferred *Pandanus* substrate (Fig. 1A,B, 1-S1B, 1-S3) (*57*). *D. yakuba* and *D.*

*santomea,* by contrast, exhibited low levels of aggression and multiple distinct, clearly recognizable

courtship actions (Fig. 1A,B). Male-male courtship in both species was visually similar as toward females

(Fig. 1G,H, 1-S4), but the two sibling species differed markedly from each other in penetrance and

expressivity. Courtship and attack between *D. yakuba* males were both infrequent, whereas *D. santomea*

males courted other males considerably more than they fought them and far more than did *D. yakuba* or

any other subgroup species (Fig. 1B,D, 1-S2D,F). High levels of male-directed courtship and low

aggression were conserved in three additional *D. santomea* strains (Fig. 1-S2C-F), were not limited to the

initial moments of an encounter when sex identification might be incomplete (Fig. 1D) (*58*, *59*), and

sometimes culminated in copulation attempts (Fig. 1A). Audio recordings (*60*) confirmed that *D.*

*santomea* males sang to males as they did to females, emitting both "pulse" and "clack" song (*61–*

*63*) with similar (though not identical) acoustic features (Fig. 1I,J, 1-S5).

Thus, under the conditions tested there exists a near-qualitative difference in intermale social

behavior between *D. melanogaster* and *D. santomea*. *D. melanogaster* and *simulans* clade relatives attack

males with varying frequencies using a conserved set of actions, whereas *D. santomea* courts both males

and females nearly indistinguishably. While widely described in other taxa (*64*), the frequent courtship

among wildtype adult males shown by *D. santomea* in naturalistic settings is, to the best of our

knowledge, novel in the *Drosophila* genus. Importantly, *D. santomea* male-directed courtship is not due

to sex misidentification, because individual flies that courted and sometimes also attacked males

immediately switched to exclusive courtship with significantly increased vigor when encountering a

female, no matter the presentation order (Fig. 1K-O).

**The abundant and monomorphic cuticular pheromone 7-tricosene is sexually attractive to *D.***

***santomea* males**

Five of the nine *melanogaster* subgroup species (*D. simulans*, *D. santomea*, *D. yakuba*, *D.*

*teissieri*, and *D. orena*) are considered pheromonally monomorphic because the same cuticular

hydrocarbon (CHC) is the most abundant compound on the cuticles of females and males (*65–67*). For all

five the defining monomorphism is of (Z)-7-tricosene (7-T) (Fig. 2A, 2-S1), a nonvolatile compound

detected via contact chemosensation (*68*, *69*). *D. santomea* courts in the dark (Fig. 2B) (*70*) and removing

CHCs in bulk from either females or males strongly diminishes their attractiveness (Fig. 2C), implying

the existence of courtship-promoting pheromone(s) shared between sexes. That 7-T is an aphrodisiac

signal for *D. santomea* males is supported by monomolecular add-back following bulk CHC removal

from *D. santomea* females (Fig. 2D), ectopic addition to intact females of species lacking endogenous 7-T

(*D. melanogaster* and others) (Fig. 2E, 2-S2), and positive correlation between 7-T presence and

courtship elicited from *D. santomea* males in a variety of heterospecific male-male pairings (Fig. 2F).

Thus, 7-T appears necessary and is sufficient to evoke sexual attraction towards males as well as females

in *D. santomea.* The inversion from 7-T suppressing courtship in *D. melanogaster* (where it is male-

specific) (*71–74*) to promoting courtship in *D. santomea* is shared with *D. yakuba* (*32*), suggesting that

the 7-T valence switch arose in tandem with monomorphism in the *D. santomea / D. yakuba* (or *yakuba*

clade) common ancestor on mainland Africa (Fig. 2G) (*75*).

**Selectively low cVA abundance on *D. santomea* males amplifies intermale courtship**

If *D. santomea* sexual attraction uses an abundant and monomorphic signal (7-T), then it would

seem unsurprising that male-directed courtship is a corollary outcome of this relatively simple pheromone

arrangement. However, intermale courtship in *D. yakuba* and other 7-T monomorphic species is rare (Fig.

2G) (*30*, *32*, *76*), hinting that *D. santomea* males have either gained an additional courtship-promoting

signal absent in the other species or lost a courtship-inhibiting signal otherwise present. We found that *D.*

*santomea* males express significantly less (Z)-11-octadecenyl acetate (cVA) than other subgroup members

(84-92% reduction, 89% less than *D. yakuba*) (Fig. 3A, 3-S1). cVA is a semi-volatile pheromone

produced in the ejaculatory bulb of male drosophilids (*67*, *77*, *78*) and has pleiotropic roles in social

behavior including promoting aggregation (*79–81*) and aggression (*82–85*) and suppressing courtship

(*86–90*), either natively on males or after transfer to females during copulation. Behavioral dose-response

curves with cVA applied onto a piece of filter paper indicated that cVA indeed suppresses courtship in *D.*

*santomea* (Fig. 3B-D), with the primary effect being to increase latency to initiation (Fig. 3C). In a choice

assay, *D. santomea* males showed nearly universal preference for courting solvent control treated males or

females over flies directly coated with cVA (Fig. 3E). Triggering fictive cVA sensation in *D. santomea*

males by phasic optogenetic activation of Or67d-expressing olfactory sensory neurons (OSNs) (Fig. 3-S2)

(*89*, *91–95*) was sufficient to interrupt natural courtship towards conspecific partners of both sexes (Fig.

3F-I), as also observed in *D. melanogaster* (*96*). Interestingly, the photostimulation light intensity needed

to pause ongoing courtship was lower with male targets than with females (Fig. 3J), consistent with some

courtship suppression even by the low endogenous amount of cVA present on *D. santomea* males.

Unlike its courtship-inhibiting function, cVA's aggression-promoting function does not appear to

be conserved in *D. santomea* males. cVA addition experiments, either onto filter paper or directly onto

live target males, showed no effect of inducing aggression (Fig. 3K,L), nor did strong and direct

photoactivation of Or67d OSNs (Fig. 3M). Pairings with unmanipulated heterospecific male partners

presenting high endogenous cVA levels also failed to significantly increase attack behavior by *D.*

*santomea* males (Fig. 3-S3). The ineffectiveness of cVA to promote aggression in *D. santomea* males may

be due to the 7-T valence switch described above (Fig. 3N-P and see Discussion).

The combination of sexual attraction by monomorphic 7-T, which likely evolved in the *yakuba*

clade common ancestor, and low levels of the male-specific anti-aphrodisiac cVA, uniquely derived in *D.*

*santomea,* thus provides a pheromonal basis for selectively elevated courtship among *D. santomea* males.

One curious additional effect was observed, presumably also due at least in part to low cVA signaling.

When paired with males of other *melanogaster* subgroup species that find cVA sexually aversive (*32*, *67*,

*89*) and show minimal sexual behavior in conspecific male-male pairs (*D. melanogaster*, *D. simulans*, *D.*

*yakuba*), *D. santomea* males elicited 200-580% higher levels of courtship and 53-91% reduced levels of

aggression (Fig. 3-S4) from the heterospecific males. Thus, *D. santomea* male attractiveness extends even

across species boundaries to multiple other *Drosophila* males.

**Central aggression circuits in *D. santomea* can evoke intense attack when photoactivated**

*D. santomea* intermale sexual behavior is frequent but does not fully saturate the time that two

males interact (Fig. 1B). Even so very little attack occurs outside of courtship periods, which in addition

to the ineffectiveness of cVA to promote aggression prompts the question of whether *D. santomea*

aggression circuitry is conserved in form and function.

Multiple aggression-promoting neurons have been identified in *D. melanogaster* males (*50*),

among which are P1/pC1 (and subsets therein) (*56*, *85*, *96–101*) and AIP (also called CL062) (*102*, *103*).

We generated transgenic reagents necessary to access P1/pC1 (referred to here as P1$^{dsx}$ to reflect the split

driver combination used) and AIP in *D. santomea* by combination CRISPR-Cas9 knock-in (*104*, *105*) and

phiC31 recombination into existing landing site lines (*106*, *107*), and tested photoactivation effects in

single or paired flies using CsChrimson as the opsin effector (*108*). AIP neurons had the expected position

and morphology in *D. santomea* males (Fig. 4A), and photoactivation in solitary flies elicited bilateral

wing elevation (i.e. threat display, the major non-contact-mediated form of aggression in male

*Drosophila*) (*41*) time-locked to the onset and offset of photostimulation (Fig. 4B-D, 4-S1). P1$^{dsx}$ neurons

were similarly present (Fig. 4E) and photostimulation evoked reliable time-locked courtship measured by

unilateral wing extension in solitary flies (Fig. 4F, 4-S2), as expected given their dual roles in courtship

and aggression (*109*).

In same-genotype pairs of *D. santomea* males both experiencing titrated photoactivations, P1$^{dsx}$

stimulation induced a mix of aggressive and courtship actions similar to the effects seen in *D.*

*melanogaster* (Fig. 4G-J, 4-S3) (*56*). Lunge attacks, the major contact-mediated aggressive action in

males (*43*), initiated during low frequency stimulations (significantly at 2 Hz) and escalated in intensity as

the stimulus frequency increased (Fig. 4H,I, 4-S3A-C). Fighting remained persistently elevated following

the final photostimulation period for the duration of the recording. High frequency stimulations (>10 Hz)

at one-minute intervals interrupted the otherwise monotonic increase in attack intensity with temporary

phases of courtship and locomotor arrest time-locked to photostimulation (Fig. 4H,J, 4-S3D,E), as also

observed in solitary flies. Mixed-genotype pairs (in which one of the two photoactivated males was

replaced with a wildtype male prepared to be passive during the assay by group-housing) confirmed that

stimulated aggression was fly-autonomous and not exclusively caused by social feedback (Fig. 4-S4A)

and revealed some contextual flexibility in its expression. P1$^{dsx}$ photoactivated flies in mixed pairs

required slightly higher stimulation frequency to initiate attack (5 instead of 2 Hz, Fig. 4-S4A vs. 4I) and

their attack intensity remained ~50-60% lower throughout the trials (Fig. 4-S4B). The delay in attack

latency allowed for an intermediate period when attack and courtship were co-expressed (Fig. 4-S4C,D),

which contrasted with the largely mutually exclusive action patterns observed in same-genotype pairs

controlled by the photostimulation pattern (Fig. 4H). This resulted in a shift from nearly equal balance

between flies exhibiting attack or courtship first to a majority of cases where courtship preceded attack

(Fig. 4-S4E).

Thus, major nodes within at least two distinct aggression-promoting circuits (P1$^{dsx}$ and AIP) are

conserved in *D. santomea* and both contact- and non-contact-mediated attack patterns can be efficiently

evoked by their photoactivation. This rules out that the usual paucity of spontaneous attack in *D.*

*santomea* male pairs under naturalistic conditions derives from an evolved loss of function of (at least

these) aggression circuits, or broadly limiting factors such as energetic or muscular deficits. P1$^{dsx}$

experiments reveal that *D. santomea* aggression can be elevated to supernormal intensity and exhibit

persistence under optimal photoactivation conditions (~23-fold higher lunge rate observed after final

photostimulation compared to spontaneous interactions, Fig. 4I vs. 1C), but the direct effect of courtship

dominates and strongly suppresses attack while P1$^{dsx}$ is highly activated (as also in *D. melanogaster*) (*56*).

An opponent whose pheromone bouquet activates *D. santomea* P1$^{dsx}$, especially a non-aggressive one,

would therefore be expected to elicit courtship instead of attack as long as the activation perdures.

**A switch from cVA sexual attraction to aversion in *D. santomea* females**

*D. santomea* has an interesting natural history (Fig. 5A-C) (*110–112*). They are endemic to the

equatorial West African island of São Tomé, located 280 km from Gabon in the Gulf of Guinea. *D.*

*yakuba* also inhabits São Tomé, and the two sibling species largely segregate by elevation along the

elevation gradient from the volcanic island's warm, dry, and open perimeter (*D. yakuba's* range) to its

cool, humid, forested summit (*D. santomea's*). A sympatric zone is formed at their mid-elevation

boundary, where both species and rare F1 hybrids can be collected that adhere to Haldane's rule for

sterility in the heterogametic sex (i.e., males) (*113*). Since *D. santomea* and *D. yakuba* males court

females of both species promiscuously (*114, 115*), male offspring sterility should pressure females

strongly to discriminate against heterospecific matings and maintain reproductive isolation (*116*). In flies

this challenge would normally be expected to be met by females detecting species-specific acoustic

features of male courtship song (*117*), but although *D. santomea* and *D. yakuba* songs are distinct (*61,*

*63*), *D. santomea* females are still able to distinguish males of each species even if they are muted by

wing amputation (*62*). These observations implicate additional species-identifying cues.

The cVA addition experiments above demonstrated that *D. santomea* males show increased

copulation latency and reduced copulation rates in the presence of high cVA (Fig. 3C,D), but we noticed

that the effects may not derive exclusively from alterations in the males' behavior. *D. santomea* females

exposed to cVA exhibited dramatically increased ovipositor extrusion (Fig. 5D, 5-S1), a sexual rejection

behavior typically seen in recently mated, unreceptive *Drosophila* females (*118–121*). Significant

rejection occurred at a lower cVA dose than was required to reduce male courtship (1 vs. 2 mg, Fig. 5D

vs. 3B), and measurements remained elevated even after normalization to the amount of male courtship

observed (Fig. 5E). Incorporating observations of wing spreading, a distinct female behavior signaling

sexual acceptance (*122*), into a combined index of *D. santomea* receptivity further captured the cVA-

dependent rejection response (Fig. 5F). Strikingly, Or67d OSN silencing via ectopic expression of the

inward rectifying potassium channel Kir2.1 (*123*) in *D. santomea* females paired with wildtype males

resulted in ~2-fold increased copulation rates compared to controls (Fig. 5G), indicating that cVA

sensation normally limits *D. santomea* female sexual responses. Unconditioned place preference tests

suggest that increased cVA aversion is likely innate (Fig. 5-S2).

The sexually appetitive effect of cVA on *D. melanogaster* females (*89*) is therefore reversed in *D.*

*santomea,* and this unexpected switch represents a third pheromone change in *D. santomea* (along with

attraction to 7-T and reduced cVA abundance in males). We hypothesize that the valence reversal may

help *D. santomea* females identify and avoid unwanted hybridization with high cVA-expressing *D.*

*yakuba* males, thereby maintaining reproductive isolation in São Tomé. This may have been preceded by

*D. santomea* male cVA reduction (making cVA-averse *D. santomea* females more sexually competitive),

or alternatively, was directly triggered by *D. yakuba* recolonization of São Tomé (*111*) and caused

compensatory male cVA reduction by sexual selection pressure (*124*) (Fig. 5H, Discussion). The latter

possibility raises a question of whether the increased intermale courtship that results from cVA reduction

is simply a costly but necessary accommodation by *D. santomea* males to appease conspecific females.

According to this view, same-sex sexual behavior (SSB) is inherently "non-reproductive" and may be

detrimental to fitness or, at best, a waste of energy (*64*). We next explored whether male-male courtship

might have any adaptive value in *D. santomea.*

**P1[dsx] activation or natural male-directed courtship induces a competitive mating advantage in *D.***

***santomea* males**

In the original comparative screen and multiple experiments conducted thereafter, paired *D.*

*santomea* males were derived from the same stock and prepared for behavior assays identically, but

nevertheless exhibited a surprising degree of courtship bias between members of the pair. That is, it was

usually the case that just one male committed most of the total courtship observed (Fig. 6A, 6-S1), similar

to biases observed during intermale aggression in *D. melanogaster* (*125*). In pairs where fighting also

took place the two actions were most often attributable to the same fly (Fig. 6B), hinting that biased

courtship and attack might both contribute to agonistic social dominance in *D. santomea.*

*D. santomea* males do not appear to be territorial (Fig. 6-S2), but we reasoned that dominance in

male-directed courtship might facilitate its female-directed counterpart due to persistent social arousal

(*126*), and therefore be a valuable action to the dominant fly. In *D. melanogaster*, pairs of males

encountering one another for the first time ten minutes after P1/pC1 photostimulation will fight

significantly more than controls (*56*). *D. santomea* female- and male-directed courtship uses the

homologous neural circuit (Fig. 6C) and, when tested in a similar assay (called here a "sliding door" for

the removable barrier separating the two flies), brief photostimulation of P1$^{dsx}$ neurons in *D. santomea*

males also triggers persistent, intense aggression after the same ten-minute delay (Fig. 6D-I, 6-S3A-E).

However, in mixed pairs between a photostimulated male and wildtype female, males switched to

exclusive courtship during the interaction phase with significantly increased vigor and decreased latency

compared to unstimulated controls (Fig. 6J-N, 6-S3F), indicating that persistent effects of P1$^{dsx}$ activation

enhance later social interactions with both sexes, and highlighting *D. santomea* males' ability to

discriminate sexes and adjust social behaviors accordingly even when highly aroused.

We recreated P1$^{dsx}$ circuit persistent activation in a naturalistic setting by taking wildtype males

through two consecutive social encounters: first in a pair with one other male during which a courtship

dominance relationship was allowed to form, and second in a trio with the same male and a newly added

female (Fig. 6O-Q). Analogously to P1$^{dsx}$ photostimulation, *D. santomea* males that dominated courtship

in the paired male setting showed significantly greater propensity to also do so toward females in the trio

phase of the experiment (Fig. 6R,S). The effect appeared to strengthen in males that expressed higher

amounts of male-directed courtship during the first phase (Fig. 6R, larger dots). Copulations were rare,

but two of the three observed were obtained by previously dominant males (Fig. 6T) (the exception was

by a male from a pair with very weak intermale dominance). Thus, male-directed courtship in *D.*

*santomea* is sufficient to induce sexual priming, in which males who have recently courted (or otherwise

experienced P1$^{dsx}$ activation) are better prepared to court and potentially also copulate with females that

soon appear.

**Convergent social behaviors and pheromones in *D. persimilis* males**

Our screen also uncovered a second species highly prone to male-directed courtship, *D. persimilis*

(Fig. 1B, 1-S1A). *D. santomea* and *D. persimilis* last shared a common ancestor ~50 million years ago

and intermale courtship appears to be selectively derived in each species (Fig. 7A). *D. persimilis* and *D.*

*santomea* both inhabit cool, high elevation, montane regions, *D. santomea* in West Africa and *D.*

*persimilis* in Western North America, and both are members of a sibling species pair with which they

geographically overlap (*52*). Importantly, males of *D. persimilis* and *D. pseudoobscura* (the sibling

species) both court females interspecifically but females show strong discrimination against cross-

hybridization (*127–129*). Like *D. santomea, D. persimilis* male-directed courtship shares visual

resemblance with the female-directed form and is most often biased to just one male within each pair

(Fig. 7B-D, 7-S1A,C,D). *D. pseudoobscura* males instead show the more genus-typical social patterns of

male-directed aggression and female-directed courtship (Fig. 7E-G, 7-S1B-D) (*130*). Finally, *D.*

*persimilis* males have also converged on the two key pheromonal features of CHC monomorphism and

low cVA expected to reduce or eliminate major barriers to male-male courtship (Fig. 7H,I) (*67, 131*).

Similar phylogenies and natural histories therefore correlate with convergent pheromones and male-male

sexual behaviors shared between these two distantly related flies.

**Discussion**

Evolutionary changes in the expression and central processing of cuticular hydrocarbons (CHCs)

have been shown to underlie phylogenetic variation in pheromonal control of male-female courtship in

*Drosophila* (*22*, *132*). Two types have been described: a change in the sex-specificity of pheromone

expression, from sexually dimorphic to monomorphic CHC patterns (*28*), and a switch in the behavioral

valence of certain CHCs (e.g., 7-T) from male courtship-inhibitory in dimorphic *D. melanogaster* (*71–74*)

to courtship-promoting in monomorphic species including *D. yakuba* (*32*) and, to a lesser extent, *D.*

*simulans* (*27*, *30*, *71*). Here we identify a novel pheromonal valence switch in *D. santomea* females, from

sexual receptivity-promoting (*89*) to receptivity-inhibiting, of cVA, a semi-volatile olfactory pheromone.

Two additional pheromone changes in *D. santomea* males, a reduction in cVA abundance relative to other

*melanogaster* subgroup species and a switch to monomorphism and sexual attraction to 7-T that is shared

with *yakuba* clade species, can explain the amplification of *D. santomea* intermale courtship. The results

illustrate how multiple changes in the behavioral valence or abundance of pheromones in both females

(cVA valence) and males (7-T valence and cVA abundance) can collectively drive evolutionary variation

in same-sex sexual behavior (SSB), as well as aggression. Our data also highlight potential social tensions

between *D. santomea* and *D. yakuba* in their overlapping habitat in West Africa and the evolutionary

impacts affecting both sibling species. A similar set of pheromonal events may also explain the difference

in male-male social behaviors observed between *D. persimilis* and *D. pseudoobscura* in North America.

**A layered evolutionary path to elevated intermale courtship in *D. santomea***

*D. melanogaster* males (*49*, *109*) and males of most *Drosophila* species surveyed here exhibit

primarily female-directed courtship and male-directed aggressive behavior. The sex-specificity of these

social behaviors in dimorphic *D. melanogaster* is controlled in part by male-specific 7-T and cVA, which

function similarly to suppress intermale courtship and promote aggression (*69*, *72*, *82*, *83*). However, *D.*

*santomea* and other *yakuba* clade species are monomorphic for 7-T (*65–67*) and males show 7-T sexual

attraction rather than aversion. This permits these species to pursue females and mate in the dark as

dimorphic species can (but 7-T monomorphic species in which males show neutral 7-T sexual responses

cannot) (*32*), and shifts the function of sex discrimination to other pheromones or sensory modalities.

Despite high 7-T abundance and monomorphism, male-directed sexual behaviors in these species are

usually minimal, except notably in *D. santomea*. The restricted specificity for elevated intermale

courtship in *D. santomea* (and hence evolutionary recency) parallels a selective reduction in the

abundance of cVA, an olfactory signal which in *D. melanogaster* suppresses male courtship (*71*, *86–90*)

and is otherwise present at high levels on males of *melanogaster* subgroup species. That this correlation

reflects causation is demonstrated by the fact that *D. santomea* courtship can be suppressed either by

addition of exogenous cVA or by direct photoactivation of olfactory sensory neurons (OSNs) expressing

the cVA receptor Or67d (*89, 91–95*) . Thus, the high level of sexual behavior among *D. santomea* males

can be explained by two, presumably sequential evolutionary changes in distinct cuticular and olfactory

pheromone signaling pathways: a change in the behavioral valence of 7-T and a reduction in the

abundance of cVA. We note that, while our data indicate that 7-T can restore male courtship towards

hexane-washed dead *D. santomea* females and promote courtship of dead females of other species which

ordinarily do not express 7-T, we cannot exclude that other components of the female CHC bouquet

and/or non-CHC pheromones contribute to *D. santomea* male attraction. We also cannot exclude a limited

role for non-chemosensory signals (*30*).

A potential evolutionary explanation for the male cVA reduction was lacking until our unexpected

discovery of a second sexual valence switch (and third pheromonal change in total), this time in the

behavioral response to cVA by *D. santomea* females. cVA promotes mating receptivity in *D. melanogaster*

(*89*) but has the opposite effect in *D. santomea*: increasing cVA abundance promotes sexual rejection and

Or67d OSN silencing increases copulation rates ~2-fold. The switch may be related to interspecific sexual

dynamics between *D. santomea* and *D. yakuba* at their geographic boundary in São Tomé. *D. yakuba*

males court *D. santomea* females vigorously (*114, 115*) and rare (sterile) F1 hybrid males can be collected

within the sympatric zone (*110–112*), so we speculate that the *D. santomea* female cVA valence switch

arose due to ongoing selection pressure to maintain reproductive isolation from *D. yakuba* (*116*). If the

model is correct, this would mean that *D. santomea* females use the high level of cVA on *D. yakuba* males

as a sex- and species-identifying cue to ward off unwanted advances from heterospecific suitors. Reduced

cVA abundance by *D. santomea* males may then have emerged by consequent sexual selection pressure

(*124*). However, two other evolutionary scenarios are also possible. In one, *D. santomea* males may have

initially reduced cVA abundance due to neutral drift or to pheromonally distinguish themselves from *D.*

*yakuba* males (*133*), which in turn may have pressured *D. santomea* females to develop cVA aversion by

sexual selection operating in the reverse direction. Alternatively, *D. santomea* female cVA aversion and

male cVA reduction may have evolved simultaneously, potentially in genetic linkage (*134*). In any of

these scenarios, the ultimate outcome would have been to maintain efficient intraspecific mating in *D.*

*santomea* but prevent hybridization with *D. yakuba*. The combination of reduced cVA and cuticular

pheromone monomorphism appears to be a generalizable mechanism for amplifying intermale courtship

because the same pattern is shared with *D. persimilis* compared to *D. pseudoobscura*.

Mechanistic explanations for the changes to cVA signaling in *D. santomea* females and males

remain to be elucidated. In *D. melanogaster*, cVA promotes female receptivity through activation of pCd

neurons (*135*). Whether behavioral inversion in *D. santomea* females is reflected in a physiological

inversion of the effect of cVA on pCd, from excitatory to inhibitory, is an interesting possibility. A

challenge also remains in resolving the genetic and biochemical nature of reduced cVA abundance in

males (e.g., decreased production vs. increased degradation), and determining whether the divergent *D.*

*santomea* and *D. persimilis* lineages achieved reductions by similar means. A biosynthetic logic for cVA

production in the ejaculatory bulb has been proposed (*136*) and one elongase identified (*137*), but

additional enzymes necessary to generate cVA from fatty acid precursors have not yet been identified.

*Cyp6a20* has been identified as a potential degradative enzyme for cVA which is expressed by non-

neuronal cells in Or67d-expressing sensillae (*44*, *47*), and may be another target of selection.

**Mechanisms for suppressing aggression in *D. santomea* males**

*D. santomea* males show little spontaneous attack. However, photoactivation of aggression-

promoting neural circuits in the central brain is sufficient to elicit multiple contact- or non-contact-

mediated agonistic actions (e.g., threat displays, lunging, tussling) with supernormal intensity, ruling out a

lack of vigor or selective atrophy of aggression circuits as explanations for the low level of spontaneous

aggressive behavior. Because *D. santomea* males can clearly discriminate conspecific sex, it can be

inferred that pheromones or behavioral cues from a *D. santomea* male are sufficient to signal "maleness,"

but are either insufficient to activate aggression circuits, or that coincident male-derived courtship-

promoting cues (e.g., 7-T) dominantly suppress attack through inhibitory influences on central circuits or

motor systems so long as the target male is present (*109*). A potentially relevant observation is that acute

inhibition of aggression by photoactivation of courtship circuits yields presumptive post-inhibitory

rebound fighting upon stimulation offset, which persists for minutes. This could explain why *D. santomea*

(and *D. melanogaster*) (*56*) males only attack each other during the light-off periods of our P1$^{dsx}$

photoactivation experiments when courtship subsides, and initiate aggression almost instantly minutes

after P1$^{dsx}$ neurons have been stimulated in the sliding door assay, when aggression circuits may still be in

the elevated rebound regime. Persistent neural effects must be flexible, however, because females are

instead met with vigorous and exclusive courtship following barrier removal.

In addition to the courtship-inhibiting effect discussed above, cVA also promotes aggression

among *D. melanogaster* males (*82–85*). At face value, this second signaling function predicts that reduced

cVA in *D. santomea* should have a corollary effect to decrease aggression, in which case cVA reduction

alone could have parsimoniously explained both the elevated courtship and reduced fighting observed in

*D. santomea* social interactions. We have so far been unable to verify this prediction, because various

experimental efforts to elevate cVA, including strong and direct photoactivation of Or67d OSNs, have

been ineffective at increasing *D. santomea* aggression (although effective at suppressing intermale

courtship). A possible explanation is suggested by the established hierarchical control of male-male social

behavior by 7-T and cVA in *D. melanogaster* (*72*). There, cVA can promote fighting but only if 7-T

signaling remains intact. In flies lacking 7-T or its receptor Gr32a, cVA has no effect to promote

aggression or to inhibit the extensive male-male courtship unmasked by genetic disruption of 7-T

signaling. Thus, suppression of intermale courtship by 7-T is likely a prerequisite for the aggression-

promoting effect of cVA in *D. melanogaster*. By extension, since in *D. santomea* 7-T no longer

suppresses courtship, according to this hierarchical model increasing cVA experimentally should be

unable to promote fighting, as is observed. This hypothesis does not exclude the possibility that thus-far

undetected evolutionary changes in aggression-promoting components of cVA central circuitry (*84*,

*126*) may contribute to reduced fighting in *D. santomea* males as well.

**Recent speciation and intermale sexual behavior**

It seems unlikely to be a coincidence that the two cases of highly expressive and penetrant same-

sex sexual behavior (SSB) discovered here in *Drosophila* are each observed in members of sibling species

pairs, and particularly in species with the more restricted geographic range. The pattern suggests that

recent speciation may predispose animals to elevated SSB expression (or, formally, high SSB may

predispose to speciation), and in fact this correlation proved significant in a phylogenetic analysis of SSB

incidence across all mammals, including the Hominidae (*138*). The findings here provide a general model

that may explain at least some of these cases: where sympatry necessitates enforcement of reproductive

isolation between interfertile species that produce unfit hybrid offspring (such as sterile F1 males) (*113*),

the challenge of interspecies identification may draw upon sensory cues that, once altered, also result in

modification of intraspecific social behaviors including SSB. If the condition of sympatry is removed, or

additional species-identifying cues are enlisted, or compensatory intraspecific mechanisms evolve, then

SSB may relax, unless it provides an adaptive value.

Such an adaptive value for SSB may have accrued in *D. santomea* for either of two reasons, not

mutually exclusive. Males performing natural courtship during male-male encounters (or artificial

courtship during direct P1$^{dsx}$ photostimulation) also initiate courtship more quickly and sustain it more

vigorously toward a female if she appears within a few minutes. Males could therefore benefit from this

sexual facilitation or "priming" effect. Alternatively, the strong bias observed in dyadic male-male

courtship interactions, and the fact that the same male that courts is also the one more likely to attack,

suggests that intermale courtship could have an adaptive function in mediating social dominance (*64*).

Interestingly, evidence in mice suggests that most episodes of male SSB in that species reflect dominance

mounting, a low-intensity form of aggressive behavior, rather than affiliative sexual behavior (*139*), and

SSB as a dominance behavior has been proposed in other taxa (*140*, *141*).

The general relationship between SSB and aggression across animals appears more complex. The

survey study in mammals (*138*) found a positive correlation between SSB and lethal male violence,

motivating the hypothesis that SSB may evolve to ease escalating intermale aggression. If this alternative

model also pertains to *Drosophila*, then the sequence and causal role of evolutionary events identified

here could be reinterpreted: rather than the key benefit of pheromone (especially cVA) changes being to

avoid cross-species hybridization, it may instead have been to effectively replace intermale aggression

with courtship, a form of social interaction carrying reduced risk of harm to individuals (*64*). In other

words, *D. santomea* males may have evolved a social strategy in which the ultimate functions of

aggression that improve fitness (e.g., securing resources, mates, etc.) can be achieved by increased male-

male courtship instead, avoiding aggression's injurious costs but maintaining its agonistic or competitive

intent (*142*). Resolving or harmonizing these alternatives, potentially by modeling or controlled selection

experiments, should help to better understand the evolutionary origins of social behavior across the

animal kingdom perhaps including in the human species.

**Main Figure Legends**

**Figure 1. Evolutionary variation in male-male social behavior within the genus *Drosophila***

**(A)** Ethograms from twenty-minute male-male interactions depicting aggressive and courtship actions scored in three representative conspecific pairs per species across the melanogaster subgroup. Nodes are bout counts of a given action type, with sizes representing frequency normalized to summed bouts scored across all actions (expert annotation). Edges indicate action transitions, with weights representing the fraction of bouts of a given action (arrow origin) for which a second action (arrow destination) occurred within one second. Eleven total actions in each ethogram are arranged in four rows: first, three aggressive threat actions (threat, pump, charge) (*102*); second, three aggressive contact-mediated actions (lunge, hold, tussle) (*48*); third, three courtship actions utilizing the wings and typically directed toward females (UWE, BWE, row) (*20*); fourth, two courtship actions involving locomotory maneuvers (circle, copulation attempt). Nodes indicating aggressive actions are filled with blue shades and courtship actions with reds. Species arrangement and divergence times according to compilation of available phylogenies (*75*, *143*, *144*). Note emergence of male-directed courtship and reduction of aggression in *D. santomea* (orange) and, to a lesser extent, its sympatric sister species *D. yakuba* (green). *D. erecta* ethogram gray coloration indicates that very few social actions of any kind were observed. Abbreviations: UWE, unilateral wing extension; BWE, bilateral wing extension; Cop. att., copulation attempt. See Table S2 for wildtype strains used in figures throughout.

**(B)** Stacked barplots showing cumulative time spent by paired males in any aggressive (blue) or courtship action (red) as a fraction of the twenty-minute total interaction. Additional Sophophora subgenus species for which male-male interactions were scored for the same action set are included. Note unusually high time dedication to male-directed courtship by *D. santomea. D. serrata* aggression scoring is inflated due to unusually long holds.

**(C,D)** Expressivity and temporal dynamics of male-male aggression (C) and courtship (D) in *D. melanogaster* (black), *D. simulans* (gray), *D. santomea* (orange), and *D. yakuba* (green). Bold traces indicate the mean fraction of frames annotated as lunge (for aggression, C) or UWE (for courtship, D) by automated classifiers in a 10-second sliding window. Envelopes represent standard error of the mean (s.e.m.). Note high aggression in *D. melanogaster* male-male pairs and high courtship in *D. santomea* throughout twenty-minute interactions (significance by Kolmogorov-Smirnov tests). See Supplemental Table S4 for statistics related to figures throughout.

**(E-H)** Video stills of *D. melanogaster* (E,F) and *D. santomea* males (G,H) interacting with conspecific female (E,G) or male targets (F,H). *D. melanogaster* court females (E, male shown in UWE) and attack

males (F, shown mid-lunge). *D. santomea* court both sexes using a similar set of motor patterns (G,H,

males shown in UWE for both) and rarely attack. Scale bars, 1 mm.

**(I,J)** Representative *D. santomea* courtship song traces showing "pulse" and "clack" emitted toward both

females (I) and males (J).

**(K)** Scheme for assessing social behavioral differences by *D. santomea* male tester flies paired

consecutively with a male then female conspecific target, or vice versa. Tester male experiences twenty

minutes with each sex.

**(L-O)** Aggression (L,N, measured as lunges per minute) and courtship (M,O, fraction of total interaction

time performing UWE) exhibited by *D. santomea* male testers towards female and male targets. (L,M)

Males presented first. (N,O) Females presented first. Data points derived from the same tester are paired

and distribution means shown as paired black dots. Note aggression towards males is never also observed

to females and significantly increased courtship rates to females, independent of the presentation order

(paired Mann-Whitney *U* tests).

**Figure 2. The abundant and monomorphic pheromone 7-tricosene is sexually attractive to *D.***

***santomea* males**

**(A)** Mirrored gas chromatograms representing male (purple, top) and female (pink, bottom) *D. santomea*

cuticular hydrocarbon (CHC) profiles. Individual measurements made on hexane extracts from ten

pooled, age-matched adult flies (3 male, 2 female replicates). Peaks representing the internal standard

(C18, 1), male-specific (Z)-11-octadecenyl acetate (cVA, 2), and monomorphic (Z)-7-tricosene (7-T, 3)

are indicated.

**(B)** Fraction of time *D. santomea* males spend courting conspecific females or males in ambient light vs.

dark during ten-minute interactions. Boxplots show full distribution range within whiskers (excluding

statistically identified outliers) and second and third quartiles within boxes, with medians in bold.

Individual data points (including outliers) overlayed as gray or, for males courting females that

successfully copulated, red dots. Note slight but non-significant decreases in courtship toward both sexes

in the dark (Mann-Whitney *U* tests), though a significant (binomial test) decrease in copulation rate with

females.

**(C)** Fraction of time *D. santomea* males spend courting conspecific dead females or males with or without

endogenous CHCs (removed by hexane washes) during ten-minute interactions. Significantly reduced

courtship without CHCs for both target sexes (Mann-Whitney *U* tests).

**(D)** Similar to (C), with 10% 7-T add-back (or hexane solvent control) to conspecific dead females after

CHC removal. 7-T treatment partially restores attractiveness to *D. santomea* male testers (lettered

statistical groupings assigned by post-hoc Dunn's tests following significant Kruskal-Wallis). Cohen's *d*

effect size for 10% 7-T addition (fourth) vs. no addition after CHC wash (second), 0.692 (95% CI -0.379-

0.821, p=0.401). Cohen's *d* effect size for solvent addition (third) vs. no addition after CHC wash

(second), 0.221 (95% CI 0.076-1.31, p=0.042). Cohen's *d* effect size for 10% 7-T addition (fourth) vs.

solvent addition (third), 0.617 (95% CI 0.069-1.17, p=0.039).

**(E)** Fraction of time *D. santomea* males spend courting dead *D. melanogaster* females (endogenous 7-T

low) perfumed with 10% 7-T or solvent control (hexane). 7-T addition elicits courtship from *D. santomea*

males. Significance by Dunn's tests.

**(F)** Fraction of time *D. santomea* males spend courting conspecific (orange) or heterospecific males (*D.*

*yakuba* green, all others black) during twenty-minute interactions. 7-T presence in each species from GC-

MS and phylogeny shown below, statistical groupings by Dunn's tests above. Note correlation between 7-

T presence and elevated *D. santomea* courtship.

**(G)** Summarized indications of 7-T monomorphism (*65*, *66*), evidence for 7-T as a sexual attractant (*32*,

*71*, *72*), and binarized male-male courtship expression in *melanogaster* subgroup species arranged by

phylogeny. "(–)" indicates dimorphic species for which 7-T is not expected to be sexually attractive but

has not yet been tested. Pheromone status of common ancestors at phylogeny branchpoints inferred by

parsimony.

**Figure 3. Selectively low cVA abundance on *D. santomea* males amplifies intermale courtship**

**(A)** cVA abundance on males of the *melanogaster* subgroup measured in single flies (5-8 per species) by

TD-GC-MS. Statistical groupings by Dunn's tests; "a.u." arbitrary units. Note low cVA level selectively

in *D. santomea* males. Data derived from (*67*) with permission from the authors.

**(B-D)** *D. santomea* sexual behavior during twenty-minute interactions with females in the presence of

increasing cVA on a piece of filter paper nearby. (B) Fraction of time spent courting. (C) Latency to first

courtship bout. (D) Latency to copulation. Pairwise statistical significance by Dunn's tests, binomial tests

for copulation rates. Note decreased courtship, increased latency, and decreased copulation rate at the

highest cVA abundance (2 mg).

**(E)** Preference index for *D. santomea* courtship towards 10% cVA vs. solvent control (acetone) perfumed

live target female or male conspecific flies during twenty-minute trio interactions. Index calculated as the

difference in time spent courting each target normalized by the sum. Significance between distributions

and between each distribution and a null median of zero (indicating no preference) determined by two-

and one-sample Mann-Whitney *U* tests, respectively. Note that *D. santomea* male testers exhibit clear

aversion to cVA-perfurmed targets of either sex. All three observed copulations occurred with the solvent

control female.

**(F)** Photomicrograph showing labeling pattern of $Or67d^{Gal4}$ in the antennal lobes (AL) of a *D. santomea*

male. Immunostaining for Gal4-dependent cytoplasmic tdTomato pseudocolored green with Bruchpilot

synaptic counterstain (using nc82 monoclonal antibody) in magenta. Uniglomerular labeling pattern in

each AL matches the expected size and position of cVA-sensitive DA1 (*145*). Scale bar, 25 μm. See Table

S3 for genotypes of transgenic strains used in figures throughout.

**(G)** Courtship rasters (red) for a single-housed *D. santomea* male carrying $Or67d^{Gal4}$ and Gal4-dependent

CsChrimson paired with a group-housed wildtype conspecific male target. Two representative sham trials

(above dash) and two photostimulation (PS) trials (below dash) are shown. Two-second PS blocks repeat

every ten seconds for a total of six activations after a one-minute baseline. PS frequencies increase in

pairs from 10 to 40 Hz within each trial and increase in intensity across trials. Note that PS immediately

interrupts spontaneous courtship by tester males in experimental but not sham trials.

**(H,I)** Quantification of (G) showing fraction of $Or67d^{Gal4}$ tester males courting males (H) or females (I)

during six-second windows surrounding PS blocks (H, purple; I, pink) or sham controls (black). Data

filtered for blocks where tester males showed spontaneous courtship within a half second of PS onset.

Data from all three frequencies were grouped since they gave similar results but separated by intensity:

120 (H) or 260 μW/mm$^2$ (I). Most tester males cease courtship by PS offset in experimental trials

(significant for both target sexes by Kolmogorov-Smirnov tests).

**(J)** Summary of courtship suppression effects in $Or67d^{Gal4}$ tester males as a function of PS intensity with

male (purple) and female (pink) targets. Baselines (horizontal dashed lines) calculated as the fraction of

spontaneous courtship bouts in sham trials lasting more than two seconds. Note strong effects with male

targets at low intensities but only at the strongest intensity with females.

**(K)** *D. santomea* male aggression during twenty-minute interactions between single-housed males in the

presence of increasing cVA on filter paper. Not significant by Kruskal-Wallis test.

**(L)** *D. santomea* male aggression towards males directly perfumed with 10% cVA or solvent control

(acetone) during twenty-minute trio interactions. Not significant by Mann-Whitney *U* test.

**(M)** Aggression in *D. santomea* males carrying $Or67d^{Gal4}$ and Gal4-dependent CsChrimson or genetic

controls during twelve-minute photoactivation trials. Pairs of group-housed males are exposed to six

thirty-second photostimulation (PS) blocks with fixed intensity and monotonically increasing frequency

(1-40 Hz), with a two-minute baseline and one-minute inter-block intervals. Measurements represent

cumulative lunge counts from all six PS blocks. Note insufficiency of $Or67d^{Gal4}$ activation to induce

attack (no significance between genotypes by Kruskal-Wallis test).

**(N-P)** Logic models for control of male social behaviors by 7-T and cVA in *D. melanogaster* (7-T

dimorphic, N), *D. yakuba* (7-T monomorphic, O), and *D. santomea* (7-T monomorphic with reduced cVA,

P). In *D. melanogaster* where 7-T inhibits intermale courtship, 7-T also gates the effect of cVA to promote

aggression (*72*). A switch in 7-T function from suppressing to stimulating courtship in monomorphic

species is predicted to reduce or eliminate the aggression-promoting effect of cVA without impacting its

courtship-suppressing effect, as evident in *D. santomea.* Intermale courtship promoted by 7-T in *D.*

*santomea* is further amplified due to reduced inhibition by cVA. Edge terminating in white box indicates a

permissive "gating" effect by 7-T on cVA-induced aggression, in addition to its direct effect.

**Figure 4. Central aggression circuits in *D. santomea* can evoke intense attack when photoactivated**

**(A)** Photomicrograph showing labeling pattern of *20E08^Gal4* in the central brain of a *D. santomea* male. Immunostaining for Gal4-dependent cytoplasmic tdTomato (green) with synaptic counterstain (nc82, magenta). Close-up (A') shows AIP somata and axons (arrowhead) from boxed region. Scale bars, 50 µm (A), 10 µm (A').

**(B)** Video still of bilateral wing elevation (threat) elicited by PS in a solitary *D. santomea* male carrying *20E08^Gal4* and Gal4-dependent CsChrimson. Scale bar, 1 mm.

**(C,D)** Quantification of PS effects in solitary *20E08^Gal4* tester males (purple) and genetic controls (black) exposed to two fifteen-second PS blocks with thirty-second baseline and inter-block interval. (C) Fraction of flies showing threat (penetrance). (D) Mean fraction of frames with flies showing threat in a five-second sliding window with s.e.m. envelopes (expressivity). Significant induction of threat during PS in *20E08^Gal4* by Mann-Whitney *U* tests.

**(E)** Photomicrograph showing *P1^dsx* split-Gal4 (*71G01^DBD;dsx^AD*) labeling in the central brain of a *D. santomea* male. Immunostaining for Gal4-dependent cytoplasmic tdTomato (green) with synaptic counterstain (nc82, magenta). Close-ups show a characteristic region of P1^dsx neuropil (lateral junction, E') and somata and axons (E'') from boxed regions. Scale bars, 50 µm (F), 10 µm (F',F'').

**(F,G)** Courtship elicited in solitary (F) and aggression in paired (G) *D. santomea* males carrying *P1^dsx* split-Gal4 and Gal4-dependent CsChrimson or a genetic control during twelve-minute photoactivation trials. In both cases group-housed males are exposed to six thirty-second PS blocks with fixed intensity and monotonically increasing frequency (1-40 Hz), with a two-minute baseline and one-minute inter-block intervals (as in H). "PS–" represents the two-minute baseline period prior to first PS. "PS+" cumulatively represents all six PS blocks (for courtship, G) or all five inter-block intervals (IBIs) and period following final PS (for aggression, F). See behavior dynamics in (H) for justification of aggression quantification during IBIs. Note significant PS induction for both actions (Dunn's tests).

**(H)** Fraction of flies showing attack (lunge, blue) and courtship (UWE, red) in same-genotype pairs of *P1^dsx* males during six PS blocks with fixed intensity and increasing frequencies. Note highly penetrant attack mostly during light-off periods punctuated by time-locked courtship during high frequency PS blocks. Video stills at right show aggressive tussling following 2 Hz PS (1) and simultaneous courtship by both tester males during 10 Hz PS (2). Scale bars, 1 mm.

**(I,J)** Expressivity and temporal dynamics of attack (I) and courtship (J) elicited by PS in *P1^dsx* male pairs (I, blue; J, red) or a genetic control (black). Significant attack induction during inter-block intervals and after final PS (I) or courtship during PS compared to control (J) by Mann-Whitney *U* tests.

**Figure 5. A switch from cVA sexual attraction to aversion in *D. santomea* females**

**(A)** Elevation map of equatorial Central and West Africa with the volcanic island of São Tomé in the Gulf of Guinea boxed. Data from (*146*) and Terrain Tiles (Mapzen, accessed Aug. 24, 2025).

**(B)** Elevation map of São Tomé. Note the central summit of Pico de São Tomé.

**(C)** Latitudinal elevation profile of Pico de São Tomé showing *D. santomea* (orange) and *D. yakuba* zones (green) separated by a hybrid zone where both pure species and F1 hybrid males are found (*112*).

**(D,E)** Behavioral dose-response curves of *D. santomea* female ovipositor extrusion (OE) to increasing cVA on filter paper during conspecific male courtship. (D) Female OE as fraction of total observation time. (E) Female OE normalized to male courtship (difference divided by sum of time spent by female in OE and time spent by male in UWE). Red dots, females that copulated. The lowest cVA dose induces an intermediate effect while both higher doses induce significant increases in OE expression (Dunn's tests).

**(F)** *D. santomea* female "receptivity index" as function of increasing cVA. For each female an index value of 1, 0, or -1 is possible. Observations of female wing spreading (WS, an acceptance behavior) and courtship-normalized OE (rejection, from E) were each binarized as present or absent (WS>0, OE(norm)>-0.25) and overall index per dose is calculated as the difference between the two binary values (WS-OE) normalized by the number of females scored. Note significantly decreased sexual acceptance again at the two higher cVA doses (Dunn's tests).

**(G)** Cumulative copulation latencies and rates by *D. santomea* tester females carrying *Or67d^{Gal4}* and Gal4-dependent Kir2.1 (black) or genetic controls (grays) grouped with an equal number of wildtype conspecific males for 45 minutes in large chambers. "BDP" refers to a Gal4 driver without labeling evident in the central brain. 24 total females tested of each genotype (two trials each of 12 single-housed males x 12 group-housed females), significance by Kolmogorov-Smirnov tests comparing latencies or binomial tests comparing rates.

**(H)** Evolutionary model for the emergence of high intermale courtship and low intermale aggression in *D. santomea,* incorporating three key pheromone changes inferred to occur prior to or following *D. santomea* speciation in São Tomé. A common sexual attractant for *D. santomea* and *D. yakuba* males (7-T) generates the behavioral potential for unfit hybridization in sympatry due to F1 hybrid male sterility (*110*). *D. santomea* females evolve sexual aversion to cVA and *D. santomea* males reduce cVA, thereby maintaining efficient intraspecific mating while also maintaining reproductive isolation from *D. yakuba* (the two events cannot be definitively ordered). Reduced male cVA, together with monomorphic 7-T, releases high intermale courtship. Reduced efficacy of cVA to promote aggression further suppresses fighting. Placement of *D. yakuba* secondary colonization after allopatric *D. santomea* speciation (instead of speciation in sympatry) from (*111*).

**Figure 6. P1<sup>dsx</sup> activation or natural male-directed courtship induces a competitive mating advantage in *D. santomea* males**

**(A)** Spontaneous bias observed between flies in male-male *D. santomea* and *D. melanogaster* pairs for attack and courtship. Indices derived from the same fly pair are connected as lines and distribution means shown as paired orange (*D. santomea)* or black dots (*D. melanogaster*). Note high courtship bias in *D. santomea.*

**(B)** Fraction of *D. santomea* male-male pairs with high attack bias where the same fly also showed majority of courtship (left), or with high courtship bias where the same fly also showed majority of attack (right). Significance compared to random chance (50%) by binomial tests.

**(C)** Courtship in the dark by single-housed *D. santomea* males carrying *P1<sup>dsx</sup>* split-Gal4 and Gal4-dependent Kir2.1 or a genetic control paired with group-housed wildtype conspecific female (left) or male targets (right). Red dots, males that copulated (only observed in controls). Significant courtship reduction in *P1<sup>dsx</sup>* tester males toward both sexes by Mann-Whitney *U* tests.

**(D)** Scheme for assessing persistent behavioral effects of *D. santomea P1<sup>dsx</sup>* split-Gal4 activation. After a one-minute baseline, isolated *P1<sup>dsx</sup>* tester males or genetic controls experience one-minute PS and a ten-minute delay. Dividers ("sliding doors") are then removed to allow paired flies (either two same-genotype males or a tester male and wildtype female) to interact spontaneously for an additional ten minutes.

**(E)** Behavior rasters for same-genotype *P1<sup>dsx</sup>* male pairs (both experienced PS before the delay) during the ten-minute interaction phase. Data from just one male per pair is shown and quantified in (F-I) for fair comparison to interactions with females below. Blue, attack (lunge); red, courtship (UWE).

**(F-I)** Quantification of (E) showing expressivity and temporal dynamics and of male-male aggression (F,H) and courtship (G,I) during the interaction phase. Note highly increased aggression and slightly increased courtship in *P1<sup>dsx</sup>* compared to genetic controls without PS (significance by Kolmogorov-Smirnov (F,G) or Mann-Whitney *U* tests (H,I)).

**(J-N)** Behavior rasters (J), temporal dynamics, and expressivity of courtship (K,M,N) and aggression (L) between *P1<sup>dsx</sup>* males or genetic controls paired with wildtype females during the interaction phase. Note highly increased courtship expression (M) and decreased courtship latency (N) in *P1<sup>dsx</sup>* without any incidence of attack.

**(O)** Scheme for determining a sexual priming effect of male-directed courtship. Two single-housed wildtype *D. santomea* males are paired for ten minutes before a wildtype female is added for an additional ten minutes (trio period). Male-directed courtship dominance is measured and identity of the dominant courter ascertained in the paired period then compared to female-directed courtship dominance in the trio.

**(P,Q)** Video stills of a male courting another male first in the paired period (P) then courting a female in the trio (Q). Scale bars, 1 mm.

**(R)** Quantification of the sexual priming effect (O-Q). Female-directed courtship dominance in trios (y-
axis) plotted as a function of male-directed courtship dominance in pairs (x-axis). Points above the x-axis
represent cases where the same male exhibited male-directed courtship dominance in pairs and female-
directed courtship dominance in trios. Dot size represents total amount of male-directed courtship in pairs
and data are filtered for pairs with at least twenty seconds of courtship during this phase (no matter the
bias). Red dots, males that copulated. Gray box, males with high levels of courtship dominance during the
paired period. Note enrichment of points above the x-axis in cases of high male-male courtship
dominance.
**(S,T)** Categorization of effects on female-directed courtship dominance (S) and copulation advantage (T)
from (R). (S) All but one male exhibiting male-directed courtship dominance in pairs (89%) also
exhibited female-directed courtship dominance in trios (significant by binomial test). (T) Two out of three
copulations observed were by the male exhibiting male-directed courtship dominance in pairs (circled red
dots in R). The exception was from a "submissive" male in which courtship bias was very weak
(uncircled red dot at bottom left in R).
698

**Figure 7. Convergent social behaviors and pheromones of *D. santomea* and *D. persimilis* males**

**(A)** Summarized indications of male-male courtship presence/absence in *Drosophila* species arranged by

phylogeny. Note sibling species *D. persimilis* (purple) and *D. pseudoobscura* (blue) in the obscura group.

**(B,C)** Video stills of *D. persimilis* males promiscuously courting conspecific female (B) and male targets

(C). Images selected to represent species-typical action selection. Scale bars, 1 mm.

**(D)** Action biases in *D. persimilis* male-male pairs. Male-directed courtship (red dots) is more common

than attack (blue dots) and often shows strong bias to one fly (boxed region). Attack and courtship

measurements from the same pair connected by thin lines. Circled "X" indicates mean value for the

corresponding action.

**(E-G)** Video stills (E,F) and action biases (G) for *D. pseudoobscura* as for *D. per*. Note female-directed

courtship (E) and biased male-directed attack (F,G). Scale bars, 1 mm.

**(H)** cVA abundance on males of the *melanogaster* subgroup and *D. persimilis* / *D. pseudoobscura* by TD-

GC-MS (*67*). Statistical groupings by Dunn's tests; "a.u." arbitrary units. Note low cVA levels in *D.*

*persimilis* and high in *D. pseudoobscura* as in *D. santomea* / *D. yakuba*.

**(I)** Summary of relevant pheromones (*67*, *76*, *131*) and social behaviors for the *melanogaster* subgroup

(excluding *D. teissieri* and *D. orena*) and the *D. persimilis* / *D. pseudoobscura* sibling species pair. CHC

monomorphism is common to multiple species but low cVA abundance and high male-male courtship are

shared only between *D. santomea* (orange) and *D. persimilis* (purple). Information compiled from the

current study and prior sources.

**Supplemental Figure Legends**

**Figure 1-S1. Additional behavior ethograms and locomotor analyses of *Drosophila* intermale social**

**interactions**

**(A)** Ethograms from twenty-minute interactions depicting aggressive and courtship actions scored in three

representative conspecific male-male pairs per species across various *Sophophora* and *Drosophila*. Nodes

are bout counts of a given action type, with sizes representing frequency normalized to summed bouts

scored across all actions (expert annotation). Edges indicate action transitions, with weights representing

the fraction of bouts of a given action (arrow origin) for which a second action (arrow destination)

occurred within one second. Nine total actions in each ethogram are arranged in five rows: first, two

aggressive threat actions (threat, pump) (*102*); second, two aggressive contact-mediated actions (lunge,

hold) (*48*); third, two additional contact-mediated aggressive actions (headbutt, barrage) (*147*); fourth,

two courtship actions utilizing the wings and typically directed toward females (UWE, BWE) (*20*); fifth,

copulation attempt. Nodes indicating aggressive actions are filled with blue shades and courtship actions

with reds, except for headbutt (white) and barrage (black). Species arrangement and divergence times

according to compilation of available phylogenies (*75*, *143*, *144*). Gray coloration indicates that very few

social actions of any kind were observed. Abbreviations: UWE, unilateral wing extension; BWE, bilateral

wing extension; Cop. att., copulation attempt. Barrage described previously in *D. pseudoobscura*

(*130*) but named here.

**(B-E)** Locomotor summaries for all fifteen species included in the behavioral screen. Per-frame

calculations of each fly's forward velocity (B), angular velocity (C), facing angle relative to the partner

(D), and inter-fly distance (E) are derived from automated tracking of both flies (*55*) in all recorded male-

male pairs. Distributions shown as violin plots with dotted means. *D. santomea,* orange; *D. yakuba*,

green.

**Figure 1-S2. Performance and usage of automated behavior classifiers for intermale social**

**interactions within the *melanogaster* subgroup and additional *D. santomea* strains**

**(A,B)** Automated behavior classifier performance for lunge (A) and unilateral wing extension (B). Three

twenty-minute representative recordings of male-male pairs for each species were manually scored for

both actions (expert annotation, same pairs as in Fig. 1A) and exposed to automated behavior classifiers

trained previously for *D. melanogaster* (*56*), implemented in JAABA (*54*). Manual (x-axis) vs. classifier-

generated annotations (y-axis) are compared in $\log_{10}$ scale, using either the number of bouts (lunge) or

their cumulative time duration (UWE). Solid lines, diagonals; dashed lines, linear fits. Adjusted $R^2$,

significance by F-tests for fits, and Pearson correlations (r) indicated at bottom right. *D. melanogaster*,

black; *D. santomea,* orange; *D. yakuba*, green; *D. simulans*, *D. mau*, *D. sec*, *D. erecta*, gray. Note near-

perfect overlap with the diagonal for lunge and strong correlation for UWE despite classifier false

positives in cases with few manual annotations (all subgroup species except *D. santomea* show very little

male-male courtship).

**(C,D)** Aggression (lunge, C) and courtship (UWE, D) measurements from automated classifiers in all

recorded male-male pairs for *melanogaster* subgroup species and additional *D. santomea* strains. *D.*

*santomea* 00 (also called STO.4) is the strain included in the behavioral screen and used throughout as

"wildtype" (National Drosophila Species Stock Center 14021.0271.00). Strains 146 (STO-CAGO 1482),

148 (STO.7), and 151 (STO.6) also derive from São Tomé from females collected between 1100 and 1500

m elevation (*111*). Boxplots show full distribution range within whiskers (excluding statistically identified

outliers) and second and third quartiles within boxes, with medians in bold. Individual data points

(including outliers) overlayed as gray dots. Lettered statistical groupings assigned by post-hoc Dunn's

tests following significant Kruskal-Wallis. Note low aggression and high courtship conserved in *D.*

*santomea* strains (orange), low aggression and courtship in the sibling species *D. yakuba* (green), and

high aggression and low courtship in *D. melanogaster* (black).

**(E,F)** Summaries of penetrance (x-axes) and expressivity (y-axes) for aggression (E) and courtship (F)

across subgroup species and *D. santomea* strains. Metrics derived from data in (C,D).

**Figure 1-S3. Intermale aggression and courtship in *melanogaster* subgroup species with host**

**specializations**

**(A,B)** Aggression (A) and courtship (B) in *D. melanogaster*, *D. sechellia*, *D. santomea* (orange), and *D.*

*erecta* male-male pairs competing over the standard apple juice substrate and other fruits based on

reported specializations (*53*). Significance for each fruit/species combination by pairwise Mann-Whitney

*U* tests to apple. Removing food source entirely significantly reduces aggression in *D. melanogaster* (*59*).

Using morinda (also called noni) significantly increases aggression but not courtship in *D. sec*. *D.*

*santomea* aggression and courtship depend only weakly if at all on food presence and source.

**Figure 1-S4. Male- and female-directed social behaviors by *D. melanogaster*, *D. simulans*, *D.***

***santomea,* and *D. yakuba* males**

**(A,B)** Ethograms representing male-male (A) and male-female (B) social behaviors for *melanogaster*

subgroup species of interest. Phylogeny also includes *D. mauritiana* and *D. sechellia* from the *simulans*

clade, *D. teissieri* from the *yakuba* clade, and *D. erecta* without associated ethograms. Female-directed

ethograms represent expert annotation for three recordings of male-female pairs per species, with male

flies similarly prepared by single-housing and interactions taking place under identical conditions. Male-

directed ethograms reproduced from Fig. 1A for visual comparison to female-directed counterparts. Note

nearly exclusive male-directed aggressive actions (blue nodes) by *D. melanogaster* and *D. simulans* males

but prevalence of male-directed courtship actions (red nodes) by *D. santomea* and *D. yakuba* (though in

*D. yakuba* total male-direcetd actions are many fewer on average than *D. santomea)*, matching the actions

taken when paired with females.

**Figure 1-S5. Acoustic features of female- and male-directed courtship song by *D. santomea* males**

**(A,B)** Representative courtship song traces by *D. santomea* males toward a female (A, pink) or male

conspecific target fly (B, purple). Key acoustic features are indicated including pulse, pulse train, inter-

pulse interval (IPI), clack, clack train, and inter-clack interval (ICI) (*63*). Traces reproduced and enlarged

from Fig. 1I,J.

**(C,D)** Video stills of song recording chambers (*60*) containing *D. santomea* males courting either a

female (C) or male (D). Isolated waveforms of single clacks and pulse trios shown below, with asterisks

indicating event peaks. Scale bars, 1 mm.

**(E-J)** Quantitative acoustic analyses of female- (pink) and male-directed *D. santomea* courtship songs

(purple). Clacks (E-G) and pulses (H-J) were identified by expert annotation from three recordings with

each sex (clack, 32-103 events per recording; pulse, 140-346 events per recording). Data shown as

"hemiviolin" plots in which female- and male-directed distributions are vertically mirrored. Distribution

means are semicircles on the appropriate side. Note general similarity between female- and male-directed

song features (significant differences based on Mann-Whitney *U* tests despite small difference

magnitudes are often because so many data points make up the underlying distributions). Analyses of

absolute event abundances were omitted since annotations captured only a fraction of events in each

recording.

**Figure 2-S1. GC-MS profiling of *D. santomea* male and female cuticular hydrocarbons (CHCs)**

**(A,B)** Absolute (A) and relative abundances (B) of 41 compounds identifiable as peaks in *D. santomea*

male (purple) and female hexane extracts (pink). Individual measurements are made on ten pooled, age-

matched adult flies (3 male, 2 female replicates). Calculation of absolute abundance uses known

concentration of the internal standard (octadecane/C18, not shown) and normalizes for number of flies

included in the extract. Relative abundances normalize to the summed absolute abundance of all

compounds excluding the standard. Male-specific (Z)-11-octadecenyl acetate (cVA, purple) and

monomorphic (Z)-7-tricosene (7-T, black) are indicated. Question marks in compound names indicate

uncertainty in chemical assignment. See Table S5 for retention times and diagnostic ions used to identify

and assign CHCs.

**Figure 2-S2. 7-T abundance in males and females of various *Drosophila* species**

**(A)** Absolute 7-T abundances on males (purple) and females (pink) *Drosophila* measured by GC-MS on

hexane extracts from ten pooled, age-matched adult flies (2-8 replicates each). Monomorphic species (*D.*

*santomea, D. yakuba*, *D. simulans*) show similar 7-T abundance between sexes while dimorphic species

show large differences (*D. erecta*, *D. melanogaster*). nd, not detected in either sex (*D. serrata*, *D.*

*ananassae*, *D. willistoni*).

**(B-D)** Fraction of time *D. santomea* males spend courting conspecific (B) or heterospecific dead females

(C,D) perfumed with 10% 7-T or solvent control (hexane). Species selected for having either low (*D.*

*erecta*) or undetectable (*D. ana*) endogenous 7-T levels (from GC-MS). *D. santomea* female targets as

positive controls. 7-T addition elicits courtship from *D. santomea* males in both heterospecific cases.

Significance by Dunn's tests.

**Figure 3-S1. Reduced cVA on *D. santomea* males compared to *D. yakuba* and *D. erecta***

**(A-C)** Mirrored gas chromatograms representing male (purple, top) and female (pink, bottom) cuticular

hydrocarbon (CHC) profiles of *D. santomea* (A), *D. yakuba* (B), and *D. erecta* (C). Individual

measurements made on hexane extracts from ten pooled, age-matched adult flies (2-3 replicates each).

Peaks representing male-specific (Z)-11-octadecenyl acetate (cVA) indicated. *D. santomea*

chromatograms reproduced from Fig. 2A for visual comparison.

**(D,E)** Absolute (D) and relative abundances (E) of cVA on *D. santomea* (orange), *D. yakuba* (green), and

*D. erecta* males. Calculation of absolute abundance uses known concentration of the internal standard

(C18) and normalizes for number of flies included in the extract. Relative abundances normalize to the

summed absolute abundance of all compounds detected excluding the standard. *D. santomea* males show

63% and 75% reduced cVA comparing absolute abundance means or 64% and 65% reductions comparing

relative abundance means to *D. yakuba* and *D. erecta*, respectively.

**Figure 3-S2. CRISPR-Cas9 HDR knock-in strategy used to generate *D. santomea Or67d$^{Gal4}$***

**(A)** Schematic diagrams of the Gal4 transgene and selection marker assembled into an Atalanta vector

(pJAT32, addgene #204297) (*148*) and *Or67d* genomic target locus shown with upstream and downstream

sgRNA cleavage sites. Transgene (above) contains two cassettes: *T2A-Gal4-stop(hsp70)* oriented natively

to the genomic target, and downstream *ie1-mCherry-stop(p10)* fluorescent marker for successful

integration (*32*) with inverted orientation. One kilobase homology arms flank both ends. *Or67d* genomic

locus (below) is targeted using sgRNAs directing double-strand breaks to immediately (15 bp)

downstream of the translational start codon and 348 bp downstream of the stop, resulting in removal of

1152 of 1167 protein coding base pairs (99%). Co-injection of the assembled Atalanta vector and nls-

Cas9-nls protein into wildtype *D. santomea* eggs gives mosaic *ie1-mCherry$^{pos}$* G0 progeny carrying *T2A-*

*Gal4* inserted in-frame with the five residual residues of *Or67d*. G1 progeny resulting from G0 crosses to

wildtype are re-screened for abdominal *ie1-mCherry* expression to ensure germline transmission. In this

strategy spatiotemporal patterns and magnitude of Gal4 expression are controlled entirely by *Or67d*

endogenous genomic regulation.

**(B)** Photomicrograph showing labeling pattern of *Or67d$^{Gal4}$* in the whole brain of a *D. santomea* male

(same fly from which close-up of antennal lobes (ALs) is shown in Fig. 3F). Immunostaining for Gal4-

dependent cytoplasmic tdTomato pseudocolored green with Bruchpilot synaptic counterstain (using nc82

monoclonal antibody) in magenta. Uniglomerular labeling pattern in each AL (arrowheads) matches the

expected size and position of cVA-sensitive DA1 (*145*). Bifurcating OSN commissures (axon bundles)

from the antennae leading into ALs can also be seen. Scale bar, 50 µm.

**Figure 3-S3. Ineffectiveness of naturally varying cVA levels to promote aggression by *D. santomea***

**males**

**(A)** Absolute cVA abundance on *Drosophila* males measured by GC-MS on hexane extracts from ten

pooled, age-matched adult flies (2-8 replicates each). Calculation uses known concentration of the

internal standard (C18) and normalizes for number of flies included in the extract. *D. santomea* (orange),

*D. yakuba* (green), and *D. erecta* measurements reproduced from Fig. 3-S1 for visual comparison to

additional species. Means indicated above and phylogeny below.

**(B)** Spontaneous aggression (lunges per minute) exhibited by *D. santomea* males toward conspecific

(orange) or heterospecific males (*D. yakuba* green, all others black) in single-housed pairs during twenty-

minute interactions. Statistical groupings by Dunn's tests shown above and phylogeny below. Note

similarly low or even decreased attack levels toward males of all species.

**(C)** Correlation between cVA abundance measured by GC-MS and aggression elicited from *D. santomea*

males. Dashed line, linear fit. Adjusted $R^2$, significance by F-test for fit, and Pearson correlation (r)

indicated at bottom right. Note marginally significant correlation driven mostly by low attack rates toward

distant species (*D. serrata*, *D. ananassae*, *D. willistoni*) with many other pheromone changes in addition

to low cVA. Three of four species with higher cVA abundance than *D. santomea* elicit less attack than

observed among *D. santomea* males (*D. simulans* is the exception but the slight increase is non-

significant). Data derived from (A,B).

**Figure 3-S4. cVA abundances and social behavior changes by *melanogaster* subgroup males toward**

***D. santomea* males**

**(A)** Relative fraction of time *D. yakuba* (green), *D. simulans*, and *D. melanogaster* males court

conspecific males (first distribution of each pair) vs. *D. santomea* males (second distribution, orange). All

measurements within each species normalized to the conspecific median and $\log_{10}$ transformed. Males of

all three *melanogaster* subgroup species show significantly increased courtship toward *D. santomea*

males by Mann-Whitney *U* tests, including *D. melanogaster* where 7-T (abundant on *D. santomea* males)

is male-specific and sexually aversive (*71–74*).

**(B)** Similar to (A) but showing paired relative distributions for aggression within and between males of

the three subgroup species. Note decreases in male-directed aggression by all three toward *D. santomea*

(significant for *D. yakuba* and *D. melanogaster* by Mann-Whitney *U* tests).

**(C)** Heatmap showing median fraction time spent courting by males in conspecific (diagonal) and

heterospecific male-male pairs (off-diagonal) among the four subgroup species. Rows ("courters") are

males whose courtship is measured (UWE). Columns ("courtees") are males targeted by courtship. Gray

entry where data not collected. Data derived from (A) and additional pairings. Horizontal dashed arrows

identify relevant comparisons between conspecific male- and *D. santomea* male-directed courtship for

visual aid.

**(D)** Similar heatmap showing median aggression (lunge rate) by male "attackers" (rows) and "attackees"

(columns). Data derived from (B) and additional pairings. Note consistently decreased aggression toward

*D. santomea* males.

**(E)** Summarized changes in courtship (first row) and aggression rates (second row) by heterospecific

males toward *D. santomea* compared to conspecific rates for each, derived from data in (A,B). Percent

increases in first row (courtship) and percent decreases in second (aggression).

**Figure 4-S1. Induction of time-locked wing threat by photoactivation of AIP neurons in solitary *D.***

***santomea* males**

**(A)** Optogenetic photoactivation scheme for solitary *D. santomea* males carrying *20E08^{Gal4}* and Gal4-

dependent CsChrimson. Group-housed *20E08^{Gal4}* tester males or genetic controls are exposed to two

fifteen-second photostimulation (PS) blocks with a thirty-second baseline and inter-block interval.

**(B)** Manually scored wing threat rasters (purple) for solitary *20E08^{Gal4}* males during the two-minute trial.

Note threat induction time-locked to PS in most flies. Genetic controls showed no threat.

**(C,D)** Locomotor dynamics in *20E08^{Gal4}* males (browns) and genetic controls (black). Per-frame

calculations of each fly's forward velocity (C) and angular velocity (D) are derived from automated

tracking (*55*). Data is smoothed by averaging within a five-second sliding window (precluding analysis of

sub-second locomotor dynamics such as "turn" and "charge") (*102*). Envelopes represent s.e.m.

**Figure 4-S2. Induction of time-locked courtship by photoactivation of $P1^{dsx}$ neurons in solitary *D.***

***santomea* males**

**(A)** Optogenetic photoactivation scheme for solitary group-housed *D. santomea* males carrying $P1^{dsx}$

split-Gal4 (*71G01$^{DBD}$;dsx$^{AD}$*) and Gal4-dependent CsChrimson. $P1^{dsx}$ tester males or a genetic control are

exposed to six thirty-second photostimulation (PS) blocks with fixed intensity and monotonically

increasing frequency as indicated, with a two-minute baseline and one-minute inter-block intervals.

**(B)** Courtship rasters (red) for $P1^{dsx}$ tester males during the twelve-minute trial. Note early but sporadic

courtship bouts that time-lock to PS blocks and increase in penetrance starting at 5 and 10 Hz.

**(C,D)** Penetrance (C) and expressivity (D) of courtship (UWE) elicited by $P1^{dsx}$ PS. Significant courtship

expression during PS blocks compared to the genetic control in (D) by Mann-Whitney *U* tests.

**Figure 4-S3. Induction of courtship and attack by photoactivation of $P1^{dsx}$ neurons in same-**

**genotype pairs of *D. santomea* males**

**(A)** Optogenetic photoactivation scheme for same-genotype pairs of group-housed *D. santomea* males

carrying $P1^{dsx}$ split-Gal4 and Gal4-dependent CsChrimson. $P1^{dsx}$ tester males or a genetic control are

exposed to six thirty-second photostimulation (PS) blocks with fixed intensity and monotonically

increasing frequency as indicated, with a two-minute baseline and one-minute inter-block intervals.

**(B,C)** Behavior rasters for $P1^{dsx}$ male pairs (B) and genetic controls (C) during the twelve-minute trial.

Blue, attack (lunge); red, courtship (UWE). Note early induction of attack which increases in penetrance

and expressivity as the trial continues. Attacks interrupted by time-locked courtship during later PS blocks

(>5 Hz). Genetic controls show little courtship or attack.

**(D,E)** Locomotor dynamics in $P1^{dsx}$ males (browns) and genetic controls (black). Per-frame calculations

of each fly's forward velocity (D) and angular velocity (E) derived from automated tracking (*55*). Data

smoothed by averaging within a ten-second sliding window. Envelopes represent s.e.m. PS time-locked

locomotor arrest similar to observed previously in *D. melanogaster* (*56*).

**Figure 4-S4. Photoactivation effects of *D. santomea* P1$^{dsx}$ neurons in mixed-genotype male pairs**

**(A)** Expressivity and temporal dynamics of attack (lunge) elicited by PS in mixed-genotype pairs between

a *P1$^{dsx}$* split-Gal4 tester male carrying Gal4-dependent CsChrimson (or genetic control) and group-housed

wildtype *D. santomea* target male. Traces show the mean fraction of frames containing lunge in a ten-

second sliding window with s.e.m. envelopes for *P1$^{dsx}$* (blue) and control (black). PS block frequency,

intensity, and timing indicated above for the twelve-minute trial. Note significant attack induction (Mann-

Whitney *U* tests) in *P1$^{dsx}$* starting after 5 Hz.

**(B)** Attack elicited during inter-bout intervals and after final PS in same-genotype *P1$^{dsx}$* male pairs (blue,

data from Fig. 4I) vs. mixed-genotype pairs between a *P1$^{dsx}$* tester and group-housed wildtype male

(orange, from A). Period means connected with solid lines and data from individual flies shown as thin

lines. Note significant right shift in mixed pairs (Mann-Whitney *U* tests) indicating higher PS frequencies

required to induce intense attack.

**(C)** Action latencies for courtship and attack exhibited by *P1$^{dsx}$* tester males in same- and mixed-genotype

pairs under identical PS trial structures. Data points derived from the same fly pair (or tester fly for mixed

pairs) are connected as lines and distribution means shown as paired dots. Attack and courtship latencies

are similar on average in same-genotype pairs whereas attack is significantly delayed in mixed pairs.

Significance by paired or unpaired Mann-Whitney *U* tests (for within- or between-genotype group

comparisons, respectively).

**(D)** Fraction of flies showing attack (lunge, blue) and courtship (UWE, red) in *P1$^{dsx}$* mixed-genotype

pairs. Note overlap of actions during middle phase of the trials (boxed).

**(E)** Relative frequencies of *P1$^{dsx}$* same- and mixed-genotype pairs showing courtship (red) or attack (blue)

first. Significance between distributions and between each distribution and a null probability of 50%

(random chance) determined by binomial tests. Note courtship precedence in mixed pairs.

**Figure 5-S1. cVA promotes *D. santomea* female sexual rejection during courtship by conspecific**

**males**

**(A)** Rasters of *D. santomea* female and male sexual behaviors during dyadic courtship interactions with

increasing cVA on a nearby piece of filter paper. For each pair, male courtship (UWE, red) is shown

above and female rejection (ovipositor extrusion, orange) or acceptance (wing spreading, cyan) below.

Timing of copulations indicated with asterisks. cVA doses indicated at top. Note increased rejection and

decreased copulation rates at 1 and 2 mg cVA.

**Figure 5-S2. Unconditioned place preference tests with cVA in *D. melanogaster* and *D. san***

**(A)** Video still illustrating the y-maze arena. Four group-housed *D. melanogaster* males are introduced

into the central bowl at the bottom and odors (10% cVA in acetone or acetone control) into ports at the

head of each arm. Vacuum pressure continuously pulls air down the arms through an outlet. Flies can

walk freely within the bowl and arms during ten-minute trials and odors are replaced into the same port

(to avoid contamination) before each trial. Scale bar, 10 mm.

**(B)** Cumulative position trace representing movement of all four flies throughout the duration of the trial.

Position density in each of multiple trials is used to calculate the odor preference index.

**(C-H)** Video stills and cumulative position traces as in (A,B) for *D. santomea* males (C,D), *D.*

*melanogaster* females (E,F), and *D. santomea* females (G,H). Scale bars, 10 mm.

**(I)** Odor preference indices by species and sex. Index calculated as the difference in fly occupancy

between the two arms normalized by the sum. Small gray dots represent single trials with 2-4 flies each

(8-14 trials and 26-44 flies per species/sex). Distributions represented as violins with means shown as

large black dots. Note significantly reduced attraction (or increased aversion) to 10% cVA in both *D.*

*santomea* males and females (orange) compared to *D. melanogaster* (black). Significance by Mann-

Whitney *U* tests.

**Figure 6-S1. Aggression and courtship biases in *melanogaster* subgroup male-male pairs**

**(A-D)** Behavior rasters for three twenty-minute conspecific male-male pairs each of wildtype *D.*

*melanogaster* (A), *D. simulans* (B), *D. yakuba* (C), and *D. santomea* (D) (same pairs as in Fig. 1A).

Annotations of six aggressive interactions by manual scoring (blues) are shown above the horizontal

dashed lines and five courtship actions (reds) beneath. Actions split by identity of the male to which they

are attributed with the one exhibiting the greater total number of actions assigned as "Male 1." *D.*

*melanogaster* and *D. simulans* males fight with some characteristic bias between flies, *D. yakuba* males

are mostly passive but can either fight or court, and *D. santomea* males show frequent and elaborate

courtship often with strong bias.

**(E)** Spontaneous action biases observed between flies in male-male pairs. Male-directed courtship (red

dots) and attack (blue) are plotted as a function of their bout abundance (x-axis) and bias indices (y-axis).

Measurements from the same pair connected by thin lines. Cases where the same fly is dominant for both

actions show connections confined to the sector above the x-axis. Circled "X" indicates mean value for

the corresponding action by color. Boxed regions indicate pairs with high attack expression and wide

range of biases in *D. melanogaster* or high courtship expression and consistently strong bias in *D.*

*santomea,* for visual aid.

1014

**Figure 6-S2. Weak supporting evidence for territoriality in paired *D. santomea* males**

**(A)** Video still of a pair of single-housed, wildtype *D. santomea* males in a large behavior chamber (40 mm wide x 50 mm long x 60 mm tall) containing an apple juice-based "food patch" (25 mm diameter) and neutral surround used to test territoriality. (A') Close-up of intermale courtship from boxed region. Scale bars, 10 mm (A), 1 mm (A').

**(B-D)** Heatmaps representing cumulative spatial patterns of all fly positions (B), attack (lunge) positions (C), and courtship (UWE) positions (D) for 17 *D. santomea* male-male pairs recorded during ten-minute interactions. Normalization in (B) to all frames, in (C) to attack frames only, and in (D) to courtship frames only. Note little sign of increased density on or around the food patch in any case, unlike previously observed for *D. melanogaster* aggression in similar assays (*59*).

**(E)** Quantification of (B-D) showing the fraction of time spent on the food patch in total, during attack, and during courtship for each individual *D. santomea* male. Normalizations as in heatmaps above. None are significantly different from having a null median of 25% (random chance, calculated as surface area of the food patch relative to the full arena) by one-sample Mann-Whitney *U* tests. Attack frames show near-significant deviation from chance, but in the opposite direction from that expected for territoriality (enrichment off the food patch).

**(F)** "Territoriality index" for *D. santomea* males calculated as the difference in time spent on the food patch between flies normalized by the sum. First distribution represents all pairs, second filtered for pairs where one of the flies showed attack dominance, and third filtered for pairs where one of the flies showed courtship dominance. None are significantly different from having a null median of zero (no territorial advantage) by one-sample Mann-Whitney *U* tests.

**Figure 6-S3. Additional behavior characterizations for the sliding door assay**

**(A)** Optogenetic photoactivation scheme for isolated *D. santomea* males carrying *P1$^{dsx}$* split-Gal4 (*71G01$^{DBD}$;dsx$^{AD}$*) and Gal4-dependent CsChrimson during the first phase of the delayed encounter ("sliding door") assay. After a one-minute baseline, two group-housed *P1$^{dsx}$* tester males on either side of a removable divider are exposed to a one-minute photostimulation (PS) block. PS is followed by a ten-minute delay before doors are opened and flies allowed to interact. Delay phase shown as split into one- and nine-minute subphases to reflect cessation of detailed courtship annotations one minute after PS offset (videos were visually scanned after this point and very little UWE was observed).

**(B)** Manually scored courtship rasters (UWE, red) for isolated *P1$^{dsx}$* tester males during the first three minutes of the trial.

**(C,D)** Expressivity (C) and penetrance (D) of courtship elicited by PS. (C) Traces show the mean fraction of frames containing UWE in a ten-second sliding window with s.e.m. envelopes for *P1$^{dsx}$* (red) and a genetic control (black, no courtship observed). (D) Fraction of flies showing courtship. Note extremely high penetrance and expressivity of induced courtship time-locked to PS in *P1$^{dsx}$*.

**(E,F)** Fraction of flies showing courtship (red) and attack (blue) during the ten-minute interaction phase after doors open. (E) *P1$^{dsx}$* in same-genotype male pairs (both experienced PS). (F) *P1$^{dsx}$* males with wildtype females. Switching target sex produces a dramatic shift from predominant aggression to exclusive courtship.

**Figure 7-S1. Aggression and courtship by *D. persimilis* and *D. pseudoobscura* males**

**(A,B)** Behavior rasters for three twenty-minute conspecific male-male pairs each of wildtype *D.*

*persimilis* (A) and *D. pseudoobscura* (B). Annotations of seven aggressive interactions by manual scoring

(blues and black) are shown above the horizontal dashed lines and five courtship actions (reds) beneath.

Actions split by identity of the male to which they are attributed with the one exhibiting the greater total

number of actions assigned as "Male 1." *D. persimilis* males court frequently including making

copulation attempts (e.g., Pair 3). *D. pseudoobscura* also show some courtship but attack more intensely

and often.

**(C,D)** Ethograms representing male- (C) and female-directed social behaviors (D) by *D. persimilis* and *D.*

*pseudoobscura* males. Male-female ethograms represent expert annotation for three recordings of male-

female pairs per species, with male flies similarly prepared by single-housing and interactions taking

place under identical conditions. Male-directed ethograms reproduced from Fig. 1-S1A for visual

comparison to female-directed counterparts. Note similarity between interactions with males and females

in *D. per*.

**(E)** Spontaneous bias observed between flies in male-male *D. persimilis* and *D. pseudoobscura* pairs for

attack and courtship. Indices derived from the same fly pair are connected as lines and distribution means

shown as paired purple (*D. per*) or blue dots (*D. pse*). Both species show strong behavioral biases.

1073

1074   **Materials and Methods**

1075

**Fly stocks**

All species, stocks, and crosses were reared at 24-25C and 50% humidity on a 12hr:12hr light

cycle with standard Caltech fly media. Details about the flies used in each experiment (species, strain,

sex, genetic background, and experimental and control genotypes) are listed in Supplemental Tables 2 and

3. Supplemental Table 2 provides identifiers for the 18 wildtype strains and lists their usage by figure.

Supplemental Table 3 provides similar information for *D. santomea* transgenic lines. All wildtype strains

were obtained from the National Drosophila Species Stock Center (Cornell College of Agriculture and

Life Science, Ithaca, NY) except *D. melanogaster* Canton-S (Anderson lab, Caltech) and three additional

*D. santomea* strains (David Stern lab, Janelia Research Campus).

1085

**Generation of transgenic and knock-in alleles in *D. santomea***

Transgenic *D. santomea* driver and hemidriver lines using enhancer fragments to regulate Gal4

expression were made using phiC31 recombination (*106*) into existing attP landing sites (*107*). White-

mutant *D. santomea* attP sites 2251 (2R) and 2253 (3R) (*107*) were targeted using Gal4 (pBPGUw),

Gal4DBD (pBPZpGAL4DBDUw), or p65AD (pBPp65ADZpUw) vectors carrying an attB recognition

site and mini-white selection marker (*149*, *150*). When available, ready-made constructs with desired

enhancer sequences were obtained from Janelia. Otherwise enhancer sequences were obtained from (*149*),

synthesized by Twist Bioscience (South San Francisco, CA) with flanking attL1/L2 recognition sites, and

cloned upstream of transcriptional control cassettes in the backbone vectors using Gateway LR reactions

(Thermo Fisher Scientific, Waltham, MA). Assembled constructs were verified by sequencing and

restriction digest, maxi-prepped using endotoxin-free kits, and cleaned using ethanol precipitation before

sending for injection. Rainbow Transgenic Flies, Inc. (Camarillo, CA) generated injection mixes

containing the construct and phiC31 integrase mRNA (and sometimes also integrase plasmid to increase

efficiency) and performed injections into *D. santomea* eggs in batches of 200-250 per desired

transformant. Viable G0 (injected) flies were collected and crossed to a white-mutant strain (*2041/w-

[3.2]*) (*107*) before screening for yellow-eyed (transgene heterozygous) flies in the G1 progeny to confirm

germline transmission. Heterozygotes were crossed together to generate transgene homozygotes as stable

lines used for experiments.

Knock-in driver lines were generated using CRISPR/Cas9-mediated homology-directed repair

(HDR) (*104*). Donor construct assembly and sgRNA selection for *D. santomea dsx-p65AD* were

performed previously (*105*) but transformants positive for the selection marker (*MHC-DsRed*) had not

been obtained. The same reagents were re-injected into *D. santomea 146/STO-CAGO 1482* (David Stern,

Janelia) in a mixture with nls-Cas9-nls protein (PNA Bio, Thousand Oaks, CA) and one transformant with

germline transmission of abdominal mCherry was successfully recovered. Crosses among G1 progeny

yielded sporadic G2 flies with the expected *dsx* genital phenotype, verifying the genomic insertion.

*D. santomea Or67d*$^{Gal4}$ was generated using the Atalanta strategy (*148*). A backbone vector with

two U6 promoters (for expression of two sgRNAs) and an "empty payload" (pJAT32) was selected as the

Gateway acceptor. Two donor sequences were synthesized by Twist Bioscience and cloned into the

acceptor in a simultaneous double-Gateway reaction. The first (flanked by attL3/L4) is a *T2A-Gal4-*

*stop(hsp70)* cassette (*Gal4-stop(hsp70)* from pBPGUw) designed to be inserted downstream from but as

close as possible to the *D. santomea Or67d (3L:LOC120447699)* translational start codon. Placement of

the adjacent homology arm ensures that insertion occurs in-frame with the residual 15 coding base pairs

so that Gal4 is properly translated. The second (flanked by attL5/L6) is an *ie1-mCherry-stop(p10)*

selection marker (*32*) with inverted orientation relative to Gal4. Both donors also contain a 1 kb

homology arm and U6-tRNA-sgRNA-tRNA array which, following LR recombination, is positioned

downstream of U6. Candidate sgRNAs were designed in Geneious Prime (Auckland, NZ) and a pair was

selected which excises 99% of *Or67d* coding sequence, resulting in a null allele (upstream

*AATGGCAAAAGTTGAGCCCG*, downstream *ACTGCTGGTTTCCAAAGGAG*). Successful Gateway

products were verified by restriction digest and tiled sequencing. Injection into *D. santomea 146/STO-*

*CAGO 1482* was performed by Rainbow Transgenic Flies using nls-Cas9-nls protein as the enzyme

source and G1 progeny were screened for germline-transmitting abdominal mCherry. Multiple

independent insertion lines were recovered and homozygosed, one of which was used throughout after

anatomical verification of uniglomerular labeling in the antennal lobes of the size and position expected

for Or67d OSNs (*145*). Note that in this strategy Gal4 expression is controlled exclusively by the *Or67d*

native genomic regulatory context rather than an exogenous promoter.

1131

**Spontaneous behavior experiments with video recording**

Experimental crosses were seeded with 10-12 virgin females and 5-6 males and flipped every 3-4

days. Experimental flies were collected mostly as virgins on the day of eclosion and reared either in

isolation (single-housed, 1 individual per vial) or in same-sex groups (group-housed, 10-20 individuals

per vial) for 5-6 days until testing. Group-housed (but not single-housed) flies were flipped into fresh

vials 1-2 days before testing. For experiments requiring unambiguous determination of individual identity

(e.g., in mixed-genotype or mixed-species male-male pairs), a small distal portion of one fly's wing was

clipped using a razor blade under light anesthesia. Whenever possible, the "target" fly (individual whose

behavior was not being measured) was selected for wing clipping. Otherwise clips were balanced between

the two interacting species or genotypes. Females were collected and group-housed as were males (mostly

as virgins on the day of eclosion, 10-20 individuals per vial) and age-matched to the males they were

tested with as courtship targets. As this was usually a period of 5-6 days, it is likely that even the few

females that were not virgins upon collection had by the time of the experiment shed residual male

pheromones potentially transferred during prior copulation (e.g., cVA). Group-housed females were also

flipped 1-2 days before testing. Housing for each experiment arranged by figure can be found in

Supplemental Tables 2 and 3 ("single" or "group" under the "Housing" heading).

Behavior experiments took place in the morning between 9:00AM (ZT0) and 1:00PM (ZT4) in a

temperature and humidity-controlled room (24C, 50% humidity) with ambient white lighting. Single,

pairs, or trios of flies interacted in 8-well acrylic circular chambers (~2.0 $cm^2$ surface area, ~2.4 $cm^3$

volume) designed and described previously (*56, 151*). Walls of the chambers were coated with Fluon (Tar

Heel Ants, Raleigh, NC) and lids with a siliconizing fluid (Thermo Fisher) to discourage flies from

climbing. Chamber floors were covered with a thin layer of freshly prepared "food" (2.5% (w/v) sucrose,

2.25% (w/v) agarose in Tropicana apple juice) and illuminated with an 855 nm backlight (Smart Vision

Lights, Muskegon, MI). Experiments testing alternative foods replaced the apple with other juices

acquired commercially (morinda, fig) or from botanical gardens (marula, pandanus), or with water as a

"no food" condition (sucrose was also omitted in this case). Flies were introduced one at a time into

chambers by gentle aspiration through a small hole in the lid. Interactions were video recorded from

above using a Point Grey Flea3 camera running at 30 Hz with a 780 nm long pass IR filter (Midwest

Optical Systems, Palatine, IL). Experiments performed in 8-well chambers are listed as "small" under the

"Chamber" heading in Supplemental Tables 2 and 3.

For copulation tests of Or67d OSN-silenced *D. santomea* females and territoriality assays of

wildtype *D. santomea* males, "large" chambers consisting of a central food patch (apple juice agarose)

and a neutral surround (bare acrylic) were used. Chambers measure 20 $cm^2$ surface area and 120 $cm^3$

volume, and the circular food patch ~4.9 $cm^2$ (25% surface area). Flies (12 females and 12 males for

copulation tests, 2 males for territoriality tests) were gently aspirated into chambers and video recorded

using identical lighting and camera settings as for small chambers described above.

1168

**1169 Optogenetic photoactivation experiments**

Flies were collected on the day of eclosion into vials containing food supplemented with 400 µM

all *trans*-Retinal (Millipore Sigma, Burlington, MA) and reared in the dark for 5-6 days, with one flip into

fresh supplemented vials 1-2 days before testing. Photoactivated flies were usually group-housed (10-20

males per vial), except in male Or67d OSN courtship suppression experiments which were single-housed

to encourage spontaneous courtship prior to stimulation. Group-housed males paired together during

behavior assays came from different vials.

Experiments took place in the small chambers using an 8-LED array setup reported previously

(*56, 151*). LEDs delivered flashing 685 nm red light in photostimulation (PS) blocks with 10 ms pulse

width, frequencies ranging from 1-40 Hz, and intensities 10-260 µW/mm$^2$, calibrated with a photodiode

power sensor (S130VC, Thorlabs, Newton, NJ) placed at the location of the behavior chambers. Exact PS

conditions (timing, frequency, intensity) for each experiment are shown in Figures and described in

Figure Legends.

"Sliding door" experiments testing persistent behavioral responses after PS were performed in 8-

well chambers fitted with vertical slits to allow metal barriers to divide each chamber in half. Barriers

remained in place during the isolation/PS phase of the experiment, which upon completion of the delay

were manually removed during a ~30 second period while video recordings were paused. Once the

barriers were removed recordings resumed for the interaction phase.

1187

**Behavior annotations and analysis**

Annotations of aggressive (A) and courtship behaviors (C) in male-male pairs from each species

for the primary screen were performed manually using action definitions from (*55*) (threat (A), charge

(A), lunge (A), hold (A), tussle (A), unilateral wing extension (C), circle (C), copulation attempt (C)),

(*102*) (pump (A)), (*20*) (bilateral wing extension (C), row (C)), and (*147*) (headbutt (A)). One additional

action observed exclusively in *D. pseudoobscura* males, "barrage," is defined as follows: *the attacking fly*

*pushes into and quickly chases the opponent while both wings are fully extended and vibrating rapidly as*

*if in flight.* This action appears to have been described once before without being named (*130*). Ethograms

were constructed from calculations of action frequencies and transition matrices using a 1-second cutoff.

For most experiments, video recordings were analyzed in an automated fashion by first tracking

the absolute and relative positions of flies in the chamber at every frame (Caltech FlyTracker,

https://kristinbranson.github.io/FlyTracker/) (*55*) and using the tracking information as input to automated

behavior classifiers developed previously for *D. melanogaster* social interactions (*56*), implemented in

JAABA (https://jaaba.sourceforge.net/) (*54*). Two classifiers were used extensively after satisfactory

performance evaluation for species within the melanogaster subgroup: "lunge" for consummatory,

contact-mediated male aggression, and "unilateral wing extension" for male courtship. Recordings

analyzed in this way are listed as "automated" under the "Behavioral analysis" heading in Supplemental

Tables 2 and 3. Manually annotated recordings, including ones analyzed for the female sexual behaviors

ovipositor extrusion (*120, 121*) and wing spreading (*122*), are instead labeled "manual." For experiments

involving a male-female pair, analysis ended at the point of copulation (if it occurred) and the latency to

copulation was used for normalization procedures rather than total interaction time.

1209

**Courtship song recordings and analysis**

Flies (*D. santomea* wildtype strain 14021-0271.00) were raised in standard laboratory conditions at 23C and 12:12 light cycle. Sexually naïve individuals were isolated within 3 hours of eclosion and single-housed for 5-7 days prior to behavioral analysis. Synchronized audio and video song recordings were made within 4 hours of lights-on in the morning using the Song Torrent system (*60*). Individual flies arranged in male-male or male-female pairs were loaded onto opposite sides of a divider in recording chambers and introduced to each other simultaneously at the beginning of 30-minute recordings.

Videos of 16 male-male and 16 male-female pairs of *D. santomea* flies were visually scanned and 3 pairs of each type with ample courtship were selected for audio analysis. Song traces were visualized and annotated in Tempo (*63*) until dozens to hundreds of both "pulse" and "clack" events had been recorded for each courting male (instead of to saturation). The exact amplitude and position of each event's peak was determined by obtaining the maximum signal intensity within a 20 ms time window centered on the annotation point. Events were assigned to being part of a "train" when their separation from another event of the same type was less than 200 ms for pulse and 400 ms for clack.

**Pheromone applications**

(Z)-11-octadecenyl acetate (cVA) and (Z)-7-tricosene (7-T) were acquired commercially from Pherobank B.V. (Wijk bij Duurstede, NL) as neat liquid formulations (i.e., without a solvent). For behavioral experiments with cVA on filter paper, 0.5-2 µl neat formulation was applied just prior to introduction of flies into the chamber. For direct treatment of live flies, cVA neat formulation was diluted to a 10% working solution in acetone (179124, Millipore Sigma), from which 0.2 µl was pipetted onto the abdomen of target flies under anesthesia 1-2 hours prior to behavioral testing, as previously (*89*). Flies that did not rouse were discarded. For direct treatment of dead flies, 0.2 µl of 10% cVA (acetone) or 7-T (hexane) working solutions was directly pipetted onto the abdomen of flies freshly killed by freezing at -80C and glued to the chamber floor using melted agarose of the same "food" composition.

**Odor place preference tests**

The olfactory "y-maze" was constructed by laser cutting multiple layers of 1/8th inch 3143 infrared (IR)-transmitting acrylic (ACRY31430.125PM, ePlastics, San Diego, CA) to shape. The bottom floor layer was finely wet sanded for texture. Chambers at the end of each arm meant for odor reservoirs and a center port for the vacuum line were blocked off with fine mesh. Top layers consisted of a ceiling of static dissipating acrylic (8774K11, McMaster-Carr, Santa Fe Springs, CA) with a rim of IR acrylic to block ambient light, and a final roof layer with small holes above the chambers to allow air to be drawn into and down the arena by the vacuum line. The layers were supported and sealed with three threaded

rods and bolts. Acrylic layers were periodically cleaned with 1% SDS, water, and isopropanol. The arena

was illuminated from below using a custom-built IR 810 nm LED lighting board and diffused with semi-

opaque white acrylic. Videos were collected with a machine vision camera (BFS-U3-04S2M-C: 0.4 MP,

Teledyne FLIR) running at 45 Hz and a Pentax 12mm 1:1.2 TV lens (FL-HC1212B-VG, Ricoh).

Flies were collected on the day of eclosion, separated by sex, and single- or group-housed for one

week until testing. For each trial, 5 µl odor (10% cVA in acetone or acetone control) was first loaded at

the top of each arm of the y-maze then 2-4 cold anesthetized males or females were aspirated into the

central bowl and the lid quickly fastened. Ten-minute recordings initiated once the flies had roused.

Cumulative position traces were obtained by first median filtering on a subset of frames to construct a

background image. OpenCV (4.6.0) was used to perform thresholding on background-subtracted frames.

The resulting "blob detection" indicated animal position per frame, and positions across the entire trial

were plotted using Matplotlib (3.4.3). Normalized occupancy in the left vs. right arm was calculated from

cumulative position traces using cutoffs including the two arms but excluding the central bowl.

1257

**Gas chromatography – mass spectrometry (GC-MS)**

Flies were collected on the day of eclosion, separated by sex, and group-housed in sets of 10 for

one week until analysis. Flies were anesthetized and transferred to 2 ml glass screw top vials (Thermo

Fisher) containing 100 µl hexane (139386, Millipore Sigma) with 10 ng/µl octadecane (C18) internal

standard (O652, Millipore Sigma). Hexane extraction was allowed to proceed for 15 minutes at room

temperature, after which extracts were transferred to glass inserts (Ibis Scientific, Clark County, NV)

positioned in new screw top vials for either immediate analysis or storage at -80C.

An AOC-2i autoinjector (Shimadzu, Kyoto, Japan) was used to inject 2 µl of each sample into a

Shimadzu QP-2020 GC-MS system (Shimadzu, Kyoto, Japan) equipped with a ZB-5MS fused silica

capillary column (30 m x 0.25 mmID, df=0.25) (Phenomenex, Torrance, CA) and helium as the carrier

gas. The injection port was maintained at a temperature of 310C and was operated in splitless mode, with

a column flow rate of 2.15 ml/min. The temperature program consisted of a 40C hold for 1 minute,

followed by a 20C/min ramp to 250C, a 5C/min ramp to 320C, and a 320C hold for 7.5 minutes. The

transfer line to the mass spectrometer was held at 320C and the ion source temperature at 230C. Electron

ionization was performed at 70 eV and MS scans were collected at 2 Hz, spanning 40-650 m/z.

Identification of cuticular hydrocarbon (CHC) and ester compounds was determined by retention index,

diagnostic ions, and comparison to previous GC-MS results from *Drosophila* (*65*, *66*, *152*, *153*). GC-MS

spectra were integrated in python using the pyteomics package (v4.4.0).

1276

1277 **Immunostaining and imaging**

1278   Whole brains were dissected in ice-cold phosphate-buffered saline (PBS) and fixed in 4%

1279 paraformaldehyde for 20 minutes at room temperature (RT). Samples were washed three times in 0.5%

1280 PBST (PBS containing 0.5% (v/v) Triton X-100) for 10 minutes at RT, then permeabilized in 2% PBST

1281 for 30 minutes at RT. Samples were blocked in 0.5% PBST with 5% normal goat serum (NGS) for 1 hour

1282 at RT or overnight at 4C. Samples were incubated with primary antibodies in blocking solution for 48-72

1283 hours at 4C (rabbit anti-DsRed 1:1000, mouse anti-Bruchpilot 1:50) followed by three 10-minute washes

1284 in 0.5% PBST. Samples were incubated in secondary antibodies (anti-rabbit IgG AF568 and anti-mouse

1285 IgG AF633, 1:1000) in blocking solution for 36-48 hours at 4C, then again washed three times in 0.5%

1286 PBST for 10 minutes at RT before mounting.

1287   Samples were mounted in VectaShield antifade medium (Vector Laboratories, Newark, CA) and

1288 imaged on an Olympus Fluoview FV3000 confocal microscope (Tokyo, JP) with inverted 30X/1.05 or

1289 60X/1.30 oil immersion objectives. Image stacks were acquired at 1024 x 1024 $px^2$ spatial resolution with

1290 1.0-1.5 μm z-step intervals. Images (referred to as "photomicrographs") are maximum intensity

1291 projections (MIPs) of image stacks generated in Fiji. Fiji was also used to pseudocolor, overlay, adjust

1292 brightness and contrast, and add scale bars to images.

1293

1294 **Quantification and statistical analysis**

1295   Fly positional tracking, automated social behavior annotations, and song analyses were performed

1296 using software implemented in MATLAB (MathWorks, Natick, MA). Odor place preference and GC-MS

1297 experiments were analyzed in python using Jupyter Notebooks (https://jupyter.org/). The tabulated

1298 outputs of these analyses were saved and exported as text files for final analyses in R (R Project for

1299 Statistical Computing) using RStudio (Posit PBC, Boston, MA). Custom scripts in R were used to

1300 calculate raw data derivations reported in figures, produce plots visualizing them, and perform statistical

1301 tests. Illustrator 2025 (Adobe, San Jose, CA) was used to adjust aesthetic features of plots and assemble

1302 figures.

1303   Measurement distributions are often shown as boxplots highlighting the median and overlayed

1304 data points. Outliers are identified in the standard way (> Q3+1.5*QR or < Q1-1.5*QR) and visually

1305 excluded from whiskers, but are retained in all quantifications (including statistical tests) and still shown

1306 as dots. Violin plots show full distribution ranges as vertically mirrored kernel density plots with mean

1307 highlighted. Statistical tests consisted mostly of non-parametric comparisons between pairs of or multiple

1308 distributions and were implemented in R. Mann-Whitney *U* tests (R base *stats* package: *wilcox.test*

1309 function) were used to compare pairs of distributions (two-sample) or an empirical distribution to a null

1310 median (one-sample), and could be paired or unpaired depending on whether the comparison involved the

1311 same individual or interacting pair. Two-sample Kolmogorov–Smirnov tests (*stats: ks.test*) were used to

compare pairs of time-evolving distributions or cumulative distributions. Kruskal-Wallis tests (*stats:*

*kruskal.test*) and post-hoc Dunn's tests (*FSA: dunnTest*) were used to compare multiple distributions and

assign lettered groupings. Binomial tests (*stats: binom.test*) were used to compare pairs of frequencies for

binary outcomes (two-sample) or an empirical frequency to a null probability (one-sample). Linear fits

and Pearson correlations were calculated using the *lm* and *cor* functions (*stats*), respectively. Benjamini-

Hochberg multiple test correction was used to adjust p-values whenever appropriate (*stats: p.adjust:*

*method='BH'*). Standardized effect sizes were calculated using Cohen's *d* (*effsize: cohen.d*). Tests shown

in each figure can be referenced in Supplemental Table 4 for details including sample sizes, sample units,

test used, whether correction was applied, adjusted p-value, and effect size. Significance levels are

reported in Figures as *<0.05, **<0.01, and ***<0.001.

1322

**Acknowledgments**

We thank the Anderson lab (Caltech) for help throughout the project, including experimental suggestions and conceptual advice. Liliana Chavarria, Gina Mancuso, Celine Chiu (Caltech) for administrative help. Arnold Sanchez (Caltech) for fly stock maintenance. Heather Dionne, Jeremy Hasseman, Elizabeth Kim, Katie Schretter, and Hiroshi Shiozaki (Janelia Research Campus) for constructs, reagents, and fly stocks. Rory Coleman (NYU) and Vanessa Ruta (Rockefeller) for constructs, fly stocks, information related to unpublished results, and comments on the manuscript. Hong Yu and Garrett Kuiken (Rainbow Transgenic Flies) for embryo injections. Kathy Musial, Gary Roberson (Huntington Botanical Gardens), and David Orr (Waimea Botanical Garden) for access to fruits for host specialization experiments. Ryan York (Arcadia Science) for help with behavioral and evolutionary analyses. Peter Andolfatto (Columbia) for access to an assembled and annotated *D. santomea* reference genome. Markus Knaden (MPI Chemical Ecology) for permission to reanalyze published TD-GC-MS data.

**Funding**

This work was supported by a Jane Coffin Childs Memorial Fund Postdoctoral Fellowship, Della Martin Foundation Postdoctoral Fellowship, and Caltech Center for Evolutionary Science grant to YO. NIH NIDA 5R37DA031389 grant and Howard Hughes Medical Institute funding to DJA.

**Author Contributions**

Conceptualization: YO, DJA

Formal Analysis: YO, DJA

Funding Acquisition: YO, DJA

Investigation: YO, THN, HS, ELB

Methodology: ELB, YD, DLS

Resources: HS, MAK, YD, DLS, JP

Software: THN, HS

Visualization: YO, DJA

Writing – Original Draft Preparation: YO

Writing – Review & Editing: YO, THN, HS, ELB, MAK, YD, DLS, JP, DJA

**Competing Interests**

The authors declare no competing interests.

1357 **Data and Materials Availability**

1358 All data are available in the main manuscript or the supplementary materials. Raw source data, analytical

1359 code, constructs and associated maps, and fly stocks will be made publicly available upon submission of

1360 the manuscript in accordance with current Caltech and HHMI policies.

1361

**References**

1. T. Clutton-Brock, Social evolution in mammals. *Science* **373**, eabc9699 (2021).

2. C. B. Stanford, The Social Behavior of Chimpanzees and Bonobos: Empirical Evidence and Shifting Assumptions. *Curr Anthropol* **39**, 399–420 (1998).

3. E. O. Wilson, The Sociogenesis of Insect Colonies. *Science* **228**, 1489–1495 (1985).

4. R. D. Alexander, The Evolution of Social Behavior. *Annu Rev Ecol Syst* **5**, 325–383 (1974).

5. S. A. West, G. A. Cooper, M. B. Ghoul, A. S. Griffin, Ten recent insights for our understanding of cooperation. *Nat Ecol Evol* **5**, 419–430 (2021).

6. N. Tinbergen, A. C. Perdeck, On the Stimulus Situation Releasing the Begging Response in the Newly Hatched Herring Gull Chick (Larus Argentatus Argentatus Pont.). *Behaviour* **3**, 1–39 (1951).

7. A. Butenandt, R. Beckmann, E. Hecker, Über den Sexuallockstoff des Seidenspinners, I. Der biologische Test und die Isolierung des reinen Sexuallockstoffes Bombykol. *Hoppe-Seylers Z für Physiol Chem* **324**, 71–83 (1961).

8. D. W. Pfaff, Luteinizing Hormone-Releasing Factor Potentiates Lordosis Behavior in Hypophysectomized Ovariectomized Female Rats. *Science* **182**, 1148–1149 (1973).

9. M. de Bono, C. I. Bargmann, Natural Variation in a Neuropeptide Y Receptor Homolog Modifies Social Behavior and Food Response in C. elegans. *Cell* **94**, 679–689 (1998).

10. L. Keller, K. G. Ross, Selfish genes: a green beard in the red fire ant. *Nature* **394**, 573–575 (1998).

11. C. F. Yang, M. C. Chiang, D. C. Gray, M. Prabhakaran, M. Alvarado, S. A. Juntti, E. K. Unger, J. A. Wells, N. M. Shah, Sexually Dimorphic Neurons in the Ventromedial Hypothalamus Govern Mating in Both Sexes and Aggression in Males. *Cell* **153**, 896–909 (2013).

12. H. Lee, D.-W. Kim, R. Remedios, T. E. Anthony, A. Chang, L. Madisen, H. Zeng, D. J. Anderson, Scalable control of mounting and attack by Esr1+ neurons in the ventromedial hypothalamus. *Nature* **509**, 627–632 (2014).

13. Z. Wu, A. E. Autry, J. F. Bergan, M. Watabe-Uchida, C. G. Dulac, Galanin neurons in the medial preoptic area govern parental behaviour. *Nature* **509**, 325–330 (2014).

14. H. Chang, S. Cassau, J. Krieger, X. Guo, M. Knaden, L. Kang, B. S. Hansson, A chemical defense deters cannibalism in migratory locusts. *Science* **380**, 537–543 (2023).

15. N. Niepoth, J. R. Merritt, M. Uminski, E. Lei, V. S. Esquibies, I. B. Bando, K. Hernandez, C. Gebhardt, S. A. Wacker, S. Lutzu, A. Poudel, K. K. Soma, S. Rudolph, A. Bendesky, Evolution of a novel adrenal cell type that promotes parental care. *Nature* **629**, 1082–1090 (2024).

16. P. M. O'Grady, R. DeSalle, Phylogeny of the Genus Drosophila. *Genetics* **209**, 1–25 (2018).

17. A. H. Sturtevant, Experiments on sex recognition and the problem of sexual selection in Drosophila. *J Anim Behav* **5**, 351–366 (1915).

18. H. T. Spieth, Courtship Behavior in Drosophila. *Annu Rev Entomol* **19**, 385–405 (1974).

19. A. W. Ewing, Functional Aspects of Drosophila Courtship. *Biol Rev* **58**, 275–292 (1983).

20. M. Cobb, K. Connolly, B. Burnet, Courtship Behaviour in the Melanogaster Species Sub-Group of Drosophila. *Behaviour* **95**, 203–230 (1985).

21. R. J. Greenspan, J.-F. Ferveur, Courtship in Drosophila. *Annu Rev Genet* **34**, 205–232 (2000).

22. R. R. H. Anholt, P. O'Grady, M. F. Wolfner, S. T. Harbison, Evolution of Reproductive Behavior. *Genetics* **214**, 49–73 (2020).

23. M. Bastock, A. Manning, The Courtship of Drosophila melanogaster. *Behaviour* **8**, 85–110 (1955).

24. J. C. Hall, The Mating of a Fly. *Science* **264**, 1702–1714 (1994).

25. D. Yamamoto, M. Koganezawa, Genes and circuits of courtship behaviour in Drosophila males. *Nat Rev Neurosci* **14**, 681–692 (2013).

26. D. A. Wheeler, C. P. Kyriacou, M. L. Greenacre, Q. Yu, J. E. Rutila, M. Rosbash, J. C. Hall, Molecular Transfer
of a Species-specific Behavior from Drosophila simulans to Drosophila melanogaster. *Science* **251**, 1082–1085
(1991).

27. J. Coyne, A. Crittenden, K. Mah, Genetics of a pheromonal difference contributing to reproductive isolation in
Drosophila. *Science* **265**, 1461–1464 (1994).

28. T. R. Shirangi, H. D. Dufour, T. M. Williams, S. B. Carroll, Rapid Evolution of Sex Pheromone-Producing
Enzyme Expression in Drosophila. *PLOS Biol* **7**, e1000168 (2009).

29. Y. Ding, A. Berrocal, T. Morita, K. D. Longden, D. L. Stern, Natural courtship song variation caused by an
intronic retroelement in an ion channel gene. *Nature* **536**, 329–332 (2016).

30. L. F. Seeholzer, M. Seppo, D. L. Stern, V. Ruta, Evolution of a central neural circuit underlies Drosophila mate
preferences. *Nature* **559**, 564–569 (2018).

31. O. M. Ahmed, A. Avila-Herrera, K. M. Tun, P. H. Serpa, J. Peng, S. Parthasarathy, J.-M. Knapp, D. L. Stern, G.
W. Davis, K. S. Pollard, N. M. Shah, Evolution of Mechanisms that Control Mating in Drosophila Males. *Cell
Reports* **27**, 2527-2536.e4 (2019).

32. R. T. Coleman, I. Morantte, G. T. Koreman, M. L. Cheng, Y. Ding, V. Ruta, A modular circuit coordinates the
diversification of courtship strategies. *Nature* **635**, 142–150 (2024).

33. R. Tanaka, Y. Hara, K. Sato, S. Kohatsu, H. Murakami, T. Higuchi, T. Awasaki, S. Kondo, A. Toyoda, A.
Kamikouchi, D. Yamamoto, Cross-species implementation of an innate courtship behavior by manipulation of the
sex-determinant gene. *Science* **389**, 747–752 (2025).

34. T. Dobzhansky, Species of Drosophila. *Science* **177**, 664–669 (1972).

35. J. A. Coyne, H. A. Orr, Patterns of Speciation in Drosophila. *Evolution* **43**, 362–381 (1989).

36. N. Jourjine, H. E. Hoekstra, Expanding evolutionary neuroscience: insights from comparing variation in
behavior. *Neuron* **109**, 1084–1099 (2021).

37. C. T. Miller, D. Gire, K. Hoke, A. C. Huk, D. Kelley, D. A. Leopold, M. C. Smear, F. Theunissen, M. Yartsev,
C. M. Niell, Natural behavior is the language of the brain. *Curr Biol* **32**, R482–R493 (2022).

38. G. L. Hunt, M. W. Hunt, Female-Female Pairing in Western Gulls (Larus occidentalis) in Southern California.
*Science* **196**, 1466–1467 (1977).

39. D. W. Mock, Siblicidal Aggression and Resource Monopolization in Birds. *Science* **225**, 731–733 (1984).

40. M. E. Jacobs, Influence of Light on Mating of Drosophila Melanogaster. *Ecology* **41**, 182–188 (1960).

41. M. A. Dow, F. von Schilcher, Aggression and mating success in Drosophila melanogaster. *Nature* **254**, 511–512
(1975).

42. K. H. Skrzipek, B. Kröner, H. Hager, Aggression bei Drosophila melanogaster— Laboruntersuchungen. *Z für
Tierpsychol* **49**, 87–103 (1979).

43. A. A. Hoffmann, A laboratory study of male territoriality in the sibling species Drosophila melanogaster and D.
simulans. *Anim Behav* **35**, 807–818 (1987).

44. H. A. Dierick, R. J. Greenspan, Molecular analysis of flies selected for aggressive behavior. *Nat Genet* **38**,
1023–1031 (2006).

45. A. C. Edwards, S. M. Rollmann, T. J. Morgan, T. F. C. Mackay, Quantitative Genomics of Aggressive Behavior
in Drosophila melanogaster. *PLOS Genet* **2**, e154 (2006).

46. S. C. Hoyer, A. Eckart, A. Herrel, T. Zars, S. A. Fischer, S. L. Hardie, M. Heisenberg, Octopamine in Male
Aggression of Drosophila. *Curr Biol* **18**, 159–167 (2008).

47. L. Wang, H. Dankert, P. Perona, D. J. Anderson, A common genetic target for environmental and heritable
influences on aggressiveness in Drosophila. *PNAS* **105**, 5657–5663 (2008).

48. S. Chen, A. Y. Lee, N. M. Bowens, R. Huber, E. A. Kravitz, Fighting fruit flies: A model system for the study of aggression. *PNAS* **99**, 5664–5668 (2002).

49. H. Dankert, L. Wang, E. D. Hoopfer, D. J. Anderson, P. Perona, Automated monitoring and analysis of social behavior in Drosophila. *Nat Methods* **6**, 297–303 (2009).

50. E. D. Hoopfer, Neural control of aggression in Drosophila. *Curr Opin Neurobiol* **38**, 109–118 (2016).

51. J. R. David, F. Lemeunier, L. Tsacas, A. Yassin, The Historical Discovery of the Nine Species in the Drosophila melanogaster Species Subgroup. *Genetics* **177**, 1969–1973 (2007).

52. S. Leigh, M. G. Ritchie, A history of studies of reproductive isolation between Drosophila pseudoobscura and D. persimilis. *Fly* **19**, 2439111 (2025).

53. T. O. Auer, M. P. Shahandeh, R. Benton, Drosophila sechellia: A Genetic Model for Behavioral Evolution and Neuroecology. *Annu Rev Genet* **55**, 1–28 (2021).

54. M. Kabra, A. A. Robie, M. Rivera-Alba, S. Branson, K. Branson, JAABA: interactive machine learning for automatic annotation of animal behavior. *Nat Methods* **10**, 64–67 (2013).

55. E. Eyjolfsdottir, S. Branson, X. P. Burgos-Artizzu, E. D. Hoopfer, J. Schor, D. J. Anderson, P. Perona, Detecting Social Actions of Fruit Flies. *ECCV*, 772–787 (2014).

56. E. D. Hoopfer, Y. Jung, H. K. Inagaki, G. M. Rubin, D. J. Anderson, P1 interneurons promote a persistent internal state that enhances inter-male aggression in Drosophila. *eLife* **4**, e11346 (2015).

57. J. Linz, A. Baschwitz, A. Strutz, H. K. M. Dweck, S. Sachse, B. S. Hansson, M. C. Stensmyr, Host plant-driven sensory specialization in Drosophila erecta. *Proc Royal Soc B* **280**, 20130626 (2013).

58. S. J. Certel, M. G. Savella, D. C. F. Schlegel, E. A. Kravitz, Modulation of Drosophila male behavioral choice. *PNAS* **104**, 4706–4711 (2007).

59. R. S. Lim, E. Eyjólfsdóttir, E. Shin, P. Perona, D. J. Anderson, How Food Controls Aggression in Drosophila. *PLOS One* **9**, e105626 (2014).

60. S. Sawtelle, L. Narayan, Y. Ding, E. Kim, E. L. Behrman, J. L. Lillvis, T. Kawase, D. L. Stern, Song Torrent: A modular, open-source 96-chamber audio and video recording apparatus with optogenetic activation and inactivation capabilities for Drosophila. *bioRxiv*, doi: 10.1101/2024.01.09.574712 (2024).

61. E. T. Watson, E. Rodewald, J. A. Coyne, The courtship song of Drosophila santomea and a comparison to its sister species D. yakuba (Diptera: Drosophilidae). *Eur J Entomol* **104**, 145–148 (2007).

62. J. E. Blyth, D. Lachaise, M. G. Ritchie, Divergence in Multiple Courtship Song Traits between Drosophila santomea and D. yakuba. *Ethology* **114**, 728–736 (2008).

63. Y. Ding, J. L. Lillvis, J. Cande, G. J. Berman, B. J. Arthur, X. Long, M. Xu, B. J. Dickson, D. L. Stern, Neural Evolution of Context-Dependent Fly Song. *Curr Biol* **29**, 1089-1099.e7 (2019).

64. N. W. Bailey, M. Zuk, Same-sex sexual behavior and evolution. *Trends Ecol Evol* **24**, 439–446 (2009).

65. J. Jallon, J. R. David, Variations in Cuticular Hydrocarbons Among the Eight Species of the Drosophila melanogaster Subgroup. *Evolution* **41**, 294–302 (1987).

66. F. Mas, J.-M. Jallon, Sexual Isolation and Cuticular Hydrocarbon Differences between Drosophila santomea and Drosophila yakuba. *J Chem Ecol* **31**, 2747–2752 (2005).

67. M. A. Khallaf, R. Cui, J. Weißflog, M. Erdogmus, A. Svatoš, H. K. M. Dweck, D. R. Valenzano, B. S. Hansson, M. Knaden, Large-scale characterization of sex pheromone communication systems in Drosophila. *Nat Commun* **12**, 4165 (2021).

68. F. Lacaille, M. Hiroi, R. Twele, T. Inoshita, D. Umemoto, G. Manière, F. Marion-Poll, M. Ozaki, W. Francke, M. Cobb, C. Everaerts, T. Tanimura, J.-F. Ferveur, An Inhibitory Sex Pheromone Tastes Bitter for Drosophila Males. *PLOS One* **2**, e661 (2007).

69. T. Miyamoto, H. Amrein, Suppression of male courtship by a Drosophila pheromone receptor. *Nat Neurosci* **11**, 874–876 (2008).

70. A. Llopart, S. Elwyn, D. Lachaise, J. A. Coyne, Genetics of a Difference in Pigmentation between Drosophila
yakuba and Drosophila santomea. *Evolution* **56**, 2262–2277 (2002).

71. J.-C. Billeter, J. Atallah, J. J. Krupp, J. G. Millar, J. D. Levine, Specialized cells tag sexual and species identity
in Drosophila melanogaster. *Nature* **461**, 987–991 (2009).

72. L. Wang, X. Han, J. Mehren, M. Hiroi, J.-C. Billeter, T. Miyamoto, H. Amrein, J. D. Levine, D. J. Anderson,
Hierarchical chemosensory regulation of male-male social interactions in Drosophila. *Nat Neurosci* **14**, 757–762
(2011).

73. R. Thistle, P. Cameron, A. Ghorayshi, L. Dennison, K. Scott, Contact Chemoreceptors Mediate Male-Male
Repulsion and Male-Female Attraction during Drosophila Courtship. *Cell* **149**, 1140–1151 (2012).

74. P. Fan, D. S. Manoli, O. M. Ahmed, Y. Chen, N. Agarwal, S. Kwong, A. G. Cai, J. Neitz, A. Renslo, B. S.
Baker, N. M. Shah, Genetic and Neural Mechanisms that Inhibit Drosophila from Mating with Other Species. *Cell*
**154**, 89–102 (2013).

75. D. A. Turissini, D. R. Matute, Fine scale mapping of genomic introgressions within the Drosophila yakuba
clade. *PLOS Genet* **13**, e1006971 (2017).

76. M. Cobb, J.-M. Jallon, Pheromones, mate recognition and courtship stimulation in the Drosophila melanogaster
species sub-group. *Anim Behav* **39**, 1058–1067 (1990).

77. F. M. Butterworth, Lipids of Drosophila: A Newly Detected Lipid in the Male. *Science* **163**, 1356–1357 (1969).

78. G. Brieger, F. M. Butterworth, Drosophila melanogaster: Identity of Male Lipid in Reproductive System.
*Science* **167**, 1262–1262 (1970).

79. R. J. Bartelt, A. M. Schaner, L. L. Jackson, cis-Vaccenyl acetate as an aggregation pheromone in Drosophila
melanogaster. *J Chem Ecol* **11**, 1747–1756 (1985).

80. P. Xu, R. Atkinson, D. N. M. Jones, D. P. Smith, Drosophila OBP LUSH Is Required for Activity of
Pheromone-Sensitive Neurons. *Neuron* **45**, 193–200 (2005).

81. S. Lebreton, V. Grabe, A. B. Omondi, R. Ignell, P. G. Becher, B. S. Hansson, S. Sachse, P. Witzgall, Love
makes smell blind: mating suppresses pheromone attraction in Drosophila females via Or65a olfactory neurons. *Sci*
*Rep* **4**, 7119 (2014).

82. L. Wang, D. J. Anderson, Identification of an aggression-promoting pheromone and its receptor neurons in
Drosophila. *Nature* **463**, 227–231 (2010).

83. W. Liu, X. Liang, J. Gong, Z. Yang, Y.-H. Zhang, J.-X. Zhang, Y. Rao, Social regulation of aggression by
pheromonal activation of Or65a olfactory neurons in Drosophila. *Nat Neurosci* **14**, 896–902 (2011).

84. I. Taisz, E. Donà, D. Münch, S. N. Bailey, B. J. Morris, K. I. Meechan, K. M. Stevens, I. Varela-Martínez, M.
Gkantia, P. Schlegel, C. Ribeiro, G. S. X. E. Jefferis, D. S. Galili, Generating parallel representations of position and
identity in the olfactory system. *Cell* **186**, 2556-2573.e22 (2023).

85. C. Gao, M. Ma, J. Chen, X. Ji, Q. Peng, Y. Pan, Social experience shapes fighting strategies in Drosophila. *eLife*
**13**, RP104212 (2025).

86. J.-M. Jallon, C. Antony, O. Benamar, Un anti-aphrodisiaque produit par les mâles Drosophila melanogaster et
transféré aux femelles lors de la copulation. *C R Acad Sc Paris* **292**, 1147–1149 (1981).

87. S. D. Mane, L. Tompkins, R. C. Richmond, Male Esterase 6 Catalyzes the Synthesis of a Sex Pheromone in
Drosophila melanogaster Females. *Science* **222**, 419–421 (1983).

88. S. Zawistowski, R. C. Richmond, Inhibition of courtship and mating of Drosophila melanogaster by the male-
produced lipid, cis-vaccenyl acetate. *J Insect Physiol* **32**, 189–192 (1986).

89. A. Kurtovic, A. Widmer, B. J. Dickson, A single class of olfactory neurons mediates behavioural responses to a
Drosophila sex pheromone. *Nature* **446**, 542–546 (2007).

90. A. Ejima, B. P. C. Smith, C. Lucas, W. van der G. van Naters, C. J. Miller, J. R. Carlson, J. D. Levine, L. C. Griffith, Generalization of Courtship Learning in Drosophila Is Mediated by cis-Vaccenyl Acetate. *Curr Biol* **17**, 599–605 (2007).

91. T. S. Ha, D. P. Smith, A Pheromone Receptor Mediates 11-cis-Vaccenyl Acetate-Induced Responses in Drosophila. *J Neurosci* **26**, 8727–8733 (2006).

92. R. Benton, K. S. Vannice, L. B. Vosshall, An essential role for a CD36-related receptor in pheromone detection in Drosophila. *Nature* **450**, 289–293 (2007).

93. W. van der G. van Naters, J. R. Carlson, Receptors and Neurons for Fly Odors in Drosophila. *Curr Biol* **17**, 606–612 (2007).

94. M. L. Schlief, R. I. Wilson, Olfactory processing and behavior downstream from highly selective receptor neurons. *Nat Neurosci* **10**, 623–630 (2007).

95. S. R. Datta, M. L. Vasconcelos, V. Ruta, S. Luo, A. Wong, E. Demir, J. Flores, K. Balonze, B. J. Dickson, R. Axel, The Drosophila pheromone cVA activates a sexually dimorphic neural circuit. *Nature* **452**, 473–477 (2008).

96. T. H. Sten, R. Li, F. Hollunder, S. Eleazer, V. Ruta, Male-male interactions shape mate selection in Drosophila. *Cell* **188**, 1486-1503.e25 (2025).

97. M. Koganezawa, K. Kimura, D. Yamamoto, The Neural Circuitry that Functions as a Switch for Courtship versus Aggression in Drosophila Males. *Curr Biol* **26**, 1395–1403 (2016).

98. K. Ishii, M. Wohl, A. DeSouza, K. Asahina, Sex-determining genes distinctly regulate courtship capability and target preference via sexually dimorphic neurons. *eLife* **9**, e52701 (2020).

99. M. Wohl, K. Ishii, K. Asahina, Layered roles of fruitless isoforms in specification and function of male aggression-promoting neurons in Drosophila. *eLife* **9**, e52702 (2020).

100. H. Chiu, E. D. Hoopfer, M. L. Coughlan, H. J. Pavlou, S. F. Goodwin, D. J. Anderson, A circuit logic for sexually shared and dimorphic aggressive behaviors in Drosophila. *Cell* **184**, 507-520.e16 (2021).

101. C. Han, Q. Peng, M. Sun, X. Jiang, X. Su, J. Chen, M. Ma, H. Zhu, X. Ji, Y. Pan, The doublesex gene regulates dimorphic sexual and aggressive behaviors in Drosophila. *PNAS* **119**, e2201513119 (2022).

102. B. J. Duistermars, B. D. Pfeiffer, E. D. Hoopfer, D. J. Anderson, A Brain Module for Scalable Control of Complex, Multi-motor Threat Displays. *Neuron* **100**, 1474-1490.e4 (2018).

103. L. Tao, D. Ayambem, V. J. Barranca, V. Bhandawat, Neurons Underlying Aggression-Like Actions That Are Shared by Both Males and Females in Drosophila. *J Neurosci* **44**, e0142242024 (2024).

104. F. Port, H.-M. Chen, T. Lee, S. L. Bullock, Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in Drosophila. *PNAS* **111**, E2967–E2976 (2014).

105. D. Ye, J. T. Walsh, I. P. Junker, Y. Ding, Changes in the cellular makeup of motor patterning circuits drive courtship song evolution in Drosophila. *Curr Biol* **34**, 2319-2329.e6 (2024).

106. A. C. Groth, M. Fish, R. Nusse, M. P. Calos, Construction of Transgenic Drosophila by Using the Site-Specific Integrase From Phage φC31. *Genetics* **166**, 1775–1782 (2004).

107. D. L. Stern, J. Crocker, Y. Ding, N. Frankel, G. Kappes, E. Kim, R. Kuzmickas, A. Lemire, J. D. Mast, S. Picard, Genetic and Transgenic Reagents for Drosophila simulans, D. mauritiana, D. yakuba, D. santomea, and D. virilis. *G3* **7**, 1339–1347 (2017).

108. N. C. Klapoetke, Y. Murata, S. S. Kim, S. R. Pulver, A. Birdsey-Benson, Y. K. Cho, T. K. Morimoto, A. S. Chuong, E. J. Carpenter, Z. Tian, J. Wang, Y. Xie, Z. Yan, Y. Zhang, B. Y. Chow, B. Surek, M. Melkonian, V. Jayaraman, M. Constantine-Paton, G. K.-S. Wong, E. S. Boyden, Independent optical excitation of distinct neural populations. *Nat Methods* **11**, 338–346 (2014).

109. D. J. Anderson, Circuit modules linking internal states and social behaviour in flies and mice. *Nat Rev Neurosci* **17**, 692–704 (2016).

110. D. Lachaise, M. Harry, M. Solignac, F. Lemeunier, V. Bnassi, M.-L. Cariou, Evolutionary novelties in islands: Drosophila santomea, a new melanogaster sister species from São Tomé. *Proc Royal Soc B* **267**, 1487–1495 (2000).

111. M. -L. Cariou, J. -F. Silvain, V. Daubin, J. -L. D. Lage, D. Lachaise, Divergence between Drosophila santomea and allopatric or sympatric populations of D. yakuba using paralogous amylase genes and migration scenarios along the Cameroon volcanic line. *Mol Ecol* **10**, 649–660 (2001).

112. A. Llopart, D. Lachaise, J. A. Coyne, An anomalous hybrid zone in Drosophila. *Evolution* **59**, 2602–2607 (2005).

113. J. B. S. Haldane, Sex ratio and unisexual sterility in hybrid animals. *J Genet* **12**, 101–109 (1922).

114. J. A. Coyne, S. Y. Kim, A. S. Chang, D. Lachaise, S. Elwyn, Sexual isolation between two sibling species with overlapping ranges: Drosophila santomea and Drosophila yakuba. *Evolution* **56**, 2424–2434 (2002).

115. J. A. Coyne, S. Elwyn, E. Rolán-Alvarez, Impact of experimental design on Drosophila sexual isolation studies: Direct effects and comparison to field hybridization data. *Evolution* **59**, 2588–2601 (2005).

116. J. A. Coyne, H. A. Orr, *Speciation* (Sinauer Associates, Inc., 2004).

117. C. A. Baker, J. Clemens, M. Murthy, Acoustic Pattern Recognition and Courtship Songs: Insights from Insects. *Annu Rev Neurosci* **42**, 1–19 (2019).

118. K. Connolly, R. Cook, Rejection Responses By Female Drosophila Melanogaster : Their Ontogeny, Causality and Effects Upon the Behaviour of the Courting Male. *Behaviour* **44**, 142–165 (1973).

119. K. Kimura, C. Sato, M. Koganezawa, D. Yamamoto, Drosophila Ovipositor Extension in Mating Behavior and Egg Deposition Involves Distinct Sets of Brain Interneurons. *PLOS One* **10**, e0126445 (2015).

120. C. Mezzera, M. Brotas, M. Gaspar, H. J. Pavlou, S. F. Goodwin, M. L. Vasconcelos, Ovipositor Extrusion Promotes the Transition from Courtship to Copulation and Signals Female Acceptance in Drosophila melanogaster. *Curr Biol* **30**, 3736-3748.e5 (2020).

121. F. Wang, K. Wang, N. Forknall, R. Parekh, B. J. Dickson, Circuit and Behavioral Mechanisms of Sexual Rejection by Drosophila Females. *Curr Biol* **30**, 3749-3760.e3 (2020).

122. M. Li, D. S. Chen, I. P. Junker, F. I. Szorenyi, G. H. Chen, A. J. Berger, A. A. Comeault, D. R. Matute, Y. Ding, Ancestral neural circuits potentiate the origin of a female sexual behavior in Drosophila. *Nat Commun* **15**, 9210 (2024).

123. R. A. Baines, J. P. Uhler, A. Thompson, S. T. Sweeney, M. Bate, Altered electrical properties in Drosophila neurons developing without synaptic transmission. *J Neurosci* **21**, 1523–31 (2001).

124. M. E. N. Majerus, The genetics and evolution of female choice. *Trends Ecol Evol* **1**, 1–7 (1986).

125. A. Yurkovic, O. Wang, A. C. Basu, E. A. Kravitz, Learning and memory associated with aggression in Drosophila melanogaster. *PNAS* **103**, 17519–17524 (2006).

126. Y. Jung, A. Kennedy, H. Chiu, F. Mohammad, A. Claridge-Chang, D. J. Anderson, Neurons that Function within an Integrator to Promote a Persistent Behavioral State in Drosophila. *Neuron* **105**, 322-333.e5 (2020).

127. D. E. Lancefield, A genetic study of crosses of two races or physiological species of Drosophila obscura. *Z für Indukt Abstamm- Vererbungslehre* **52**, 287–317 (1929).

128. E. Mayr, Th. Dobzhansky, Experiments on Sexual Isolation in Drosophila IV. Modification of the Degree of Isolation Between Drosophila Pseudoobscura and Drosophila Persimilis and Sexual Preferences in Drosophila Prosoltans. *PNAS* **31**, 75–82 (1945).

129. M. A. Noor, Speciation driven by natural selection in Drosophila. *Nature* **375**, 674–675 (1995).

130. L. Partridge, A. Hoffmann, J. S. Jones, Male size and mating success in Drosophila melanogaster and D. pseudoobscura under field conditions. *Anim Behav* **35**, 468–476 (1987).

131. M. A. F. Noor, J. A. Coyne, Genetics of a difference in cuticular hydrocarbons between Drosophila pseudoobscura and D. persimilis. *Genet Res* **68**, 117–123 (1996).

132. R. J. V. Roberts, S. Pop, L. L. Prieto-Godino, Evolution of central neural circuits: state of the art and
perspectives. *Nat Rev Neurosci* **23**, 725–743 (2022).

133. B. R. Grant, P. R. Grant, Songs of Darwin's finches diverge when a new species enters the community. *PNAS*
**107**, 20156–20163 (2010).

134. S. Enge, C. Mérot, R. Mozūraitis, V. Apšegaitė, L. Bernatchez, G. A. Martens, S. Radžiutė, H. Pavia, E. L.
Berdan, A supergene in seaweed flies modulates male traits and female perception. *Proc Royal Soc B* **290**,
20231494 (2023).

135. C. Zhou, Y. Pan, C. C. Robinett, G. W. Meissner, B. S. Baker, Central Brain Neurons Expressing doublesex
Regulate Female Receptivity in Drosophila. *Neuron* **83**, 149–163 (2014).

136. J.-M. Jallon, A few chemical words exchanged by Drosophila during courtship and mating. *Behav Genet* **14**,
441–478 (1984).

137. T. Chertemps, L. Duportets, C. Labeur, C. Wicker-Thomas, A new elongase selectively expressed in
Drosophila male reproductive system. *Biochem Biophys Res Commun* **333**, 1066–1072 (2005).

138. J. M. Gómez, A. Gónzalez-Megías, M. Verdú, The evolution of same-sex sexual behaviour in mammals. *Nat
Commun* **14**, 5719 (2023).

139. T. Karigo, A. Kennedy, B. Yang, M. Liu, D. Tai, I. A. Wahle, D. J. Anderson, Distinct hypothalamic control of
same- and opposite-sex mounting behaviour in mice. *Nature* **589**, 258–263 (2021).

140. F. A. Issa, D. H. Edwards, Ritualized Submission and the Reduction of Aggression in an Invertebrate. *Curr
Biol* **16**, 2217–2221 (2006).

141. S. M. Lane, A. E. Haughan, D. Evans, T. Tregenza, C. M. House, Same-sex sexual behaviour as a dominance
display. *Anim Behav* **114**, 113–118 (2016).

142. J. M. Smith, G. R. Price, The Logic of Animal Conflict. *Nature* **246**, 15–18 (1973).

143. T. Gregg, M. Hahn, Drosophila 25 species phylogeny, *figshare* (2017).
https://doi.org/10.6084/m9.figshare.5450602.v1.

144. D. E. Miller, C. Staber, J. Zeitlinger, R. S. Hawley, Highly Contiguous Genome Assemblies of 15 Drosophila
Species Generated Using Nanopore Sequencing. *G3* **8**, 3131–3141 (2018).

145. A. Depetris-Chauvin, D. Galagovsky, I. W. Keesey, B. S. Hansson, S. Sachse, M. Knaden, Evolution at
multiple processing levels underlies odor-guided behavior in the genus Drosophila. *Curr Biol* **33**, 4771-4785.e7
(2023).

146. D. Runfola, A. Anderson, H. Baier, M. Crittenden, E. Dowker, S. Fuhrig, S. Goodman, G. Grimsley, R. Layko,
G. Melville, M. Mulder, R. Oberman, J. Panganiban, A. Peck, L. Seitz, S. Shea, H. Slevin, R. Youngerman, L.
Hobbs, geoBoundaries: A global database of political administrative boundaries. *PLOS One* **15**, e0231866 (2020).

147. S. P. Nilsen, Y.-B. Chan, R. Huber, E. A. Kravitz, Gender-selective patterns of aggressive behavior in
Drosophila melanogaster. *PNAS* **101**, 12342–12347 (2004).

148. D. L. Stern, E. Kim, E. L. Behrman, The Janelia Atalanta plasmids provide a simple and efficient
CRISPR/Cas9-mediated homology directed repair platform for Drosophila. *bioRxiv*, 2023.06.17.545412 (2024).

149. B. D. Pfeiffer, A. Jenett, A. S. Hammonds, T.-T. B. Ngo, S. Misra, C. Murphy, A. Scully, J. W. Carlson, K. H.
Wan, T. R. Laverty, C. Mungall, R. Svirskas, J. T. Kadonaga, C. Q. Doe, M. B. Eisen, S. E. Celniker, G. M. Rubin,
Tools for neuroanatomy and neurogenetics in Drosophila. *PNAS* **105**, 9715–9720 (2008).

150. B. D. Pfeiffer, T.-T. B. Ngo, K. L. Hibbard, C. Murphy, A. Jenett, J. W. Truman, G. M. Rubin, Refinement of
Tools for Targeted Gene Expression in Drosophila. *Genetics* **186**, 735–755 (2010).

151. H. K. Inagaki, Y. Jung, E. D. Hoopfer, A. M. Wong, N. Mishra, J. Y. Lin, R. Y. Tsien, D. J. Anderson,
Optogenetic control of Drosophila using a red-shifted channelrhodopsin reveals experience-dependent influences on
courtship. *Nat Methods* **11**, 325–332 (2014).

152. J. Y. Yew, K. Dreisewerd, H. Luftmann, J. Müthing, G. Pohlentz, E. A. Kravitz, A New Male Sex Pheromone
and Novel Cuticular Cues for Chemical Communication in Drosophila. *Curr Biol* **19**, 1245–1254 (2009).

153. L. M. Dembeck, K. Böröczky, W. Huang, C. Schal, R. R. H. Anholt, T. F. C. Mackay, Genetic architecture of
natural variation in cuticular hydrocarbon composition in Drosophila melanogaster. *eLife* **4**, e09861 (2015).

1672

**Table S1. Key reagents and resources**

| Reagent | Source | Identifier | RRID or CAS |
|---|---|---|---|
| *D. san BDPG4U-Gal4@2253* (mw) | This study | | |
| *D. san 20E08-Gal4@2253* (mw) | This study | | |
| *D. san Or67d-Gal4* (ie1-mCherry) | This study | | |
| *D. san 71G01-Gal4DBD@2251* (mw) | This study | | |
| *D. san dsx-p65AD* (MHC-DsRed) | This study | | |
| *D. san UAS-nlsTdT@2253* (mw) | 10.1101/2025.05.06.652433 | | |
| *D. san UAS-GCaMP7s-2A-tdT@2283* (mw) (introgressed 10 generations from *D. yak* ) | Hiroshi Shiozaki, David Stern | | |
| *D. san UAS-CsChrimson-tdT@2252* (mw) | Yun Ding, David Stern | | |
| *D. san UAS-CsChrimson-tdT@2253* (mw) | Yun Ding, David Stern | | |
| *D. san UAS-EGFP-Kir2.1@2174* (mw) | 10.1016/j.cub.2019.02.019 | | |
| pBPGUw | Addgene | 17575 | Addgene_17575 |
| pBPZpGAL4DBDUw | Addgene | 26233 | Addgene_26233 |
| pBPp65ADZpUw | Addgene | 26234 | Addgene_26234 |
| pJAT32 | Addgene | 204297 | Addgene_204297 |
| Gateway LR Clonase II Enzyme mix | Thermo Fisher Scientific | 11791020 | |
| NucleoBond Xtra Maxi EF | Takara Bio | 740424 | |
| Cas9 with dual NLS | PNA Bio | CP01 | |
| anti-DsRed (rabbit) | Takara Bio | 632496 | AB_10013483 |
| anti-Bruchpilot (mouse) | Developmental Studies Hybridoma Bank | nc82 | AB_2314866 |
| Goat anti-rabbit IgG (H+L) Alexa Fluor 568 | Thermo Fisher Scientific | A-11011 | AB_143157 |
| Goat anti-mouse IgG (H+L) Alexa Fluor 633 | Thermo Fisher Scientific | A-21050 | AB_2535718 |
| Normal Goat Serum | Jackson ImmunoResearch Labs | 005-000-121 | AB_2336990 |
| Triton X-100 | Millipore Sigma | X100 | 9036-19-5 |

| | | | |
|---|---|---|---|
| Paraformaldehyde, 16% solution, EM grade | Electron Microscopy Sciences | 15710 | 30525-89-4 |
| VectaShield Antifade Mounting Medium | Vector Laboratories | H-1000 | AB_2336789 |
| all *trans*-Retinal | Millipore Sigma | R2500 | 116-31-4 |
| Hydrocarbon Soluble Siliconizing Fluid | Thermo Fisher Scientific | TS-42800 | 474-02-4 |
| Fluon (PTFE-30) | Tar Heel Ants | | 9002-84-0 |
| (Z)-7-Tricosene (7-T) | Pherobank B.V. | 12457 | 52078-57-6 |
| (Z)-11-octadecenyl acetate (cVA) | Pherobank B.V. | 10421 | 6186-98-7 |
| Octadecane (C18) | Millipore Sigma | O652 | 593-45-3 |
| Hexane | Millipore Sigma | 139386 | 110-54-3 |
| Acetone | Millipore Sigma | 179124 | 67-64-1 |
| Tropicana apple juice | Amazon | B007SNJDXM | |
| Virgin noni juice | Amazon | B082S4Q6C7 | |
| Caramelized fig syrup | Amazon | B0DHY7K9RP | |
| Marula (*Sclerocarya birrea*) | Waimea Botanical Garden (O'ahu, HI) | | |
| Pandanus (*Pandanus furcatus*) | Huntington Botanical Gardens (San Marino, CA) | | |
| *D. san* genome assembly (Prin_Dsan_1.1) | NCBI RefSeq | GCF_016746245.2 | |
| Geneious Prime | Biomatters | | SCR_010519 |
| Caltech FlyTracker | 10.1007/978-3-319-10605-2_50 | | SCR_027431 |
| JAABA | 10.1038/nmeth.2281 | | SCR_027430 |
| Tempo | 10.1016/j.cub.2019.02.019 | | SCR_027432 |
| Fiji | GitHub | | SCR_002285 |
| MATLAB | MathWorks | | SCR_001622 |
| Jupyter Notebook | GitHub | | SCR_018315 |
| R | R Project for Statistical Computing | | SCR_001905 |
| RStudio | Posit PBC | | SCR_000432 |
| Illustrator 2025 | Adobe | | SCR_010279 |

**A** *D. san courting female*
clack train
clack
pulse train
pulse
inter-clack interval
inter-pulse interval

**B** *D. san courting male*
clack train
clack
pulse
pulse train
100 ms

**C** ♂ > ♀
*Clack*
*Pulse*
10 ms

**D** ♂ > ♂
*Clack*
*Pulse*
10 ms

**E** ns
Clack amplitude (mV)
♀ ♂

**F** ***
Inter-clack interval (ms)
♀ ♂

**G** ns
No. clacks per train
♀ ♂

**H** ***
Pulse amplitude (mV)
♀ ♂

**I** ***
Inter-pulse interval (ms)
♀ ♂

**J** ns
No. pulses per train
♀ ♂

**A**

7-tricosene (7-T)

cVA

D. san male
D. san female

CHC abs. abundance (ng/fly)

**B**

7-tricosene (7-T)

cVA

D. san male
D. san female

CHC rel. abundance (% total)

**A**

Atalanta payload

Genomic target

*D. san 3L:LOC120447699*

1 kb

① Co-inject Atalanta plasmid & NLS-Cas9 protein into eggs

② Screen G1 progeny for abdominal mCherry

OR Gal4 knock-in

| *MAKVE GSGEGR...EENPGP MKLLSS...* | | |
|---|---|---|
| Or67d | T2A | Gal4 |

Legend

☐ CDS
☐ promoter
⬡ hsp70 terminator
⬡ p10 terminator

↱ translational start codon
■ T2A self-cleaving peptide
◀ loxP recombination site
△ sgRNA cleavage site
HA homology arm (1kb)

**B**

*D. san Or67d^Gal4^ male*

AL    AL

**A**

20 Hz, 200 µW/mm²

30s  15s  30s  15s  30s

**B**

Wing threat

**C**

20E08 > CsChrimson
Control

Forward velocity (mm/s)

0          1          2 min

**D**

20E08 > CsChrimson
Control

Angular velocity (deg/s)

270
180
90
0

0          1          2 min

**A**

cVA   0          0.5          1          2 mg

Pair 1  ← Copulation
Pair 2
Pair 3

Unilateral wing extension (male court)
Ovipositor extrusion (female reject)
Wing spread (female accept)

1 min