Investigating the resistome, taxonomic composition, and mobilome of bacterial communities in hospital wastewaters of Metro Manila using a shotgun metagenomics approach

Jiaan Carlo E. Santos[1,5], Daphne Janelyn L. Go[3,5,6], Robert D. Unciano[4,5], Paul K. Yu[8], Angelyn R. Lao[2,5], Ma. Luisa D. Enriquez[1,5,7], Llewelyn M. Espiritu[1,5], and Anish M.S. Shrestha[3,5,6]

[1]Department of Biology, College of Science, De La Salle University, Manila
[2]Department of Mathematics and Statistics, College of Science, De La Salle University, Manila
[3]Department of Software Technology, College of Computer Studies, De La Salle University, Manila
[4]Institute of Arts and Sciences, Far Eastern University, Manila
[5]Systems & Computational Biology Unit, Center for Natural Sciences and Environmental Research, De La Salle University, Manila
[6]Bioinformatics Lab, Advanced Research Institute for Informatics, Computing, and Networking, De La Salle University, Manila
[7]Research & Biotechnology Division, St. Luke's Medical Center
[8]Bioinformatics and Genomics Program, Pennsylvania State University College of Medicine, Hershey, PA 17033, USA

Abstract

We profiled antibiotic resistance genes, antibiotic-resistant bacteria, and mobile genetic elements in untreated hospital wastewater from three tertiary hospitals in Metro Manila using shotgun metagenomic sequencing. The resistome profiles revealed high abundances of resistance genes against sulfonamides (*sul1*, *sul2*), aminoglycosides (*aadS*), and macrolides/streptogramins (*msrE*, *mphE*). To the best of our knowledge, this study provides the first report of antiseptic resistance genes (*qacEΔ1*, *qacL*) in hospital wastewater in the Philippines. Resistance genes of critical concern were also detected, including those conferring resistance to β-lactams (*bla*$_{OXA}$, *bla*$_{TEM}$, *bla*$_{GES}$, *bla*$_{NDM}$, *bla*$_{KPC}$), colistins (*mcr-5*), and tetracyclines (*tet(C)*, *tet(A)*, *tet(L)*, *tet(M)*). Bacterial composition was dominated by the phylum Pseudomonadota, with high abundances of the genera *Stenotrophomonas*, *Rhodococcus*, and *Pseudomonas*, while ESKAPEE pathogens were detected at lower levels. A wide diversity of mobile genetic elements associated with multiple drug classes, including antiseptic and disinfectant resistance genes was observed. Overall, this study demonstrates the utility of shotgun metagenomics for comprehensive profiling of the resistome, bacterial composition, and mobilome in hospital wastewaters and provides supplemental data to support AMR surveillance and mitigation in the Philippine healthcare system.

Keywords: metagenomics, antibiotic resistance, bioinformatics, hospital wastewater, resistome

1 Introduction

Antimicrobial resistance (AMR) is one of the most pressing global health threats. In 2019, an estimated 1.27 million deaths were directly attributed to bacterial AMR and 4.9 million deaths associated with it [2]. It is predicted that the economic burden of AMR-related healthcare costs will reach USD 1 trillion annually by 2050 [34]. The Philippines, a lower-middle-income country (LMIC), is among those disproportionately affected with an estimated 15,700 deaths attributed directly to resistant infections and 56,700 deaths associated with AMR in 2019 [2]. Effective surveillance methods are critical for tracing and understanding AMR risks, including the spread of antibiotic resistance genes (ARGs), discovery of novel resistance mechanisms, and identification of bacterial pathogens carrying these genes. The Philippine Department of Health established the Antimicrobial Resistance Surveillance Program (ARSP) in 1988 to monitor clinically significant antibiotic-resistant bacteria (ARB) from clinical samples. This program has reported an alarming increase in AMR in recent years. In 2023, the program documented rising resistance trends, for example, to carbapenems in *E. coli* and *K. pneumoniae*; to ceftazidime, piperacillin, carbapenems, and possible multidrug-resistance in *P. aeruginosa*; and to ceftazidime, carbapenems, piperacillin, and cotrimoxazole in *A. baumannii* [3].

Laboratories participating in the ARSP utilize culture-dependent methods for antibiotic susceptibility testing on clinically significant species. However, there remains a need for more comprehensive approaches in addressing the full scope of AMR. Shotgun metagenomics provides a powerful approach to analyze the genetic material of microorganisms, including unculturable bacteria, eliminating the biases inherent in culture-dependent methods. This approach enables comprehensive identification and profiling of microorganisms, ARGs, and mobile genetic elements (MGEs) that mediate transfer and spread of resistance genes.

Hospitals are recognized hotspots for AMR due to the frequent use of antibiotics in treating several pathogens. Hospital wastewater (HWW) in particular, contains a variety of waste materials including human waste, medical waste, hazardous chemicals, and antibiotic residues. This environment fosters a high concentration and diversity of ARGs, and bacterial pathogens, enabling mutations and horizontal gene transfer (HGT) between ARBs, making HWW an important medium for capturing a snapshot of AMR activity [92]. Indeed, several studies worldwide have applied shotgun metagenomics for AMR profiling in HWW (see e.g. [37], [55], [61], [47], [77]).

We employed shotgun metagenomics to profile the relative abundances of ARGs, bacterial composition, and MGEs in untreated HWW from three tertiary hospitals in Metro Manila, Philippines. We identified a broad range of ARGs, including those against aminoglycosides, macrolides, β-lactams, tetracyclines, sulfonamides, antiseptics and disinfectants. Notably, this is the first report of antiseptic and disinfectant resistance genes (*qacEΔ1* and *qacL*) in HWW in the Philippines. Our findings demonstrate that bacterial composition and ARG profiles in Philippine tertiary hospitals are similar to those observed globally. This research provides valuable data, describing the diversity and prevalence of ARGs and clinically relevant pathogens in healthcare-associated wastewater that inform future strategies for AMR surveillance and mitigation in the Philippines.

Sample ID	Hospital	Date of collection
A1, A2	A	May 2023
A3, A4	A	June 2023
A5, A6	A	July 2023
B1, B2	B	May 2023
B3, B4	B	June 2023
B5, B6	B	July 2023
C1, C2	C	May 2023
C3, C4	C	June 2023
C5, C6	C	July 2023

Table 1: Sample information

2 Methods

2.1 HWW sampling

HWW samples were collected biweekly over a three-month period (May to July 2023) between 8:00 and 11:00 AM from three hospitals (Hospital A, Hospital B, and Hospital C) located in Metro Manila. Untreated HWW samples were obtained from the collection points immediately prior to wastewater treatment, using sterile collection bottles with a total volume of 1000 mL. Samples A1-A6 were collected from Hospital A, from a collection pipe right before the primary collection tank of the treatment facility. Samples B1-B6 were collected from hospital B, from a manhole directly before the primary collection tank of hospital B's treatment facility. Samples C1-C6 were collected from hospital C from a collection chute right before the primary treatment tank of their treatment facility. Sample information is summarized in Table 1.

The surface of the sample bottles was sterilized with ethyl alcohol, placed in a zip bag, labeled, transferred on ice (2–4°C), and processed within 5 hours after collection. The collected HWW samples were subjected to vacuum filtration using 0.22 μm polyethersulfone (PES, Merck Millipore) membrane filters connected to a vacuum filtration assembly (AGAP Laboratories) powered by a portable vacuum pump. For each sample, 100 - 150 mL of HWW was filtered until clogging of the membrane occurred. The used filters were then transferred into 15 mL sterile conical centrifuge tubes (Falcon$^{\text{TM}}$) and stored at 2–4°C until DNA extraction.

2.2 DNA extraction

Bacterial DNA was extracted using the DNeasy$^{®}$ PowerWater$^{®}$ extraction kit (Qiagen, 2020) per the manufacturer's protocol. The quality and quantity of the extracted DNA were evaluated using a NanoDrop 2000c spectrophotometer, yielding >5 ng/μL concentration, purity ratios (A260/280) >1.8, and volumes >10 μL. DNA integrity was checked via electrophoresis (Vivantis) on a 1.2% (w/v) agarose gel run at 100 V for 30 minutes. All samples met the quality control criteria for high-quality DNA extracts. The DNA samples were resuspended in an elution buffer and stored at $-80°$C prior to sequencing. For sequencing, 20 μL aliquots of each sample were placed in 2 mL sterile microcentrifuge tubes sealed with parafilm, and submitted for sequencing.

2.3 DNA sequencing

A total of 18 DNA extracts were subjected to Nextera XT library preparation and shotgun metagenomic sequencing using the Illumina Novaseq6000 platform at the Macrogen Laboratories, South Korea. Shotgun metagenomic sequencing generated an average of 8.8 Gbp of raw sequence data per sample with a read length of 150 bp paired-end reads.

2.4 Quality control of sequence data

Quality assessment of the raw sequence data was performed using FastQC (v 0.12.1) [FastQC link] and MultiQC (v 1.14) [25]. Trimming of adapter sequences, primers, and low-quality reads was performed using Cutadapt (v 4.2) [48], with parameters provided by the sequencing facility based on the Nextera XT library preparation protocol. The following settings were used: `-a CTGTCTCTTATACACATCT -A CTGTCTCTTATACACATCT` for forward and reverse adapter sequences, respectively, `-O 10` to remove adapters with a minimum overlap of 10 bases, `-m 30` to discard reads shorter than 30 bases, and `-q 20` to trim bases with a quality score below 20.

2.5 ARG identification and quantification

Bowtie2 (v 2.5.1) [39] was used to align the sequencing reads against the Comprehensive Antibiotic Resistance Database (CARD) (v 3.2.6, accessed June 2023) [1]. The alignment was performed using the command `bowtie2 -x CARD-db -1 reads1 -2 reads2 -D 20 -R 3 -N 1 -L 20 -i S,1,0.50`, which corresponds to Bowtie2's very sensitive preset.

For an ARG gene G, its number of copies per genome in a sample [89] was estimated as follows:

$$n_G = \frac{150 \times r_G/l_G}{B \times \text{ average bacterial genome size}},$$ (1)

where r_G is the number of reads mapped to G, l_G is the nucleotide length of G, B is the number of bases in the sample, and 150 refers to the length of the sequence reads. The relative abundance of G in that sample, denoted by a_G, was then computed using:

$$a_G = \frac{n_G}{\sum_H n_H},$$ (2)

where the summation is across all ARG H in the sample.

All resistance genes in the CARD database were clustered into their corresponding drug classes. These drug classes categorize ARGs based on their mechanisms of action and the bacterial pathways they target. The clustering was done using the Antibiotic Resistance Ontology (ARO) numbers associated with each resistance gene. For a sample, the total count of ARGs belonging to a drug class D, denoted by n_D, was calculated using Equation 3.

$$n_D = \sum_{G \in D} n_G$$ (3)

The relative abundance of D in that sample, denoted by a_D, was then computed using:

$$a_D = \frac{n_D}{\sum_T n_T}$$ (4)

where the summation is across all Drug Classes T in the sample.

2.6 Bacterial Taxonomy Identification and Quantification

Taxonomic identification of bacterial communities was performed with Kraken 2 [84] using Standard Kraken 2 16GB Database. The relative abundances of various microbial genera were estimated using Bracken [44].

2.7 MGE Identification and Quantification

MGEs were identified by aligning the reads to three reference databases: ISFinder database for insertion sequences (IS) [72], PlasmidFinder database for plasmid replicons [15], and INTEGRALL database for integrases [52]. The alignment was performed using Bowtie2 (v2.5.1) with its very sensitive preset: `bowtie2 -x database_index -1 reads1 -2 reads2 -D 20 -R 3 -N 1 -L 20 -i S,1,0.50`, where `database_index` corresponds to the Bowtie2 index generated from the ISFinder, PlasmidFinder, or INTEGRALL databases.

The relative abundance of an IS I in a sample, denoted by a_I was computed using:

$$a_I = \frac{r_I/l_I}{\sum_H r_H/l_H},\tag{5}$$

where r_I is the number of reads mapped to I, l_I is the nucleotide length of I, and the summation is across all insertion sequences H in the sample.

The relative abundances of plasmid replicons in PlasmidFinder and integrases in INTEGRALL were computed similarly.

3 Results and Discussion

3.1 *sul1*, *sul2*, *aadS*, *msrE*, *mphE*, *qacEΔ1*, and *qacL* were the predominant ARGs across all HWW samples.

Figure 1: Relative abundances of the top ARGs from Hospital A (A), Hospital B (B), and Hospital C (C). ARGs not part of the top were grouped as "Others."

We observed a wide diversity of ARGs across all samples. Figure 1 shows the relative abundances of the top ARGs found at each hospital. The hospitals exhibited similar sets of high-abundance ARGs, which include *sul1*, *sul2*, *aadS*, *msrE*, *mphE*, *qacEΔ1*, and *qacL*.

Of these, *sul1* and *sul2* confer sulfonamide resistance by encoding alternative dihydropteroate synthase (DHPS) enzymes that reduce antibiotic efficacy [80]. These plasmid-mediated genes are

frequently identified in clinical isolates belonging to the genera *Acinetobacter*, *Escherichia*, *Klebsiella*, *Stenotrophomonas*, and *Staphylococcus* [1]. The prevalence of *sul* genes has been documented globally, including in LMICs such as South Africa and India, where they are predominant in HWW [77, 22].

The streptomycin resistance gene *aadS* encodes an aminoglycoside acetyltransferase that inactivates aminoglycoside antibiotics [75, 1]. This gene is typically phenotypically silent in wild-type *Bacteroides*, with expression induced only by a chromosomal mutation [73]. Associated with the phylum Bacteroidota, *aadS* is commonly found on IS and transposons, indicating its potential role in HGT [79, 63]. Its presence has been documented in wastewater systems, including untreated HWWs from South Korea, Mexico, and several member states of the European Union [67, 28, 64].

The plasmid-associated *msrE* and *mphE* encode an ABC-F subfamily protein and a macrolide phosphotransferase respectively, which protect against and inactivate streptogramin and macrolide antibiotics. Studies have reported that both genes confer higher levels of resistance against second-generation macrolides compared to first-generation macrolides, underscoring the potential influence of increased clinical use of second-generation macrolides in recent years [24, 16]. Both ARGs are frequently associated with *Acinetobacter*, *Pseudomonas*, *Escherichia*, and *Klebsiella* among others [8, 1]. The prevalence of both genes has been documented in HWWs across various countries, including the Czech Republic and Scotland (UK) [37, 61]

The *qacE∆1* and *qacL* genes confer resistance to quaternary ammonium compounds (QAC) widely used as surfactants, antimicrobials in disinfectants, and cleaning products[5]. Both genes have been mainly associated with *Acinetobacter*, *Klebsiella*, *Pseudomonas*, and *Stenotrophomonas* among others [1, 38]. Both ARGs are typically plasmid-encoded, enhancing their potential for HGT and environmental dissemination [43]. Their presence has been documented in HWWs from several countries, including China, Israel, and South Africa [82, 88, 62, 22].

3.2   Aminoglycoside, macrolide, β-lactam, tetracycline, and sulfonamide-associated ARGs had the highest combined relative abundances.

Figure 2: Top 10 drug classes based on combined relative abundances of associated ARGs identified in Hospital A (A), Hospital B (B), and Hospital C (C). Drug classes not part of the top 10 were grouped as "Others."

We consolidated the relative abundances of the ARGs based on the drug classes they are reported to be associated with in the CARD database, as shown in Equation 4. Figure 2 shows the top 10 drug classes with the consolidated abundances. The most abundant consolidated drug classes include aminoglycosides, macrolides, β-lactams, tetracyclines, and sulfonamides.

Among these, aminoglycosides are commonly used to treat severe Gram-negative bacterial infections, acting by binding to the 30S ribosomal subunit and inhibiting protein synthesis. Resistance is often mediated by ARGs encoding aminoglycoside-modifying enzymes (AMEs), including *aph(3")-Ib, aph(6)-Id, aac(6')-Ib7, aac(6')-IIa,* and *aac(6')-II* which were observed in our samples. These ARGs are plasmid-mediated increasing the likelihood of HGT across bacterial species. [83, 81, 42, 29]. The WHO classifies aminoglycosides as Critically Important Antimicrobials, highlighting the urgent need to preserve their clinical efficacy and prevent the potential spread of resistance [85].

Macrolides are commonly used to treat pneumonia and manage cystic fibrosis infections caused by multidrug-resistant, hospital-acquired *Pseudomonas aeruginosa* [31]. Multiple ARGs conferring resistance to macrolides were detected in our samples, including efflux pumps genes (*smeE, mefC, axyY, mexK, smeD*), ribosomal target modification genes (*erm(46)*), and antibiotic inactivation genes (*mphG, ereD, eraA2, mphF*). The widespread use of macrolides in both clinical and agricultural settings has contributed to the increasing prevalence of resistance among pathogenic organisms [57, 40, 33]. Like aminoglycosides, macrolides are classified by the WHO as Critically Important Antimicrobials. [85].

β-lactam antibiotics are among the most widely used agents to combat a broad spectrum of Gram-positive and Gram-negative bacterial infections. They act by inhibiting cell wall synthesis through binding to essential penicillin-binding proteins[14]. We detected bla_{OXA}, bla_{GES}, bla_{TEM}, bla_{KPC}, and bla_{NDM} resistance genes in our samples. β-lactams, particularly 3^{rd}- and 4^{th}-generation cephalosporins, are classified as Highest Priority Critically Important Antimicrobials, as their proper use is essential to preserve effectiveness against the increasing threat of multidrug-resistant *Enterobacterales* worldwide [85].

Tetracyclines are broad-spectrum antibiotics that are commonly used to treat and manage various infections. They act by inhibiting the 30S ribosomal subunit, thereby halting bacterial growth and replication [17]. We identified several tetracycline resistance genes, including flavin-dependent monooxygenase (*tet*(X)), major facilitator superfamily efflux pump genes (*tet*(G), *tet*(C), *tet*(39), *tet*(A), *tet*(L)), and ribosomal protection proteins genes (*tet*(M), *tet*(O), *tet*(Q), *tet*(W)). The WHO classifies tetracyclines as Highly Important Antimicrobials, emphasizing the need for continuous surveillance to preserve their clinical efficacy [85].

Sulfonamides are used to treat a range of bacterial infections, including urinary tract infections, respiratory diseases, and opportunistic infections in immunocompromised patients[58]. They act as competitive inhibitors of DHPS in the bacterial folate synthesis pathway, thereby inhibiting their growth and development [80]. Specifically, the ARGs (*sul1, sul2,* and *sul4*) were detected in our samples.

3.3 ARGs of critical concern were observed in all HWW samples.

Previous studies have classified certain ARGs as being of "critical concern" based on their abundance, mobility, and host pathogenicity [90, 71]. Multiple ARGs of critical concern were identified across the three hospitals as shown in Figure 3. ARGs of "critical concern" are further classified into 4 categories (Highest Priority, High Priority, Highly Important, or Important) reflecting their clinical relevance and alignment with the WHO list of medically important antimicrobials for human medicine [85, 90, 71].

Multiple β-lactam resistance genes classified as Highest Priority ARGs were detected across all hospitals. The presence of these ARGs may be attributed to the frequent clinical use of β-lactam antibiotics, which corresponds to the increasing resistance rates among Gram-negative bacilli isolates from Philippine hospitals [3]. National surveillance studies have frequently reported β-lactam resistance genes such as bla_{AIM}, bla_{OXA}, and bla_{TEM}, whereas bla_{NDM} and bla_{KPC} have been detected

Figure 3: ARGs of critical concern. Black boxes indicate the presence of ARGs for each hospital A, B, and C.

in a few clinical isolates [3]. Of concern, bla_{NDM}, a carbapenemase that confers resistance to most β-lactam antibiotics, was first identified in India and has since spread globally [26]. Similarly, bla_{KPC}, first identified from a *K. pneumoniae* isolate in the United States, has been documented in multiple countries including France and Israel [10, 53, 54]. The detection of both bla_{NDM} and bla_{KPC} in our samples raises concern regarding the transmission of these clinically significant ARGs and warrants further investigation at the national level.

The mobile colistin resistance gene (*mcr-5*) was also identified in our samples. Colistin serves as a last-resort treatment against multidrug-resistant Gram-negative infections and is classified as a Highest Priority Critically Important Antimicrobial, highlighting the need to preserve its clinical effectivity [76, 85]. Although clinical studies in the Philippines have primarily reported *mcr-1*, the detection of *mcr-5* in our study is noteworthy due to its higher resistance activity and greater potential for HGT, underscoring the need for continued surveillance of this resistance gene [91, 3].

Tetracycline ARGs, classified as Highly Important, were detected across all hospitals. We identified genes encoding major facilitator superfamily antibiotic efflux pumps (*tet(39)*, *tet(A)*, *tet(L)*) and tetracycline-resistant ribosomal protection proteins (*tet(M)*, *tet(O)*, *tet(Q)*, *tet(W)*). Previous studies in the Philippines have reported the prevalence of ARGs *tet(A)*, *tet(C)*, *tet(M)*, and *tet(L)* in *E. coli* and *P. aeruginosa* isolates, highlighting the similarities of our results with national surveillance efforts [3, 69].

3.4 Antiseptic and disinfectant resistance genes (*qacEΔ1* and *qacL*) were identified for the first time in Philippine HWW samples.

To the best of our knowledge, this is the first report of *qacEΔ1* and *qacL* genes in HWW from the Philippines. Both genes confer resistance to QACs, which are commonly used as disinfectants and cleaning agents[5]. The increased use of QAC-containing products during the COVID-19 pandemic has resulted in greater release of these compounds into the environment [93, 9]. The extensive use of QAC-based disinfectants in hospitals during the pandemic may have contributed to the high abundance of *qacEΔ1* and *qacL* resistance genes in our samples.

The presence of these genes in hospital wastewater has also been documented in China, Israel, Australia, and South Africa [82, 88, 62, 22, 70]. The detection of antiseptic and disinfectant resistance

genes in our samples underscores the need for further investigation in the Philippines to evaluate the potential risks of antiseptic resistance in both clinical and environmental contexts.

3.5 Phyla Pseudomonadota, Actinomycetota, and Bacteroidota dominated the HWW microbial community.

Figure 4: Relative abundance of the top phyla from Hospital A (A), Hospital B (B), and Hospital C (C). Phyla not part of the top were grouped as "Others."

We assessed the bacterial taxonomic composition in our samples. Pseudomonadota (synonym Proteobacteria), Actinomycetota (synonym Actinobacteria), and Bacteroidota (synonym Bacteroidetes) were the most abundant phyla in all HWW samples (Figure 4). These were also reported to be most abundant in HWW samples from other LMICs such as Burkina Faso, Benin, and India [47, 77], highlighting similarities in phylum-level bacterial composition to our HWW samples. Members of Actinomycetota and Bacteroidota are known to be the major constituents of the human gut microbiome and are present in wastewaters [4, 55]. Members of Pseudomonadota are known human pathogens harboring various resistance genes against aminoglycosides, sulfonamides, β-lactams, and tetracyclines [56, 68, 51]. Under the phylum Pseudomonadota, ESKAPEE pathogens were detected at relatively lower abundances in our samples. These pathogens are responsible for a wide range of human infections and include strains that exhibit resistance to most antibiotics, prompting their classification as "high-priority pathogens" by the WHO [86, 21, 23, 59, 50].

Figure 5: Relative abundance of the top genera from Hospital A (A), Hospital B (B), and Hospital C (C). Genera not part of the top were grouped as "Others."

Among the genera under these three phyla, *S.*, *Rhodococcus*, and *Pseudomonas* were consistently among the most abundant across all samples (Figure 5). *Stenotrophomonas* are opportunistic pathogens commonly infecting humans, with known multidrug-resistant species such as

Stenotrophomonas maltophilia [30, 20, 12, 11]. This genus has been documented in various hospital plumbing systems, including sinks, showers, and toilets, which likely contributes to their presence in HWWs [87]. Similarly, reports in India, Singapore and the UK have also identified the genus in HWW [77, 55, 18]. *Rhodococcus* is typically found in soil, water, and animal hosts with a relatively low prevalence in hospital environments. Interestingly, in our study this genus was detected at high abundance in HWW samples. Previous reports of human infections caused by several *Rhodococcus* species indicate their increasing clinical relevance and potential establishment within hospital settings [6, 13, 19]. Similarly, a study in China has detected the presence of *Rhodococcus* in HWWs [94], while other studies have identified them from patient samples [66, 78]. *Pseudomonas* are clinically significant opportunistic pathogens capable of causing a wide range of human infections in hospital settings. They are frequently reported to exhibit resistance to most antibiotics, especially against β-lactams. [23, 21].

Other highly abundant genera included *Achromobacter*, *Brecundimonas*, *Acidovorax*, *Microbacterium*, *Mycolicibacterium*, *Nocardoides*, *Rhodanobacter*, and *Thermomonas*. These genera have been previously described as components of HWW bacterial communities that shape the HWW environment and contribute to the removal of organic and inorganic compounds from wastewaters [77, 47, 18, 45, 32]. Interestingly, *Mycolicibacter* and *Pseudoxanthomonas* were present at high abundance in our samples. Both genera are recognized as environmental contaminants, and their members are typically non-pathogenic to humans. However, recent studies have reported certain species with the potential to cause human infections which may lead to severe diseases [36, 7, 46].

3.6   Diverse MGEs were detected across all HWW samples.

Figure 6: The relative abundance of the top IS from Hospital A (A), Hospital B (B), and Hospital C (C). IS not part of the top were grouped as "Others."

We identified a broad diversity of MGEs of IS and transposons, integrons, and plasmids. Among the most abundant IS and transposons (Figure 6), we observed the presence of IS1071, ISPasp1, IS1247, ISVsa3, ISAba1, and ISStma11. These IS are known to carry ARGs against β-lactams, tetracyclines, aminoglycosides, sulfonamides, macrolides, and disinfecting agents among others. They are known to be associated with the genera *Klebsiella*, *Pseudomonas*, *Stenotrophomonas*, *Rhodococcus*, and *Mycobacterium* among others. [65, 41, 35, 27]. Class 1 integrons (intI1) were the most abundant type of integrons detected in our samples. These integrons are frequently associated with Gramnegative ARBs in clinical settings. Class 1 integrons often contain ARGs conferring resistance to β-lactams, aminoglycosides, sulfonamides, and disinfectants [60]. We have also identified a diversity of plasmid types that carry ARGs including IncQ2 (NCBI accession FJ696404), IncP6 (NCBI

Figure 7: The relative abundance of the top plasmid replicons from Hospital A (A), Hospital B (B), and Hospital C (C). Plasmid replicons not part of the top were grouped as "Others."

accession JF785550), repUS43 (NCBI accession CP003584), Col440II (NCBI accession CP023921), Col(pHAD28) (NCBI accession KU674895) and ColRNAI (NCBI accession DQ298019) among others (Figure 7). These plasmids can carry multiple ARGs against β-lactams, tetracyclines, carbapenems, macrolides, and disinfectants among others [60, 74, 49].

3.7 Our data supplement the findings of the Philippine Antibiotic Resistance Surveillance Program

The Philippine Department of Health (DOH) established the Antimicrobial Resistance Surveillance Program (ARSP) in 1988, to monitor the prevalence and trends of antimicrobial resistance in selected bacterial pathogens. The ARSP reports phenotypic resistance rates of 11 species (*S. pneumoniae*, *H. influenzae*, *S. enterica*, *Shigella*, *V. cholerae*, *N. gonorrhoeae*, *S. aureus*, *Enterococcus*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*, and *A. baumannii*) against commonly used antibiotics [3].

ARBs such as *E. coli*, *K. pneumoniae*, *P. aeruginosa*, and *A. baumannii* among others showed resistance rates above 50% against drug classes of aminoglycosides, β-lactams, tetracyclines, and sulfonamides among others [3]. The detection of abundant ARGs in our samples conferring resistance to aminoglycosides, β-lactams, tetracyclines, and sulfonamides supplements ARSP reports by providing insights into genotypic resistance mechanisms.

The ARSP reports on clinically significant bacterial species that play a role in the spread of AMR in hospital settings. The presence of other ARBs such as *Rhodococcus*, *Mycolicibacter*, and *Pseudoxanthomonas* in our samples underscores the potential of shotgun metagenomics to identify underexplored ARBs for AMR surveillance. These organisms may also contribute to AMR dissemination and represent possible emerging pathogens warranting further investigation.

4 Conclusion

Our study investigated the diversity and abundance of ARGs, bacterial taxa, and MGEs in HWW in three hospitals in Metro Manila, Philippines. ARGs conferring resistance to critically important antibiotics including β-lactams, colistin, and tetracyclines were present in our samples. Interestingly, this is the first report of antiseptic and disinfectant resistance genes (*qacEΔ1*, *qacL*) in HWW in the Philippines. Notable bacterial genera such as *Stenotrophomonas*, *Rhodococcus*, *Mycolicibacter*, and *Pseudoxanthomonas* were detected across all sampling sites. MGEs associated with clinically important drug classes were also detected. These findings are consistent with global studies across

various economic settings, suggesting shared challenges in AMR management. Our HWW resistome profiles provide a more comprehensive view of ARGs and bacterial pathogens circulating in healthcare environments and can supplement data generated by national surveillance programs.

Overall, our results highlight the pressing need for strengthened antimicrobial stewardship and targeted control strategies within healthcare systems. The detection and characterization of ARGs of critical concern, emerging bacterial pathogens, and MGEs are critical to deepen our understanding of AMR interaction in hospital settings. These insights provide valuable evidence to inform national policy, enhance AMR monitoring, and support the development of more effective interventions to mitigate the growing threat of AMR in the country.

5 Acknowledgments

The authors express their gratitude to Dr. Evelina Lagamayo, Ms. Sherill Tesalona, and Mr. Joemarie Malana for their valuable assistance in obtaining ethics clearances for the three hospitals. We also extend our thanks to Mr. Elmer Abanzado, Engr. Sherwin Mina, and Mr. Emerson Vergara for their support in the collection of samples. We also acknowledge the members of the e-ASIA JRP group for their insights and constructive discussions that greatly contributed to this study.

6 Funding

This research was funded by the Department of Science and Technology Philippine Council for Health Research and Development (DOST - PCHRD) through East Asia Science and Innovation Area Joint Research Program (e-Asia JRP). The sponsors had no role in study design, data collection and analysis, decision to publish or manuscript preparation.

7 Availability of data and materials

The datasets generated during the current study are available in the DDBJ Sequence Read Archive under the accession number PRJDB37674 [currently on Hold-Until-Published arrangement]. The computational pipeline used for data analysis is available at the following URL: `https://github.com/bioinfodlsu/attack_amr_pipeline`.

References

[1] B. P. Alcock, A. R. Raphenya, T. T. Y. Lau, K. K. Tsang, M. Bouchard, A. Edalatmand, W. Huynh, A.-L. V. Nguyen, A. A. Cheng, S. Liu, S. Y. Min, A. Miroshnichenko, H.-K. Tran, R. E. Werfalli, J. A. Nasir, M. Oloni, D. J. Speicher, A. Florescu, B. Singh, M. Faltyn, A. Hernandez-Koutoucheva, A. N. Sharma, E. Bordeleau, A. C. Pawlowski, H. L. Zubyk, D. Dooley, E. Griffiths, F. Maguire, G. L. Winsor, R. G. Beiko, F. S. L. Brinkman, W. W. L. Hsiao, G. V. Domselaar, and A. G. McArthur. CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Res.*, 48(D1):D517–D525, 2020.

[2] Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet*, 399(10325):629–655, Feb. 2022.

[3] Antimicrobial Resistance Surveillance Program (ARSP). Annual report, 2023. URL `https://arsp.com.ph/arsp-2023-annual-report-data-summary`.

[4] L. Anto and C. N. Blesso. Interplay between diet, the gut microbiome, and atherosclerosis: Role of dysbiosis and microbial metabolites on inflammation and disordered lipid metabolism. *The Journal of Nutritional Biochemistry*, 105:108991, 03 2022. doi: 10.1016/j.jnutbio.2022.108991.

[5] W. A. Arnold, A. Blum, J. Branyan, T. A. Bruton, C. C. Carignan, G. Cortopassi, S. Datta, J. DeWitt, A.-C. Doherty, R. U. Halden, H. Harari, E. M. Hartmann, T. C. Hrubec, S. Iyer, C. F. Kwiatkowski, J. LaPier, D. Li, L. Li, J. G. Muñiz Ortiz, and A. Salamova. Quaternary ammonium compounds: A chemical class of emerging concern. *Environmental Science & Technology*, 57(20), 2023. doi: 10.1021/acs.est.2c08244.

[6] H. Baba, T. Nada, K. Ohkusu, T. Ezaki, Y. Hasegawa, and D. L. Paterson. First case of bloodstream infection caused by *Rhodococcus erythropolis*. *Journal of Clinical Microbiology*, 47 (8):2667–2669, 2009. doi: 10.1128/JCM.00294-09.

[7] K. A. Bernard, A. Vachon, A. L. Pacheco, T. Burdz, D. Wiebe, D. R. Beniac, S. L. Hiebert, T. Booth, D. A. Doyle, P. Lawson, and A.-M. Bernier. *Pseudoxanthomonas winnipegensis* sp. nov., derived from human clinical materials and recovered from cystic fibrosis and other patient types in canada, and emendation of *Pseudoxanthomonas spadix* Young et al. 2007. *International Journal of Systematic and Evolutionary Microbiology*, 70(12):6313–6322, 2020. ISSN 1466-5034. doi: 10.1099/ijsem.0.004533.

[8] G. A. Blackwell and R. M. Hall. The *tet39* determinant and the *msrE-mphE* genes in *Acinetobacter* plasmids are each part of discrete modules flanked by inversely oriented *p dif* (xerc-xerd) sites. *Antimicrobial Agents and Chemotherapy*, 61(8), 2017. doi: 10.1128/aac.00780-17.

[9] J. M. Boyce. Quaternary ammonium disinfectants and antiseptics: tolerance, resistance and potential impact on antibiotic resistance. *Antimicrobial Resistance & Infection Control*, 12:32, 2023. doi: 10.1186/s13756-023-01241-z.

[10] S. Bratu, M. Mooty, S. Nichani, D. Landman, C. Gullans, B. Pettinato, U. Karumudi, P. Tolaney, and J. Quale. Emergence of kpc-possessing *Klebsiella pneumoniae* in Brooklyn, New York: Epidemiology and recommendations for detection. *Antimicrobial Agents and Chemotherapy*, 49(7):3018–3020, 2005. doi: 10.1128/aac.49.7.3018-3020.2005.

[11] J. S. Brooke. *Stenotrophomonas maltophilia*: an emerging global opportunistic pathogen. *Clinical Microbiology Reviews*, 25(1):2–41, 2012. doi: 10.1128/cmr.00019-11.

[12] J. S. Brooke. Advances in the microbiology of *Stenotrophomonas maltophilia*. *Clinical Microbiology Reviews*, 34, 05 2021. doi: 10.1128/cmr.00030-19.

[13] E. Brown and E. Hendler. Rhodococcus peritonitis in a patient treated with peritoneal dialysis. *American Journal of Kidney Diseases*, 14(5):417–418, 1989. ISSN 0272-6386. doi: https://doi.org/10.1016/S0272-6386(89)80177-5.

[14] K. Bush and P. A. Bradford. β-lactams and β-lactamase inhibitors: An overview. *Cold Spring Harbor Perspectives in Medicine*, 6(8):a025247, 2016. doi: 10.1101/cshperspect.a025247.

[15] A. Carattoli, E. Zankari, A. García-Fernández, M. Voldby Larsen, O. Lund, L. Villa, F. Møller Aarestrup, and H. Hasman. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrobial Agents and Chemotherapy*, 58(7):3895–3903, 2014. doi: 10.1128/AAC.02412-14.

[16] Q. Chen, W. Lu, D. Zhou, G. Zheng, H. Liu, C. Qian, W. Zhou, J. Lu, L. Ni, Q. Bao, A. Li, T. Xu, and H. Xu. Characterization of two macrolide resistance-related genes in multidrug-resistant *Pseudomonas aeruginosa* isolates. *Polish Journal of Microbiology*, 69(3):349–356, 2020. doi: 10.33073/pjm-2020-038.

[17] I. Chopra and M. Roberts. Tetracycline antibiotics: Mode of action, applications, molecular biology, and epidemiology of bacterial resistance. *Microbiology and Molecular Biology Reviews*, 65(2):232–260, 2001. doi: 10.1128/mmbr.65.2.232-260.2001.

[18] C. Cruz-Cruz, J. Gaytán-Cervantes, C. González-Torres, A. E. Nolasco-Rojas, M. Loyola-Cruz, L. Delgado-Balbuena, J. Delgado-Balbuena, M. Paredes-Mendoza, M. C. Tamayo-Ordóñez, Y. d. J. Tamayo-Ordoñez, E. M. Durán-Manuel, A. Rojas-Bernabé, C. A. Jiménez-Zamarripa, O. Sosa-Hernández, O. A. García-Hernández, E. Ocharan-Hernández, P. B. Zárate-Segura, E. González-Terreros, D. A. Ramírez-Villanueva, C. C. Calzada-Mendoza, and J. M. Bello-López. Profiling of bacterial communities of hospital wastewater reveals clinically relevant genera and antimicrobial resistance genes. *Microorganisms*, 13(6), 2025. ISSN 2076-2607. doi: 10.3390/microorganisms13061316.

[19] P. L. DeMarais and F. E. Kocka. Rhodococcus meningitis in an immunocompetent host. *Clinical Infectious Diseases*, 20(1):167–169, 1995. doi: 10.1093/clinids/20.1.167.

[20] Y. Deng, X.-F. Han, Z.-M. Jiang, L.-Y. Yu, Y. Li, and Y.-Q. Zhang. Characterization of three *Stenotrophomonas* strains isolated from different ecosystems and proposal of *Stenotrophomonas mori* sp. nov. and *Stenotrophomonas lacuserhaii* sp. nov. *Frontiers in Microbiology*, 13, 12 2022. doi: 10.3389/fmicb.2022.1056762.

[21] S. P. Diggle and M. Whiteley. Microbe profile: *Pseudomonas aeruginosa*: Opportunistic pathogen and lab rat. *Microbiology*, 166:30–33, 10 2019. doi: 10.1099/mic.0.000860.

[22] M. D. Ekwanzala, J. B. Dewar, and M. N. B. Momba. Environmental resistome risks of wastewaters and aquatic environments deciphered by shotgun metagenomic assembly. *Ecotoxicology and Environmental Safety*, 197:110612, 07 2020. doi: 10.1016/j.ecoenv.2020.110612.

[23] A. Elfadadny, R. F. Ragab, M. AlHarbi, F. Badshah, E. Ibáñez-Arancibia, A. Farag, A. O. Hendawy, Patricio, M. Aboubakr, S. A. Zakai, and W. M. Nageeb. Antimicrobial resistance of *Pseudomonas aeruginosa*: navigating clinical impacts, current resistance trends, and innovations in breaking therapies. *Front. Microbiol*, 15, 04 2024. doi: 10.3389/fmicb.2024.1374466.

[24] R. Ero, V. Kumar, W. Su, and Y. Gao. Ribosome protection by abc-f proteins—molecular mechanism and potential drug design. *Protein Science*, 28:684–693, 03 2019. doi: 10.1002/pro.3589.

[25] P. Ewels, M. Magnusson, S. Lundin, and M. Käller. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, 32(19):3047–3048, 2016.

[26] N. Farhat and A. U. Khan. Evolving trends of new delhi metallo-betalactamase (ndm) variants: A threat to antimicrobial resistance. *Infection, Genetics and Evolution*, 86:104588, 12 2020. doi: 10.1016/j.meegid.2020.104588.

[27] N. Fayad, M. Kallassy Awad, and J. Mahillon. IS982 and kin: new insights into an old is family. *Mobile DNA*, 11:24, 2020. doi: 10.1186/s13100-020-00221-z.

[28] M. Galarde-López, M. E. Velazquez-Meza, E. E. Godoy-Lozano, B. A. Carrillo-Quiroz, P. Cornejo-Juárez, A. Sassoé-González, A. Ponce-de León, P. Saturno-Hernández, and C. M. Alpuche-Aranda. Presence and persistence of ESKAPEE bacteria before and after hospital wastewater treatment. *Microorganisms*, 12(6):1231, 2024. doi: 10.3390/microorganisms12061231.

[29] S. B. Georghiou, M. Magana, R. S. Garfein, D. G. Catanzaro, A. Catanzaro, and T. C. Rodwell. Evaluation of genetic mutations associated with Mycobacterium tuberculosis resistance to amikacin, kanamycin and capreomycin: A systematic review. *PLoS ONE*, 7(3):e33275, 2012. doi: 10.1371/journal.pone.0033275.

[30] R. Ghosh, S. Chatterjee, and N. C. Mandal. Stenotrophomonas. *Elsevier eBooks*, pages 427–442, 01 2020. doi: 10.1016/b978-0-12-823414-3.00020-4.

[31] L. Goltermann, K. L. Andersen, H. K. Johansen, S. Molin, and R. La Rosa. Macrolide therapy in *Pseudomonas aeruginosa* infections causes ul4 ribosomal protein mutations leading to high-level resistance. *Clinical Microbiology and Infection*, 28(12):1594–1601, 2022. doi: 10.1016/j.cmi.2022.08.003.

[32] S. J. Green, O. Prakash, P. Jasrotia, W. A. Overholt, E. Cardenas, D. Hubbard, J. M. Tiedje, D. B. Watson, C. W. Schadt, S. C. Brooks, and J. E. Kostka. Denitrifying bacteria from the genus Rhodanobacter dominate bacterial communities in the highly contaminated subsurface of a nuclear legacy waste site. *Applied and Environmental Microbiology*, 78(4):1039–1047, 2012. doi: 10.1128/AEM.06435-11.

[33] N. Johnston, T. Mukhtar, and G. Wright. Streptogramin antibiotics: Mode of action and resistance. *Current Drug Targets*, 3:335–344, 08 2002. doi: 10.2174/1389450023347678.

[34] O. B. Jonas, A. Irwin, F. C. J. Berthe, F. G. Le Gall, and P. V. Marquez. Drug-resistant infections : a threat to our economic future (vol. 2) : final report (vol 2). hnp/agriculture global antimicrobial resistance initiative washington, d.c. : World bank group. `http://documents.worldbank.org/curated/en/323311493396993758/final-report`, 2017.

[35] M. Krauland, L. Harrison, D. Paterson, et al. Novel integron gene cassette arrays identified in a global collection of multi-drug resistant non-typhoidal *Salmonella enterica*. *Current Microbiology*, 60(3):217–223, 2010. doi: 10.1007/s00284-009-9527-3.

[36] S. F. Kuo and C. H. Lee. An oil refinery worker at kaohsiung, with *Pseudoxanthomonas kaohsiungensis* bloodstream infection presenting as chronic pericarditis and masquerading as tuberculosis pericarditis. *Journal of Microbiology, Immunology and Infection*, 51(4):575–577, 2018. doi: 10.1016/j.jmii.2017.12.003.

[37] I. Kutilova, M. Medvecky, P. Leekitcharoenphon, P. Munk, M. Masarikova, L. Davidova-Gerzova, I. Jamborova, V. Bortolaia, S. J. Pamp, and M. Dolejska. Extended-spectrum β-lactamase-producing *Escherichia coli* and antimicrobial resistance in municipal and hospital wastewaters in Czech Republic: Culture-based and metagenomic approaches. *Environ. Res.*, 193(110487):110487, 2021. doi: 10.1016/j.envres.2020.110487.

[38] D. Kücken, H. Feucht, and P. Kaulfers. Association of *qacE* and *qacEΔ1* with multiple resistance to antibiotics and antiseptics in clinical isolates of gram-negative bacteria. *FEMS Microbiology Letters*, 183(1):95–98, 2000. doi: 10.1111/j.1574-6968.2000.tb08939.x.

[39] B. Langmead and S. L. Salzberg. Fast gapped-read alignment with bowtie 2. *Nat. Methods*, 9 (4):357–359, 2012.

[40] K. D. Lenz, K. E. Klosterman, H. Mukundan, and J. Z. Kubicek-Sutherland. Macrolides: From toxins to therapeutics. *Toxins*, 13(5):347, 2021. doi: 10.3390/toxins13050347.

[41] G. L. Lewis, R. J. Fenton, E. N. Moriyama, J. D. Loy, and R. A. Moxley. Association of ISVsa3 with multidrug resistance in *Salmonella enterica* isolates from cattle (Bos taurus). *Microorganisms*, 11(3):631, 2023. doi: 10.3390/microorganisms11030631.

[42] X.-Z. Li, P. Plésiat, and H. Nikaido. The challenge of efflux-mediated antibiotic resistance in gram-negative bacteria. *Clinical Microbiology Reviews*, 28(2):337–418, 2015. doi: 10.1128/cmr. 00117-14.

[43] X. Liu, L. Gong, E. Liu, C. Li, Y. Wang, and J. Liang. Characterization of the disinfectant resistance genes qaceδ1 and cepA in carbapenem-resistant *Klebsiella pneumoniae* isolates. *American Journal of Tropical Medicine and Hygiene*, 110:136–141, 01 2024. doi: 10.4269/ajtmh.23-0247.

[44] J. Lu, F. P. Breitwieser, P. Thielen, and S. L. Salzberg. Bracken: estimating species abundance in metagenomics data. *PeerJ Comput. Sci.*, 3(e104):e104, 2017.

[45] Y. Ma, J. Wang, Y. Liu, X. Wang, B. Zhang, W. Zhang, T. Chen, G. Liu, L. Xue, and X. Cui. Nocardioides: "specialists" for hard-to-degrade pollutants in the environment. *Molecules*, 28 (21):7433, 2023. doi: 10.3390/molecules28217433.

[46] K. Manika, F. Kontos, A. Papavasileiou, D. Papaventsis, M. Sionidou, and I. Kioumis. Severe pulmonary disease caused by *Mycolicibacter kumamotonensis*. *Emerging Infectious Diseases*, 27(3):962–964, 2021. doi: 10.3201/eid2703.191648.

[47] M. A. Markkanen, K. Haukka, K. M. M. Pärnänen, V. T. Dougnon, I. J. O. Bonkoungou, Z. Garba, H. Tinto, A. Sarekoski, A. Karkman, A. Kantele, and M. P. J. Virta. Metagenomic analysis of the abundance and composition of antibiotic resistance genes in hospital wastewater in Benin, Burkina Faso, and Finland. *mSphere*, 8(1):e0053822, Feb. 2023.

[48] M. Martin. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.*, 17(1):10, May 2011.

[49] E. A. Miller, E. Elnekave, C. Flores-Figueroa, A. Johnson, A. Kearney, J. Munoz-Aguayo, K. A. Tagg, L. Tschetter, B. P. Weber, C. A. Nadon, D. Boxrud, R. S. Singer, J. P. Folster, and T. J. Johnson. Emergence of a novel *Salmonella enterica* serotype reading clonal group is linked to its expansion in commercial turkey production, resulting in unanticipated human illness in North America. *mSphere*, 5(2), 2020. doi: 10.1128/msphere.00056-20.

[50] W. R. Miller and C. A. Arias. ESKAPE pathogens: antimicrobial resistance, epidemiology, clinical impact and therapeutics. *Nature Reviews Microbiology*, 22:598–616, 2024. doi: 10. 1038/s41579-024-01054-w.

[51] C. D. Moon, W. Young, P. H. Maclean, A. L. Cookson, and E. N. Bermingham. Metagenomic insights into the roles of proteobacteria in the gastrointestinal microbiomes of healthy dogs and cats. *MicrobiologyOpen*, 7, 06 2018. doi: 10.1002/mbo3.677.

[52] A. Moura, M. Soares, C. Pereira, N. Leitão, I. Henriques, and A. Correia. INTEGRALL: a database and search engine for integrons, integrases and gene cassettes. *Bioinformatics*, 25(8): 1096–1098, 2009. doi: 10.1093/bioinformatics/btp105.

[53] T. Naas, P. Nordmann, G. Vedel, and C. Poyart. Plasmid-mediated carbapenem-hydrolyzing β-lactamase kpc in a *Klebsiella pneumoniae* isolate from France. *Antimicrobial Agents and Chemotherapy*, 49(10):4423–4424, 2005. doi: 10.1128/aac.49.10.4423-4424.2005.

[54] S. Navon-Venezia, I. Chmelnitsky, A. Leavitt, M. J. Schwaber, D. Schwartz, and Y. Carmeli. Plasmid-mediated imipenem-hydrolyzing enzyme kpc-2 among multiple carbapenem-resistant *Escherichia coli* clones in Israel. *Antimicrobial Agents and Chemotherapy*, 50(9):3098–3101, 2006. doi: 10.1128/aac.00438-06.

[55] C. Ng, M. Tay, B. Tan, T.-H. Le, L. Haller, H. Chen, T. H. Koh, T. M. S. Barkham, and K. Y.-H. Gin. Characterization of metagenomes in urban aquatic compartments reveals high prevalence of clinically relevant antibiotic resistance genes in wastewaters. *Front. Microbiol.*, 8: 2200, 2017. doi: 10.3389/fmicb.2017.02200.

[56] T. K. Nielsen, P. D. Browne, and L. H. Hansen. Antibiotic resistance genes are differentially mobilized according to resistance mechanism. *GigaScience*, 11, 2022. doi: 10.1093/gigascience/giac072.

[57] N. A. Nor Amdan, N. A. Shahrulzamri, R. Hashim, and N. Mohamad Jamil. Understanding the evolution of macrolides resistance: A mini review. *Journal of Global Antimicrobial Resistance*, 38:368–375, 2024. doi: 10.1016/j.jgar.2024.07.016.

[58] A. Ovung and J. Bhattacharyya. Sulfonamide drugs: structure, antibacterial property, toxicity, and biophysical interactions. *Biophysical Reviews*, 13:259–272, 03 2021. doi: 10.1007/s12551-021-00795-9.

[59] N. Oyenuga, J. F. Cobo-Díaz, A. Alvarez-Ordóñez, and E.-A. Alexa. Overview of antimicrobial resistant ESKAPEE pathogens in food sources and their implications from a one health perspective. *Microorganisms*, 12:2084, 10 2024. doi: 10.3390/microorganisms12102084.

[60] S. R. Partridge, S. M. Kwong, N. Firth, and S. O. Jensen. Mobile genetic elements associated with antimicrobial resistance. *Clinical Microbiology Reviews*, 31(4), 2018. doi: 10.1128/cmr.00088-17.

[61] M. R. Perry, H. C. Lepper, L. McNally, B. A. Wee, P. Munk, A. Warr, B. Moore, P. Kalima, C. Philip, A. M. de Roda Husman, F. M. Aarestrup, M. E. J. Woolhouse, and B. A. D. van Bunnik. Secrets of the hospital underbelly: Patterns of abundance of antimicrobial resistance genes in hospital wastewater vary by specific antimicrobial and bacterial family. *Front. Microbiol.*, 12:703560, 2021. doi: 10.3389/fmicb.2021.703560.

[62] M. L. Petrovich, A. Zilberman, A. Kaplan, G. R. Eliraz, Y. Wang, K. Langenfeld, M. Duhaime, K. Wigginton, R. Poretsky, D. Avisar, and G. F. Wells. Microbial and viral communities and their antibiotic resistance genes throughout a hospital wastewater treatment system. *Frontiers in Microbiology*, 11, 02 2020. doi: 10.3389/fmicb.2020.00153.

[63] D. N. Pham and M. Li. Comparative resistomics analysis of multidrug-resistant Chryseobacteria. *Environmental Microbiology Reports*, 16(3):e13288, 2024. doi: 10.1111/1758-2229.13288.

[64] K. M. M. Pärnänen, C. Narciso-da Rocha, D. Kneis, T. U. Berendonk, D. Cacace, T. T. H. Do, C. Elpers, D. Fatta-Kassinos, I. Henriques, T. Jaeger, A. Karkman, J. L. Martinez, I. Michael, K. O'Sullivan, S. Rodriguez-Mozaz, T. Schwartz, H. Sheng, H. Sorum, R. D. Stedtfeld, J. M. Tiedje, E. Topp, I. Vaz-Moreira, M. P. J. Virta, E. Wellington, H. Sørum, and C. M. Manaia. Antibiotic resistance in European wastewater treatment plants mirrors the pattern of clinical antibiotic resistance prevalence. *Science Advances*, 5(3):eaau9124, 2019. doi: 10.1126/sciadv. aau9124.

[65] J. Qian, Z. Wu, L. Ma, Y. Zhu, Y. Zhang, M. Li, X. Guo, and C. Liu. Metagenomic insights into correlation of microbiota and antibiotic resistance genes in the worker-pig-soil interface: A one health surveillance on Chongming Island, China. *Journal of Hazardous Materials Advances*, 18:100648, 2025. ISSN 2772-4166. doi: https://doi.org/10.1016/j.hazadv.2025.100648.

[66] N. Ranganath, M. A. Mendoza, R. Stevens, D. Kind, N. Wengenack, and A. Shah. *Rhodococcus* infection: a 10-year retrospective analysis of clinical experience and antimicrobial susceptibility profile. *Journal of Clinical Microbiology*, 62, 03 2024. doi: 10.1128/jcm.01537-23.

[67] S. Raza, H. Shin, H.-G. Hur, and T. Unno. Higher abundance of core antimicrobial resistant genes in effluent from wastewater treatment plants. *Water Research*, 208:117882, 2022. ISSN 0043-1354. doi: 10.1016/j.watres.2021.117882.

[68] G. Rizzatti, L. R. Lopetuso, G. Gibiino, C. Binda, and A. Gasbarrini. Proteobacteria: a common factor in human diseases. *BioMed Research International*, 2017:1–7, 11 2017. doi: 10.1155/2017/9351507.

[69] D. M. Salvador-Membreve and W. L. Rivera. Predominance of bla_{TEM} and $tetA$ genes in antibiotic-resistant *Escherichia coli* isolates from Laguna Lake, Philippines. *Journal of Water, Sanitation and Hygiene for Development*, 11:814–823, 07 2021. doi: 10.2166/washdev.2021.067.

[70] S. A. Sapula, A. Amsalu, J. J. Whittall, B. J. Hart, N. L. Siderius, L. Nguyen, C. Gerber, J. Turnidge, and H. Venter. The scope of antimicrobial resistance in residential aged care facilities determined through analysis of *Escherichia coli* and the total wastewater resistome. *Microbiology Spectrum*, 11(6):e00731–23, 2023. doi: 10.1128/spectrum.00731-23.

[71] X. Shuai, Z. Zhou, L. Zhu, C. Achi, Z. Lin, Z. Liu, X. Yu, J. Zhou, Y. Lin, and H. Chen. Ranking the risk of antibiotic resistance genes by metagenomic and multifactorial analysis in hospital wastewater systems. *Journal of hazardous materials*, 468:133790–133790, 04 2024. doi: 10.1016/j.jhazmat.2024.133790.

[72] P. Siguier, J. Perochon, L. Lestrade, J. Mahillon, and M. Chandler. ISFinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Research*, 34, 01 2006. ISSN 0305-1048. doi: 10.1093/nar/gkj014.

[73] C. J. Smith, C. Owen, and L. Kirby. Activation of a cryptic streptomycin-resistance gene in the *Bacteroides erm* transposon, *Tn4551*. *Molecular Microbiology*, 6:2287–2297, 08 1992. doi: 10.1111/j.1365-2958.1992.tb01404.x.

[74] K. Sugita, K. Aoki, K. Komori, T. Nagasawa, Y. Ishii, S. Iwata, and K. Tateda. Molecular analysis of $bla_{\text{kpc-2}}$-harboring plasmids: Tn*4401a* interplasmid transposition and Tn*4401a*-carrying ColRNAI plasmid mobilization from *Klebsiella pneumoniae* to *Citrobacter europaeus* and *Morganella morganii* in a single patient. *mSphere*, 6(6), 2021. doi: 10.1128/mSphere. 00850-21.

[75] I. Sultan, S. Rahman, A. T. Jan, M. T. Siddiqui, A. H. Mondal, and Q. M. R. Haq. Antibiotics, resistome and resistance mechanisms: A bacterial perspective. *Frontiers in Microbiology*, 9: 2066, 2018. ISSN 1664-302X. doi: 10.3389/fmicb.2018.02066.

[76] A. Talat, A. Usmani, and A. U. Khan. Detection of *E. coli* incx1 plasmid-mediated *mcr-5.1* gene in an Indian hospital sewage water using shotgun metagenomic sequencing: A first report. *Microbial Drug Resistance*, 28:759–764, 06 2022. doi: 10.1089/mdr.2021.0338.

[77] A. Talat, K. S. Blake, G. Dantas, and A. U. Khan. Metagenomic insight into microbiome and antibiotic resistance genes of high clinical concern in urban and rural hospital wastewater of northern India origin: A major reservoir of antimicrobial resistance. *Microbiol. Spectr.*, 2023. doi: 10.1128/spectrum.04102-22.

[78] M. Torres-Tortosa, J. Arrizabalaga, J. L. Villanueva, J. Gálvez, M. Leyes, M. E. Valencia, J. Flores, J. M. Peña, E. Pérez-Cecilia, C. Quereda, G. A. para el estudio de las Enfermedades Infecciosas, and G. de estudio de SIDA of the Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Prognosis and clinical evaluation of infection caused by *Rhodococcus equi* in HIV-infected patients: a multicenter study of 67 cases. *Chest*, 123(6):1970–1976, 2003. doi: 10.1378/chest.123.6.1970.

[79] M. E. Velazquez-Meza, M. Galarde-López, P. Cornejo-Juárez, M. Bobadilla-del Valle, E. Godoy-Lozano, E. Aguilar-Vera, B. A. Carrillo-Quiroz, A. Ponce de León-Garduño, C. Velazquez Acosta, and C. M. Alpuche-Aranda. Bacterial communities and resistance and virulence genes in hospital and community wastewater: Metagenomic analysis. *International Journal of Molecular Sciences*, 26(5), 2025. doi: 10.3390/ijms26052051.

[80] M. Venkatesan, M. Fruci, L. A. Verellen, T. Skarina, N. Mesa, R. Flick, C. Pham, R. Mahadevan, P. Stogios, and A. Savchenko. Molecular mechanism of plasmid-borne resistance to sulfonamide antibiotics. *Nature Communications*, 14, 07 2023. doi: 10.1038/s41467-023-39778-7.

[81] J.-I. Wachino, Y. Doi, and Y. Arakawa. Aminoglycoside resistance. *Infectious Disease Clinics of North America*, 34(4):887–902, 2020. doi: 10.1016/j.idc.2020.06.002.

[82] C. Wang, P. Cai, Y. Guo, and Z. Mi. Distribution of the antiseptic-resistance genes $qacE\Delta1$ in 331 clinical isolates of *Pseudomonas aeruginosa* in China. *Journal of Hospital Infection*, 66: 93–95, 05 2007. doi: 10.1016/j.jhin.2007.01.012.

[83] N. Wang, J. Luo, F. Deng, Y. Huang, and H. Zhou. Antibiotic combination therapy: A strategy to overcome bacterial resistance to aminoglycoside antibiotics. *Frontiers in Pharmacology*, 13, 02 2022. doi: 10.3389/fphar.2022.839808.

[84] D. E. Wood, J. Lu, and B. Langmead. Improved metagenomic analysis with Kraken 2. *Genome Biol.*, 20(1):257, 2019.

[85] World Health Organization. WHO's List of Medically Important Antimicrobials: A Risk Management Tool for Mitigating Antimicrobial Resistance Due to Non-Human Use, 2024. URL

`https://www.who.int/publications/i/item/9789240086813`. Licence: CC BY-NC-SA 3.0 IGO.

[86] World Health Organization. WHO bacterial priority pathogens list, 2024: Bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance, 2024. URL `https://www.who.int/publications/i/item/9789240093461`. Licence: CC BY-NC-SA 3.0 IGO.

[87] J. Wu, T. P. Thompson, N. H. O'Connell, K. McCracken, J. Powell, B. F. Gilmore, C. P. Dunne, and S. A. Kelly. Extended-spectrum β-lactamase-producing bacteria from hospital wastewater pipes: isolation, characterization and biofilm control using common disinfectants. *Journal of Hospital Infection*, 156:34–49, 2025. ISSN 0195-6701. doi: 10.1016/j.jhin.2024.11.013.

[88] W.-Y. Xie, Y.-T. Wang, J. Yuan, W.-D. Hong, G.-Q. Niu, X. Zou, X.-P. Yang, Q. Shen, and F.-J. Zhao. Prevalent and highly mobile antibiotic resistance genes in commercial organic fertilizers. *Environment International*, 162:107157, 04 2022. doi: 10.1016/j.envint.2022.107157.

[89] X. Yin, X. Chen, X.-T. Jiang, Y. Yang, B. Li, M. H.-H. Shum, T. T. Y. Lam, G. M. Leung, J. Rose, C. Sanchez-Cid, T. M. Vogel, F. Walsh, T. U. Berendonk, J. Midega, C. Uchea, D. Frigon, G. D. Wright, C. Bezuidenhout, R. C. Picão, S. Z. Ahammad, P. H. Nielsen, P. Hugenholtz, N. J. Ashbolt, G. Corno, D. Fatta-Kassinos, H. Bürgmann, H. Schmitt, C.-J. Cha, A. Pruden, K. Smalla, E. Cytryn, Y. Zhang, M. Yang, Y.-G. Zhu, A. Dechesne, B. F. Smets, D. W. Graham, M. R. Gillings, W. H. Gaze, C. M. Manaia, M. C. M. van Loosdrecht, P. J. J. Alvarez, M. J. Blaser, J. M. Tiedje, E. Topp, and T. Zhang. Toward a universal unit for quantification of antibiotic resistance genes in environmental samples. *Environ. Sci. Technol.*, 57(26):9713–9721, 2023.

[90] A. N. Zhang, J. M. Gaston, C. L. Dai, S. Zhao, M. Poyet, M. Groussin, X. Yin, L.-G. Li, M. C. M. van Loosdrecht, E. Topp, M. R. Gillings, W. P. Hanage, J. M. Tiedje, K. Moniz, E. J. Alm, and T. Zhang. An omics-based framework for assessing the health risk of antimicrobial resistance genes. *Nature Communications*, 12:4765, 08 2021. doi: 10.1038/s41467-021-25096-3.

[91] H. Zhang, Z. Zong, S. Lei, S. Srinivas, J. Sun, Y. Feng, M. Huang, and Y. Feng. A genomic, evolutionary, and mechanistic study of mcr-5 action suggests functional unification across the mcr family of colistin resistance. *Advanced Science*, 6, 04 2019. doi: 10.1002/advs.201900034.

[92] S. Zhang, J. Huang, Z. Zhao, Y. Cao, and B. Li. Hospital wastewater as a reservoir for antibiotic resistance genes: A meta-analysis. *Front. Public Health*, 8:574968, Oct. 2020.

[93] G. Zheng, T. F. Webster, and A. Salamova. Quaternary ammonium compounds: Bioaccumulation potentials in humans and levels in blood before and during the Covid-19 pandemic. *Environmental Science & Technology*, 55(21):14689–14698, 2021. doi: 10.1021/acs.est.1c01654.

[94] L. Zhu, L. Yuan, X.-Y. Shuai, Z.-J. Lin, Y.-J. Sun, Z.-C. Zhou, L.-X. Meng, F. Ju, and H. Chen. Deciphering basic and key traits of antibiotic resistome in influent and effluent of hospital wastewater treatment systems. *Water Research*, 231:119614, 03 2023. doi: 10.1016/j.watres.2023.119614.