Synergistic Enhancement of Acetamiprid Toxicity by Herbicide and Fungicide: Triple Mixture Outperforms Binary Combination

Jan Erik Sedlmeier[1*], Emma Gamer[1], Birgit Höglinger[2], Frank Walker[2], Carsten A Brühl[3], Georg Petschenka[1,4*]

[1]Department of Applied Entomology, University of Hohenheim, Stuttgart, Germany
[2]Institute of Phytomedicine, University of Hohenheim, Stuttgart, Germany
[3]Community Ecology & Ecotoxicology, University of Kaiserslautern-Landau, Landau, Germany
[4]KomBioTa – Center for Biodiversity and Integrative Taxonomy, University of Hohenheim and State Museum of Natural History, Stuttgart, Germany

[*]**Corresponding authors**
Jan Erik Sedlmeier
Department of Applied Entomology, Institute of Phytomedicine, University of Hohenheim, Stuttgart, Germany; jan.sedlmeier@uni-hohenheim.de
Georg Petschenka
Department of Applied Entomology, Institute of Phytomedicine, University of Hohenheim, Stuttgart, Germany; georg.petschenka@uni-hohenheim.de

Author ORCiDs
Jan Erik Sedlmeier: 0000-0002-4694-3894
Emma Gamer: 0009-0003-4241-7840
Frank Walker: 0009-0001-3738-4840
Carsten A. Brühl: 0000-0003-1332-535X
Georg Petschenka: 0000-0002-9639-3042

Keywords
synergism, terbuthylazine, fluopyram, neonicotinoids, non-target, pesticide mixtures

Acknowledgments
We would like to thank Sarah Rißmann for her consistent support throughout the laboratory work and insect rearing. We would also like to thank the Graduate Academy at the University of Hohenheim for making this project possible by awarding a scholarship as part of the Baden-Württemberg Graduate Support Programme to JES.

Abstract

Pesticides are considered one of the main drivers of insect decline, and complex pesticide mixtures are frequently detected in our landscapes. However, the risk posed by mixtures of active substances is not systematically assessed. In this study using acute contact dose-response testing, we show that binary and triple combinations of relevant surrogates for major pesticide classes can lead to synergistic toxic effects on the non-target insect *Oncopeltus fasciatus*. First, we tested the individual active substances acetamiprid (insecticide), fluopyram (fungicide) and terbuthylazine (herbicide), detecting acute toxic effects only for acetamiprid. However, the combination of acetamiprid and terbuthylazine significantly increased toxicity compared to acetamiprid alone and the combination of acetamiprid and fluopyram with the effective dose 50 being 2.7 and 2.3 times lower respectively. Strikingly, a triple mixture of all active substances exceeded the toxicity of any other treatment. While the exact mechanisms remain unclear, we speculate that increased physiological stress contributes to the observed synergistic effects, as terbuthylazine and fluopyram have been shown to interfere, though to varying extents, with cellular energy metabolism, oxidative balance, detoxification and neurotransmission in several non-target organisms. These results are concerning and highlight the ecotoxicological uncertainty posed by traces of complex pesticide mixtures present in the air, soil, water and plants across landscapes. We therefore propose a systematic assessment of realistic exposure scenarios reflecting both agricultural practices and observed environmental mixtures.

1. Introduction

Pesticide contamination of the environment caused by intensified agriculture is considered to be one of the major drivers of the unprecedented decline in insect biomass, species richness and abundance (Geiger et al. 2010; Sánchez-Bayo and Wyckhuys 2019; Wagner et al. 2021; Dicks et al. 2021). Several recent studies document widespread contamination of landscapes with mixtures of agrochemicals (Silva et al. 2023; Brühl et al. 2024; Mauser et al. 2025). These mixtures, present in different matrices (e.g. soil, vegetation, water), consist of varying combinations of active substances, often representing several major pesticide classes, such as insecticides, herbicides, and fungicides, simultaneously. Such mixtures are found not only in agricultural areas but also in nature conservation zones and even on remote mountain peaks. A recent study found 140 individual mixtures of active substances (containing two or more components) in soil, vegetation and water samples from the Upper Rhine Valley in Germany (Mauser et al. 2025). In these mixtures, individual detected amounts of active substances add up to considerable total concentrations (see e.g. Brühl et al. 2024; Honert et al. 2025; Mauser et al. 2025). Another study detected a total of 109 individual pesticides in the ambient air across Germany. A median of 19 substances was detected per sampling site, with a maximum of 36 substances. The sampling sites included forests and national parks and the substances detected were herbicides, fungicides, insecticides, repellents, growth

regulators, and metabolites of the aforementioned substances (Kruse-Plaß et al.
2021). Based on these measurements, Gandara et al. (2024) exposed *Drosophila*
melanogaster larvae to nine pesticides at environmentally relevant concentrations.
Although this mixture did not include any insecticides, this treatment resulted in
behavioral changes and reduced egg-laying capacity of females. These results
suggest that pesticide mixtures at sublethal, environmentally relevant concentrations
can impact the fitness of insect populations.

Current risk assessment of plant protection products in the EU focuses on individual
active substances and formulated products. Some mixtures are tested as part of the
formulation evaluation process. These formulations may contain more than one
active substance (e.g. two fungicides with different modes of action for resistance
management). In addition, synergistic effects of certain combinations of pesticide
classes, such as neonicotinoids or pyrethroids in combination with azole fungicides,
on pollinator toxicity are known and therefore considered in application regulations
(i.e. tank-mix regulations). However, there is currently no systematic assessment of
the ecological risk posed by mixtures of active substances. This is alarming given the
425 active substances currently approved in the EU (European Commission n.d.)
and the resulting potential combinations, as well as the numerous mixtures that have
been detected in the environment. Similarly, most ecotoxicological research is also
focused on the effects of individual substances with a bias towards pollinators and
neglecting herbivorous insects. The result is a major knowledge gap regarding
combined toxicity of active substances towards non-target insects (Tosi et al. 2022).

In this study, we address this research gap by testing the effects of a combination of
three active ingredients as surrogates for the major pesticide classes of (1)
insecticides, (2) herbicides and (3) fungicides.

As representatives, we chose the active substances acetamiprid (insecticide, CAS:
135410-20-7), fluopyram (fungicide, CAS: 658066-35-4) and terbuthylazine
(herbicide, CAS: 5915-41-3). The substances were chosen based on their
agricultural but also ecotoxicological relevance as detailed below.
Acetamiprid is a neurotoxic neonicotinoid insecticide that can act as a contact or
systemic insecticide. It binds to the post-synaptic nicotinic acetylcholine receptor,
disrupting neural transmission (IRAC Class 4A; IRAC n.d.). We chose acetamiprid
because it is the last member of the controversial group of neonicotinoids still
registered for field use in the EU and used in arable crops, fruit- and wine growing.
Although it is less harmful to pollinators than the previously banned neonicotinoids
imidacloprid, thiacloprid, clothianidin and thiamethoxam, recent studies have raised
concerns about the environmental safety of acetamiprid regarding non-target insect
toxicity (Varga-Szilay and Tóth 2022; Sedlmeier et al. 2025). In addition, acetamiprid
was detected in environmental samples in Europe, including protected areas (Brühl
et al. 2024; Mauser et al. 2025). It is also known that a combination of neonicotinoids
and certain fungicides can result in synergistic toxicity, meaning that the combined
effect exceeds the sum of the individual toxic effects. This is particularly true for

fungicides from the class of sterol biosynthesis inhibitors (SBIs; Sgolastra et al.
2017; Schuhmann et al. 2022). However, a recent study by Manzer et al. (2024)
demonstrated that even non-SBI fungicides, can exhibit synergistic effects when
combined with acetamiprid, leading to enhanced toxicity in non-target insects.

Among the agrochemical residues detected in landscapes, the fungicide fluopyram
and the herbicide terbuthylazine were often the most frequently detected
substances. In addition, fluopyram is often found at comparably high concentrations
in vegetation, soil and water (Honert et al. 2025; Mauser et al. 2025).
Fluopyram inhibits the enzyme succinate dehydrogenase (SDH) in the mitochondria
of eucaryotic cells. SDH is a membrane-bound protein complex that is part of the
citric acid cycle and the respiratory chain (also referred to as complex II). It is
therefore directly involved in the electron transport chain of the inner mitochondrial
membrane (FRAC Class C2; Sun et al. 2005; FRAC 2025). Besides its function as a
fungicide, it is also used as a nematicide, with the same mode of action (Schleker et
al. 2022). Terbuthylazine blocks electron transport in photosystem II of plant cells,
leading to an accumulation of reactive oxygen that ultimately destroys cell
membranes (HRAC Class C1; HRAC 2025).
We used the large milkweed bug *Oncopeltus fasciatus* (Dallas, 1852) (Hemiptera:
Heteroptera: Lygaeinae) as a model for a herbivorous non-target insects to test for
individual and combined effects of these three active substances in dose-response
assays. *O. fasciatus* is closely associated with milkweed (*Asclepias*, Apocynaceae),
on whose seeds it feeds to sequester cardiac glycosides (Duffey et al. 1978).
Nevertheless, it is also capable of feeding on unrelated, non-toxic plants (Ralph
1976). It can be easily reared in the laboratory on sunflower seeds throughout the
year, allowing for a large sample size. In addition, this study does not aim to replicate
a specific local non-target scenario, but to investigate the fundamental issue of
combined toxicity of several classes of pesticides.

We hypothesized that combinations of an insecticide with either a herbicide, a non-
SBI fungicide, or both, exert synergistic acute toxicity on the non-target insect
Oncopeltus fasciatus, leading to effects stronger than those of the individual
compounds alone.

2. Methods

2.1 Bug rearing

O. fasciatus were reared in a climate chamber (APT.line[TM] KBF-ICH, BINDER
GmbH, Tuttlingen, Germany) under long day conditions (16/8h light/dark cycle) at 27
°C and 60%rH. Nymphs were kept in 5.8 l plastic boxes covered with an insect
mesh. The bottom of the box was covered with kitchen paper. Tap water was
provided in a 15 ml centrifuge tube with a cotton plug in the lid and organic sunflower
seeds were provided ad libitum as food. A piece of cotton wool was provided for egg
laying.

2.2 Active substances

Analytical grade standards of acetamiprid, fluopyram and terbuthylazine were obtained commercially (acetamiprid: HPC Standards GmbH, Borsdorf, Germany; fluopyram/terbuthylazine: Merck KGaA, Darmstadt, Germany). The compounds were always stored in the dark at 4 °C in their original packaging.

The effect of the three active substances on *O. fasciatus* nymphs was first tested individually (see *Individual substance applications*). In parallel, all possible binary mixtures of the substances and the triple mixture were tested (*Mixed substance applications*).

2.3 Individual substance applications

For each active substance (a.s.), a permitted field concentration of a.s. derived from approved formulated products in the Germany (as of May 2023, see Supplementary Information Table 1) was used as the highest concentration in the dose-response assays. In crop protection, the amount of permitted plant protection product is fixed to a specific rate per hectare, but the water volume used for dilution can vary within a specified range (e.g. 200 to 400 liters per hectare). For fluopyram and terbuthylazine, the lowest volume of water has been used in the calculation to imitate the highest permitted amount of a.s./volume. However, the upper volume limit was used for acetamiprid because acetone was to be used as the common solvent in all assays and acetamiprid has a low reported solubility in acetone (~ 0.16 g/l).

For each a.s., a stock solution equivalent to the permitted field concentration was prepared. Fluopyram and terbuthylazine were dissolved in pure acetone, while acetamiprid was dissolved in a 9:1 acetone:deionized water mixture. The resulting stock concentrations were 83.2 ng/µl (acetamiprid), 990.0 ng/µl (fluopyram) and 3694.0 ng/µl (terbuthylazine). Detailed information on the calculations and volumes used is provided in Supplement Information Tables 2-4. Stock solutions were stored in transparent glass vials sealed with metal screw caps and kept in the dark at 4 °C. A dose-range series of six concentrations was prepared for each a.s., beginning with the stock solution and diluting each subsequent concentration by a factor of three (see table 1). In addition, solvent controls were prepared for each a.s. (fluopyram/terbuthylazine: pure acetone; acetamiprid: 9:1 acetone:deionized water). Dilutions were made by pipetting 250 µl of the respective stock into 1.5 ml ND8 amber glass screw-neck vials (LABSOLUTE—TH GEYER) pre-filled with 500 µl of pure acetone. This step was repeated to generate the full dose-range series, which was prepared once per substance and used for all individual substance assays. Each vial was thoroughly vortexed immediately after every dilution step.

Table 1 Dilution series of acetamiprid, fluopyram and terbuthylazine used in the *O. fasciatus* individual substance dose-response assays. The highest dose corresponds to the permitted field concentrations of terbuthylazine and fluopyram, based on approved formulated products in Germany (as of May 2023). For

208 acetamiprid, the field-use concentration could not be reached because of its lower
209 solubility in acetone compared to water, resulting in a lower applied concentration.

Field Concentration Equivalent [%]	Active Substance [ng/µl]		
	Acetamiprid	Fluopyram	Terbuthylazine
100	83.2	990.0	3694.0
33.3	27.7	329.7	1230.1
11.1	9.2	109.9	410.0
3.7	3.1	36.6	136.7
1.24	1.0	12.3	45.8
0.41	0.3	4.1	15.1
0.0	0.0	0.0	0.0

For each dose-response assay, L4 nymphs (i.e. 4th instar nymphs) of *O. fasciatus*
were isolated from the rearing box and transferred to Petri dishes. Six nymphs were
randomly assigned to one of the six treatments (i.e. concentrations) and placed in
individual 90 mm Petri dishes. Immediately before topical pesticide application, the
nymphs were anesthetized with CO_2. Then, 1 µl of a.s. solution (or pure solvent for
the controls) was applied dorsally on the abdomen of each nymph using a Hamilton
repeating dispenser with a 50 µl Hamilton syringe (Hamilton Bonaduz AG,
Switzerland). All a.s. solutions were thoroughly vortexed immediately before drawing
the solution into the syringe. Application started with the control followed by
increasing concentrations of the a.s. solutions. The syringe was rinsed six times with
pure acetone between concentrations. After the droplet had completely evaporated
from the bug's body surface and the nymphs had recovered from anesthesia, each
bug was transferred to an individual 55 mm petri dish containing filter paper at the
bottom, sunflower seeds and water in a 0.5 ml Eppendorf tube plugged with cotton
wool. All Petri dishes within the same a.s. assay were randomized and placed in the
same climate chamber under the conditions described above. The number of
concentrations, individuals and cohorts tested are detailed in Table 2.

Table 2 Overview of *O. fasciatus* assays performed for individual active
substances assays. The concentrations tested are listed in Table 1. For the
acetamiprid individual assay, 12 individuals per concentration were tested in the first
cohort, but only 6 in the second due to greater initial availability of test individuals.
This reduced sample size was maintained for the fluopyram and terbuthylazine
assays. Abbreviations: A = acetamiprid, F = fluopyram, T = terbuthylazine. The

Assay (Individual)	Acetamiprid	Fluopyram	Terbuthylazine	Number of Concentrations Within Dose Range	Individuals Tested per Concentration	Total Number of Individuals Tested	Number of Cohorts
A	dose range	–	–	7	12/6	126	2
F	–	dose range	–	7	6	84	2
T	–	–	dose range	7	6	84	2

234 concentrations tested are listed in Table 1.

2.4 Mixed substance applications

Dose-response assays with a.s. mixtures were conducted for the following
combinations: (1) acetamiprid + fluopyram (AF), (2) acetamiprid + terbuthylazine
(AT), (3) acetamiprid + fluopyram + terbuthylazine (AFT, Table 3). Each assay
consisted of the six concentration levels of acetamiprid already used in the individual
experiments, to which the highest concentration of either fluopyram (990.0 ng/ µl),
terbuthylazine (3694.0 ng/ µl) or a combination of the two from the individual
experiments was added. The concentrations of fluopyram and terbuthylazine were
selected because they correspond to the maximum levels permitted for field
application in Germany and showed no effect on *O. fasciatus* in the individual dose-
response assays (see Results). A solvent only control (pure acetone) was also
included.

To prepare the binary substance combinations (AF and AT), an acetamiprid stock
solution was first made at twice the concentration used in the individual substance
assay (166.4 ng/µl, i.e. 2 x the permitted field concentration), maintaining a 9:1 ratio
of acetone and deionized water. A dose-range series of six concentrations, starting
from and including the acetamiprid stock solution, was then prepared as described
for the individual substance assays. Consequently, all six concentrations in this
dilution series were initially twice as high as those required for testing. In parallel,
separate stock solutions of fluopyram and terbuthylazine were prepared in pure
acetone at twice their respective maximum permitted field concentrations (i.e. 1980.0
and 7388.0 ng/µl). For each dose of the binary mixtures, 50 µl of one of the six
acetamiprid dilutions was pipetted into an empty 1.5 ml ND8 amber glass vial,
followed by 50 µl of the fluopyram or terbuthylazine stock solution. Mixing equal
volumes (2x 50 µl) yielded the same dose range of acetamiprid as was used in the
single substance assay combined with the maximum permitted field concentration of
either fluopyram or terbuthylazine.

For the triple combination (AFT), an acetamiprid stock solution was first prepared at
three times the concentration used in the individual substance assay (249.6 ng/ul,
i.e. 3 x permitted field concentration), maintaining a 9:1 acetone-to-deionized water
ratio. A six-point dose-range series, starting from and including this stock solution,
was then prepared as described for the individual substance assays. As a result, all
concentrations in this series were initially three times higher than the final required
levels. In parallel, fluopyram and terbuthylazine stock solutions were prepared in
pure acetone at 3 x their maximum field concentrations (2970.0 ng/µl and 11,082.0
ng/µl, respectively). For each dose, 50 µl of each of the six acetamiprid dilutions was
pipetted into an empty 1.5 ml ND8 amber glass vial, followed by 50 µl each of the
fluopyram and terbuthylazine stock solutions. Mixing equal volumes (3 x 50 µl)
yielded the intended acetamiprid dose range, each combined with the maximum
permitted field concentration of fluopyram and terbuthylazine.

The double- and triple-concentrated acetamiprid dose-range series, along with the
fluopyram and terbuthylazine stock solutions, were used for all assays and stored in
the dark at 4°C. Mixtures of A + F, A + T or A + F + T were freshly prepared for each
assay.
Topical application of pesticide solutions on L4 *O. fasciatus* nymphs was carried out
as described for the single substance assays above. Again, Petri dishes containing
L4 nymphs were randomized within the same assay and maintained in the climate
chamber under the same conditions. Each concentration in the mixture assays (and
a control of pure acetone) was tested on six individuals per acetamiprid
concentration and each experiment was replicated three times (n = 126 in total).
In addition to the dose-range series of acetamiprid supplemented with fluopyram or
terbuthylazine, a mixture containing only fluopyram and terbuthylazine (each at
100% field application rate) was tested in a supplementary experiment with reduced
replication. A volume of 10 µl of each double-concentrated stock solution (200% field
concentration) was pipetted into a 1.5 ml ND8 glass vial, resulting in a 100% field
concentration FT mixture. Six L4 nymphs were treated with 1 µl of the mixture and
six with 1 µl of pure acetone as a control. The application was conducted as
described for the individual assays above. Petri dishes containing nymphs were
randomized and stored in the climate chamber under the conditions described
above.
The number of concentrations, individuals and cohorts tested in all mixed substance
assays are detailed in Table 3.
297
298 **Table 3 Overview of *O. fasciatus* assays performed for mixed active substance**
299 **assays.** A "fixed dose" indicates that the corresponding maximum permitted field
300 concentration was used. Abbreviations: A = acetamiprid, F = fluopyram, T =
301 terbuthylazine.

Assay (mixture)	Acetamiprid	Fluopyram	Terbuthylazine	Number of Concentrations Within Acetamiprid Dose Range	Individuals Tested per Concentration	Total Number of Individuals Tested	Number of Cohorts
AF	dose range	fixed dose	–	7	6	126	3
AT	dose range	–	fixed dose	7	6	126	3
AFT	dose range	fixed dose	fixed dose	7	6	126	3
FT	–	fixed dose	fixed dose	–	6	12	1

2.5 Evaluation of toxic effects

Nymphs were evaluated 96 hours after application. Petri dishes were removed from
the climate chamber and placed in the laboratory. Each nymph was classified into
one of three categories: dead, affected or unaffected. Classification was aided by
gently touching the abdomen with a fine brush. Healthy individuals typically
responded with a flight reaction or flinching. Affected individuals reacted slowly, if at
all, often only moving a few legs or antennae, or leaning to one side. Nymphs that

appeared dead (typically lying on their back with bent legs and dark discoloration)
were further tested with the brush to confirm the absence of any response. This
evaluation scheme was applied consistently across all assays. Raw data for all
treatments is provided in Supplementary Data.

2.6 Statistical analysis and graphical representation

Using R (version 4.3.0; R Core Team 2023) and R-Studio (version 2023.6.1.524;
Posit team 2023) and the 'drc' package (Ritz et al. 2015), a four-parameter log-
normal cumulative dose-response model, assuming a binomial distribution, was fitted
to the response data at 96 h post application for the data from the individual
acetamiprid assay and from the mixed assays AF and AT.
The following data were not included in the model: (1) data from the individual
fluopyram and terbuthylazine assays as no effects were observed; (2) data from the
fluopyram and terbuthylazine mixed assay (FT) as no effects were observed; (3)
data from the acetamiprid, fluopyram and terbuthylazine triple mixed assay (AFT) as
the effects were too strong across the dose range (see Results: Mixed substance
assays).
The lower asymptote of the dose-response model was set to '0' because in all
included assays, none of the control animals showed any effects or mortality. The
upper asymptote was set to '1'. The *plot()* function was used to generate dose-
response curves. Model fit was assessed by visual inspection of the regression
curve and comparison of the Akaike Information Criterion (AIC) values. LD_{50} values
(and respective 95% confidence intervals) were derived from the model using the
ED() function. The *compParm()* function was used to assess differences in the dose
required to affect 50% of the tested population (ED_{50} value) The function compares
individual curves (i.e., ED_{50} values) within the model and performs approximate t-
tests with $\alpha = 0.05$. Note that all derived ED_{50} values, including those from the mixed
assays, refer to the respective acetamiprid concentration in ng/µl. The full R-code is
provided in Supplementary Information – Supplementary Code.
Figure 1, including the dose response curves and mean data points from the AFT
treatment, was created in R and later edited in Inkscape (version 1.4; Inkscape
Project 2020). Figure 2 was created with JMP® Student Edition (version 18.2.2; JMP
Statistical Discovery LLC 2025) and later edited with Inkscape.

3 Results

3.1 Individual substance assays

Oncopeltus fasciatus nymphs showed high sensitivity to the insecticide acetamiprid
96 hours after application. At all concentrations tested, dead or affected nymphs
were observed. However, at no concentration 100% mortality was observed in the
two cohorts tested. The derived ED_{50} for acetamiprid is 1.61 ng/µl (combined '*dead*
or affected' data) based on a multiple treatment cumulative dose-response model
with a log-normal distribution (Fig. 1).

No effects were observed across concentrations in the single-substance dose-
response assays with fluopyram and terbuthylazine 96 hours after application, and
no dose-response curves were fitted. However, there were isolated cases of
mortality in both experiments. Two nymphs died in the fluopyram assay, one at
33.3% maximum field concentration equivalent (1st cohort) and one at 11.1% (2nd
cohort). In the terbuthylazine assay, four nymphs died, two in the control (one in
each cohort), one at 3.7% field concentration equivalent (1st cohort) and one at
1.24% (2nd cohort). However, in both assays, there were no nymphs categorized as
affected in addition to those that died, supporting that these were random mortality
events and these values were not considered further.

3.2 Mixed substance assays

The effect of the AF mixed treatment (insecticide + fungicide) on the nymphs
compared to the individual acetamiprid application was not significantly different
based on the cumulative dose-response model followed by an ED_{50} comparison
(t=0.58, p=0.563; Fig. 1). However, sensitivity towards the AT mixed application
(insecticide + herbicide) compared to both acetamiprid alone and the AF treatment
was significantly higher based on the ED_{50} comparison (A-AT: t=2.73, p=0.06; AF-AT:
t=2.79, p=0.005). The ED_{50} of AT (0.59 ng/µl) was approximately 2.7-fold lower than
that of acetamiprid alone (1.61 ng/µl), and 2.3-fold lower than that of AF (1.36 ng/µl).
The strongest effects were observed in the AFT treatment (insecticide + fungicide +
herbicide). However, the data could not be included in the dose-response model
because the observed effects were too pronounced across the entire dose range. At
the lowest tested dose (0.34 ng acetamiprid /µl), the mean response was 83.3%,
exceeding the inflection point of a typical dose-response curve. As a result, no curve
could be fitted. Nonetheless, this suggests that the ED_{50} for the AFT treatment would
be well below 0.34 ng/µl.

Derived ED_{50} Values

Treatment	ED_{50} [ng/µl]	95% CI (lower - upper)
A	1.61[a]	0.92 - 2.30
AF	1.36[a]	0.87 - 1.84
AT	0.59[b]	0.36 - 0.82
AFT	<< 0.34	—

Pairwise ED_{50} Comparisons

Treatment Contrast	Estimate	Standard Error	t-value	p-value
A-AF	0.25	0.43	0.58	0.563
A-AT	1.01	0.37	2.73	**0.006**
AF-AT	0.76	0.27	2.79	**0.005**

Fig. 1 *Oncopeltus fasciatus* dose-response curves for individual and mixture assays and comparison of derived ED_{50} values. A**:** acetamiprid individual assay, AF: acetamiprid + fluopyram, AT: acetamiprid + terbuthylazine, AFT: acetamiprid + fluopyram + terbuthylazine. Dose-response relationships were modelled simultaneously using a four-parameter log-normal function. Black symbols represent treatment-specific mean responses per concentration, with grey error bars indicating standard errors (SE). SE indicators were offset in cases of overlap to facilitate interpretation. Red data points (and respective SE indicators) represent mean responses to the AFT treatment but were not included in the dose-response model due to convergence failure. Derived ED_{50} values and respective 95% confidence intervals (CI) for each treatment included in the model are listed in the top right table. For AFT, the ED_{50} is reported as << 0.34 because 83.3% mean response was observed already at the lowest tested concentration (0.34 ng/µl), indicating that the actual ED_{50} must lie well below this value. LD_{50} values derived from the model were compared using the *compParm* function of the *drc* package in R. LD_{50} values of treatments not sharing a common letter are significantly different. A statistical report on the pairwise comparisons is given in the bottom right table with bolt p-values indicating significant differences at α = 0.05. All numbers presented in the adjacent tables are rounded. For all assays, N = 18 biological replicates (nymphs) per concentration

When differentiating between dead and merely affected individuals, mortality in the AFT and AT assay was proportionately higher at lower acetamiprid doses than in the other treatments (Fig. 2). Strikingly, in the AFT treatment, even at the lowest concentration, 67% of nymphs died within 96 hours, compared to 17% in the AT, 0% in the AF, and 4% in the assay with acetamiprid only.
No effects on *O. fasciatus* nymphs were observed in the combined FT treatment (fungicide + herbicide, each at 100% field concentration, see Supplementary Data) 96 hours after application. All nymphs were unaffected in both the treatment and the control group.

Fig. 2 Response of *O. fasciatus* nymphs to individual and mixed substance treatments containing acetamiprid. Proportions of response types are indicated by different colors for each acetamiprid concentration tested. Numbers inside or left of the bars indicate percentage response (rounded). For all assays, N = 18 biological replicates (nymphs) per concentration

4 Discussion

To test the hypothesis that a combination of an insecticide with either a herbicide, a non-SBI fungicide, or both exerts synergistic acute toxicity on the non-target insect *O. fasciatus*, we investigated mixtures of acetamiprid, terbuthylazine, and fluopyram, all of which are widely used in agriculture. First, we assessed each substance individually in dose-response assays, followed by binary mixtures and finally a combined mixture of all three.

4.1 Synergistic Effects

Previous studies on the combined effects of major pesticide classes have primarily focused on interactions between insecticides and fungicides, with most research being conducted on bees (Tosi et al. 2022). A well-documented example is the synergistic interaction between neonicotinoids such as acetamiprid and SBI fungicides. SBIs have been shown to interfere with insect detoxification pathways (e.g., P450 enzymes), thereby enhancing the toxicity of insecticides (Sgolastra et al. 2017; Schuhmann et al. 2022). In Germany, this has led to additional regulations regarding tank mixtures of certain pesticides in order to protect bees (BVL 2018). However, recent evidence suggests that synergistic interactions may also occur with non-SBI fungicides (Manzer et al. 2024), although the underlying mechanisms remain less well understood. Our results support and extend this perspective.

While our individual substance assays revealed acute toxicity only for the insecticide acetamiprid and not for the fungicide fluopyram (non-SBI fungicide) or the herbicide terbuthylazine, certain mixtures of these compounds induced synergistic effects on the non-target insect *O. fasciatus*. This was observed with the binary mixture of acetamiprid and terbuthylazine, as well as with the mixture of all three compounds. Formal synergy modelling was not possible due to the absence of individual activity of the added compounds and the lack of a complete dose-dose matrix. However, given the significant reduction in ED_{50} (A: 1.61 ng/µl, AT: 0.59 ng/µl) and the drastic effects observed in the triple combination (AFT), although neither fluopyram nor terbuthylazine had an effect on their own, we interpret the enhanced acute toxicity as resulting from a synergistic effect.

These findings support existing indications that synergistic interactions may not be limited to specific pesticide groups such as SBI fungicides. Rather, they could emerge more broadly, including from combinations involving herbicides, a pesticide class less commonly associated with synergism. In addition, complex mixtures (i.e., those containing three or more compounds) appear to be especially problematic due to their unpredictable interaction potential. In the following sections, we focus on

terbuthylazine and fluopyram individually in the context of their combinations with
acetamiprid. We then discuss their potential joint contribution to the pronounced
toxicity observed in the triple mixture.

4.2 Terbuthylazine

Herbicides have been shown to exert direct detrimental effects on non-target
organisms, including invertebrates (Wan et al. 2025). Moreover, by eliminating food
plants, herbicides also impose indirect negative impacts on invertebrate populations
(Bohan et al. 2005). However, the mode of action of terbuthylazine does not initially
suggest any potential effect on insects, since the target is exclusively found in plants.
It blocks electron transport in the photosystem II complex by binding to the QB-
binding site of the D1 protein, thereby interrupting the electron transport chain.
However, although comprehensive data are limited, there are individual reports of
adverse effects of terbuthylazine on both invertebrates and vertebrates.
Terbuthylazine can affect the antioxidant defense systems of fish and induce
oxidative stress (Semren et al. 2018). A terbuthylazine-containing herbicide
formulation caused acute toxic effects on several soil invertebrates, including
enchytraeids and mites, despite the pure active ingredient showing no toxicity under
the same conditions (Salminen et al. 1996). Notably, no insects were included in this
study.
Aside from our study, we are not aware of any published evidence indicating that
terbuthylazine can increase the toxicity of insecticides in insects. However,
synergistic toxicity has been reported for another s-triazine herbicide, atrazine, in the
midge *Chironomus tentans*, when mixed with organophosphate insecticides
(Pape□Lindstrom and Lydy 1997). The authors speculated that this effect may be
linked to an enhanced metabolic activation of the organophosphates, as most
organophosphorus insecticides would require metabolic activation to effectively
inhibit the acetylcholinesterase. Since s-triazine herbicides like terbuthylazine were
shown to induce the cytochrome P450 machinery (Kao et al. 1995), the authors
suggested that such herbicides might accelerate the metabolization of
organophosphates and thus results in increased toxicity.

However, acetamiprid does not require metabolic activation to effectively inhibit the
nicotinic acetylcholine receptor. Given the known role of cytochrome P450 enzymes
in the metabolism of acetamiprid (Iwasa et al. 2004), and the evidence that triazine
herbicides like terbuthylazine can interact with detoxification systems in various
organisms, it is plausible that co-application could overburden shared detoxification
pathways, even if direct metabolic competition remains to be demonstrated in
insects. In summary, the synergistic toxicity of the AT mixture in *O. fasciatus* may be
explained by a combination of stressors induced by the co-applied herbicide.

4.3 Fluopyram

The fungicide fluopyram targets the mitochondrial SDH, a key enzyme of the
respiratory chain and ATP production that is not specific to fungi (Sun et al. 2005).

Unlike other SDH-inhibitor (SDHI) fungicides, fluopyram is also used to affect plant-
parasitic nematodes, acting via the same mode of action as in fungi (Schleker et al.
2022). Strikingly, the SDH of *Apis mellifera* was also shown to be sensitive to
fluopyram (Bénit et al. 2019) suggesting that it could directly impair cellular
metabolism and energy production in insects.

In addition, a recent study conducted on the nematode *Caenorhabditis elegans*
showed that fluopyram not only inhibits SDH, but also triggers oxidative stress,
intestinal damage, and cellular apoptosis (Liu et al. 2022). The authors observed
negative impacts on mobility, feeding, reproduction, and longevity in *C. elegans*.
Adverse effects are also reported for other non-target organisms. Fluopyram was
proven to be harmful to soil microbial communities (Zhang et al. 2014). In fish,
fluopyram exposure can induce altered behavior and metabolic disruption despite a
low mitochondrial complex II inhibition in vitro. However, reduced
acetylcholinesterase activity was detected in the muscle cells (Bouly et al. 2024). In
addition, Zhou et al. (2025) suggest a potential chronic toxicity to fish.

To our knowledge, no synergistic enhancement of insecticide toxicity by fluopyram
has been reported to date. However, a recent study by Manzer et al. (2024)
demonstrated a synergistic increase in acetamiprid toxicity in honeybees when
larvae were orally exposed to a mixture of two other fungicides inhibiting
mitochondrial electron transport: the SDHI boscalid (targeting mitochondrial complex
II) and dimoxystrobin (targeting mitochondrial complex III). This combination led to
increased adult mortality compared to acetamiprid alone, whereas exposure to the
fungicide mixture alone had no effect. Interestingly, this synergy was only observed
in a low-dose mixture, but not at higher concentrations.

In our study, fluopyram alone did not cause any measurable acute toxic effects in *O.*
fasciatus nymphs, and the combination with acetamiprid did not significantly increase
toxicity compared to the acetamiprid-only treatment. It remains possible that the
experimental design did not capture delayed or developmental-stage-specific effects.
Notably, in the study by Manzer et al. (2024), increased mortality was observed only
20-30 days after emergence, following larval-stage exposure.

Although no significant synergistic toxicity was detected in our binary AF treatment,
potential interactions between fluopyram and acetamiprid cannot be ruled out. While
acetamiprid selectively activates nicotinic acetylcholine receptors, fluopyram has
been shown to inhibit the AChE in vertebrates, which may also apply to
invertebrates. Such inhibition could lead to endogenous acetylcholine accumulation
and overstimulation of cholinergic pathways. In combination with acetamiprid, this
may result in dual interference within the cholinergic system, plausibly increasing
toxicity. Additionally, the potential inhibition of mitochondrial respiration in
invertebrates by fluopyram could reduce the availability of cellular energy, which
could compromise detoxification mechanisms. As energy availability is essential for
cellular detoxification processes, such as biotransformation and the active transport

of xenobiotics, mitochondrial disruption could indirectly increase susceptibility to
insecticides.

While these scenarios remain speculative, it is notable that the proportion of
unaffected (i.e., neither dead nor affected) nymphs declined more steeply with
increasing acetamiprid concentrations in the AF mixture than in the acetamiprid-only
treatment (Fig. 2). Although our dose-response analysis focused on dead and
affected individuals, this trend was observed visually but not statistically analyzed,
suggesting a potential interaction not captured by ED_{50} values alone. However,
significantly increased toxicity in mixtures including fluopyram was only observed in
the triple mixture AFT.

4.4 Complex mixtures

The significantly higher toxicity observed in the triple mixture AFT is likely due to the
combined physiological stress imposed by all three co-applied pesticides. Previous
studies have shown that terbuthylazine and fluopyram can interfere with critical
biological processes in several non-target organisms, including cellular energy
metabolism, oxidative balance, detoxification capacity and neurotransmission (see
above). While many of these pesticide-induced effects have not been confirmed
directly in insects, their presence in a broad range of organisms suggests that they
may also be observed in insects. Taken together, the potential co-occurrence of
neurotoxicity, metabolic disturbances and oxidative stress could overwhelm cellular
defense mechanisms. Therefore, the pronounced increase in toxicity observed in the
AFT treatment, compared to all other treatments, may be explained by the collapse
of critical physiological systems, including ATP-dependent processes such as
detoxification and cholinergic regulation.

It is important to note that our study only assessed acute toxicity. Potential sublethal
effects, such as behavioral or physiological alterations, impaired development or
communication, and reproductive deficits, were not investigated but are commonly
reported, including for pesticide mixtures (Müller 2018; Gandara et al. 2024). The
potential for complex mixtures to cause disproportionately stronger sublethal effects
than binary combinations has not yet been systematically addressed and demands
further investigation.

4.5 Realistic Scenarios and Risk Assessment

Although this study was not designed to reflect a specific local exposure scenario,
but rather to address the general question of combined toxicity across pesticide
classes, the combination of the tested substances is not merely hypothetical. There
are realistic conditions under which such mixtures can be encountered by non-target
organisms.

Firstly, it is noteworthy that ecotoxicological studies of environmental samples, such
as soil, vegetation, water and crop plants, conducted in recent years have detected

all three of the substances tested here across all sample types (Honert et al. 2025;
Mauser et al. 2025).
Secondly, environmental measurements clearly show that the tested combinations
that showed synergistic effects (AT, AFT) are indeed present in the landscape. Soil
samples from agricultural areas in Rhineland-Palatinate (Germany) analyzed by
Honert et al. (2025) contained the a.s. combination AT in topsoil. The combination
AFT was detected only once. However, FT was often found in combination with one
or more other neonicotinoids, such as thiamethoxam, clothianidin, or imidacloprid.
These local examples of potential exposure to hazardous a.s. combinations clearly
indicate a landscape-level risk. Brühl et al. (2021) further demonstrate this risk by
finding that insects in nature conservation zones in Germany were contaminated, on
average, with 16.7 pesticides. While this finding is alarming, it is not unexpected,
given that contamination of areas surrounding fields – which in some cases might be
conservation zones – is generally anticipated (Rautmann et al. 2001). Given the
results of our study, these findings are alarming and demand further attention from
research, legislation and risk assessment.
582
While EU regulations acknowledge the importance of assessing cumulative and
synergistic effects in the context of simultaneous or sequential applications, current
risk assessment procedures still predominantly focus on individual active substances
(EFSA PPR 2015). As a result, the ecological risks arising from the potential for
multiple substances to co-occur in the environment forming complex mixtures remain
insufficiently addressed. Such mixtures may arise in-field due to common plant
protection practices and the long environmental persistence of some substances
(e.g., fluopyram, DT90 = 833 days in soil in field at 20 °C; Lewis et al. 2016), or off-
field via various input pathways (e.g., spray drift; Honert et al. 2025; Mauser et al.
2025).
Investigating combined pesticide effects is inherently challenging: the number of
possible compound combinations is vast, and the range of potential concentration
scenarios is practically infinite. Our results show that complex mixtures are a major
source of uncertainty in ecotoxicological assessment due to their unpredictable
interaction potential. Although environmental pesticide concentrations are often
comparatively low, our study demonstrates at the mechanistic level that interactions
among active substances may occur more broadly than previously assumed, making
interactions at lower environmental doses plausible. In addition, a recent study
demonstrated that environmentally relevant combinations of pesticides at realistic
concentrations do exert adverse effects on non-target insects (Gandara et al. 2024).
Moreover, a.s. interactions may manifest differently across species, further
increasing the complexity of this field. We therefore propose testing realistic
exposure scenarios based on (1) plant protection practices, including crop type,
application method, and degradation behavior, and (2) environmental monitoring of
actual a.s. combinations. These tests should consider both likely individual

concentrations and potentially affected non-target organisms, depending on the
environmental matrix (soil, plant, water, air).

We believe a reformed ecotoxicological assessment is urgently needed, given that
evidence, such as this study, increasingly suggests environmental pesticide mixtures
may be influencing the ongoing biodiversity crisis.

References

Bénit P, Kahn A, Chretien D, et al (2019) Evolutionarily conserved susceptibility of
the mitochondrial respiratory chain to SDHI pesticides and its consequence
on the impact of SDHIs on human cultured cells. PLOS ONE 14:e0224132.
https://doi.org/10.1371/journal.pone.0224132

Bohan DA, Boffey CWH, Brooks DR, et al (2005) Effects on weed and invertebrate
abundance and diversity of herbicide management in genetically modified
herbicide-tolerant winter-sown oilseed rape. Proceedings of the Royal Society
B: Biological Sciences 272:463–474. https://doi.org/10.1098/rspb.2004.3049

Bouly L, Jacquin L, Chapeau F, et al (2024) Fluopyram SDHI pesticide alters fish
physiology and behaviour despite low *in vitro* effects on mitochondria.
Ecotoxicology and Environmental Safety 288:117400.
https://doi.org/10.1016/j.ecoenv.2024.117400

Brühl CA, Bakanov N, Köthe S, et al (2021) Direct pesticide exposure of insects in
nature conservation areas in Germany. Sci Rep 11:24144.
https://doi.org/10.1038/s41598-021-03366-w

Brühl CA, Engelhard N, Bakanov N, et al (2024) Widespread contamination of soils
and vegetation with current use pesticide residues along altitudinal gradients
in a European Alpine valley. Commun Earth Environ 5:1–9.
https://doi.org/10.1038/s43247-024-01220-1

BVL (2018) Neue Vorgaben zum Bienenschutz bei Tankmischungen von Insektiziden
mit Fungiziden. In: Bundesamt für Verbrucherschutz und
Lebensmittelsicherheit.
https://www.bvl.bund.de/SharedDocs/Fachmeldungen/04_pflanzenschutzmitte
l/2018/2018_11_12_Fa_Bienenschutz_Tankmischung_Insekt_Fung.html.
Accessed 25 Aug 2025

Dicks LV, Breeze TD, Ngo HT, et al (2021) A global-scale expert assessment of
drivers and risks associated with pollinator decline. Nat Ecol Evol 5:1453–
1461. https://doi.org/10.1038/s41559-021-01534-9

Duffey SS, Blum MS, Isman MB, Scudder GGE (1978) Cardiac glycosides: A
physical system for their sequestration by the milkweed bug. Journal of Insect
Physiology 24:639–645. https://doi.org/10.1016/0022-1910(78)90127-0

EFSA PPR P on PPP and their R (PPR) (2015) Scientific Opinion addressing the
state of the science on risk assessment of plant protection products for non-
target arthropods. EFSA Journal 13:3996.
https://doi.org/10.2903/j.efsa.2015.3996

European Commission (n.d.) EU Pesticides Database (v3.3) - Active substances.
https://ec.europa.eu/food/plant/pesticides/eu-pesticides-
database/start/screen/active-substances. Accessed 1 Apr 2025

FRAC (2025) SDHI fungicides. In: Fungicide Resistance Action Committee.
https://www.frac.info/frac-teams/working-groups/sdhi-fungicides/. Accessed 22
July 2025

Gandara L, Jacoby R, Laurent F, et al (2024) Pervasive sublethal effects of
agrochemicals on insects at environmentally relevant concentrations. Science
386:446–453. https://doi.org/10.1126/science.ado0251

Geiger F, Bengtsson J, Berendse F, et al (2010) Persistent negative effects of
pesticides on biodiversity and biological control potential on European
farmland. Basic and Applied Ecology 11:97–105.
https://doi.org/10.1016/j.baae.2009.12.001

Honert C, Mauser K, Jäger U, Brühl CA (2025) Exposure of insects to current use
pesticide residues in soil and vegetation along spatial and temporal
distribution in agricultural sites. Sci Rep 15:1817.
https://doi.org/10.1038/s41598-024-84811-4

HRAC (2025) Global Herbicide Classification Lookup. In: Herbicide Resistance
Action Committee. https://hracglobal.com/tools/classification-lookup.
Accessed 22 July 2025

IRAC (2025) Acetamiprid. In: Insecticide Resistance Action Committee. https://irac-
online.org/active-ingredient/acetamiprid/. Accessed 22 July 2025

Iwasa T, Motoyama N, Ambrose JT, Roe RM (2004) Mechanism for the differential
toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. Crop
Protection 23:371–378. https://doi.org/10.1016/j.cropro.2003.08.018

Kao LM, Wilkinson CF, Brattsten LB (1995) In-vivo effects of 2,4-D and atrazine on
cytochrome P-450 and insecticide toxicity in southern armyworm (Spodoptera
eridania) larvae. Pesticide Science 45:331–334.
https://doi.org/10.1002/ps.2780450407

Kruse-Plaß M, Hofmann F, Wosniok W, et al (2021) Pesticides and pesticide-related
products in ambient air in Germany. Environ Sci Eur 33:114.
https://doi.org/10.1186/s12302-021-00553-4

Lewis KA, Tzilivakis J, Warner DJ, Green A (2016) An international database for
pesticide risk assessments and management. Human and Ecological Risk
Assessment: An International Journal 22:1050–1064.
https://doi.org/10.1080/10807039.2015.1133242

Liu Y, Zhang W, Wang Y, et al (2022) Oxidative stress, intestinal damage, and cell
apoptosis: Toxicity induced by fluopyram in Caenorhabditis elegans.
Chemosphere 286:131830.
https://doi.org/10.1016/j.chemosphere.2021.131830

Manzer S, Thamm M, Hilsmann L, et al (2024) The neonicotinoid acetamiprid
reduces larval and adult survival in honeybees (*Apis mellifera*) and interacts
with a fungicide mixture. Environmental Pollution 360:124643.
https://doi.org/10.1016/j.envpol.2024.124643

Mauser KM, Wolfram J, Spaak JW, et al (2025) Current-use pesticides in vegetation,
topsoil and water reveal contaminated landscapes of the Upper Rhine Valley,
Germany. Commun Earth Environ 6:166. https://doi.org/10.1038/s43247-025-
02118-2

Müller C (2018) Impacts of sublethal insecticide exposure on insects — Facts and
knowledge gaps. Basic and Applied Ecology 30:1–10.
https://doi.org/10.1016/j.baae.2018.05.001

Pape☐Lindstrom PA, Lydy MJ (1997) Synergistic toxicity of atrazine and
organophosphate insecticides contravenes the response addition mixture
model. Environmental Toxicology and Chemistry 16:2415–2420.
https://doi.org/10.1002/etc.5620161130

Ralph CP (1976) Natural food requirements of the large milkweed bug,Oncopeltus
fasciatus (Hemiptera: Lygaeidae), and their relation to gregariousness and
host plant morphology. Oecologia 26:157–175.
https://doi.org/10.1007/BF00582894

Rautmann D, Streloke M, Winkler R (2001) New basic drift values in the
authorization procedure for plant protection products. Workshop on Risk
Assessment and Risk Mitigation Measures (WORMM), vol 13 3–141

Salminen J, Eriksson I, Haimi J (1996) Effects of Terbuthylazine on Soil Fauna and
Decomposition Processes. Ecotoxicology and Environmental Safety 34:184–
189. https://doi.org/10.1006/eesa.1996.0062

Sánchez-Bayo F, Wyckhuys KAG (2019) Worldwide decline of the entomofauna: A
review of its drivers. Biological Conservation 232:8–27.
https://doi.org/10.1016/j.biocon.2019.01.020

Schleker ASS, Rist M, Matera C, et al (2022) Mode of action of fluopyram in plant-
parasitic nematodes. Sci Rep 12:11954. https://doi.org/10.1038/s41598-022-
15782-7

Schuhmann A, Schmid AP, Manzer S, et al (2022) Interaction of Insecticides and
Fungicides in Bees. Frontiers in Insect Science 1:

Sedlmeier JE, Grass I, Bendalam P, et al (2025) Neonicotinoid insecticides can pose
a severe threat to grassland plant bug communities. Commun Earth Environ
6:1–12. https://doi.org/10.1038/s43247-025-02065-y

Semren TŽ, Žunec S, Pizent A (2018) Oxidative stress in triazine pesticide toxicity: a
review of the main biomarker findings. Archives of Industrial Hygiene and
Toxicology 69:109–125. https://doi.org/10.2478/aiht-2018-69-3118

Sgolastra F, Medrzycki P, Bortolotti L, et al (2017) Synergistic mortality between a
neonicotinoid insecticide and an ergosterol-biosynthesis-inhibiting fungicide in
three bee species. Pest Management Science 73:1236–1243.
https://doi.org/10.1002/ps.4449

Silva V, Gai L, Harkes P, et al (2023) Pesticide residues with hazard classifications
relevant to non-target species including humans are omnipresent in the
environment and farmer residences. Environment International 181:108280.
https://doi.org/10.1016/j.envint.2023.108280

Sun F, Huo X, Zhai Y, et al (2005) Crystal Structure of Mitochondrial Respiratory
Membrane Protein Complex II. Cell 121:1043–1057.
https://doi.org/10.1016/j.cell.2005.05.025

Tosi S, Sfeir C, Carnesecchi E, et al (2022) Lethal, sublethal, and combined effects
of pesticides on bees: A meta-analysis and new risk assessment tools.
Science of The Total Environment 844:156857.
https://doi.org/10.1016/j.scitotenv.2022.156857

Varga-Szilay Z, Tóth Z (2022) Is acetamiprid really not that harmful to bumblebees
(Apidae: Bombus spp.)? Apidologie 53:2. https://doi.org/10.1007/s13592-022-
00909-6

Wagner DL, Grames EM, Forister ML, et al (2021) Insect decline in the
Anthropocene: Death by a thousand cuts. Proceedings of the National
Academy of Sciences 118:e2023989118.
https://doi.org/10.1073/pnas.2023989118

Wan N-F, Fu L, Dainese M, et al (2025) Pesticides have negative effects on non-
target organisms. Nat Commun 16:1360. https://doi.org/10.1038/s41467-025-
56732-x

Zhang Y, Xu J, Dong F, et al (2014) Response of microbial community to a new
fungicide fluopyram in the silty-loam agricultural soil. Ecotoxicology and
Environmental Safety 108:273–280.
https://doi.org/10.1016/j.ecoenv.2014.07.018

Zhou L, Li Y, He C, et al (2025) Toxicity, bioaccumulation and metabolism of
fluopyram in zebrafish (*Danio rerio*). Aquatic Toxicology 285:107420.
https://doi.org/10.1016/j.aquatox.2025.107420

Data Availability

All data generated in this study are provided in the supplementary materials.

Statements and Declarations

Funding

Open Access funding enabled and organized by Projekt DEAL.

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Contributions

J.E.S., E.G. and G.P designed the experiments. E.G. and J.E.S. conducted the experiments. B.H. and F.W. developed the active substance mixing strategies in a common solvent based on chemical properties. J.E.S. and E.G. performed the analysis of the experiments. J.E.S. wrote the original draft of the manuscript and created the figures. G.P. supervised the research and edited the manuscript.