Ubiquitination regulates granulostasis and DRiP accumulation in SGs under heat stress

via the E3 ligase MKRN2

Emmanuel Amzallag1,2, Yehuda M. Danino1,2, Valentina Secco 3, Serena Carra 3, Eran

Hornstein1,2#

1Department of Molecular Genetics, Weizmann Institute of Science, Israel.

2Department of Molecular Neuroscience, Weizmann Institute of Science, Israel.

3 Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio

Emilia Modena, Italy.

Corresponding Author: Eran.Hornstein@weizmann.ac.il

Abstract

Stress granules (SGs) are transient cytoplasmic biomolecular condensates that play a role in the

cellular response to proteotoxic stress. It has been previously shown that ubiquitination regulates

SG dynamics; however, the specific mechanisms by which ubiquitin affects SGs are not fully

understood. Here, using proximity proteomics, we discover that the engagement of several E3

ubiquitin ligases in SGs is dependent on UBA1 activity. A detailed study of the E3 ubiquitin-protein

ligase Makorin 2 (MKRN2) demonstrated that it is localized to SGs in a manner dependent on

active ubiquitination. MKRN2 promotes both the proper formation of SGs and their disassembly

following stress recovery, by preventing the accumulation of defective ribosomal products (DRiPs)

within SGs. Therefore, MKRN2 is a novel regulator of SGs that mediates the maintenance of

granulostasis, suggesting that the localization of a subset of E3 ligases into SGs is linked to their

capacity to ubiquitinate target proteins.

Introduction

Stress granules (SGs) are biomolecular condensates that form via a process of liquid-liquid phase

separation (LLPS) of biomacromolecules (1). SG assembly is initiated by a variety of stressors,

including heat stress, oxidative stress, osmotic stress, and ER stress, during which the cell

activates the integrated stress response (ISR) (2). Phosphorylation of eIF2α initiates the ISR,

resulting in polysome disassembly and global shutdown of translation. This releases mRNA

molecules from ribosomes and engages a variety of RNA-binding proteins (RBPs) to form

multivalent interactions with both RNA and with other proteins via RNA-binding domains (RBDs)

and intrinsically disordered domains (IDD), respectively, resulting in the formation of discrete SGs.

Upon recovery from stress, SGs rapidly disassemble, and their dissolution allows mRNAs to be

redirected to translation (1).

However, SGs can fail to disassemble and persist for longer time, affecting the cells' ability to

restore translation, as well as the retro-translocation in the nucleus of RBPs (1). Several factors

can delay SG disassembly including the accumulation of misfolded and damaged proteins and

the impairment of the protein quality control system. The latter refers to molecular chaperones

and protein degradation systems that cooperates to detect unfolded/misfolded proteins and

promote their clearance (3). Amongst the misfolded and potentially toxic proteins that can

accumulate inside SGs changing their dynamic liquid-like properties are disease-linked mutated

RBPs such as TDP-43 or FUS or hnRNPA1 (4–6), as well as defective ribosomal products

(DRiPs), which are truncated proteins released from disassembling polyribosomes. DRiPs are

rapidly polyubiquitinated for degradation by the ubiquitin-proteasome system as well as by

autophagy (7). Amongst the chaperones that participate in the maintenance of SG dynamics are

the ATP-dependent chaperone complexes HSP70/HSPB8/BAG3 and VCP/UFD1L/PLAA. These

chaperone complexes reduce DRiP accumulation inside SGs (8–10). Accordingly, the activity of

HSP70, VCP, and also the autophagic machinery, thus helps surveil stress-induced DRiPs and

maintain SGs functioning properly, in a process termed granulostasis (10).

Until now, the maintenance of granulostasis under stress has been largely associated with the

activity of specific molecular chaperones, namely the HSP70/HSPB8/BAG3 and

VCP/UFD1L/PLAA complexes (8,9). The ubiquitin-proteasome system (UPS) is a major

component of cellular protein degradation and mediates the conjugation of ubiquitin onto target

molecules marked for proteasomal degradation. While ubiquitination and SUMOylation, have

been shown to impact SG dynamics as well (11–13) it is unknown if they regulate DRiPs

accumulation in SGs.

Active ubiquitination is required for proper SG disassembly (11,14–17); however, the specific E3

ligases that maintain proper SG dynamics and granulostasis are only partially known. For

example, the E3 ligase TRIM21 controls G3BP1 ubiquitination and SG condensation and the E3

ligase HECT regulates SG disassembly (16,17). However, whether specific E3 ligases regulate

the clearance of DRiPs from SGs has not been yet explored.

Here, we report ubiquitination-dependent changes in SG proteomic composition that led to the

identification of Makorin 2 (MKRN2) as an important SG-engaged E3 ligase that regulates SG

dynamics and the sequestration of stress-induced DRiPs into SGs.

Results

Ubiquitination affects stress granule dynamics and composition

It was recently shown that active ubiquitination is required for proper SG disassembly

(11,14,15,18). However, a detailed characterization of how SGs and the UPS interact has not

been achieved. Using G3BP1-mCherry expressing U2OS cells (in which the endogenous G3BP1

and 2 genes were knocked out) (19) we studied the full SG assembly and disassembly cycle,

induced by heat stress (HS; 43°C followed by stress recovery at 37°C; Fig. 1A,1B). Inhibition of

the ubiquitin-activating enzyme UBA1 (ubiquitin-like modifier-activating enzyme 1) by TAK243 (1

µM) facilitated SG assembly, while delaying their dissolution (Fig. 1A-C, Supplementary fig. 1).

Treatment with TAK243 coupled with HS caused SGs to nucleate and begin assembling earlier

than in HS-treated cells (Fig.1B, 1C). SG disassembly following recovery from heat stress, at 37

°C, was also delayed (Fig.1A, B, D, Supplementary fig. 1).

To determine whether active ubiquitination affects SG liquidity, we performed fluorescence

recovery after photobleaching (FRAP) on cells under HS, following UBA1 inhibition. Inhibition of

ubiquitination delayed fluorescence recovery dynamics of G3BP1-mCherry, relative to HS treated

cells (Figure 1F, Supplementary fig. 6A). These observation indicate that ubiquitination is required

for maintaining SG liquidity, consistent with previous reports (14).

To characterize the composition of SGs and identify players of the ubiquitination pathway that are

involved in maintaining SG dynamics, we used APEX proximity labeling coupled to mass

spectrometry (APEX-MS). G3BP1/2 KO U2OS cells expressing G3BP1-APEX were used to label

SG proteins, while U2OS cells expressing nuclear export signal (NES)-APEX were used to label

the surrounding cytoplasm). Heat stress was induced (43°C; 90 min), UBA1 was inhibited with

TAK243, and proximity labeling was induced with biotin phenol (500 µM) and a short hydrogen

peroxide pulse (1mM, 60 seconds). Biotin-labelled proteins were isolated with streptavidin

magnetic beads and analyzed by mass spectrometry. Samples without biotin-phenol were used

to identify and subtract non-APEX-derived biotinylation background. Peptide reads were mapped

to proteins with the MaxQuant search engine (v1.6.6.0). Relative to the NES cytoplasmic control,

6200 proteins interacting with G3BP1-APEX were identified, including previously-reported SG

proteins (Fig. 2A), providing confidence that our protocol effectively isolated SG proteomes. To

understand which proteins are associated with SGs in a ubiquitination-dependent manner, we

next compared the HS-induced SG proteomes without or with TAK243 (Fig. 2B, Supplementary

fig. 2A,B). This provided a list of proteins either enriched (83 proteins) or depleted (215 proteins)

from SGs upon inhibition of ubiquitination (Student's T-test, FDR<0.05; Fig. 2C). Amongst the

proteins that associated with SGs in a ubiquitin-dependent manner we report HSPA12, DNAJA1,

and the VCP cofactors, UFD1L, NSFL1C, FAF1, and PLAA. Interestingly, these chaperones were

previously implicated in promoting SG disassembly by preventing aberrant accumulation of

stress-damaged proteins (9,14). Ubiquitin hydrolases and E3 ligases were also associated with

SGs in a ubiquitination-dependent manner, including the RNA-binding ubiquitin ligases (RBULs)

MKRN2, CNOT4, TRIM25, and ZNF598. We orthogonally validated the ubiquitin-dependent

interaction of the most SG-depleted E3 ligase, MKRN2, by immunofluorescence in U2OS cells

fixed after 90 min HS, without or with TAK243 (Fig. 2D, 2E). In agreement with APEX-MS results,

endogenous MKRN2 exhibited decreased SG partitioning in response to TAK243 treatment (Fig.

2E).

The E3 ligase MKRN2 affects SG dynamics

While the human genome encodes only 2 E1s and ~30 E2 enzymes, ~600 E3 ligases are known,

conferring substrate specificity in targeting specific proteins and cellular pathways (20). To

determine whether ubiquitin-sensitive RBULs ligases control SG dynamics, we knocked down

CNOT4, ZNF598, and MKRN2 in U2OS cells that were subjected to HS, followed by recovery (43

°C, 90 min -> 37 °C). Knockdown of CNOT4, ZNF598, or MKRN2 all impacted SG dynamics to

varying degrees (Supplementary Fig. 5A, B). MKRN2 depletion affected both SG assembly and

disassembly, with cells forming smaller SGs that took longer to disassemble upon recovery

(Figure 3,4). ZNF598 similarly affected SG size but had a smaller impact on disassembly. CNOT4

had a minimal effect on SG assembly, with a modest effect on their disassembly (Supplementary

Fig. 5A, B). Based on our mass spectrometry data, indicating MKRN2 as the most highly SG-

depleted E3 upon TAK243 treatment and its strongest impact on SG dynamics, we focused

mechanistic studies on MKRN2.

Using SG live imaging, we investigated the consequences of MKRN2 KD on SG dynamics under

HS. SG assembly under heat stress typically involves a reduction in SG numbers over time, as

smaller condensates coalesce to form larger but fewer SGs (8,21–23). The knockdown (KD) of

MKRN2 resulted in a higher number of SGs at 90 min of HS (Fig. 3A, B, D) and the average size

of SGs was almost half as small compared to the control (Fig. 3C, EI Supplementary fig. 3). These

data suggest that SGs fail to coalesce and mature upon MKRN2 KD. Having established that

MKRN2 depletion affects SG number and size, we then measured their liquid-like properties by

performing FRAP experiments, using the mobility of G3BP1 as readout. Consistently, FRAP

analysis demonstrated that MKRN2 KD resulted in a slight increase in SG liquidity (Fig. 3F,

Supplementary fig. 6B), that could be interpreted as associated to decreased coalescence into

dense and mature SGs (23,24).

Then, to further test the impact of MKRN2 on SG dynamics, we overexpressed MKRN2–GFP and

we observed that it was recruited to SG (Fig. 3G). We next quantified SG ubiquitin content in

response to MKRN2-GFP overexpression, using an antibody that recognizes free, mono- and

poly-ubiquitinated proteins (FK2). MKRN2-GFP increased the partitioning of ubiquitin into SGs,

relative to neighboring cells that did not express MKRN2-GFP in the same polyclonal mosaic

culture (Fig. 3G, 3H). Together these data indicate that MKRN2 regulates the assembly dynamics

of SGs by promoting their coalescence during HS and can increase SG ubiquitin content.

We next tested the impact of MKRN2 on SG disassembly following stress removal. MKRN2 KD

delayed the disassembly of SGs (Fig. 4A, 4B; Supplementary fig. 4). The effect was pronounced

30 and 60 minutes after cessation of HS (Fig. 4C, 4D). Therefore, MKRN2 is required for

maintaining proper SG dynamics under HS.

MKRN2 regulates accumulation of DRiPs in SGs

DRiPs are truncated peptides that are prone to aggregation and DRiPs ubiquitination affects their

solubility and clearance (7,8,25) . Proteostasis impairment causes aberrant DRiP accumulation

inside SGs and impairs SG disassembly. This suggests the idea that MKRN2 may ubiquitinate a

subset of DRiPs, avoiding their accumulation inside SGs and facilitating their disposal. To test the

idea that ubiquitin impacts a subset of DRiPs in SGs, we induced SG formation by HS in absence

or presence of the UBA1 inhibitor TAK243. Concomitantly we treated the cells with O-propargyl

puromycin (OP-puro, 25uM), which is incorporated into nascent polypeptide chains, causing

polysome disassembly and the emergence of DRiPs. We observed increased partitioning of OP-

puro labelled DRiPs in SGs upon inhibition of ubiquitination relative to control (Fig. 5A, 5B). These

data suggest that active ubiquitination prevents the accumulation of DRiPs inside SGs. To

understand whether a fraction of DRiPs is ubiquitinated by MKRN2, we then performed a similar

analysis in proficient and MKRN2-deficient cells. We measured increased partitioning of OP-puro

labelled DRiPs in SGs upon knockdown of MKRN2, relative to control (Fig. 5C, 5D). Therefore,

we conclude that MKRN2 ubiquitinates a subset of DRiPs, avoiding their accumulation inside

SGs.

Together our study provides evidence for a new role for ubiquitination and for the E3 ligase

MKRN2 in the control of SG dynamics and DRiPs turnover.

Discussion

In this study, we characterize the impact of ubiquitination on SG proteomic composition during

heat stress. Using APEX proximity labeling and mass spectrometry, we demonstrate that

ubiquitination regulates SG functions under heat stress by controlling the recruitment of several

molecular regulators of proteostasis into SGs.

Specifically, ubiquitination regulates the localization of several proteostasis-related proteins to

SGs, including four VCP cofactors (UFD1L, PLAA, NSFL1C, and FAF1), two of which (UFD1L

and PLAA) have been shown to regulate SG dynamics and sequestration of DRiPs (8). This

suggests that ubiquitin-binding VCP cofactors are recruited to SGs through their interaction with

ubiquitinated proteins within these condensates. Furthermore, the localization of the heat shock

protein HSP70 (HSPA12B) and its co-chaperone DNAJA1 to SGs is also regulated by

ubiquitination. This finding is consistent with similar behavior observed under sodium arsenite

stress (17) and is likely to influence the accumulation of DRiPs in SGs (9).

Several E3 ubiquitin ligases were depleted from SGs in the absence of UBA1 activity, namely

MKRN2, ZNF598, CNOT4, TRIM25, and TRIM26, whereas the E3 RNF20 was enriched in SGs.

A detailed study of the E3 ligase MKRN2 demonstrated ubiquitination-dependent partitioning of

endogenous MKRN2 into SGs. This suggests that the localization of a subset of E3 ligases into

SGs is linked to their capacity to ubiquitinate target proteins.

Pan et al. reported enrichment of several HECT-domain E3 ligases in SGs following inhibition of

ubiquitination by TAK243 (17), suggesting that this process is specific and potentially regulated

in a stress- or protein-specific manner. SG E3 ligases that depend on UBA1 activity are RBULs,

and their capacity to bind RNA may be particularly intriguing for SG E3s.

Endogenous and ectopically expressed MKRN2 were both localized to SGs. MKRN2 has a

functional role in regulating SG dynamics under heat stress, and its knockdown results in more

numerous and smaller SGs, as well as a significant delay in SG disassembly following recovery

from stress.

We also show that UBA1 activity and MKRN2 regulates a subset of DRiP homeostasis in SGs.

Thus, ubiquitination is required for preventing excessive DRiPs accumulation in SGs under heat

stress. We further show that the RBUL MKRN2 similarly prevents accumulation of DRiPs in SGs

suggesting that it may execute ubiquitin-mediated extraction of DRiPs from SGs. We note that

our experiments cannot delineate whether ubiquitination of DRiPs by MKRN2 occurs inside SGs

or in the cytoplasm. However, the finding that MKRN2 is enriched in SGs under stress coupled

with the fact that its overexpression increases SG ubiquitin content, suggests that MKRN2

controls DRiP clearance from SGs via polyubiquitination of SG-localized substrates. In summary,

MKRN2 is the first E3 ligase shown to play a role in DRiP extraction from SGs. The phenotype of

smaller, more numerous SGs concomitant with increased DRiP partitioning, also reported upon

inhibition of autophagy and VCP function (8), suggests that MKRN2 may act in the same pathway.

We note that the SG disassembly phenotype upon MKRN2 knockdown is less pronounced than

that observed with TAK243 treatment. Therefore, MKRN2 may target a subset of DRiPs.

consistent with the role of additional SG-localized E3 ligases, such as ZNF598 or CNOT4, that

regulate SG dynamics. Overall, our results support the conclusion that proper SG dynamics

depend on clearance of DRiPs from these condensates and portray MKRN2 as a novel

proteostasis factor regulating this process.

Limitations: The direct MKRN2 substrates and ubiquitin-chain types (K63/K48) are currently

unknown. In addition, it is interesting to understand if DRiP solubility is dependent on

ubiquitination or on specific E3 ligases, such as MKRN2.

In sum, global ubiquitination promotes the recruitment of key SG-engaged chaperones, including

HSP70 and VCP, as well as a subset of E3 ubiquitin ligases. We identify the E3 ligase MKRN2

as a novel regulator of SG dynamics and proteostatic function. Clearance of DRiPs from SGs

requires active ubiquitination and is mediated in part by MKRN2. These insights refine our

understanding of how SGs coordinate the triage of stress-damaged proteins with condensate

dynamics during proteotoxic stress. Ultimately, defining ubiquitination-dependent regulation of SG

composition and function may illuminate new therapeutic avenues for restoring proteostasis in

diseases that involve protein aggregation.

Methods

Mammalian cell culture

U2OS cells were cultured in growth media consisting of high glucose DMEM (Sartorius 01-052-

1A), 10% FBS (Sigma F7524) and 1% penicillin-streptomycin-amphotericin B (Sartorius 03-033-

1B) in a 37°C incubator supplemented with 5% CO_2. To induce heat stress, cells were placed

into a dedicated incubator set to 43°C supplemented with 5% CO_2. For UBA1 inhibition, cells

were pre-treated with 1uM TAK243 (Med Chem Express; Cat# HY-100487) prior to heat stress.

For MKRN2 KD, cells were transfected with MKRN2 siRNA (Dharmacon siGENOME

SMARTpool; M-006960-01-0005) using Lipofectamine RNAiMAX (Cat#: 13778075) and

cultured for 72 hours before analysis.

Molecular cloning

Molecular cloning of MKRN2-GFP was performed using In-Fusion® Snap Assembly (Cat#

638947; Takara Clontech) and inserted into pcDNA4.0 vector with puromycin resistance.

G3BP1-mCherry cells were transfected with Lipofectamine 2000 (Thermo Fisher Scientific, Cat#

11668027) cultured with 5ng/ml puromycin for 7 days.

Microscopy

SG live imaging was performed in a Zeiss CD7 widefield microscope equipped with an Axiocam

702 CMOS camera using a 20X air objective with an additional 2X post-magnification (effective

magnification of 40X, excitation wavelength 590nm). G3BP1-mCherry cells were seeded into

optical 24-well plates (Cellvis P24-1.5H-N) and placed into the heated microscope incubator

chamber (37°C, 5% CO_2). After one cycle of imaging, the microscope incubator temperature

was increased to 43°C. After 90 min of heat stress, the temperature was lowered back down to

37°C to image SG disassembly. SG number and area was analysed using the surfaces feature

in Imaris software. Immunofluorescent staining was performed on 12mm round glass coverslips

inside 24-well plates. Immunofluorescent images were captured in a Zeiss LSM900 confocal

microscope using a 63X oil objective lens.

FRAP

G3BP1-mCherry cells were seeded into 3.5mm dishes with a central coverslip (MatTek P35GC-

1.5-14-C). Cells were imaged in a Leica SP8 STED microscope using a 100X objective. To

induce heat stress, the microscope incubator temperature was increased to 43°C and an

objective heater was attached to the objective and heated to 43°C. Cells were left in the heated

microscope incubator chamber for 60 minutes, and FRAP was performed for 30 minutes

thereafter. To photobleach SGs, a 561nm laser set to max power was used to bleach entire

SGs. In order to monitor fluorescence recovery, images were collected every 400ms for 63

seconds. Fluorescence recovery data was collected using the Leica LasX software. Analysis

corrected for background fluorescence intensity and normalized to the pre-bleach SG intensity

to calculate the fraction of fluorescence recovery at each time point.

APEX proximity labelling

G3BP1-APEX or NES-APEX U2OS (Marmor-Kollet *et al.*) cells were seeded into T125 flasks

with 50ng/ml tetracycline (Sigma T7660) and allowed to incubate for 24h before performing

APEX experiment. Four technical repeats were used for each condition. Following pre-treatment

with TAK243, cells were supplemented with 500uM Biotin-Tyramide (Iris Biotech 41994-02-9)

for 90 minutes to allow for labelling activity. Labelling was then induced by addition of H_2O_2 (JT

Baker 2186-01) for 60 seconds. APEX activity was extinguished with quenching solution (10mM

sodium azide (Sigma cat number S2002), 10mM sodium ascorbate (Alfa Aesar A17759), 5mM

Trolox (Sigma 238813) in PBS. Cells were then scraped in PBS, centrifuged at 3000xg for 10

minutes at 4°C and the cell pellet was lysed in ice-cold RIPA buffer (50mM Tris-HCl pH8,

150mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS) supplemented with cOmplete

Protease Inhibitor Cocktail (Roche, 4693116001) and PhosSTOP (Roche, 4906837001).

Lysates were centrifuged at 15000xg for 10 minutes at 4°C and protein concentration was

quantified using Bradford assay (Bio-Rad 5000006). Streptavidin magnetic beads (Thermo

Fisher cat number 88817) were incubated with the protein extracts at a ratio of 100ul beads per

500ug protein extract overnight at 4°C with rotation.

Liquid chromatography and mass spectrometry

Samples were subjected to on-bead tryptic digestion. The resulting peptides were analysed

using nanoflow liquid chromatography (nanoAcquity) coupled to high resolution, high mass

accuracy mass spectrometry (Q-Exactive Plus). Each sample was analysed on the instrument

separately in a random order in discovery mode. Raw data were processed with MaxQuant

v1.6.6.0. The data were searched with the Andromeda search engine against the human

proteome database, appended with common lab protein contaminants and the default

modifications. Quantification was based on the LFQ method, based on unique/all peptides.

Proteomics statistical analysis

ProteinGroups output table from MaxQuant was imported into Perseus for analysis. First,

contaminants were removed. Raw intensity values were $\log_2$ transformed and missing values

imputed for proteins with at least four valid values using an artificial normal distribution with a

downshift of 1.8 standard deviations and a width of 0.3 of the original ratio distribution. To

identify proteins non-specifically bound to streptavidin beads, student's t test with FDR <=0.05

was used to compare each sample to respective no BP control samples. These non-specific

binders were filtered out and proteins were then compared between G3BP1-APEX and NES-

APEX baits. LFQ values were $\log_2$ transformed and missing values were imputed for proteins

with at least two valid values in at least one group using an artificial normal distribution with a

downshift of 1.8 standard deviations and a width of 0.3 of the original ratio distribution. Proteins

were then compared between the G3BP1-APEX bait to the NES-APEX bait using student's t

test with FDR <=0.05 to determine which proteins were significantly enriched in proximity to

G3BP1 compared to NES. G3BP1-APEX values were then normalized by the mean of their

corresponding NES-APEX values. The NES-normalized values were then compared between

the TAK243 vs DMSO treated samples using student's t test with FDR <=0.05.

Statistical analysis

Statistical analysis of live imaging data was done using repeated-measures ANOVA test with

contrast analysis in R. Pairwise comparisons were done using Welch's t test in Graphpad Prism.

Statistical p values < 0.05 were considered significant.

Labeling of nascent peptides with OP-puro and high content imaging-based assay

Cells were grown on polylysine-coated glass coverslip. Newly synthesized proteins were

labelled by incubating the cells with 25 mM O-Propargyl-puromycin (OP-puro) for the indicated

time points and CuAAC detection of OP-puro labelled DRiPs was performed as previously

described, using Alexa Fluor 594 azide (A10270, Life Technologies) (PMID: 22160674). Cells

were next processed for immunofluorescence microscopy. Blocking and incubation with primary

and secondary antibodies were performed in PBS containing 3% BSA and 0.1% Triton X-100.

DRiPs enrichment inside SGs was measured using the Scan^R Analysis software (Olympus).

SGs were segmented based on G3BP1 signal using edge detection algorithm. The mean

fluorescence intensity of OP-puro labelled DRiPs was detected in each detected SG and was

additionally measured in an area surrounding each SG. The relative enrichment of OP-puro

labelled DRiPs in individual SGs was calculated as a ratio of mean fluorescence intensity inside

the SG divided by mean intensity in the region surrounding the SG, as previously described

(PMID: 27570075).

Contributions statement

E.H. and E.A. oversaw the conceptualization of this study. E.A. and Y.M.D. performed the APEX

proteomics experiment. Mass spectrometry was performed at the de Botton Institute for Protein

Profiling from the Nancy and Stephen Grand Israel National Center for Personalized Medicine (G-

INCPM). E.H. and E.A. designed the study and had unrestricted access to all data. E.A. performed

and analyzed all live imaging experiments. E.A. performed immunofluorescence staining,

microscopy and analysis of ubiquitin and MKRN2. E.A. performed all FRAP experiments. V.S.

and S.C. performed the O-propargyl puromycin fluorescent staining, microscopy and analysis of

DRiP partitioning in stress granules. E.A. and E.H participated in drafting the initial manuscript

draft and E.H., E.A. and S.C. reviewed and revised the draft. All authors agreed to submit the

manuscript, read and approved the final draft, and take full responsibility for its content, including

the accuracy of the data and its statistical analysis.

Acknowledgments

Eran Hornstein is the Mondry Family Professorial Chair and Head of the Nella and Leon Benoziyo

Centre for Neurological Diseases and of the Andi and Larry Wolfe Centre for Neuroimmunology

and Neuromodulation. The authors thank Dr. Yishai Levin, Dr. Alon Savidor, Corine Katina at the

de Botton Institute for Protein Profiling from the Nancy and Stephen Grand Israel National Center

for Personalized Medicine (G-INCPM) for performing mass spectrometry. The authors thank

Yoseph Addadi, Inna Goliand and Tatiana Smirnova from the de Picciotto Cancer Cell

Observatory In memory of Wolfgang and Ruth Lesser of the Moross Integrated Cancer Center in

the department of Life Science Core Facilities, Weizmann Institute of Science for help with

performing live imaging and FRAP. The authors thank Prof. Itay Koren for his advice on our study.

The authors thank CIGS (University of Modena microscopy facility) and Dr. Jonathan Vinet for

technical support for confocal microscopy image acquisition and analysis. Special thanks to Iddo

Magen, Yahel Cohen and Aviad Siany from the Hornstein Lab for fruitful discussion and advice.

Conflict of interests

The authors declare that they have no conflict of interest.

Funding

Eran Hornstein lab is funded by the Andi and Larry Wolfe Centre for Neuroimmunology and

Neuromodulation; the Binational Science Foundation (BSF); Association Française Contre les

Myopathies (AFM) grants 24882, 28680; Muscular Dystrophy Association (MDA) grant 1280000;

Target ALS; Israel Science Foundation (ISF 3497/21, 424/22); ALS Canada; Minna-James-

Heineman Stiftung through, Minerva Foundation, with funding from the Federal German Ministry

for Education and Research; Robert Packard Center for ALS Research at Johns Hopkins; McGill

University; EU - ERA-Net; Radala Foundation for ALS Research; Additional support generously

provided by the Kekst Family Institute for Medical Genetics. Weizmann SABRA - Yeda-Sela -

WRC Program, the Estate of Emile Mimran, and The Maurice and Vivienne Wohl Biology

Endowment. Nella and Leon Benoziyo Center for Neurological Diseases, Goldhirsh-Yellin

Foundation. Dr. Sydney Brenner and friends, Weizmann - Center for Research on

Neurodegeneration. Redhill Foundation – Sam and Jean Rothberg Charitable Trust Dr. Dvora

and Haim Teitelbaum Endowment Fund and Weizmann – Institute for Artificial Intelligence -

Acceleration Grant. Target ALS; Israel Science Foundation (ISF 3497/21, 424/22), CReATe

Consortium. Carra lab was supported by the Armenise-Harvard and AirAlzh Foundations (AHA

Mid-Career Award 2022), the AriSLA Foundation (SUMOsolvable).

References

1. Protter DSW, Parker R. Principles and properties of stress granules. Trends Cell Biol. 2016
Sept;26(9):668–79.

2. Bond S, Lopez-Lloreda C, Gannon PJ, Akay-Espinoza C, Jordan-Sciutto KL. The integrated
stress response and phosphorylated eukaryotic initiation factor 2α in neurodegeneration. J
Neuropathol Exp Neurol. 2020 Feb 1;79(2):123–43.

3. Hipp MS, Kasturi P, Hartl FU. The proteostasis network and its decline in ageing. Nat Rev
Mol Cell Biol. 2019 July;20(7):421–35.

4. Gordon D, Dafinca R, Scaber J, Alegre-Abarrategui J, Farrimond L, Scott C, et al. Single-
copy expression of an amyotrophic lateral sclerosis-linked TDP-43 mutation (M337V) in
BAC transgenic mice leads to altered stress granule dynamics and progressive motor
dysfunction. Neurobiol Dis. 2019 Jan;121:148–62.

5. Baron DM, Kaushansky LJ, Ward CL, Sama RRK, Chian R-J, Boggio KJ, et al.
Amyotrophic lateral sclerosis-linked FUS/TLS alters stress granule assembly and
dynamics. Mol Neurodegener. 2013 Aug 31;8(1):30.

6. Gui X, Luo F, Li Y, Zhou H, Qin Z, Liu Z, et al. Structural basis for reversible amyloids of
hnRNPA1 elucidates their role in stress granule assembly. Nat Commun. 2019 May
1;10(1):2006.

7. Schubert U, Antón LC, Gibbs J, Norbury CC, Yewdell JW, Bennink JR. Rapid degradation
of a large fraction of newly synthesized proteins by proteasomes. Nature. 2000 Apr
13;404(6779):770–4.

8. Seguin SJ, Morelli FF, Vinet J, Amore D, De Biasi S, Poletti A, et al. Inhibition of autophagy,
lysosome and VCP function impairs stress granule assembly. Cell Death Differ. 2014
Dec;21(12):1838–51.

9. Ganassi M, Mateju D, Bigi I, Mediani L, Poser I, Lee HO, et al. A Surveillance Function of
the HSPB8-BAG3-HSP70 Chaperone Complex Ensures Stress Granule Integrity and
Dynamism. Mol Cell. 2016 Sept 1;63(5):796–810.

10. Alberti S, Mateju D, Mediani L, Carra S. Granulostasis: Protein quality control of RNP
granules. Front Mol Neurosci. 2017 Mar 27;10:84.

11. Maxwell BA, Gwon Y, Mishra A, Peng J, Nakamura H, Zhang K, et al. Ubiquitination is
essential for recovery of cellular activities after heat shock. Science. 2021 June
25;372(6549):eabc3593.

12. Marmor-Kollet H, Siany A, Kedersha N, Knafo N, Rivkin N, Danino YM, et al.
Spatiotemporal Proteomic Analysis of Stress Granule Disassembly Using APEX Reveals
Regulation by SUMOylation and Links to ALS Pathogenesis. Mol Cell. 2020 Dec
3;80(5):876-891.e6.

13. Verde EM, Antoniani F, Mediani L, Secco V, Crotti S, Ferrara MC, et al. SUMO2/3
conjugation of TDP-43 protects against aggregation. Sci Adv. 2025 Feb
21;11(8):eadq2475.

14. Gwon Y, Maxwell BA, Kolaitis R-M, Zhang P, Kim HJ, Taylor JP. Ubiquitination of G3BP1
mediates stress granule disassembly in a context-specific manner. Science. 2021 June
25;372(6549):eabf6548.

15. Tolay N, Buchberger A. Comparative profiling of stress granule clearance reveals
differential contributions of the ubiquitin system. Life Sci Alliance [Internet]. 2021 May;4(5).
Available from: http://dx.doi.org/10.26508/lsa.202000927

16. Yang C, Wang Z, Kang Y, Yi Q, Wang T, Bai Y, et al. Stress granule homeostasis is
modulated by TRIM21-mediated ubiquitination of G3BP1 and autophagy-dependent
elimination of stress granules. Autophagy. 2023 July;19(7):1934–51.

17. Pan CR, Knutson SD, Huth SW, MacMillan DWC. µMap proximity labeling in living cells
reveals stress granule disassembly mechanisms. Nat Chem Biol. 2025 Apr;21(4):490–500.

18. Tolay N, Buchberger A. Role of the ubiquitin system in stress granule metabolism. Int J Mol
Sci. 2022 Mar 26;23(7):3624.

19. Kedersha N, Panas MD, Achorn CA, Lyons S, Tisdale S, Hickman T, et al. G3BP-Caprin1-
USP10 complexes mediate stress granule condensation and associate with 40S subunits. J
Cell Biol. 2016 Mar 28;212(7):845–60.

20. Clague MJ, Heride C, Urbé S. The demographics of the ubiquitin system. Trends Cell Biol.
2015 July;25(7):417–26.

21. Chernov KG, Barbet A, Hamon L, Ovchinnikov LP, Curmi PA, Pastré D. Role of
microtubules in stress granule assembly. J Biol Chem. 2009 Dec;284(52):36569–80.

22. Hu S, Zhang Y, Yi Q, Yang C, Liu Y, Bai Y. Time-resolved proteomic profiling reveals
compositional and functional transitions across the stress granule life cycle. Nat Commun.
2023 Nov 27;14(1):7782.

23. Van Treeck B, Parker R. Principles of stress granules revealed by imaging approaches.
Cold Spring Harb Perspect Biol. 2019 Feb 1;11(2):a033068.

24. Wheeler JR, Matheny T, Jain S, Abrisch R, Parker R. Distinct stages in stress granule
assembly and disassembly. Elife [Internet]. 2016 Sept 7;5. Available from:
http://dx.doi.org/10.7554/elife.18413

25. Qian S-B, Princiotta MF, Bennink JR, Yewdell JW. Characterization of rapidly degraded
polypeptides in mammalian cells reveals a novel layer of nascent protein quality control. J
Biol Chem. 2006 Jan 6;281(1):392–400.

A

	No stress	90 min heat stress	60 min recovery
DMSO			
TAK243			

B Heat stress (43°C) | Recovery (37°C)

SG number/cell number vs Time (min)

DMSO

TAK243

****

C SG number/cell number

DMSO | TAK243

**

D SG number/cell number

DMSO | TAK243

**

E Pre-bleach | 0s | 10s | 30s

F Fluorescence recovery vs Time (s)

DMSO

TAK243

****

Figure 1. Ubiquitination affects stress granule dynamics and biophysical properties. A. Representative images of U2OS cells overexpressing G3BP1-mCherry with G3BP1/2 KO background, without (DMSO) or with TAK243 (1uM) after 90 min of heat stress at 43 °C (HS) or during recovery at 37 °C for 1h. Scale bar = 20uM. **B.** Quantification of SG number/cell number as a function of time. Data were normalized internally to the time point with maximal SG assembly. ~1000 SG counted per experimental repeat. Nine fields per condition. Data from one experiment out of three experimental repeats (Supplementary fig. 1). Two-way ANOVA with repeated measures. **C.** Quantification of SG number/cell number at 15 min of HS. Average of three experimental repeats shown. Welch's t-test. **D.** Quantification of SG number/cell number at 60 min recovery. Average of three experimental repeats shown. Welch's t-test. **E.** A temporal image series from a FRAP experiment on G3BP1-mCherry U2OS cells under heat stress. Arrow indicates photobleached SG. Scale bar = 5uM. **F.** Quantification of fluorescence recovery after photobleaching in U2OS cells without or with TAK243. Data from one experimental repeat out of three is shown (Supplementary fig. 6). ~20 SGs were counted per experiment. Two-way ANOVA with repeated measures. P value * <0.05; ** <0.01, ****<0.0001.

Figure 2. Ubiquitination affects stress granule proteomic composition. A. Volcano plot of proteins enriched at the proximity of G3BP1-APEX relative to NES-APEX cytoplasm control (x-axis = Log2 fold change) and statistical significance (y-axis = Log10 p-value). G3BP1/2 KO cells expressing G3BP1-APEX were used to label SG proteins. Induction media contained Biotin phenol, 500uM, and H2O2, 1mM. Labeling activity was quenched 60 seconds after H2O2 addition. **B.** Volcano plot displaying the ubiquitin-dependent enrichment in SG proteins. 83 proteins enriched and 215 proteins depleted from SGs following TAK243 treatment. (x-axis = LogFC; y-axis = log10 p-value). Several E3 ligases, ubiquitin hydrolases, and chaperones (shown in red) are depleted. **C.** Table showing proteostasis factors depleted from SGs upon inhibition of ubiquitination. **D.** Representative images of G3BP1-mCherry cells treated with TAK243 and stained for endogenous MKRN2. **E.** Quantification of the partition coefficient of MKRN2 in SGs in cells treated with DMSO or TAK243. The proportion of SGs with PC>1.5 is shown. Average of three experimental repeats, ~1000 SGs per repeat. Welch's two-sided t-test, p-value * = <0.05; **<0.01, ****<0.0001.

Figure 3. MKRN2 promotes SG assembly and increases SG ubiquitin content.

A. Representative images of G3BP1-mCherry cells with siMKRN2 vs siCtrl before HS and at 90 min HS at 43 °C. **B.** Quantification of SG number/cell number from 30 min to 90 min HS at 43 °C. Data were normalized internally to the time point with maximal SG assembly. ~1000 SG counted per experiment. Nine fields per condition. Data from one experimental repeat out of five independent experiments. Two-way ANOVA with repeated measures. **C.** Quantification of SG number/cell number at 90 min HS. Average of five experimental repeats shown. Welch's t-test. **D.** Average SG area over the 90 min HS at 43 °C. Data from one experiment out of five experimental repeats. (Supplementary fig. 3). Two-way ANOVA with repeated measures. **E.** Average SG area at 90 min HS at 43 °C. Data were internally normalized to siCtrl. The average of five experimental repeats is shown. One-sample t-test. **F.** Quantification of fluorescence recovery after photobleaching in U2OS cells with either siMKRN2 or siCTRL. Data from one experimental repeat out of three is shown. (Supplementary fig. 6). ~20 SGs were counted per experiment. Two-way ANOVA with repeated measures. **G.** Confocal micrographs of G3BP1-mCherry/MKRN2-GFP U2OS cells stained for poly-ubiquitin (FK2). **H.** FK2 partition coefficient quantification in SGs from either MKRN2-GFP+ vs MKRN2-GFP- cells within the same coverslip. The proportion of SGs with PC>1.5 is shown. Average of 5 experimental repeats shown. >1000 SGs per repeat. p-value * = <0.05; **<0.01, ****<0.0001.

A

	No stress	90 min heat stress	30 min recovery
siCTRL			
siMKRN2			

B

siMKRN2
****
siCTRL

SG number/cell number vs Time in recovery (min)

C

30 min recovery

60 min recovery

SG number/cell number

siMKRN2 ** siCTRL

D

30 min recovery

60 min recovery

%cells with SGs

siMKRN2 ** siCTRL

Figure 4. MKRN2 promotes SG disassembly. A. Representative images of G3BP1-mCherry cells with siMKRN2 vs siCtrl before HS, at 90 min HS at 43 °C and at 30 min recovery at 37 °C. **B.** Live imaging quantification of SG number/cell number during recovery at 37 °C. Data were internally normalized to 90 min of HS. 9 sites per condition. Data of one experiment out of five experimental repeats. Two-way ANOVA with repeated measures. **C.** Quantification of SG number/cell number at 30 min or 60 min recovery for siMKRN2 vs siCTRL. Average of five experimental repeats shown. Welch's t test. **D.** Quantification of percent cells with SGs at 30 min or 60 min recovery for siMKRN2 vs siCTRL. Average of five experimental repeats shown. Welch's t-test. p-value * = <0.05; **<0.01, ****<0.0001.

A

mCherry-G3BP1   DRiPs   merge

HS + OP-puro

TAK-243 + HS + OP-puro

B

%SGs with DRiP PC>1.5

**

DMSO   TAK243

C

mCherry-G3BP1   DRiPs   merge   zoom

siRNA control: HS + OP-puro

siRNA MKRN2: HS + OP-puro

D

%SGs with DRiP PC>3

**

siMKRN2   siCTRL

Figure 5. Ubiquitination is required for preventing DRiP accumulation in SGs. A. Representative images of DMSO vs TAK243 G3BP1-mCherry U2OS cells treated with 25 µM OP-puro to label DRiPs and heat-stressed at 43 °C for 60 min to induce SGs. **B.** Quantification of the partition coefficient of DRiPs in SGs from cells treated with either DMSO or TAK243. The proportion of SGs with PC>1.5 is shown. Welch's t-test. **C.** Representative images of siMKRN2 vs siCTRL G3BP1-mCherry U2OS cells treated with 25 µM OP-puro to label DRiPs and heat-stressed at 43 °C. **D.** Quantification of the partition coefficient of DRiPs in SGs from cells treated with either siMKRN2 or siCTRL. The proportion of SGs with PC>1.5 is shown. Welch's t-test. **<0.01

Supplementary figure 1. SG disassembly live imaging repeats with TAK243. Live imaging quantification of SG number/cell number during recovery at 37°C. Data were internally normalized to 90 min of HS. 9 sites per condition. Three experimental repeats shown. Two-way ANOVA with repeated measures. ****<0.0001.

A

G3BP1-APX TAK243	G3BP1-APX DMSO	NES-APX TAK243	NES-APX DMSO	No biotin
R1 R2	R1 R2	R1 R2	R1 R2	

G3BP1-APX TAK243	G3BP1-APX DMSO	NES-APX TAK243	NES-APX DMSO	No biotin
R3 R4	R3 R4	R3 R4	R3 R4	

B

G3BP1-APX TAK243	G3BP1-APX DMSO	NES-APX TAK243	NES-APX DMSO
R1 R2	R1 R2	R1 R2	R1 R2

G3BP1-APX TAK243	G3BP1-APX DMSO	NES-APX TAK243	NES-APX DMSO	No biotin
R1 R2	R1 R2	R1 R2	R1 R2	

Supplementary figure 2. APEX samples. A. Anti-Biotin Western blot for samples used in APEX proteomics experiment. **B.** Ant-ubiquitin (FK2) Western blot for samples used in APEX proteomics experiment.

Supplementary figure 3: SG assembly live imaging repeats with MKRN2 KD. Live imaging quantification of SG area during HS at 43°C. 9-12 sites per condition. Four experimental repeats shown (fifth repeat shown in Fig. 3C). Two-way ANOVA with repeated measures. ****<0.0001.

Supplementary figure 4. SG disassembly live imaging repeats with MKRN2 KD. Live imaging quantification of SG number/cell number during recovery at 37°C. Data were internally normalized to 90 min of HS. 9-12 sites per condition. Four experimental repeats shown (fifth repeat shown in Fig. 4B). Two-way ANOVA with repeated measures. ****<0.0001.

Supplementary figure 5. Live imaging with MKRN2, ZNF598 and CNOT4. A. SG number/cell number during stress recovery (normalized to 90 min HS). **B.** Average SG area over time in HS.

A

B

Supplementary figure 6: FRAP repeats. A. Quantification of fluorescence recovery after photobleaching in TAK243 vs control cells. Two experimental repats shown (third repeat in Fig. 1F). ~20 SGs per repeat. **B.** Quantification of fluorescence recovery after photobleaching in TAK243 vs control cells. Two experimental repats shown (third repeat in Fig. 3F). ~20 SGs per repeat. Two way ANOVA with repeated measures. ****<0.0001.