Loss of PIMREG impairs double-strand break signaling and repair

Maria Vitória de Rizzo Gasparini[1,a]; Laís do Carmo[1,a]; Giovanna Gonçalves de Almeida[1];

Mariana Siqueira Lacerda Mamede[1]; Isabela Spido Dias[1]; Jennifer Ann Black[1,2]; Vanessa

Cristina Arfelli[3]; Yun-Chien Chang[4]; Stephanie Heinzlmeir[4]; Bernhard Kuster[4], Philipp A.

Greif[3,5,6]; Leticia Fröhlich Archangelo[1,*]

[1] Department of Cellular and Molecular Biology and Pathogenic Bioagents, Ribeirão Preto

Medical School, University of São Paulo (FMRP-USP), Ribeirão Preto, São Paulo, Brazil.

[2] University of Glasgow Centre for Parasitology, School of Infection and Immunity, Sir

Graeme Davies Building, 120 University Place, Glasgow, G12 8TA, United Kingdom

[3] Laboratory for Experimental Leukemia and Lymphoma Research, Munich University

Hospital, Ludwig-Maximilians University (LMU), Munich, Germany

[4] Chair of Proteomics and Bioanalytics, School of Life Sciences Weihenstephan, Technical

University of Munich (TUM), Freising, Germany

[5] German Cancer Consortium (DKTK), partner site Munich

[6] German Cancer Research Center (DKFZ), Heidelberg, Germany

[a] These authors contributed equally to this work

*Corresponding Author: Leticia Fröhlich Archangelo

Department of Cellular and Molecular Biology and Pathogenic Bioagents, Ribeirão Preto

Medical School, University of São Paulo,

Av Bandeirantes 3900, 14049-900 Ribeirão Preto-SP, Brazil. Tel: +55 (16) 33153118; fax:

+ 55 (16) 3315 0728

E-mail: leticiafa@fmrp.usp.br (L.F. Archangelo)

Abstract

Glioblastoma (GBM) is the most common primary brain tumor in adults. It is an aggressive

type of tumor with an average overall survival of 12-15 months post-diagnosis. Standard

treatment involves surgical resection, followed by radiotherapy and chemotherapy with

temozolomide (TMZ). However, resistance to treatment is frequently observed. PIMREG

is a marker of proliferation highly expressed across different tumor types, with particularly

elevated levels in GBM. Our previous study revealed that PIMREG silencing sensitizes

GBM cells to TMZ treatment and affects ATM activation, a key apical kinase involved in

DNA damage signaling. Given the emerging significance of PIMREG in DNA damage

response, we sought to explore PIMREG's role in DNA damage repair and signaling in

GBM cells exposed to genotoxic agents. Here we assessed the functionally enriched

pathways associated with high *PIMREG* expression in GBM patient samples, analyzed the

interactome of PIMREG upon treatment of GBM cells with TMZ and evaluated the impact

of PIMREG on DNA damage signaling alongside its function in the repair of double-strand

breaks. Our findings reveal that PIMREG functions as a direct substrate for ATM/ATR

kinases, interacts with key DNA repair proteins upon TMZ treatment and actively facilitates

the cellular response to chemotherapy- and radiotherapy-induced DNA damage. Taken

together, PIMREG may function during DNA damage signaling, specifically during the

repair of double strand breaks via homologous recombination in GBM cells. In all, our data

suggested high PIMREG expression may contribute to the resistance of GBM cells to

treatment.

48
49 **Key Words:** CATS, FAM64A, GBM, DSBs, HR, TMZ, IR

1. Introduction

Diffuse gliomas are the most common primary malignant brain tumors [1]. Among gliomas, GBM grade 4 is the most aggressive subtype, characterized by rapid progression and poor prognosis [2,3]. The median survival time for patients with GBM is approximately 12 to 15 months following diagnosis, with a 5-years survival rate of 9.8 % [4]. The standard treatment consists of surgical resection followed by radiotherapy with ionizing irradiation (IR) and chemotherapy with TMZ [5].

Treatment with IR and TMZ is based on inducing DNA damage in tumor cells, which promotes cytotoxicity and cell death. Whereas IR mostly induces double-strand breaks (DSBs) in the DNA through radiolysis and formation of reactive oxygen species (ROS) [6], TMZ primarily alkylates DNA bases. O6-methylguanine (O6-MeG) is the most cytotoxic lesion and is repaired by the direct repair mechanism mediated by the O6-methylguanine-DNA methyltransferase (MGMT) [7]. Thus, hypermethylation of the promoter region of the *MGMT* gene predicts response to TMZ treatment [8]. In this scenario, inefficient repair of O6-MeG leads to futile cycles of mismatch repair (MMR), replication fork collapse and ultimately DSB formation [9].

DSBs are highly cytotoxic DNA lesions that threaten genome stability if unresolved. In response, eukaryotic organisms possess a multitude of pathways orchestrated through a network of proteins to tackle these severe lesions. DSBs however, are primarily resolved through the activation of two main pathways: homologous recombination (HR) [10] and non-homologous end joining (NHEJ) [11]. Overcoming DSB lesions through DNA repair is a critical mechanism for acquiring resistance to therapy. In fact, elevated activities of HR and NHEJ are observed in patient-derived GBM samples with acquired TMZ resistance, suggesting that these pathways play a central role in the development of treatment resistance [12].

PICALM-interacting mitotic regulator (*PIMREG*) plays a well-established role in cell cycle

progression, specifically in proliferation and chromosome segregation during cell division

[13–15]. Additional functions have been proposed for PIMREG in the developing central

nervous system (CNS) [13], cell differentiation [15–17], fetal cardiomyocyte cell cycle

progression [18,19], and in signaling pathways such as JAK-STAT3 [17], NF-κB [20], and

β-catenin [21,22]. PIMREG expression levels strongly correlate with cellular proliferation

in both normal and malignant cells [13].

The contribution of PIMREG in promoting malignant phenotypes has been primarily

addressed in breast cancers, where multiple functional studies have linked PIMREG to cell

migration, metastasis, epithelial-mesenchymal transition, enhanced stemness features, and

aggressiveness of breast cancer cells [20,22,23]. In another hormone-sensitive cancer, such

as prostate cancer (PCa), PIMREG was described as an oncogenic factor whose expression

became activated through direct binding of the androgen receptor (AR) to the *PIMREG*

promoter, leading to proliferation, colony formation, migration, invasion, and cell cycle

progression of PCa cells [24].

The oncogenic role of PIMREG in cancer development has been shown *in vivo* by three

elegant studies. Jiang and colleagues demonstrated that overexpression of PIMREG

promotes breast cancer aggressiveness via constitutive activation of NF-κB signaling

pathway. Mechanistically, PIMREG overexpression sustains constitutive activation of NF-

κB signaling by competing with IkB for interaction, thereby disrupting the NF-κB/IκBα

negative feedback loop. In cells expressing high levels of PIMREG, such as breast cancer

cells, PIMREG interacts with the p65 subunit of NF-κB within the nucleus, thereby

stabilizing p65, and enhancing its binding to NF-κB response elements, promoting

activation of its target genes and consequently hyperactivation of NF-κB signaling that

correlates with cancer development and progression [20].

The PIMREG-mediated activation of the NF-κB pathway is also implicated in promoting

cisplatin resistance in triple-negative breast cancer (TNBC) both *in vivo* and *in vitro* [25].

For instance, SOD1-high fibroblasts produce exosomal miR-3960 that targets the kinase

BRSK2 in TNBC cells. BRSK2 is responsible for phosphorylating PIMREG at Ser16

leading to PIMREG ubiquitination and subsequent degradation. Due to miR-3960-mediated

downregulation of BRSK2 , PIMREG Ser16 phosphorylation becomes reduced thereby

increasing protein stability. Consequently, NF-κB signaling is activated correlating with

cisplatin resistance in TNBC cells. Conversely, decreased PIMREG levels due its

phosphorylation of Ser16, enhances cisplatin sensitivity in TNBC cells [25].

In a third study, Xu and colleagues revealed PIMREG's critical role in inflammation-

associated tumorigenesis. Mice lacking *Pimreg* exhibited impaired differentiation of Th17

lymphocytes, leading to reduced proinflammatory activity typically associated with these

cells. PIMREG is essential for IL-6–mediated activation of STAT3 and regulates Th17 cell

differentiation by modulating STAT3 transcriptional activity. As a result, PIMREG plays a

key role in the development of Th17-driven inflammatory diseases, including autoimmune

encephalomyelitis, colitis and colitis-associated carcinogenesis in animal models [17].

Consistent with its role as a proliferation marker, *PIMREG* is highly expressed in tumor

tissues compared to normal tissue samples. Additionally, high expression of *PIMREG* in

patient samples correlated with worse prognosis and reduced overall survival across various

types of cancer [26,27]; including breast cancer [20,22,23,28]; pancreatic cancer [29]; lung

cancer [30]; prostate cancer [24]; head and neck squamous cell carcinoma [31]; clear cell

renal cell carcinoma [32]; gastric cancer [33]; melanoma [34] and gliomas [35].

We have previously shown that GBM exhibits the highest level of *PIMREG* expression

amongst 25 types of cancer. As well as showing that PIMREG silencing reduced malignant

characteristics of GBM cells, such as cell cycle progression, proliferation and migration

capacity, we demonstrated that treating these cells with TMZ increased the levels of

PIMREG protein. In addition, PIMREG silencing rendered GBM cells more sensitive to

TMZ exposure and interfered with ATM activation, a key apical kinase that orchestrates

the cellular response to DSBs. We hypothesized that PIMREG plays a role in DNA damage

response (DDR) and therefore could contribute to TMZ resistance [35].

The mechanism by which PIMREG participates in the DDR in GBM cells is unclear. To

begin addressing these knowledge deficits and gain further insight into PIMREG function

and relevance in GBM, we tested for functionally enriched pathways related to elevated

PIMREG expression in GBM patient samples and analyzed the correlation between

PIMREG expression and those of DDR-related genes. We investigated the interactome of

PIMREG upon treatment of GBM cells with TMZ and evaluated the impact of PIMREG

on DNA damage signaling alongside its function in the repair of DSBs. Our data reveal that

PIMREG interacts with proteins involved mainly in DDR and RNA metabolism following

TMZ treatment in GBM cells. Additionally, PIMREG serves as a substrate for ATM/ATR

and plays a role in DNA damage signaling and the repair of DSBs. These findings further

suggest the importance of elevated expression of PIMREG in GBM maintenance and its

possible contribution to the treatment resistance in GBM patients.

2. Material and methods

2.1. Public data analysis

GBM RNAseq-based mRNA expression (RSEM) data were downloaded from TCGA-PanCancer Atlas_2018 (592 samples) using cBioPortal for Cancer Genomics [36].

Gene set enrichment analysis was performed using GSEA version 4.3.2 [37]. Results were regarded as significant with a p-value < 0.05 and 25 % false discovery rate (FDR) threshold (q-value <0.25).

Genes coding for DDR proteins, whose expression covariates with *PIMREG* in GBM patient samples were plotted in a correlation curve using GraphPad Prism 9.

PIMREG mRNA expression data, along with *MGMT* promoter methylation status, were also obtained from the merged cohorts of Brain Lower Grade Glioma (LGG) (TCGA, PanCancer Atlas, 514 samples) and GBM (TCGA, PanCancer Atlas, 592 samples). Among the 1106 available samples, 635 contained information regarding the methylation status of the *MGMT* promoter.

2.2. Cell lines

The GBM human cell line T98G was obtained from the American Type Culture Collection (ATCC, Manassas, USA) and U87MG from the Banco de Células do Rio de Janeiro (BCRJ, Rio de Janeiro, Brazil). The reporter cell lines HeLa EJ5-GFP and U2OS DR-GFP were kindly provided by Dr. Valeria Valente (School of Pharmaceutical Science, UNESP, Brazil). The cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM, Gibco) supplemented with 10 % of fetal bovine serum (FBS, Gibco) and 1 % of the antibiotic penicillin/streptomycin (Gibco), at 37 °C and 5 % of CO_2. All cell lines were tested for mycoplasma with MycoAlert PLUS Mycoplasma Detection Kit (Lonza). T98G were authenticated by STR matching analysis using the GenePrint 10 system (Promega,

Madison, WI, USA) and 3730 DNA Analyzer - Applied Biosystems (Applied Biosystems,

Foster City, CA, USA).

2.3. Cell treatment with genotoxic agents

Temozolomide (TMZ, Sigma-Aldrich # T2577 and Selleckchem S1237) were diluted in

DMSO at a stock concentration of 50 mM and 200 mM, respectively. Stocks were stored

in 500 μL aliquots at - 80 ºC.

GBM cells were treated with 300 μM TMZ or vehicle (DMSO) for 24 h, for Western blot

analysis, cell cycle, apoptosis, immunoprecipitation and immunofluorescence experiments.

For IR treatment, GBM cells were irradiated using the PRIMUS linear accelerator (Siemens

Healthineers, Brazil) with a 6 MV photon beam at the Hospital das Clínicas - Ribeirão Preto

Medical School. The administered dose was 4 Gy. Cells were collected 0.5 h and 1 h post-

irradiation for Western blot analysis and immunofluorescence experiments. The

concentration and time used for the TMZ and IR treatments were defined as such to induce

DNA damage signaling but not cell cycle arrest or apoptosis. Cell cycle progression and

viability were monitored in all GBM cell samples treated with genotoxic agents

(Supplementary Fig. S1C, S1D and Fig. S2D).

2.4. Plasmid construction

For genome editing, single guide RNAs (sgRNAs) targeting exon 2 of *PIMREG* were

designed   using   the   Benchling   CRISPR   sgRNA   design   tool

(https://www.benchling.com/crispr). The 20 bp oligos contained the restriction sites CACC

and AAAC at the 5'end of the sense and antisense oligos, respectively, as described in [38].

Annealed oligos were cloned into the PX458 vector (Addgene plasmid #48138) digested

with the *BpiI* restriction enzyme (Thermo Fisher Scientific) generating the PX458-

PIMREG-KO vectors (KO-1, KO-2, KO-3 and KO-4). The target sequences are as follows:

sgRNA-1   -   sense: GAGAGATCTCCGGCGCACGG   and   antisense:

CCGTGCGCCGGAGATCTCTC; sgRNA-2 - sense: CTTCTCGGTGGCAGAACATG

and antisense: CATGTTCTGCCACCGAGAAG; sgRNA-3 - sense:

CCATCAGGAGACCTCTGTAG and antisense: CTACAGAGGTCTCCTGATGG;

sgRNA-4 - sense: TCAACCTCAACCTCCGCGCA and antisense:

TGCGCGGAGGTTGAGGTTGA.

2.5. Generation of CRISPR/Cas9 edited monoclonal cell lines

The GBM T98G cells and the reporter cell lines HeLa EJ5-GFP and U2OS DR-GFP were

edited to generate PIMREG knockout cell lines (PIMREG KO-1, KO-2, KO-3, KO-4). 2 -

3×10^5 cells were seeded at 6-well plates and transfected with 2.5 µg of PX458-PIMREG-

KO vectors and Lipofectamine 2000 according to manufacturer's instructions

(ThermoFisher). Efficiently transfected cells were single sorted by Fluorescence-Activated

Cell Sorting (FACS) based on GFP expression. The individual cells were cultured in 96

well plates for clonal expansion and generation of the monoclonal cell lines. PIMREG KOs

were confirmed by Western blot analysis and genomic sequencing of the edited *PIMREG*

locus.

The presence of phenotypic variability among the different PIMREG KO clones, is a

common limitation inherent to studies relying on single-cell CRISPR/Cas9-generated KOs.

Particularly intriguing, was the unexpected re-expression of PIMREG protein after

extended culture of the GBM T98G PIMREG KO-2 cell line. This finding prompted us to

routinely verify PIMREG absence by immunofluorescence in all KO cell lines throughout

the course of the experiments. Importantly, the PIMREG KO-2 cell line that recovered

PIMREG protein expression (KO-2 recPIMREG) was subsequently used as a control in

experiments analyzing γH2AX fluorescence intensity in PIMREG KO and WT cells

following IR exposure.

2.6. Cell cycle analysis

1×10^6 cells were fixed in 70 % ethanol for 30 min on ice, washed with PBS, treated with RNAse A solution (100 µg/ml) and stained with 100 µg/ml propidium iodide (PI). Cell cycle analysis was performed using FACS Symphony A1 (BD Biosciences) and FlowJo software version 10. Statistical analyses were performed with Two-way ANOVA, using GraphPad Prism 9.

2.7. Cell viability analysis

Cell viability was assessed with either Annexin V or Zombie NIR staining. For the Annexin V staining, cells were washed with the Annexin V Binding Buffer (BD Pharmingen™) and incubated with 5 µl Annexin V reagent (BD Pharmingen™) for 15 min at RT followed by 2.5 µL of PI solution (Miltenyi Biotec) (100 µg/mL). For the Zombie NIR staining, cells were incubated with Zombie NIR reagent (BioLegend) (1:400) for 15 min, washed with PBS, fixed with 2 % paraformaldehyde (PFA) for 15 min and resuspended in 1 % glycine/PBS. Cell viability was analyzed using FACS Symphony A1 (BD Biosciences) and FlowJo software version 10.

2.8. Immunoblotting and Immunoprecipitation

Cells were lysed in lysis buffer with protease inhibitors (10 mM EDTA, 100 mM Tris-HCl pH 7.5; 1 % Triton X-100, 10 mM Sodium Pyrophosphate ($Na_2P_2O_7$), 100 mM Sodium Fluoride (NaF), 10 mM Sodium Orthovanadate (Na_3VO_4); 2 mM PMSF). Protein extracts were separated by electrophoresis on 12 % or on a gradient 2.5-12.5 % SDS polyacrylamide (SDS-PAGE) gel and transferred to a nitrocellulose membrane (GE Healthcare Life Science). Membranes were blocked with 1 % BSA or 5 % low fat milk and probed with a specific antibody, followed by incubation with a secondary antibody conjugated to horseradish peroxidase (HRP). Protein signals were detected by chemiluminescence and the images were captured by ChemiDoc™ Imaging System (Bio-Rad). Primary antibodies

were: anti-PIMREG (anti-CATS 2C4) (1:10) (Archangelo et al., 2008), anti-β-Actin

(ThermoFisher Scientific, # AM4302 and Santa Cruz Biotechnology sc-1616) (1:10000

and 1:2000, respectively), anti-α-tubulin (Santa Cruz Biotechnology sc-5286) (1:1000),

anti-phospho-H2AX (S139) (Cell Signaling Technology, #2577 and #80312) (1:500 and

1:1000, respectively), anti-phospho-ATM (S1981) (Sigma-Aldrich, clone 10H11.E12)

(1:1000), anti-ATM (Sigma-Aldrich, clone MAT3-4G10/8) (1:1000), anti-phospho-DNA-

PKcs (S2056) (Cell Signaling Technology, #68716) (1:1000), anti-DNA-PKcs (Cell

Signaling Technology, #38168) (1:1000), anti-phospho-CHK1 (S345) (Cell Signaling

Technology, #2348) (1:1000), anti-CHK1 (Cell Signaling Technology, #2360) (1:1000),

anti-phospho-KAP1 (S824) (Abcam, ab243870) (1:1000), anti-KAP1 (Abcam, ab109545)

(1:1000) and anti-phospho-ATM/ATR substrate (Phospho-(Ser/Thr) ATM/ATR Substrate;

Cell Signaling Technology, #2851) (1:1000). The secondary antibodies were anti-mouse

(Santa Cruz Biotechnology, sc-2064) (1:3000), anti-rat (KPL, 52200364) (1:3000), anti-

goat (Santa Cruz Biotechnology, sc-2354) (1:3000) and anti-rabbit (Abcam, ab6721)

(1:3000). The blots were stripped with Restore Western Blot Stripping Buffer (Thermo

Scientific, YE366939) before reprobing. Quantitative analyses of the optical intensities of

the protein bands were determined with UN-SCAN-IT gel 6.1 software (Silk Scientific Inc.,

UT) and normalized by control protein.

For immunoprecipitation, 170 - 290 μg of cellular extracts were incubated overnight with

50 μl of monoclonal rat anti-PIMREG (anti-CATS 2C4), 1:20 (v/v) rabbit anti-phospho-

ATM/ATR substrate antibodies and respective isotype controls (1.6 μg of rat IgG Santa

Cruz Biotechnology, sc-2026 and 21 μg of rabbit IgG, Jackson ImmunoResearch

Laboratories, Inc., 011-000-003) at 4 ºC, under rotation. The next day, 40 μl of Protein G-

Agarose (Roche) was added and incubated further for 3 h. Beads were extensively washed

with lysis buffer (50 mM Tris-HCl pH 8, 150 mM NaCl, 0.1 % SDS, 1 % IGEPAL, and 0.5

% sodium deoxycholate), resuspended in SDS sample buffer, boiled, and separated on a 12

% SDS-PAGE gel. Subsequent Western blot analysis was carried out with the antibodies

anti-PIMREG (anti-CATS 2C4) and anti-phospho-ATM/ATR substrate.

2.9. Immunofluorescence

Cells grown on coverslips were treated with TMZ and IR, fixed and permeabilized by

sequential incubation in 1 % PFA/ 0.5 % Triton X-100 followed by 1 % PFA/ 0.3 % Triton

X-100 and 0.5 % methanol for 20 min and blocked for 1 h with blocking solution (0.5 %

Triton X-100, 1 mM EDTA, 1 mg/mL BSA, and 0.03 % FBS) at RT. Coverslips were

incubated for 1 h with the primary antibodies diluted in blocking solution: anti-PIMREG

(Anti-CATS-2C4) (1:10) and anti-phospho-H2AX (S139) (Cell Signaling Technology,

#80312) (1:100). Secondary Alexa Fluor-conjugated antibodies Alexa Fluor anti-rat 488

(Invitrogen, A-11006) (1:500), Alexa Fluor anti-mouse 555 (Invitrogen, A21325) (1:600)

and Alexa Fluor anti-rabbit 594 (Thermo Fisher Scientific, A32754) (1:600) were used for

detection of the primary antibodies. Nuclei were counterstained with 300 nM DAPI solution

and coverslips were mounted on slides using Fluoromount-G™ Mounting Medium

(Invitrogen).

Images were acquired using a Leica SP5 confocal microscope and a Leica CTR 6000

Fluorescence Microscope equipped with LAS AF software, employing 100x magnification

in z-stack mode. γH2AX fluorescence intensity was quantified using ImageJ Fiji software

(version 2.14). Corrected Total Cell Fluorescence (CTCF) was calculated for each nucleus

(CTCF = IntDen - (mean background × area)). A minimum of 75 cells were analysed per

condition per experiment.

For colocalization, line scan analysis was performed using the "plot profile" tool in ImageJ

Fiji (version 2.14). Fluorescence intensity values (gray values) across distance (μm) for

each channel were plotted as line graphs in GraphPad Prism 9.

2.10. DNA repair assay

The frequency of HR and NHEJ were estimated using U2OS DR-GFP and HeLa EJ5-GFP reporter cells [39,40]. 3 x 10^5 HeLa EJ5-GFP and 1 x 10^5 U2OS DR-GFP cells WT and PIMREG KOs (KO-1, -2, -3 and -4) were seeded at 6-well plates and transfected with 2.5 μL of the *I-SceI* endonuclease expressing plasmid, which creates the DSBs (pCBASceI; Addgene Plasmid #26477). Cells were collected 48 h after transfection and the percentage of GFP-positive cells was determined by flow cytometry using FACS Symphony A1 or Accuri C6 (BD Biosciences) and FlowJo software. Dead cells were excluded from the analysis using Zombie NIR (BioLegend) staining at proportion 1:400. Fifty thousand events were acquired. The percentage of GFP-positive cells of each sample was normalized by the mean value of WT GFP-positive cells used as internal control for each experiment.

2.11. Mass spectrometry analysis

PIMREG was immunoprecipitated from cellular extracts of U87MG (900 μg) and T98G (500 μg) cells treated with TMZ and DMSO using the anti-PIMREG (anti-CAST2C4) antibody and protein G magnetics beads (PureProteome™ Protein G Magnetic Beads, Merck). Normal rat IgG antibody (Santa Cruz Biotechnology) was used as isotype control. The experiments were performed in triplicates (Supplementary Fig. S1).

Immunoprecipitated proteins were reduced with 10 mM dithiothreitol (DTT) at 37 °C for 45 min, then alkylated with 55 mM chloroacetamide at RT for 30 min in the dark. Samples were mixed with NuPAGE LDS sample buffer (Thermo Fisher Scientific), incubated at 70 °C for 10 min, and loaded onto a NuPAGE Bis-Tris gel by electrophoresis at 200 V for 5 min, allowing proteins to migrate approximately 1 cm into the gel. Gels were fixed in 2 % acetic acid and 40 % methanol and stained with Coomassie blue. Gel bands were excised, transferred to a 96-well plate, and destained with 50 mM triethylammonium bicarbonate (TEAB) in 50 % ethanol for 2 h at RT followed by 1 h at 55 °C. Gel pieces were dehydrated

with 100 % ethanol for 10 min, rehydrated in 5 mM TEAB for 20 min, and dehydrated again

with 100 % ethanol for 10 min.

For digestion, 20 µl of trypsin solution (10 ng/µl in 5 mM TEAB) was added to each gel

piece and incubated at 4 °C for 15 min. Excess of trypsin solution was removed and

replaced with 20 µl of 5 mM TEAB, and digestion was performed overnight at 37 °C.

Peptides were extracted by sequential incubations: two rounds with 20 µl of 1 % formic

acid (FA) for 30 min each, followed by extraction with 20 µl of 60 % acetonitrile (ACN)

containing 0.1 % FA for 30 min. Extracted peptides were dried and stored at -20 °C until

LC-MS analysis.

Peptides were analyzed using an Orbitrap Fusion Lumos mass spectrometer (Thermo Fisher

Scientific) coupled to a Dionex UltiMate 3000 RSLCnano system (Thermo Fisher

Scientific). Peptide separation was performed at a flow rate of 300 nL/min on a 50-min

linear gradient from 4% to 32% solvent B (0.1 % FA and 5 % DMSO in ACN) in solvent

A (0.1 % FA and 5 % DMSO in water). The MS was operated in positive ion mode using

data-dependent acquisition (Top20 method). Full MS scans (MS1) were acquired in the

Orbitrap from 360 to 1300 m/z at a resolution of 60,000 in profile mode, with an automatic

gain control (AGC) target of 4e5 and a maximum injection time (maxIT) of 50 ms. For

MS/MS (MS2) acquisition, precursor ions were isolated with a 1.7 Th window, fragmented

by higher-energy collisional dissociation (HCD) at a normalized collision energy (NCE) of

30 %, and analyzed in the Orbitrap at a resolution of 30,000 in centroid mode (AGC target

of 2e5 and maxIT of 50 ms). Dynamic exclusion was set to 20 s.

MS raw files were processed using MaxQuant (version 1.6.17.0) and searched against the

UniProtKB human reference proteome (downloaded on 2019-07-15) with default

parameters. Carbamidomethylation of cysteine was set as a fixed modification, and

oxidation of methionine as well as N-terminal protein acetylation were defined as variable

modifications. Protein groups labeled as "reverse" or "potential contaminant" were

excluded. Protein abundance was estimated based on iBAQ values, and missing values were

imputed with a constant value corresponding to a log2 iBAQ intensity of 10 to enable

downstream statistical analysis. Differential expression analysis was performed using a

two-sample t-test, and p-values were adjusted for multiple testing using the Benjamini-

Hochberg procedure.

The comparison between PIMREG immunoprecipitates vs. IgG under vehicle (DMSO)

conditions (PIMREG IP vs. IgG) identified the PIMREG-interacting partners (i.e.

significantly upregulated proteins), considered as the *bona fide* PIMREG interactome. The

comparison between PIMREG immunoprecipitates from TMZ vs. vehicle (TMZ vs.

DMSO) revealed TMZ-induced alterations in PIMREG interactions and gave rise to the

TMZ-regulated interactome. The identified proteins were filtered by significance based on

$\log_2$ fold change cutoff > 1 (PIMREG IP vs. IgG) and $\log_2$ fold change cutoff > 1 and < -1

(TMZ vs. DMSO) and adjusted *p*-value < 0.05. Data was analyzed using R software

(version 4.3.2) and Excel (Microsoft Office).

Upregulated proteins in the TMZ-regulated interactome are proteins that interact with

PIMREG only under TMZ treatment or show increased association under treatment

whereas the downregulated proteins are those with reduced or lost binding upon TMZ

treatment.

2.12. Gene Ontology and network analyses

Gene Ontology (GO) analysis was carried out using the EnrichR platform

(https://maayanlab.cloud/Enrichr/). Terms with adjusted *p*-value < 0.05 were considered

statistically significant.

PIMREG and its interacting proteins were searched for interaction networks using the

STRING database (https://string-db.org). Interactions determined experimentally,

370 registered in curated databases, extracted from text mining or coexpression data were

371 considered, using a medium confidence (0.400) interaction score. Network was clustered

372 using the MCL clustering algorithm, with an inflation parameter of 3.

3. Results

3.1. GSEA analysis implicates *PIMREG* in DNA repair and splicing

PIMREG expression correlates with genes associated with cell cycle, DNA replication and DNA repair in GBM [35]. To dissect the relationship between high *PIMREG* expression and GBM further, we performed a GSEA analysis testing for functionally enriched pathways that correlate with high expression of *PIMREG* in GBM patient samples (TCGA, PanCancer Atlas, 592 samples). Our analysis revealed 392 enriched gene sets related to high *PIMREG* expression using the HALLMARK (17 gene sets), KEGG (65 gene sets) and REACTOME (310 data sets) databases (Supplementary Table 1). In support of previous results [35], our analysis uncovered correlations between high *PIMREG* expression and the gene sets from several DNA repair pathways (33 out of 392 gene sets), including base excision repair (BER), nucleotide excision repair (NER), mismatch repair (MMR), double-strand break repair (DSB repair), homology directed repair (HDR), Fanconi anemia (FA), and UV-response (Fig. 1A, Supplementary Table 2), indicating putative functions for PIMREG across multiple DNA repair pathways in GBM.

Moreover, gene sets pertaining to RNA metabolism, such as spliceosome, RNA polymerase, RNA transport, RNA degradation, processing of capped intron-containing pre-mRNA and mRNA splicing were also abundant, accounting for 12.5 % of all gene sets (49 out of 392), suggesting an unforeseen function for *PIMREG* in RNA splicing and processing (Fig. 1B, Supplementary Table 3).

In line with putative functions for PIMREG in DDR, we performed a correlation analysis using data from the GBM patient samples uncovering a strong positive correlation between *PIMREG* transcript abundance and the abundance of transcripts from essential genes required for DSBs recognition and repair, including *H2AX*, *RAD51*, *RAD51AP1*, and *FANCD2* (Fig. 1C).

Since the methylation status of the *MGMT* promoter is a well-established predictive

biomarker for response to TMZ treatment in gliomas [41], *PIMREG* expression was

additionally assessed in glioma patient samples stratified by the methylation status of the

MGMT promoter. The analysis revealed that samples with unmethylated *MGMT* promoters,

(MGMT proficient) possessed significantly higher levels of *PIMREG* expression compared

to samples with methylated *MGMT* promoters (Fig. 1D), suggesting that PIMREG is more

abundant in the group of patients with lower response to TMZ.

Altogether, these findings support our previous analyses, suggesting wide roles for

PIMREG during the DDR but also in RNA metabolism in GBM.

3.2. PIMREG interactome in response to TMZ treatment

We have previously shown that treating GBM cells with TMZ triggers an increase in

PIMREG protein levels, as cells accumulate DNA lesions. We hypothesized that PIMREG

accumulation is required to mitigate DNA lesions induced by TMZ treatment [35].

To gain insight into the functional implications of PIMREG accumulation in response to

TMZ treatment, PIMREG was immunoprecipitated from GBM cells treated with TMZ and

its interactome identified by mass spectrometry (Supplementary Fig. S1).

To ensure reproducibility, only the proteins with peptides detected in all three replicates and

filtered by significance based on $\log_2$ fold change cutoff > 1 (PIMREG IP vs. IgG) or $\log_2$

fold change cutoff > 1 and < -1 (TMZ vs. DMSO) and adjusted *p*-value < 0.05, were

considered and identified as PIMREG interactors.

In U87MG cells, a total of 36 proteins were identified as PIMREG interacting partners (Fig.

2A and Supplementary Table 4), whereas 29 proteins were identified as TMZ-regulated

PIMREG interactors (Fig. 2B and Supplementary Table 5). In T98G cells, 8 proteins were

identified as PIMREG interacting partners (Fig. 2C and Supplementary Table 6) while 4 were

identified as TMZ-induced interacting proteins (Fig. 2D and Supplementary Table 7).

To better understand the molecular functions of PIMREG, we analyzed its protein interaction

network using the STRING database. Both upregulated and downregulated interacting

proteins were used for this analysis. This approach strongly implicated PIMREG in splicing

and ribosome biology, as protein clusters related to these functions were identified in both

PIMREG interactome and TMZ-regulated interactome of U87MG cells, albeit with distinct

sets of interactors. Additionally, a cluster of proteins related to NADH dehydrogenase

activity and one unidentified cluster of PIMREG and TRIP13 were observed in the TMZ-

regulated interactome only, whereas a cluster related to immunoglobulin complex was

observed in the PIMREG interactome only (Fig. 2E and 2F). For the T98G cells, a cluster of

proteins related to the Polycomb complex was observed within the TMZ-regulated PIMREG

interactors (Fig. 2G), whilst no cluster was formed among the proteins identified in the

PIMREG interactome.

GO analysis of the proteins identified as the TMZ-regulated interactors of PIMREG in

U87MG retrieved 10 ontological annotations whereas the proteins identified in PIMREG

interactome and TMZ-regulated interactome of T98G, retrieved 16 and 50, respectively.

These ontological annotations were classified according to their related biological processes

(BP) (Fig. 2H, 2I and 2J; Supplementary Tables 8, 9 and 10). The BP terms obtained for

TMZ-regulated PIMREG interacting proteins in U87MG cells were related to RNA

splicing/processing, cell division/proliferation and mitochondrial metabolism (Fig. 2H;

Supplementary Table 8). The BP terms obtained for PIMREG interactome in T98G cells were

related to insulin pathway, transcription/mRNA processing, immune response, protein

transport, and cell signaling (Fig. 2I; Supplementary Table 9). Lastly, the BP terms described

for TMZ-regulated PIMREG interacting proteins identified in the T98G cells were related to

cell differentiation/development, transcription/translation, macromolecule metabolic process,

telomere maintenance, IFN signaling, ribosome biogenesis, apoptosis, DNA replication and

cell division. Notably, the BP related to DDR exhibited the highest number of ontological

annotations, which included telomere maintenance- (GO:1904507), intrinsic apoptotic-

(GO:0008630), mitotic check-point (GO:0044773) signaling pathways in responses to DNA

damage as well as regulation of DNA repair (GO:0006282) and DSB repair via homologous

recombination (GO:0000724), underscoring the role of PIMREG in the DDR of T98G cells

treated with TMZ (Fig. 2J; Supplementary Table 10).

3.3. PIMREG is a substrate of ATM/ATR in GBM cells treated with TMZ

PIMREG is a phospho-protein [42] containing residues, which lie within the S/TQ motif, a

known target site for ATM and ATR phosphorylation [43]. To ask if PIMREG is a substrate

of ATR/ATM, we precipitated endogenous PIMREG from TMZ- and DMSO-treated as well

as untreated T98G GBM cells and blotted the precipitated protein against the phospho-

ATM/ATR substrate antibody (Fig. 3A) used to detect proteins containing the phosphorylated

substrate motif of ATM/ATR. Under all conditions we identified the precipitated PIMREG

protein (Fig. 3A, lower panel, lanes 3, 6 and 9). In the reciprocal immunoprecipitation

experiment, precipitated phospho-ATM/ATR substrates (Fig. 3B, upper panel) were blotted

against anti-PIMREG antibody (Fig. 3B, lower panel) and the PIMREG protein could be

detected by anti-PIMREG among the precipitated phospho-ATM/ATR substrates in TMZ-

treated cells (Fig. 3B, lower panel, lane 9). These results strongly suggest that PIMREG is a

direct substrate for ATM/ATR kinases *in vivo.*

3.4. PIMREG deletion interferes with γH2AX accumulation in GBM cells

To investigate the role of PIMREG in the cellular response to DNA damage, we generated

PIMREG knockout GBM T98G cell lines (PIMREG KO-1, KO-2, KO-3 and KO-4) and

measured γH2AX levels by Western blot to monitor DNA damage and signaling.

Quantification of γH2AX band intensity revealed that PIMREG KOs exhibited higher levels

of γH2AX than the wild-type (WT) cells (Fig. 4A and 4B; Supplementary Fig. 2A),

suggestive of higher basal DNA damage levels and/or genomic instability in the absence of

PIMREG when compared to controls.

We next examined the relative amount of DNA damage in the PIMREG KOs treated with

TMZ and IR to control cells. As expected, γH2AX signal increased in WT cells treated with

both TMZ and IR compared to DMSO or non-irradiated (NI) control cells (Fig. 4C and 4E,

middle panels).

Although PIMREG KOs exhibit higher levels of basal γH2AX than WT cells, when the

amount of γH2AX of TMZ- or IR-treated cells was normalized by those of DMSO or NI

(TMZ/DMSO or IR/NI), comparable levels of γH2AX accumulation were observed in

PIMREG KOs and WT treated with TMZ, with the exception of PIMREG KO-4 which

exhibited reduced levels of γH2AX accumulation (Fig. 4D; Supplementary Fig. 2B).

In PIMREG KOs cells treated with IR, reduced levels of γH2AX accumulation were

observed across all clones (Fig. 4F; Supplementary Fig. 2C). Taken together, these results

suggest that although PIMREG KO cells exhibit high levels of basal γH2AX, its absence

compromises the accumulation of γH2AX induced following exposure to genotoxic stress.

To complement these results, immunofluorescence experiments were performed to

examine the fluorescence intensity of γH2AX in the nuclei of PIMREG KO and WT GBM

cells treated with TMZ or exposed to IR. As shown in Figure 5, TMZ and IR treatments

enhanced γH2AX signal in both PIMREG KOs and WT cells, compared to DMSO or non-

irradiated (NI) control cells. However, the overall amount of γH2AX fluorescence intensity

appeared significantly lower in the PIMREG KOs compared to WT cells (Fig. 5), with the

exception of PIMREG KO-3 exposed to IR which exhibited similar levels of γH2AX

accumulation compared to WT. Interestingly, in one PIMREG KO cell line (KO-2), where

PIMREG protein expression was unexpectedly restored (KO-2-recPIMREG) (Fig. 5C and

5E), IR exposure led to the opposite effect, with markedly increased γH2AX accumulation

(Fig. 5D). At present we do not know the underlying mechanism of PIMREG recovery.

Nevertheless, the data suggest that the presence of PIMREG could rescue the impaired

DNA damage response phenotype observed in its absence, which is supported by the fact

that the same KO-2 cell line, prior to recovery of PIMREG (Fig. 5A), exhibited significant

reduction of γH2AX following TMZ treatment (Fig. 5B).

Taken together, our results demonstrate that PIMREG deletion impairs γH2AX signaling

in GBM cells under conditions of genotoxic stress.

3.5. PIMREG deletion compromises DNA damage signaling

We next asked whether PIMREG KO affects the activation of additional DNA

damage response signaling molecules following treatment with to TMZ or IR (Fig. 6;

Supplementary Fig. S3). Besides ATM, we analyzed molecules representing different steps

of the DNA damage signaling cascade. These include KAP1, which facilitate DSB repair

in both euchromatin and heterochromatin [44]; CHK1, a critical cell-cycle checkpoint

regulator; and DNA-PKc, a central component of DSB signaling pathway that orchestrates

NHEJ-mediated repair [45]

Western blot analysis and quantification of phosphorylated protein band intensities

revealed no differences in the activation of KAP1, CHK1 or DNA-PKc between PIMREG

KOs and WT cells after exposure to TMZ (Fig. 6A-6F). In contrast, PIMREG KO cells

exposed to IR showed a consistent trend of reduced ATM, KAP1 and CHK1 activation

compared to WT cells. This reduction was most evident at 0.5 h post-irradiation (Fig. 6G-

6L). Activation of DNA-PKc was unaffected in PIMREG KOs cells relative to WT

following IR exposure (Fig. 6M and 6N). Thus, loss of PIMREG alters DDR signaling in

irradiated GBM cells.

3.6. PIMREG deletion interferes with the repair of DSBs

ATM and γH2AX are key regulators of DSBs signaling cascade to promote repair [45,46]. Given ATM/ATR can phosphorylate PIMREG and that its deletion reduces both γH2AX and, putatively, may alter wider DDR signaling, we sought to investigate the impact of PIMREG deletion on DSB repair.

To this end, we performed DNA repair assays using HR (U2OS DR-GFP) and NHEJ (HeLa EJ5-GFP) reporter cell lines [39,40] in which PIMREG was deleted. In the DR-GFP and EJ5-GFP reporter systems, the expression of GFP protein is restored upon HR and NHEJ repair, respectively.

The absence of PIMREG significantly impaired DNA repair by HR in the U2OS DR-GFP PIMREG KOs (KO-1, KO-2, KO-3 and KO-4). Although the reduction of the HR-mediated repair was not statistically significant in the PIMREG KO-2 cell line, this trend was consistently observed across the three independent experiments (Fig. 7A). Additionally, the absence of PIMREG was associated with an increase in DNA repair by the NHEJ pathway in one out of four HeLa EJ5-GFP PIMREG KOs (KO-4), with a similar trend being observed in the PIMREG KO-3 cell line (Fig. 7B). Together, these findings suggest that PIMREG plays a role in promoting HR-mediated DSBs repair, while its absence may favor repair by NHEJ pathway.

4. Discussion

This study greatly expands our understanding of PIMREG, redefining it from a proliferation marker to an integrated component of the DDR in GBM. Our findings reveal that PIMREG functions as a direct substrate for ATM/ATR kinases, interacts with key DNA repair proteins, and actively facilitates the cellular response to chemotherapy- and radiotherapy-induced DNA damage. By demonstrating its role in signaling and repairing of DSBs, we consolidate PIMREG as a key player in maintaining genomic integrity and possibly, promoting treatment resistance.

Pathway enrichment analysis performed using GBM patient samples revealed putative functions for PIMREG across multiple DNA repair pathways and in RNA splicing and processing, finding supported by our mass spectrometry data. Specifically, we uncovered in our PIMREG interactome associations with key DNA repair factors involved in HR-mediated resolution of DSB, including YY1 in T98G cells, and TRIP13 and PRMT5 in U87MG.

YY1 is a transcription factor, which complex with chromatin-remodeling factors and is essential for HR-based DNA repair [47]. TRIP13 is an ATPase involved in early DNA damage sensing and ATM signal amplification [48] thereby contributing to the promotion of HR [49]. PRMT5, an arginine methyltransferase with described roles in splicing and stemness in GBM [50] is also a key regulator of HR-mediated DSB repair [51]. Notably, such associations between PIMREG and DDR-related proteins were only observed in TMZ-regulated interactomes, suggesting that these interactions or the modulation of binding of these proteins to PIMREG are specific to TMZ exposure, and thus DNA damage, further evidencing roles for PIMREG in DSB repair by HR.

In both TMZ-regulated PIMREG interactome and PIMREG interactome under control conditions, we identified putative interactions between PIMREG and proteins

related to RNA splicing and processing. To the best of our knowledge, this is the first report

linking PIMREG to RNA metabolism. Such observations provide a strong foundation for

future investigation into PIMREG's role in RNA metabolism in GBM. Moreover, the

difference in the splicing network observed between PIMREG interactome under control

condition or induced by TMZ treatment evidences a dynamic role in RNA processing that

may be influenced in response to the genotoxic stress. It is tempting to speculate that this

shift in the splicing network relates to a non-canonical function of PIMREG in DDR-

specific splicing. Mounting evidence supports crosstalk between RNA splicing and the

DDR [52–57], for example, the splicing factor SF3B2, a putative interactor of PIMREG

identified in this study, has been previously linked to HR-mediated repair of DSB [58]. In

support, we have recently reported UHMK1, a known PIMREG interacting protein [42], as

a splicing regulatory kinase [59]. In addition to phosphorylating PIMREG and several

mRNA related proteins, UHMK1 also regulates the phosphorylation of critical DDR

proteins, including RAD54L [60], THRAP3, BCLAF1 [61], and TPT1 [62].

Another PIMREG-associated function observed from TMZ-regulated PIMREG

interactome was related to NADH dehydrogenase activity and mitochondrial metabolism,

suggestive of a potential role for PIMREG in mitochondrial regulation under genotoxic

stress. Though metabolic reprogramming is a known mechanism of TMZ resistance in

GBM [63], how PIMREG contributes in this context remains to be investigated.

We observed distinct PIMREG protein interacting profiles between the two GBM

cell lines. Though both cell lines support TMZ-induced PIMREG interactions with DDR-

related proteins, interactions related to RNA splicing and mitochondrial metabolism

appeared exclusive to U87MG cells. Such differences likely reflect the distinct genetic

background of the two cell lines: T98G cells are MGMT-proficient and carry *TP53*

mutation, while U87MG cells are MGMT-deficient and *TP53* wild-type, features that

enable distinct response to TMZ treatment [64].

Surprisingly, several proteins previously reported to interact with PIMREG were

not identified in our dataset, including PICALM [65], UHMK1 [42], PrPC [66], ROD1

[67], NuRD complex [14], STAT3 [17], NFkB/p65 [20], and BRSK2 [25]. A plausible

explanation is that these interactions were originally identified in different cell lines.

In summary, our proteomic findings expand the current understanding of

PIMREG's interaction network in established and novel cellular processes under both

physiological and genotoxic stress contexts of GBM cells.

PIMREG has two amino acid residues, serine at position 137 (S137) and threonine

at position 114 (T114) that could be sites for ATM/ATR phosphorylation [43]. Our data

shows that PIMREG is recognized by the phospho-ATM/ATR substrate antibody,

suggestive of PIMREG as a direct substrate of these DDR kinases. This is further supported

by the identification of S137 and T114 of PIMREG as phosphosites in proteomic studies

enriched for phospho-ATM/ATR substrates and curated into PhosphoSitePlus® database

(http:// www.phosphosite.org). Notably, phosphorylated PIMREG was detected under

DMSO- and TMZ-treated, as well as in untreated conditions. Therefore, whether

ATM/ATR-mediated phosphorylation of PIMREG is specifically induced in response to

DNA damage or occurs as part of a broader mechanism remains to be determined.

PIMREG deficient cells in the absence of genotoxic stress exhibited elevated levels

of γH2AX, suggesting PIMREG importance for genomic stability. Conversely, upon

exposure to TMZ or IR, these cells overall exhibited an impaired γH2AX response.

Quantification of γH2AX fluorescence intensity confirmed reduced γH2AX in both TMZ

and IR treated KO cells. However, under IR treatment, in which three out of four PIMREG

KO cell lines were assessed, reduced levels of γH2AX were observed exclusively in one

PIMREG KO cell line.

Unexpectedly, the PIMREG KO-2 cell line restored PIMREG protein expression

after extended culturing. Remarkably, upon genotoxic stress, PIMREG KO-2 that regained

PIMREG (KO-2-recPIMREG) displayed increased levels of γH2AX, effectively reversing

the phenotype observed in early passages and highlighting the heterogeneity among the KO

cell lines. The mechanism behind PIMREG re-expression in the KO-2 cell line remains

unclear. However, given that the T98G cells employed for genome editing, are highly

polyploid cells [68], it is plausible that one or more *PIMREG* alleles would lie within

chromatin regions that were inaccessible to Cas9 at the time of editing. Over time, selective

pressures during cell culture may have led to chromatin remodeling, enabling reactivation

of a previously silenced, unedited allele and restoration of PIMREG expression.

However, this study lacks a formal rescue experiment in which PIMREG is

reintroduced exogenously into KO cells. Such an experiment would definitively confirm

that the observed phenotype is specifically due to the loss of PIMREG and not potential

off-target effect.

Reduced activation of ATM (Ser1981) in both PIMREG KO cells exposed to IR

and in PIMREG knockdown cells treated with TMZ [35] supports roles for PIMREG in the

ATM/ATR and γH2AX signaling under damage. Given γH2AX represents a key early

event in DSB signaling [45], PIMREG may also act during the early stages of the cell

response to damage. Consistent with this, we demonstrated that KAP1 activation was

significantly impaired in PIMREG KOs cells exposed to IR. KAP1 is a protein involved in

chromatin condensation. Following DSB induced by IR, ATM phosphorylates KAP-1 at

S824 at the site of damage, where it colocalizes with γH2AX foci and rapidly spreads

throughout the chromatin [69]. Phosphorylated KAP1 facilitates DSB repair in both

euchromatin and heterochromatin [44].

Importantly, we demonstrated that loss of PIMREG significantly impacts DSB repair by reducing HR-mediated repair whereas NHEJ repair was only mildly enhanced in two out four KO cell lines. Notably, DNA-PKc activation, a central orchestrator of NHEJ, was unaffected by PIMREG absence after IR exposure, further supporting specific roles for PIMREG during the signaling and/or activation of HR. Finally, PIMREG's influence on HR may be partially explained by its expression pattern throughout the cell cycle. PIMREG levels become elevated during S and G2, sharply declining as cells enter G1 [13]. HR is predominantly active in S and G2 phases, due to the availability of a sister chromatid for repair, thus this temporal expression profile may underlie PIMREG's preferential role in supporting HR over NHEJ.

Overall, our findings support a role for PIMREG in facilitating DSB signaling and repair following damage, however it is important to note that a degree of phenotypic heterogeneity was observed across treatments (TMZ and IR) and amongst KO clones. The origin of this variability may be a result of off-target effects or the clonal differences arising during the monoclonal selection of the KOs.

We observed a distinct difference in the activation of the DNA signaling molecules in PIMREG KO cells treated with TMZ or exposed to IR. These differences are likely attributable to the nature and timing of DNA damage induced by each agent. IR rapidly induces DSBs whereas TMZ primarily causes base alkylation. The formation of secondary DSBs after TMZ treatment depends on one or more rounds of DNA replication with futile MMR repair [9]. It is conceivable that our cells, treated with TMZ for 24h, harbor significant lower levels of DSBs at the time point analyzed, perhaps explaining why western blot analysis failed to detect substantial changes in DSB signaling in TMZ-treated PIMREG KO cells. In contrast, by immunofluorescence analysis, we detected a reduction in γH2AX intensity in TMZ-treated KO cells. Notably, γH2AX fluorescence intensity was

approximately 100-fold higher in IR-treated KO-1 cells compared to those treated with

TMZ, highlighting the stark difference in DSB burden between the two treatments. These

findings strongly implicate PIMREG as a DDR factor, specifically operating to support

HR-directed repair of DSBs in GBM cells. A key question emerging from this study is

whether PIMREG participates in these pathways directly or exerts its effects indirectly

through broader cellular processes.

Given such broad functionality for PIMREG, our data is more suggestive of

PIMREG as an indirect contributor to the DDR, either through the regulation of upstream

or parallel processes essential for an effective response to genotoxic stress. PIMREG's

multifaceted roles are therefore especially important to understand in the context of cancer.

Especially given the consistent overexpression of PIMREG in many, if not all, types of

cancer, suggesting dysregulation of PIMREG activity may promote tumor progression and

treatment resistance, not only in GBM but in other cancer types.

In conclusion, our findings provide new evidence that PIMREG plays a role in DSB

signaling and HR repair highlighting PIMREG as a potential contributor to therapy

resistance and a promising target for future cancer therapies.

Credit authorship contribution statement

Maria Vitória de Rizzo Gasparini: Investigation, Conceptualization, Visualization, Formal

analysis, Writing - Original Draft. **Laís do Carmo:** Investigation, Visualization, Formal

analysis, Writing - Original Draft. **Giovanna Gonçalves de Almeida**: Investigation,

Visualization, Formal analysis, Writing - Original Draft. **Mariana Siqueira Lacerda**

Mamede: Investigation. **Isabela Spido Dias**: Investigation. **Jennifer Ann Black:**

Supervision, Writing - Review & Editing. **Vanessa C Arfelli:** Validation, Supervision,

Writing - Review & Editing. **Yun-Chien Chang:** Investigation, Formal analysis, Writing -

Review & Editing. **Stephanie Heinzlmeir:** Supervision, Writing - Review & Editing.

Bernhard Kuster: Resources, Writing - Review & Editing. **Philipp A. Greif:** Supervision,

Resources, Writing - Review & Editing. **Leticia Fröhlich Archangelo:** Conceptualization,

Supervision, Writing - Original Draft, Project administration, Funding acquisition.

Declaration of competing interests

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Data availability

The raw mass spectrometry data, MaxQuant search results, and the Swiss-Prot reference

database used for analysis have been deposited to the ProteomeXchange Consortium via

the MassIVE partner repository with the dataset identifier PXD069394 (MassIVE ID:

MSV000099465).

Supporting information

This article contains supporting information.

Acknowledgements

We thank Dr. Jeremy A. Squire for the English review and Patrícia Vianna Bonini, Camila

Menezes Bonaldo (FACS) and Leandro Federiche Borges (IR) for technical assistance.

Marlon Fontes Rocha and Dr. Larissa Dias da Cunha for helping with the FACS analysis.

PAG acknowledges support by the Wilhelm Sander-Stiftung (Förderantrag Nr. 2014.162.3).

This research was supported by the Fundação de Amparo à Pesquisa de São Paulo (FAPESP;

2019/26035-1 and 2022/08648-9 to LFA). FAPESP supports the LMMC (2004/08868-0).

References

[1]   Q.T. Ostrom, M. Price, C. Neff, G. Cioffi, K.A. Waite, C. Kruchko, J.S. Barnholtz-Sloan, CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2016-2020, Neuro Oncol, 25 (2023) IV1–IV99.

[2]   D.N. Louis, A. Perry, P. Wesseling, D.J. Brat, I.A. Cree, D. Figarella-Branger, C. Hawkins, H.K. Ng, S.M. Pfister, G. Reifenberger, R. Soffietti, A. Von Deimling, D.W. Ellison, The 2021 WHO classification of tumors of the central nervous system: A summary, Neuro Oncol, 23 (2021) 1231–1251.

[3]   Q.T. Ostrom, M.L. Shoaf, G. Cioffi, K. Waite, C. Kruchko, P.Y. Wen, D.J. Brat, J.S. Barnholtz-Sloan, J.B. Iorgulescu, National-level overall survival patterns for molecularly-defined diffuse glioma types in the United States, Neuro Oncol, 25 (2023) 799–807.

[4]   R. Stupp, M.E. Hegi, W.P. Mason, M.J. van den Bent, M.J. Taphoorn, R.C. Janzer, S.K. Ludwin, A. Allgeier, B. Fisher, K. Belanger, P. Hau, A.A. Brandes, J. Gijtenbeek, C. Marosi, C.J. Vecht, K. Mokhtari, P. Wesseling, S. Villa, E. Eisenhauer, T. Gorlia, et al., Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial, Lancet Oncol, 10 (2009) 459–466.

[5]   R. Stupp, W.P. Mason, M.J. van den Bent, M. Weller, B. Fisher, M.J.B. Taphoorn, K. Belanger, A.A. Brandes, C. Marosi, U. Bogdahn, J. Curschmann, R.C. Janzer, S.K. Ludwin, T. Gorlia, A. Allgeier, D. Lacombe, J.G. Cairncross, E. Eisenhauer, R.O. Mirimanoff, Radiotherapy plus Concomitant and Adjuvant Temozolomide for Glioblastoma, New England Journal of Medicine, 352 (2005) 987–996.

[6]   J.A. Nickoloff, N. Sharma, C.P. Allen, L. Taylor, S.J. Allen, A.S. Jaiswal, R. Hromas, Roles of homologous recombination in response to ionizing radiation-induced DNA damage, Int J Radiat Biol, 99 (2023) 903–914.

[7]   H. Erasimus, M. Gobin, S. Niclou, E. Van Dyck, DNA repair mechanisms and their clinical impact in glioblastoma, Mutat Res Rev Mutat Res, 769 (2016) 19–35.

[8]   M.E. Hegi, A.-C. Diserens, T. Gorlia, M.-F. Hamou, N. De Tribolet, M. Weller, J.M. Kros, J.A. Hainfellner, W. Mason, L. Mariani, E.C. Bromberg, P. Hau, R.O. Mirimanoff, J.G. Cairncross, R.C. Janzer, R. Stupp, MGMT Gene Silencing and Benefit from Temozolomide in Glioblastoma, 2005.

[9]   W.P. Roos, B. Kaina, DNA damage-induced cell death: From specific DNA lesions to the DNA damage response and apoptosis, Cancer Lett, 332 (2013) 237–248.

[10]   W.D. Wright, S.S. Shah, W.D. Heyer, Homologous recombination and the repair of DNA double-strand breaks, Journal of Biological Chemistry, 293 (2018) 10524–10535.

[11]   H.H.Y. Chang, N.R. Pannunzio, N. Adachi, M.R. Lieber, Non-homologous DNA end joining and alternative pathways to double-strand break repair, Nat Rev Mol Cell Biol, 18 (2017) 495–506.

[12]   D.J. Laverty, S.K. Gupta, G.A. Bradshaw, A.S. Hunter, B.L. Carlson, N.M. Calmo, J. Chen, S. Tian, J.N. Sarkaria, Z.D. Nagel, ATM inhibition exploits checkpoint defects and ATM-dependent double strand break repair in TP53-mutant glioblastoma, Nat Commun, 15 (2024).

[13]   L.F. Archangelo, P.A. Greif, M. Hölzel, T. Harasim, E. Kremmer, G.K.H. Przemeck, D. Eick, A.J. Deshpande, C. Buske, M.H. de Angelis, S.T.O. Saad, S.K. Bohlander,

The CALM and CALM/AF10 interactor CATS is a marker for proliferation, Mol
Oncol, 2 (2008) 356–67.
[14] W.M. Zhao, J.A. Coppinger, A. Seki, X.L. Cheng, J.R. Yates, G. Fang, RCS1, a
substrate of APC/C, controls the metaphase to anaphase transition, Proc Natl Acad
Sci U S A, 105 (2008) 13415–13420.
[15] I. Barbutti, J.M. Xavier-Ferrucio, J.A. Machado-Neto, L. Ricon, F. Traina, S.K.
Bohlander, S.T. Olalla Saad, L.F. Archangelo, CATS (FAM64A) abnormal
expression reduces clonogenicity of hematopoietic cells, Oncotarget, 7 (2016)
68385–68396.
[16] K. Hashimoto, A. Kodama, M. Ohira, M. Kimoto, R. Nakagawa, Y. Usui, Y. Ujihara,
A. Hanashima, S. Mohri, Postnatal expression of cell cycle promoter Fam64a causes
heart dysfunction by inhibiting cardiomyocyte differentiation through repression of
Klf15, IScience, 25 (2022).
[17] Z.S. Xu, H.X. Zhang, W.W. Li, Y. Ran, T.T. Liu, M.G. Xiong, Q.L. Li, S.Y. Wang,
M. Wu, H.B. Shu, H. Xia, Y.Y. Wang, FAM64A positively regulates STAT3 activity
to promote Th17 differentiation and colitisassociated carcinogenesis, Proc Natl Acad
Sci U S A, 116 (2019) 10447–10452.
[18] K. Hashimoto, A. Kodama, T. Honda, A. Hanashima, Y. Ujihara, T. Murayama, S.I.
Nishimatsu, S. Mohri, Fam64a is a novel cell cycle promoter of hypoxic fetal
cardiomyocytes in mice, Sci Rep, 7 (2017) 1–17.
[19] P. Locatelli, M.N. Belaich, A.E. López, F.D. Olea, X. Martín Uranga Vega, C.S.
Giménez, J. Alejandro Simonin, M. del Rosario Bauzá, M.G. Castillo, L.A.
Cuniberti, A. Crottogini, X.S. Carolina Cerrudo, X.D. Pablo Ghiringhelli, U.M.
Vega, Novel insights into cardiac regeneration based on differential fetal and adult
ovine heart transcriptomic analysis, Am J Physiol Heart Circ Physiol, 318 (2020)
994–1007.
[20] L. Jiang, L. Ren, X. Zhang, H. Chen, X. Chen, C. Lin, L. Wang, N. Hou, J. Pan, Z.
Zhou, H. Huang, D. Huang, J. Yang, Y. Liang, J. Li, Overexpression of PIMREG
promotes breast cancer aggressiveness via constitutive activation of NF-κB
signaling, EBioMedicine, 43 (2019) 188–200.
[21] D. Wang, A. Hu, H. Peng, D. Li, L. Zhang, Tumor-promoting function of PIMREG
in glioma by activating the β-catenin pathway, 3 Biotech, 11 (2021).
[22] J. Zhang, L. Qian, J. Wu, D. Lu, H. Yuan, W. Li, X. Ying, S. Hu, Up-regulation of
FAM64A promotes epithelial-to-mesenchymal transition and enhances stemness
features in breast cancer cells, Biochem Biophys Res Commun, 513 (2019) 472–478.
[23] Z. Yao, X. Zheng, S. Lu, Z. He, Y. Miao, H. Huang, X. Chu, C. Cai, F. Zou,
Knockdown of FAM64A suppresses proliferation and migration of breast cancer
cells, Breast Cancer, 26 (2019) 835–845.
[24] Y. Zhou, L. Ou, J. Xu, H. Yuan, J. Luo, B. Shi, X. Li, S. Yang, Y. Wang, FAM64A
is an androgen receptor-regulated feedback tumor promoter in prostate cancer, Cell
Death Dis, 12 (2021).
[25] K. Li, H. Lin, A. Liu, C. Qiu, Z. Rao, Z. Wang, S. Chen, X. She, S. Zhu, P. Li, L.
Liu, Q. Wu, G. Wang, F. Xu, S. Li, SOD1-high fibroblasts derived exosomal miR-
3960 promotes cisplatin resistance in triple-negative breast cancer by suppressing
BRSK2-mediated phosphorylation of PIMREG, Cancer Lett, 590 (2024).
[26] S. Hu, H. Yuan, Z. Li, J. Zhang, J. Wu, Y. Chen, Q. Shi, W. Ren, N. Shao, X. Ying,
Transcriptional response profiles of paired tumor-normal samples offer novel
perspectives in pan-cancer analysis, Oncotarget, 8 (2017) 41334–41347.

[27] H. Zhu, X. Hu, Y. Ye, Z. Jian, Y. Zhong, L. Gu, X. Xiong, Pan-Cancer Analysis of PIMREG as a Biomarker for the Prognostic and Immunological Role, Front Genet, 12 (2021).

[28] C. Zhang, Y. Han, H. Huang, L. Min, L. Qu, C. Shou, Integrated analysis of expression profiling data identifies three genes in correlation with poor prognosis of triple-negative breast cancer, Int J Oncol, 45 (2014) 2025–2033.

[29] Y. Jiao, Z. Fu, Y. Li, W. Zhang, Y. Liu, Aberrant FAM64A mRNA expression is an independent predictor of poor survival in pancreatic cancer, PLoS One, 14 (2019).

[30] K. Mizuno, K. Tanigawa, N. Nohata, S. Misono, R. Okada, S. Asai, S. Moriya, T. Suetsugu, H. Inoue, N. Seki, Fam64a: A novel oncogenic target of lung adenocarcinoma regulated by both strands of mir99a (mir-99a-5p and mir-99a-3p), Cells, 9 (2020) 1–22.

[31] X. Zhao, H. Chen, Y. Qiu, L. Cui, FAM64A promotes HNSCC tumorigenesis by mediating transcriptional autoregulation of FOXM1, Int J Oral Sci, 14 (2022).

[32] H. Yao, F. Lyu, J. Ma, F. Sun, G. Tang, J. Wu, Z. Zhou, PIMREG is a prognostic biomarker involved in immune microenvironment of clear cell renal cell carcinoma and associated with the transition from G1 phase to S phase, Front Oncol, 13 (2023).

[33] W. jing Yun, L. Zhang, N. Yang, Z. guo Cui, H. mao Jiang, M. wen Ha, D. yong Yu, M. zhen Zhao, H. chuan Zheng, FAM64A aggravates proliferation, invasion, lipid droplet formation, and chemoresistance in gastric cancer: A biomarker for aggressiveness and a gene therapy target, Drug Dev Res, 84 (2023) 1537–1552.

[34] X. Wei, Y. Jiang, T. Xia, J. Du, Oncogenic role and prognostic significance of PIMREG in melanoma, Transl Cancer Res, 14 (2025) 1070–1084.

[35] R.B. Serafim, C. Cardoso, V.C. Arfelli, V. Valente, L.F. Archangelo, PIMREG expression level predicts glioblastoma patient survival and affects temozolomide resistance and DNA damage response, Biochim Biophys Acta Mol Basis Dis, 1868 (2022).

[36] Cerami et al., The cBio Cancer Genomics Portal: An Open Platform for Exploring Multidimensional Cancer Genomics Data, Cancer Discov., 2 (2012) 401–404.

[37] A. Subramanian, P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, A. Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, J.P. Mesirov, Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles, 2005.

[38] F.A. Ran, P.D. Hsu, J. Wright, V. Agarwala, D.A. Scott, F. Zhang, Genome engineering using the CRISPR-Cas9 system, Nat Protoc, 8 (2013) 2281–2308.

[39] A.J. Pierce, M. Jasin, in Mouse Embryonic Stem Cells, Molecular Toxicology Protocols. Methods in Molecular Biology, 291 (2005).

[40] N. Bennardo, A. Cheng, N. Huang, J.M. Stark, Alternative-NHEJ is a mechanistically distinct pathway of mammalian chromosome break repair, PLoS Genet, 4 (2008).

[41] W. Wick, M. Weller, M. Van Den Bent, M. Sanson, M. Weiler, A. Von Deimling, C. Plass, M. Hegi, M. Platten, G. Reifenberger, MGMT testing - The challenges for biomarker-based glioma treatment, Nat Rev Neurol, 10 (2014) 372–385.

[42] L.F. Archangelo, P.A. Greif, A. Maucuer, V. Manceau, N. Koneru, C.L. Bigarella, F. Niemann, M.T. dos Santos, J. Kobarg, S.K. Bohlander, S.T.O. Saad, The CATS (FAM64A) protein is a substrate of the Kinase Interacting Stathmin (KIS), Biochim Biophys Acta Mol Cell Res, 1833 (2013) 1269–1279.

[43] S.-T. Kim, D.-S. Lim, C.E. Canman, M.B. Kastan, Substrate Specificities and Identification of Putative Substrates of ATM Kinase Family Members*, 1999.

[44] D. White, I.U. Rafalska-Metcalf, A. V. Ivanov, A. Corsinotti, H. Peng, S.C. Lee, D. Trono, S.M. Janicki, F.J. Rauscher, The ATM substrate KAP1 controls DNA repair in heterochromatin: Regulation by HP1 proteins and serine 473/824 phosphorylation, Molecular Cancer Research, 10 (2012) 401–414.

[45] A.N. Blackford, S.P. Jackson, ATM, ATR, and DNA-PK: The Trinity at the Heart of the DNA Damage Response, Mol Cell, 66 (2017) 801–817.

[46] W.M. Bonner, C.E. Redon, J.S. Dickey, A.J. Nakamura, O.A. Sedelnikova, S. Solier, Y. Pommier, γH2AX and cancer, Nat Rev Cancer, 8 (2008) 957–967.

[47] S. Wu, Y. Shi, P. Mulligan, F. Gay, J. Landry, H. Liu, J. Lu, H.H. Qi, W. Wang, J.A. Nickoloff, C. Wu, Y. Shi, A YY1-INO80 complex regulates genomic stability through homologous recombination-based repair, Nat Struct Mol Biol, 14 (2007) 1165–1172.

[48] H. Jeong, M. Wie, I.J. Baek, G. Sohn, S.H. Um, S.G. Lee, Y. Seo, J. Ra, E.A. Lee, S. Kim, B.G. Kim, R.A. Deshpande, T.T. Paull, J.S. Han, T. Kwon, K. Myung, TRIP13 Participates in Immediate-Early Sensing of DNA Strand Breaks and ATM Signaling Amplification through MRE11, Cells, 11 (2022).

[49] C.S. Clairmont, P. Sarangi, K. Ponnienselvan, L.D. Galli, I. Csete, L. Moreau, G. Adelmant, D. Chowdhury, J.A. Marto, A.D. D'Andrea, TRIP13 regulates DNA repair pathway choice through REV7 conformational change, Nat Cell Biol, 22 (2020) 87–96.

[50] P. Sachamitr, J.C. Ho, F.E. Ciamponi, W. Ba-Alawi, F.J. Coutinho, P. Guilhamon, M.M. Kushida, F.M.G. Cavalli, L. Lee, N. Rastegar, V. Vu, M. Sánchez-Osuna, J. Coulombe-Huntington, E. Kanshin, H. Whetstone, M. Durand, P. Thibault, K. Hart, M. Mangos, J. Veyhl, et al., PRMT5 inhibition disrupts splicing and stemness in glioblastoma, Nat Commun, 12 (2021) 1–17.

[51] T.L. Clarke, M.P. Sanchez-Bailon, K. Chiang, J.J. Reynolds, J. Herrero-Ruiz, T.M. Bandeiras, P.M. Matias, S.L. Maslen, J.M. Skehel, G.S. Stewart, C.C. Davies, PRMT5-Dependent Methylation of the TIP60 Coactivator RUVBL1 Is a Key Regulator of Homologous Recombination, Mol Cell, 65 (2017) 900-916.e7.

[52] L. Shkreta, B. Chabot, The RNA splicing response to DNA damage, Biomolecules, 5 (2015) 2935–2977.

[53] S.C. Lenzken, A. Loffreda, S.M.L. Barabino, RNA splicing: A new player in the DNA damage response, Int J Cell Biol, (2013).

[54] A. Montecucco, G. Biamonti, Pre-mRNA processing factors meet the DNA damage response, Front Genet, 4 (2013).

[55] M. Tresini, D.O. Warmerdam, P. Kolovos, L. Snijder, M.G. Vrouwe, J.A.A. Demmers, W.F.J. Van Ijcken, F.G. Grosveld, R.H. Medema, J.H.J. Hoeijmakers, L.H.F. Mullenders, W. Vermeulen, J.A. Marteijn, The core spliceosome as target and effector of non-canonical ATM signalling, Nature, 523 (2015) 53–58.

[56] C. Naro, P. Bielli, V. Pagliarini, C. Sette, The interplay between DNA damage response and RNA processing: The unexpected role of splicing factors as gatekeepers of genome stability, Front Genet, 6 (2015).

[57] A.S. Bader, B.R. Hawley, A. Wilczynska, M. Bushell, The roles of RNA in DNA double-strand break repair, Br J Cancer, 122 (2020) 613–623.

[58] R. Prados-Carvajal, A. López-Saavedra, C. Cepeda-García, S. Jimeno, P. Huertas, Multiple roles of the splicing complex SF3B in DNA end resection and homologous recombination, DNA Repair (Amst), 66–67 (2018) 11–23.

[59] V.C. Arfelli, Y.-C. Chang, J.W. Bagnoli, P. Kerbs, F.E. Ciamponi, L.M. da S. Paz, S. Pankivskyi, J. de Matha Salone, A. Maucuer, K.B. Massirer, W. Enard, B. Kuster,

P.A. Greif, L.F. Archangelo, UHMK1 is a novel splicing regulatory kinase, Journal
of Biological Chemistry, 299 (2023) 103041.

[60] J.M. Mason, K. Dusad, W.D. Wright, J. Grubb, B. Budke, W.D. Heyer, P.P. Connell,
R.R. Weichselbaum, D.K. Bishop, RAD54 family translocases counter genotoxic
effects of RAD51 in human tumor cells, Nucleic Acids Res, 43 (2015) 3180–3196.

[61] J. Vohhodina, E.M. Barros, A.L. Savage, F.G. Liberante, L. Manti, P. Bankhead, N.
Cosgrove, A.F. Madden, D.P. Harkin, K.I. Savage, The RNA processing factors
THRAP3 and BCLAF1 promote the DNA damage response through selective
mRNA splicing and nuclear export, Nucleic Acids Res, 45 (2017) 12816–12833.

[62] J. Zhang, S.M. De Toledo, B.N. Pandey, G. Guo, D. Pain, H. Li, E.I. Azzam, Role
of the translationally controlled tumor protein in DNA damage sensing and repair,
Proc Natl Acad Sci U S A, 109 (2012).

[63] H.Y. Li, Y.H. Feng, C.L. Lin, T.I. Hsu, Mitochondrial Mechanisms in
Temozolomide Resistance: Unraveling the Complex Interplay and Therapeutic
Strategies in Glioblastoma, Mitochondrion, 75 (2024).

[64] M. Hermisson, A. Klumpp, W. Wick, J. Wischhusen, G. Nagel, W. Roos, B. Kaina,
M. Weller, O6-methylguanine DNA methyltransferase and p53 status predict
temozolomide sensitivity in human malignant glioma cells, J Neurochem, 96 (2006)
766–776.

[65] L.F. Archangelo, J. Gläsner, A. Krause, S.K. Bohlander, The novel CALM interactor
CATS influences the subcellular localization of the leukemogenic fusion protein
CALM/AF10, Oncogene, 25 (2006) 4099–4109.

[66] J. Satoh, S. Obayashi, T. Misawa, K. Sumiyoshi, K. Oosumi, H. Tabunoki, Protein
microarray analysis identifies human cellular prion protein interactors, Neuropathol
Appl Neurobiol, 35 (2009) 16–35.

[67] T.F. Brazão, J. Demmers, W. Van Ijcken, J. Strouboulis, M. Fornerod, L. Romão,
F.G. Grosveld, A new function of ROD1 in nonsense-mediated mRNA decay, FEBS
Lett, 586 (2012) 1101–1110.

[68] G.H. Stein, T98G: An anchorage-independent human tumor cell line that exhibits
stationary phase G1 arrest in vitro, J Cell Physiol, 99 (1979) 43–54.

[69] Y. Ziv, D. Bielopolski, Y. Galanty, C. Lukas, Y. Taya, D.C. Schultz, J. Lukas, S.
Bekker-Jensen, J. Bartek, Y. Shiloh, Chromatin relaxation in response to DNA
double-strand breaks is modulated by a novel ATM-and KAP-1 dependent pathway,
Nat Cell Biol, 8 (2006) 870–876.

[70] T. Aparicio, R. Baer, J. Gautier, DNA double-strand break repair pathway choice
and cancer, DNA Repair (Amst), 19 (2014) 169–175.

Fig. 1: GBM samples with *PIMREG* high expression are enriched for DDR and RNA metabolism. (**A**) GSEA analysis reveals enriched gene sets related to DDR and (**B**) RNA metabolism. (**C**) Correlation plot between the transcript levels of *PIMREG* and the DDR coding genes, *H2AX*, *RAD51*, *RAD51AP1* and *FANCD2* in GBM patient samples. (ρ) Spearman's correlation coefficient and (r) Pearson's correlation coefficient. (**D**) *PIMREG* expression in glioma patients classified according to methylation status of the *MGMT* promoter region. (****) *p*-value < 0.0001, Mann-Whitney two-tailed test. RSEM: RNAseq by Expectation-Maximization.

Fig. 2: PIMREG interactome of GBM cells treated with TMZ and vehicle (DMSO). (A-D) Volcano plot graphs of PIMREG interacting proteins in U87MG (A, B) and T98G (C, D) cells. **(B, D)** TMZ-regulated PIMREG interactors (TMZ vs. DMSO). Upregulated proteins (red dots) interact with PIMREG only under TMZ treatment or show increased association under treatment. Downregulated proteins (blue dots) exhibit reduced or lost binding upon TMZ treatment. **(A, C)** PIMREG interacting proteins under vehicle (DMSO) conditions (PIMREG IP vs. IgG). Proteins with adjusted p-value < 0.05 and $\log_2$ fold change > 1 (PIMREG IP vs. IgG) or $\log_2$ fold change < -1 and >1 (TMZ vs. DMSO) were considered significant (highlighted as blue or red dots). **(E)** Interaction network of proteins identified as PIMREG interactors in U87MG cells (PIMREG IP vs. IgG). **(F)** Interaction network of the proteins identified as TMZ-regulated PIMREG interactors

in U87MG and (**G**) in T98G cells. Each node represents one protein. The proteins interacting with each other are represented in clusters. Node colors indicate proteins that have the same function within the clusters. Lines connecting the nodes represent known interactions (cyan: from curated databases; magenta: experimentally determined), predicted interactions (green: gene neighborhood; red: gene fusions; blue: gene co-occurrence) and other types of interactions (black: co-expression; yellow: text mining; light blue: protein homology). (**H, I, J**) Biological processes (BP) in each group of PIMREG interacting proteins. Number of ontological terms grouped as a BP (graph bars) is depicted in the x axis. Number of genes involved in each BP is shown next to each bar. The bars in gray represent previously described functions of PIMREG. The hatched bars represent the putative new functions of PIMREG.

Fig. 3: PIMREG is a substrate of ATM/ATR. Lysates of T98G cells untreated (UT) or treated with TMZ and vehicle (DMSO) for 24 h were precipitated with (**A**) anti-PIMREG antibody and isotype control (IgG). Western blot analysis with anti-PIMREG antibody shows the precipitated protein in the upper panel. Precipitated PIMREG is indicated with the arrowhead (lanes 3, 6 and 9). Immunoblotting of the precipitated protein with anti-phospho-ATM/ATR substrate antibody is shown in the lower panel. Asterisks (*) indicate the precipitated PIMREG protein detected with the anti-phospho-ATM/ATR substrate antibody. (**B**) ATM/ATR substrates immunoprecipitated using anti-phospho-ATM/ATR substrate antibody were immunoblotted with the same antibody (upper panel) and anti-PIMREG antibody (lower panel). Asterisk (*) indicates the precipitated phospho-ATM/ATR substrates detected with anti-PIMREG antibody (lane 9). As input control, lanes 1, 4 and 7 were loaded with 20 % of the protein extract used for each precipitation reaction. UT: untreated.

Fig. 4: γH2AX accumulation in PIMREG KO cells. (**A**) Western blot analysis of protein extracts from T98G WT and PIMREG KO (KO-1, KO-2, KO-3 and KO-4) cells. Membranes were blotted with anti-PIMREG, anti-phospho-H2AX (S139) antibodies. Actin was used as a loading control. (**B**) Bar graph showing densitometric quantification of γH2AX bands, normalized to their corresponding endogenous actin levels (γH2AX/actin). Results are shown as mean ± SD (standard deviation) of three independent experiments. Red dashed line indicates γH2AX levels in WT cells for comparison. (**C**) Western blot analysis of protein extracts derived from T98G WT and PIMREG KOs treated with TMZ or vehicle (DMSO) for 24 h. (**D**) Bar graph showing γH2AX accumulation expressed as the ratio of γH2AX levels (γH2AX/actin) in TMZ-treated cells relative to their corresponding DMSO control (TMZ/DMSO). Results are shown as mean ± SD (standard deviation) of five independent experiments. Red dashed line indicates the γH2AX accumulation in WT cells upon TMZ treatment. (**E**) Western blot analysis of protein extracts from T98G WT and PIMREG KO cells following IR exposure. Samples were collected at 0.5 and 1h post-irradiation. Tubulin was used as a loading control. (**F**) Bar graph showing γH2AX accumulation expressed as the ratio of γH2AX levels (γH2AX/tubulin) in IR-treated cells relative to their corresponding non-irradiated controls (IR/NI). Results are shown as mean ± SD (standard deviation) of four independent experiments. Red dashed line indicates γH2AX accumulation in WT cells following IR exposure. Representative western blots are shown. (*) p-value < 0.05, One-way ANOVA test.

Fig. 5.: PIMREG deficiency impairs nuclear accumulation of γH2AX in GBM cells treated with genotoxic agents. Representative immunofluorescence analysis of T98G WT and PIMREG KO cells treated with (**A**) TMZ or vehicle (DMSO) for 24 h and (**C**) exposed to IR (4 Gy) and collected at 0.5 h post-irradiation. PIMREG is shown in green (Alexa Fluor 488), γH2AX in red (Alexa Fluor 555) and nuclei (DAPI) in cyan. Scale bar: 10 µM. (**B, D**) Fluorescence intensity quantification of nuclear γH2AX, quantified in at least 75 cells per condition. Data are from 3 to 5 independent experiments. The horizontal bars indicate the median intensity. (******) p-value < 0.01, (*******) p-value < 0.001 and (********) p-value < 0.0001, One-way ANOVA test followed by the post-test Kruskal-Wallis was applied. WT: wild-type; KO: knockout; KO-2-recPIMREG: KO-2 cells that recovered PIMREG's expression; NI: non-irradiated. (**E**) Western blot analysis of PIMREG in the same samples in (C and D): T98G WT, PIMREG KOs (KO-1 and KO-3) and KO-2-recPIMREG. Membranes were blotted with anti-PIMREG and actin was used as a loading control.

Fig. 6: PIMREG KO affects activation of DNA damage response signaling molecule following IR exposure. (**A, C, E**) Western blot analysis of protein extracts derived from T98G WT and PIMREG KOs treated with TMZ or vehicle (DMSO) for 24 h. Membranes were blotted with anti-Phospho-KAP1 (Ser824) and KAP1 (A), anti-Phospho CHK1 (Ser345) and CHK1 (C) and anti-Phospho-DNA-PKc (Ser2056) and DNA-PKc (E) antibodies. Tubulin was used as a loading control. (**B, D, F**) Bar graph showing densitometric quantification of phosphorylated protein band intensities normalized to total protein levels. Protein activation is expressed as the ratio of the phosphorylation levels in TMZ-treated cells relative to their corresponding DMSO control (TMZ/DMSO). Results are shown as mean ± SD (standard deviation) of three independent experiment. Red dashed line indicates the protein activation in WT cells upon TMZ treatment. (**G, I, K, M**) Western blot analysis of protein extracts from T98G WT and PIMREG KO cells (KO-1, KO-2, KO-3, and KO-4) following exposure to IR (4Gy). Samples were collected at 0.5 and 1h post-irradiation. Membranes were blotted with anti-Phospho ATM (Ser1981) and ATM (G), anti-Phospho-KAP1 (Ser824) and KAP1 (I), anti-Phospho CHK1 (Ser345) and CHK1 (K), anti-Phospho-DNA-PKc (Ser2056) and DNA-PKc (M) antibodies. Tubulin was used as a loading control. (**H, J, L, N**) Bar graph showing densitometric quantification of phosphorylated protein band intensities normalized to total protein levels. Protein activation is expressed as the ratio of the phosphorylation levels in IR samples relative to their corresponding non-irradiated controls (IR/NI). Results are shown as mean ± SD (standard deviation) of four independent experiment. Red dashed line indicates the protein activation in WT cells at the indicated time points following IR exposure. (*) p-value < 0.05 and (**) p-value < 0.01, One-way ANOVA test. WT: wild-type; KO: knockout; NI: non-irradiated.

Fig. 7: DNA repair reporter assay. The graphs show the percentage of PIMREG WT and PIMREG KO (KO-1, KO-2, KO-3, and KO-4) cells that repaired the DSBs. GFP reports the repair events via (**A**) HR in U2OS DR-GFP and (**B**) NHEJ in HeLa EJ5-GFP cells. The percentage of GFP-positive cells was normalized to the mean percentage of WT control cells within each experiment. Experiments were performed at least three times in triplicates. Fifty thousand events were acquired. Sample sizes for HR repair (A): WT (n = 5), KO-1 (n = 5), KO-2 (n = 3), KO-3 (n = 4), KO-4 (n = 5). Sample sizes for NHEJ repair (B): WT (n = 9), KO-1, KO-2, KO-3 and KO-4 (n = 7). (*) p-value < 0.05, (***) p-value < 0.001 and (****) p-value < 0.0001, Brown-Forsythe and Welch ANOVA.