Iron Deficiency Impairs Mitochondrial Energetics and Early Axonal Growth and
Branching in Developing Hippocampal Neurons

Author list: Daniel C. Mendez[1,2], Karishma Devgun[1], Timothy R. Monko[1], Luke H.
Carlson[1], Daniel J. Mickelson[1], Lorene M. Lanier[3], Michael K. Georgieff[1], & Thomas W.
Bastian[1*]

Author affiliations

1 Department of Pediatrics, School of Medicine, University of Minnesota, Minneapolis,
MN, 55455

2 Graduate Program in Neuroscience, University of Minnesota, Minneapolis, MN, 55455

3 Department of Neuroscience, University of Minnesota, Minneapolis, MN, 55455

30 Corresponding author. *email address:* bastian@umn.edu (Thomas W. Bastian)

Abstract:

Each stage of neuronal development (i.e., proliferation, differentiation, migration, neurite outgrowth and synapse formation) requires functional and highly coordinated metabolic activity to ultimately ensure proper sculpting of complex neural networks. Energy deficits underlie many neurodevelopmental, neuropsychiatric and neurodegenerative diseases implicating mitochondria as a potential therapeutic target. Iron is necessary for neuronal energy output through its direct role in mitochondrial oxidative phosphorylation. Iron deficiency (ID) reduces mitochondrial respiratory and energy capacity in developing hippocampal neurons, causing permanently simplified dendritic arbors and impaired learning and memory. However, the effect of ID on early axonogenesis has not been explored. We used an embryonic mixed-sex primary mouse hippocampal neuron culture model of developmental ID to evaluate mitochondrial respiration and dynamics and effects on axonal morphology. At 7 days in vitro (DIV), ID impaired mitochondrial oxidative phosphorylation capacity and stunted growth of both the primary axon and branches, without affecting branch number. Mitochondrial motility was not altered by ID, suggesting that mitochondrial energy production --- not trafficking --- underlie the axon morphological deficits. These findings provide the first link between iron-dependent neuronal energy production and early axon structural development and emphasize the importance of maintaining sufficient iron during gestation to prevent the negative consequences of ID on brain health across the lifespan.

Significance Statement:

This study used a primary mouse hippocampal neuron culture model of iron deficiency to address an important gap in knowledge of how disruption of iron-regulated mitochondrial activities affects axonal development. After axon initiation but prior to rapid dendrite outgrowth, iron chelation reduced mitochondrial oxidative phosphorylation capacity and stunted the growth of the primary axon and branches but without affecting branch number. Mitochondrial motility was not altered in iron-deficient axons, indicating that reduced neuronal energetic capacity and not impaired axonal mitochondrial trafficking may underlie these morphological deficits. Both iron and mitochondrial dyshomeostasis underlies many neurodevelopmental, neuropsychiatric, and neurodegenerative disorders, which can have origins during the period of fetal-neonatal development when rapid axon growth/branching occurs. This study highlights the importance of advancing knowledge on the effects of mitochondrial deficits in early life as it pertains to optimizing brain health throughout the lifespan.

Introduction:

Iron-deficiency (ID) is the most common nutrient deficiency globally and affects 40-50% of pregnant women and children (Gao et al., 2025; Lozoff & Georgieff, 2006). Fetal and neonatal brain development is a period of high iron and energy requirements (Fretham et al., 2011; Georgieff et al., 2018; Radlowski & Johnson, 2013). ID during this early life period acutely impairs brain development and function and increases the long-term risk for cognitive deficits, schizophrenia, ADHD and depression (Insel et al., 2008; Konofal et al., 2004; Lozoff et al., 2013; Lukowski et al., 2010; Schmidt et al., 2014). The hippocampal region of the brain, involved in learning and memory, is rapidly developing during this iron-sensitive period and is particularly iron and energy demanding. Fetal-neonatal ID impairs hippocampal-mediated learning and memory in children, adolescents, and adults even after early postnatal iron repletion (Fretham et al., 2011; Georgieff et al., 2018; Radlowski & Johnson, 2013). Rodent models of early-life ID have demonstrated reduced myelination of axons and dendritic complexity of hippocampal neurons (Bastian et al., 2016, 2022; Brunette et al., 2010; Carlson et al., 2010; Fretham et al., 2012; Greminger & Mayer-Pröschel, 2015; Jorgenson et al., 2003; Perng et al., 2021), which may underlie the persistent deficits in synaptic function and learning and memory behaviors following recovery from fetal-neonatal ID. In contrast, very little is known about the effects of fetal-neonatal ID on axonal structural development.

Iron is critical for cellular energy production through mitochondrial oxidative phosphorylation by providing heme and/or iron-sulfur clusters for cytochrome c and all four protein complexes of the electron transport chain (ETC), as well as succinate dehydrogenase and aconitase in the tricarboxylic acid (TCA) cycle (Bastian et al., 2020). We have previously shown that early-life ID reduces hippocampal neuron mitochondrial respiration and adenosine-5'-triphosphate (ATP) production throughout the period of rapid dendritic arborization (Bastian et al., 2016, 2019, 2020, 2022; de Deungria et al., 2000; Monko et al., 2024; Rao et al., 2003). Neuronal ID also reduces mitochondrial size, motility, and density in the dendritic compartment of hippocampal neurons (Bastian et al., 2019). Thus, we proposed ID-induced dendrite growth and branching deficits are caused by impaired mitochondrial energetics and quality control (Bastian, 2019; Bastian et al., 2016, 2019, 2020). Despite the body of knowledge on mitochondrial and dendritic structure, no studies have addressed the effect of developmental ID on axonal growth and mitochondrial capacity. This is important because misguided axonal growth can contribute significantly to structural and functional deficits in the hippocampus.

During fetal brain development, neuronal polarization begins with axon differentiation and subsequent rapid growth and branching of axons to reach their permanent targets (Dotti et al., 1988). Neurites destined for dendrite specification undergo rapid growth and branch beginning later in development (late gestation - first 2 years after birth in humans and first 2-3 weeks in rodents) (Semple et al., 2013). Mitochondrial localization and distribution within neuronal sub compartments are critical for neuronal polarization

through regulation of ATP production, calcium buffering, reactive oxygen species (ROS)
signaling, iron homeostasis and apoptosis (Bradke & Dotti, 1997; Bastian, 2019; Lewis
et al., 2013). Inasmuch, the homeostatic balance of mitochondrial dynamics (e.g.,
bidirectional mitochondria trafficking vs. anchoring) is also important for meeting the
metabolic demands of developing axons (Cheng et al., 2022; Cheng & Sheng, 2021;
Sheng, 2014). We hypothesized that ID could impair axon development due to iron's
direct role in mitochondrial respiration and the effect of ID on mitochondrial trafficking in
dendrites (Bastian et al., 2020). The effects of ID on mitochondrial activity during early
neuronal maturation (prior to rapid dendrite outgrowth) remain under-explored.

In the current study, we showed that ID impairs mitochondrial respiration and ATP
production rate during the period of rapid axon growth/branching in our hippocampal
neuronal culture model of early-life ID. Contrary to our prior dendrite studies, no
changes in axon mitochondrial motility were observed in iron-deficient neurons during
early axonogenesis. Morphological analysis of iron-deficient axons showed simplified
complexity, characterized by stunted growth of both the primary axon and branches.
Together, these findings suggest iron-dependent mitochondrial energy production as a
key determinant of proper neuronal structural development.

Materials and Methods:

ANIMALS

Timed pregnant CD1 mice were purchased from Charles River Laboratories. The mice were delivered on embryonic day (E) 13 or 14 and housed in a constant temperature and humidity-controlled facility with food and water administered ad libitum. All animal procedures were performed in facilities accredited by the Association for Assessment and Accreditation of Laboratory Animal Care and in alignment with guidelines and standards established in the *National Institutes of Health's Guide for the Care and Use of Laboratory Animals.* The protocols for animal procedures were also approved by The University of Minnesota Institutional Animal Care and Use Committee prior to commencement.

CELL CULTURE.

The primary hippocampal neuron culture model of developmental ID used in this study has been previously characterized in long-term studies of mitochondrial function and dendrite growth (Bastian et al., 2016, 2019, 2022; Monko et al., 2024). Briefly, the hippocampi of mixed-sex E16 CD1 mouse embryos were pooled, followed by dissociation and plating at in 6-well plates (125,000 cells/well), 96-well plates (5,000 cells/well), or 24-well XFe microwell plates (8,000 cells/well) coated with poly-D-lysine/laminin. Neuronal plating media [Minimal Essential medium (Corning #15-010-CV), 10mM HEPES (Invitrogen 15630-080), 10mM Sodium Pyruvate (Gibco 11360-070), 0.5mM Glutamine (Invitrogen #25030-164), 12.5µm glutamate (Sigma 128430-100g), 10% fetal bovine serum (Sigma #F6178-500ml), and 0.6% glucose (Fisher MT25037Cl)] was exchanged 2-4 hours after plating cells with glia-conditioned neural growth media [Neurobasal (Gibco #12348-017), 1x B27 (Gibco #17504-044), 0.5mM Glutamine (Invitrogen # 25030-164) and 1x penicillin/streptomycin (Invitrogen #CX30324)]. Time of plating is designated as 0 days *in vitro* (DIV).

At 3 DIV, ID was induced with 5µM of the extracellular iron chelator deferoxamine (DFO, Cayman Chemical #14595) for 24-well XFe (Agilent Technologies) and 96-well plate or 9µM DFO for 35mm dish (Bastian et al., 2016, 2019, 2022; Monko et al., 2024). Concomitantly, 5-fluoro-2'-deoxyuridine (5-FU, Sigma #F0503-100MG) was administered to inhibit glia proliferation during DFO-treatment. Iron-sufficient control cultures received an equal volume of vehicle (ddH2O) in fresh unconditioned neuronal growth medium. As shown previously (Bastian et al., 2016, 2019, 2022; Monko et al., 2024), these DFO doses produce a similar degree of neuronal ID at 11 and 18 DIV as observed in the developing brains of rodent ID models (Carlson et al., 2007, 2009) and human iron-deficient newborn brains (Petry et al., 1992). They do not affect neuronal zinc status (Bastian et al 2016). Assays were performed at 7 DIV since by this time in hippocampal neuronal cell cultures the axonal initial segment (AIS) has been well formed (2-4 DIV) and this point marks the beginning of rapid axonal branching and the

initial period of synapse formation, which is critical for proper development of complex
neural networks (Grabrucker et al., 2009; Sahu et al., 2019).

INTRACELLULAR ATP ASSAY.

Intracellular ATP concentrations were quantified following manufacturer's instructions
(Promega Cell Glo Titer 2.0) and as reported previously (Bastian et al., 2019, 2022;
Monko et al., 2024). Prior to the assay, 1μM Hoechst 33342 was added to each well
and live nuclei imaging was performed to normalize ATP concentrations to cell density
as we have described (Bastian et al., 2022). Luminescence was measured with a
BioTek Synergy LX Multimode Plate Reader. The ATP concentrations of each well were
determined based on standard curve readings and then normalized to cell count from
live Hoechst imaging (see normalization described below).

ATP RATE ASSAY FOR BIOENERGETICS.

The Seahorse XF ATP Rate Assay was performed according to the manufacturer's
instructions and as described previously (Bastian et al., 2022), to assess relative ATP
contribution from either oxidative phosphorylation or glycolysis. In brief, XF Assay
Medium (XF DMEM, pH 7.4, with final concentrations of 1mM XF Sodium Pyruvate,
2mM XF Glutamine, and 10mM XF Glucose) was prepared and sterile filtered either the
day before and stored at 4°C or prepared the same day. On the day of assay at 7 DIV,
neuronal medium was replaced with XF Assay Medium containing either 5μm DFO or
vehicle. Working dilutions of compounds, Oligomycin (ATP synthase inhibitor) and
Antimycin A/Rotenone (Complex III and I inhibitors, respectively) were made to achieve
a final concentration of 1.5μM and 0.5μM after injection, respectively, in XF Assay
Medium. To de-gas the plate of CO_2, the cell culture microwell plate was placed
immediately in a 37°C non-CO_2 incubator for 30 minutes (min) prior to the start of the
bioenergetics assay. Oxygen consumption rates (OCR) and extracellular acidification
rates (ECAR) were measured simultaneously and normalized to cell density as done
previously and described below (Bastian et al., 2022).

INTRACELLULAR AND ATP RATE CELL DENSITY NORMALIZATION

To normalize intracellular ATP and Seahorse OCR and ECAR measurements to cell
density, neuron cultures were treated with 1uM Hoescht 33342 (Cayman Chemical,
15547) for live nuclear imaging and placed in humidity and temperature-controlled
incubation (5% CO2 and 37°C) for live cell imaging using Zeiss Celldiscoverer 7 and
Zeiss Axiocam 506 mono camera with the following specifications: 5x Plan-Apochromat
objective (NA=0.35), LED illumination for oblique contrast transmitted light (725nm) and
fluorescence (385nm), and fluorescent quad bandpass filter set 90HE (beamsplitter:
RQFT 405 + 493 + 575 + 653nm, emission: QPB: 425/30 + 514/30 + 592/25 +
709/100nm). Data for intracellular and ATP rate assays were normalized to the number

of cells in the respective well. Images were segmented, classified, measured, and
mapped to experimental data in batch using tools from our Python package, [napari-
ndev] (T. Monko & Bastian, 2025). Images were loaded into Python using
[bioio](https://github.com/bioio-devs/bioio) and [pyclesperanto] (Haase et al., 2023).
GPU-accelerated algorithms were used to pre-process by first reducing noise (median)
and then subtracting background (tophat). Images were segmented with pyclesperanto
by using [Otsu's] (Otsu, 1979), thresholding algorithm on gaussian blurred images
followed by separation of neighboring cells with a seeded watershed, created by a
voronoi diagram of peak spot detection of a gaussian blurred image. To detect live and
dead cells among the segmented nuclei, we trained Random Forest machine learning
object classifiers using napari-ndev and [accelerated pixel and object classifiers] (Haase
et al., 2022). Segmented nuclei were predicted for nuclei classification in batch with
napari-ndev. Finally, segmented images were measured with napari-ndev tools using
either [scikitimage] (Van Der Walt et al., 2014) or [Nyxus]
(https://github.com/PolusAI/nyxus) as the label measurement backend. Image
metadata, including well position in the respective culture plate, was used to map data
to treatment conditions and other experimental metadata using napari-ndev's
PlateMapper. Data was grouped and summarized according to this metadata.
Intracellular ATP data values were relativized to the average control value for each
unique culture experiment and log transformed before statistical analysis.

MITOCHONDRIAL MOTILITY LIVE IMAGING.

Prior to plating, cells were co-nucleofected with an mcherry-Mito-7 (Addgene # 55102)
and "humanized" recombinant green fluorescent protein (hrGFP) plasmid driven by a
CMV enhanced chicken B-actin promoter (Zolotukhin et al., 1996), using a Lonza
Amaxa II and Mouse Neuron Nucleofector Kit (Cat. No. VAPG-100), as described
previously (Bastian et al., 2016; Penrod et al., 2015). Nucleofected cells were mixed 1:1
with non-nucleofected cells and plated in 96-well plates at a seeding density of 5,000
cells/well. At 7 DIV, time-lapse imaging was performed using a ZEISS Celldiscoverer 7
with environmental control set to 37°C and 5% CO_2. mCherry-positive neurons (n = 11-
12 neurons per group) were imaged in 3-channels (oblique, mcherry and hrGFP) every
1 second (s) for a duration of 1.7 min for one independent time-lapse experiment or 3.5
s for 5 min for a second independent time-lapse experiment with a ZEISS Axiocam 506
mono camera with the following specifications: 50x Plan-Apochromat objective (NA=1.2)
with water immersion and 0.5x magnification, LED illumination for oblique contrast
transmitted light (725nm), and fluorescence (590nm & 470nm), and fluorescent quad
bandpass filter set 90HE or triple bandpass filter set 91HE (beamsplitter RTFT 450 +
538 + 610; emission filter TBP 467/24 + 555/25 + 687/145).

The images were converted to TIFF files and then organized according to the
hierarchical organization described in the KymoAnalyzer manual (Neumann et al.,
2017). Axonal segments between 50-100μm were traced anterogradely to the axonal

growth cone. Kymographs for each neuron generated were then traced and processed
through semi-automated Fiji plugins of the KymoAnalyzer plugin package version 1.01
(Neumann et al., 2017; Schindelin et al., 2012). Data for the various mitochondrial
parameters were then organized by category, mean and median compiled for individual
mitochondria for each imaged axon, then analyzed and graphed in PRISM.

IMMUNOCYTOCHEMISTRY OF AXONAL PROCESSES.

Cells nucleofected with hrGFP were mixed 1:1 with non-nucleofected cells and plated in
a 6-well plate (125,000 cells/well). This allowed sparse neuronal labeling to trace
individual neurons. At 7 DIV, cultures were fixed and processed for
immunocytochemistry (ICC) in 6-well plates as described (Bastian 2016, 2017, 2019,
2022). Briefly, cultures were incubated with a mouse monoclonal primary antibody
against MAP2 (Abcam, Catalogue: ab11268) at a 1:500 dilution for 1 hour at room
temperature (RT) or overnight at $4°C$ to specifically label the dendrites. Cells were then
incubated with a secondary donkey anti-mouse Alexa Fluor® (AF) 647 antibody
(Jackson Immunoresearch, Catalogue:715-605-151) diluted at 1:50 for 1 hour at RT.
DAPI stain was used to visualize the nucleus. Individual GFP-positive neurons were
imaged using a ZEISS Celldiscoverer 7 and Zeiss Axiocam 506 mono camera with 20x
Plan-Apochromat objective (NA=0.7) at 0.5x magnification and fluorescent filter sets
described above. Tiled images were stitched using ZEISS Zen Software and exported
as OME-TIFF for morphology analysis.

AXONAL MORPHOLOGY ANALYSIS.

The images were analyzed using Fiji and the Simple Neurite Tracer plugin to determine
the structural effect of iron deficiency on axon length and branching as done previously
for dendrites (Bastian et al., 2016, 2022). The center of the soma was identified and
marked. In addition to being much longer, thinner and more highly branched than
dendrites at this stage, the axon was identified as being GFP-positive and MAP2-
negative. The primary axon was determined as the longest and straightest axonal
process, with secondary branches emanating directly from the primary axon and all
other branches considered tertiary axon branches. Sholl analysis was performed to
determine the number of crossing points as a function of distance from soma (overall
complexity), and the number of branches and length of primary axon and branches
were quantified.

mRNA EXPRESSION ANALYSIS.

TRIzol cell lysates from hippocampal cultures at 7 DIV were pooled from 8 wells of a 96-
well plate or 1 well of a 6-well plate and RNA was extracted with a Monarch® Spin RNA
Cleanup kit (New England Biosciences, catalogue: T2030L). Purity and concentration of
samples was determined using a nanodrop spectrophotometer (catalogue #: ND-1000)
and 100 ng of RNA diluted in nuclease-free water (20µL reaction volume) was reverse

transcribed to synthesize cDNA for quantitative real-time PCR (qPCR) analysis using a
cDNA synthesis kit (Applied Biosciences, Catalogue: AB#4387406). TaqMan probe for
the *transferrin receptor 1* (*Tfr1*) gene (Thermo, Catalogue: 4331882, Mm00441941_m1)
was used as a measure of cellular iron status as we have described (Bastian et al.,
2016, 2019, 2022; Monko et al., 2024). TaqMan probes for *TATA binding protein* (*Tbp*)
(Thermo, Catalogue: 4331182, Mm00446971_m1), *ribosomal protein s18* (*Rps18*)
(Thermo, Catalogue: 4331182, Mm02601777_g1), and *glyceraldehyde-3-phosphate*
dehydrogenase (*Gapdh*) (Thermo, Catalogue: 4331182, Mm99999915_g1) genes were
used as reference genes for normalization. Briefly, qPCR was performed in triplicate on
2.5ng cDNA using the LUNA master mix (NEB #M3004) adapted from the
manufacturer's protocol for a 10μL reaction volume. An Applied Biosystems
QuantStudio 3 qPCR machine (catalogue: A28132) was used with a denaturation step
at 95℃ for 1 min., PCR amplification step at 95℃ for 15 sec. and extension at 60℃ for
30 sec per cycle (40 cycles total). Relative gene expression for all genes of interest was
calculated relative to an internal calibrator cDNA sample. GeNorm was used to
determine the stability of *Tbp*, *Rps18*, and *Gapdh* and their geometric mean was used
for normalization of *Tfr1* by ΔΔCt method. Relative *Tfr1* gene expression data from
DFO-treated cells was normalized to the average of all IS samples for each unique
neuron culture preparation.

STATISTICAL ANALYSIS APPROACH.

Graphpad Prism 10 was used to carry out univariate statistical analyses with α=0.05 to
determine statistically significant differences between groups. Unpaired t-test or Mann-
Whitney was used for parametric or non-parametric data, respectively for statistical
analysis of ATP concentrations, Seahorse respiratory parameters, and mitochondrial
motility. All datasets were tested for normality (Anderson-Darling, D'Agostino-Pearson
omnibus, Shapiro-Wilk, & Kolmogorov-Smirnov) and equal variance (F-test) prior to
subjecting to parametric or non-parametric tests. If data was normally distributed but
had unequal variance, an unpaired t-test with Welch's correction was applied. To adjust
for multiple comparisons, the Benjamini-Hochberg method of FDR correction was used
on resulting p-values from Mann-Whitney tests for all mitochondrial motility parameters.
Statistics for Sholl Analysis were computed using repeated-measures 2-factor ANOVA
to compare group differences with respect to distance from the soma. Multivariate
analysis was coded using Python Jupyter Lab Notebook using exported data from all
Sholl Analysis parameters. Prior to analysis, Levene's test showed equal variance but
non-normal distribution of the Sholl dataset with or without log-transformation. A Box's
M test also showed differences in covariance matrices between IS and ID groups.
Though assumption violations were present due to non-normal distribution and
difference in covariance matrices, a multivariate analysis of variance (MANOVA) was
still performed which included statistical tests by Pillai's trace and Wilk's lambda. Pillai's
trace is a MANOVA statistical approach robust to violations of normality and covariance

particularly for datasets with large and balanced sample sizes, while Wilk's lambda is

more sensitive to these assumption violations and outliers (Appolus EE & Okoli CN,

2022). Thus, both statistical results are reported.

Results:

Neuronal reduction of total intracellular ATP after iron chelation during early axonogenesis.

To evaluate 7 DIV neuronal iron status in our hippocampal neuron culture model - as done previously at 11, 18, and 21 DIV (Bastian et al., 2016, 2022; Monko et al., 2024) - mRNA levels of the *transferrin receptor 1* gene (*Tfr1*), which encodes the surface receptor chiefly responsible for cellular iron uptake and is a sensitive indicator of neuronal iron status, were measured by qPCR (Carlson et al., 2007, 2009). Relative mRNA expression of *Tfr1* significantly increased in ID hippocampal neuronal cultures by 1.83-fold in a 6-well vessel using a 9µM DFO dosage and 1.32-fold increase in the 96-well format using a 5µM DFO dosage (Figure 1B). Total intracellular ATP (p = 0.0276) was significantly decreased by 18% in 5uM DFO-treated neurons compared to IS (Figure 1C), consistent with previous work demonstrating reduced energy status at later stages of neuronal development (Bastian et al., 2019, 2022; Monko et al., 2024). ATP measurements were normalized to cell density which was not significantly different between IS and ID wells (p=0.8938).

Effects of iron chelation on neuronal energetics during axonogenesis.

Mitochondrial respiration by oxidative phosphorylation was impaired in 7 DIV ID neurons across several energetic respiration states compared to IS neurons. Specifically, OCR associated with basal respiration (p = 0.0108; Figure 2B), ATP-linked respiration (p = 0.0246; Figure 2C) and non-mitochondrial respiration (p=0.0122; Figure 2D) were decreased by 27%, 25%, and 47%, respectively. This was associated with trending decreases in mitochondrial ATP production rate (p = 0.0926) and glycolytic ATP production rate (p=0.1075) and a 26% lower Total ATP production rate (p=0.0152) in ID compared to IS neurons (Figure 2E-G). ATP rate index (p=0.6092) was not different between IS and ID neurons (Figure 2H), which is consistent with previous observations (Bastian et al., 2022). All bioenergetic read-out measurements were normalized to cell density, which was not significantly different between groups (p=0.6367).

Iron chelation does not affect mitochondrial motility during axonogenesis.

Since mitochondrial trafficking to sites of high energy demand is required to metabolically support developing neuronal processes and is itself an ATP-consuming process, and because we saw mitochondrial motility differences in dendritic compartment of hippocampal neurons due to ID (Bastian et al., 2019), mitochondrial motility was assessed in the axonal compartment (Millecamps & Julien, 2013). Kymograph analysis on time-lapse images (Movie 1A-B) of iron-sufficient and iron-deficient axons (Figure 3 A-B) showed no significant difference in any of the motility parameters (Table 1) including net velocity, segmental velocity (segments of mitochondria motion excluding pauses), density, total pause frequency in retrograde and anterograde direction and percent time in motion (Figure 4A-D, Table 1). However,

assessment of sub-categorized data showed a 38.2% decrease in segmental velocity in
the retrograde direction in iron-deficient compared to iron-sufficient neurons (p = 0.0314;
Figure 4B). Specifically, mitochondria moving in the anterograde direction towards the
axon terminal paused 57% more in iron-deficient neurons than iron-sufficient neurons (p
= 0.0064) (Figure 4D). Ultimately, when p-values were adjusted for false discovery rate
(FDR) using the Benjamini-Hochberg method, no significant difference was observed for
any of the parameters assessed. Similar trends and no significant differences were
found for axonal mitochondrial motility parameters when assessed in the 9µm DFO
dose/35mm dish culture system (data not shown).

Iron chelation during axonogenesis impairs axonal growth and branching.

To assess if ID affects early axon growth and branching, semi-automated tracing of the
entire axonal arbor from individual neurons was performed (Figure 5A). Sholl analysis of
axonal tracings revealed significantly decreased (p≤0.05) number of axonal crossings
between 440µm-850µm from the soma in iron-deficient compared to iron-sufficient
hippocampal neurons, suggesting impaired axonal growth and/or branching (Figure 5B).
To determine whether ID specifically impaired axonal outgrowth or *de novo* axon
branching, the number and length of axonal processes were quantified. Consistent with
the Sholl analysis, the primary (p=0.0003) and total length (p=0.0169) of the axonal
arbor (primary segment + branches) was reduced in iron-deficient neurons (Figure
5B,C). ID did not alter the number of branches or the total branch length (Figure 6B, C),
however, ID significantly reduced the average branch length (p=0.0012) compared to IS
neurons (Figure 6A), suggesting impaired axon linear growth but not branch
formation/retraction. To further explore this, a MANOVA was performed using statistical
tests by Wilk's lambda and Pillai's trace (Supplemental Table 1). Data are tabulated
(Supplemental Table 1) ranking each modeled multivariate interaction by F-value
(highest to lowest) with its corresponding raw and FDR-adjusted p-value and effect size
(η^2 from Wilk's and Pillai's). Significant differences between IS and ID for all interactions
were reported from the MANOVA except for interaction between intersection and total
branch length (p = 0.0508) and intersections and number of branches (p = 0.0866),
(Supplemental Table 1). The strongest interaction was seen between radius and
primary axon length (p = 0.0000). Significant group-dependent differences in covariance
between number of branches and primary axon length (p = 0.0077) and number of
branches and average branch length (p = 0.0167) was captured in the multivariate
analysis.

Pairwise interactions are displayed as a pairplot with overlaid correlation values (R^2) for
IS and ID neurons (Supplemental Figure 1). Comparisons for regressions statistics
between IS and ID neurons for each pairwise interactions was also assessed and
displayed as a heatmap scaled by FDR-adjusted p-values but no differences were
observed (Supplemental Figure 2). A corresponding table (Supplemental Table 2)
displays the resulting linear regression statistics for each pairwise comparison
(Supplemental Figure 2), listing each x-and y-variable pair, effect size, upper and lower

confidence intervals, R^2 values, raw p-values and FDR adjusted p-values comparing
between IS and ID neurons (Supplemental Table 2). The resulting linear regression
statistics for each pairwise comparison (Supplemental Figure 1), was also tabulated for
each x-and y-variable pair, slope, intercept, R^2 values, raw p-values and FDR adjusted
p-values and listed for IS and ID neurons, independently (Supplemental Table 3). The
strongest pair association with significant differences was shown between branch length
and total axon length (IS: R^2 = 0.9103, p = 0.0000, ID: R^2 = 0.9481, p = 0.0000) and
total branch length and number of branches (IS: R^2 = 0.5636, p = 0.0000 , ID: R^2 =
0.6646, p = 0.0000) in both ID and IS neurons. Pairwise association between number
of branches and primary axon length was insignificant and not strongly correlated for
both IS and ID neurons (IS: R^2 =0.6248 , p = 0.7719, ID: R^2 = 0.1982, p = 0.44107).

Discussion:

Optimal brain circuit formation depends on proper establishment of neuronal polarity, a process that is specific and unique to neurons, and includes both axonogenesis and dendritogenesis (M. Zhang et al., 2019). Early-life ID impairs neurocircuit formation and function, particularly in circuits involving hippocampal pyramidal neurons (e.g., learning and memory) (Brunette et al., 2010; Carlson et al., 2010; Fretham et al., 2011, 2012; Georgieff et al., 2018; Jorgenson et al., 2003, 2005; Radlowski & Johnson, 2013). There has been extensive work done previously providing evidence of hippocampal neuron dendrite growth and branching deficits during early-life ID (Bastian et al., 2016, 2022; Brunette et al., 2010; Carlson et al., 2010; Fretham et al., 2012; Jorgenson et al., 2003). A simplified dendritic arbor provides a post-synaptic explanation for the neurocircuit behavioral impairments caused by fetal-neonatal ID. However, the effects of ID on the pre-synaptic, axonal side (outside of myelination deficits (Lozoff et al., 2006; Todorich et al., 2009) remain largely unknown. Axonogenesis is an energetically demanding process and thus energy production deficits often cause morphological aberrations in axonal development (Harris & Attwell, 2012). Neuronal ID reduced basal and ATP-linked mitochondrial respiration, ATP production rate, and neuronal ATP concentrations, suggesting a whole neuron level reduction in energy capacity of ID hippocampal neurons at 7DIV, during this period of rapid axon growth and branching. This is consistent with previous observations of impaired mitochondrial energetic capacity in the same neuron culture model, during the period of rapid dendritic growth and arborization (11-18DIV) (Bastian et al., 2016, 2019, 2022). This finding suggested that impaired iron-dependent mitochondrial function may also lead to axonal structural deficits, prior to the onset of rapid dendrite growth and branching, and influencing the subsequent dendrite growth/branching process.

It is important to advance knowledge on presynaptic ID effects as previous studies have shown that activity dependent axonal signaling directly influences hippocampal dendritic development through dendritic localization of neurotransmitter receptors, which are critical for activity-dependent dendrite growth and branching (Paus, 2023; Yamashita, Aoki, et al., 2016; Yamashita, Yamane, et al., 2016). If this signaling is disrupted, this results in reduced dendritic branching complexity (Paus, 2023; Yamashita, Aoki, et al., 2016; Yamashita, Yamane, et al., 2016). Neuronal network connections in hippocampus are important for synaptic plasticity such as with CA1 pyramidal dendrites which are influenced by connections with CA3 axons, signifying the importance of understanding the coupling activity between dendrite and axons for circuit development in ID conditions (Paus, 2023; Shepherd & Harris, 1998; Yamashita, Aoki, et al., 2016; Yamashita, Yamane, et al., 2016). In our primary embryonic mouse hippocampal neuron culture model of fetal-neonatal ID, iron chelation begins after axon initiation, which allows us to examine effects on axonal growth and branching without potential confounding effects on initial axon formation. ID caused gross axon morphological deficits with a reduction in overall axon complexity at 440μm-850μm from the soma,

which was caused by a reduction in axon and branch length but not branch number.
Multivariate analysis revealed group-dependent differences in covariance between
primary axon length to number of branches and the average branch length to number of
branches not captured in the univariate data. This would indicate that while the mean
number of branches are not affected by the effects of ID at this stage of axonal
development, the stochiometric relationship between axon or branch length and number
of branches is altered by ID (i.e. length of axons does not correlate with higher number
of branches). However, within group pairwise regression statistic comparisons for
variable interactions between length variables (primary axon and average branch
length) and number of branches revealed no significant differences. Thus, our findings
suggest that iron-dependent mitochondrial energetics during early axonogenesis are
critical for axonal growth and elongation but not for branch number as summarized in
illustration for Figure 7. Stunted elongation of the primary axon and branch length
indicates that the proper structural development of neurons requires sufficient iron much
earlier than previously thought and has implications for axons facing the challenge of
reaching distant targets during neonatal brain development. Whether this decrease in
axonal complexity in ID hippocampal neurons occurs during *in vivo* early axonogenesis
is an important unanswered question that has critical implications for overall
developmental circuitry deficits and long-term propagation dynamics (Ofer et al., 2017).

Since anterograde trafficking of cargo is crucial to meet the demands of developing
neuronal processes (Millecamps & Julien, 2013), decreased mitochondrial motility was
hypothesized to contribute to the stunted axonal growth due to ID. Consistent with
unchanged dendrite mitochondrial density at 11 DIV (Bastian et al., 2019), mitochondrial
density was unaltered in the terminal (50-100μm from beginning of growth cone) of 7
DIV iron-deficient axons. Previous work with our hippocampal neuron culture model of
ID has shown that at 11 DIV, ID reduces average mitochondrial speed due to increased
pause frequency and decreases anterograde motion of moving mitochondria to energy-
demanding terminal processes of developing dendrites, ultimately leading to reduced
mitochondrial density in terminal dendrites at 18DIV (Bastian et al., 2019). However,
during hippocampal neuron axonogenesis, significant differences in mitochondrial
motility were not observed between IS and ID neurons. Given ID effects on
mitochondrial energy production during axon growth and the known role of ATP in
axonal mitochondrial motility (Bastian, 2019; Lewis et al., 2013; Sheng, 2014, 2017),
this finding was unexpected. These data suggest that axonal mitochondria trafficking is
prioritized over other ATP-dependent processes and indicates that there are potential
compensatory mechanisms to maintain trafficking during neuronal ID (Cheng et al.,
2022; Sheng, 2014; Wang et al., 2021). Overall, these findings imply that the
developmental impact of ID on axon growth is due to impaired mitochondrial
bioenergetics and ATP production but not reduced mitochondrial trafficking to the
growth cone (Figure 7).

Although iron's role in mitochondrial energetics is critical to cellular health and
development, iron is necessary for many additional important cellular processes that

could contribute to impaired axon development during ID such as regulation of gene
transcription, neurotransmitter synthesis, and myelination (Hare et al., 2013). The
reduced non-mitochondrial respiration observed in iron-deficient neurons likely indicates
reduced activity of other iron-dependent oxygen consuming enzymes (e.g., Tet
dioxygenases, lysine histone demethylases (Kdms), Prolyl hydroxylase, desaturases,
etc.). Tet and Kdm enzymes catalyze the demethylation of DNA and histones,
respectively, and are responsible for iron-dependent epigenetic regulation of
transcription for key neurodevelopmental genes (e.g., Bdnf) (Barks et al., 2018; Tran et
al., 2015). In addition, the iron-dependent enzyme, prolyl hydroxylase, is necessary for
the regulation of hypoxia inducible factor 1 α (HIF1α), a regulator of many metabolic and
neurodevelopmental genes (Hare et al., 2013; Xu et al., 2017; D. L. Zhang et al., 2014).
Iron-containing desaturase enzymes are responsible for the production of mono- and
poly-unsaturated fatty acids, key metabolic substrates and components of neuronal
membranes (Bazinet & Layé, 2014). Future work is needed to explore the prioritization
of iron to these enzymes during neuronal ID and their relative contribution to iron-
dependent neuron structural development.

An important unanswered question is why ID-induced mitochondrial energy deficits
impair axonal growth but not branch number (Figure 7). Tao *et al* showed mitochondrial
ATP dependent branch formation in immature hippocampal neurons with stationary
mitochondria present at branch points (Tao et al., 2014). In early sensory axons,
impairment of neural growth factor (NGF) mediated mitochondrial respiration impaired
actin but not microtubule polymerization indicating NGF mediated axonal branching
requires local ATP regulation of actin dynamics (Smith & Gallo, 2018; Spillane et al.,
2013). Since we examined only whole cell mitochondrial respiration and ATP
production, it is unclear whether the local ATP concentration near axon branch points
was maintained or not. Since it is known that stationary mitochondria along neuronal
processes are important for branch formation, it is possible that *de novo* axonal
branching does not necessarily depend on mitochondrial ATP production but rather
other mitochondrial activities (e.g. calcium buffering). Since we did not observe a
change in axonal mitochondrial motility or density, one explanation is that branch
number is maintained because ID does not alter mitochondrial localization near branch
points. Future studies with temporal and spatial resolution of ATP dynamics targeted to
subcellular regions of interest (e.g., branch points and growth cone) are necessary to
better understand the relationship between mitochondrial energetics and axonal
growth/branching during ID (Marvin et al., 2024). In addition, many other mechanisms
(e.g., mRNA trafficking, actin, and microtubule polymerization, and recruitment of other
organelles) are also involved in determining branch formation (Lewis et al., 2013; Smith
& Gallo, 2018) and will need to be explored during neuronal ID.

Most pre-clinical studies on the effects of fetal-neonatal ID on brain development have
focused on postnatal neurodevelopmental processes (e.g., dendritic arborization,
myelination, and synapse formation). ID-induced axon structural deficits have
implications for the proper neurocircuit sculpting of the brain starting during embryonic

development. Translationally, these findings provide biological plausibility for shifting the
focus from treating iron-deficient children to monitoring and management of maternal
iron status throughout pregnancy (Georgieff et al., 2018). Indeed, several clinical
studies demonstrate that maternal iron status during pregnancy predicts the risk of
offspring developing autism spectrum, attention deficit hyperactive, and schizophrenia
disorders (reviewed in (Georgieff et al., 2018; Maxwell & Rao, 2022); all thought to be
disorders of circuit construction. This is further supported by studies demonstrating that
gestational ID alters the balance of excitatory to inhibitory neurons in the embryonic
brain. The long-term implications of impaired iron-dependent neurodevelopment are
further underscored by studies showing that early-life ID causes Parkinsonian-like
neurodegeneration (Matak et al., 2016), highlighting the need for early interventions.
More efforts have been focusing on using and developing mitochondrial targeted
therapies to treat neurodegenerative and neuropsychiatric disorders, thus showing a
converging interest at optimizing mitochondrial health (Andreyev et al., 2024; Callegari
et al., 2025; Włodarczyk et al., 2018). Improved understanding of the interface between
ID and mitochondrial function in early-life neuronal development could bring about
preventative solutions and interventions to optimize brain health for populations at risk
of ID as well as those suffering from seemingly unrelated brain-health disorders with
potential origins in iron dysregulation. Altogether, our observations underscore the need
to devote future efforts to understanding the effects of ID (and other common nutrient
disruptions) on embryonic brain development processes to understand the stage of
development that interventions are required to prevent the long-lasting neurobehavioral
deficits caused by fetal-neonatal ID.

Figures and Tables:

Figure 1: ID decreases total intracellular ATP concentrations in 7 DIV hippocampal neurons during axonogenesis. A) Illustration of hippocampal neuron culture experimental design (Created in BioRender). Primary hippocampal neuronal cultures from E16 mice were treated with 5-FU and with or without DFO at 3 DIV. **B)** At 7DIV, quantitative real-time PCR (qPCR) was performed for *Tfrc*, the gene coding for the main neuronal iron uptake protein, to validate iron status of treatment groups (n= 6 independent wells per group pooled from two unique culture preparations). *Tfr1* mRNA levels were calculated relative to the geometric mean of three reference genes (*Tbp*, *Rps18*, and *Gapdh*) and relative change in *Tfr1* mRNA is compared from experiments in 6-well and 96-well vessels. **C)** Total intracellular ATP concentration was measured and normalized to cell count per well (n = 32-36 independent wells per group). Statistical significance was determined on log10-scaled data by unpaired t-test with or without Welch's correction for qPCR in 96-well and 6-well, respectively, comparing ID induced DFO dose to IS neurons and displayed on linear scale. The parameteric unpaired t- test was used on relativized and on log10-scaled intracellular ATP assay for determining significance between IS versus ID neurons. Statistical significance is denoted with asterisk (*) for p-values ≤ 0.05. For both B and C individual data points representing one independent replicate are calculated relative to the average for the IS group and are shown along with the mean ± SD. Relative data are shown on a log scale to accurately reflect the magnitude of changes.

**Figure 2: ID impairs mitochondrial respiratory capacity in 7 DIV hippocampal
neurons.** Primary hippocampal neuronal cultures from E16 mice were treated with 5-FU
and with or without DFO at 3 DIV after axon specification and assessed for their
respiratory capacity using ATP Rate Assay. **A)** Kinetic plot of real-time OCR normalized
to cell count (pmol/min/1000 cells) measured before and following injections of
Oligomycin and Antimycin A/Rotenone (AA/Rot). **B-D)** Normalized respiration
parameters for non-mitochondrial respiration, basal respiration, and ATP-linked
respiration were derived from the real-time OCR data. **E-F)** OCR and ECAR data was
used to calculate the ATP production rate from either mitochondrial oxidative
phosphorylation or glycolysis. **G)** The sum of both ATP production rates was used to
calculate the total ATP production rate. **H)** The ATP Rate Index was calculated by the
ratio of mitochondrial ATP production (mito ATP) rate over the glycolytic ATP (glyco
ATP) production rate. Data were pooled from three unique culture preparations (n=11-
14 independent wells per group). Data points for values of independent wells with the
mean ± SD are displayed on each plot. An asterisk (*) indicates a statistically significant
difference (p≤0.05) by parametric unpaired-test or non-parametric unpaired Mann-
Whitney test for each respiration parameter.

Figure 3: Axonal mitochondrial motility kymograph analysis in 7DIV hippocampal neurons. Primary hippocampal neuronal cultures from E16 mice were treated with 5-FU and with or without DFO at 3 DIV. At 7 DIV, time-lapse imaging (1s or 3.5s intervals for 1.7 or 5min) was performed to capture motility of fluorescently labelled mitochondria in axonal compartment of neurons transfected with mcherry-Mito-7. **A)** Illustration of axonal region of interest analyzed and mitochondrial trafficking activity (created in BioRender). B) Representative still frame (t=0 min) of fluorescently labelled axonal mitochondria is shown with a representative kymograph of IS and ID. To better visualize the traced axonal path, the images were straightened using ImageJ "Straightened" tool. Tracks in kymographs represent individual mitochondria over time (x-axis) and direction (y-axis). Manually traced tracks are color-coded by category of direction of mitochondrial motion relative to the soma and are defined as retrograde (magenta), reversing (yellow), anterograde (cyan), and stationary (blue). Scale bars represent 5μm.

Movie 1: Representative movies of IS and ID mitochondrial motility in terminal axon processes. Movies show time-lapse imaging of mCherry-Mito-7 positive mitochondria in 7 DIV **A)** IS and **B)** ID axonal terminal processes from E16 hippocampal neuronal cultures treated with 5-FU and DFO at 3 DIV. The mitochondria for this representative movie were imaged every 1s for 1.7 min and movies are set to 30 frames per second. Images were straightened using the Fiji "Straightened" function to better visualize the region of interest in the axon terminal. Scale bars represent 5µm.

653 **Table 1: Definitions of mitochondrial motility parameters**

Table 1. Axonal mitochondrial motility parameters		
Parameter	Definition	Figure
Mitochondrial Densities		
Anterograde density	# of mitochondria moving in anterograde direction with or without pausing per µm axon	3C
Reversal density	# of mitochondria with at least one change in motion directionality per µm axon	3C
Reversal density	# of mitochondria moving in retrograde direction with or without pausing per µm axon	-----
Moving density	sum of # of moving mitochondria per µm axon for all motions (anterograde, reversal, and retrograde)	-----
Stationary density	sum of # of stationary mitochondria per per µm axon	3C
Total density	Total number of mitochondria per µm axon (moving and stationary mitochondria)	
Average Speed and Net Velocity		
Average Speed	Average speed of individual mitochondria including pauses, with or without net velocities, and all directionalities (anterograde, reversal, and retrograde)	3A
Net velocity	Average net velocity of all moving mitochondria with net motion () irrespective of directionality	3A
Net anterograde velocity	Average net velocity of all moving mitochondria with net motion () in the anterograde direction	3A
Net retrograde velocity	Average net velocity of all moving mitochondria with net motion () in the retrograde direction	3A
Segmental Velocity		
Average segmental speed	Average of all segmental speeds (anterograde and retrograde) per mitochondrion regardless of direction	3B
Anterograde segmental velocity	Average of combined velocity segments in anterograde direction	3B
Retrograde segmental velocity	Average of combined velocity segments in retrograde direction	3B
Segmental Run Length		
Average run length	Average length of combined segmental runs regardless of direction per mitochondrion	-----
Anterograde run length	Average length of segmental run in the anterograde direction	-----
Retrograde run length	Average length of segmental run in the retrograde direction	-----
Percent Time of Motion		
Percent time moving	Total percent time mitochondria are in motion (anterograde or retrograde), excluding pauses	3D
Percent time anterograde motion	Total percent time moving mitochondria spend in anterograde direction	3D
Percent time retrograde motion	Total percent time moving mitochondria spend in retrograde direction	3D
Percent time pausing	Total percen time mitochondria spend pausing	3D
Pause Frequency		
Total pause frequency	Average number of pause per moving mitochondria regardless of direction (anterograde or retrograde) between pauses	3D
Anterograde pause frequency	# of pauses per mitochondria between movements in the anterograde direction	3D
Retrograde pause frequency	# of pauses per mitochondria between movements in the retrograde direction	3D
Reversal pause frequency	# of pauses per mitochondria between reversal movements	3D
Pause Duration		
Total pause duration	Average duration of all pauses between all time segments in motion for all directionalities (anterograde, reversal and retrograde) per mitochondria	3D
Anterograde pause duration	Average duration of all pauses between time spent by mitochondria moving in anterograde direction per moving mitochondria	3D
Retrograde pause duration	Average duration of all pauses between time spent by mitochondria moving in retrograde direction per moving mitochondria	3D
Reversal pause duration	Average duration of all pauses between time spent by mitochondria reversing direction per moving mitochondria	3D

Figure 4: ID does not alter mitochondrial motility in 7DIV axons. A) Average speed for all moving mitochondria (regardless of direction or net displacement) was calculated. The net velocity represents any mitochondria (IS: n=98; ID: n= 70) displaying net movement in the anterograde or retrograde direction and displayed as a scalar value (speed μm/s). **B)** The average segmental speed was calculated for each mitochondria only when in active motion (excluding pauses) (IS: n=98; ID: n=70). All individual moving segments were pooled and sub-categorized by anterograde (IS: n=486; ID: n=364) and retrograde (IS: n=391; ID: n=267) directionalities to calculate average segmental speed. All values are represented in scalar form (μm/s). **C)** The total density of mitochondria per axonal length and the density of sub-populations of mitochondria that showed overall anterograde, reversal, or retrograde movement were plotted for IS and ID axons (IS: n=11; ID: n=12). **D)** The pause frequency (PF) of all pausing instances (IS: n=98; ID: n=70) of individual mitochondria for IS and ID was plotted. The PF of individual mitochondria with overall anterograde (IS: n=84; ID: n=58), reversing (IS: n=58; ID: n=31), or retrograde (IS: n=72; ID: n=43) directionality was also extracted and plotted. Time duration in seconds (s) for all pauses (IS: n=79; ID: n=49) was calculated and subdivided for anterograde (IS: n=32; ID: n=26), reversing (IS: n=21; ID: n=7), and retrogade (IS: n=26; ID n=16) directions. Total percent time in motion for all moving mitochondria regardless of direction (excluding pauses) was calculated and subdivided for anterograde, reversing and retrograde (IS: n=98; ID: n=70) directions. The percent time mitochondria spend pausing was also calculated and plotted (IS: n=98; ID: n=70). All scatter plots display the median with the interquartile range from data pooled from two unique neuron culture preparations. An unpaired Mann-Whitney test was performed for all mitochondrial motility, density, and population parameters to compare the medians of the non-parametric datasets with alpha threshold = 0.05. Statistically significant discoveries (*) were determined after correction with FDR analysis using Benjamini-Hochberg method (q=5%).

Figure 5: ID impairs hippocampal neuron axonal growth. Hippocampal neuronal cultures isolated from E16 mice were nucleofected with an hrGFP-expressing plasmid at time of plating. Cultures were then treated with 5-FU and with or without DFO at 3 DIV after axon specification. At 7 DIV, the neuronal cultures were fixed and processed for immunocytochemistry against MAP2 (dendrite specific marker). Individual hrGFP-positive neurons were imaged using a Zeiss Celldiscoverer 7 and Zeiss Axiocam 506 mono camera with a 20x (NA 0.7) objective and 0.5x magnification. The hrGFP-positive and MAP2-negative axon was distinguished from MAP2/hrGFP double-positive dendrites to complete axonal tracing using FIJI and Simple Neurite Tracer. **A)** Two representative grayscale images are shown of iron-sufficient and -deficient axons with corresponding hrGFP (neuron filler), MAP2 (dendrites), merged hrGFP/MAP2/DAPI channels, and skeletonized tracing. **B)** Sholl-curve (mean±SEM; n=100 neurons). An asterisk (*) indicates statistical significance at a given distance from the soma by repeated measures 2-way ANOVA and Tukey's post-hoc test. Univariate morphological measurements were calculated from simple neurite tracer analysis of **C)** primary axon length and **D)** total axon length. Data points representing individual neurons (n=100) are displayed with Mean ± SD. An asterisk (*) indicates statistical significance via Welch's t-test (p-values ≤ 0.05).

A

hrGFP	MAP2	Merged	Trace

Iron-Sufficient

Iron-Deficient

B Axon Sholl Analysis

C Primary Axon Length

D Total Axon Length

707 **Figure 6: Iron deficiency impairs hippocampal neuron axonal branch growth but**
708 **not formation.** Axonal traces (see Fig 5) were used to measure and calculate **A)**
709 average branch length, **B)** total branch length, and **C)** number of branches per axon.
710 Data points representing individual neurons (n=100) are displayed with Mean ± SD. An
711 asterisk (*) indicates statistical significance via Welch's t-test (p-values ≤ 0.05).

721 **Figure 7: Summary of ID effects on axonal mitochondria during axonogenesis:**
722 Axonal mitochondrial trafficking and branch number appears to be prioritized in ID
723 neurons over energy production and axon growth**.**

Author Contributions:

D.C.M., K.D., and T.R.M. and T.W.B wrote the first draft of the paper; D.C.M., T.W.B, M.K.G., and L.M.L. designed the experiments and edited the paper; D.C.M., K.D., L.H.C., D.J.M. and T.W.B. performed the experiments; D.C.M., K.D., T.W.B. and T.R.M. analyzed the data; The paper was read and approved by all authors before submission.

Funding Sources and Disclosures:

The work from these studies was supported by the National Institute of Health (NIH) Grant R01HD094809 to M.K.G., NIH R21HD106043 to T.W.B. and the Howard Hughes Medical Institute (HHMI) Gilliam Fellows program award GT15879 to T.W.B. and D.C.M. T.R.M. was supported by an NIH T32 training grant (T32HL007062).

Conflict of Interest:

The authors declare no conflict of interest to report.

Code and Data Availability:

Acknowledgements:

With utmost gratefulness, we acknowledge the support of other lab members of the Bastian, Georgieff and Lanier laboratories for their assistance with preparation of cell cultures, troubleshooting and valuable feedback.

References:

Andreyev, A. Y., Yang, H., Doulias, P. T., Dolatabadi, N., Zhang, X., Luevanos, M., Blanco, M., Baal, C., Putra, I., Nakamura, T., Ischiropoulos, H., Tannenbaum, S. R., & Lipton, S. A. (2024). Metabolic Bypass Rescues Aberrant S-nitrosylation-Induced TCA Cycle Inhibition and Synapse Loss in Alzheimer's Disease Human Neurons. *Advanced Science*, *11*(12). https://doi.org/10.1002/advs.202306469

Appolus EE, & Okoli CN. (2022). A Robust Comparison Powers of Four Multivariate Analysis of Variance Tests. *European Journal of Statistics and Probability*, *10*(1), 11–20. https://doi.org/10.37745/ejsp.2013

Barks, A., Fretham, S. J. B., Georgieff, M. K., & Tran, P. V. (2018). Early-Life Neuronal-Specific Iron Deficiency Alters the Adult Mouse Hippocampal Transcriptome. *Journal of Nutrition*, *148*(10), 1521–1528. https://doi.org/10.1093/jn/nxy125

Bastian, T. W. (2019). Potential mechanisms driving mitochondrial motility impairments in developing iron-deficient neurons. *Journal of Experimental Neuroscience*, *13*. https://doi.org/10.1177/1179069519858351

Bastian, T. W., Rao, R., Tran, P. V., & Georgieff, M. K. (2020). The Effects of Early-Life Iron Deficiency on Brain Energy Metabolism. In *Neuroscience Insights* (Vol. 15). SAGE Publications Ltd. https://doi.org/10.1177/2633105520935104

Bastian, T. W., von Hohenberg, W. C., Georgieff, M. K., & Lanier, L. M. (2019). Chronic energy depletion due to iron deficiency impairs dendritic mitochondrial motility during hippocampal neuron development. *Journal of Neuroscience*, *39*(5), 802–813. https://doi.org/10.1523/JNEUROSCI.1504-18.2018

Bastian, T. W., Von Hohenberg, W. C., Kaus, O. R., Lanier, L. M., & Georgieff, M. K. (2022). Choline Supplementation Partially Restores Dendrite Structural Complexity in Developing Iron-Deficient Mouse Hippocampal Neurons. *Journal of Nutrition*, *152*(3), 747–757. https://doi.org/10.1093/jn/nxab429

Bastian, T. W., Von Hohenberg, W. C., Mickelson, D. J., Lanier, L. M., & Georgieff, M. K. (2016). Iron Deficiency Impairs Developing Hippocampal Neuron Gene Expression, Energy Metabolism, and Dendrite Complexity. *Developmental Neuroscience*, *38*(4), 264–276. https://doi.org/10.1159/000448514

Bazinet, R. P., & Layé, S. (2014). Polyunsaturated fatty acids and their metabolites in brain function and disease. In *Nature Reviews Neuroscience* (Vol. 15, Issue 12, pp. 771–785). Nature Publishing Group. https://doi.org/10.1038/nrn3820

Brunette, K. E., Tran, P. V., Wobken, J. D., Carlson, E. S., & Georgieff, M. K. (2010). Gestational and neonatal iron deficiency alters apical dendrite structure of CA1 pyramidal neurons in adult rat hippocampus. *Developmental Neuroscience*, *32*(3), 238–248. https://doi.org/10.1159/000314341

Callegari, S., Kirk, N. S., Gan, Z. Y., Dite, T., Cobbold, S. A., Leis, A., Dagley, L. F., Glukhova, A., & Komander, D. (2025). Structure of human PINK1 at a mitochondrial TOM-VDAC array. *Science*. https://doi.org/10.1126/science.adu6445

Carlson, E. S., Fretham, S. J. B., Unger, E., O'Connor, M., Petryk, A., Schallert, T., Rao, R., Tkac, I., & Georgieff, M. K. (2010). Hippocampus specific iron deficiency alters competition and cooperation between developing memory systems. *Journal of Neurodevelopmental Disorders*, *2*(3), 133–143. https://doi.org/10.1007/s11689-010-9049-0

Carlson, E. S., Stead, J. D. H., Neal, C. R., Petryk, A., & Georgieff, M. K. (2007). Perinatal iron deficiency results in altered developmental expression of genes mediating energy metabolism and neuronal morphogenesis in hippocampus. *Hippocampus*, *17*(8), 679–691. https://doi.org/10.1002/hipo.20307

Carlson, E. S., Tkac, I., Magid, R., O'Connor, M. B., Andrews, N. C., Schallert, T., Gunshin, H., K.Georgieff, M., & Petryk, A. (2009). Iron is essential for neuron development and memory function in mouse hippocampus 1-3. *Journal of Nutrition*, *139*(4), 672–679. https://doi.org/10.3945/jn.108.096354

Cheng, X. T., Huang, N., & Sheng, Z. H. (2022). Programming axonal mitochondrial maintenance and bioenergetics in neurodegeneration and regeneration. In *Neuron* (Vol. 110, Issue 12, pp. 1899–1923). Cell Press. https://doi.org/10.1016/j.neuron.2022.03.015

Cheng, X. T., & Sheng, Z. H. (2021). Developmental regulation of microtubule-based trafficking and anchoring of axonal mitochondria in health and diseases. In *Developmental Neurobiology* (Vol. 81, Issue 3, pp. 284–299). John Wiley and Sons Inc. https://doi.org/10.1002/dneu.22748

Deungria, M., Rao, R., Wobken, J. D., Luciana, M., Nelson, C. A., & Georgieff, M. K. (2000). Perinatal Iron Deficiency Decreases Cytochrome c Oxidase (CytOx) Activity in Selected Regions of Neonatal Rat Brain. In *Pediatric Research*. https://doi.org/https://doi.org/10.1203/00006450-200008000-00009

Dotti, C. G., Sullivan, C. A., & Banker, G. A. (1988). The Establishment of Polarity by Hippocampal Neurons in Culture. In *The Journal of Neuroscience: Vol. I* (Issue 4).

Fretham, S. J. B., Carlson, E. S., & Georgieff, M. K. (2011). The role of iron in learning and memory. *Advances in Nutrition*, *2*(2), 112–121. https://doi.org/10.3945/an.110.000190

Fretham, S. J. B., Carlson, E. S., Wobken, J., Tran, P. V., Petryk, A., & Georgieff, M. K. (2012). Temporal manipulation of transferrin-receptor-1-dependent iron uptake identifies a sensitive period in mouse hippocampal neurodevelopment. *Hippocampus*, *22*(8), 1691–1702. https://doi.org/10.1002/hipo.22004

Gao, Q., Zhou, Y., Chen, Y., Hu, W., Jin, W., Zhou, C., Yuan, H., Li, J., Lin, Z., & Lin, W. (2025). Role of iron in brain development, aging, and neurodegenerative diseases. *Annals of Medicine*, *57*(1), 2472871. https://doi.org/10.1080/07853890.2025.2472871

Georgieff, M. K., Ramel, S. E., & Cusick, S. E. (2018). Nutritional influences on brain development. In *Acta Paediatrica, International Journal of Paediatrics* (Vol. 107, Issue 8, pp. 1310–1321). Blackwell Publishing Ltd. https://doi.org/10.1111/apa.14287

Grabrucker, A., Vaida, B., Bockmann, J., & Boeckers, T. M. (2009). Synaptogenesis of hippocampal neurons in primary cell culture. *Cell and Tissue Research*, *338*(3), 333–341. https://doi.org/10.1007/s00441-009-0881-z

Greminger, A. R., & Mayer-Pröschel, M. (2015). Identifying the threshold of iron deficiency in the
central nervous system of the rat by the auditory brainstem response. *ASN Neuro*, *7*(1), 1–
10. https://doi.org/10.1177/1759091415569911

Hare, D., Ayton, S., Bush, A., & Lei, P. (2013). A delicate balance: Iron metabolism and
diseases of the brain. In *Frontiers in Aging Neuroscience* (Vol. 5, Issue JUL).
https://doi.org/10.3389/fnagi.2013.00034

Harris, J. J., & Attwell, D. (2012). The energetics of CNS white matter. *Journal of Neuroscience*,
32(1), 356–371. https://doi.org/10.1523/JNEUROSCI.3430-11.2012

Insel, B. J., Schaefer, C. A., Mckeague, I. W., Susser, E. S., & Brown, A. S. (2008). Maternal
Iron Deficiency and the Risk of Schizophrenia in Offspring. In *Arch Gen Psychiatry* (Vol. 65,
Issue 10).

Jorgenson, L. A., Wobken, J. D., & Georgieff, M. K. (2003). Perinatal Iron Deficiency Alters
Apical Dendritic Growth in Hippocampal CA1 Pyramidal Neurons. *Developmental
Neuroscience*, *25*(6), 412–420. https://doi.org/10.1159/000075667

Kim, J. W., Byun, M. S., Yi, D., Lee, J. H., Jeon, S. Y., Ko, K., Joung, H., Jung, G., Lee, J. Y.,
Sohn, C. H., Lee, Y. S., Kim, Y. K., & Lee, D. Y. (2021). Blood Hemoglobin, in-vivo
Alzheimer Pathologies, and Cognitive Impairment: A Cross-Sectional Study. *Frontiers in
Aging Neuroscience*, *13*. https://doi.org/10.3389/fnagi.2021.625511

Konofal, E., Lecendreux, M., Arnulf, I., & Mouren, M.-C. (2004). *Iron Deficiency in Children With
Attention-Deficit/Hyperactivity Disorder*.

Lewis, T. L., Courchet, J., & Polleux, F. (2013). Cellular and molecular mechanisms underlying
axon formation, growth, and branching. In *Journal of Cell Biology* (Vol. 202, Issue 6, pp.
837–848). https://doi.org/10.1083/jcb.201305098

Lozoff, B., Beard, J., Connor, J., Felt, B., Georgieff, M., & Schallert, T. (2006). *Long-Lasting
Neural and Behavioral Effects of Iron Deficiency in Infancy*.

Lozoff, B., & Georgieff, M. K. (2006). Iron Deficiency and Brain Development. *Seminars in
Pediatric Neurology*, *13*(3), 158–165.

Lozoff, B., Smith, J. B., Kaciroti, N., Clark, K. M., Guevara, S., & Jimenez, E. (2013). Functional
significance of early-life iron deficiency: Outcomes at 25 years. *Journal of Pediatrics*,
163(5), 1260–1266. https://doi.org/10.1016/j.jpeds.2013.05.015

Lukowski, A. F., Koss, M., Burden, M. J., Jonides, J., Nelson, C. A., Kaciroti, N., Jimenez, E., &
Lozoff, B. (2010). Iron deficiency in infancy and neurocognitive functioning at 19 years:
Evidence of long-term deficits in executive function and recognition memory. *Nutritional
Neuroscience*, *13*(2), 54–70. https://doi.org/10.1179/147683010X12611460763689

Marvin, J. S., Kokotos, A. C., Kumar, M., Pulido, C., Tkachuk, A. N., Yao, J. S., Brown, T. A., &
Ryan, T. A. (2024). iATPSnFR2: A high-dynamic-range fluorescent sensor for monitoring
intracellular ATP. *Proceedings of the National Academy of Sciences of the United States of
America*, *121*(21). https://doi.org/10.1073/pnas.2314604121

Matak, P., Matak, A., Moustafa, S., Aryal, D. K., Benner, E. J., Wetsel, W., & Andrews, N. C. (2016). Disrupted iron homeostasis causes dopaminergic neurodegeneration in mice. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(13), 3428–3435. https://doi.org/10.1073/pnas.1519473113

Maxwell, A. M., & Rao, R. B. (2022). Perinatal iron deficiency as an early risk factor for schizophrenia. In *Nutritional Neuroscience* (Vol. 25, Issue 10, pp. 2218–2227). Taylor and Francis Ltd. https://doi.org/10.1080/1028415X.2021.1943996

Millecamps, S., & Julien, J. P. (2013). Axonal transport deficits and neurodegenerative diseases. In *Nature Reviews Neuroscience* (Vol. 14, Issue 3, pp. 161–176). https://doi.org/10.1038/nrn3380

Monko, T. R., Tripp, E. H., Burr, S. E., Gunderson, K. N., Lanier, L. M., Georgieff, M. K., & Bastian, T. W. (2024a). Cellular Iron Deficiency Disrupts Thyroid Hormone Regulated Gene Expression in Developing Hippocampal Neurons. *Journal of Nutrition*, *154*(1), 49–59. https://doi.org/10.1016/j.tjnut.2023.11.007

Monko, T. R., Tripp, E. H., Burr, S. E., Gunderson, K. N., Lanier, L. M., Georgieff, M. K., & Bastian, T. W. (2024b). Cellular Iron Deficiency Disrupts Thyroid Hormone Regulated Gene Expression in Developing Hippocampal Neurons. *Journal of Nutrition*, *154*(1), 49–59. https://doi.org/10.1016/j.tjnut.2023.11.007

Neumann, S., Chassefeyre, R., Campbell, G. E., & Encalada, S. E. (2017). KymoAnalyzer: a software tool for the quantitative analysis of intracellular transport in neurons. *Traffic*, *18*(1), 71–88. https://doi.org/10.1111/tra.12456

Ofer, N., Shefi, O., & Yaari, G. (2017). Branching morphology determines signal propagation dynamics in neurons. *Scientific Reports*, *7*(1). https://doi.org/10.1038/s41598-017-09184-3

Paus, T. (2023). Tracking Development of Connectivity in the Human Brain: Axons and Dendrites. In *Biological Psychiatry* (Vol. 93, Issue 5, pp. 455–463). Elsevier Inc. https://doi.org/10.1016/j.biopsych.2022.08.019

Penrod, R. D., Campagna, J., Panneck, T., Preese, L., & Lanier, L. M. (2015). The presence of cortical neurons in striatal-cortical co-cultures alters the effects of dopamine and BDNF on medium spiny neuron dendritic development. *Frontiers in Cellular Neuroscience*, *9*(July), 1–14. https://doi.org/10.3389/fncel.2015.00269

Perng, V., Li, C., Klocke, C. R., Navazesh, S. E., Pinneles, D. K., Lein, P. J., & Ji, P. (2021). Iron Deficiency and Iron Excess Differently Affect Dendritic Architecture of Pyramidal Neurons in the Hippocampus of Piglets. *Journal of Nutrition*, *151*(1), 235–244. https://doi.org/10.1093/jn/nxaa326

Petry, C. D., Eaton, M. A., Wobken, J. D., Mills, M. M., Johnson, D. E., & Georgieff, M. K. (1992). *Iron deficiency of liver, heart, and brain in newborn infants of diabetic mothers*. https://doi.org/https://doi.org/10.1016/s0022-3476(05)82554-5

Radlowski, E. C., & Johnson, R. W. (2013a). Perinatal iron deficiency and neurocognitive development. In *Frontiers in Human Neuroscience* (Issue SEP). Frontiers Media S. A. https://doi.org/10.3389/fnhum.2013.00585

Radlowski, E. C., & Johnson, R. W. (2013b). Perinatal iron deficiency and neurocognitive
development. In *Frontiers in Human Neuroscience* (Issue SEP). Frontiers Media S. A.
https://doi.org/10.3389/fnhum.2013.00585

R., Rao, R., Tkac, I., Townsend, E. L., Gruetter, R., & Georgieff, M. K. (2003). Nutritional
Neurosciences Perinatal Iron Deficiency Alters the Neurochemical Profile of the
Developing. In *J. Nutr* (Vol. 133, pp. 3215–3221).
https://doi.org/https://doi.org/10.1093/jn/133.10.3215

Sahu, M. P., Nikkilä, O., Lagas, S., Kolehmainen, S., & Castrén, E. (2019). Culturing primary
neurons from rat hippocampus and cortex. *Neuronal Signaling*, *3*(2).
https://doi.org/10.1042/NS20180207

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V.,
Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for
biological-image analysis. In *Nature Methods* (Vol. 9, Issue 7, pp. 676–682).
https://doi.org/10.1038/nmeth.2019

Schmidt, R. J., Tancredi, D. J., Krakowiak, P., Hansen, R. L., & Ozonoff, S. (2014). Maternal
intake of supplemental iron and risk of autism spectrum disorder. *American Journal of
Epidemiology*, *180*(9), 890–900. https://doi.org/10.1093/aje/kwu208

Semple, B. D., Blomgren, K., Gimlin, K., Ferriero, D. M., & Noble-Haeusslein, L. J. (2013). Brain
development in rodents and humans: Identifying benchmarks of maturation and
vulnerability to injury across species. In *Progress in Neurobiology* (Vols. 106–107, pp. 1–
16). https://doi.org/10.1016/j.pneurobio.2013.04.001

Sheng, Z. H. (2014). Mitochondrial trafficking and anchoring in neurons: New insight and
implications. In *Journal of Cell Biology* (Vol. 204, Issue 7, pp. 1087–1098). Rockefeller
University Press. https://doi.org/10.1083/jcb.201312123

Sheng, Z. H. (2017). The Interplay of Axonal Energy Homeostasis and Mitochondrial Trafficking
and Anchoring. In *Trends in Cell Biology* (Vol. 27, Issue 6, pp. 403–416). Elsevier Ltd.
https://doi.org/10.1016/j.tcb.2017.01.005

Shepherd, G. M. G., & Harris, K. M. (1998). Three-Dimensional Structure and Composition of
CA3 --> CA1 Axons in Rat Hippocampal Slices: Implications for Presynaptic Connectivity
and Compartmentalization. *Journal of Neuroscience*.

Smith, G. M., & Gallo, G. (2018). The role of mitochondria in axon development and
regeneration. In *Developmental Neurobiology* (Vol. 78, Issue 3, pp. 221–237). John Wiley
and Sons Inc. https://doi.org/10.1002/dneu.22546

Spillane, M., Ketschek, A., Merianda, T. T., Twiss, J. L., & Gallo, G. (2013). Mitochondria
Coordinate Sites of Axon Branching through Localized Intra-axonal Protein Synthesis. *Cell
Reports*, *5*(6), 1564–1575. https://doi.org/10.1016/j.celrep.2013.11.022

Tao, K., Matsuki, N., & Koyama, R. (2014). AMP-activated protein kinase mediates activity-
dependent axon branching by recruiting mitochondria to axon. *Developmental
Neurobiology*, *74*(6), 557–573. https://doi.org/10.1002/dneu.22149

Todorich, B., Pasquini, J. M., Garcia, C. I., Paez, P. M., & Connor, J. R. (2009).
Oligodendrocytes and myelination: The role of iron. In *GLIA* (Vol. 57, Issue 5, pp. 467–
478). https://doi.org/10.1002/glia.20784

Tran, P. V, Kennedy, B. C., Lien, Y.-C., Simmons, R. A., & Georgieff, M. K. (2015). *Fetal iron*
deficiency induces chromatin remodeling at the Bdnf locus in adult rat hippocampus.
https://doi.org/10.1152/ajpregu.00429.2014.-Fetal

Wang, B., Huang, M., Shang, D., Yan, X., Zhao, B., & Zhang, X. (2021). Mitochondrial Behavior
in Axon Degeneration and Regeneration. In *Frontiers in Aging Neuroscience* (Vol. 13).
Frontiers Media S.A. https://doi.org/10.3389/fnagi.2021.650038

Włodarczyk, A., Wiglusz, M. S., & Cubała, W. J. (2018). Ketogenic diet for schizophrenia:
Nutritional approach to antipsychotic treatment. *Medical Hypotheses*, *118*, 74–77.
https://doi.org/10.1016/j.mehy.2018.06.022

Xu, M.-M., Wang, J., & Jun-Xia, X. (2017). Regulation of iron metabolism by hypoxia-inducible
factors 1 Hypoxia-inducible factors (HIFs). *Acta Physiologica Sinica*, *69*(5), 598–610.
https://doi.org/10.13294/j.aps.2017.0054

Yamashita, N., Aoki, R., Chen, S., Jitsuki-Takahashi, A., Ohura, S., Kamiya, H., & Goshima, Y.
(2016). Voltage-gated calcium and sodium channels mediate Sema3A retrograde signaling
that regulates dendritic development. *Brain Research*, *1631*, 127–136.
https://doi.org/10.1016/j.brainres.2015.11.034

Yamashita, N., Yamane, M., Suto, F., & Goshima, Y. (2016). TrkA mediates retrograde
semaphorin 3A signaling through plexin A4 to regulate dendritic branching. *Journal of Cell*
Science, *129*(9), 1802–1814. https://doi.org/10.1242/jcs.184580

Zhang, D. L., Ghosh, M. C., & Rouault, T. A. (2014). The physiological functions of iron
regulatory proteins in iron homeostasis - an update. In *Frontiers in Pharmacology: Vol. 5*
JUN. Frontiers Research Foundation. https://doi.org/10.3389/fphar.2014.00124

Zhang, M., Ergin, V., Lin, L., Stork, C., Chen, L., & Zheng, S. (2019). Axonogenesis Is
Coordinated by Neuron-Specific Alternative Splicing Programming and Splicing Regulator
PTBP2. *Neuron*, *101*(4), 690-706.e10. https://doi.org/10.1016/j.neuron.2019.01.022

Zolotukhin, S., Potter, M., Hauswirth, W. W., Guy, J., & Muzyczka, N. (1996). A "Humanized"
Green Fluorescent Protein cDNA Adapted for High-Level Expression in Mammalian Cells.
In *JOURNAL OF VIROLOGY* (Vol. 70, Issue 7). https://journals.asm.org/journal/jvi