Insights into the potential functions of nucleoside diphosphate kinases in *Waddlia chondrophila* encoded by two *ndk* paralogs

Giti Ghazi-Soltani, Carole Kebbi-Beghdadi, Simone E. Adams, Gilbert Greub[*]

Center for Research on Intracellular Bacteria, Institute of Microbiology, University Hospital Center and University of Lausanne, Lausanne, Switzerland

[*] Corresponding author:

Prof. Gilbert Greub: gilbert.greub@chuv.ch

ABSTRACT

The *Chlamydiota* phylum consists of obligate intracellular bacteria, including well-known pathogens and emerging environmental species, with diverse host ranges and metabolic capabilities. Among these bacteria, the *ndk* gene, which encodes nucleoside diphosphate kinase, is present in variable copy numbers. While most chlamydial species carry a single copy of the *ndk* gene, some species have two copies. In *W. chondrophila*, both *ndk* paralogs retain conserved kinase motifs but differ in subcellular localization, suggesting divergent functional roles. *WcNdk1* is confined to the inclusion and likely supports nucleotide metabolism, while *WcNdk2* localizes to the host nucleus, perinuclear space, and Golgi apparatus, implying involvement in host interaction. Azidothymidine (AZT), a known Ndk inhibitor, was found to impair the growth of *W. chondrophila*, possibly by targeting WcNdk2 due to its role in host interaction. Our findings shed light on the potential functions of *ndk* paralogs in *W. chondrophila* and support a role for WcNdk2 in host manipulation and pathogenesis. This research offers a starting point for deeper functional analyses as genetic manipulation techniques for this organism become more accessible.

Keywords: *Waddlia chondrophila*, Nucleoside diphosphate kinase (Ndk), AZT inhibition, Subcellular localization, Protein trafficking

INTRODUCTION

The *Chlamydiota* phylum encompasses several families of obligate intracellular bacteria, including five main families beyond the well-known *Chlamydiaceae*: *Parachlamydiaceae*, *Waddliaceae*, *Simkaniaceae*, *Rhabdochlamydiaceae*, and *Criblamydiaceae*. Members of these families share a chlamydia-like developmental cycle, which reflects their adaptation to intracellular environments and reliance on host cell machinery for survival.

W. chondrophila belongs to the family *Waddliaceae* and is known as an abortigenic agent in ruminants. It was first isolated from aborted bovine fetus[1,2] and serological studies showed a strong association between anti-*Waddlia* antibodies and bovine abortion[3]. In humans, *W. chondrophila* seropositivity is significantly associated with adverse pregnancy outcomes[4–6]. This pathogen was also detected in respiratory tract samples of patients with pneumonia and children with bronchiolitis[7,8].

The *W. chondrophila* developmental cycle is biphasic and is divided into three major phases[9,10]. In the early phase (0-8 hours post-infection, hpi) elementary bodies (EBs) enter in host cells and differentiate into reticulate bodies (RBs). This phase is followed by a proliferation phase (8-24 hpi) during which the number of RBs increases exponentially through binary fission. In the late phase (24-48 hpi) RBs revert into EBs which lyse the host cell to release infectious progeny. Under environmental or antibiotic stress, the developmental cycle is arrested and this leads into an enlarged, non-dividing RB-like form called Aberrant Body (AB)[11]. While *W. chondrophila* significantly impacts both animal and human health, the specific molecular and genomic pathways, particularly those involving developmental and regulatory proteins, remain largely unexplored. Functional characterization of proteins, such as nucleoside diphosphate kinase (Ndk), could help us to better understand the biology and pathogenicity of this pathogen and its adaptation inside host cells.

Ndk is a highly conserved enzyme that plays a central role in maintaining nucleotide pools. The maintenance of intracellular NTP pools is vital for the survival and proliferation of all living organisms. Ndk regulates nucleotide pools through its autophosphorylation and phosphotransferase activities. This ensures that NTP and dNTP levels remain balanced for optimal cellular function.

Beyond its housekeeping functions, Ndk is implicated in various cellular processes, including cell differentiation and development in eukaryotes and prokaryotes[12–15]. Additionally, Ndk is involved in the regulation of gene expression[16–18]. Due to its ability to bind ATP, Ndk is

involved in inhibiting extracellular ATP (eATP) and disrupting purinergic signaling in immune

and inflammatory responses[19–21].

We previously reported *W. chondrophila* transcriptomic data comparing genes expression in

RBs (24h pi) and EBs (72h pi)[22]. In this study *ndk* was among the genes differentially expressed

between the two developmental stages. Despite the well-established functions of Ndk across

various organisms, its specific function in the *Chlamydiae* remains uncharacterized. Based on

these observations, we hypothesized that this enzyme could play an important role in the

development and pathogenesis of *Chlamydia*-like organism. In this study, we characterize the

two Ndk proteins in *W. chondrophila* to better understand their potential role in the bacterium's

development and pathogenicity. This study provides an overall view of Ndk proteins in the

Chlamydiota phylum and offers a foundational characterization of *W. chondrophila* Ndks. Our

results pave the way for more detailed research once genetic tools become available for *W.*

chondrophila.

RESULTS

Evolution of the *ndk* operon in the *Chlamydiota* phylum

Similar to other organisms, bacteria of the *Chlamydiota* phylum possess at least one copy of

the *ndk* gene. A second copy (*ndk2*) is present in members of the *Parachlamydiaceae*,

Waddliaceae, and *Criblamydiaceae* families (Figure 1A). Interestingly, another *ndk2* forms an

operon with the ancestral *ndk1* gene. This operon in majority of species is further associated

with *pabA*, which encodes para-aminobenzoate synthase, and is involved in the folate

biosynthesis pathway. Comparisons of gene-based and species-based phylogenetic trees

(Figure 1B and 1C) suggest that the duplication of *ndk2* most likely originated from a single

duplication in the common ancestor of *Parachlamydiaceae*, *Waddliaceae*, and

Criblamydiaceae; however, a scenario of ancestral duplication with subsequent loss in

Simkaniaceae and *Chlamydiaceae* cannot be excluded, though it is less parsimonious.

High sequence and structural conservation of Ndk across species

The multiple sequence alignment shows a high degree of sequence conservation of Ndk

proteins from various bacterial and eukaryotic species (Figure 2A). Specifically, the His-Gly-

Ser-Asp (HGSD) motif, which is the enzyme's active site[23], is well conserved across all aligned

sequences. The highly conserved histidine residue inside this motif transiently receives the

phosphate group during autophosphorylation reaction and is, therefore, critical for the catalytic

function of Ndk. In addition to the HGSD motif, other conserved residues are observed in

regions associated with nucleotide binding and structural stability, many of which are enriched

in hydrophobic amino acids. These residues likely contribute to the proper folding and

enzymatic efficiency of Ndk.

W. chondrophila contains two copies of *ndk*: *wcndk1 (*locus *wcw_1543*) and *wcndk2* (locus

wcw_1545), with a sequence similarity of 54% (Figure 2B). The most striking difference

between the two *W. chondrophila ndk* paralogs is the presence of a predicted signal peptide of

17 amino acids in WcNdk2, which is absent in WcNdk1.

The sequence similarity matrix of *W. chondrophila* Ndk proteins, WcNdk1 and WcNdk2, and

their counterparts from other bacterial and eukaryotic species shows differing degrees of

conservation as shown in Figure 2B. WcNdk1 shares the highest sequence identity with *E. coli*

Ndk (EcNdk) and *Chlamydia trachomatis* Ndk (CtNdk) (67 % and 66%, respectively). In

contrast, WcNdk2 exhibits slightly lower sequence similarity across species, which may

indicate functional divergence of the two Ndks following gene duplication. Interestingly, both

W. chondrophila Ndk homologs display relatively high sequence identity (~50%) with human

Ndk (HsNdk), which suggests the conservation of core functional domains across vast

evolutionary distance. Sequence similarity matrix of Ndk proteins across different species is

shown in Figure S2.

WcNdk1 has 143 residues forming a polypeptide chain and adopting a very similar fold to Ndk

protein from other origins including human Ndk[24,25] (Figure 2C). One Ndk unit has α/β

domains comprising a four-stranded antiparallel β-sheet and two connecting α-helices. This

high degree of sequence and structural conservation between eukaryotes and the pathogens

may enable them to interfere with host cellular process by mimicking host Ndk.

Expression profile of *W. chondrophila* Ndks during infection

Temporal expression levels of gene and protein across different developmental stages may help

us understand their stage-specific roles and regulatory mechanisms. Since each of these phases

has unique biological features, we can infer a protein's function by mapping both transcript

and protein levels across these stages.

We quantified the expression profiles of *W. chondrophila ndks*, at both the transcript and

protein level during host cell infection (Figure 3). *wcndk1* and *wcndk2* transcripts were highly

abundant at early time points (3-8 hpi) as revealed by RT-qPCR, then the mRNA level declined

sharply to a minimum at 32 hpi, with a moderate surge toward the end of the cycle (48 hpi),

(Figure 3A). In parallel, no WcNdk1 or WcNdk2 protein could be detected by western blot of

the *W. chondrophila*-infected cells at 3 to 8 hpi (Figure 3B and 3C). Both proteins became

clearly detectable by 24 hpi, showing the highest observed signal intensity at this time point,

followed by a reduction to approximately 30% of that level by 48 hpi. To further assess protein

expression at early stages, we performed immunofluorescence microscopy. In contrast to

Western blot results, WcNdk1 and WcNdk2 signals were already detectable from 0 to 8 hpi,

colocalizing with intracellular *W. chondrophila* (Figure S1). Together, these results suggests

that both WcNdk paralogs are expressed throughout the developmental cycle, with

transcription peaking early. While protein expression begins early, overall accumulation or

detection sensitivity peaks around 24 hpi, followed by a decline toward the end of the cycle.

Subcellular localization of WcNdk1 and WcNdk2 in *C. trachomatis* heterologous
expression

While signal peptides are typically associated with protein secretion, previous studies have

demonstrated that Ndk can be secreted in certain bacterial species despite lacking a canonical

signal sequence[26–28]. Moreover, the secretion of both WcNdk1 and WcNdk2 into the host cell

cytoplasm is experimentally shown[29], suggesting that WcNdk2 signal peptide may not

contribute to secretion. The presence of this signal peptide suggests that WcNdk2 may be

translocated to a specific subcellular compartment and possibly confers a distinct biological

role to WcNdk2 compared to WcNdk1. To assess this possibility, we overexpressed a V5-

tagged version of WcNdk1 and WcNdk2 in *C. trachomatis* and studied their subcellular

localization by immunofluorescence and confocal microscopy (Figure 4). In this heterologous

expression assay, WcNdk1 was exclusively detected inside the inclusion, closely associated

with *C. trachomatis*, and no signal was observed in the host cell cytoplasm. Although this does

not rule out secretion of WcNdk1, it suggests limited accumulation or detection outside the

inclusion. In contrast, WcNdk2 was detected within both chlamydial inclusions and the host-

cell nucleus, indicating its trafficking into the nucleus. To further investigate the nuclear

localization of WcNdk2, we cloned the gene into the pBOMBL vector[30] and overexpressed the

protein in *C. trachomatis*. Upon infection of McCoy cells with the transformed bacteria,

inclusions were smaller, and bacterial growth was impaired compared to controls. Due to these

growth defects, we could not confirm the nuclear localization of WcNdk2 in this system.

Subcellular localization of WcNdk1 and WcNdk2 in transfected HEK293T cells

To further confirm the observed subcellular localization of WcNdk1 and WcNdk2 in another

heterologous system, we transiently expressed the V5-tagged WcNdk1 and WcNdk2 in

HEK293T cells and analyzed their distribution by immunofluorescence microscopy. WcNdk1

was found exclusively in the cytoplasm absent from cell compartments such as nucleus or

Golgi (Figure 5), whereas WcNdk2 showed two mutually exclusive distribution patterns. In

some cells the V5 signal was found throughout the nucleus, while in others, it was localized to

perinuclear-Golgi regions, with no nuclear signal (Figure 5).

The Ndk inhibitor, Azidothymidine (AZT), inhibits the growth of *W. chondrophila*

Since *W. chondrophila* is not currently amenable to genetic modifications, we employed

chemicals to inhibit Ndk function in this pathogen. AZT (3′-azido-3′-deoxythymidine), also

known as zidovudine, is a known Ndk inhibitor[14,31–35]. This pharmaceutical, primarily

approved for the treatment of human immunodeficiency virus (HIV), inhibits viral reverse

transcription through chain termination[36]. This compound is structurally a thymidine analogue,

where the 3'-hydroxyl group on the deoxyribose sugar is replaced by an azido group (N_3)

(Figure 6A).

The inhibition of Ndk from *Aspergillus flavus* by AZT is documented, where AZT forms a

strong hydrogen bond with key active site residues of Ndk (Arg-104, His-117 and Asp-120)

and inhibits its enzymatic activity[14]. Since these residues are highly conserved in Ndks of

diverse organisms, including WcNdk1 and WcNdk2, we hypothesized that AZT could also

serve as an effective inhibitor of Ndks in *W. chondrophila* (Figure 2A).

To test this hypothesis, we treated both uninfected and *W. chondrophila*-infected cells with

increasing concentrations of AZT following infection and measured cell lysis resulting from

bacterial proliferation using propidium iodide (PI) staining. PI binds to the DNA of lysed cells

and is used to assess plasma membrane integrity and cell death. In the absence of AZT or at

low concentrations (1 µg/ml) *W. chondrophila*-infected cultures exhibited a sharp increase in

cell mortality over time, reaching maximal levels by 144 hpi. Treatment with AZT at

concentrations above 25 µg/ml significantly suppressed infection-induced cell death. A delay

in cell mortality was observed with the intermediate concentration of 5 µg/ml (Figure 6B, right

panel). In contrast, in uninfected cells, cell lysis remained comparable across all AZT

concentrations, showing no significant difference from the no-AZT control. Therefore, the

slight increase in cell death observed over time is due to normal cell aging and culture

dynamics, and not due to the AZT effect (Figure 6B, left panel). These results indicate that

AZT did not exhibit adverse effects on mammalian host cells and provided a protective effect

against *W. chondrophila*-induced cell death in a dose-dependent manner.

Only early AZT treatment prevents *W. chondrophila*–mediated host cell lysis

To define the temporal window in which AZT exhibits its inhibitory effect on *W. chondrophila*

growth, *W. chondrophila*-infected cell monolayers were treated with 25 µg/ml AZT at 0, 3, 8,

24, 32, or 48 hpi and the propidium iodide (PI) uptake was monitored at 24, 48, 72, and 144

hpi (Figure 6C). When AZT was added at 0, 3, or 8 hpi, cell mortality remained minimum

(approximately 25% at 144 hpi), comparable to uninfected controls, indicating complete

protection against bacterial-induced lysis. In contrast, administering AZT at 24 hpi or later

failed to prevent cell death, resulting in maximum cell lysis, comparable to the no-AZT control.

This indicates that once RB replication is well established, AZT cannot reverse the course of

infection. Taken together, these results suggest that AZT must be applied before 24 hpi,

coinciding with maximum RB replication and inclusion expansion, to fully prevent *W.*

chondrophila–induced cytotoxicity, whereas later administration results in reduced

effectiveness.

AZT induced the production of aberrant bodies (ABs) in *W. chondrophila*

To investigate the cellular-level effects of AZT treatment on *W. chondrophila* infection, we

performed confocal microscopy on *W. chondrophila*-infected McCoy cells treated with 25

µg/ml AZT at 0 hpi. In untreated cultures, *W. chondrophila* underwent normal intracellular

development, leading to host cell lysis at the end of infection. At 8 hpi, *W. chondrophila* EBs

were observed attached to the host cell surface and internalized in both AZT-treated and

untreated cultures indicating that AZT does not interfere with bacterial attachment or entry

(Figure 6D). By 24 hpi, inclusions had formed in both conditions. However, in AZT-treated

cells, these inclusions failed to expand, and only a limited number of bacteria were detected

within them. This suggests that although EB-to-RB differentiation and initial rounds of

replication may occur, bacterial proliferation is subsequently arrested. In addition, the *W.*

chondrophila RBs appeared slightly enlarged in AZT-treated cultures, consistent with the

formation of aberrant bodies (ABs). These abnormal conditions persisted until 72 hpi,

indicating a disruption of the normal developmental cycle likely resulting in bacterial

persistence.

Effect of AZT on *W. chondrophila* growth is probably correlated with WcNdk2 activity

As mentioned earlier, *W. chondrophila* encodes two *ndk* paralogs, organized in single operon

with *pabA* positioned in between them. To investigate which Ndk copy mediates susceptibility

to AZT, and whether this susceptibility is linked to the presence of the *pabA* gene, we selected

representative *Chlamydiota* species with different *ndk* operon configurations and compared

their response to AZT treatment. The selected species include *C. trachomatis*, which carries a

single *ctndk* gene and lacks *pabA*; *Simkania negevensis,* which also has a single *snndk* gene

but retains *pabA* elsewhere in the genome; *Estrella lausannensis*, which harbors a two-gene

elndk1–elndk2 operon without *pabA*; and *W. chondrophila*, which contains the full *wcndk1–*
pabA–wcndk2 operon (Figure 7, bottom panel). To evaluate the effect of AZT on these species,
cell monolayers were infected, treated with 25 µg/ml at 0 hpi and fixed at 48 hpi before
immunostaining and observation under confocal microscopy. As shown in figure 7, *C.*
trachomatis and *S. negevensis* exhibited normal intracellular development in presence of AZT,
indicating no significant growth inhibition. In contrast, *E. lausannensis*, like *W. chondrophila*,
displayed enlarged intracellular structures resembling aberrant bodies. This suggest that AZT
inhibitory effect depends on the presence of *ndk2*, regardless of the presence of *ndk1* and *pabA*.

DISCUSSION

The diversity of *ndk* copy number within members of the *Chlamydiota* phylum reveals lineage-
specific adaptations with potential metabolic and pathogenic consequences. While most
Chlamydiales species possess a single *ndk* gene, a subset, including members of the
Parachlamydiaceae, *Waddliaceae*, and *Criblamydiaceae*, harbors a second copy (*ndk2*), likely
arising from a gene duplication event. The observed gene duplication may have provided a
selective advantage by providing increasing metabolism flexibility, which could, in part,
account for the comparatively faster growth of *W. chondrophila* and *E. lausannensis* in
mammalian cells relative to bacteria of *Chlamydiaceae*. Such an expansion of metabolic
capacities might also represent an adaptation facilitating persistence and replication within
free-living protists, suggesting that these lineages have evolved mechanisms enabling survival
across diverse host environments. This is in line with their larger genome size (>2 Mb),
expanded metabolic genes and their reduced dependence on host-derived metabolites. On the
other side, the absence of the second copy of the *ndk* gene in *Rhabdochlamydiaceae*,
Simkaniaceae and *Chlamydiaceae* coincides with their adaptations to more specific host niches
and reduced metabolic pathways. The *pabA* gene is also absent in all members of the
Chlamydiaceae family. In this family, enzymes from other biosynthetic pathways are recruited
to meet their folate requirements[37]. The mechanisms driving the evolution of this operon in
certain chlamydial families remains unknown. Further studies are needed to clarify the
operon's functional role and its evolutionary pathways in these families.

Both *W. chondrophila ndk* paralogs retain the universally conserved HGSD active-site motif
and surrounding hydrophobic residues essential for autophosphorylation and
phosphotransferase activities. This conservation suggests that both Ndk1 and Ndk2, despite

their apparent different subcellular localizations, exert their functions, at least partially, through phosphorylation mechanisms.

The temporal expression patterns of *wcndk1* and *wcndk2* provide insights into their potential biological roles and regulation during the *W. chondrophila* developmental cycle. The organization of *wcndk1* and *wcndk2* within a single operon explains their synchronized expression profiles. The apparent mid-cycle surge in Ndk proteins at 24 hpi coincides with the maximal bacterial replication, an increased demand for NTPs, and an increased host-manipulation activities that are essential for intracellular growth.

Despite sharing conserved kinase domains, our study showed significant differences in subcellular localization of WcNdk1 and WcNdk2. This divergence in their localizations suggests functional specializations and may indicate that the two proteins utilize divergent cellular trafficking mechanisms during *W. chondrophila* infection.

In *C. trachomatis* expression system, WcNdk1 was confined to the inclusion. This restricted distribution suggests that WcNdk1 may primarily serve bacterial intracellular functions within the inclusion. It is also possible that WcNdk1 is secreted transiently or at levels insufficient to be detected via immunofluorescence. In contrast, WcNdk2 exhibited nuclear localization both in *C. trachomatis* and HEK293T expression systems. This nuclear localization is consistent with previous studies on other pathogens, where Ndk was shown to bind host DNA and regulate gene expression through DNA cleavage[17,18]. Such putative nuclear activity suggests a potential role for WcNdk2 in reprogramming host transcriptional responses to favor bacterial survival. It would be of great interest to identify the specific host genes targeted by WcNdk2, as this could uncover previously unrecognized mechanisms by which *W. chondrophila* manipulates host cell function and could provide broader insights into its pathogenesis.

When expressed in HEK293T cells, WcNdk2 localized not only to the nucleus but also to the perinuclear-Golgi region. Bacterial Ndks, such as *Porphyromonas gingivalis* Ndk, are found in the perinuclear area of the host cells. This localization has been linked to the modulation of host purinergic signaling via $P2X_7$ receptors[28], implying that the perinuclear localization of WcNdk2 could be a biologically relevant phenomenon rather than nonspecific aggregation. Unlike the perinuclear accumulation of bacterial Ndks, Golgi localization represents a novel observation not previously reported for any bacterial species. One plausible explanation for an association of WcNdk2 with the Golgi is that, after translocation into the host cytosol, WcNdk2 enters the classical ER–Golgi secretory pathway and is packaged into Golgi-derived vesicles for extracellular release to hydrolyze host extracellular ATP (eATP) and subvert purinergic

signaling. Another possible explanation for the Golgi recruitment of WcNdk2 is that it might

play a role in modulating vesicle trafficking between the Golgi and the bacterial inclusion.

Although *W. chondrophila* does not redirect sphingomyelin transport from the Golgi to its

inclusion[38], it may still influence other aspects of vesicular trafficking, altering vesicle

formation or fusion processes, to benefit bacterial survival or replication.

AZT is a well-established inhibitor of Ndk[14,31–35]. In our study, AZT treatment of *W.*

chondrophila-infected cells protected them from bacteria-induced cell lysis and bacterial

growth was arrested in the replication phase. Our comparative data across *Chlamydiales*

species suggest that WcNdk2 is most likely the target for AZT. The inhibition of WcNdk2 by

AZT appears to interfere with a function that is essential for bacterial survival. Our results

demonstrated that WcNdk2 localizes to host cell compartments, including the nucleus,

perinuclear region, and Golgi apparatus in a heterologous expression system. Such host-

directed localization implies that secretion of WcNdk2 facilitates its accessibility to AZT and

may be critical for its effector functions within the host cell. Thus, bacterial growth arrest may

result from a disruption of Ndk2-dependent pathogenic mechanisms required for successful

replication, proliferation and host-cell manipulation. In contrast, WcNdk1, which

predominantly localizes within the inclusion, and is more likely involved in maintaining

bacterial nucleotide pools, appears less susceptible to AZT, possibly due to limited drug access

to the inclusion compartment. These findings collectively support a model in which the

inhibition of host-targeted, secreted WcNdk2, rather than inclusion-confined WcNdk1, is

central to AZT's antimicrobial activity against *W. chondrophila*.

Based on these observations, we propose a functional model for the two *W. chondrophila* Ndks,

as illustrated in Figure 8. Together, these findings highlight multifunctional nature of Ndk

proteins and shed light on the distinct roles of *W. chondrophila* Ndks. However, the lack of

genetic manipulation tools in *W. chondrophila* currently impedes direct investigation of Ndk

function via gene deletion or mutagenesis. Although localization studies using heterologous

expression systems suggest that WcNdk2 may interact with host cells, the possibility of

overexpression artifacts or mis-localization cannot be entirely ruled out. Additionally, although

AZT is a well-established Ndk inhibitor with no other known bacterial targets, direct in vitro

evidence confirming WcNdk2 as the AZT target is still missing. In vitro ATPase assays are

essential to validate this interaction. In the absence of such studies the proposed functions

remain speculative. The advancement of genetic tools would be essential to validate the

proposed roles of WcNdk1 and WcNdk2 in *W. chondrophila* metabolism and host interaction.

CONCLUSION

Given its diverse activities, Ndk has attracted interest across multiple disciplines, including microbiology, cell biology, and drug development. In this study, we introduced chlamydial Ndk as a multifunctional protein with roles extending beyond nucleotide metabolism, highlighting its potential involvement in host–pathogen interactions. These findings not only enhance our understanding of Ndk biology in the *Chlamydiota* phylum but also point to Ndk2 as a potential therapeutic target, opening new avenues for dissecting host manipulation strategies in obligate intracellular bacteria.

MATERIALS AND METHODS

Phylogenetic and structural analysis

Species-based and *ndk* gene phylogenies were retrieved from the *Chlamydia* Database[39]. Trees were trimmed using FigTree (v1.4.4) to retain only relevant *Chlamydiales* species. The Ndk protein sequences across different taxa were retrieved from the National Center for Biotechnology Information (NCBI). The Multiple sequence alignment of Ndk proteins across eukaryotes and prokaryotes was generated using the ClustalW tool on UGENE and the results were visualized with EPSript https://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi. The 3D structure of the protein was reconstructed using Phyre2[40] and visualized using Jmol v.16.2.15.

Cell culture and *W.chondrophila* infection

McCoy and HEK293T cells were maintained at 37°C in 5% CO_2 in DMEM GlutaMAX (Gibco), supplemented with 10% fetal bovine serum (FBS; Gibco). *W. chondrophila* was propagated in *Acanthamoeba castellanii* at 25°C. At the time of infection, lysed *W. chondrophila*-infected amoebae were filtered through a 5 mm syringe filter (Life science) to remove amoebal debris. The bacterial solution was then used to infect the host cells at a dilution of 1:1000, which was optimum for infection. To synchronize the infection, the infected cells were centrifuged at 1790 x g for 10 minutes. They were then incubated at 37°C in 5% CO_2 for 15 minutes. Following incubation, the inoculum was then replaced with fresh medium.

Purification of recombinant His-tagged WcNdk1 and WcNdk2 for antiserum production

The *E. coli* strain BL21 containing pET28a-*wcndk1-6xHis* or pET28a-*wcndk2-6xHis* was grown to an OD_{600} of 0.5. The culture was induced using 1 mM Isopropyl β-D-thiogalactopyranoside (IPTG, Applichem) and incubated at 37°C for 4 hours to express the recombinant protein. Bacterial pellets were resuspended in native lysis buffer (50 mM

NaH$_2$PO$_4$, 300 mM NaCl, and 10 mM imidazole, pH 8). The resuspended cells were lysed

using a combination of methods: three cycles of freeze (ethanol-dry ice bath) and thaw (at

37°C), followed by chemical lysis with 1 mg/ml lysozyme and sonication. The recombinant

protein was purified under native condition using Ni-NTA agarose beads (Qiagen) according

to the manufacturer's instructions. Briefly, bacterial lysates were incubated with Ni-NTA resin

for 2 hours at 4 °C with gentle rotation. The protein-bound resin was then loaded onto poly-

prep chromatography columns (Bio-Rad Laboratories), washed with wash buffer (50 mM Tris-

HCl, 300 mM NaCl, 20 mM imidazole, pH 8.0), and eluted with elution buffer containing

250 mM imidazole. The purified protein was dialyzed using Slide-A-Lyzer Dialysis Cassette

(Thermo scientific, 2,000 MWCO) overnight against PBS to remove imidazole. The

concentration of the protein was determined using Bradford's reagent with BSA (Bio-Rad

Laboratories) as a standard. Following protein purification, antisera were produced using

immunization services offered by Eurogentec SA (Seraing, Belgium).

Western Blot

McCoy cells were seeded at a density of 1 x 10^6 per T-25 flask one day before infection and

infected with *W. chondrophila* as described above. Infected cells were harvested by scraping at

the specified hpi and a fraction of the culture was saved for extraction and quantification of the

genomic DNA (see below). The cell suspension was pelleted at 1790 x g for 10 minutes, then

washed twice with PBS, and finally resuspended in 0.5 ml of 1x Laemmli sample buffer. An

equal volume of each lysate was resolved on 12% SDS-PAGE precast gels (Bio-Rad

Laboratories) and transferred onto an Amersham Protran nitrocellulose membrane (Cytiva).

The membrane was blocked in saturation buffer (10 mM Tris-Base, 150 mM NaCl, and 0.05%

Tween 20) containing 5% nonfat dried milk (AppliChem) for 2 hours at room temperature

(RT). Blots were probed overnight at 4°C with primary antibodies diluted in saturation buffer

with 0.5% non-fat dried milk. After three washes with saturation buffer containing 0.5% non-

fat dried milk, blots were incubated with secondary antibodies (horseradish peroxidase-

conjugated anti-rabbit IgG; Promega, or anti-mouse IgG; Bio-Rad Laboratories) for 1 hour at

Room temperature and processed using the Amersham ECL detection system (GE Healthcare).

Western blot bands intensities were quantified using EvolutionCapt edge x64 software and

normalized to the corresponding bacterial genome copy number. Graphs were generated using

GraphPad v. 10.4.1.

Extraction of genomic DNA and quantification of *W. chondrophila* genome copy number

Genomic DNA from collected samples was extracted using the Wizard SV Genomic DNA
purification system (Promega) according to the manufacturer's instructions. The extracted
DNA served as a template for qPCR with iTaq Universal Probes Supermix (Bio-Rad
Laboratories) to quantify the *W. chondrophila* bacterial population in 1 μl of culture. The 16S
copy numbers were determined using a standard curve, which was generated from serial
dilutions of a plasmid containing one copy of the 16S rRNA gene.

RNA extraction and cDNA synthesis

W. chondrophila-infected McCoy cell monolayers or uninfected control cells were harvested
at indicated hpi by scraping and centrifugation (5000 x g, 10 minutes). Following
centrifugation cells were lysed directly in TRIzol Reagent (Invitrogen). Total RNA was
extracted by chloroform separation and recovery of the aqueous phase ($12,000 \times g$, 15 min,
4 °C). RNA was precipitated with isopropanol, washed with 75 % ethanol, air-dried, and
resuspended in RNase-free water. To remove genomic DNA, samples were treated with
RNase-free DNase I (Invitrogen) following the manufacturer's instructions. cDNA was
synthesized using the GoScript Reverse Transcription System (Promega) with random primers.
Reverse transcription was performed at 42°C for 60 minutes, followed by enzyme inactivation
at 70°C for 15 minutes. All cDNAs were diluted 1:5 prior to qPCR.

RT-qPCR

The qPCR reactions were set up in a final volume of 20 μl containing 4 μl of cDNA template,
the appropriate concentration of each primer, and 1X iTaq Universal SYBR Green Supermix
(Bio-Rad Laboratories). All samples were run in duplicates, and no-template controls were
included for each primer pair to assess non-specific amplification. The fluorescent reporter
signal was normalized against the internal reference dye (ROX) signal. qPCR was carried out
on QuantStudio 3 (Applied Biosystems) using the following thermal program. A single cycle
of DNA polymerase activation for 3 min at 95 °C, followed by 45 amplification cycles of 15 s
at 95 °C (denaturing step) and 1 min at 60 °C (annealing and extension step). Gene expression
data were normalized to *W. chondrophila* 16 rRNA gene, which served as the internal reference
gene. Quantitative data from qRT-PCR experiments were collected from at least three
independent replicates. The primers used for RT-qPCR are listed in Table S1.

Application of AZT on *W. chondrophila* culture and cell death assessment

McCoy cells were seeded at a density of 1×10^4 cells per well in 96-well plates (Corning).
Cells were infected with *W. chondrophila* or left uninfected as a control. To monitor cell lysis,

Propidium Iodide (PI, 7 µg/mL, Sigma-Aldrich) was added to the growth medium. AZT was

purchased from TOCRIS (Cat. No.:4150) and was added to the wells at concentrations ranging

from 0 to 250 µg/mL in six technical replicates per condition. Depending on the experimental

design, AZT was added at the time of infection (0 hpi) or at later time points (3, 8, 24, 32, and

48 hpi). PI fluorescence was measured at specified time points post-infection using a FLUOstar

Omega plate reader (BMG LABTECH; excitation: 540 nm; emission: 640 nm). To define

100% cell lysis, 0.1% Triton X-100 was added to control wells prior to PI reading.

Immunofluorescence assay

McCoy cells were cultured on glass coverslips placed in 24-well-plates and infected with

various bacterial species. At the indicated time points, cells were fixed with ice-cold methanol

for 5 minutes at 4°C, followed by three washes with phosphate-buffered saline (PBS). Fixed

cells were permeabilized and blocked for 30 minutes using a blocking solution containing

0.1% saponin, 10% fetal bovine serum (FBS), and 0.04% sodium azide in PBS. Blocked cells

were incubated with primary antibodies for 2 hours at RT. After three washing steps with

blocking solution, cells were incubated for 1 hour with Alexa Fluor 488-conjugated secondary

antibodies (Invitrogen, 1:1000 dilution) to stain bacteria. DAPI (Invitrogen, 1:3000 dilution)

and Texas Red-conjugated Concanavalin A (Invitrogen) were used to stain nuclei and

carbohydrates, respectively. Cells were then washed three times with PBS and briefly rinsed

with purified water before mounting. Coverslips were mounted on glass slides with Mowiol 4-

88 and stored in the dark until imaging using Zeiss LSM 900 confocal laser scanning

microscope (Carl Zeiss Microscopy GmbH).

pGL2 plasmid construction

The genes *wcndk1* and *wcndk2* were amplified from *W. chondrophila* genomic DNA using

primers with 5′ overhangs compatible with the pGL2 backbone. All PCR products included a

C-terminal V5 epitope tag to facilitate detection. PCR products were purified using the

QIAquick PCR Purification Kit (Qiagen). The native *C. trachomatis*–derived pGL2 vector

(11,786 bp; β-lactamase selection marker) was linearized by Age I digestion (NEB) and

recovered from agarose gels with the QIAquick Gel Extraction Kit (Qiagen). Purified inserts

and linearized vector were assembled via In-fusion cloning using 5X In-fusion Snap assembly

Master Mix (Takara). Assemblies were incubated at 50 °C for 15 min. Transformations were

performed using *E. coli* dam⁻ dcm⁻ cells. Transformants were selected on LBA supplemented

with Carbenicillin (50 µg/mL). Positive clones were screened by colony PCR, and successful

insertions were confirmed by Sanger sequencing using both vector- and insert-specific primers.

C. trachomatis transformation

C. trachomatis transformation

Transformation of *C. trachomatis* was performed with minor modifications to previously

described methods[41]. 1 x 10^6 McCoy cells were seeded in 6-well plates and cultured overnight.

2.5 x 10^6 plasmid-free *C. trachomatis* serovar L2 was incubated with 2 µg of sequence-verified

plasmid DNA for 30 minutes. The quantity of plasmid-free *C. trachomatis* serovar L2 (EBs),

resuspended in 300µl Tris-CaCl$_2$ buffer (10 mM Tris, 50 mM CaCl2, pH 7.4), and incubated

with 2 µg of sequence-verified plasmid DNA for 30 minutes. 1 mL Hank's balanced salt

solution (HBSS, Gibco) was then added to each reaction. This mixture was added to McCoy

cells in a 6-well plate after removing the medium. The infection was carried out by

centrifugation at 400 × g for 15 minutes at room temperature and incubation at 37°C for 15

minutes. Then, the inoculum was removed, and cells were incubated with 2 mL of DMEM

medium supplemented with 10% FBS and 1 µg/mL cycloheximide for 8 hours at 37°C and 5%

CO_2. After this incubation, the medium was replaced with DMEM containing 10% FBS, 1

µg/mL cycloheximide, and penicillin (0.6 mg/mL) to select for transformed bacteria. Cells

were passaged every 48 hours, and the development of fluorescent inclusions were monitored

until they were clearly observed and established. The transformed *C. trachomatis* was

harvested and titrated by IFU assay and stored at -80°C. For experiments requiring induction,

50 ng/ml anhydrotetracycline (aTc, Sigma-Aldrich) was added at the time of infection,

whereas, in control conditions, the inducer was excluded.

Gateway cloning for mammalian expression in pDEST47

Gateway cloning for mammalian expression in pDEST47

For generation of pDEST47 expression plasmids, attB-flanked primers were used to amplify

the C-terminal V5-tagged version of *wcndk1* and *wcndk2*. Entry clones were created by BP

recombination (Gateway BP Clonase, Invitrogen), following the manufacturer's protocol. The

resulting pDONR201-*ndk-V5* entry clones were selected on appropriate antibiotic. Destination

vectors were generated via LR recombination of the entry clones with pDEST47 using LR

Clonase (Gateway, Invitrogen), according to the supplier's instructions. Final constructs were

transformed and validated as described above, with plasmid DNA prepared from overnight

cultures using the GeneJET plasmid Miniprep kit (Thermo Scientific).

Transfection in HEK293T cells

Transfection in HEK293T cells

Glass coverslips in 24-well plates were coated with poly-L-lysine (Sigma-Aldrich) at a final

concentration of 100 µg/ml, incubated for 30 minutes at room temperature, washed three times

with PBS, and air-dried for 3 hours. HEK293T cells were then seeded on coated coverslips at

2×10^5 cells per well and incubated overnight at 37 °C, 5% CO_2. HEK293T cells were

transfected 24 hours post-seeding with the pDEST47 plasmids using the lipofectamine 3000

transfection kit (Invitrogen). DNA (3 µg) was diluted in 150 µL serum-free DMEM with 6 µL

P3000 reagent. Separately, 4.5 µL Lipofectamine 3000 (Invitrogen) was diluted in 150 µL

DMEM. The two mixtures were combined, incubated for 15 minutes at room temperature, and

50 µL was added dropwise per well. Cells were fixed 24 hours post-transfection with ice-cold

methanol for downstream immunofluorescence analysis.

Figure1: Phylogeny of *Chlamydiota* phylum and the *ndk* gene operon evolution. (A) Species phylogeny based on 32 single-copy orthologous genes. (B) Simplified representation of the species phylogeny highlighting major clades. (C) Phylogeny based on the *ndk* gene, showing its evolutionary relationship among *Chlamydiales* species.

Figure 2: Ndk sequence and structure conservation. (A) Multiple sequence alignment of Ndk proteins across eukaryotes and prokaryotes, generated using ClustalW tool on UGENE and visualized with https://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi. Conserved residues are highlighted in red with the highly conserved H residue, essential for catalytic activity, marked with an asterisk. Af: *Aspergillus flavus*, An: *A. niger*, Hs: *Homo sapiens*, Mt: *Mycobacterium tuberculosis*, Ec: *Escherichia coli,* Ct: *C. trachomatis*, Sn: *Simkania negevensis*, Rp: *Rhabdochlamydia porcellionis,* Wc: *W. chondrophila*, El: *Estrella lausannensis,* Cs: *Criblamydia sequanensis,* Pa: *Parachlamydia acanthamoebae,* Nsp: *Neochlamydia sp.*, Pae: *Pseudomonas aeruginosa*, Ho: *Halofilum ochraceum*, Lp: *Legionella pneumophila* (B) Amino acid sequence similarity matrix comparing Ndk proteins from *W. chondrophila* (WcNdk1 and WcNdk2), *Chlamydia trachomatis* (CtNdk), *Escherichia coli* (EcNdk), *Mycobacterium tuberculosis* (MtNdk), and human Ndk (HsNdk). (C) Predicted 3D structure of WcNdk1, illustrating its conserved α/β fold characteristic of the Ndk family. 3D structure reconstructed using Phyre2 and visualized using Jmol v.16.2.15. The H177, highlighted in red, is a critical histidine residue and positioned within the active site.

Figure 3: Temporal expression analysis of *ndk* paralogs in *W. chondrophila* during infection. (A) Quantitative RT-PCR analysis of *ndk1* and *ndk2* transcript levels at different hours post-infection (hpi), normalized to 16S rRNA. Transcript abundance is expressed relative to the 24 hpi time point, which was set as the reference (1.0). (B) Relative protein levels of WcNdk1 and WcNdk2 quantified by Western blot, normalized to bacterial genome copy number. Protein expression is shown relative to 24 hpi, set as 100%. (C) Representative Western blot images showing WcNdk1 and WcNdk2 expression across the infection time course. Data are shown as mean ± SD of 3 independent experiments.

Figure 4: Subcellular localization of WcNdk1 and WcNdk2 overexpressed in *C. trachomatis*. *C. trachomatis* was transformed with plasmids expressing either WcNdk1 or WcNdk2 tagged with a V5 epitope. Infected McCoy cells were fixed at 24 hpi. GFP (green) indicates transformed *C. trachomatis*, anti-V5 (red) detects WcNdk proteins, and DAPI (blue) stains host cell nuclei and bacterial DNA. White arrows point to V5 signal within host nuclei.

Figure 5: Subcellular distribution of WcNdk1 and WcNdk2 in HEK293T Cells. HEK293T cells were transiently transfected with C-terminal V5-tagged WcNdk1 and WcNdk2 and stained with the Golgi marker GM130 (green), anti-V5 antibody (red), and DNA (DAPI, blue). High-magnification insets highlight the co-localization of WcNdk2 (red) with the Golgi (green) or the perinuclear region.

Figure 6: AZT inhibits *W. chondrophila* growth in infected host cells. (A) Chemical structure of azidothymidine (AZT). (B) AZT treatment of uninfected host cells, serving as a negative control (left panel) and treatment of infected host cells at concentrations ≥25 μg/mL (right panel). (C) Effect of AZT addition on *W. chondrophila*-infected cultures at various time points after infection. (D) Confocal microscopy of *W. chondrophila*-infected cells treated or not treated with AZT. In untreated conditions, *W. chondrophila* formed typical intracellular inclusions, whereas AZT treatment resulted in enlarged, aberrant structures. Bacteria are labeled using anti-*W. chondrophila* antibody (green); host cell nuclei are stained with DAPI (blue); and the host structures are marked with Concanavalin A (red).

Figure 7: AZT sensitivity is restricted to *Chlamydiales* species encoding *ndk2*. Representative confocal micrographs of McCoy cells infected with *C. trachomatis*, *S. negevensis*, *W. chondrophila* or *E. lausannensis*, treated or not treated with 25 µg/ml AZT. The infected cells were fixed and stained 48 hpi. The lower panel shows the *ndk* operon arrangement: *C. trachomatis* and *S. negevensis*, each carries a single *ndk* gene. *S. negevensis* retains *pabA* elsewhere in the genome. *W. chondrophila* carries the two *ndk* genes with a *pabA* in the middle. *E. lausannensis* encodes a two-gene *ndk1–ndk2* operon but lacks *pabA*. Bacteria are stained in green using species-specific antibodies, host cytoplasm is labeled with Concanavalin A (red), and nuclei are stained with DAPI (blue). *ctndk*: *C. trachomatis ndk1*, *snndk1*: *S. negevensis ndk1*, *elndk1*: *E. lausannensis ndk1*, *elndk2*: *E. lausannensis ndk2*.

Figure 8: Proposed model for WcNdk1 and WcNdk2 functions in *W. chondrophila*. This schematic illustrates the distinct subcellular localizations and putative roles of WcNdk1 (green) and WcNdk2 (red), during infection of a host cell. A: WcNdk1 is predominantly retained within the bacterial inclusion, where it sustains basic nucleotide metabolism and "housekeeping" functions. B: A fraction of WcNdk1 is secreted across the inclusion membrane into the host cytosol, however, the specific functions of cytosolic WcNdk1 remain undefined. C: WcNdk2 is observed accumulating within the host nucleus. Its import into the nucleus could occur via one or more of the following mechanisms: a deacetylation event allowing passive nuclear import; free diffusion through nuclear pores owing to its small molecular size; or interaction with host proteins that escort WcNdk2 into the nucleus. Inside host nucleus, WcNdk2 could bind directly to DNA leading to up- or downregulation of host genes involved in immunity or apoptosis. The precise gene targets remain unknown. D: WcNdk2 traffics through Golgi/ER-derived vesicles and perinuclear region. This localization may be related to the packaging of WcNdk2 into vesicles destined for secretion to subvert host purinergic signaling through hydrolyzes of eATP into ADP. E: The association of WcNdk2 with Golgi apparatus implies its involvement in vesicle trafficking and acquisition of host-derived metabolites necessary for expansion of the inclusion and bacterial survival.

FUNDING

This work was supported by the Swiss National Science Foundation (SNSF) [Grant No. 197768].

REFERENCES

1. Dilbeck PM, Evermann JF, Crawford TB, et al. Isolation of a previously undescribed rickettsia from an aborted bovine fetus. *J Clin Microbiol*. 1990;28(4):814-816

2. Henning K, Schares G, Granzow H, et al. *Neospora caninum* and *Waddlia chondrophila* strain 2032/99 in a septic stillborn calf. *Veterinary Microbiology*. 2002;85(3):285-292. doi:10.1016/S0378-1135(01)00510-7

3. Dilbeck-Robertson P, McAllister MM, Bradway D, Evermann JF. Results of a new serologic test suggest an association of *Waddlia chondrophila* with bovine abortion. *J Vet Diagn Invest*. 2003;15(6):568-569. doi:10.1177/104063870301500609

4. Baud D, Thomas V, Arafa A, Regan L, Greub G. *Waddlia chondrophila*, a Potential Agent of Human Fetal Death. *Emerg Infect Dis*. 2007;13(8):1239-1243. doi:10.3201/eid1308.070315

5. Baud D, Goy G, Osterheld MC, Borel N, Vial Y, pospischil A, Greub G. *Waddlia chondrophila*: from bovine abortion to human miscarriage. *Clin Infect Dis*. 2011;52(12):1469-1471. doi:10.1093/cid/cir205

6. Baud D, Regan L, Greub G. Emerging role of Chlamydia and Chlamydia-like organisms in adverse pregnancy outcomes. *Curr Opin Infect Dis*. 2008;21(1):70-76. doi:10.1097/QCO.0b013e3282f3e6a5

7. Haider S, Collingro A, Walochnik J, Wagner M, Horn M. Chlamydia-like bacteria in respiratory samples of community-acquired pneumonia patients. *FEMS Microbiol Lett*. 2008;281(2):198-202. doi:10.1111/j.1574-6968.2008.01099.x

8. Goy G, Croxatto A, Posfay-Barbe KM, Gervaix A, Greub G. Development of a real-time PCR for the specific detection of *Waddlia chondrophila* in clinical samples. *Eur J Clin Microbiol Infect Dis*. 2009;28(12):1483-1486. doi:10.1007/s10096-009-0804-7

9. de Barsy M, Greub G. *Waddlia chondrophila*: from biology to pathogenicity. *Microbes and Infection*. 2013;15(14):1033-1041. doi:10.1016/j.micinf.2013.09.010

10. K Kebbi-Beghdadi C, Cisse O, Greub G. Permissivity of Vero cells, human pneumocytes and human endometrial cells to *Waddlia chondrophila*. *Microbes and Infection*. 2011;13(6):566-574. doi:10.1016/j.micinf.2011.01.020

11. Scherler A, Jacquier N, Kebbi-Beghdadi C, Greub G. Diverse Stress-Inducing Treatments cause
Distinct Aberrant Body Morphologies in the Chlamydia-Related Bacterium, *Waddlia*
chondrophila. *Microorganisms*. 2020;8(1):89. doi:10.3390/microorganisms8010089

12. Timmons L, Shearn A. Role of AWD/Nucleoside Diphosphate Kinase in Drosophila Development.
J Bioenerg Biomembr. 2000;32(3):293-300. doi:10.1023/A:1005545214937

13. Masoudi N, Fancsalszky L, Pourkarimi E, et al. The NM23-H1/H2 homolog NDK-1 is required for
full activation of Ras signaling in C. elegans. *Development*. 2013;140(16):3486-3495.
doi:10.1242/dev.094011

14. Wang Y, Wang S, Nie X, et al. Molecular and structural basis of nucleoside diphosphate kinase–
mediated regulation of spore and sclerotia development in the fungus Aspergillus flavus. *J Biol*
Chem. 2019;294(33):12415-12431. doi:10.1074/jbc.RA119.007505

15. Muñoz-Dorado J, Inouye S, Inouye M. Nucleoside diphosphate kinase from Myxococcus xanthus.
II. Biochemical characterization. *Journal of Biological Chemistry*. 1990;265(5):2707-2712.
doi:10.1016/S0021-9258(19)39859-X

16. Postel EH, Berberich SJ, Rooney JW, Kaetzel DM. Human NM23/Nucleoside Diphosphate Kinase
Regulates Gene Expression through DNA Binding to Nuclease-Hypersensitive Transcriptional
Elements. Published online 2000:8.

17. Levit MN, Abramczyk BM, Stock JB, Postel EH. Interactions between Escherichia coli nucleoside-
diphosphate kinase and DNA. *J Biol Chem*. 2002;277(7):5163-5167. doi:10.1074/jbc.M11117020

18. Kumar P, Verma A, Saini AK, et al. Nucleoside diphosphate kinase from Mycobacterium
tuberculosis cleaves single strand DNA within the human c-myc promoter in an enzyme-catalyzed
reaction. *Nucleic Acids Res*. 2005;33(8):2707-2714. doi:10.1093/nar/gki568

19. Coutinho-Silva R, Ojcius DM. Role of extracellular nucleotides in the immune response against
intracellular bacteria and protozoan parasites. *Microbes Infect*. 2012;14(14):1271-1277.
doi:10.1016/j.micinf.2012.05.009

20. Kolli BK, Kostal J, Zaborina O, Chakrabarty AM, Chang KP. *Leishmania*-released nucleoside
diphosphate kinase prevents ATP-mediated cytolysis of macrophages. *Molecular and Biochemical*
Parasitology. 2008;158(2):163-175. doi:10.1016/j.molbiopara.2007.12.010

21. Yilmaz Ö, Yao L, Maeda K, et al. ATP scavenging by the intracellular pathogen Porphyromonas
gingivalis inhibits P2X7-mediated host-cell apoptosis. *Cellular Microbiology*. 2008;10(4):863-
875. doi:10.1111/j.1462-5822.2007.01089.x

22. Ardissone S, Scherler A, Pillonel T, Martin V, Kebbi-Beghdadi C, Greub G. Transcriptional
Landscape of *Waddlia chondrophila* Aberrant Bodies Induced by Iron Starvation. *Microorganisms*.
2020;8(12):1848. doi:10.3390/microorganisms8121848

23. Arumugam M, Ajitkumar P. Histidine 117 in the His-Gly-Ser-Asp motif is Required for the
Biochemical Activities of Nucleoside Diphosphate Kinase of Mycobacterium smegmatis. *Open*
Biochem J. 2012;6:71-77. doi:10.2174/1874091X01206010071

24. Singh S, Yadav PK, Singh AK. In-silico structural characterization and phylogenetic analysis of
575   Nucleoside diphosphate kinase: A novel antiapoptotic protein of Porphyromonas gingivalis.
576   *Journal of Cellular Biochemistry*. 2023;124(4):545-556. doi:10.1002/jcb.30389

577   25. Janin J, Dumas C, Moréra S, et al. Three-Dimensional Structure of Nucleoside Diphosphate Kinase.
578   *J Bioenerg Biomembr*. 2000;32(3):215-225. doi:10.1023/A:1005528811303

579   26. Kamath S, Chen ML, Chakrabarty AM. Secretion of Nucleoside Diphosphate Kinase by Mucoid
580   Pseudomonas aeruginosa 8821: Involvement of a Carboxy-Terminal Motif in Secretion. *J*
581   *Bacteriol*. 2000;182(13):3826-3831.

582   27. Neeld D, Jin Y, Bichsel C, et al. Pseudomonas aeruginosa injects NDK into host cells through a
583   type III secretion system. *Microbiology (Reading)*. 2014;160(Pt 7):1417-1426.
584   doi:10.1099/mic.0.078139-0

585   28. Atanasova K, Lee J, Roberts J, Lee K, Ojcius DM, Yilmaz Ö. Nucleoside-Diphosphate-Kinase of
586   P. gingivalis is Secreted from Epithelial Cells In the Absence of a Leader Sequence Through a
587   Pannexin-1 Interactome. *Scientific Reports*. 2016;6(1). doi:10.1038/srep37643

588   29. Kebbi-Beghdadi C, Pilloux L, Martin V, Greub G. Eukaryotic Cell Permeabilisation to Identify
589   New Putative Chlamydial Type III Secretion System Effectors Secreted within Host Cell
590   Cytoplasm. *Microorganisms*. 2020;8(3):361. doi:10.3390/microorganisms8030361

591   30. Ouellette, S. P., Blay, E. A., Hatch, N. D. & Fisher-Marvin, L. A. CRISPR Interference To Inducibly
592   Repress Gene Expression in Chlamydia trachomatis. Infect Immun 89, e00108-21. doi:
593   10.1128/IAI.00108-21

594   31. Valenti D, Barile M, Quagliariello E, Passarella S. Inhibition of nucleoside diphosphate kinase in
595   rat liver mitochondria by added 3′-azido-3′-deoxythymidine. *FEBS Letters*. 1999;444(2-3):291-
596   295. doi:10.1016/S0014-5793(99)00071-X

597   32. Schaertl S, Konrad M, Geeves MA. Substrate Specificity of Human Nucleoside-diphosphate
598   Kinase Revealed by Transient Kinetic Analysis. *Journal of Biological Chemistry*.
599   1998;273(10):5662-5669. doi:10.1074/jbc.273.10.5662

600   33. Gallois-Montbrun S, Schneider B, Chen Y, et al. Improving Nucleoside Diphosphate Kinase for
601   Antiviral Nucleotide Analogs Activation. *Journal of Biological Chemistry*. 2002;277(42):39953-
602   39959. doi:10.1074/jbc.M206360200

603   34. Xu Y, Sellam O, Moréra S, et al. X-ray analysis of azido-thymidine diphosphate binding to
604   nucleoside diphosphate kinase. *Proceedings of the National Academy of Sciences*.
605   1997;94(14):7162-7165. doi:10.1073/pnas.94.14.7162

606   35. Wambaugh MA, Shakya VPS, Lewis AJ, Mulvey MA, Brown JCS. High-throughput identification
607   and rational design of synergistic small-molecule pairs for combating and bypassing antibiotic
608   resistance. *PLOS Biology*. 2017;15(6):e2001644. doi:10.1371/journal.pbio.2001644

609   36. Furman PA, Barry DW. Spectrum of antiviral activity and mechanism of action of zidovudine. An
610   overview. *Am J Med*. 1988;85(2A):176-181.

37. Adams NE, Thiaville JJ, Proestos J, et al. Promiscuous and Adaptable Enzymes Fill "Holes" in the
Tetrahydrofolate Pathway in Chlamydia Species. *mBio*. 2014;5(4):e01378-14.
doi:10.1128/mBio.01378-14

38. Dille S, Kleinschnitz EM, Kontchou CW, Nölke T, Häcker G. In contrast to *Chlamydia trachomatis*,
Waddlia chondrophila grows in human cells without inhibiting apoptosis, fragmenting the Golgi
apparatus, or diverting post-Golgi sphingomyelin transport. *Infect Immun*. 2015;83(8):3268-3280.
doi:10.1128/IAI.00322-15

39. Pillonel T, Tagini F, Bertelli C, Greub G. ChlamDB: a comparative genomics database of the
phylum Chlamydiae and other members of the Planctomycetes-Verrucomicrobiae-Chlamydiae
superphylum. *Nucleic Acids Research*. 2020;48(D1):D526-D534. doi:10.1093/nar/gkz924

40. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. The Phyre2 web portal for protein
modeling, prediction and analysis. *Nat Protoc*. 2015;10(6):845-858. doi:10.1038/nprot.2015.053

41. Mueller KE, Wolf K, Fields KA. *Chlamydia trachomatis* transformation and allelic exchange
mutagenesis. *Curr Protoc Microbiol*. 2017;45:11A.3.1-11A.3.15. doi:10.1002/cpmc.31

625