3' Nucleotide Asymmetry Directs miRNA Strand Selection

Jeffrey C. Medley[1], Sumire Kurosu Moriya[1], Huiwu Ouyang[2], Heather Crawshaw[1], Sarah Y. Zhang[1], Ganesh Panzade[1,3], Will J. Sydzyik[1], Joel T. Sydzyik[1,4], Mira Bhandari[1,5], Christopher M. Hammell[2] and Anna Zinovyeva[1]

[1] Division of Biology, Kansas State University. Manhattan, KS.
[2] Cold Spring Harbor Laboratory. Cold Spring Harbor, NY.
[3] Laboratory of Human Retrovirology and Immunoinformatics, Frederick National Laboratory for Cancer Research, Frederick, MD.
[4] University of Kansas School of Medicine, Kansas City, KS.
[5] Department of Molecular Genetics and Cell Biology, University of Chicago. Chicago, IL.

Abstract

During miRNA biogenesis, double-stranded precursors are processed into duplexes comprising functionally distinct strands. The guide strand directs the miRNA-induced silencing complex (miRISC) to complementary mRNA targets, while the passenger strand is degraded. As each strand has different molecular targets, strand selection effectively determines the target repertoire of the miRISC. Previous studies suggest that 5' nucleotide identity and thermodynamic asymmetries determine miRNA strand choice in vitro. However, these guidelines cannot fully explain the observed strand preference of all miRNAs in vivo. Here, we demonstrate a conserved role for 3' nucleotide asymmetry in facilitating miRNA strand selection in *Caenorhabditis elegans* and human HEK293T cells. Our data show that a favorable 3' nucleotide on miRNA passenger strands can promote selection of the opposing guide strand. We propose that terminal nucleotide asymmetries on both strands of miRNA duplexes promote accurate strand selection in vivo.

29 **Main**
30
The biogenesis of miRNAs is integral to their biological activity, as the precise sequence of
mature miRNAs must be preserved to maintain sequence-specific interactions with mRNA
targets[1,2]. Primary miRNA (pri-miRNA) transcripts are genomically encoded and processed
within the nucleus by the microprocessor complex to produce stem-loop miRNA precursors
(pre-miRNA)[3–6]. Following nuclear export, Dicer cleaves pre-miRNAs to generate partially
double-stranded ~22 nucleotide duplexes that typically contain two-nucleotide overhangs at
their 3' ends[7–12]. The miRNA duplex is subsequently loaded into an Argonaute protein, which
selectively anchors a single, mature strand to initiate formation of the miRNA induced silencing
complex (miRISC)[13,14]. Although the strands originating from either the 5' (5p) or 3' (3p) arms of
the pre-miRNA are competent for Argonaute loading[15–17], asymmetric strand selection usually
leads to a dominant guide strand and a less abundant passenger strand that is rapidly degraded
after release from Argonaute[18]. For some miRNAs, strand selection occurs in a more
symmetrical fashion where both strands appear to function as mature miRNAs[19,20]. Differences
in miRNA strand usage have also been observed in different cellular, tissular and
developmental contexts, suggesting that strand selection is a regulated process[19,21–24]. As
miRNA-dependent gene regulation is sequence dependent, alternative miRNA strand choice is
expected to lead to broad changes in target selection. Consistently, aberrant miRNA strand
selection has also been reported in several human diseases including cancers, although the
mechanisms leading to alternative strand selection in human disease are not fully understood[18].
50
Argonaute plays a direct role in strand selection and favors loading the duplex end containing a
favorable 5' nucleotide (U>A>C>G) and lower thermodynamic stability in vitro[16,17,25,26]. The 5'
nucleotide of the guide strand is embedded within a pocket in the MID domain of Argonaute,
which has a strong preference for uracil[13,14,25]. Several residues within the Argonaute MID and
PIWI domains directly contact the phosphate backbone of the guide strand[13,14] and may serve
as a sensor for thermodynamic asymmetry[17]. The twin-drive model postulates that miRNA
strand selection can be predicted by modeling the relative effects of 5' nucleotide identity and
thermodynamic asymmetry[17]. However, this model fails to explain strand selection of ~25% of
miRNAs that have unfavorable 5' nucleotides and more stable 5' ends[18], suggesting that
additional asymmetries influence miRNA strand selection in vivo.
61
Here, we used genome editing to alter endogenous *C. elegans* miRNAs and evaluated the
relative contributions of 5' nucleotide identity and thermodynamic asymmetry in vivo. Consistent
with previous findings, we show that a 5' uracil is favorable for stand choice, and that unpaired
5' nucleotides are strongly selected regardless of nucleotide identity. We were surprised to find
that duplexes with symmetrical 5' nucleotides and thermodynamic end stabilities underwent
asymmetric strand selection, suggesting that additional features outside of the 5' end contribute
to strand choice. We identified a strong bias for cytosine at the 3' end of passenger strands in *C.*
elegans. Mutating the 3' nucleotide of symmetrical duplexes was sufficient to reverse strand
selection, suggesting that 3' nucleotide identity influences miRNA strand asymmetry in vivo. In
duplexes with asymmetric 5' ends, 3' nucleotide asymmetry had a lesser effect on strand
choice, indicating that various nucleotide asymmetries coordinately drive miRNA strand
selection. By expressing exogenous miRNA variants in human HEK293T cells, we show that 3'
nucleotide asymmetry plays a conserved role in miRNA strand selection, although a 3' cytosine
on passenger strands was less preferred in human cells. Collectively, our results support a
model where nucleotide asymmetries on both strands of miRNA duplexes promote asymmetric
strand selection.
78
79 **Results**

80
Characterization of *C. elegans* miRNA duplexes

82
To better understand how 5' nucleotide identity and thermodynamic asymmetries are linked to
miRNA strand selection, we profiled the duplex characteristics of *C. elegans* miRNAs
(Supplementary Table 1). The preferred guide strand is expected to have a favorable 5'
nucleotide and lower 5' end stability than the corresponding passenger strand (Fig. 1A).
Consistent with previous observations[27], the majority of *C. elegans* guide strands have a 5'
uracil, whereas passenger strands lack an obvious 5' nucleotide bias (Fig. 1B). To examine
thermodynamic asymmetry, we used free energy values of *C. elegans* miRNA duplex ends from
a previous study[28]. As expected, the average stability of duplex ends corresponding to guide
strands was lower than passenger strands (Fig. 1C, Supplementary Table 1). We observed
thermodynamic asymmetry when considering the first through fourth terminal nucleotides, with
the largest asymmetry observed for three terminal nucleotides (Fig. 1C). Thus, strand selection
of *C. elegans* miRNAs is correlated to 5' nucleotide identity and thermodynamic asymmetry.

95
A favorable 5' nucleotide is not sufficient to drive miRNA strand selection in vivo

97
To address whether 5' nucleotide identity and thermodynamic asymmetry directly influence
miRNA strand choice in vivo, we used genome editing to alter the duplex properties of
endogenous *C. elegans* miRNAs. We first examined a series of mutations in the *mir-58a*, *mir-1*
and *mir-2* miRNAs that primarily impacted the identity of 5' nucleotides (Fig. 2, Supplementary
Table 2). Both strands of the miR-58 duplex have a favorable 5' uracil, whereas only the guide
strands of miR-1 and miR-2 have a 5' uracil (Fig. 2). We used RNAcofold to determine how
each miRNA variation affected the thermodynamic stability of the terminal three nucleotides for
each duplex end[28] (Supplementary Table 3). For all three miRNAs, the guide duplex end has a
lower stability than the passenger duplex end (Fig. 2). Based on predictions from the twin-drive
model that considers 5' nucleotide identity and thermodynamic asymmetry[17], the four miR-58
mutations (M1-M4) should switch strand selection, the miR-1-M1 mutation should retain normal
strand choice, and the miR-2-M1 mutation should choose both duplex strands indiscriminately
leading to equal accumulation of both strands (Supplementary Table 4).

111
To investigate how miRNA mutations affect strand selection, we performed small RNA
sequencing in L4-staged animals (Supplementary Tables 5 and 6), a developmental stage
where miR-58, miR-1 and miR-2 have relatively high expression[29]. The miR-58-M1 and miR-58-
M2 mutations change the 5' uracil of the guide strand to less favorable nucleotides (adenine and
guanine respectively), which results in the 5' nucleotide of the guide strand becoming unpaired,
thereby modestly reducing duplex end stability (Fig. 2A). In the miR-58-M3 mutation, we
introduced a compensatory mutation in the passenger strand that restores base pairing to the 5'
adenine variant (Fig. 2A). Our analysis revealed that miR-58-M1 and miR-58-M2 retain normal
strand choice, while miR-58-M3 accumulated both duplex strands at near equal levels (Fig. 2A).
As the primary difference between the miR-58-M1 and miR-58-M3 mutations is the base-pairing
status of the 5' nucleotide, these results suggest that unpaired 5' nucleotides are highly
favorable for strand selection and can override a preferrable 5' nucleotide on the other strand.
Consistently, the miR-58-M4 mutation, which alters the 3' end of the guide strand to disrupt
base-pairing of the 5' nucleotide on the passenger strand, appeared to select the passenger
strand although total miR-58 levels were reduced (Fig. 2A).

127
Both miR-1-M1 and miR-2-M1 mutations change a terminal C:G pair on the passenger end of
the duplex into a U:G wobble pair, such that both duplex strands have a symmetrical 5' uracil
and similar thermodynamic end stabilities (Figs. 2B and 2C). For the miR-1-M1 mutation, we

observed some accumulation of passenger strand, however, the miR-1-M1 mutation also
appeared to affect mature miR-1 levels (Fig. 2B). On the other hand, we found that the miR-2-
M1 mutation caused a reversal of miR-2 strand selection (Fig. 2C). Given that the miR-2-M1
mutation has symmetrical 5' nucleotide identity and thermodynamic end stability (Fig. 2C), this
suggests that additional factors likely drive asymmetric strand selection of miR-2. Notably, our
data show that the twin-drive model poorly predicted the actual strand preference of these
miRNA mutants (Supplementary Table 4). We conclude that 5' nucleotide identity directly
influences miRNA strand selection in vivo, although favorable 5' nucleotides are not sufficient to
drive asymmetric strand choice, particularly when the 5' nucleotide of one strand is unpaired.
140
141 **Thermodynamic asymmetry is largely dispensable for miRNA strand choice**
142
We next examined mutations affecting thermodynamic asymmetry of *mir-58a*, *mir-1* and *mir-2*
(Fig. 3). The miR-58-M5, miR-58-M6 and miR-58-M7 mutations reduce thermodynamic end
stability without altering 5' nucleotide identity (Fig. 3A). The twin-drive model[17] predicted each of
these mutations, except for miR-58-M6 and miR-1-M2, to switch strands (Supplementary Table
4). Small RNA sequencing of L4-staged animals (Supplementary Tables 5 and 6) showed that
the symmetrical miR-58-M5 and miR-58-M6 mutants asymmetrically selected the canonical
miR-58 guide strand (Fig. 3A). Furthermore, while miR-58-M7 mutants accumulated some
passenger strands, they did not switch strand preference, despite the passenger duplex end
having lower stability (Fig. 3A). This observation suggested factors outside 5' nucleotide identity
and thermodynamic asymmetry promote strand selection of miR-58.
153
The more egregious miR-58-M8, miR-58-M9 and miR-58-M10 mutations reduce the stability of
the passenger duplex end while also introducing an unfavorable 5' nucleotide to the guide
strand (Fig. 3A). The miR-58-M8 and miR-58-M10 mutations both reversed strand selection,
demonstrating that there is not a barrier to selecting the miR-58 passenger strand (Fig. 3A).
While the miR-58-M9 mutation did not appear to switch strands, we observed a substantial
reduction in mature miR-58 levels (Fig. 3A), which was associated with a significant increase in
pri-mir-58 levels, suggesting ineffective processing of *pri-mir-58* (Extended Data Fig. 1A). We
did not observe significant changes in the levels or strand selection of the *mir-58* family
members *mir-80*, *mir-81* or *mir-82*[30], supporting our observations are specific to *mir-58*
(Extended Data Fig. 2).
164
The miR-1-M2, miR-1-M3 and miR-1-M4 mutations affect thermodynamic asymmetry while also
introducing a 5' uracil to the passenger strand; the 5' nucleotide of the miR-1 guide strand is
also mutated to a less favorable cytosine in miR-1-M4 mutants (Fig. 3B). We found that the
miR-1-M2 and miR-1-M3 mutations both preferred selection of the canonical miR-1 guide
strand, while miR-1-M4 asymmetrically selected the passenger strand (Fig. 3B). We observed a
reduction in mature miR-1 levels in each of the miR-1 mutants that we generated in this study
(Figs. 2B and 3B), possibly due to decreased *mir-1* expression as *pri-mir-1* levels were
significantly reduced in miR-1-M4 mutants (Extended Data Fig. 1B). The miR-2-M2 and miR-2-
M3 mutations incorporate a favorable 5' uracil to the miR-2 passenger strand, while also
introducing cytosines to the 5' end of the miR-2 guide strand that increase the stability of the
guide duplex end (Fig. 3C). The miR-2-M2 and miR-2-M3 mutations both resulted in highly
asymmetric selection of the miR-2 passenger strand and appeared to increase mature miR-2
levels (Fig. 3C). We observed modest effects on *pri-mir-2* levels in miR-2 mutants, suggesting
that any effects on mature miR-2 levels likely occur downstream of miRNA biogenesis
(Extended Data Fig. 1C). Notably, the twin-drive model poorly predicted the strand preference of
most miRNA mutants (Supplementary Table 4), suggesting additional factors contribute to
strand selection in vivo. Absolute quantification of select miRNA mutants agreed with our small

RNA sequencing analysis (Extended Data Fig. 3). Further, we observed phenotypes consistent
with loss of guide strand activity in strand switching mutants (Extended Data Fig. 4). In the *mir-*
58 family mutant background, the miR-58-M8, miR-58-M9 and miR-58-M10 mutations caused a
reduction in adult body size, consistent with loss of *mir-58* function[30] (Extended Data Fig. 4A-B).
Similarly, our *mir-1* and *mir-2* mutants showed resistance to levamisole and aldicarb
respectively, as previously reported for loss of *mir-1*[31] and *mir-2*[32] function (Extended Data Fig.
4C-D).
189
To address whether miRNA biogenesis was affected in miRNA mutants, we quantified how
each mutation affected isomiR populations. With a few exceptions, most of the miR-58 variants
did not significantly affect the distribution of isomiRs for guide or passenger strands compared
to controls, suggesting that miRNA mutations do not affect production of the expected mature
miRNA (Extended Data Fig. 5, Supplementary Tables 7 and 8). In miR-58-M4, miR-58-M9 and
miR-58-M10 mutants, a four-base truncation of the 3' end of the guide strand was more
abundant than the canonical guide sequence without affecting isomiRs for passenger strands
(Extended Data Fig. 5, Supplementary Tables 7 and 8). Each of these three mutations contain
the same CA:UC mutation towards the 3' end of the guide strand (Supplementary Table 2),
which may impact processing of these mutants, as we observed for miR-58-M9 (Extended Data
Fig. 1A), or could adversely affect the stability of the 3' end of the miRNA within the Argonaute
PAZ domain[13,14,33,34] thereby leading to trimming of exposed 3' ends[35,36]. Similarly, the miR-1-M4
mutation accumulated a one-base truncation of the 3' end of the canonical guide strand without
affecting passenger strand isomiRs, while isomiR populations appeared unaffected for other
miR-1 mutants (Extended Data Fig. 6A-B, Supplementary Table 9). The isomiR populations of
miR-2 appeared to be more dynamic, including two abundant 3' truncations of passenger
strands in wildtype animals (Extended Data Fig. 6C-D, Supplementary Tables 10 and 11).
Interestingly, passenger strand isomiRs were reduced in miR-2 mutants compared to wildtype,
suggesting increased precision of generating the expected mature sequence (Extended Data
Fig. 6C-D). The miR-2-M2 and miR-2-M3 mutants, which asymmetrically select the miR-2
passenger strand (Fig. 3F), also altered the isomiR frequency of the miR-2 guide strand
(Extended Data Fig. 6C-D). Thus, while most of the miRNA mutants do not significantly affect
isomiR populations, changes in isomiR populations were more frequently observed for
passenger strands of highly asymmetric miRNAs. It is possible that certain isomiRs have altered
duplex properties that promote their own strand selection, thereby leading to their increased
representation relative to the corresponding canonical miRNA, without impacting the actual rate
of isomiR formation. For example, in miR-2-M2 mutants, the most abundant guide strand isomiR
was a truncation of the 5' cytosine resulting in a more favorable 5' adenine (Extended Data Fig.
6C-D). Collectively, our data support a model where the identity and base-pairing status of 5'
nucleotides promotes miRNA strand selection but is not sufficient to explain strand preference in
vivo.
221
222   **Identification of duplex asymmetries associated with miRNA strand choice**
223
As we observed asymmetric strand selection for duplexes with symmetrical 5' ends (Figs. 2 and
3), we hypothesized that additional duplex features influence miRNA asymmetry in vivo. As we
were unable to identify additional asymmetries within the first four nucleotides on the 5' end of
guide and passenger strands (Fig. 4A), we next turned our attention to the 3' end of duplexes
and identified two nucleotide biases (Fig. 4B). The first bias was an adenine in the third-to-final
position of miRNA passenger strands, which was noticeably enriched among the 30 highest
expressed miRNAs in L4-staged animals (Fig. 4B) that represent more than 90% of total miRNA
reads and have more confident passenger strand annotations[37] (Fig. 4C). As miRNA duplexes
typically contain single stranded 2-nucleotide 3' overhangs[38,39], this adenine bias likely reflects

base paring to the preferred 5' uracil of guide strands. The second bias was the presence of a 3'
terminal cytosine on miRNA passenger strands, which was less frequently observed on guide
strands and nearly absent from the 3' ends of highly expressed miRNAs (Fig. 4B). As the levels
of many miRNAs are developmentally regulated, we next examined whether this 3' cytosine bias
was present throughout animal development or specific to the L4-stage. Using an existing small
RNA sequencing dataset[29], we quantified the observed 3' nucleotide identity of miRNA guide
and passenger strands throughout *C. elegans* development (Fig. 4D). Throughout animal
development, we found that very few guide strands ended with a 3' cytosine, whereas cytosine
was among the most frequent 3' nucleotides of passenger strands (Fig. 4D). We hypothesized
that 3' nucleotide asymmetry is associated with miRNA strand choice throughout *C. elegans*
development.
244
245 **A 3' cytosine on miRNA passenger strands promotes strand selection in *C. elegans***
246
To examine whether 3' nucleotide identity influences miRNA strand selection in vivo, we created
an additional set of *mir-58* mutations that alter 3' nucleotide identity in the wildtype or
symmetrical miR-58-M6 duplexes (Fig. 5, Supplementary Table 2). The passenger strand of
miR-58 has a 3' cytosine and the guide strand has a 3' uracil (Fig. 5A). We generated mutants
where the 3' nucleotide of both strands were swapped, or the 3' nucleotide of one strand was
mutated to produce duplexes containing symmetrical 3' nucleotides (Fig. 5A). While swapping
the 3' nucleotide of the wild type miR-58 duplex in miR-58-M14 mutants was not sufficient to
reverse strand selection, swapping the 3' nucleotides of the miR-58-M6 duplex in miR-58-M16
mutants reversed strand preference (Figure 5A, Supplementary Tables 5 and 6). This
observation suggests a novel role for 3' nucleotide identity in directing miRNA strand choice.
Interestingly, in the miR-58-M6 background, we observed near equal levels of both guide and
passenger strands for the miR-58-M17 mutant where both strands have 3' cytosine, but
asymmetric selection of the miR-58 guide strand in miR-58-M18 mutants where both strands
have 3' uracil (Fig. 5A). Thus, a specific preference for 3' cytosine on passenger strands
appears to regulate strand selection of miR-58.
262
We next examined 3' nucleotide mutations of *mir-2*, which has a 3' guanine on the passenger
strand and a 3' cytosine on the guide strand (Fig. 5B). To minimize the effect of 5' nucleotide
identity, we mutated the miR-2-M1 duplex where both miR-2 strands have a symmetrical 5'
uracil (Fig. 5B, Supplementary Table 1). The miR-2-M4 mutation switches the 3' nucleotides of
each miR-2-M1 strand and the miR-2-M5 mutations mutates the 3' nucleotide of the miR-2-M1
passenger strand so that both strands have a symmetrical 3' guanine (Fig. 5B). The miR-2-M1
mutation reversed strand preference of miR-2 (Figs. 2B and 5B) presumably due to the 3'
cytosine on the canonical miR-2 guide strand (passenger strand of miR-2-M1) (Fig. 5B). This
passenger strand accumulation in miR-2-M1 was relieved by mutation of the 3' cytosine on the
miR-2 guide strand in miR-2-M4 and miR-2-M5 mutants, consistent with 3' cytosine directing
strand selection of mIR-2 (Fig. 5B). Analysis of miR-2 (Extended Data Fig. 6C-D,
Supplementary Tables 10 and 11) and miR-58 isomiRs (Extended Data Fig. 5, Supplementary
Tables 7 and 8) indicated that the 3' nucleotide variants were processed correctly. Furthermore,
introducing a 3' cytosine to either strand of the miR-48 duplex (Supplementary Table 2) led to a
small accumulation of the opposing strand without affecting miR-48 processing (Extended Data
Fig. 7, Supplementary Table 12). Similarly, introducing a 3' cytosine to the guide strand of miR-
71 (Supplementary Table 2) promoted selection of the miR-71 passenger strand without
affecting isomiR populations (Extended Data Fig. 7, Supplementary Table 13), indicating that 3'
nucleotide identity influences the strand selection of a wide range of miRNAs in vivo.
282

The nucleotide composition of 3' overhangs may influence strand asymmetry, as we observed
modest differences in strand selection when we swapped 3' nucleotides or the dinucleotide
overhangs of miR-58 duplexes (Extended Data Fig. 8). However, we did not identify obvious
nucleotide asymmetries beyond the terminal nucleotide of 3' ends, so we chose to focus on the
role of 3' nucleotide identity in this study (Fig. 4B). We next asked whether 3' nucleotide identity
might be directly recognized by Argonaute. To address this, we mutated a putative 3' nucleotide
binding pocket[17] that is positioned near the 5' nucleotide binding pocket in the MID domain[13,14,25]
of the *C. elegans* miRNA-associated Argonaute ALG-1 (Extended Data Fig. 9A-B). Although
small RNA sequencing did not identify miRNA strand switching in ALG-1[7A] mutants, we
observed accumulation of several miRNA passenger strands (Extended Data Fig. 9C,
Supplementary Table 6), suggesting that the interaction between 3' nucleotides and Argonaute
may, at least in part, promote accurate strand selection. Collectively, our results suggest that a
3' cytosine on miRNA passenger strands promotes guide strand selection and is sufficient to
direct strand selection of miRNAs with symmetrical 5' duplex ends.
297
298   **3' nucleotide identity plays a conserved role in directing miRNA choice**
299
We next asked whether 3' nucleotide identity influences miRNA strand selection in an
evolutionarily conserved fashion. We used the shERWOOD UltramiR system[40] to introduce a
set of miR-58 variants into human HEK293T cells (Fig. 6A). The human genome does not
encode a miR-58 homolog, which allowed us to track the strand preference of the exogenous
miR-58 variants. We first transfected the wildtype miR-58 and symmetrical miR-58-M6 duplexes
as well as their respective 3' nucleotide swapped variants miR-58-M14 and miR-58-M16 (Fig.
6A). Our small RNA sequencing (Supplementary Table 14) showed that the canonical miR-58
guide strand is asymmetrically selected, like what is observed in *C. elegans* (Fig. 6A). However,
unlike in *C. elegans*, both strands of the symmetrical miR-58-M6 duplex appear to be selected
at near equal frequency (Fig. 6A), suggesting there may be an inherent difference in miRNA
strand selection between *C. elegans* and humans. Furthermore, we found that passenger strand
levels were reduced in the 3' nucleotide swapped variants miR-58-M14 and miR-58-M16 that
each have a 3' cytosine on the miRNA guide strand (Fig. 6A), suggesting 3' cytosine is less
favored in human cells. Each of the miR-58 variants appeared to be processed correctly based
on isomiR analysis (Extended Data Fig. 10, Supplementary Table 15). However, an additional
set of mutations that we designed in a modified version of the human miR-339 duplex were not
processed as expected (Extended Data Fig. 10, Supplementary Table 15), despite a modest
effect on strand asymmetry by those mutations (Supplementary Table 14).
318
To characterize the 3' nucleotide preference in human cells, we created several miR-58-M6
duplexes comprising 16 possible 3' nucleotide combinations for each duplex strand (Fig. 6B). To
quantify the levels of each miR-58 variant, we performed absolute quantification for each duplex
strand (Fig. 6C, Supplementary Table 16). We found that a 3' guanine on the guide (3p) strand
of miR-58-M6 and a 3' uracil on the passenger (5p) strand of miR-58-M6 were most favorable
for guide strand selection (Fig. 6B). Notably, when each miR-58-M6 strand contained a 3' uracil
or adenine, the 3p strand was highly favored whereas when each strand contained a 3' guanine
or cytosine the 5p strand was asymmetrically selected (Fig. 6B). These observations suggest
that precursor origin may influence miRNA strand choice, or that additional duplex features,
such as 3' overhang composition, may play a prominent role in directing miRNA strand choice in
human cells. We were unable to identify an obvious nucleotide bias at the 3' end of human
miRNA duplexes (Fig. 6C-D, Supplementary Table 17), consistent with the idea that additional
duplex features may influence strand asymmetry in human cells. Collectively, our results
indicate that 3' nucleotide asymmetry plays an evolutionarily conserved role in regulating miRNA
strand choice. While it is unclear how 3' nucleotide asymmetry is recognized, we propose two

alternative hypotheses that are not mutually exclusive, where the 3' nucleotide is directly bound
by Argonaute or by an accessory protein to facilitate guide strand selection and passenger
strand removal (Fig. 6E).
337
338   **Discussion**
339
How does 3' nucleotide identity promote miRNA strand choice? One possibility is that
Argonaute directly binds to the 5' nucleotide of the guide strand[13,14,25] and the 3' nucleotide of
the passenger strand to initiate duplex unwinding (Fig. 6E). Consistent with this idea, we found
that mutating a putative 3' nucleotide binding pocket on Argonaute[17] led to an accumulation of
some miRNA passenger strands (Extended Data Fig. 9C). It is possible that duplex binding
triggers a conformational change of Argonaute that promotes passive wedging by its N-terminal
domain[55]. Further, mutations in *C. elegans* ALG-1 and its human homolog Ago2 affect miRNA
strand asymmetry[56,57], although it is currently unclear whether those mutations are connected to
recognition of 3' nucleotide identity. An alternative, but not mutually exclusive hypothesis is that
Argonaute interacts with additional factors that are responsible for recognizing 3' nucleotide
identity (Fig. 6E). For example, the Dicer-interacting proteins TRBP and PACT regulate miRNA
strand selection in humans[58–60], and Dicer can reposition the miRNA duplex after dicing[59], which
could indicate that Dicer may pass miRNA duplexes to Argonaute in a favorable orientation.
Notably, the *C. elegans* genome lacks obvious homologs for both TRPB and PACT, which could
at least partially explain some of the differences in strand selection and 3' nucleotide
preferences that we observed between *C. elegans* and human cells. It is also possible that the
protein composition of different strand selection complexes might regulate miRNA strand choice
or play a role in sorting duplexes to specific Argonaute proteins that may have different 3'
nucleotide preferences. Although 3' nucleotide identity appears to play a conserved role in
miRNA strand selection, it is worth noting that human miRNAs, unlike *C. elegans* miRNAs, lack
an obvious 3' nucleotide asymmetry. This observation highlights the value of model systems,
such as *C. elegans*, in understanding the basic mechanisms of miRNA strand selection, of
which our current understanding remains incomplete.
363
While both regulation and dysregulation of miRNA strand asymmetry have previously been
reported (see review[18]), current models of miRNA strand selection lack a dynamic component to
explain context-dependent changes in strand preference[17]. As 3' isomiRs are typically more
frequent than 5' isomiRs[29,41,42], changes in 3' isomiRs reported in developmental[29,43–45],
cellular[46,47] or disease[48] contexts may underly miRNA strand dynamicity. Some 3' modifications,
such as miRNA tailing and trimming[36,49], likely occur after strand selection and are not
necessarily expected to affect miRNA strand choice. Other 3' modifications, such as adenylation
or uridylation, often target pre-miRNA molecules[42,50–52] and are more likely to influence strand
asymmetry. Consistently, uridylation of human miR-324 leads to altered strand selection,
although uridylation of miR-324 may primarily influence Dicer-dependent processing[53].
Intriguingly, cytidylation of miRNA precursors has been observed in Arabidopsis[54] and the miR-
83-3p guide strand is cytidylated in *C. elegans*[49]. As our data indicates that a 3' cytosine on
miRNA passenger strands is favorable for strand selection, cytidylation of one strand could
promote selection of the other duplex strand. However, modification of miRNA passenger
strands may be difficult to detect due to their rapid degradation in cells. Additional work will be
needed to fully understand how 3' nucleotide are recognized and how changes in 3' nucleotide
identity may regulate miRNA strand selection across different contexts and potentially disrupt
strand selection in human disease.
382
383   **Methods**
384

C. elegans **Strains and Genetic Analysis**

A complete list of *C. elegans* strains used in this study is provided in Supplementary Table 18. All strains were derived from the wild-type N2 genetic background. Strains were maintained at 20°C under standard growth conditions[61]. Nematode growth medium (NGM) plates that were seeded with *Escherichia coli* OP50 were used as a food source. For developmental stage-specific analysis, animals were synchronized using standard alkaline hypochlorite treatment[62,63] and harvested when most (>90%) of the larval population reached the desired developmental stage. For body length measurements, adult animals were imaged on a Leica DM6 upright microscope. Adobe illustrator was used to quantify body length by measuring the path length of a curved line drawn through the center of the animal from the head to base of the tail. For drug assays, young adult worms were added to plates containing 200 µM levamisole[31] or 1mM aldicarb[32] and scored for paralysis at indicated time intervals. Paralysis was defined as the inability for worms to move in response to a touch stimulus from a platinum wire.

Profiling miRNA Duplex Characteristics

Mature *C. elegans* and human miRNA sequences were retrieved from miRbase v 22.1[37]. Guide and passenger strand annotations were determined based on which strand had the most reported reads on miRbase. 5' nucleotide information was compiled for 5p and 3p strands. RNAcofold from the ViennaRNA Package 2.0[64] was used for duplex folding and to calculate thermodynamic end stability at 20°C as previously described[28]. Predictions of miRNA strand selection were calculated using twin-drive model of strand selection as previously described[17].

CRISPR/Cas9 Genome Editing

Streptococcus pyogenes Cas9 (IDT, Alt-R® S.p. Cas9 Nuclease V3), single-stranded repair templates (IDT, custom ultramers), crRNA and tracrRNA were commercially sourced. A full list of oligonucleotides used in this study is provided in Supplementary Table 19. Microinjection and screening were performed using *dpy-10* co-conversion as previously described[65]. Cas9 was injected at a final concentration of 2.65 µM and all injection mixes included 200 µM KCl and 7.5 µM HEPES [pH 7.4]. All crRNAs were injected at a final concentration of 50 µM, except *dpy-10* crRNA was injected at a final concentration of 5 µM. Equimolar tracrRNA was included in all injection mixes. Single stranded DNA oligonucleotides were used to facilitate homology-directed repair and were included at final concentrations of 6 µM or 3 µM (*dpy-10*).

RNA Extraction and Isolation

Immediately after sample collection, samples were centrifuged and flash-frozen in liquid nitrogen and stored at -80°C until ready for use. Pellets were thawed in 250 µL RNase-free water (final volume). 1 mL of Trizol (Invitrogen) was added and the mixture was continuously vortexed for 10 minutes. Afterwards, 212 µL of chloroform was added and vortexed for an additional minute. The aqueous and organic layers were separated by centrifugation at 19000 x g for 10 minutes at 4°C. The aqueous phase was then recovered and re-extracted using equal volume of phenol:chloroform:isoamyl alcohol (25:24:1, VWR). The RNA was precipitated for 2 hours at -20°C using an equal volume of isopropanol. 1.8 µL of Glycoblue (Invitrogen) was added as a co-precipitant. The RNA pellets were washed using 80% (v/v) ethanol and allowed to air-dry on ice for 10 minutes. The dried pellets were dissolved in RNase-free water and stored at -80°C. RNA concentration was determined using a Nanodrop spectrophotometer.

Small RNA Sequencing and Analysis

For preparation of small RNA libraries, total RNA was first size selected using a 15% denaturing gel using UreaGel (National Diagnostics) reagents. Single-stranded 18mer (AGCTGCCAGCAACTCTAA) and 26mer (TAGTATGTAGCTGCCAGCAACTCTAA) were combined at a final concentration of 5 µM and used as a molecular ladder in an empty lane. Gels were soaked in 1XTBE solution containing 1X SYBR Gold stain (Invitrogen) for 10 minutes and subsequently visualized under UV light. Small RNAs migrating between the 16mer and 26mer markers were excised from the gel and soaked in 750 µL of TE buffer supplemented with 0.3 M NaCl overnight while shaking at 550 rpm at 7°C. The solution was transferred to a 0.45 µm column and centrifuged at 4°C for 10 minutes at 2500 x g. Small RNAs were precipitated for 2 hours at -20°C using equal volume of isopropanol and 10% sample volume 3M NaOAc. 1.5 µL of Glycoblue (Invitrogen) was used as a co-precipitant. The small RNA pellets were washed once using 80% (v/v) ethanol and allowed to air-dry on ice for 10 minutes. The dried pellets were dissolved in 11 µL of RNase-free water. Size-selected small RNA samples were used to make libraries using the Nextflex Small RNA-Seq Kit v3 (PerkinElmer) or the Nextflex Small RNA-Seq Kit v4 (Revvity) according to manufacturer instructions. Please note that the Nextflex Small RNA-Seq Kit v3 (PerkinElmer) was discontinued during this study. We separated small RNA sequencing data acquired from the v3 (Supplementary Table 5) and v4 kits (Supplementary Table 6) and only made direct comparisons for samples that were sequenced using the same kit, as we observed differences in normalized read counts for several miRNAs when using different kits. All small RNA libraries were sequenced at the Kansas State University Integrated Genomics Facility using an Illumina NextSeq 500 instrument at 75 cycles.

Small RNA sequence data were processed and analyzed as previously described[29] The cutadapt tool[66] was used to process and trim the sequencing reads. The 3' adapter sequences were first removed using the parameters "-a ATCTCGTATGCCGTCTTCTGCTTGX" and demultiplexed using the parameters "e 1 -a file$:barcodes.fasta", where the "barcodes.fasta" contains a list of barcode sequences in fasta format. The 3' primer sequence was then removed from each read using the parameters "-a TGGAATTCTCGGGTGCCAAGGAACTCCAGTCAC". For the Nextflex Small RNA-Seq Kit v3, each end of the processed sequencing reads contained four-nucleotide randomers that were removed using cutadapt. The final four nucleotides of each sequencing read were trimmed using the parameters "-u -4". The first four nucleotides were trimmed using the parameters "-u 4 -m17 -M29", which also removed processed reads that were shorter than 17 or longer than 19 nucleotides. Read quality was examined using the fastqc tool[67]. Reads were aligned to a list of mature *C. elegans* miRNAs using bowtie[68]. The following command was used for bowtie alignments: "-x miR -a --best --strata --norc -v 3", where "miR" is a bowtie build assembled using a fasta file containing all *C. elegans* mature miRNA sequences obtained from miRbase. The samtools tool[69] was used to generate, sort and index bam files to generate read count data. A custom bash script was used to identify and quantify isomiR abundance from fastq files. Normalization (reads per million) was performed by multiplying the number of mapped reads for individual miRNA loci by the total number of mapped miRNA reads divided by one million.

qPCR and Absolute Quantification

The QuantiNova SYBR Green RT-PCR kit (Qiagen) was used for standard qPCR reactions according to manufacturer instructions. Final reaction volumes were adjusted to 5 µL and 100 ng of total *C. elegans* RNA was used as a template. Absolute quantification of miRNAs using Taqman was performed as previously described[70]. Standard curves were first made using known quantities (0.6ng/µL, $0.6x10^{-2}$ ng/µL, $0.6x10^{-4}$ ng/µL, $0.6x10^{-6}$ ng/µL, and $0.6x10^{-8}$ ng/µL)

of commercially purchased (IDT) single-stranded RNA oligonucleotides prepared from a 2 µg/µL
stock. All reactions were carried out at a final volume of 10 µL using the Taqman™ MicroRNA
Reverse Transcription Kit (Thermofisher) and Taqman™ Universal Master Mix II, no UNG
(Thermofisher) according to manufacturer instructions. Custom or predesigned Taqman™ Small
RNA Assays (ThermoFisher) were used at 1X final concentration. 1 ng of total *C. elegans* RNA
was used as a template for reverse transcription (10 µL reaction volume) and 0.67 µL of the
cDNA product was used for experimental Taqman assays. The ABI 7500 Real-Time PCR
System (Applied Biosystems) was used to run all qPCR and Taqman reactions. A full list of
oligonucleotides and Taqman assays used in this study is provided in Supplementary Table 19.
496
UltramiR Plasmid Construction

A full list of plasmids used in this study is provided in Supplementary Table 20. Each miRNA
variant was cloned into the pLMN-ZsGreen-Neomycin-shRNA vector (transOMIC) and
transformed into NEB$^®$ Stable Competent *E. coli* (High Efficiency) according to manufacturer
instructions. The pJM1 through pJM7 plasmids were created by Gibson assembly of
Gblock gene fragments (IDT) (Supplementary Table 19) and HpaI/MluI (NEB) digested pLMN-
ZsGreen-Neomycin-shRNA vector using the Gibson Assembly$^®$ Cloning Kit (NEB) according to
manufacturer instructions. The pJM10 through pJM23 plasmids were created by performing
site-directed mutagenesis of the pJM2 plasmid to mutate the nucleotides corresponding to the 3'
end of the 5p or 3p strands using the Q5$^®$ Site-Directed Mutagenesis Kit (NEB) according to
manufacturer instructions. The primers used for site-directed mutagenesis are listed in
Supplementary Table 19. All clones were verified by Sanger sequencing prior to transfection
510
Human Cell Culture and Transfection

Human HEK 293T cells were seeded in 12-well plates containing 1 mL DMEM with 10% FBS
and transfected when reaching ~90% confluency. In one tube 1 µg of UltramiR plasmid was
added to 100 µL of Opti-MEM™ (Thermofisher) and vortexed. In a second tube, 5 µL of 1
mg/mL polyethylenimine (PEI) reagent was added to 100 µL of Opti-MEM™ (Thermofisher),
vortexed and incubated at room temperature for 5 min. The contents of the second tube were
subsequently pipetted into the first tube, vortexed and incubated at room temperature for 15
min. The mixture was then carefully added to the cell media dropwise. After 6 hours, the cell
media was changed with fresh warmed DMEM with 10% FBS. 48 hours post-transfection, cells
were imaged for ZsGreen expression (Supplementary Figure 1), harvested in 1 mL of Trizol
(Invitrogen), frozen and stored at -80ºC until use.
523
Statistical Analysis

526   R statistical software was used to calculate all statistics. All *P*-values were calculated using two-
527   tailed *t*-tests assuming equal variance among samples. All statistics are presented as mean ±
528   one standard deviation unless otherwise specified.
529
Data Availability

532   Small RNA sequencing data files will be made available in the GEO database upon final
533   publication. All strains and plasmids used in this study are available upon request. All other data
534   supporting the findings of this study are available from the corresponding author on reasonable
535   request.
536
Acknowledgements

538
We thank members of the Zinovyeva lab for helpful discussions. We thank Hui Tian, Sarah
Coffey and Katherine Adams (Hwang) for technical assistance in generating some worm strains,
and Alina Akhunova from the Kansas State Integrated Genomics Facility for technical expertise
in generating small RNA sequencing data. We are grateful to Anne Hart for sharing *mir-2*
mutant strains. Some of the strains used in this study were provided by the *Caenorhabditis*
Genetics Center (CGC) which is funded by NIH Office of Research Infrastructure Programs
(P40 OD010440). This work was supported by grants from the National Institutes of Health
(R35GM124828 to A.Z. and R01GM117406 to C.M.H.), the Kansas INBRE program (P20
GM103418) and the Johnson Cancer Center at K.S.U. J.C.M was supported by a fellowship
from the NIH (F32GM148040).

549
550 **Author Contributions**
551
552 J.C.M and A.Z. conceptualized and designed the experiments. A.Z and C.M.H supervised the
553 research and acquired funding. J.C.M, S.K, H.O, H.C, S.Z, W.S, J.S and M.B carried out the
554 experiments. J.C.M and A.Z. performed data analysis and interpreted results. J.C.M and G.P
555 performed the required bioinformatics. J.C.M wrote the original manuscript draft, and all other
556 authors edited the manuscript.
557
558 **Competing Interests**
559
560 The authors declare no competing interests.

Figure Legends

Figure 1. Duplex characteristics of *C. elegans* miRNAs. (A) Duplex features associated with miRNA strand selection. The guide (miR) end of typically contains a favorable 5' nucleotide and lower thermodynamic stability than the passenger (miR*) duplex end. The guide strand is shown in maroon, and the passenger strand is shown in gray. U:A pairs are illustrated by an open circle, G:U wobble pairs are indicated by an open circle with a vertical line and G:C pairs are represented by closed circles. The sizes of bulges and mismatched nucleotides are drawn to scale. (B) Quantification of 5' nucleotide identity for *C. elegans* miRNA guide (maroon) and passenger strands (gray). n is given as the number of miRNA duplexes quantified (C) Quantification of the free energy of miRNA duplex ends corresponding to guide (maroon) and passenger (gray) duplex ends. Free energy values were calculated by in silico unwinding[28] of the terminal 1:4 nucleotides of each duplex end.

Figure 2. 5' nucleotide identity influences miRNA strand selection in vivo. (Left) Schematic representations of (A) miR-58, (B) miR-1 and (C) miR-2 duplex mutations that affect 5' nucleotide identity. (A-C) The guide strand is shown in maroon, and the passenger strand is shown in gray. Mutated nucleotides are indicated by red text. U:A pairs are illustrated by an open circle, G:U wobble pairs are indicated by an open circle with a vertical line and G:C pairs are represented by closed circles. Free energy values (kcal/mol) of the terminal three nucleotides (boxed) are shown on each duplex end. The less stable duplex end is marked with an asterisk and highlighted in a light red box. RNAcofold was used to predict duplex structures at 20°C. (Right) Quantification of small RNA sequencing results for (A) miR-58, (B) miR-1 and (C) miR-2. Data are presented as mean ± standard deviation, with each dot representing quantification from a single library using the NEXTFLEX v3 Small RNA Sequencing Kit. Maroon bars correspond to miRNA guide strands and gray bars represent miRNA passenger strands. Normalization was performed by dividing raw read counts by total mapped miRNA reads and multiplying by one million to acquire reads per million (rpm) values.

Figure 3. Thermodynamic asymmetry is dispensable for miRNA strand selection. (Left) Schematic representations of (A) miR-58, (B) miR-1 and (C) miR-2 duplex mutations that affect the relative thermodynamic stability of each duplex end. (A-C) The guide strand is shown in maroon, and the passenger strand is shown in gray. Mutated nucleotides are indicated by red text. U:A pairs are illustrated by an open circle, G:U wobble pairs are indicated by an open circle with a vertical line and G:C pairs are represented by closed circles. Free energy values (kcal/mol) of the terminal three nucleotides (boxed) are shown on each duplex end. The less stable duplex end is marked with an asterisk and highlighted in a light red box. RNAcofold was used to predict duplex structures at 20°C. (Right) Quantification of small RNA sequencing results for (A) miR-58, (B) miR-1 and (C) miR-2. Data are presented as mean ± standard deviation, with each dot representing quantification from a single library using the NEXTFLEX v3 Small RNA Sequencing Kit. Maroon bars correspond to miRNA guide strands and gray bars represent miRNA passenger strands. Normalization was performed by dividing raw read counts by total mapped miRNA reads and multiplying by one million to acquire reads per million (rpm) values.

Figure 4. *C. elegans* miRNA duplex asymmetries. (A) Sequence logo representation of 5' nucleotide frequencies for *C. elegans* miRNA guide (left) and passenger (right) strands. (B) Sequence logo representation of 3' nucleotide frequencies for *C. elegans* miRNA guide (left) and passenger (right) strands. Asterisk indicates the third nucleotide from the 3' duplex end of the passenger strand that is adjacent to the 3' dinucleotide overhang and is expected to base pair to the 5' nucleotide of the guide strand. (A-B) Sequences are given in 5'-3' orientation, with

the 5' or 3' terminal nucleotides indicated by a bulbed line above. Sequence logos were created
using the weblogo3 server and colors were modified using Adobe illustrator. (C) Cumulative
expression of *C. elegans* miRNAs at L4-stage. Each dot represents one miRNA (guide and
passenger strands), plotted in order of most abundant to least abundant. The 10 most abundant
miRNAs account for ~65% and the 30 most abundant miRNAs represent >90% of all miRNA
sequencing reads at the L4 stage. (D) Quantification of 3' nucleotide frequency for miRNA guide
(left) and passenger (right) strands throughout *C. elegans* development. Stacked bar graphs
represent the normalized abundance of canonical miRNA guide or passenger strands (not
including isomiRs) sorted by terminal 3' nucleotide identity. Schematic (top) illustrates an
example of a miRNA duplex loaded into Argonaute (gray) and the arrowhead indicates the
position of the 3' nucleotide of the guide (left) or passenger (right) strand. Developmental small
RNA sequencing data were acquired from a previous study[29].
624
Figure 5. A 3' cytosine on miRNA passenger strands is favorable for guide strand choice in *C.*
elegans. (Left) Schematic representations of (A) miR-58 and (B) miR-2 duplex mutations that
affect 3' nucleotide identity. (A-B) The guide strand is shown in maroon, and the passenger
strand is shown in gray. The nucleotide composition of dinucleotide overhang are shown for
each strand. Note that the guide strand of miR-2 has an atypical trinucleotide overhang at its 3'
end. Mutated nucleotides are indicated by red text. U:A pairs are illustrated by an open circle,
G:U wobble pairs are indicated by an open circle with a vertical line and G:C pairs are
represented by closed circles. RNAcofold was used to predict duplex structures at 20°C. (Right)
Quantification of small RNA sequencing results for (A) miR-58 and (B) miR-2. Data are
presented as mean ± standard deviation, with each dot representing quantification from a single
library using the NEXTFLEX v4 Small RNA Sequencing Kit. Maroon bars correspond to miRNA
guide strands and gray bars represent miRNA passenger strands. Normalization was performed
by dividing raw read counts by total mapped miRNA reads and multiplying by one million to
acquire reads per million (rpm) values.
639
Figure 6. 3' nucleotide identity influences miRNA strand asymmetry in human HEK293T cells.
(A) (Left) Schematic representations of exogenous miR-58 duplexes transfected into HEK293T
cells that alter 3' nucleotide identity. The canonical miR-58 guide strand (from *C. elegans*) is
shown in maroon, and the passenger strand is shown in gray. The nucleotide composition of
dinucleotide overhang are shown for each strand. U:A pairs are illustrated by an open circle,
G:U wobble pairs are indicated by an open circle with a vertical line and G:C pairs are
represented by closed circles. RNAcofold was used to predict duplex structures at 37°C. (Right)
Quantification of small RNA sequencing results for transfected miR-58 variants. Data are
presented as mean ± standard deviation, with each dot representing quantification from a single
library using the NEXTFLEX v4 Small RNA Sequencing Kit. Maroon bars correspond to
canonical miRNA guide strands and gray bars represent canonical miRNA passenger strands.
Normalization was performed by dividing raw read counts by total mapped miRNA reads and
multiplying by one million to acquire reads per million (rpm) values. (B) Absolute quantification
of miR-58-M6 duplex variants comprising each combination of 3' nucleotides on each duplex
strand. Heatmap illustrates calculated ln(5p/3p) strand ratios where a deeper maroon color
indicates preference for the miR-58-M6-3p (guide) strand. The non-transformed 5p/3p strand
ratios are given in each box. (C) Sequence logo representation of 5' nucleotide frequencies for
human miRNA guide (left) and passenger (right) strands. (D) Sequence logo representation of
3' nucleotide frequencies for human miRNA guide (left) and passenger (right) strands. (C-D)
Sequences are given in 5'-3' orientation, with the 5' or 3' terminal nucleotides indicated by a
bulbed line above. Sequence logos were created using the weblogo3 server and colors were
modified using Adobe illustrator. (E) Hypothetical models of how 3' nucleotide identity influences
miRNA strand selection. (Top) The MID domain of Argonaute may directly interact with the 5'

nucleotide of the guide strand and the 3' nucleotide of the passenger strand. Such interaction
may induce a conformational change that promotes duplex wedging and release of the
passenger strand. (Bottom) Alternatively, additional factors may interact with miRNA 3'
nucleotides to form a strand selection complex. Dissolution of such complex could facilitate
guide strand retention within Argonaute and passenger strand removal.
668
Extended Data Figure 1. Quantification of (A) *pri-mir-58*, (B) *pri-mir-1* and (C) *pri-mir-2* levels in
L4-staged miRNA mutants by qRT-PCR. (A-C) Each dot represents one qRT-PCR data point.
Each sample was performed in duplicate and at least three independent replicates were
included for each mutation. Data were first normalized to α-tubulin (*tba-1*) and subsequently
normalized to appropriate PAM or wildtype (WT) controls. The dashed horizontal line indicates
the self-normalized average for appropriate controls. Boxes range from the first to third quartile
of the data. The thick bar represents the statistical median. Lines extend to the minimum and
maximum data point excluding outliers that were defined as 1.5 times the interquartile range.
Statistical analysis: *p<0.05, ***p<0.001.
678
Extended Data Figure 2. Analysis of miR-58 family miRNAs in miR-58 mutants. (A) Sequence
alignment of the *C. elegans* miR-58 miRNA family members. (B-C) Quantification of mature
miR-80 levels in L4-staged animals normalized to PAM controls. Libraries were created with the
(B) Nextflex v3 small RNA sequencing kit or the (C) Nextflex v4 small RNA sequencing kit. (D-
E) Quantification of mature miR-81 levels in L4-staged animals normalized to PAM controls.
Libraries were created with the (D) Nextflex v3 small RNA sequencing kit or the (E) Nextflex v4
small RNA sequencing kit. (F-G) Quantification of mature miR-82 levels in L4-staged animals
normalized to PAM controls. Libraries were created with the (F) Nextflex v3 small RNA
sequencing kit or the (G) Nextflex v4 small RNA sequencing kit. (B-G) Data are presented as
mean ± standard deviation, with each dot representing quantification from a single library.
Maroon bars correspond to canonical miRNA guide strands and gray bars represent canonical
miRNA passenger strands.
691
Extended Data Figure 3. Absolute quantification of miRNA levels in representative miRNA
mutants using miRNA Taqman assays. (A-C) Standard curves created using known amounts of
guide and passenger strand RNA for (A) miR-58, (B) miR-1 and (C) miR-2. Linear regression
and R2 coefficients are indicated for guide (maroon) and passenger (gray) strands. Each dot
represents an independently prepared Taqman assay. (D-E) Absolute quantification of both
miR-58 strands in (D) L4-staged or (E) embryonic miR-58 mutants. (F-G) Absolute quantification
of both miR-1 strands in (F) L4-staged or (G) embryonic miR-1 mutants. (H-I) Absolute
quantification of both miR-58 strands in (H) L4-staged or (I) embryonic miR-52 mutants. (D-I)
Data are presented as mean ± standard deviation. Each sample was assayed in triplicate and at
least three independent replicates were included for each mutation. Each dot represents
quantification from a single Taqman assay. Maroon bars correspond to canonical miRNA guide
strands and gray bars represent canonical miRNA passenger strands.
704
Extended Data Figure 4. Phenotypic analysis of miRNA mutants. (A) Representative images of
a (top) *mir-80; mir-81 mir-82* triple miR-58 family mutant and (bottom) *mir-80; mir-58; mir-81 mir-*
82 quadruple family mutant illustrates reduced adult body size associated with loss of miR-58
family function. (B) Quantification of adult body length in miR-58 mutants in the *mir-80; mir-81*
mir-82 genetic background. Each dot represents one animal. Boxes range from the first to third
quartile of the data. The thick bar represents the statistical median. Lines extend to the
minimum and maximum data point excluding outliers that were defined as 1.5 times the
interquartile range. (C) Percent of miR-1 mutant animals paralyzed on plates containing 200 μM
aldicarb over time. Data are presented as mean ± standard deviation. (D) Percent of miR-2

mutant animals paralyzed on plates containing 1 mM aldicarb over time. Data are presented as
mean ± standard deviation. (B-D) Statistical analysis: ***p<0.001.
716
Extended Data Figure 5. Analysis of miR-58 isomiRs in miR-58 mutants. (A-B) Heatmaps
illustrating relative distribution of miR-58-3p (canonical guide strand) isomiRs in miR-58
mutants. Data are from small RNA sequencing data of L4-staged mutants that were created
using the (A) Nextflex v3 or (B) Nextflex v4 small RNA sequencing kits. (C-D) Heatmaps
illustrating relative distribution of miR-58-5p (canonical passenger strand) isomiRs in miR-58
mutants. Data are from small RNA sequencing data of L4-staged mutants that were created
using the (C) Nextflex v3 or (D) Nextflex v4 small RNA sequencing kits. (A-D) Data are
presented as the percent occurrence of a given isomiR within each individual mutant and do not
reflect possible changes in miRNA abundance. Each isomiR is illustrated on the left. The
maroon (guide) and gray (passenger) bars indicates the expected miR-58 sequence. Truncated
isomiRs are indicated by negative values corresponding to the length of the truncation.
Nucleotide extensions are depicted by the corresponding letters to each duplex end, which may
include templated or non-templated (uridylation and adenylation) extensions.
730
Extended Data Figure 6. Analysis of miR-1 and miR-2 isomiRs in miRNA mutants. Heatmaps
illustrate relative distribution of (A) miR-1-3p (canonical guide strand) or (B) miR-1-5p (canonical
passenger strand isomiRs in miR-1 mutants. (A-B) Data are from small RNA sequencing data of
L4-staged mutants. Heatmaps depict (C) miR-2-3p (canonical guide strand) and (D) miR-2-5p
(canonical passenger strand) isomiR distribution in miR-2 mutants. (C-D) Data are from small
RNA sequencing data of L4-staged mutants that were created using the (left) Nextflex v3 or
(right) Nextflex v4 small RNA sequencing kits. (A-D) Data are presented as the percent
occurrence of a given isomiR within each individual mutant and do not reflect possible changes
in miRNA abundance. Each isomiR is illustrated on the left. The maroon (guide) and gray
(passenger) bars indicates the expected miRNA sequence. Truncated isomiRs are indicated by
negative values corresponding to the length of the truncation. Nucleotide extensions are
depicted by the corresponding letters to each duplex end, which may include templated or non-
templated (uridylation and adenylation) extensions. Note that pre-miR-2 contains a templated
uracil immediately downstream of the 3' nucleotide of the mature miR-2 guide and passenger
strands, preventing disambiguation of non-templated or templated addition of uracil to either
strand.
747
Extended Data Figure 7. Analysis of miR-48 and miR-71 isomiRs in miRNA mutants.
Heatmaps illustrate relative distribution of (A) miR-48-5p (canonical guide strand), (B) miR-48-
3p (canonical passenger strand), (C) miR-71-5p (canonical guide strand) and (D) miR-71-3p
(canonical passenger strand) in miR-48 or miR-71 mutants. (A-D) Data are presented as the
percent occurrence of a given isomiR within each individual mutant and do not reflect possible
changes in miRNA abundance. Each isomiR is illustrated on the left. The maroon (guide) and
gray (passenger) bars indicates the expected miRNA sequence. Truncated isomiRs are
indicated by negative values corresponding to the length of the truncation. Nucleotide
extensions are depicted by the corresponding letters to each duplex end, which may include
templated or non-templated (uridylation and adenylation) extensions. Note that pre-miR-48
contains a templated adenine immediately downstream of the 3' nucleotide of the mature miR-
48 guide strand and a templated 3' uracil downstream immediately downstream of passenger
strand, preventing disambiguation of non-templated or templated additions. Similarly, pre-miR-
71 contains a templated uracil immediately downstream of the 3' nucleotide of the mature miR-
71 guide strand. (E-F) (Bottom) Schematic representations of (E) miR-48 and (F) miR-71 duplex
mutations that affect 3' nucleotide identity. The guide strand is shown in maroon, and the
passenger strand is shown in gray. The nucleotide composition of dinucleotide overhangs are

shown for each strand. Mutated nucleotides are indicated by red text. U:A pairs are illustrated
by an open circle, G:U wobble pairs are indicated by an open circle with a vertical line and G:C
pairs are represented by closed circles. RNAcofold was used to predict duplex structures at
20°C. (Top) Quantification of small RNA sequencing results for (E) miR-48 and (B) miR-71.
Data are presented as mean ± standard deviation, with each dot representing quantification
from a single library using the NEXTFLEX v4 Small RNA Sequencing Kit. Maroon bars
correspond to miRNA guide strands and gray bars represent miRNA passenger strands.
Normalization was performed by dividing raw read counts by total mapped miRNA reads and
multiplying by one million to acquire reads per million (rpm) values.
774
Extended Data Figure 8. The composition of dinucleotide overhangs may influence miRNA
strand selection. (A-B) (Left) Schematic representations of miR-58 duplex mutations that affect
the nucleotide composition of dinucleotide overhangs. Mutations were made in a (A) wildtype
miR-58 duplex background or (B) the miR-58-M7 mutant duplex. The guide strand is shown in
maroon, and the passenger strand is shown in gray. Mutated nucleotides are indicated by red
text. U:A pairs are illustrated by an open circle, G:U wobble pairs are indicated by an open circle
with a vertical line and G:C pairs are represented by closed circles. RNAcofold was used to
predict duplex structures at 20°C. (Right) Quantification of small RNA sequencing results for
each miR-58 duplex strand. Data are presented as mean ± standard deviation, with each dot
representing quantification from a single library using the NEXTFLEX v3 Small RNA
Sequencing Kit. Maroon bars correspond to miRNA guide strands and gray bars represent
miRNA passenger strands. Normalization was performed by dividing raw read counts by total
mapped miRNA reads and multiplying by one million to acquire reads per million (rpm) values.
788
Extended Data Figure 9. Functional analysis of ALG-1[7A] mutants. (A) Location of alanine
substitutions in ALG-1[7A] mutants. Mutations are indicated by red text. The location of the ALG-1
5' nucleotide binding pocket[13,14] and putative 3' nucleotide binding pocket[17] are bolded. (B)
Physical location of 5' nucleotide binding pocket (yellow) and putative 3' nucleotide binding
pocket (red) on the Alphafold predicted structure of ALG-1 (AF-Q3LTR7-F1). The MID (dark
purple) and PIWI (light purple) domains are marked. ChimeraX v1.9 was used for structure
visualization. Inset highlights relative position of the 5' nucleotide binding pocket and putative 3'
nucleotide binding pocket within the RNA-binding groove of ALG-1 (B') and rotated 180° (B").
(C) Differential expression analysis of ALG-1[7A] mutants compared to wildtype (N2) controls was
performed using the 'DESeq2' package for R. Each dot represents one miRNA strand.
Statistical significance was considered at p<0.01. Upregulated miRNAs are labelled and colored
pink while downregulated miRNAs are labelled and colored blue.
801
Extended Data Figure 10. Analysis of isomiRs from miRNAs transfected into human HEK293T
cells. Heatmaps illustrate relative distribution of (A) miR-58a-3p (*C. elegans* guide strand), (B)
miR-58a-5p (*C. elegans* passenger strand), (C) miR-339-syn-3p (expected guide strand) and
(D) miR-339-syn-5p (expected passenger strand) in miR-58 or miR-339-syn mutants. The miR-
339-syn mutants carry an additional internal nucleotide substitution that is not expected to
affected duplex structure (Supplementary Table 15), which allowed us to distinguish exogenous
and endogenous miR-339 variants. (A-D) Data are presented as the percent occurrence of a
given isomiR within each individual mutant and do not reflect possible changes in miRNA
abundance. Each isomiR is illustrated on the left. The maroon (guide) and gray (passenger)
bars indicates the expected miRNA sequence. Truncated isomiRs are indicated by negative
values corresponding to the length of the truncation. Nucleotide extensions are depicted by the
corresponding letters to each duplex end, which may include templated or non-templated
(uridylation and adenylation) extensions. Note that the Ultramir scaffold contains a templated

uracil immediately downstream of the 3' nucleotide of the 3p, preventing disambiguation of non-
templated or templated uridylation for the 3p strand.
817
818   **Supplementary Figure 1.** ZsGreen expression in transfected human HEK293T cells.
819   Representative images of HEK293T cells expressing ZsGreen prior to harvesting. (A) Images of
820   samples corresponding to Figure 6B. (B) Images of samples corresponding to Figure 6C.

References

1. Kim, H., Lee, Y.-Y. & Kim, V. N. The biogenesis and regulation of animal microRNAs. *Nat. Rev. Mol. Cell Biol.* 1–21 (2024) doi:10.1038/s41580-024-00805-0.

2. Shang, R., Lee, S., Senavirathne, G. & Lai, E. C. microRNAs in action: biogenesis, function and regulation. *Nat. Rev. Genet.* **24**, 816–833 (2023).

3. Denli, A. M., Tops, B. B. J., Plasterk, R. H. A., Ketting, R. F. & Hannon, G. J. Processing of primary microRNAs by the Microprocessor complex. *Nature* **432**, 231–235 (2004).

4. Gregory, R. I. *et al.* The Microprocessor complex mediates the genesis of microRNAs. *Nature* **432**, 235–240 (2004).

5. Lee, Y., Jeon, K., Lee, J., Kim, S. & Kim, V. N. MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J.* **21**, 4663–4670 (2002).

6. Lee, Y. *et al.* The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**, 415–419 (2003).

7. Grishok, A. *et al.* Genes and Mechanisms Related to RNA Interference Regulate Expression of the Small Temporal RNAs that Control C. elegans Developmental Timing. *Cell* **106**, 23–34 (2001).

8. Hutvágner, G. *et al.* A Cellular Function for the RNA-Interference Enzyme Dicer in the Maturation of the let-7 Small Temporal RNA. *Science* **293**, 834–838 (2001).

9. Jiang, F. *et al.* Dicer-1 and R3D1-L catalyze microRNA maturation in Drosophila. *Genes Dev.* **19**, 1674–1679 (2005).

10. Ketting, R. F. *et al.* Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in C. elegans. *Genes Dev.* **15**, 2654–2659 (2001).

11. Knight, S. W. & Bass, B. L. A Role for the RNase III Enzyme DCR-1 in RNA Interference and Germ Line Development in Caenorhabditis elegans. *Science* **293**, 2269–2271 (2001).

12. Saito, K., Ishizuka, A., Siomi, H. & Siomi, M. C. Processing of Pre-microRNAs by the Dicer-1–Loquacious Complex in Drosophila Cells. *PLoS Biol.* **3**, e235 (2005).

13. Elkayam, E. *et al.* The Structure of Human Argonaute-2 in Complex with miR-20a. *Cell* **150**, 100–110 (2012).

14. Schirle, N. T. & MacRae, I. J. The Crystal Structure of Human Argonaute2. *Science* **336**, 1037–1040 (2012).

15. Nykänen, A., Haley, B. & Zamore, P. D. ATP Requirements and Small Interfering RNA Structure in the RNA Interference Pathway. *Cell* **107**, 309–321 (2001).

16. Schwarz, D. S. *et al.* Asymmetry in the Assembly of the RNAi Enzyme Complex. *Cell* **115**,
199–208 (2003).

17. Suzuki, H. I. *et al.* Small-RNA asymmetry is directly driven by mammalian Argonautes. *Nat*
Struct Mol Biol **22**, 512–521 (2015).

18. Medley, J. C., Panzade, G. & Zinovyeva, A. Y. microRNA strand selection: Unwinding the
rules. *Wiley Interdiscip. Rev.: RNA* **12**, e1627 (2021).

19. Humphreys, D. T. *et al.* Complexity of Murine Cardiomyocyte miRNA Biogenesis, Sequence
Variant Expression and Function. *PLoS ONE* **7**, e30933 (2012).

20. Okamura, K. *et al.* The regulatory activity of microRNA* species has substantial influence on
microRNA and 3′ UTR evolution. *Nat. Struct. Mol. Biol.* **15**, 354–363 (2008).

21. Ghildiyal, M., Xu, J., Seitz, H., Weng, Z. & Zamore, P. D. Sorting of Drosophila small
silencing RNAs partitions microRNA* strands into the RNA interference pathway. *RNA* **16**, 43–
56 (2010).

22. Kuchen, S. *et al.* Regulation of MicroRNA Expression and Abundance during
Lymphopoiesis. *Immunity* **32**, 828–839 (2010).

23. Ro, S., Park, C., Young, D., Sanders, K. M. & Yan, W. Tissue-dependent paired expression
of miRNAs. *Nucleic Acids Res.* **35**, 5944–5953 (2007).

24. Yang, J.-S. *et al.* Widespread regulatory activity of vertebrate microRNA* species. *RNA* **17**,
312–326 (2011).

25. Frank, F., Sonenberg, N. & Nagar, B. Structural basis for 5′-nucleotide base-specific
recognition of guide RNA by human AGO2. *Nature* **465**, 818–822 (2010).

26. Khvorova, A., Reynolds, A. & Jayasena, S. D. Functional siRNAs and miRNAs Exhibit
Strand Bias. *Cell* **115**, 209–216 (2003).

27. Warf, M. B., Johnson, W. E. & Bass, B. L. Improved annotation of C. elegans microRNAs by
deep sequencing reveals structures associated with processing by Drosha and Dicer. *RNA* **17**,
563–577 (2011).

28. Medley, J. C. & Zinovyeva, A. In silico unwinding of Caenorhabditis elegans microRNA
duplexes to evaluate thermodynamic end stabilities improves predictions of microRNA strand
selection. *bioRxiv* 2025.06.26.661848 (2025) doi:10.1101/2025.06.26.661848.

29. Panzade, G., Li, L., Hebbar, S., Veksler-Lublinsky, I. & Zinovyeva, A. Global profiling and
annotation of templated isomiRs dynamics across Caenorhabditis elegans development. *RNA*
Biol. **19**, 928–942 (2022).

30. Alvarez-Saavedra, E. & Horvitz, H. R. Many Families of C. elegans MicroRNAs Are Not
Essential for Development or Viability. *Curr Biol* **20**, 367–373 (2010).

31. Simon, D. J. *et al.* The MicroRNA miR-1 Regulates a MEF-2-Dependent Retrograde Signal
at Neuromuscular Junctions. *Cell* **133**, 903–915 (2008).

32. O'Hern, P. J. *et al.* Decreased microRNA levels lead to deleterious increases in neuronal M2
muscarinic receptors in Spinal Muscular Atrophy models. *eLife* **6**, e20752 (2017).

33. Lingel, A., Simon, B., Izaurralde, E. & Sattler, M. Nucleic acid 3′-end recognition by the
Argonaute2 PAZ domain. *Nat. Struct. Mol. Biol.* **11**, 576–577 (2004).

34. Wang, Y., Sheng, G., Juranek, S., Tuschl, T. & Patel, D. J. Structure of the guide-strand-
containing argonaute silencing complex. *Nature* **456**, 209–13 (2008).

35. Ameres, S. L. & Zamore, P. D. Diversifying microRNA sequence and function. *Nat. Rev.*
Mol. Cell Biol. **14**, 475–488 (2013).

36. Han, J. & Mendell, J. T. MicroRNA turnover: a tale of tailing, trimming, and targets. *Trends*
Biochem. Sci. **48**, 26–39 (2023).

37. Kozomara, A., Birgaoanu, M. & Griffiths-Jones, S. miRBase: from microRNA sequences to
function. *Nucleic Acids Res.* **47**, D155–D162 (2018).

38. Creugny, A., Fender, A. & Pfeffer, S. Regulation of primary microRNA processing. *FEBS*
Lett. **592**, 1980–1996 (2018).

39. Han, J. *et al.* The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev.*
18, 3016–3027 (2004).

40. Knott, S. R. V. *et al.* A Computational Algorithm to Predict shRNA Potency. *Mol. Cell* **56**,
796–807 (2014).

41. Newman, M. A., Mani, V. & Hammond, S. M. Deep sequencing of microRNA precursors
reveals extensive 3′ end modification. *RNA* **17**, 1795–1803 (2011).

42. Wyman, S. K. *et al.* Post-transcriptional generation of miRNA variants by multiple nucleotidyl
transferases contributes to miRNA transcriptome complexity. *Genome Res* **21**, 1450–61 (2011).

43. Fernández-Pérez, D., Brieño-Enríquez, M. A., Isoler-Alcaraz, J., Larriba, E. & Mazo, J. del.
MicroRNA dynamics at the onset of primordial germ and somatic cell sex differentiation during
mouse embryonic gonad development. *Rna* **24**, 287–303 (2018).

44. Fernandez-Valverde, S. L., Taft, R. J. & Mattick, J. S. Dynamic isomiR regulation in
Drosophila development. *RNA* **16**, 1881–1888 (2010).

45. Oliveira, A. C. de *et al.* MicroRNA Transcriptomes Reveal Prevalence of Rare and Species-
Specific Arm Switching Events During Zebrafish Ontogenesis. *Evol. Bioinform. Online* **20**,
11769343241263230 (2024).

46. Londin, E. *et al.* Analysis of 13 cell types reveals evidence for the expression of numerous
novel primate- and tissue-specific microRNAs. *Proc. Natl. Acad. Sci.* **112**, E1106–E1115 (2015).

47. Smith, C. M. & Hutvagner, G. A comparative analysis of single cell small RNA sequencing
data reveals heterogeneous isomiR expression and regulation. *Sci. Rep.* **12**, 2834 (2022).

48. Telonis, A. G. *et al.* Knowledge about the presence or absence of miRNA isoforms
(isomiRs) can successfully discriminate amongst 32 TCGA cancer types. *Nucleic Acids Res.* **45**,
2973–2985 (2017).

49. Vieux, K.-F. *et al.* Screening by deep sequencing reveals mediators of microRNA tailing in
C. elegans. *Nucleic Acids Res* **49**, gkab840- (2021).

50. Burroughs, A. M. *et al.* A comprehensive survey of 3′ animal miRNA modification events and
a possible role for 3′ adenylation in modulating miRNA targeting effectiveness. *Genome Res.*
20, 1398–1410 (2010).

51. Heo, I. *et al.* Lin28 Mediates the Terminal Uridylation of let-7 Precursor MicroRNA. *Mol. Cell*
32, 276–284 (2008).

52. Heo, I. *et al.* Mono-Uridylation of Pre-MicroRNA as a Key Step in the Biogenesis of Group II
let-7 MicroRNAs. *Cell* **151**, 521–532 (2012).

53. Kim, H. *et al.* A Mechanism for microRNA Arm Switching Regulated by Uridylation. *Mol Cell*
78, 1224-1236.e5 (2020).

54. Song, J. *et al.* Prevalent cytidylation and uridylation of precursor miRNAs in Arabidopsis.
Nat. Plants **5**, 1260–1272 (2019).

55. Kwak, P. B. & Tomari, Y. The N domain of Argonaute drives duplex unwinding during RISC
assembly. *Nat Struct Mol Biol* **19**, 145–151 (2012).

56. Duan, Y. *et al.* Modeling neurodevelopmental disorder-associated human AGO1 mutations
in Caenorhabditis elegans Argonaute alg-1. *Proc. Natl. Acad. Sci.* **121**, e2308255121 (2024).

57. Zinovyeva, A. Y., Veksler-Lublinsky, I., Vashisht, A. A., Wohlschlegel, J. A. & Ambros, V. R.
Caenorhabditis elegans ALG-1 antimorphic mutations uncover functions for Argonaute in
microRNA guide strand selection and passenger strand disposal. *Proc. Natl. Acad. Sci.* **112**,
E5271–E5280 (2015).

58. Kim, Y. *et al.* Deletion of human tarbp2 reveals cellular microRNA targets and cell-cycle
function of TRBP. *Cell Reports* **9**, 1061–74 (2014).

59. Noland, C. L. & Doudna, J. A. Multiple sensors ensure guide strand selection in human
RNAi pathways. *RNA* **19**, 639–648 (2013).

60. Wilson, R. C. *et al.* Dicer-TRBP Complex Formation Ensures Accurate Mammalian
MicroRNA Biogenesis. *Mol. Cell* **57**, 397–407 (2015).

61. Brenner, S. The Genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

955 62. Hibshman, J. D., Webster, A. K. & Baugh, L. R. Liquid-culture protocols for synchronous
956 starvation, growth, dauer formation, and dietary restriction of Caenorhabditis elegans. *STAR*
957 *Protoc.* **2**, 100276 (2021).

958 63. Porta-de-la-Riva, M., Fontrodona, L., Villanueva, A. & Cerón, J. Basic Caenorhabditis
959 elegans methods: synchronization and observation. *J. Vis. Exp. : JoVE* e4019 (2012)
960 doi:10.3791/4019.

961 64. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol. Biol. : AMB* **6**, 26 (2011).

962 65. Paix, A., Folkmann, A., Rasoloson, D. & Seydoux, G. High Efficiency, Homology-Directed
963 Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein Complexes.
964 *Genetics* **201**, 47–54 (2015).

965 66. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
966 *EMBnetJ.* **17**, 10–12 (2011).

967 67. Andrews. FastQC: A Quality Control Tool for High Throughput Sequence Data. *Available*
968 *online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/* (2010).

969 68. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient
970 alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25–R25 (2009).

971 69. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
972 2079 (2009).

973 70. Donnelly, B. F. *et al.* The developmentally timed decay of an essential microRNA family is
974 seed-sequence dependent. *Cell Rep.* **40**, 111154–111154 (2022).

A

miR
miR*

U:A pair ○
G:U pair ◑
G:C pair ●

Favorable 5'nt
Lower Stability

Unfavorable 5'nt
Higher Stability

5' nucleotide: U>A>C>G

B

miRNA Count (n=190)

■ miR ■ miR*

5' Nucleotide Identity

C

Average ΔG (kcal/mol)

■ miR ■ miR*

Terminal Nucleotides

Figure 1

Figure 3

Figure 4

Figure 5

Figure 6

Extended Data Figure 2

Extended Data Figure 5

Extended Data Figure 6

Extended Data Figure 7

A

	5' nt (guide)	3' nt (passenger)

ALG-1 PMFKYLKQNYSGIQLVVVVL**PGKTP**VYAEVKRVGDTVLGIATQCVQAKNAI**RTTPQTL**SNLCLKMNVKLGGVNSI – 720
ALG-1⁷ᴬ PMFKYLKQNYSGIQLVVVVL**PGKTP**VYAEVKRVGDTVLGIATQCVQAKNAI**AAAAAAA**SNLCLKMNVKLGGVNSI – 720

ALG-1⁷ᴬ: R697A, T698A, T699A, P700A, Q701A, T702A, L703A

B

ALG-1 (AF-Q3LTR7-F1)

B' B''

C

Extended Data Figure 9

A

GFP siRNA control	**hsa-miR-339-syn** (pJM1)	**cel-miR-58α-M6** (pJM2)	**cel-miR-58α-M16** (pJM3)
hsa-miR-339-syn-5pC (pJM4)	**hsa-miR-339-syn-3pC** (pJM5)	**cel-miR-58α** (pJM6)	**cel-miR-58α-M14** (pJM7)

B

miR-58-M6_5pC-3pU (pJM2)	**miR-58-M6_5pU-3pC** (pJM3)	**miR-58-M6_5pA-3pU** (pJM10)	**miR-58-M6_5pU-3pU** (pJM11)
miR-58-M6_5pG-3pU (pJM12)	**miR-58-M6_5pC-3pA** (pJM13)	**miR-58-M6_5pA-3pA** (pJM14)	**miR-58-M6_5pU-3pA** (pJM15)
miR-58-M6_5pG-3pA (pJM16)	**miR-58-M6_5pG-3pU** (pJM17)	**miR-58-M6_5pA-3pG** (pJM18)	**miR-58-M6_5pU-3pG** (pJM19)
miR-58-M6_5pG-3pG (pJM20)	**miR-58-M6_5pA-3pC** (pJM21)	**miR-58-M6_5pC-3pC** (pJM22)	**miR-58-M6_5pG-3pC** (pJM23)

Supplementary Figure 1