1 **Repeated Disuse Atrophy Imprints a Molecular Memory in Skeletal Muscle:**

2 **Transcriptional Resilience in Young Adults and Susceptibility in Aged Muscle**

7 Daniel C. Turner[1,2]*, Truls Raastad[1], Max Ullrich[1], Stian F. Christiansen[1], Hazel Sutherland[3], James

8 Boot[4,5], Eva Wozniak[4], Charles Mein[4], Emilie Dalbram[6], Jonas T. Treebak[6], Daniel J. Owens[3], David C.

9 Hughes[7], Sue C. Bodine[7], Jonathan C. Jarvis[3], Adam P. Sharples[1]*.

[1] Department for Physical Performance, Norwegian School of Sport Sciences, Oslo, Norway.

[2] Department of Molecular Medicine, University of Pavia, Italy.

[3] Research Institute for Sport and Exercise Sciences, Liverpool John Moores University (LJMU),

Liverpool, United Kingdom.

[4] Genome Centre, Queen Mary University of London (QMUL), London, United Kingdom.

[5] Francis Crick Institute, London, United Kingdom.

[6] Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical

Sciences, University of Copenhagen (UCPH), Copenhagen, Denmark.

[7] Aging and Metabolism Research Program, Oklahoma Medical Research Foundation (OMRF),

Oklahoma City, Oklahoma, United States.

37 *Corresponding authors

38 Adam P. Sharples: adams@nih.no

39 Daniel C. Turner: daniel.turner@unipv.it

Abstract

Disuse-induced muscle atrophy is common after illness, injury, or falls, and becomes increasingly frequent with ageing. However, whether skeletal muscle retains a "memory" of disuse remains unknown. We investigated repeated lower-limb immobilization in young adults and a refined aged rat model of repeated disuse, integrating physiology, multi-omics, immunohistochemistry, biochemistry, and primary human cell analyses. In young human muscle, a second period of disuse elicited an attenuated transcriptional response in oxidative and mitochondrial pathways, suggesting a protective molecular memory, although this did not translate into attenuated atrophy. In contrast, aged muscle exhibited a detrimental molecular memory, with greater atrophy, exaggerated suppression of aerobic metabolism genes/pathways despite recovery of these genes after initial disuse, depletion of NAD^+ and mitochondrial DNA, and activation of proteasomal, extracellular matrix and DNA-damage pathways. Repeated disuse induced DNA hypermethylation and downregulation of aerobic metabolism and mitochondrial genes across species. *NR4A1* remained suppressed into recovery after initial disuse in young human muscle, whereas acetylcholine receptor subunit genes (e.g., *CHRNA1*, *CHRND*) were epigenetically primed in humans and further upregulated with repeated atrophy in both young and aged muscle. *NMRK2,* an NAD^+ biosynthesis gene, was among the most downregulated genes after both periods of atrophy in humans, and nicotinamide riboside (NR) improved myotube size in primary human muscle cells derived post-atrophy. These findings reveal that disuse atrophy imprints a molecular memory in skeletal muscle - attenuated transcriptional resilience in young adults yet exaggerated susceptibility in aged muscle.

Keywords: muscle memory, DNA methylation, aerobic metabolism, NAD^+ metabolism, *NR4A* family, AChR, *NMRK2*, nicotinamide riboside

Introduction

Age-related skeletal muscle loss (termed sarcopenia) is one of the main contributors to falling injury [1, 2, 3], often resulting in prolonged periods of disuse. Consequently, repeated falls in the elderly become increasingly common, leading to recurrent loss of skeletal muscle and thus increased risk of frailty, morbidity, hospital admittance, reduced quality of life and ultimately earlier mortality [2, 4, 5, 6, 7]. Restoration of skeletal muscle following periods of disuse is also impaired in aged muscle [5, 8, 9, 10, 11, 12], further increasing the risk of repeated falls [2, 3]. Despite the well-characterised skeletal muscle wasting (atrophy) response to a single period of limb immobilisation [8, 9, 10, 13, 14, 15, 16, 17, 18, 19, 20, 21], bed rest [22, 23, 24, 25, 26] and spaceflight [27, 28, 29], whether muscle atrophy is exacerbated upon repeated disuse in both adult and aged muscle is currently unknown.

Skeletal muscle "memory" refers to the differential tissue response to repeated exposure to the same (or similar) environmental stimuli [30, 31]. While it is evident that skeletal muscle exhibits a 'positive' memory of anabolic stimuli, resistance [32, 33, 34, 35, 36, 37, 38, 39] and endurance [34, 40] training in both humans and rodents, it is unknown whether skeletal muscle possesses a ´negative´ memory of disuse-induced muscle atrophy. Specifically, whether an earlier encounter of disuse renders the muscle more susceptible to further wasting if reencountered in the future following injury reoccurrence or repeated falling in the elderly.

Epigenetics is considered a key mechanism underpinning skeletal muscle memory, whereby retention of epigenetic modifications in skeletal muscle are either; a) modified following exercise training and retained during detraining, leading to sustained elevations in gene transcription [33, 34, 36, 37, 39, 40] and protein abundance [38] and/or; b) enhanced upon later retraining, resulting in amplified changes in gene expression [33, 34, 36, 37, 39, 40]. Among the various epigenetic modifications, DNA methylation is one of the most prominent emerging regulators of skeletal muscle memory. DNA methylation undergoes genome-wide alterations and modulates the transcriptional response to acute exercise and chronic training in divergent pathways according to the mode of exercise performed [33, 36, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55]. Moreover, DNA methylation is dynamic and can both precede and modulate the time course of transient gene expression in skeletal muscle in the minutes to hours following a single bout of exercise (reviewed in [56]). However, some DNA methylation changes are long-lasting, particularly those induced by chronic training, thus providing a key mechanism mediating skeletal muscle memory of exercise [31, 33, 34, 36, 37, 39, 40].

Despite the epigenetic role in modulating gene transcription in response to positive stimuli, studies examining the DNA methylation response to negative stimuli such as muscle disuse atrophy remain scarce. Nevertheless, there is clear evidence of a dysregulated methylome in skeletal muscle of aged [57, 58, 59, 60, 61] and diseased populations (e.g., type-II diabetes [42, 51, 62] and cancer [48]), in which greater aerobic fitness (higher $\dot{V}O_{2max}$) [57], higher physical activity, or exercise training can recalibrate the nuclear [39, 48, 57, 58, 59] and mitochondrial methylome [63] toward epigenetic signatures resembling that of youthful

muscle. Following disuse atrophy, hypermethylation and reduced expression of targeted genes such as *PPARGC1A* after nine days of bed rest in human muscle [64], and *Nos1* after acute immobilization in mice [65] have been reported. Our group have also demonstrated hypomethylation and increased expression of E3 ligases (*Murf1* and *MAFbx/Fbxo32*), the acetylochline receptor subunit (*Chrna1*), and the histone deacetylase (*Hdac4*) following tetrodotoxin (TTX)-included disuse atrophy in young adult rat muscle [66]. To date, however, most studies have predominantly focussed on targeted gene methylation with limited data available on genome-wide methylation. A recent study combined methylome and transcriptome analyses after 2 weeks of disuse atrophy in young adult humans (via lower limb casting) and reported predominance of global hypermethylation that was associated with a larger number of downregulated versus upregulated genes [39]. However, leg casting-induced disuse was preceded by resistance training, and therefore it is unknown whether the such alterations occur following disuse atrophy without prior resistance training that is more representative of the general population [39]. A recent study analysing genome-wide DNA methylation in purified muscle stem cells (MuSC) following injury in adult mice demonstrated sustained differential methylation long after recovery [67], whereas myoblasts retain DNA methylation changes after repeated inflammatory exposure *in vitro* [68]. However, no studies have investigated the combined methylome and transcriptome following repeated atrophy to determine whether skeletal muscle imprints a ´negative´ memory of disuse-induced atrophy in both young adult and aged muscle.

Therefore, we sought to determine whether skeletal muscle exhibits a 'negative memory' of disuse atrophy. We hypothesized that prior episodes of disuse atrophy would induce persistent molecular imprints that exacerbate subsequent repeated atrophy, particularly in aged muscle. Due to frailty and therefore ethical issues associated with repeated atrophy in elderly humans; to test this, we combined comprehensive physiological, histological and molecular analyses in parallel young adult human and aged animal models of disuse atrophy, recovery and repeated atrophy. Here, we demonstrate that skeletal muscle retained a molecular memory of disuse. In young adults, repeated immobilization triggered a predominantly resilient and 'protective' transcriptional memory, characterised by attenuated transcriptional changes in aerobic metabolism and mitochondrial pathways following repeated disuse, despite full restoration of skeletal muscle after recovery. However, this did not lead to attenuated muscle loss after repeated atrophy. In contrast, aged muscle exhibited a detrimental susceptibility, characterised by an exaggerated suppression of aerobic metabolism and mitochondrial pathways, mtDNA loss, and muscle wasting despite recovery of these gene pathways after initial disuse. Proteasomal, extracellular matrix (ECM), and DNA-damage associated pathways were uniquely elevated in aged muscle to repeated disuse. Epigenomic profiling revealed conserved DNA hypermethylation in aerobic and mitochondrial gene networks across species to atrophy and repeated atrophy, alongside identification of locus-specific memory genes, *NR4A1* and AChR subunit genes. Finally, NAD$^+$ salvage with nicotinamide riboside (NR) improved myotube size in human primary MuSC derived post-atrophy.

Methods

Human ethics and participant information

Ethical approval was granted by the Regional Committees for Medical and Health Research Ethics

(REK, Application ID 227924). The study was registered with the Norwegian Agency for Shared

Services in Education and Research (Sikt) under reference number 627824. Participants were

familiarised with the study intervention and the possible associated risks. Informed written consent and

completion of an approved health and pre-biopsy medical questionnaire were obtained prior to

participation. Ten young adults, males (n = 7) and females (n = 3) living in Oslo, Norway participated in

the study. Participants were free from any known injury, illness and disease and had no previous history

or predisposition to deep vein thrombosis (DVT). Participant baseline characteristics are presented in

Table 1. All human data collection was conducted at the Norwegian School of Sport Sciences.

Table 1. Young adult human baseline characteristics. Values presented as mean ± standard deviation

(SD).

	Male (n = 7)	**Female** (n = 3)	**All** (n = 10)
Age (yrs)	27.1 ± 4.1	23.7 ± 3	26.5 ± 4.1
Height (cm)	181.7 ± 2.7	167.9 ± 7.5	177.5 ± 7.9
Weight (kg)	93.4 ± 16	62 ± 12.6	84 ± 20.9
Body mass index, BMI (kg/m^2)	28.4 ± 5.3	21.8 ± 2.6	26.4 ± 5.5
Total Lean body mass, LBM (kg)	64.8 ± 4.4	41.5 ± 7.2	57.8 ± 12.3
Fat mass, FM (kg)	24.8 ± 13.1	18.3 ± 5.7	22.3 ± 11.4
Fat free mass, FFM (kg)	68.2 ± 4.5	43.9 ± 7.3	60.9 ± 12.8
Body fat, BF (%)	25.4 ± 9.4	29.1 ± 4.5	26.5 ± 8.2
Immobilised leg lean mass, LM (kg)	11.4 ± 1.2	7.4 ± 1.5	10.2 ± 2.3
Non-immobilised leg LM (kg)	11.5 ± 1.3	7.1 ± 1.6	8.9 ± 2.5

Assessment of physical activity levels and dietary intake in humans

The International Physical Activity Questionnaire Short Form (IPAQ-SF) was used to assess short-term

physical activity levels prior to the study [68]. Participants also completed an additional non-validated

questionnaire including questions related to the specific type of sport (e.g., team, endurance,

speed/power) or activity (e.g. cardiorespiratory fitness, strength, concurrent), duration of participation

and competition level (i.e. recreational, national, international) to assess long-term physical activity

levels.

Human study design

In a repeated measures design, participants underwent 2 weeks (wks) (14 days/d) disuse atrophy (via

unilateral limb immobilization) → ≈ 7 wks (49 ± 5 d) recovery → 2 wks repeated disuse atrophy (**Figure**

1A). During immobilisation, participants were fitted with a knee brace (DonJoy® X-Act ROM,
Enovis/DJO LLC, USA) fixed at 30° knee flexion to prevent any weight bearing of the non-dominant leg.
Crutches were provided to enable ambulation without weight bearing on the braced immobilised leg.
Following immobilisation, the knee brace and crutches were removed and participants returned to their
normal ambulatory activity for ≈ 7 wks to allow recovery. Finally, the same non-dominant leg underwent
a second period of 2 wks of immobilisation (repeated atrophy). Before immobilisation, participants were
trained using the crutches and knee brace and were instructed to only remove the knee brace for
showering and bedtime. Daily exercises (e.g., ankle rotation/dorsiflexion/plantarflexion) were
encouraged to reduce the risk of DVT[69] . There were no reported cases throughout the present study.
Before and after each immobilization, muscle biopsies and measurements of SkM size and strength
were conducted in the morning (± 1 hr between timepoints) under fasted conditions following the same
testing order (**Figure 1A**). Participants were also advised to refrain from strenuous exercise and alcohol
consumption 24 hrs prior to muscle biopsy and testing procedures. Muscle biopsies and the testing
protocol are detailed below. Upon completion of the study intervention, for ethical reasons, all
participants were provided with the opportunity to undertake supervised resistance training programme
(3 d/wk for 7 wks consisting of 2 lower body and 1 upper body sessions each wk) to restore any loss of
SkM. Two-four weeks of strength training is sufficient for restoring SkM following a similar duration of
atrophy in young adult humans [8, 11, 70].

Human skeletal muscle biopsy procedures

Muscle biopsies were obtained from the immobilised leg according to our procedures described
elsewhere [48, 49]. Briefly, local anaesthesia (Xylocaine with 10 mg/mL lidocaine + 5 µg/mL adrenaline,
AstraZeneca, Södertälje, Sweden) was administered to the biopsy area and muscle tissue (133 ± 47
mg excluding muscle tissue for immunohistochemical analysis, see below) was obtained from the mid-
belly (10% proximal of the mid-thigh) of the *m. vastus lateralis (VL)* muscle using a modified Bergström
technique (6 mm Pelomi-needle, Albertslund, Denmark) with manual suction (i.e., 50 mL syringe
connected to the Bergström needle). For immunohistochemical (IHC) analysis, a proportion (≈ 50 mg)
of muscle tissue was embedded in optimal cutting temperature (OCT) compound (CellPath, Chemi-
Teknik AS, Oslo, Norway) ensuring myofibres were aligned in parallel before freezing in pre-cooled
(i.e., above liquid nitrogen, LN_2) isopentane (Sigma-Aldrich, Oslo, Norway) placed on dry ice and stored
at −80 °C. Remaining tissue was rinsed in pre-cooled (4 °C) saline containing 0.9% NaCl (Braun,
Melsungen, Germany) and dissected to remove visible fat, connective tissue and blood (vessels).
Tissue intended for RNA, DNA and protein analysis was immediately weighed, transferred to nuclease-
free Eppendorf tubes (Invitrogen™ RNase-free Microfuge Tubes, Fisher Scientific, Oslo, Norway),
snap-frozen in LN_2 and stored at −80 °C. Remaining tissue used for *in vitro* experiments was submerged
in pre-cooled (4 °C) transfer media (Hams F-10 medium that includes 1 mM L-glutamine/LG, 0.2% heat
inactivated/hiFBS, 100 U/mL penicillin, 100 µg/mL streptomycin/PS and 2.5 µg/mL amphotericin-B/AB,
Fisher Scientific) for subsequent primary cell isolations detailed below.

Assessment of body composition in humans

Height (cm) and body weight (kg) were measured using a digital stadiometer and scale, respectively (Seca, Hamburg, Germany). Whole-body composition and leg lean mass (LLM) were assessed by dual-energy x-ray absorptiometry (DEXA; Lunar iDXA, enCORE software version 18; GE Healthcare, Chicago, USA). Total LLM was defined as all lean mass located distally from the collum femoris and inferior pubic ramus. A triangular segment depicting the pelvic area and a midline separating left and right legs were constructed as previously described [71]. Specifically, the horizontal line of the triangle segment transversed slightly above the two iliac crests, and the two angled lines crossed the collum femoris, meeting just below the genital area [71]. An additional line crossing the intercondylar eminence separated the upper and lower leg. All images were analysed single-blinded by three experienced personnel, and the average values were reported. The coefficient of variation (CV) for test-retest reliability was 1%.

Assessment of muscle cross-sectional area (mCSA) in humans

Muscle cross-sectional area (mCSA) of the *m. rectus femoris (RF)* and *VL* muscles were assessed using panoramic B-mode ultrasound scanning using a linear transducer (L18-5) fixed to an ultrasound system (MACH 30; Hologic SuperSonic Imagine™, Aix-en-Provence, France). During scanning, participants lay supine (knee angle at 0 °C) with their leg muscles relaxed and feet strapped to a stable box to prevent any hip rotation. Three measurements per muscle (*RF* and *VL* each scanned separately) were measured and analysed by following a transverse trajectory line drawn across the mid-thigh (i.e., 50% of the distance between the greater trochanter and distal end of the muscle) using a marker pen, demonstrating a low <2.5% CV across all timepoints (RF baseline 2.4 ± 1.1%, atrophy 1.6 ± 1%, recovery 2.1 ± 1.1%, repeated atrophy 1.8 ± 1.6%, VL baseline 1.7 ± 1.3%, atrophy 1.3 ± 0.7%, 1.8 ± 1% recovery, 1.6 ± 0.7% repeated atrophy). Only acquired images free from visible artefacts/stitching were considered acceptable for analysis. Participants were encouraged to redraw the trajectory line during the immobilization and recovery periods to ensure accurate probe repositioning during subsequent timepoints. A transparent plastic sheet was also used to mark the trajectory and other clear landmarks (e.g., scars, patella and other marks) to provide additional accuracy during rescanning. Finally, previous images were displayed during rescanning to confirm image acquisition accuracy between timepoints. All images were analysed single-blinded using ImageJ2 with Bio-Format plugin (version 2.14.0/1.54f, U. S. National Institutes of Health, Bethesda, Maryland, USA) [57,58] at the end of the intervention. One participant was excluded from all mCSA analyses due to inaccurate repositioning and/or visible imaging artefacts.

Assessment of muscle strength in humans

Voluntary maximal Isometric knee extension and flexion torque (Nm) was measured at 90, 60 and 30° knee flexion using a dynamometer (HUMAC NORM; Computer Sports Medicine inc, Massachusetts,

USA). Participants were first familiarised with the protocol one week prior to baseline testing. Prior to

strength testing, participants performed a standardised warm-up consisting of 5 mins low intensity

cycling, 5 mins stretching and 3 × submaximal efforts (50-80%). Following the warm-up, participants

performed 5 × maximal repetitions at each knee flexion angle (in order of 90, 60 and 30°). Verbal

encouragement was provided during each repetition. The greatest torque produced was considered

their maximum voluntary contraction (MVC). If the difference in torque between the first and second

strongest repetition was >5%, participants performed an additional repetition. MVC was acquired on

MP150 (Biopac, Goleta, CA, USA) with a sampling frequency of 1000 Hz and analysed using

AcqKnowledge software (version 4.4.2, Biopac Systems, Inc., Knivsta, Sweden).

Animals and study design

Ethical approval was obtained, and experimental procedures were conducted with the permissions

within a project license granted (PA693D221) under the British Home Office Animals (Scientific

Procedures) Act 1986. Aged (23 ± 1 months) male Fischer 344 rats (Charles River) weighing 354 ± 54

g were housed at 20°C and 45% relative humidity on an alternating 12 hr light and 12 hr dark cycle,

with food and water available *ad libitum*. Animals were assigned to 1 of 4 groups (*n* = 3 per group)

including a control group (sham control) and 3 experimental groups (atrophy, recovery and repeated

atrophy). All experimental groups received tetrodotoxin (TTX) to the common peroneal nerve (CPN),

preventing contraction of the ankle dorsiflexor muscles. TTX binds to the voltage-gated sodium channel

in the nerve cell membranes and so prevents the conduction of action potentials. Therefore, muscles

innervated by the blocked CPN cannot be activated to contract. This induces disuse atrophy of the hind

limb dorsiflexor muscles i.e., the tibialis anterior (TA) and extensor digitorum longus (EDL), and is

advantageous vs. denervation or hindlimb unloading as it allows normal conduction in the tibial nerve

to achieve plantarflexion, and therefore animals can undertake locomotion for daily living (**Figure 2A**)

[59–61]. The atrophy and recovery groups received a single infusion of TTX (6-7 d) followed by 9 d of TTX

abstinence (e.g., TTX → recovery), whereas the repeated atrophy group received a second TTX

infusion (5-6 d) following recovery (e.g., TTX → recovery → repeated TTX) (**Figure 2A**). The sham

control group consisted of *n* = 1 for each timepoint (TTX/recovery/repeated TTX) that received the same

surgical procedures (detailed below) as the experimental groups in the absence of TTX administration.

Pre- and post-intervention animal weights for each group are presented in **supplementary Table 1** and

muscle weights are provided in **Supplementary Figure 3.**

Animal experimental procedures

Surgical and TTX-administration procedures are described elsewhere [66, 72]. Briefly, animals were first

anaesthetized via inhalation of a gaseous mixture of 3% isoflurane in oxygen for induction and 1–2%

for maintenance prior to subcutaneous implantation of a mini osmotic pump (Mini Osmotic Pump 2002;

Alzet, Cupertino, CA) in the scapular region. Silicone tubing attached to the mini osmotic pump was

then passed under the skin to the site of the left common peroneal nerve (CPN). A second incision was

made laterally through the skin, just proximal to the knee joint and through the biceps femoris muscle
(posterior compartment of the thigh) to enable access to the CPN. A silicone cuff extending from the
silicone tubing was placed around the nerve. All incisions were closed in layers. The mini osmotic pump
(Mini Osmotic Pump 2002; Alzet, Cupertino, CA) delivered TTX to the CPN. The osmotic pump
successfully delivered 0.5 µl/hr TTX (350 mg/mL in sterile 0.9% saline), continuously blocking ankle
dorsiflexion for a 7d period after which the TTX was exhausted, while maintaining normal voluntary
plantarflexion via the tibial nerve. TTX was not administered to the right limb, and it served as a
contralateral control for each individual animal. For the repeated disuse atrophy experiments, the rats
were anaesthetized briefly after the recovery period. The empty osmotic pump was removed through
an interscapular incision and replaced with a new TTX-loaded pump. The welfare and mobility of the
rats were checked daily by animal welfare staff.
321

Human muscle derived cell (MuSC) experimental procedures

Isolation of human muscle derived cells (MuSC) and subsequent cell culture procedures are previously
described by our group [59, 72]. Briefly, pre- and post-atrophy muscle biopsies (*n* = 4 of pre- and post-
biopsies) were immediately transferred to a sterilised class II culture hood (Thermo Scientific MSC-
advantage) in low serum transfer media (components described above). Visible fat and connective
tissue were removed, and remaining muscle tissue was washed 3 × PBS (1× with added PS and AB),
before undergoing repeated rounds of scissor mincing in trypsin (0.5%)-EDTA (0.2%) and centrifugation
(340 × *g* for 5 min at 24 °C) to promote cell migration. The supernatant, cell pellet and remaining non-
homogenised tissue were seeded into separate T25 pre-gelatinised flasks containing 5 mL growth
media (GM: Ham's F-10, 10% hiFBS, 10% heat inactivated new born calf serum/hiNBCS, 2 mM LG,
PS and AB) and left unagitated in a humidified incubator (37 °C, 5%, VWR ILCO 180 Premium) for 5-7
d to promote cell growth. GM was changed every 2-3 d thereafter until sufficient cell attachment and
growth (<80% confluency within cell colonies) was achieved. MuSC were transferred and sub-cultured
in T75 flasks to further increase cell yield and ensure sufficient MuSC numbers for subsequent
experiments described below.

To determine the effects of disuse *in vivo* on differentiation capacity *in vitro*, both pre- and post-atrophy
MuSC (7 × 10^4 cells/mL in 2 mL of GM) were seeded onto pre-gelatinised six-well plates and left until
≈ 80% confluency was attained (taking between 24 and 48 hrs). Cells were then washed 3 × PBS (1×)
and media was switched to low serum media (DM: same components as GM with lower 2% hiFBS) to
permit differentiation over 10 days (d). An additional 1 mL DM was added at 72 hrs and media was
switched to DM containing either 0.5 mM nicotinamide riboside (NR, ChromaDex Inc, USA), previously
used by out group [73] or control (H_2O) for both pre- and post-atrophy conditions. Cell pellets derived from
4 × wells at 0 hrs, 7 and 10 d (+/- NR supplementation) were snap-frozen and stored at -80 °C for later
copurification of DNA (2 ± $0.19_{A260/A280}$/0.14 ± $0.14_{A260/A230}$) and RNA (2.05 ± $0.03_{A260/A280}$/1.86 ±
$0.48_{A260/A230}$) using the AllPrep RNA/DNA Mini Kit (Qiagen). The remaining 2 × wells were treated for
subsequent immunocytochemistry analysis described below. There was no difference in the proportion

of myoblasts (42 ± 22% versus 45 ± 26%, *p* = 0.62) and fibroblast (58 ± 22% versus 56 ± 26%, *p* =

0.62) in pre- and post-atrophy MuSC, respectively. Thus, MuSC were matched prior to differentiation

experiments. All MuSC were between passage 3 and 6 to avoid issues of cellular senescence [74].

Immuno-histo/cytochemistry, imaging and analysis

Serial sections at either 8 µm (human VL muscle) or 10 µm (mid-belly of the rat TA muscle) thickness

were made at −20 °C (CM1860, Leica Biosystems GmbH, Heidelberger, Germany), mounted on

microscope slides (Fisherbrand™ Superfrost™ Plus Microscope Slides, Fisher Scientific), air-dried for

30 mins and stored at −80 °C. When labelling muscle tissue samples, samples were first air-dried for

30 mins prior to blocking in 2% bovine serum albumin (BSA, Sigma-Aldrich) in 1× PBS (10 mM

phosphate buffer, 3 mM KCl, pH 7.4, Merck, Oslo, Norway) with 0.05% added tween (PBS-T, Tween®

20, VWR, Oslo, Norway) and 1% fat-free dry milk (Sigma-Aldrich) for 1 hr at room temperature (RT).

Muscle cross-sections were then counterstained for dystrophin (1:500; ab15277, Abcam, Cambridge,

UK), myosin heavy chain (MHC) type I (1:100; BA-F8, DSHB Supernatant, Schiaffino, S, Iowa, USA)

and MHC type IIA (1:100; SC-71, DSHB Supernatant, Schiaffino, S, Iowa, USA) diluted in 0.05% PBS-

T and 1% milk overnight at 4 °C (**Table 2**). After overnight incubation, samples were labelled with goat

anti-rabbit (GaR) IgG H+L (1:200; Alexa Fluor™ 488, cat. #A-11008 or Alexa Fluor™ 633, cat. #A-

21070, Thermo Fisher Scientific, Waltham, MA, USA), goat anti-mouse (GaM) IgG1 (1:200; Alexa

Fluor™ 488, cat. #A-21121) and/or GaM IgG2b (1:200; Alexa Fluor™ 594, cat. #A-21145) diluted in

0.05% PBS-T and 1% milk for 1 hr and RT (**Table 2**). Slides were washed 3 × 5 mins in 0.05% PBS-T

between each blocking and AB labelling steps. Finally, coverslips were mounted using ProLong™ Gold

Antifade Mountant with 4′,6-diamidino-2-phenylindole/DAPI (cat. #P36935, Thermo Fisher Scientific)

and left to air-dry at RT overnight prior to imaging. For imaging, tile scan images (1024×1024 frame

size) of whole cross-sections were acquired using a ZEISS LSM 800 (with Airyscan) confocal

microscope (ZEISS, Oberkochen, Germany) equipped with a 10×/0.3 objective and subsequently

processed using either ZEN 2 (blue edition, version 2.6) or ImageJ2 (version 2.14.0/1.54f, U. S. National

Institutes of Health, Bethesda, Maryland, USA) software. Automated analysis of myofibre cross-

sectional area (fCSA) and type was conducted using MyoVision 2.0 software [59]. Issues surrounding

muscle tissue processing in the human muscle (e.g., biopsy size, freeze damage) resulted in variable

sample numbers for average (type I and II combined; *n* = 8 for baseline/atrophy, *n* = 7 for recovery, *n* =

6 for repeated atrophy) and type I/II (*n* = 7 baseline, *n* = 8 atrophy, *n* = 6 recovery, *n* = 5 repeated

atrophy) myofibre analysis. The total number of fibres analysed from human biopsies was 398 ± 331

(average), 149 ± 94 (type I) and 281 ± 77 (type II). The total number of fibres analysed from whole rat

muscle was 6297 ± 2067 (average), 280 ± 109 (type I), 1237 ± 546 (type IIA), 69 ± 48 (type I/IIA), 4712

± 1613 (type IIX/IIB).

For characterising myogenic (% myoblasts and fibroblasts) proportions and myotube differentiation,

MuSC were washed 3 × TBS (1×), fixed in 4% PFA for 15 mins, incubated in blocking/permeabilization

solution (5% goat serum/GS and 0.2% Triton X-100 in 1× TBS) for 1 hr at room temperature (RT). Cells

were then washed and incubated overnight (4 °C on a rocker) in Ab solution (2% GS and 0.2% Triton X-100 in 1× TBS) containing anti-desmin (1:50; ab15200, Abcam) and TE-7 (1:100; CBL271, Merck) primary Ab for counterstaining myoblasts and fibroblasts, respectively. After overnight incubation, cells were washed 3 × TBS and incubated at RT for 3 hrs (4 °C on a rocker) in secondary Ab solution containing anti-rabbit IgG TRITC (1:75; T6778, Sigma-Aldrich) for myoblasts and GaM Alexa Fluor® 488 (1:500; ab150117, Abcam) for fibroblasts. Finally, cells were washed 3 × TBS, prior to counterstaining nuclei using DAPI (300 nM; D1306, Thermo Fisher Scientific) for 30 mins at RT. DAPI was removed and stained cells were washed and stored in TBS for later image acquisition. The same microscope and acquisition settings used for imaging tissue sections above were used when imaging MuSC (2 × duplicate wells per condition/timepoint) except for acquiring nine (3 × 3) tile scan images. The ImageJ2 cell counter was used to quantify the number of myoblasts and fibroblasts. The polygon tool was used to outline the sarcolemma of all myotubes (cells with ≥3 nuclei) for assessing myotube area (μm^2).

Table 2. Primary (1º) and secondary (2º) antibodies used for immunohistochemistry (IHC) and Western blot (WB) procedures.

Application	Antibody	Supplier (cat. #)	Concentration	Species
IHC (1º)	anti-dystrophin	Abcam (ab15277)	1:500	Human/Rat
	BA-F8/anti-MHCI	DSHB	1:100	Human/Rat
	SC-71/anti-MHCIIA	DSHB	1:100	Rat
	Anti-Desmin	Abcam (ab15200)	1:50	Human
	TE-7 Anti-Fibroblasts	Merck (CBL271)	1:200	Human
	DAPI	Thermo Fisher (P36935)	in mountant	Human/Rat
	DAPI (*for MuSC*)	Thermo Fisher (D1306)	300 nM	Human
WB (1º)	OXPHOS	Abcam (ab110413)	1:500	Human/Rat
	CS	Abcam (ab96600)	1:4000	Human/Rat
IHC (2º)	GaR IgG (H+L) (Alexa Fluor 488)	Thermo Fisher (A-11008)	1:200	Human
	GaM IgG (H+L) (Alexa Fluor 488)	Abcam (ab150117)	1:500	Human
	GaR IgG (H+L) (Alexa Fluor 633)	Thermo Fisher (A-21070)	1:200	Rat
	GaM IgG1 (Alexa Fluor 488)	Thermo Fisher (A-21121)	1:200	Rat
	GaM IgG2b (Alexa Fluor 594)	Thermo Fisher (A-21145)	1:200	Human/Rat
	anti-rabbit TRITC	Sigma-Aldrich (T6778)	1:75	Human
WB (2º)	GaM IgG (H+L) HRP	Thermo Fisher (31430)	1:30000	Human/Rat
	GaR IgG (H+L) HRP	CST (7074)	1:3000	Human/Rat

DSHB = Developmental Studies Hybridoma Bank, CST = Cell Signalling Technology.

RNA extraction from muscle tissue

Skeletal muscle tissue derived from humans (33 ± 6 mg) and animals (27 ± 4 mg) was added to 2 mL homogenizing mix tubes (Fisherbrand™, Fisher Scientific) containing 600 µl Buffer RLT (Qiagen) with added 1% β-Mercaptoethanol (β-ME) before transferring to a tissue homogenizing instrument (MagNA Lyser; Roche) for subsequent homogenization (3 × 45 s at 6000 rpm, placing samples on ice for 5 mins between each disruption). Following homogenization, RNA was purified using a RNeasy® Mini kit (Qiagen) and treated with DNase (RNAse-Free DNase Set, Qiagen) as per the manufacturer's instructions. RNA quantity and purity for both human (128 ± 175 ng/mg tissue, $1.9 ± 0.1_{A260/A280}/2.1 ± 0.2_{A260/A230}$) and rat (406 ± 215 ng/mg tissue, $2.1 ± 0_{A260/A280}/2.2 ± 0.2_{A260/A230}$) samples were quantified via spectrophotometry (QIAxpert®, software version 2.4, Qiagen) and concentrations confirmed via fluorometry (Qubit®, Qiagen). RNA integrity prior to sequencing was confirmed via Agilent Bioanalsyer 2100 with RIN scores 7.22 + 0.36 for human and 8.29 + 0.62 for rat samples (detailed below).

qPCR for targeted mRNA analysis of aged rat muscle tissue and human MuSC

Methods for mRNA expression analysis via qPCR by our group have been explained elsewhere [40, 48, 49, 59]. Briefly, a one-step QuantiFast SYBR Green kit (Qiagen) was used for preparation of PCR reactions. Each 10 µl reaction contained 4.75 µl experimental sample (7.36 ng/µl totalling 35 ng per reaction), 0.075 µl of both forward and reverse primer (100 µM stock) of the genes of interest (primer details are provided in **Table 3**), 0.1 µl of reverse transcriptase and 5 µl of SYBR Green Master Mix (totalling reactions). A thermocycler (Rotorgene 3000Q, Qiagen) was used to amplify target transcripts. Reverse transcription was initiated with a hold at 50 °C for 10 mins (cDNA synthesis) and 5 mins at 95 °C (transcriptase inactivation and initial denaturation), before 40 cycles of; 95 °C for 10 s (denaturation) followed by 60 °C for 30 s (annealing and extension). Only intended targets (via BLAST search) were identified, thus yielding a single peak after melt curve analysis. All relative gene expression was quantified using the comparative Ct ($^{\Delta\Delta}$Ct) method [75]. For *Nmrk2* expression analysis of aged animal muscle tissue, C_T values (mean of technical duplicates) for each condition (atrophy/recovery/repeated atrophy) were relativised to pooled C_T values from the control limb ($n = 9$) and reference gene, *Pol2ra* (19.27 ± 0.41, 2.12% CV). For temporal mRNA analysis of *MRF* and *NMRK* genes in human MuSC, C_T values (mean of technical duplicates) from pre- and post-atrophy MuSC at 7 and 10 d (+/- NR) were relativised to own 0 hr C_T value and pooled reference gene, *RPL13A* (14.90 ± 0.43, 2.87% CV). To determine the effects of NR supplementation, C_T values from 10 d +NR were relativised 10 d -NR and pooled *RPL13A* values for both pre- and post-atrophy timepoints. Primers were designed using Basic Local Alignment Search Tool (BLAST; http://blast.ncbi.nlm.nih.gov/) and Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/) and purchased from Sigma.

     Table 3: qPCR primers

Species	Gene	Primer sequence (5' - 3')	Length (bp)	Reference
Human	**RPL13A**	F: GGCTAAACAGGTACTGCTGGG	105	Turner *et al.,* [59]
		R: AGGAAAGCCAGGTACTTCAACTT		
	MYOD	F: GCCACAACGGACGACTTCTA	117	Turner *et al.,* [59]
		R: GTGCTCTTCGGGTTTCAGGA		
	MYOG	F: TCCCAGATGAAACCATGCCC	103	Turner *et al.,* [59]
		R: AGGCCCCTGCTACAGAAGTA		
	NMRK1	F: AAATGTTGCTCCATACCATCCC	70	newly designed
		R: CTGCAGCTCCTAGTAATGCT		
	NMRK2	F: AGCTCTTCCGTGAAGTCCTG	62	newly designed
		R: TGATTCCTGGGAGCGGTTCA		
Rat	**Pol2ra**	F: GCTGGACCTACTGGCATGTT	102	Seaborne *et al.,* [72]
		R: ACCATAGGCTGGAGTTGCAC		
	Nmrk2	F: GCCGTATGAGGAATGCAAGC	90	newly designed
		R: ATGGGCCATACGTGACCATC		

RNA-seq and associated bioinformatic analysis

Total RNA (100 ng) from each sample was used to prepare mRNA libraries using the NEBNext mRNA isolation kit in conjunction with the NEBNext Ultra II Directional RNA Library preparation kit (New England Biolabs, Massachusetts, USA), and samples were randomised before preparation. Fragmentation of isolated mRNA prior to first strand cDNA synthesis was carried out using incubation conditions recommended by the manufacturer for an insert size of 300 bp (94 °C for 10 minutes). 13 cycles of PCR were performed for final library amplification. Resulting libraries were quantified using the Qubit 4.0 spectrophotometer (Life Technologies, California, USA) and average fragment size assessed using the Agilent 4200 Tapestation with D1000 screentape (Agilent Technologies, Waldbronn, Germany). A final sequencing pool was created for each project using equimolar quantities of each sample with compatible indexes. 59 bp paired-end reads were generated for each library using the Illumina NextSeq 2000 in conjunction with the NextSeq 2000 P3 100 cycle kit (Illumina Inc., Cambridge, UK). Resulting FastQ files were imported and processed in Partek Flow® software (version 12.4.0; Partek, Inc. Missouri, USA) to filter contaminants (rDNA/tRNA/mtrDNA) using Bowtie 2 (version 2.2.5) and align reads to the hg38 *Homo sapiens* (release 39) and rn7 *Rattus norvegicus* genome assemblies using Spliced Transcripts Alignment to a Reference (STAR) alignment tool (version 2.7.3a) [76]. Total number of pre/post-aligned reads was 35/34 million and 49/49 million for human (65 bp average read length) and rat (59 bp average read length) samples, respectively. Phred scores were >33 for both human and rat RNA samples, thus trimming and filtering of reads was not required. Normalised differential gene expression analysis was performed using *DESeq2* median of ratios [77]. Principal component analysis (PCA), cluster heatmaps and volcano plots were created in R studio [78]. Venn diagrams were redrawn from the open-source DeepVen web application[79]. Pathway (KEGG and REACTOME) and gene ontology (GO; cellular component/CC, molecular function/MF and biological

process/BP) performed via the Database for Annotation, Visualization, and Integrated Discovery (DAVID) knowledgebase [80] and associated figures created in R studio. Custom background gene lists of detected genes following *DESeq2* normalization were used for functional annotation analysis. For human analysis, normalised reads for each timepoint (atrophy/recovery/repeated atrophy) were relativised to baseline. For rat samples, all experimental conditions in the left limb (atrophy/recovery/repeated atrophy) were relativised to pooled right contralateral non-surgical limb. Given there were no DEGs detected between sham and pooled contralateral control limb samples, we excluded sham samples from further analysis and pooled contralateral non-surgical limb samples for the control group. Genes with a ≥20% fold change (FC) and false discovery rate (FDR) of ≤5% are considered differentially expressed genes (DEGs). Data are presented as either $\log_2$ or FC ± confidence intervals (CI). Genes with negative changes presented as signed FC (-1/ratio). To perform temporal clustering of the identified DEGs and DMRs (described below), self-organising map (SOM) profiling was conducted using Partek Genomics suite, as previously described [38, 40, 46, 59]. SOM analysis plots a standardised mean (z-score normalisation) shifting the mean to a value of 0 and scaling to a standard deviation of 1. Therefore, by subtracting the mean of the group this centres the data around zero and then dividing by the standard deviation, scales the data so the spread is consistent, allowing removal of the absolute differences and focusing on relative changes allowing comparison of gene transcripts and DMRs on a common scale. The mapping then clusters DEGs and DMRs based on similar temporal profiles allowing visualisation of temporal signatures over time relative to the group mean. Also, this allows the temporal signature of the DEGs and DMRs on common genes to be compared on the same scale over the time-course of the intervention.

DNA extraction from muscle tissue

Skeletal muscle tissue derived from humans (21 ± 3 mg) and animals (12 ± 2 mg) was added to 2 mL homogenizing mix tubes (Fisherbrand™, Fisher Scientific) containing 180 μl Buffer ATL (Qiagen, Hilden, Germany) before transferring to a tissue homogenizing instrument (MagNA Lyser; Roche, Basel, Switzerland) for subsequent homogenization (3 × 45 s at 6000 rpm, placing samples on ice for 5 mins between each disruption). Following homogenization, 20 μl proteinase K (Qiagen) was added and samples were incubated at 56°C until complete lysis (≈ 30-60 mins) before continuing with DNA purification using a DNeasy® Blood & Tissue kit (Qiagen) as per the manufacturer's instructions. DNA quantity and quality for both human (251 ± 47 ng/mg tissue, 2 ± 0 $_{A260/A280}$/2.1 ± 0.1 $_{A260/A230}$) and rat (1427 ± 426 ng/mg tissue, 2 ± 0 $_{A260/A280}$/2 ± 0.1 $_{A260/A230}$) samples were quantified using spectrophotometry (NanoDrop™ 2000; Thermo Fisher Scientific) and concentrations confirmed via fluorometry (Qubit®, Qiagen). DNA integrity was confirmed prior to sequencing via Agilent Bioanalsyer 2100 with DIN scores 7.11 ± 0.17 for human and 7.22 ± 0.23 for rat samples.

Reduced Representation Bisulfite Sequencing (RRBS): Library preparation and sequencing

RRBS libraries were prepared from 100 ng of genomic DNA using the Diagenode Premium RRBS v2

library preparation kit, as per the manufacturer's protocol. For human samples, two pools of 8 samples,

and three pools of 7 samples were prepared after qPCR quantification of adapter-ligated sample and

prior to bisulphite conversion. 14 cycles of final library amplification were performed after assessment

of bisulphite converted library pool by qPCR. For rat samples RREM (reduced-representation enzymatic

methyl-seq) libraries were prepared using 100 ng of genomic DNA in conjunction with the NEBNext

Enzymatic Methyl-Seq kit (New England Biolabs, Massachusetts, USA). Briefly, DNA samples were

digested using MspI restriction enzyme followed by end-repair and adapter ligation. A two-step

enzymatic conversion was then performed by Oxidation of 5mC and 5hmC using TET2 and

glucosylation of 5hmC using T4-BGT. Deamination of unmodified cytosines was subsequently

performed using APOBEC. Converted samples were then uniquely barcoded and amplified by PCR

prior to sequencing. Libraries were sequenced using the Illumina NextSeq 2000 with P3 SBS 100-cycle

sequencing kit for human, and with P4 XLEAP 100-cycle sequencing kit for rat, both generating 50 bp

paired-end reads.

RRBS quality control, trimming, alignment and methylation calling

Reduced representation bisulfite sequencing (RRBS) libraries were processed using a standard

workflow. Raw base-call files were demultiplexed using Illumina `bcl2fastq /2.19.1`, and unique

molecular identifiers (UMIs) were extracted using `umi_tools` [81]. Reads were adapter and quality-

trimmed (minimum phred score 30) with `trim_galore` (https://doi.org/10.5281/zenodo.5127898). in

RRBS mode, reads less than 15 bp after trimming were discarded. Trimmed reads were aligned to the

human reference genome (hg38) or rat genome Rn6 using `bismark/0.22.1` [82] with `bowtie2`, with

the PBAT option enabled. PCR duplicates were removed with `deduplicate_bismark` in a UMI-aware

manner. CpG methylation calls were obtained using `bismark_methylation_extractor`.

RRBS: Identification of differentially methylated positions (DMPs) and regions (DMRs)

Downstream analysis was performed in R. Per-CpG coverage files were imported into `methylKit` [83]

with sites of low (<10 reads) or extreme coverage (top 0.1%) removed and outlier samples excluded.

For humans, data from each sample was then merged based on only CpGs present in at least 6

samples. Each of our groups contained 7-8 samples after outlier removal, making this threshold a

minimum of 75% of samples and giving a total of ≈ 1.8 million total CpGs analysed. For rats, there were

9 samples (3 per experimental condition of atrophy, recovery and repeated atrophy) and 9 samples for

the contralateral control limbs, therefore data from each sample was merged based on CpGs present

in all samples (100%), giving a total of 29K CpGs analysed. Differential methylation was identified at

individual CpGs (DMPs) and then 1-kb tiled regions (DMRs) using `calculateDiffMeth`. The

percentage methylation of each CpG was calculated using the percMethylation function in methylKit

and during differential analysis the mean percentage methylation of each CpG was calculated for each

group. For DMRs, mean percentage methylation for each experimental group was calculated by

averaging the percentage methylation of all CpGs within a given DMR across all samples from the given

group. For humans, we performed 6 pairwise comparisons of: 1) Atrophy vs. Baseline, 2) Recovery vs.

Baseline, 3) Repeated Atrophy vs. Baseline, 4) Recovery vs. Atrophy, 5) Repeated Atrophy vs.

Recovery, 6) Repeated Atrophy vs. atrophy. For rat analysis we performed: 1) Atrophy vs. Control, 2)

Recovery vs. Control, 3) Repeated atrophy vs. Control, 4) Recovery vs. Atrophy, 5) Repeated Atrophy

vs. Recovery, 6) Repeated atrophy vs. atrophy. For the rats, as with the transcriptome data, control was

determined as the pooled right/non-surgical limb (contralateral control limb = 9). Differentially

methylated CpGs and regions were annotated to genes and regulatory features for humans using

TxDb.Hsapiens.UCSC.hg38.knownGene(https://doi.org/10.18129/B9.bioc.TxDb.Hsapiens.UCSC.hg3

8.knownGene),   org.Hs.eg.db   (https://doi.org/10.18129/B9.bioc.org.Hs.eg.db)   and   annotatr

(https://doi.org/10.18129/B9.bioc.annotatr) and in rats using TxDb.Rnorvegicus.UCSC.rn6.refGene

(https://doi.org/doi:10.18129/B9.bioc.TxDb.Rnorvegicus.UCSC.rn6.refGene)   and   rg.Rn.eg.db

(https://doi.org/doi:10.18129/B9.bioc.org.Rn.eg.db). Default settings were used unless otherwise

specified. Only DMRs with FDR ≤ 0.05 were considered significant. SOM clustering (described above)

was performed on significant DMRs using percentage difference methylation as this allowed

methylation values at baseline (human) and control (rat) to enable temporal profiling.

Mitochondrial DNA content

Mitochondrial DNA (mtDNA) content was assessed via real time quantitative polymerase chain reaction

(RT-qPCR). PCR primers encoding for nuclear DNA (nDNA) markers; β$_2$ microglobulin (*B2M*),

hemoglobin subunit beta *(HBB)*, tubulin-specific chaperone A (*Tbca*), glyceraldehyde 3-phosphate

dehydrogenase (*Gapdh*), and mtDNA markers; mitochondrial 16S rRNA (*MT-RNR2*), D-loop region

(*DLOOP*), cytochrome B (*CYTB*), mitochondrially encoded 16S rRNA (*mt-Rnr2*) and 12S ribosomal

RNA (*mt-Rnr1),* were used to determine nuclear and mitochondrial DNA copy number, respectively.

Primer information is presented in **Table 4.** For qPCR, 20 ng DNA (9.6 µl at 2.08 ng/µl in nuclease-free

H$_2$O) was added to 10 µl SYBR$^®$ Green Mastermix (Qiagen) and 0.2 µl primers (100 µM stock, 1 µM

final concentration) to ensure 20 µl reaction volume. DNA was then amplified using a Rotor-Gene Q 5-

Plex HRM thermocycler (Qiagen) with supporting software (version 2.3.1). Thermal cycle settings

consisted of 3 mins at 95 °C (initial denaturation), followed by 40 cycles of 10 s at 95 °C, 30 s at 62 °C

(annealing) and final extension of 10 s at 95 °C. Melt curve analysis was performed at 0.5 °C increments

from 65 to 95 °C. Mitochondrial DNA content was reported as the ratio of mtDNA (*16S, CYTB* and/or

DLOOP for human; *mt-Rnr2* and/or *mt-Rnr1* for rat) to pooled nDNA (*B2M/HBB* for human; *Tbca/Gapdh*

for rat) determined using the ^{ΔΔ}Ct method [75, 84].

Table 4. Primers used for encoding for nuclear DNA (nDNA) and mitochondrial DNA (mtDNA) markers for assessment of mtDNA content.

Species	Gene	Primer sequence (5' - 3')	Length (bp)	Reference
Human	**B2M**	F: TGCTGTCTCCATGTTTGATGTATCT	86	Lapatto *et al.,* [85]
	(nDNA)	R: TCTCTGCTCCCCACCTCTAAGT		
	HBB	F: CAGGTACGGCTGTCATCAGTTAG	185	Lapatto *et al.,* [85]
	(nDNA)	R: CATGGTGTCTGTTTGAGGTTGCT		
	DLOOP	F: CATCTGGTTCCTACTTCAGGG	106	Lapatto *et al.,* [85]
	(mtDNA)	R: CCGTGAGTGGTTAATAGGGTG		
	CYTB	F: GCCTGCCTGATCCTCCAAAT	132	Lapatto *et al.,* [85]
	(mtDNA)	R: AAGGTAGCGGATGATTCAGCC		
	MT-RNR2	F: GGGGCGACCTCGGAGCAGAA	213	Lapatto *et al.,* [85]
	(mtDNA)	R: ATAGCGGCTGCACCATCGGGA		
Rat	**Tbca**	F: TGTCCCACCTGTTTGTTCTCT	50	Kemper *et al.,* [86]
	(nDNA)	R: GGTCCTTGGACACCTAGACG		Arduini *et al.,* [87]
	Gapdh	F: GGAAAGACAGGTGTTTTGCA	129	Stevanovic-Silva *et al.,* [88]
	(nDNA)	R: AGGTCAGAGTGAGCAGGACA		
	mt-Rnr1	F: CTCAAGACGCCTTGCCTAGC	131	Kemper *et al.,* [86]
	(mtDNA)	R: TCGTATGACCGCGGTGGCT		Arduini *et al.,* [87]
	mt-Rnr2	F: AGCTATTAATGGTTCGTTTGT	132	Stevanovic-Silva *et al.,* [88]
	(mtDNA)	R: AGGAGGCTCCATTTCTCTTGT		

Protein extraction and Western blot

Skeletal muscle tissue derived from humans (21 ± 6 mg) and rats (38 ± 14 mg) was added to 2 mL homogenizing mix tubes (Fisherbrand™, Fisher Scientific) containing either 1 mL RIPA buffer (Thermo Fisher Scientific) with added protease and phosphatase inhibitors (Halt™ inhibitor cocktail, Thermo Fisher Scientific) for human samples or sucrose lysis buffer (250 mM sucrose, 50 mM Tris pH 7.5, 1 mM EDTA, 1 mM EGTA, 1% Triton™ X-100, Sigma-Aldrich) with added inhibitors (Thermo Scientific™ Pierce Mini Tablets, Thermo Fisher Scientific) for rat samples. Tubes were then transferred to a MagNA Lyser instrument (Roche) for subsequent homogenization (3 × 45 s at 6000 rpm, placing samples on ice for 5 mins between each disruption). Following homogenization, protein concentration was quantified for both human (73 ± 13 µg/mg) and rat (84 ± 18 µg/mg) tissue via the bicinchoninic acid (BCA) method (Pierce™ BCA protein assay kit, Thermo Fisher Scientific) and a FLUOstar Omega microplate reader (BMG LABTECH, Ortenberg, Germany) with supporting software (Omega version 1.3). For Western blotting, protein aliquots were first diluted in 4× Laemmli Sample Buffer (Bio-Rad) with added 0.5 M Dithiothreitol (DTT, Bio-Rad) and heated at either 45°C (oxidative phosphorylation, OXPHOS) or 70°C (citrate synthase, CS) for 10 mins. Following denaturation, 20 or 25 µg for all protein samples (in 15 µL for human and 20 µL for rat) and 5 µL protein marker (10-250 kDa, Precision Plus Protein™ All Blue Prestained Protein Standards, Bio-Rad) were loaded into pre-cast gels (12 or 15 well,

4–20% Mini-PROTEAN® TGX Stain-Free™, Bio-Rad) before undergoing protein separation via electrophoresis (200 volts/V for 30 mins). The current was monitored (A) pre- and post-electrophoresis. Following electrophoresis, gels were activated under UV light ChemiDoc MP Imaging System (Bio-Rad) and proteins transferred to PVDF nitrocellulose membranes (0.2 μm, Bio-Rad) using transfer stacks (Bio-Rad) and a Trans-Blot Turbo Transfer System (Bio-Rad). Following protein transfer, membranes were blocked (5% milk in 1% PBS-T for 1 hr at RT) and subsequently labelled with primary antibodies against mitochondrial OXPHOS complexes (CI-NDUFB8, CII-SDHB, CIII-UQCRC2, CIV-MTCO1, CV-ATP5A) (1:500, cat. #ab110413, Abcam) and CS (1:4000, cat. #ab96600, Abcam). Primary Ab were diluted in 1% milk in 1% PBS-T and incubated on a roller overnight at 4 °C. After overnight incubation, OXPHOS and CS proteins were labelled with secondary antibodies, GaM IgG (H+L) HRP (1:30000, cat. #31430, Thermo Fisher Scientific) and GaR IgG (H+L) HRP (1:3000, cat. #7074, Cell Signaling Technology), respectively. Membranes were washed (3 × 5 mins in 1% PBS-T) between blocking and antibody labelling steps. Finally, membranes were incubated in enhanced chemiluminescence (ECL) HRP substrate (SuperSignal™ West Dura Extended Duration Substrate, Thermo Scientific) for 5 mins at RT to enable visualisation of membrane-bound proteins via chemiluminescence using a ChemiDoc MP system (Bio-Rad) and analysis of band intensities using supporting software (Image Lab 6.1, Bio-Rad). For human samples, all timepoints for each participant were loaded on the same gel and repeated to ensure duplicate samples across two separate gels. Band intensities were relativised to own baseline sample for subsequent timepoints (atrophy/recovery/repeated atrophy) and the average across the 2 gels was reported. One human recovery sample was excluded from all analysis due to low detection of protein. Thus, there was n = 8 (rather than n = 9) samples for recovery timepoint. Similarly for aged animal samples, the left/surgical and right/non-surgical limbs for each animal and condition (excluding repeated atrophy as the protein lysates were low quality and quantity) were loaded on the same gel across two separate gels. Band intensities of left/surgical limb were relativised to thier own right/non-surgical limb for each condition (sham/atrophy/recovery). Control samples consisting of a mixture of all human or rat samples were loaded in duplicate within each gel to ensure consistent sample loading. There was no CII-SDHB band detected in 1 out of the 2 gels for both human and rat samples. Therefore, data is derived from only one gel for this protein.

NAD$^+$ measurements

NAD$^+$ levels were measured using an enzymatic cycling assay [89]. Briefly, 11 ± 1 mg (human) and 8 ± 2 mg (rat) muscle were lysed in 400 μl 0.6 M perchloric acid (HClO$_4$) and homogenized using steel beads and a TissueLyser II (Qiagen). Following homogenisation, samples were centrifuged for 5 min at 13,000 g. The supernatant was transferred to new tubes and neutralized in 100 mM Na$_2$HPO$_4$ (1:300 for human, 1:800 for rat). For the assay, 100 μl of diluted extract was pipetted into a white 96-well plate after which 100 μl of reaction mix was added. The reaction mix contained: 100 mM Na$_2$HPO$_4$, 10 μM flavinmononucleotide, 2 % ethanol, 90 U/mL alcohol dehydrogenase, 130 mU/mL diaphorase, 2.5 μg/mL resazurin, and 10 mM nicotinamide. A Hidex Sense Microplate Reader (Hidex, Finland) was used to measure the fluorescence of accumulated resorfurin (Ex 544 nm/ Em 580 nm) for 30 min

immediately after the reaction mix was added. Samples were run in duplicates and NAD$^+$ content was

calculated from the increase in fluorescence of a dilution series of NAD$^+$. Values were normalized to

protein concentration by dissolving the resultant pellets from the previous centrifugation step in 0.2 M

NaOH via heating to 95°C. Protein concentrations were determined using the bicinchoninic acid (BCA)

assay (Pierce™ BCA protein assay kit, Thermo Fisher Scientific).

Statistical Analysis

A Shapiro-Wilk´s test of normality was first performed for all physiological (e.g., body weight, SkM mass,

strength, mCSA and fCSA), protein abundance, mtDNA content and NAD$^+$ measurements. A repeated

(for human) or between (for aged animals) measures one-way analysis of variance (ANOVA) with

Tukey´s HSD post hoc detected significant ($p \leq 0.05$) differences whenever data was normally

distributed. Whenever data was non-normally distributed, a Friedman (for human) or Krustal-Wallis (for

aged animals) test with follow-up Dunn´s post hoc was carried out. For human myofibre (fCSA and fibre

type) analysis, a mixed-effect model was performed due to differences in sample number across

timepoints. For animal body weight (individual conditions), myofibre, mtDNA and NAD$^+$ content

analyses, a two-way ANOVA (within subject factor: limb, between subject factor: condition/timepoint)

with Tukey´s HSD post hoc was performed. For mean animal body weight across all conditions, an

unpaired t-test was performed. All statistical analysis was conducted in GraphPad Prism software for

macOS (version 10.4.1, Massachusetts, USA) and associated figures were created using GraphPad or

R studio for macOS (version 4.4.3). All data are presented as mean ± standard deviation (SD) unless

otherwise stated in the figure legends. RNA-seq and RRBS data statistical analyses are described

above.

Results

Atrophy and repeated atrophy results in loss of leg lean mass, skeletal muscle size and strength in
young adult humans, with full recovery after initial disuse

In young adult humans, the first period of disuse-induced atrophy (hereafter referred to as "atrophy")

resulted in a significant -3.6% reduction in total leg lean mass of the immobilised limb ($p = 0.005$, **Figure**

1B). Upon returning to normal habitual activity (i.e., "recovery"), mass was fully restored to pre-atrophy

baseline levels ($p = N.S$ recovery vs. baseline). The second period of disuse (i.e., "repeated atrophy")

led to a similar significant -3.2% loss compared to previous recovery ($p = 0.004$, **Figure 1B**). Further

assessment of *vastus lateralis (VL)* and *rectus femoris (RF)* anatomical muscle cross-sectional area

(mCSA) in the immobilised limb revealed a -9% reduction in *VL* mCSA after initial atrophy ($p = 0.0003$),

that was fully restored during recovery (recovery vs. baseline, *p = N.S,* **Figure 1C**)*. Vastus lateralis*

mCSA was also significantly reduced after repeated atrophy relative to both baseline (-10.6%, $p =$

0.0003) and previous recovery (-8.9%, $p = 0.002$, **Figure 1C**). *Rectus Femoris* mCSA was only

significantly reduced after repeated atrophy relative to recovery (-3.5%, $p = 0.02$, **Figure 1D**). There

were no significant changes in the non-immobilised limb for total lean leg mass (**Supplementary Figure 1A-F)** or *VL* and *RF* mCSA (p = N.S, **Supplementary Figure 1 G-J**). There was however a significant increase in lower leg lean mass of the non-immobilised limb during both periods of disuse periods possibly due to a greater reliance on plantarflexor and dorsflexor muscles to support balance and locomotion during immobilization of the opposite limb (**Supplementary Figure 1F**).

Concomitant with the loss of leg lean mass and quadriceps muscle size was a significant reduction in knee extension strength (assessed via MVC) at 90° (-14.3%, p = 0.02) and 60° (-12.5%, p = 0.02) knee flexion following initial atrophy that was also fully restored during recovery (p = N.S) (**Figure 1E & F**). Repeated atrophy induced a similar loss of strength at 90° (-13.8%, p = 0.02) and 60° (-14.3%, p = 0.002) following recovery (**Figure 1E & F**). Despite this loss of muscle strength at 60° and 90°, there were no significant changes in knee extension MVC at 30° knee flexion (p = N.S, **Figure 1G**). Concurrent reductions in size and strength also resulted in impaired muscle quality (e.g., specific force by normalizing MVC to mCSA) (**Supplementary Figure 2A-I**). Specifically, MVC at 90° knee flexion when normalised to *RF* mCSA was significantly reduced after atrophy (-12.8%, p = 0.05, **Supplementary Figure 2D),** where repeated atrophy led to the greatest loss of specific force at 60° when normalised to *VL* (-7.1%, p = 0.03), *RF* (-12.2%, p = 0.04) and *VL+RF* mCSA combined (-8.2%, p = 0.02, **Supplementary Figure 2B & E & H** respectively). Finally, repeated atrophy led to significantly reduced knee flexion torque at 90° (p = 0.03*,* **Supplementary Figure 2J**). Mean MVC at 60° and 30° knee flexion also demonstrated the same trend but did not reach statistical significance (p = N.S, **Supplementary Figure 2 K & L**)

At the myofibre level, IHC analysis revealed an average reduction in fibre type-specific cross-sectional area (fCSA) after atrophy which approached significance when pooling fast and slow fibres (**Figure 1H-K**). However, due to limited fibre number in the cross-sections from fewer samples in specific conditions, we also used the open-source "FibeRtypeR" [90] web application to predict fibre type proportions using RNA-seq gene lists, with this analysis suggesting a significantly greater proportion of type II fibres at baseline (60% type II vs. 40% type I, p = 0.01), atrophy (67% type II vs. 33% type I, p < 0.001) and repeated atrophy (63% type II vs. 37% type I, p = 0.003) with no significant difference after recovery (61% type II vs. 39% type I, p = 0.05) (**Figure 1L**). Overall, demonstrating a reduction in the number type I fibres and an increase in type-II fibres with both atrophy and repeated atrophy and a return to baseline proportions with recovery (**Figure 1L**). From these gene lists, we next interrogated all essential and regulatory fast (*MYL1, MYLPF*), slow (*MYL2, MYL3, MYL6B*) and fetal myosin light chains (*MYL6*) as well as adult (*MYH7, MYH2, MYH1*) and developmental (*MYH3*, *MYH8*) myosin heavy chains (**Figure 1M & N)**. On average, all 6 myosin light chains decreased after atrophy, were fully restored after recovery and then reduced again after repeated atrophy. The slow *MYL2* and *MYL6B* genes were significantly downregulated after atrophy (**Figure 1M**). A similar temporal profile was observed for slow(er) adult *MYH7* (type I) and *MYH2* (type IIA) genes that both decreased after atrophy and repeated atrophy whereas fast *MYH1* (type IIX) increased after both periods of atrophy (**Figure 1N**). Overall, suggesting a predominant downregulation of slow MYL and MYHC´s after repeated atrophy and an increase in type IIX myosin, thus suggestive of a shift towards a faster phenotype in response to disuse.

Finally, expression of embryonic *MYH3* and perinatal *MYH8* both significantly increased after repeated
atrophy (**Figure 1N**).

Figure 1. Young adult human intervention study design (created using BioRender.com) **(A).** Changes in leg lean mass **(B),**
Vastus Lateralis (*VL*) mCSA **(C)**, *Rectus Femoris* (*RF*) mCSA **(D)** and strength/MVC across knee flexion angles 90, 60 and 30°
(E, F, G respectively) after disuse (atrophy), recovery and repeated disuse (repeated atrophy). Significant reductions in muscle
size and strength were observed after disuse atrophy, restored during recovery and reduced to a similar extent after repeated
atrophy. *RF* mCSA only occurred after repeated disuse **(D).** Bar graphs depict relative change (%) from the previous timepoint
(i.e., atrophy versus baseline, repeated atrophy versus recovery). *N* = 10 for leg lean mass **(A)** and strength **(E-J)**; *n* = 9 for mCSA
(B-D). Myofibre analyses **(H-M)** revealed a non-significant reduction in fCSA **(H-K)**. *N* = 8 baseline and atrophy, *n* = 6 recovery,
n = 5 repeated atrophy. **(H)** Representative fluorescence microscopic image (10× magnification; magenta = dystrophin/myofibre
membrane, yellow = MHCl/type I myofibres, cyan = DNA, non-staining = putative type II myofibres; Scale bar within tile scan
image = 500 μm, scale bar within magnified inset image = 100 μm). FibeRtypeR[90] analysis suggested a larger proportion of
fast/type II versus slow/type I fibres at baseline, atrophy and repeated atrophy with no significant differences at recovery **(L)**.
Despite no significant reduction in fCSA, there was reduced mRNA levels of slow isoform *MYL* and *MYH* genes after atrophy and
repeated atrophy as well as increased fast type IIX (*MYH1*) and developmental *MYH* genes after repeated atrophy **(M & N)**. *N* =
9 for each timepoint derived from RNA-seq data. *$p \leq 0.05$, **$p \leq 0.01$, **$p \leq 0.001$, ****$p \leq 0.0001$.

Repeated TTX-induced disuse atrophy results in a reduction in dorsiflexor muscle mass and fibre cross-
sectional area (fCSA) and a lack of recovery in aged rat muscle

In aged animals, there was no significant reduction in body weight in controls, atrophy or recovery
animals (*p = N.S*) (**Supplementary Table 1**). However, a significant reduction in body weight after
repeated atrophy (-18.1%, *p* = 0.02) as well as mean post-interventional versus pre-surgical body
weight when pooling all conditions (-11.7%, *p* = 0.003, **Supplementary Table 1**). Thus, muscle weights
were normalised to respective body weights for each condition throughout. TTX administration to the
common peroneal nerve (CPN) (**Figure 2A**) resulted in a non-significant loss of TA muscle weight
during initial atrophy (-4.4%, *p* = 0.6) and a significant loss after repeated TTX-induced atrophy (-29.8%,
p = 0.002, **Figure 2B**). There was also a significant reduction in EDL muscle weight following repeated
atrophy (-31.1%, *p* = 0.01). Previously published work by our group reported sufficient recovery of
muscle mass following TTX-induced atrophy in young adult rats [66]. However, despite TTX-cessation
and a return to normal dorsiflexion in aged rats during the recovery period, TA muscle loss continued
to occur after recovery (-17.9%, *p* = 0.01, **Figure 2B**), despite shorter TTX exposure and longer active
recovery than young adult rats [66]. There was no significant reduction in plantarflexor (soleus, plantaris
and gastrocnemius) muscle weights, thus emphasising precise selective silencing of the dorsiflexor
muscles using TTX to the common peroneal nerve (*p = N.S*, **Supplementary Figure 3A-D**). At the
transcript level, there was increased expression of developmental myosin heavy chain (*Myh3/8*) genes
across all conditions (**Figure 2C**), whereas adult fast (*Myh1/4*), slow (*Myh7*) myosin heavy chains, as
well as essential (*Myl3*) and regulatory (*Mylpf/2*) myosin light chains significantly decreased after both
periods of disuse (**Figure 2D**). Interestingly, these returned to control levels upon recovery despite
continued muscle loss during TTX cessation (**Figures 2C & D**) suggesting that *Myh1/4/7* and *Myl2/3/pf*
genes were sensitive to the TTX-disuse stimulus. Conversely, *Myh2*, *Myl1* and *Myl4* gene expression
significantly increased following TTX cessation only. Overall, persistent elevation of developmental
Myh3 and *Myh8* mRNA expression coupled with significant reductions in adult fast and slow *Myh* and
Myl isoforms occurred after both periods of atrophy. Coinciding with reductions in muscle weight and
Myh and *Myl* expression, IHC analysis (**Figure 2E**) showed a significant loss in average (type I & II

combined, **Figures 2F & K**), type I (**Figures 2G & L**) and type IIA (**Figures 2H & M**) myofibre cross-sectional area (fCSA) after repeated atrophy. There was also a significant reduction in pure type I (**Figure 2G & L**) and hybrid I/IIA (**Figure 2I & N**) fCSA after initial TTX-induced atrophy, thus supporting the transcriptional expression data that slow fibres were sensitive to TTX-induced disuse. When relativising left TTX to right control limb, there were no significant changes in fCSA across all fibre types (**Supplementary Figures 3 F-J**). Finally, there was a greater contribution of type IIA and reduced proportion of IIX/IIB fibres after recovery (**Figure 2P**) as well as a leftward shift and larger frequency of smaller (≈ <1000 μm^2) predominantly slower fibres in the left TTX versus right control limb (**Figures 2Q-U**).

Figure 2. (**A**) Schematic representation of repeated disuse model in aged animals (created with BioRender.com). CPN = common peroneal nerve. (**B**) Reduction in TA muscle weight during atrophy and repeated atrophy with lack of recovery (presented as % change of the left TTX versus right control limb normalised to body weight). (**C**) Myosin heavy chain transcriptional analysis revealed significant changes in developmental (*Myh3/8*) and adult (*Myh7/2/1/4*) isoforms after atrophy and repeated atrophy (**D**) Essential (*Myl1/3/4/6*) and regulatory (*Mylpf/2*) myosin light chains were also differentially expressed after atrophy, recovery and/or repeated atrophy yet most were restored during recovery, despite continued muscle loss (**E**) Representative fluorescence microscopic image of whole TA muscle section labelled with dystrophin (magenta), MHCI (yellow), MHCIIA (blue), DNA (cyan). Non-staining represents putative type IIX/IIB myofibres. Tile scan images acquired at 10× magnification (scale bar within scan image = 500 μm, scale bar within magnified inset image = 100 μm). (**F-O**) There were significant reductions in fCSA of predominantly slow fibres after atrophy (pure type I and I/IIA hybrid fibres) and repeated atrophy (average, type I and IIA fibres). Symbol shapes represent individual animals. (**P**) Proportional fibre type analysis revealed greater contribution of type IIA and

reduced number of IIX/IIB after recovery versus control in the left surgical limb. (**Q-U**) Fibre size distribution analysis demonstrates

a leftward shift and larger frequency of smaller (<1000 μm^2) fibres in the left TTX versus right control limb. *$p \leq 0.05$, **$p \leq 0.01$,

***$p \leq 0.001$. fCSA = myofibre cross-sectional area.

Atrophy and repeated atrophy in adult human muscle predominantly alters transcript expression of

genes associated with aerobic metabolism, mitochondrial function, skeletal muscle structure and

function

In young adult humans, the largest changes in gene expression occurred after initial atrophy, where

most genes were downregulated after both atrophy and repeated atrophy (**Figures 3A-F**). PCA analysis

revealed a larger overlap between baseline and recovery as well as between atrophy and repeated

atrophy timepoints, suggesting a closer transcriptomic profile between both periods of physical activity

(baseline and recovery) and inactivity (atrophy and repeated atrophy) (**Figure 3B**). Among the 15,096

genes detected, there were 4-5-fold more DEGs after initial atrophy (381 DEGs) than after repeated

atrophy (103 DEGs vs. baseline, 82 DEGs vs. recovery) with predominant downregulation of genes

after both atrophy (63%) and repeated atrophy (77% vs. baseline, 85% vs. recovery) (**Figures 3C-F**).

Following recovery from initial atrophy, more genes were upregulated (58%) (**Figure 3E**). Venn diagram

analysis demonstrated a large proportion of overlapping downregulated (45% of total repeated atrophy

DEGs vs. baseline and recovery) or upregulated (53%) genes after both periods of atrophy (**Figure**

3G). During recovery, there were no detected DEGs relative to baseline, with 536 DEGs following

recovery from initial atrophy which were all unique from atrophy and repeated atrophy comparisons

(**Figures 3E & G**).

Interrogation of the most differentially expressed genes across comparisons revealed that *NMRK2,* an

NAD⁺ biosynthesis gene, was among the top-most downregulated DEGs after atrophy (-8.4 FC, FDR

most significant), ranked second to the nuclear receptor, *NR4A3* (-11.5 FC, **Figures 3C & H**). *NMRK2*

was also the most downregulated gene after repeated atrophy versus baseline (-5.0 FC) and when

compared with recovery (-5.6 FC), as well as the most upregulated DEG after recovery versus atrophy

(9.2 FC**, Figures 3C & H).** Interestingly, after repeated atrophy only, the most upregulated genes were

developmental myosin heavy chain *MYH8* (14.2 FC vs. baseline, 9.8 FC vs. recovery)*,* tropomyosin-

binding subunit *TNNT2* (8.3 FC vs. recovery) and NMJ nicotinic acetylcholine receptor (nAChR) subunit

genes, *CHRNA1* (5.4 FC versus baseline) and *CHRND* (4.0 FC versus baseline) (**Figure 3D & F**).

Another AchR subunit gene, *CHRNG* (2.86 FC versus baseline), was one of the most upregulated by

average FC (across all timepoints) only after repeated atrophy (**Figure 3H**).

Such transcriptional perturbations across the transcriptome corresponded to significant downregulation

of enriched pathway and gene ontology terms related to aerobic metabolism, respiration (e.g., NAD⁺ /

NADH metabolism, TCA cycle, ETC, ATP synthesis) and mitochondrial structure, function and turnover

after both periods of disuse, with these genes returning to baseline levels after recovery (**Figures 3I &**

J and **Figure 3K, cluster 1).** Furthermore, enrichment of upregulated genes (including AchR subunits

and *MYH8 and TNNT2* highlighted above) were associated with muscle-specific terms including

acetylcholine receptor signalling, synaptic transmission, membrane depolarisation and skeletal muscle

contraction (**Figure 3I and J**).

Young adult skeletal muscle demonstrates a transcriptional protection against repeated atrophy

Temporal gene clustering analysis (SOM) revealed most genes decreased (57%, cluster 1) and

increased (41%, cluster 2) after atrophy, returned to baseline levels after recovery, and were less

susceptible to changes after repeated disuse (**Figure 3K**). This represents a transcriptional attenuation,

resulting in a smaller magnitude of change to repeated atrophy stimuli for both downregulated (cluster

1) and upregulated (cluster 2) genes. Based on SOM standardised means, there was a 40% smaller

magnitude in downregulation of genes in cluster 1 and an 72% smaller magnitude in upregulation of

genes in cluster 2 after repeated atrophy compared with initial atrophy. Enriched pathway and GO terms

associated with the response of these downregulated genes to both periods of disuse (cluster 1) were

the same aerobic metabolism, respiration and mitochondrial-related terms identified across global

downregulated genes described above (i.e. NAD^+ / NADH dehydrogenase activity, TCA cycle, ETC,

ATP synthesis and mitochondrial structure, assembly and protein turnover pathway genes, **Figures 3L**

& M). Genes that were upregulated after atrophy and increased less after repeated atrophy (cluster 2,

Figure 3K), were enriched in terms related to amino acid transport, transcription coregulator/coactivator

activity and chromatin binding (**Figure 3L & M**). In contrast, genes associated with these

terms/pathways were also downregulated during recovery from atrophy (**Figure 3J**). Therefore, the

fewer number of total DEGs coupled with less susceptible alterations to repeated versus initial atrophy

suggests a transcriptionally protective memory of earlier atrophy.

SOM cluster temporal analysis also identified 11 downregulated genes (*NR4A1, NR4A3, PRKAG2,*

RP9, CIITA, CORO2B, ENSG00000242299, RPL35, SNRPA1, UBE3D, RPL29P11) after atrophy that

remained downregulated during recovery, despite restoration of muscle mass (cluster 3, **Figure 3K**).

This included nuclear receptor genes, *NR4A1* and *NR4A3,* as well as *PRKAG2* (AMPK gamma2),

where *NR4A3* was among the top ranked downregulated genes within enriched metabolic processes.

Also, ribosomal genes (*RPL35, RPL29P11*) that also demonstrated downregulated enrichment at the

pathway/term level after atrophy (**Figure 3J**). Suggesting a memory of atrophy at the transcriptional

level in these genes via retention of gene expression changes from initial disuse even after the recovery

period.

Finally, 6 genes (*TNNT2, MYH3, DCLK1, ZBED6CL, CDC73, WDR44*) were upregulated after atrophy,

remained elevated during recovery, with even larger increases in expression after repeated atrophy

(cluster 4, **Figure 3K**). This included *TNNT2* (top ranked upregulated gene as discussed above),

embryonic myosin heavy chain (*MYH3*), *DLK1* associated with regeneration of muscle following injury,

and *ZBED6CL* also linked with muscle development. Therefore, whilst most genes demonstrated

transcriptional protection after repeated atrophy, a small number of genes related to muscle

development and regeneration (together with the AchR subunit genes discussed above) demonstrated

exaggerated responses to repeated atrophy in adult human muscle. The retained upregulation and exaggerated increases after repeated disuse also suggests a memory of earlier atrophy at the transcriptional level in these genes.

the top 25 DOWN- (blue) or UP-regulated (orange) genes according to average fold-change across all timepoints (**ii**). Over-representation analysis (ORA) of the top 20 enriched (**J**) pathway (KEGG, top 4 panels; REACTOME, bottom 4 panels) and (**K**) gene ontology (GO) terms for both DOWN- and UP-regulated DEGs across all comparisons ("Atrophy vs. Baseline", "Repeated Atrophy vs. Baseline", "Recovery vs. Atrophy", "Repeated Atrophy vs. Recovery"). (**K**) Self-Organizing Maps (SOM) composed of 761 genes that were significantly differentially expressed in at least one comparison, illustrating gene cluster temporal profiles across the time-course of atrophy, recovery and repeated atrophy. Cluster 1 & 3 (blue) represent genes that were DOWN-regulated following the first period of atrophy. Cluster 2 & 4 (orange) depict UP-regulated genes after initial atrophy. (**L**) Pathway and (**M**) GO analysis revealed enriched terms for clusters 1 and 2 only, in which most genes were either DOWN- (Cluster 1) or UP-regulated (Cluster 2) after atrophy, recovered upon reloading and were less susceptible to changes after repeated atrophy. Cluster 3 represents downregulated genes after atrophy, with retained reduction during recovery and repeated atrophy. Cluster 4 represents upregulated genes after atrophy, retained during recovery, followed by greater increases in expression after repeated atrophy. BP = biological process, CC = cellular component, MF = molecular function.

Repeated atrophy in aged muscle results in downregulation of aerobic metabolism, NAD⁺/NADH and muscle-specific genes and upregulation of genes associated with ECM, protein turnover and DNA damage

Alongside the loss of TA muscle mass during atrophy, recovery and repeated atrophy in aged animals, there were considerably large transcriptomic perturbations after atrophy and recovery with the greatest changes detected after repeated atrophy (**Figures 4A-I**). PCA analysis showed similar clustering of TTX-induced atrophy and repeated atrophy, and distinct clustering after TTX cessation-induced recovery (**Figures 4B & C**). There was also considerable overlap of the sham surgical and pooled non-surgical controls (**Figure 4B**). Given there were also no DEGs detected between sham and pooled contralateral control limb samples, suggesting the surgery had no significant impact on gene transcription, we excluded sham samples from further analysis and pooled contralateral (right) non-surgical limb samples for the control group. Among the 11,284 genes detected post-normalization, there was a similar number of DEGs in response to atrophy (2106 DEGs) and recovery (2008 DEGs) with 2-fold more DEGs detected after repeated atrophy (3937 DEGs) vs. controls. The proportion of downregulated DEGs was 56% after atrophy (**Figure 4D**), 53% after recovery and 56% (vs. control) or 58% (vs. recovery) after repeated atrophy (**Figure 4F-H)**. However, despite a similar number of total DEGs after atrophy and recovery, the majority of downregulated (90%) and upregulated (74%) DEGs were unique after TTX cessation (**Figure 4J**), as evidenced by the distinct PCA clustering of recovery samples (**Figure 4C**). Also, whilst the 2-fold greater number of DEGs after repeated atrophy versus initial atrophy may be partially explained by the lack of recovery during the no TTX period, when relativised to previous recovery, the number of DEGs after repeated atrophy (3198 DEGs) was still 1.5-fold more than initial atrophy (2016 DEGs) (**Figure 4H**), that corresponded to a -11.9% reduction in muscle mass after repeated atrophy versus recovery (**Figure 2B**). There were 79 DEGs after repeated versus initial atrophy in aged animals with a larger number of downregulated (84%) versus upregulated (16%) genes (**Figure 4I**). Most of these were shared across both periods of atrophy (2461 DEGs/75% downregulated, 3935 DEGs/61% upregulated), but several downregulated (1294 DEGs/25%) and upregulated (1550/39%) DEGs that were unique to repeated atrophy (**Figure 4J**).

Regarding gene-specific alterations, the top downregulated DEGs included genes associated with NMJ
function and stability (*Ache, Unc5a, Ntf3, Nfasc*), mitochondria and metabolism (*Park7, Perm1, Nfil3,*
Gpd1, Acss1, Mc5r, Dhrs7c, G0s2, Fbp2), muscle structure and contraction (*Lrrc38, Ky, Mylk2, Rgn*)
as well as DNA damage (*Fanca, Ddit4l2*) (**Figure 4K).** Interestingly, the myokine, *Apln*, important for
angiogenesis, muscle regeneration and ATP synthesis was among the top 5 downregulated genes after
initial TTX-induced atrophy (-9.7 FC, **Figure 4D**) (that also decreased after repeated atrophy) and was
within the top 25 downregulated genes in young adult humans after atrophy (**Figure 3H**). Concerning
the top upregulated genes, the muscle specific fuso-gene, *Mymk* significantly increased after both
atrophy (12.8 FC*,* **Figure 4D**) and repeated atrophy (18.6 FC, **Figure 4F**), implying satellite cell fusion
in attempt to repair and/or offset TTX-induced muscle wasting. Also, among the top-ranked upregulated
genes after initial atrophy was tropomyosin-binding, *Tnnt2* (11.8 FC*,* **Figure 4D)**. Embryonic *Myh8* was
one of the most upregulated DEGs after recovery (14.3 FC, **Figure 4E**) and was among the top-ranked
upregulated DEGs in young adult human muscle after repeated atrophy (**Figures 3D & F**). Further,
NMJ AChR subunit genes, *Chrnd,* and *Chrna1* were both ranked among the top upregulated genes in
aged animal muscle after both atrophy (*Chrnd* = 5.8 FC*, Chrna1* = 7 FC) and to a greater extent
repeated atrophy (*Chrnd* = 12.5 FC*; Chrna1* = 16 FC) whilst *Chrng* significantly increased only after
repeated atrophy (**Figure 4K**). Such AChR genes were also similarly upregulated in young adult human
muscle following disuse (**Figure 3D & H**). Collectively, genes *Tnnt2, Myh8, Chrna1, Chrnd* and *Chrng*,
increased in response to disuse-induced muscle wasting, irrespective of age, species and atrophy
model utilised.

Enrichment analysis of downregulated genes revealed similar pathways and GO terms after both
atrophy and repeated atrophy in aged muscle. Specifically, terms related to metabolic processes and
aerobic respiration (e.g., NAD binding / NADH dehydrogenase activity, TCA cycle, ETC and ATP
synthesis) as well as muscle-specific terms (e.g., muscle contraction, cytoskeleton in muscle cells and
sarcomere organisation) (**Figure 4L & M**). Such terms were also enriched in downregulated DEGs after
atrophy and repeated atrophy in young adult human muscle. In contrast, most of these terms were
enriched in upregulated DEGs during recovery from atrophy, suggesting they recovered after the initial
disuse, despite continued muscle wasting after TTX cessation. This suggests that genes were TTX-
sensitive during disuse rather than simply reflecting the continued muscle wasting observed during
recovery and a possible attempt to restore muscle during return to activity. This was also observed in
young adult human muscle, where expression of genes within these pathways, alongside skeletal
muscle size and strength, returned to baseline levels during recovery. Upregulated pathway enrichment
that occurred after both atrophy and repeated atrophy was evident within ribosomal structure and
subunit genes (**Figure 4L**). Interestingly, genes associated with proteasomal core and accessory
complex, ECM turnover, ubiquitin-mediated proteolysis and DNA damage pathways only increased
after repeated atrophy (**Figure 4L-M**).

Aged muscle demonstrates an exaggerated transcriptional response following repeated atrophy in
genes that are fully restored during recovery
977
Temporal clustering analysis by SOM profiling revealed the majority of genes that were downregulated
(40% cluster 3) or upregulated (35%, cluster 2) after atrophy, were restored to control levels after TTX
cessation during recovery, followed by exaggerated changes after repeated disuse (**Figure 4N**).
Specifically in cluster 3, downregulated genes after atrophy that returned to control levels during
recovery (despite continued loss of muscle mass), followed by greater reductions during repeated
atrophy, were enriched in: aerobic metabolism, TCA cycle, NAD binding/NADH dehydrogenase activity,
ETC and ATP synthesis in the mitochondria as well as in muscle structure and function (cluster 3)
(**Figures 4O & P**). Based on SOM standardised means, on average there was a 95% larger magnitude
of decreased expression of genes within cluster 3 after repeated atrophy compared with initial atrophy.
Therefore, genes in these metabolic pathways demonstrated, on average, a 95% more exaggerated
downregulation of transcripts in response to repeated atrophy, whereas young adult human muscle
demonstrated less susceptible attenuated transcriptional response in genes within the same pathways
following repeated atrophy. This suggests a negative, detrimental and exaggerated transcriptional
response to repeated atrophy in aged muscle. Importantly, these genes still recovered to control levels
during recovery, despite continued muscle wasting in aged animals. Therefore, the exaggerated
response to repeated atrophy in aged muscle was not simply due to the lack of recovery but rather an
exaggerated response to repeated atrophy having experienced earlier TTX-induced disuse atrophy.
995
Genes within cluster 2 (**Figure 4N**) that were upregulated after atrophy, returned to control levels during
recovery and were upregulated to a greater extent during repeated atrophy, were enriched in
proteosome, ECM turnover, and DNA damage pathways in aged muscle. Based on SOM standardised
means, there was an average 141% larger magnitude in upregulation of genes in cluster 2 after
repeated atrophy compared to the initial atrophy. These genes/terms were unique to aged muscle, also
demonstrating recovery of gene expression despite continued muscle wasting during the recovery
period, followed by exaggerated transcriptional and physiological responses to repeated atrophy.
1003
A proportion of genes also decreased (11%, cluster 1) or increased (14%, cluster 4) to a lesser extent
during atrophy with the largest change occurring during recovery, followed by a return to atrophy levels
during repeated atrophy (**Figure 4N**). Genes in both clusters 1 & 4 were enriched in inflammatory
pathways, ECM turnover, collagen assembly and matrix organisation (**Figure 4O & P**), suggestive that
changes in inflammatory and ECM genes were associated with continued muscle wasting in aged
muscle, despite TTX cessation.
1010

Figure 4. (**A**) Schematic depicting transcriptomic (via RNA-seq) analysis of repeated TTX-induced disuse atrophy in aged animals (created using BioRender.com). Principal component analysis (PCA) including (**B**) and excluding (**C**) sham surgery group. (**D-I**) Volcano plots depicting the number of differentially expressed genes (DEGs) that were either DOWN- (blue) or UP-regulated (orange) for each pairwise comparison. Labelled genes within volcano plots represent the top 5 DEGs according to greatest fold-change with FDR ≤ 0.05. There were no DEGs detected in sham versus pooled contralateral control limb, suggesting the surgery had no significant impact on gene transcription. The sham group was therefore excluded for all downstream analyses and pooled contralateral limb was used as the control. (**J**) Heatmap of all DEGs in at least one comparison (*i*). Magnified heatmap displaying the top 50 DOWN- (blue) and top 50 UP-regulated (orange) genes (by gene symbol) by average fold-change across all pairwise comparisons (*ii*). (**K**) Venn diagrams depicting the number of unique and common DEGs across all conditions for both UP- (top) and DOWN-regulated (bottom) DEGs. Over-representation analysis (ORA) of the top 20 enriched (**L**) pathway (KEGG, top 4 panels; REACTOME, bottom 4 panels) and (**M**) gene ontology (GO) terms for both DOWN- and UP-regulated DEGs across all comparisons ("Atrophy vs. Control", "Recovery vs. Control", "Repeated Atrophy vs. Control", "Recovery vs. Atrophy", "Repeated Atrophy vs. Recovery"). (**N**) Self-Organizing Maps (SOM) composed of a total 6379 genes that were DEGs in at least one comparison, illustrating gene cluster temporal profiles across the time-course of repeated disuse atrophy. Profiles 1 & 3 represent genes that were DOWN-regulated following the first period of atrophy whereas profiles 2 & 4 depict UP-regulated genes after initial atrophy. (**L**) Pathway and (**M**) GO analysis revealed enriched terms across all cluster profiles where most genes were

1028

1029 *Young adult rat skeletal muscle recovers from TTX-induced atrophy after longer TTX exposure and*
1030 *shorter active recovery than aged rats herein*

1031

In both young adult human and aged rat muscle, initial and repeated atrophy both resulted in
downregulation of genes associated with aerobic respiration and mitochondria terms. During return to
habitual activity however, these pathways recovered both in young adult humans and in aged rat
muscle, despite differential physiological responses. Given the optimal recovery observed in young
adult humans and the lack of recovery in aged animals, we compared the physiological and
transcriptomic response of aged rats to young adult rat muscle using the same TTX-induced atrophy
and recovery model [66]. Previously published work by our group reported sufficient recovery of muscle
mass following TTX-induced atrophy despite longer TTX exposure and shorter active recovery [66].
Specifically, young adult rats received 2 wks of TTX exposure to the common peroneal nerve (versus
1 wk in aged rats in the present study), resulting in an average -51% and -69% loss of TA muscle mass
and fCSA, respectively (**Figures 5A, B & C**). Interestingly, 1 wk active recovery (versus 1.5 wks in aged
animals) recovered TA mass and fCSA by 53 and 63%, respectively relative to the previous 2 wks
atrophy period (**Figures 5A & C**). This physiological response contrasts with aged animals herein,
where TA muscle weight and fCSA continued to decrease during a longer active recovery, despite less
atrophy observed following shorter TTX exposure (**Figures 5B-C**). Together with similar recovery of
genes within the same pathways in young adult human muscle and aged rat muscle, this suggests that
the lack of recovery in aged animals is dependent on age and not insufficient time to recover. In support
of this, overlapping transcriptome datasets from young and aged animals revealed a considerable
degree of overlap of DEGs after atrophy (77% upregulated and 72% downregulated DEGs) with a much
smaller overlap after recovery (34% upregulated and 26% downregulated) (**Figures 5D-H**).

1052
1053 **Figure 5.** (**A**) Schematic of study design (created using BioRender.com) of TTX-induced atrophy (3, 7 and 14 days/d) and subsequent recovery (7
1054 d) in young adult rats (Fisher et al.,) [66]. There was a loss of TA muscle mass (**B**) and fCSA (**C**) at 3, 7 and 14 d atrophy (-7 to -18%, -29 to -42%
1055 and -51 to -69%, respectively). In contrast to aged animals, muscle mass and fCSA was restored by 53 and 63%, respectively following 7 d recovery
1056 in young adult animals (**B & C** redrawn from Fisher et al., [66]). (**D**) Overlap of transcriptome in young adult and aged rat muscle revealed many
1057 common UP- (**E & F**) and DOWN-regulated (**G & H**) DEGs after atrophy and a larger proportion of unique genes after recovery.

1058

1059 *Integrative DNA methylome and transcriptome analysis reveals epigenetically regulated gene*
1060 *expression after repeated atrophy related to NAD$^+$ metabolism, aerobic metabolism and mitochondrial*
1061 *structure & protein turnover pathways as well as NMJ genes*

1062

Retained and enhanced transcriptional responses to repeated hypertrophic stimuli have been
associated with sustained DNA methylation alterations, thus proposing an important mechanism of
muscle memory. Therefore, we next determine whether detected DEGs after atrophy, recovery and
repeated atrophy were also associated with alterations in DNA methylation. Therefore, we performed
DNA methylome analysis in both young adult human and aged animal skeletal muscle in response to
atrophy, recovery and repeated atrophy and integrated these data with the transcriptome datasets
described above. Repeated periods of disuse atrophy and recovery led to differentially methylated DNA

regions (DMRs) that were inversely associated with the DEGs identified in both young (**Figures 6A-D**)

and aged (**Figures 6E-H**) skeletal muscle. These genes are termed **D**ifferentially **m**ethylated **DEGs**

(DmDEGs). In young adult humans, 71 DMRs on 57 genes (from cluster 1 above, **Figure 3K**) were

identified as DmDEGs with an inverse relationship of hypermethylation and decreased expression.

These DmDEGs were hypermethylated and downregulated after atrophy and repeated atrophy, with

both DNA methylation and gene expression returning to baseline levels after recovery where skeletal

muscle was restored (cluster 1, **Figures 6B*i* & C*i***). ORA analysis identified these DmDEGs as enriched

in NAD$^+$ metabolism, ETC, ATP synthesis, mitochondrial structure and mitochondrial ribosome

pathways, demonstrating that the enriched terms of genes within cluster 1 of the transcriptomic analysis

above (downregulated with atrophy, return to baseline levels during recovery and less susceptible

changes during repeated atrophy, **Figure 3K**) were also epigenetically regulated in response to disuse

atrophy and repeated atrophy. Of note, *NMRK,2* a gene involved in NAD$^+$ biosynthesis was among the

top downregulated gene after both periods of disuse in young adult humans), was among the

epigenetically regulated genes in this profile. Specifically, *NMRK2* was hypermethylated and

downregulated during both periods of atrophy and was hypomethylated with increased transcription

towards baseline levels during recovery. In addition to these hypermethylated and downregulated

genes, there were 189 DMRs on 101 genes that were hypomethylated and upregulated after both

atrophy and repeated atrophy (cluster 2, **Figures 6B*ii* & C*ii* & C*iii***). There was no significantly enriched

pathway or GO terms for this cluster, thus suggestive that most of these genes likely have divergent

molecular, cellular and biological roles. However, several interesting hypomethylated and upregulated

genes emerged from this cluster. Specifically, the AChR genes, *CHRND* and *CHRNA1* (that were

upregulated after atrophy and/or repeated atrophy across both human and rat muscle transcriptomes,

Figure 3D-H), demonstrated the largest hypomethylation and increased expression after repeated

atrophy (**Figures 6B*ii* and C*ii***). These genes were among the few DmDEGs in young human muscle

that demonstrated increased susceptibility and exaggerated responses to repeated atrophy. Finally,

genes *NR4A1* and *PRKAG2* (AMPK gamma2) were downregulated after atrophy (despite no change in

DNA methylation) followed by hypermethylation and sustained suppression of transcription during

recovery and repeated atrophy (cluster 3, **Figure 6B*iii***). Thus, suggestive that the retained reductions

in expression of these genes during recovery is likely due to associated hypermethylation, despite

restoration of skeletal muscle upon returning to habitual activity. The full DNA methylome data to

repeated atrophy in young adult human muscle is also available in the supplementary files

(**Supplementary Figure 4).**

1102

In aged rat muscle, integrative analysis of DNA methylome and transcriptome (**Figure 6E**) revealed 41

DMRs on 33 unique genes were hypermethylated and downregulated after atrophy (cluster 1, **Figure**

6Fi & Gi), hypomethylated and upregulated upon TTX cessation during the recovery period, and finally

hypermethylated and downregulated after repeated atrophy. As with the young human muscle, these

genes were associated with aerobic metabolism/respiration, electron transport and mitochondrial

components (**Figure 6H**). Integrative analysis also revealed 40 DMRs on 28 unique genes (cluster 2,

Figure 6Fii and 6Gii) that were hypomethylated and upregulated after atrophy, hypermethylated after

recovery, and hypomethylated and upregulated again after repeated atrophy. ORA analysis revealed
no significantly enriched pathways or GO terms for this cluster. There were also 26 DMRs on 19 unique
genes (cluster 3, **Figure 6Fiii & Giii**) that were hypermethylated and downregulated after atrophy,
hypermethylated yet upregulated during recovery, and hypomethylated and upregulated after repeated
atrophy, with no significantly enriched pathways or GO terms. Interestingly, 43 DMRs on 21 unique
genes (cluster 4, **Figure 6F*iv* & G*iv*)** demonstrated hypermethylation and downregulation after atrophy,
with retained hypermethylation and a continued decrease in gene expression during recovery, followed
by subsequent hypomethylation and increased gene expression after repeated atrophy. These genes
were enriched in JAK-STAT signalling genes (cluster 4, **Figure 6H**) and therefore demonstrated
retained epigenetic and transcriptional signatures after the initial atrophy that were retained during
recovery yet were oppositely hypomethylated and upregulated after repeated atrophy.

1121

Despite the differential physiological response to repeated atrophy and recovery, both young adult
humans (cluster 1, **Figures 6B*i*, C*i*, D)** and aged animals (cluster 1, **Figure 6F*i*, G*i*, H**) demonstrated
associated DNA methylation and gene transcription of genes enriched within the same pathway and
GO terms related to aerobic metabolism/respiration, electron transport and mitochondria (**Figures 6D**
& H). However, the majority of these DmDEGs were specific to young human or aged rat muscle
(**Figure 6I**), despite enrichment of the same metabolic pathways across species and models of atrophy
(**Figure 6J**). Further cross-species comparison of the shared transcriptomic and epigenetic responses
to repeated disuse in young adult human and age animal muscle via UpSet plots (**Supplementary**
Figure 5), identified several common up (**Supplementary Figure 5A**) and downregulated DEGs
(**Supplementary Figure 5B**) as well as hypo- (**Supplementary Figure 5C**) and hyper-methylated
DMRs (**Supplementary Figure 5D**). It was also apparent that out of the 761 DEGs used for the SOM
temporal profiling in human muscle, 54% (414 out of 761) of the genes were also differentially
expressed in the aged rat muscle (**Supplementary Figure 5E**). Pathway (**Supplementary Figure 5F**)
and GO (**Supplementary Figure 5G**) analyses suggested these shared genes were also related to
oxidative metabolism, NAD$^+$, TCA cycle, ETC, ATP synthesis and mitochondrial function. Among these
shared DEGs, there were 406 DMRs on 189 shared DEGs in young adult human muscle
(**Supplementary Figure 5H)** that were associated with oxidative metabolism, ETC, ATP synthesis and
mitochondria (**Supplementary Figure 5I & J**). In aged animals, these there were 56 DMRs on 16
shared DEGs (**Supplementary Figure 5K**) that were related to mitochondria and RNA splicing
(**Supplementary Figure 5L**). Overall, these data demonstrate epigenetic regulation of shared genes
associated with oxidative metabolism across species in response to atrophy, recovery and repeated
atrophy. The full DNA methylome data to repeated atrophy in aged rat muscle is also available as
supplementary information (**Supplementary Figure 6).**

Figure 6. Integrative DNA methylome and transcriptome analyses in young adult human (**A-D**) identified 57 hypermethylated and

downregulated genes after both periods of disuse that were returned to baseline levels after recovery (cluster 1, **Bi & Ci**); 101

hypomethylated and upregulated genes after both periods of disuse that also returned to baseline levels after recovery (cluster

2, **Bii & Cii**); 2 downregulated/hypermethylated genes after initial atrophy that were retained during recovery (cluster 3, **Biii & Ciii**). There were significant pathway and GO terms for cluster 1 in young human muscle of hypermethylated/downregulated genes after atrophy, restored during recovery, and hypermethylated/downregulated after repeated atrophy enriched in aerobic metabolism/respiration, NAD$^+$ activity, ETC, ATP synthesis, mitochondrial and ribosome terms (**D**). Integrative analyses in aged rat muscle (**E-G**) revealed 33 genes (**Fi & Gi,** cluster 1) hypermethylated/downregulated after atrophy, hypomethylated/upregulated upon TTX cessation during recovery, and hypermethylated/downregulated after repeated atrophy; 28 unique genes (**Fii & Gii,** cluster 2) were hypomethylated/upregulated after atrophy, hypermethylated/restored after recovery, and hypomethylated/upregulated again after repeated atrophy (**Fiii & Giii,** cluster 2); 19 unique genes (cluster 3) that were hypermethylated/downregulated after atrophy, hypermethylated yet upregulated during recovery, and hypomethylated/upregulated after repeated atrophy; 21 unique genes (**Fiv and Giv,** cluster 4) were hypermethylated/downregulated after atrophy, with retained hypermethylation and a continued decrease in gene expression during recovery followed by subsequent hypomethylation and increased gene expression after repeated atrophy. Significant enriched pathway and GO terms (**H**) for cluster 1 were related to aerobic metabolism/respiration, electron transport and specifically mitochondrial structure (for cluster 1) and JAK-STAT enrichment (in cluster 4). Average DMR methylation is depicted in aged animal heatmaps whenever there is >1 DMR for a given gene (**G**). Average mRNA expression is also presented where there were multiple transcript variants (**G**). UpSet plot analysis of young adult human and aged rat muscle gene clusters revealed only a few epigenetically regulated genes that overlapped between both models (**I**). However, there were 9 overlapping enriched pathway / GO terms between human and animal muscle (**J**). Note that 'DMR Methylation' is standardised percentage DMR methylation derived from SOM profiling analysis and 'Gene Expression' is representative of standardised LS means from SOM profiling analysis (**B & F**). BP = biological process, CC = cellular component, MF = molecular function.

Repeated atrophy in aged muscle leads to significant transcriptional perturbations of NAD+ biosynthesis genes resulting in reduced NAD+ content

Genes associated with NAD/NADH metabolism were enriched as downregulated and hypermethylated after atrophy and repeated atrophy in both young adult (**Figure 7A-D)** and aged rat (**Figure 7E-I)** muscle. Moreover, the NAD$^+$ biosynthetic gene, *NMRK2,* was the among the top downregulated genes after both periods of atrophy with inversely associated DNA across the time course of atrophy, recovery and repeated atrophy. Genes enriched in these pathways were also restored during the recovery period. Targeted mRNA analysis (via qPCR) of *Nmrk2* in aged rat muscle revealed a non-significant reduction in *Nmrk2* expression after atrophy, followed by a significant increase during recovery and a larger significant reduction after repeated atrophy (**Figure 7E & G**). Given the predominant role of *NMRK2/Nmrk2* in synthesising NAD$^+$ coupled with over-representation of NAD/NADH metabolism related enrichment terms for downregulated genes during both periods of atrophy, we next interrogated gene expression of other NAD$^+$ biosynthesis genes and whether such alterations corresponded with a reduction in NAD$^+$ content in both young adult (**Figure 7A-D**) and aged muscle (**Figure 7E-I**). In young adult humans, while most of these NAD$^+$ genes were downregulated after both periods of atrophy, *NMRK2* was the only significantly downregulated DEG out of the 18 genes analysed (**Figures 7A-C**). Furthermore, *NMRK2* was the most upregulated gene across all DEGs after recovery from atrophy in humans (**Figure 3E**) and was among the identified DmDEGs that was downregulated less after repeated versus initial atrophy. Given *NMRK2* was the only significantly downregulated NAD$^+$ gene, this resulted in no significant changes in NAD$^+$ content (**Figure 7D**). In contrast, half of the 14 NAD$^+$ genes assessed in aged animals were significantly differentially expressed where most genes were downregulated after atrophy, either unchanged/restored or increased after recovery, followed by greater

1193 downregulation after repeated atrophy (**Figure 7E-G**), thus resulting in significantly reduced absolute
1194 NAD$^+$ content in the left (L) TTX versus right (R) control limb after repeated atrophy (**Figure 7H & I**).
1195

1196 *Human post-atrophy MuSC display greater differentiation capacity, responsiveness to NR*
1197 *supplementation as well as lower mtDNA content relative to pre-atrophy MuSC*
1198

Given *NMRK2* was among the top downregulated genes after atrophy and repeated atrophy in humans,
and is phosphorylated (and thus activated) by the vitamin B$_3$ derivative, nicotinamide riboside (NR), we
next supplemented muscle derived cells (MuSC) derived from pre- and post-atrophy human biopsies
with NR in an attempt to restore potentially impaired *NMRK1/2* expression and myogenesis *in vitro*
(**Figure 7J-M**). Although *Nmrk2* expression and NAD$^+$ content was significantly decreased after disuse
in aged animals, we could not isolate MuSC to perform such experiments in aged muscle. Similar
proportions of myoblast (desmin+) to fibroblasts (TE7+) were observed between pre- and post-atrophy
MuSC, thus negating possible effects of cell populations (**Figure 7L**). Despite limited myotube formation
in the MuSC, likely due to a higher proportion of fibroblasts ($\approx$ 60%) versus myoblasts ($\approx$ 60%) (**Figures**
7K & L), post-atrophy MuSC were more responsive to NR supplementation, where myotube size
significantly increased in NR treated post-atrophy cells when compared to 7d differentiation (p = 0.03,
Figure 7M). This was supported by lower *MYOD (early differentiation)* and greater *MYOG*
(fusion/differentiation) gene expression in NR-supplemented MuSC, coupled with greater (albeit non-
significant *NMRK1* and *NMRK2* expression (**Figures 7N-Q**). Interestingly, mtDNA content
demonstrated lower mean (albeit non-significant) mtDNA in post- versus pre-atrophy MuSC (**Figures**
7R-U). Collectively, post-atrophy MuSC were more responsive to NR supplementation that altered
NMRK and *MYOG* expression and increased myotube size.
1216
1217

Figure 7. mRNA expression of NAD$^+$ biosynthesis genes and corresponding NAD$^+$ content in young adult human (**A-D**) and aged rat (**E-I**) skeletal muscle. (**A & E**) Schematic representation of Preiss-Handler and salvage NAD$^+$ biosynthetic pathways (redrawn from Damgaard & Treebak, 2023 [91] using BioRender.com). Key metabolites are depicted as black text. Genes encoding NAD$^+$ biosynthesis enzymes are shown in either orange (if non significantly expressed) or red text (significantly downregulated after atrophy or repeated atrophy). In young adult humans, *NMRK2* was the only significantly downregulated DEG out of the 18 NAD$^+$ genes (**B & C**), which did not lead to significant alterations in NAD$^+$ content (**D**). In aged animals, most of the 14 NAD+ genes assessed were significantly differentially expressed after atrophy and repeated atrophy and were restored upon recovery (**F & G**), resulting in significantly reduced absolute (**H**) NAD$^+$ content in the left (L) TTX versus right (R) control limb after repeated atrophy. Relative NAD$^+$ content (**I**). Nicotinamide riboside (NR) supplementation (0.5 mM) in human pre- and post-atrophy MuSC (*n* = 4) (J-K). (**K**) Fluorescence microscopic tile scan (3 × 3) images of MuSC labelled for myoblasts (desmin, red), fibroblasts (TE-7, green) and DNA (DAPI, blue), acquired at 10× magnification (scale bar = 200 μm). (**L**) Myoblasts (≈ 40%) and fibroblast (≈ 60%) proportions were similar between pre- and post-atrophy MuSC. NR supplementation resulted in significantly increased myotube area at 10 versus 7 d (*p* = 0.03) in post-atrophy MuSC only (**M**).

Mitochondrial gene expression and DNA content is significantly impaired after repeated atrophy in young adult human muscle which is exacerbated in aged animal muscle

In addition to enrichment of NAD/NADH metabolism related genes, the majority of genes were also associated with mitochondrial structure, function, assembly and turnover in both young adult human and rodent muscle. Specifically, genes involved in aerobic respiration (and thus ETC and oxidative phosphorylation) and mitochondrial structural components were hypermethylated and downregulated after both periods of atrophy, with the majority of these genes returning to baseline levels in both human and rat muscle, despite restoration of muscle in humans and continued wasting in aged rats. Furthermore, these genes demonstrated further reductions in transcription in aged rat muscle, whereas these genes were less susceptible to reductions in young adult human muscle following repeated disuse. Given these data, we next assessed gene expression of mitochondrial fission, fusion, transport and biogenesis genes-specific genes, as well as mRNA and protein abundance of ETC protein complexes I-V. Finally, we assessed mtDNA copy number and citrate synthase protein abundance as markers of mitochondrial content/density. In young adult human muscle (**Figure 8A**), the majority of the 66 genes profiled across ETC complexes I-V were significantly suppressed after both periods of atrophy where most genes either returned to baseline levels or increased after recovery (**Figure 8B**). This resulted in significant changes occurring after both atrophy and repeated atrophy in 24 (out of 66) ETC genes; within complex I (7 out of 32 genes), III (3 out of 8 genes), IV (8 out of 13 genes), and V (6 out of 9 genes) with non-significant changes in CII (**Figure 8B**). Concerning genes related to mitochondrial fission and mitophagy (*FIS1, DRP1*), fusion (*MFN1/2, OPA1*), transport (*SLC25A51, SLC25A4)* and biogenesis (*PPARGC1A, TFAM*), 8 out of 9 of these genes non-significantly decreased after atrophy and repeated atrophy, all of which returned to baseline levels or non-significantly increased after recovery (**Figure 8C**). There was a significant reduction in *PPIF*, a mitochondrial membrane gene controlling calcium overload and fatigue resistance in muscle (**Figure 8C**). Given the large transcriptional perturbations in ETC-specific genes across all 5 complexes in both atrophy periods, we next assessed ETC (I-V) protein abundance. The reductions in gene expression did not correspond to significant changes in ETC protein complex abundance (**Figure 8D**). Interestingly however, mtDNA content (**Figure 8E**) and citrate synthase protein abundance (**Figure 8F**) significantly decreased after repeated atrophy only in young adult humans. Collectively, these data suggest that the large transcriptional perturbations of genes enriched in mitochondrial-specific pathways and GO terms culminated in impaired mitochondrial content and density following repeated disuse only in young adults despite restoration of these parameters and skeletal muscle during recovery.

In aged animals, most genes across all ETC complexes were differentially expressed (**Figure 8H & I**).

Specifically, 57% (38 out of 67) of genes encoding complex´s I (13 out of 30), II (4 out of 4), III (7 out of

9), IV (6 out of 14) and V (8 out of 10 genes) were significantly downregulated after atrophy and/or

repeated atrophy (**Figure 8H & I**). Similarly to young adult humans, the majority of these genes

recovered during TTX cessation despite continued loss wasting during return to activity. Furthermore,

Sdhb, Cox6a1, Cox8a and Atp5pf were significantly upregulated during recovery (**Figure 8I**). In contrast

to young adult humans, aged rat muscle demonstrated a greater average reduction across all 67 genes

following repeated atrophy (-1.72 FC) versus the initial atrophy period (-1.52 FC), despite a similar

average FC between recovery and control (0.1 FC). Mitochondrial fusion (*Mfn1/*2), transport (*Slc25a4*)

and biogenesis (*Ppargc1a*) genes also significantly decreased after atrophy and/or repeated atrophy

whereas *Tfam* significantly increased after repeated atrophy (**figure 8J**). Most genes recovered during

TTX cessation, and *Opa1* and *Ppif* genes both significantly increased during recovery (**Figure 8J**). As

with the ETC genes described above, there was a greater average reduction across all genes (excluding

Tfam which increased) after repeated atrophy (-2.23 FC) compared to previous atrophy (-1.56 FC)

relating to a 43% larger reduction after repeated atrophy compared to initial atrophy, where most genes

recovered upon returning to activity (average 0.72 FC) (**Figure 8I & J**). At the protein level, there were

no significant changes in protein abundance of ETC complexes I-V nor citrate synthase across the

conditions analysed (e.g., sham, atrophy and recovery respectively, **Figure 8L)**. Finally, there was a

reduction in mtDNA content during atrophy (-18%), recovery (-41%) and the greatest reduction after

repeated atrophy (-50%) when *Mt-Rnr1* and *Mt-Rnr2* genes were combined (**Figure 8M).**

1292

1293

1294

Young adult human

A

B *nDNA encoded genes*
Complex I

Complex II *Complex III* *Complex IV* *Complex V*

C Mitochondrial fission, fusion, transport & biogenesis genes

D *Total Protein* *OXPHOS*

CI-NDUFB8 CII-SDHB

CIII-UQCRC2 CIV-MTCO1 CV-ATP5A

E *mtDNA content*

MT-RNR2 DLOOP

CYTB pooled

F *Total Protein* *Citrate Synthase*

Aged animal

H

I *nDNA encoded genes*
Complex I

Complex II *Complex III* *Complex IV* *Complex V*

J Mitochondrial fission, fusion, Transport & biogenesis genes

K *OXPHOS*

CI-NDUFB8 CII-SDHB

Total Protein

CIII-UQCRC2 CIV-MTCO1 CV-ATP5A

L *Citrate Synthase*

Total Protein

M *mtDNA content*

Mt-Rnr1 Mt-Rnr2 pooled

 Figure 8. Mitochondrial response to repeated periods of disuse in young adult human muscle (**A-G**) and aged rat muscle (**H-M**) skeletal muscle. (**A**
& H) Schematic depicting a specific focus on mitochondrial content as well as mitochondrial activity, fusion and fission processes (created using
BioRender.com). (**B & I**) Temporal transcriptional response of ETC complex I-V genes where most genes significantly decreased after atrophy
and/or repeated atrophy, and recovered upon returning to normal activity in both adult humans (**B**) and aged animals (**I**). A similar transcriptional
response occurred for mitochondrial fission, fusion, transport and biogenesis-related genes across species (**C & J**). There were no significant
changes in ETC complex (I-V) protein abundance in both human (**D**) and rat (**K**) muscle. However, mtDNA content and citrate synthase protein
significantly decreased after repeated atrophy in young adult humans (**E & F**). Despite no significant changes in citrate synthase across the
sham/atrophy/repeated atrophy conditions analysed in aged animals (**L**), there was a reduction in mtDNA content in response to atrophy, recovery
and repeated atrophy with the largest loss of -50% after repeated atrophy (**M**). Gene expression data presented as mean (signed) FC ± absolute
95% CI (absolute upper and lower limits). *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$.

Discussion

Summary of main findings

Skeletal muscle exhibits a molecular memory of repeated disuse-induced atrophy. In young adult humans, repeated immobilization induced a comparable loss of skeletal muscle with full recovery between periods of disuse. Repeated disuse triggered a *protective* molecular resilience in which transcriptional perturbations of genes associated with oxidative metabolism and mitochondria (structure and function) were attenuated during recurrent atrophy. In contrast, aged muscle displayed a *detrimental* molecular susceptibility, characterised by further suppression of genes within similar pathways following repeated disuse, despite transcriptional recovery and continued muscle wasting upon returning to normal habitual activity. Proteasomal, ECM, and DNA damage associated genes were uniquely elevated following repeated atrophy in aged muscle. Methylome analyses revealed conserved epigenetic regulation of aerobic metabolism and mitochondrial genes across species, along with locus-specific memory genes including *NR4A1* and acetylcholine receptor subunit genes. Notably, nicotinamide riboside (NR) supplementation in human primary MuSC partially restored *NMRK1/2* expression and imrpoved myotube size in MuSC derived post-atrophy.

Molecular memory of repeated disuse-induced skeletal muscle atrophy

Skeletal muscle can retain an epigenetic 'memory' of prior growth, as demonstrated by persistent hypomethylation during detraining in humans and mice [33, 36, 37], and amplified gene expression upon retraining [33, 34, 36, 37]. Our findings now extend this concept to a clinically relevant negative stimulus. Specifically, disuse-induced atrophy which commonly occurs after injury, hospitalization, bed rest, surgery, or spaceflight. Crucially, we identified an age-dependent molecular memory: a *protective* transcriptional attenuation in young muscle, where repeated disuse elicited blunted transcriptional responses, and a *detrimental* memory in aged muscle, where repeated atrophy exaggerated suppression of NAD⁺, aerobic metabolism and mitochondrial gene pathways.

Attenuated transcriptional memory in young adult humans and a detrimental exaggerated transcriptional memory in aged animal muscle

In young adult human muscle, genes suppressed during initial atrophy largely recovered and were less perturbed by repeated disuse, indicating a transcriptional protection or resilience. These responses

predominantly involved oxidative metabolism and mitochondrial pathways, including TCA cycle, NAD^+/NADH, ETC, ATP synthesis. In contrast, aged animal muscle showed the opposite pattern, in which these same pathways were further suppressed during repeated atrophy despite transcriptional recovery after the initial disuse, accompanied by unique activation of proteasomal, ECM, and DNA damage gene pathways after repeated atrophy. These signatures are consistent ageing-associated skeletal muscle deficits in metabolic regulation, proteostasis, NMJ dysfunction (discussed below) and mitochondrial function [10, 19, 92]. Importantly, expression of these gene pathways recovered even though muscle mass continued to decline in aged muscle, confirming that the exaggerated transcriptional and physiological responses reflected a detrimental molecular memory rather than incomplete recovery of these molecular pathways.

Mitochondrial vulnerability and loss of mtDNA with repeated disuse atrophy

Repeated disuse caused significant loss of mtDNA that was not evident after the initial atrophy period in either young adults or aged animals, with the most pronounced decline observed in aged muscle. These findings align previous bed rest studies demonstrating broad downregulation of mitochondria and reduced content/respiration during unloading [24, 64, 93]. Our transcriptomic data also revealed widespread suppression of electron transport chain and mitochondrial genes during atrophy and repeated atrophy, followed by recovery upon reloading/a return to activity. Although aged muscle exhibited the greatest mitochondrial vulnerability, young muscle also exhibited a marked loss of mtDNA loss after repeated atrophy, despite the transcriptional protection discussed above. This suggests that repeated disuse increases susceptibility to mitochondrial loss, raising the question as to whether further atrophic episodes of disuse atrophy could drive even young muscle toward the detrimental molecular program observed in aged muscle.

A similar epigenetic program in both species represses aerobic metabolism genes

Integrative methylome and transcriptome analyses revealed differentially methylated, differentially expressed genes (DmDEGs) enriched in oxidative metabolism and mitochondrial structure/turnover in both humans and rats. These genes were hypermethylated and downregulated during atrophy and repeated atrophy, returning toward baseline on recovery. This extends prior observations of disuse-linked methylation (e.g., nNOS hypermethylation [65], *Chrna1* hypomethylation/upregulation [66]) and contrasts with the hypomethylating effects of exercise training, higher physical activity and aerobic fitness associated with a 'younger' skeletal muscle methylome [36, 37, 39, 48, 57, 58, 59, 60, 63].

Neuromuscular signalling pathways in response to repeated atrophy: Epigenetic priming of the AChR subunit genes

Repeated disuse increased expression of acetylcholine receptor subunits, *CHRNA1, CHRND* in young human muscle, alongside hypomethylation at these loci, and similarly elevated *Chrna1, Chrnd, and Chrng* expression in aged rat muscle. Although potentially compensatory, such extra-junctional expression, especially of the fetal-type gamma subunit (*Chrng*), is typically linked to NMJ dysfunction

rather than stabilization [10]. In young muscle, this response emerged only after repeated atrophy,
coinciding with the strongest hypomethylation, suggesting an epigenetic memory within AChR genes.
Aged muscle demonstrated over a doubling in the upregulation of *Chrna1* (16-fold) and *Chrnd* (12.5-
fold) after repeated disuse compared with initial disuse (*Chrna1* 7-fold, *Chrnd*-5.8 fold), and only
significant elevation of *Chrng* after repeated atrophy. Whilst we did not detect hypomethylation of these
genes in aged rat muscle, prior studies have also reported *Chrna1* hypomethylation in young adult rats
utilised the same model of disuse [66]. Therefore, despite general transcriptional resilience, young muscle
still exhibited exaggerated AChR gene responses to repeated atrophy. Further, the exaggerated
response in aging muscle is consistent with aging studies associating poor recovery with NMJ instability
and ER stress [10, 26, 94], and imply that repeated stress, even in young muscle, may epigenetically prime
these genes, heightening vulnerability to future disuse.

NR supplementation partially restored NMRK2 levels and increased myotube size in human MuSC
derived post-atrophy

NMRK2, a key NAD^+ salvage enzyme, was among most downregulated genes after both periods of
atrophy in young adult humans and was significantly downregulated in aged rat muscle after repeated
atrophy only, with broader NAD^+/NADH metabolism pathway repression across species, most
pronounced in aged muscle. In primary human MuSC derived post-atrophy, nicotinamide riboside (NR)
partially restored *NMRK2* and *MYOG* expression and improved myotube size, suggesting NAD^+
salvage as a potential target. However, whilst this provides initial *in-vitro* evidence, preclinical work
shows intravenous NR elevates muscle NAD^+, while oral NR trials report mixed efficacy [91, 95]. Rigorous
clinical studies in older, immobilization-prone populations are essential before considering NR as a
viable therapeutic intervention.

ECM remodelling during recovery and repeated atrophy in aged muscle

1397
Only aged muscle upregulated genes linked to extracellular matrix (ECM), collagen organization,
collagen binding and sarcomere structure during recovery from initial disuse. This suggests structural
remodelling as a compensatory response, which may stabilize tissue but also increase stiffness [96, 97], a
hallmark of muscle aging, and therefore perhaps impair functional recovery. Further, ECM turnover,
proteasomal pathways, and ubiquitin-mediated proteolysis were exclusively enriched after repeated
atrophy in aged muscle, reinforcing the concept of compromised resilience to repeated disuse with age.

1404
1405 *The NR4A genes are important metabolic regulators of repeated disuse atrophy in young human muscle*

1406
NR4A1, a key energy metabolism gene, was hypermethylated and remained downregulated even after
full recovery in young muscle, while *NR4A3* showed the largest reduction of all altered genes (ranked
1st out of 384). Temporal profiling confirmed both genes retained suppression post-atrophy, indicating
transcriptional memory. NR4A genes are highly responsive to exercise and inactivity [98]; *NR4A1* is
oppositely hypomethylated and upregulated after high-intensity exercise [46], and *NR4A3* overexpression

in rats enhances mitochondrial content, ATP production, and type-IIA & IIX muscle fibre protein content

[99, 100, 101]. Knockdown of *NR4A3* impairs glucose metabolism and protein synthesis in skeletal muscle

[102]. Collectively, these findings nominate *NR4A* genes as candidate targets to improve recovery and

protect against future disuse atrophy.

Limitations

This study provides a comprehensive physiological and molecular analysis of repeated disuse atrophy

across young adult human and aged rat skeletal muscle. However, several limitations should be

acknowledged. 1) Interpretation of molecular protection vs. atrophy outcomes: While we observed

transcriptional attenuation in young human muscle following repeated disuse, this did not translate into

attenuated muscle mass loss. The term "protective molecular memory" refers specifically to reduced

transcriptional responsiveness in metabolic pathways, not an attenuated loss in skeletal muscle size

due to repeated atrophy. Further, whilst there was a predominance of molecular attenuation in young

human muscle to repeated atrophy, there were also more susceptible losses, such as further mtDNA

content and exaggerated increases in AChR subunit gene expression (discussed above). However, this

raises the interesting question as to whether further atrophic episodes of disuse atrophy could drive

even young muscle toward the detrimental program observed in aged muscle. 2) Coss-species

comparisons and protocol differences: Although both models were analysed independently before

integration, differences in species, disuse protocols (immobilization vs. TTX), recovery durations, and

age must be considered when interpreting cross-species comparisons. Nonetheless, the conserved

transcriptional and epigenetic responses across species, particularly in metabolic, mitochondrial and

NAD$^+$ pathways, are notable and warrant further investigation. 3) Recovery protocol and activity

monitoring: In aged rats, molecular pathways recovered during TTX cessation despite continued muscle

wasting, suggesting that the transcriptional response was TTX-atrophy dependent rather than a

consequence of incomplete recovery. Further, we present a comparison of the aged rats with young

adult rats [66] that recover half of their muscle mass following a longer period of TTX-atrophy (and thus

greater muscle wasting than aged animals herein) in a shorter recovery period. However, whilst rat

activity was confirmed by the animals being able to perform dorsiflexion, activity levels during recovery

was not directly measured. Future studies should incorporate behavioural tracking or respirometry in

animals and make further control for nutrition and activity during human immobilization considering

wearables (accelerometers) or bed-rest protocols to better characterize recovery dynamics and

disentangle molecular recovery from functional outcomes. 4) Methodological constraints and data

integration: The integration of multi-omics data across species and models introduces complexity. While

individual analyses are robust, interpretation of age-related effects must consider differences. For

example, due to limited fibre number in the cross-sections from human samples we also inferred fibre

proportions using MYH and MYL gene expression data. Further, ETC protein data were not available

for repeated atrophy in aged rats, and ETC protein changes were not statistically significant despite

transcriptomic, NAD$^+$ content and mtDNA evidence of impairment. 5) Translational scope and *in vitro*

validation: NR supplementation partially restored *NMRK2* expression and myotube size in post-atrophy

human MuSCs, suggesting a potential therapeutic avenue. However, these findings are limited to *in*

vitro experiments in young adult cells. Further studies are needed to validate NAD$^+$ salvage
interventions in aged muscle and *in vivo* models of repeated disuse. 6) Epigenetic interpretation and
functional relevance: While DNA methylation changes were inversely associated with gene expression
in key metabolic pathways, functional consequences of these epigenetic modifications remain to be
fully elucidated. Future work should explore whether these methylation signatures directly modulate
transcriptional memory or serve as biomarkers of prior disuse.

Future mechanistic and clinical directions

We propose mechanistic and translational studies that address testable hypothesis, such as: (i) Does
enhancing NAD$^+$ salvage attenuate the maladaptive transcriptional response to repeated disuse in aged
muscle? ii) Does exercise 'preconditioning' [103, 104] or training reprogram methylation signatures to
prevent negative molecular memory in the nuclear [39, 57] and mitochondrial genome [63]? (iii) does targeted
activation of *NR4A* family and *NMRK2* genes during atrophy or recovery prevent or offset disuse-
induced atrophy and/or promote recovery from atrophy?

Conclusion

Disuse atrophy can be remembered at the molecular level in skeletal muscle. It is predominantly
protective in young skeletal muscle yet detrimental in aged skeletal muscle. This memory is partially
epigenetically encoded at the single locus level and converges in aerobic metabolism, mitochondrial
and NAD$^+$ pathways, and nominates molecular targets to preserve muscle mass and function after
recurrent immobilization or disuse.

1470
1471 **Acknowledgements**
1472

This research was funded by the Research Council of Norway (RCN) (314157) awarded to principal
investigator Adam P. Sharples, post-doc fellow Daniel C. Turner and co-investigators Sue Bodine,
Jonathan C. Jarvis, David C. Hughes, Daniel J. Owens, Truls Raastad and Olivier Seynnes. We thank
Olivier Seynnes for his critical expertise and input into the funding application, training & supervision of
post-doctoral researcher for the design, testing and analysis of the human experiments and associated
physiological data acquisition and analysis (muscle size/CSA & strength). We thank Jostein Hallèn, Siri
T. Dalen, Hege N. Ødemark and Emil Løvstakken (NIH) for assistance with muscle biopsies and tissue
processing; Josephine A. Eyrich (UCPH) for assisting with the measurements of NAD$^+$ content; Rune
Enger, as the medical lead for the human biopsies; Enovis/DJO LLC for kindly donating the knee braces
and to; ChromaDex Inc. for providing nicotinamide riboside for *in vitro* studies. Finally, sincere gratitude
to all participants who volunteered to participate in this study that included two periods of muscle
wasting.

1485
1486
1487

Author contributions

Adam P. Sharples conceived the project, secured funding as principal investigator, and led the human experimental design, ethical approvals, data/finance management, dissemination, and oversaw analysis of all human and rat experiments, and supervised postdoctoral fellow Daniel C. Turner. Daniel C. Turner conducted most of the experimental work, including recruitment, human testing, biopsy processing, wet lab work, data collection, analysis, visualization, dissemination and manuscript drafting. Jonathan C. Jarvis supervised and undertook the animal intervention studies with Hazel Sutherland and provided vital input into the funding application and expertise in experimental design, analysis and interpretation. Sue C. Bodine, David C. Hughes, Daniel J Owens, and Truls Raastad were co-investigators in the funding application and heavily involved in study design, biopsy sampling, interpretation of results, and manuscript preparation. Stian F. Christiansen undertook experimental work for the mtDNA analysis in MuSC. Jonas T. Treebak and Emilie D. Kellezi undertook NAD$^+$ content experiments. Eva Wozniak, Charles Mein and James Boot undertook RNA-seq and RRBS/RREM wet lab experiments and pre-processing of RNA-seq, RRBS/RREM and bioinformatics. Max Ullrich assisted in omics analysis and visualization. All authors contributed significantly to the final manuscript.

Declarations and conflicts

The authors declare no conflicting interests. ChromaDex provided the nicotinamide riboside for *in-vitro* experiments but had no influence on the study.

Data availability

RNA-seq and RRBS/RREM data will be available via the Gene Expression Omnibus (GEO) repository (or equivalent once accepted for publication. Any other source of processed data is available on reasonable request to the corresponding authors.

Software and Code availability

All analysis code used for preprocessing, normalization, differential expression/methylation, enrichment and visualization of the data is freely available and references and links have been provided in the methods. All code that is not freely available is available on reasonable request to the corresponding authors.

References

1529 **References**

1. Hughes VA, *et al.* Longitudinal muscle strength changes in older adults: influence of muscle mass, physical activity, and health. *J Gerontol A Biol Sci Med Sci* **56**, B209-217 (2001).

2. Cruz-Jentoft AJ, Landi F, Topinkova E, Michel JP. Understanding sarcopenia as a geriatric syndrome. *Current opinion in clinical nutrition and metabolic care* **13**, 1-7 (2010).

3. Beaudart C, Zaaria M, Pasleau F, Reginster J-Y, Bruyère O. Health Outcomes of Sarcopenia: A Systematic Review and Meta-Analysis. *PLOS ONE* **12**, e0169548 (2017).

4. Fried LP, *et al.* Frailty in Older Adults: Evidence for a Phenotype. *The Journals of Gerontology: Series A* **56**, M146-M157 (2001).

5. Rejc E, *et al.* Loss of maximal explosive power of lower limbs after 2 weeks of disuse and incomplete recovery after retraining in older adults. *J Physiol* **596**, 647-665 (2018).

6. Pisot R, *et al.* Greater loss in muscle mass and function but smaller metabolic alterations in older compared with younger men following 2 wk of bed rest and recovery. *J Appl Physiol (1985)* **120**, 922-929 (2016).

7. Kramer IF, *et al.* Extensive Type II Muscle Fiber Atrophy in Elderly Female Hip Fracture Patients. *J Gerontol A Biol Sci Med Sci* **72**, 1369-1375 (2017).

8. Suetta C, *et al.* Effects of aging on human skeletal muscle after immobilization and retraining. *J Appl Physiol (1985)* **107**, 1172-1180 (2009).

9. Baehr LM, West DWD, Marshall AG, Marcotte GR, Baar K, Bodine SC. Muscle-specific and age-related changes in protein synthesis and protein degradation in response to hindlimb unloading in rats. *J Appl Physiol (1985)* **122**, 1336-1350 (2017).

10. Baehr LM, *et al.* Age-related deficits in skeletal muscle recovery following disuse are associated with neuromuscular junction instability and ER stress, not impaired protein synthesis. *Aging (Albany NY)* **8**, 127-146 (2016).

11. Hvid L, *et al.* Effects of aging on muscle mechanical function and muscle fiber morphology during short-term immobilization and subsequent retraining. *J Appl Physiol (1985)* **109**, 1628-1634 (2010).

12. Reidy PT, *et al.* Aging impairs mouse skeletal muscle macrophage polarization and muscle-specific abundance during recovery from disuse. *Am J Physiol Endocrinol Metab* **317**, E85-e98 (2019).

13. Bodine SC, *et al.* Identification of ubiquitin ligases required for skeletal muscle atrophy. *Science* **294**, 1704-1708 (2001).

14. Dirks ML, Wall BT, Snijders T, Ottenbros CL, Verdijk LB, van Loon LJ. Neuromuscular electrical stimulation prevents muscle disuse atrophy during leg immobilization in humans. *Acta Physiol (Oxf)* **210**, 628-641 (2014).

15. Kilroe SP, Fulford J, Jackman SR, LJC VANL, Wall BT. Temporal Muscle-specific Disuse Atrophy during One Week of Leg Immobilization. *Med Sci Sports Exerc* **52**, 944-954 (2020).

16. Jameson TSO, *et al.* Muscle damaging eccentric exercise attenuates disuse-induced declines in daily myofibrillar protein synthesis and transiently prevents muscle atrophy in healthy men. *Am J Physiol Endocrinol Metab* **321**, E674-e688 (2021).

17. Wall BT, Dirks ML, Snijders T, Senden JMG, Dolmans J, van Loon LJC. Substantial skeletal muscle loss occurs during only 5 days of disuse. *Acta Physiologica* **210**, 600-611 (2014).

18. Wall BT*, et al.* Short-term muscle disuse lowers myofibrillar protein synthesis rates and induces anabolic resistance to protein ingestion. *Am J Physiol Endocrinol Metab* **310**, E137-147 (2016).

19. Franchi MV*, et al.* Previous short-term disuse dictates muscle gene expression and physiological adaptations to subsequent resistance exercise. *J Physiol* **603**, 3725-3753 (2025).

20. Brocca L*, et al.* Human skeletal muscle fibre contractile properties and proteomic profile: adaptations to 3 weeks of unilateral lower limb suspension and active recovery. *J Physiol* **593**, 5361-5385 (2015).

21. Campbell EL*, et al.* Skeletal muscle adaptations to physical inactivity and subsequent retraining in young men. *Biogerontology* **14**, 247-259 (2013).

22. Dirks ML*, et al.* One Week of Bed Rest Leads to Substantial Muscle Atrophy and Induces Whole-Body Insulin Resistance in the Absence of Skeletal Muscle Lipid Accumulation. *Diabetes* **65**, 2862-2875 (2016).

23. Bamman MM*, et al.* Impact of resistance exercise during bed rest on skeletal muscle sarcopenia and myosin isoform distribution. *J Appl Physiol (1985)* **84**, 157-163 (1998).

24. Noone J*, et al.* The impact of 60 days of -6° head down tilt bed rest on mitochondrial content, respiration and regulators of mitochondrial dynamics. *J Physiol* **603**, 3777-3794 (2025).

25. Ringholm S*, et al.* Bed rest reduces metabolic protein content and abolishes exercise-induced mRNA responses in human skeletal muscle. *Am J Physiol Endocrinol Metab* **301**, E649-658 (2011).

26. Motanova E*, et al.* Neuromuscular junction instability with inactivity: morphological and functional changes after 10 days of bed rest in older adults. *J Physiol*, (2025).

27. Edgerton VR*, et al.* Human fiber size and enzymatic properties after 5 and 11 days of spaceflight. *J Appl Physiol (1985)* **78**, 1733-1739 (1995).

28. Murgia M*, et al.* Spaceflight on the ISS changed the skeletal muscle proteome of two astronauts. *NPJ Microgravity* **10**, 60 (2024).

29. Murgia M*, et al.* Signatures of muscle disuse in spaceflight and bed rest revealed by single muscle fiber proteomics. *PNAS Nexus* **1**, pgac086 (2022).

30. Sharples AP, Stewart CE, Seaborne RA. Does skeletal muscle have an 'epi'-memory? The role of epigenetics in nutritional programming, metabolic disease, aging and exercise. *Aging cell* **15**, 603-616 (2016).

31. Sharples AP, Turner DC. Skeletal muscle memory. *Am J Physiol Cell Physiol* **324**, C1274-c1294 (2023).

32. Bruusgaard JC, Johansen IB, Egner IM, Rana ZA, Gundersen K. Myonuclei acquired by overload exercise precede hypertrophy and are not lost on detraining. *Proc Natl Acad Sci U S A* **107**, 15111-15116 (2010).

33. Wen Y, Dungan CM, Mobley CB, Valentino T, von Walden F, Murach KA. Nucleus Type-Specific DNA Methylomics Reveals Epigenetic "Memory" of Prior Adaptation in Skeletal Muscle. *Function (Oxf)* **2**, zqab038 (2021).

34. Weidenhamer CJ, Huang YH, Natua S, Kalsotra A, Hernández-Saavedra D. Muscle memory of exercise optimizes mitochondrial metabolism to support skeletal muscle growth. *Am J Physiol Cell Physiol* **329**, C1239-c1255 (2025).

35. Egner IM, Bruusgaard JC, Eftestøl E, Gundersen K. A cellular memory mechanism aids overload hypertrophy in muscle long after an episodic exposure to anabolic steroids. *J Physiol* **591**, 6221-6230 (2013).

36. Seaborne RA, *et al.* Human Skeletal Muscle Possesses an Epigenetic Memory of Hypertrophy. *Scientific Reports (Nature)* **8**, 1898 (2018).

37. Turner DC, Seaborne RA, Sharples AP. Comparative Transcriptome and Methylome Analysis in Human Skeletal Muscle Anabolism, Hypertrophy and Epigenetic Memory. *Scientific reports* **9**, 4251 (2019).

38. Hulmi JJ, *et al.* Human skeletal muscle possesses both reversible proteomic signatures and a retained proteomic memory after repeated resistance training. *J Physiol*, (2025).

39. Blocquiaux S, *et al.* Recurrent training rejuvenates and enhances transcriptome and methylome responses in young and older human muscle. *JCSM Rapid Communications* **5**, 10-32 (2022).

40. Pilotto AM, *et al.* Human skeletal muscle possesses an epigenetic memory of high-intensity interval training. *Am J Physiol Cell Physiol* **328**, C258-c272 (2025).

41. Sexton CL, *et al.* Skeletal Muscle DNA Methylation and mRNA Responses to a Bout of Higher versus Lower Load Resistance Exercise in Previously Trained Men. *Cells* **12**, (2023).

42. Rowlands DS, *et al.* Multi-omic integrated networks connect DNA methylation and miRNA with skeletal muscle plasticity to chronic exercise in Type 2 diabetic obesity. *Physiological Genomics* **46**, 747-765 (2014).

43. Edman S, *et al.* The 24-hour molecular landscape after exercise in humans reveals *MYC* is sufficient for muscle growth. *EMBO reports* **25**, 5810-5837 (2024).

44. Von Walden F, *et al.* The myonuclear DNA methylome in response to an acute hypertrophic stimulus. *Epigenetics* **15**, 1151-1162 (2020).

45. Chambers TL, *et al.* Methylome–proteome integration after late-life voluntary exercise training reveals regulation and target information for improved skeletal muscle health. *The Journal of Physiology* **603**, 211-237 (2025).

46. Maasar MF, *et al.* The Comparative Methylome and Transcriptome After Change of Direction Compared to Straight Line Running Exercise in Human Skeletal Muscle. *Front Physiol* **12**, 619447 (2021).

47. Jacques M, *et al.* Methylome and proteome integration in human skeletal muscle uncover group and individual responses to high-intensity interval training. *Faseb j* **37**, e23184 (2023).

48. Gorski PP, *et al.* Aerobic exercise training resets the human skeletal muscle methylome 10 years after breast cancer treatment and survival. *Faseb j* **37**, e22720 (2023).

49. Gorski PP, Turner DC, Iraki J, Morton JP, Sharples AP, Areta JL. Human skeletal muscle methylome after low-carbohydrate energy-balanced exercise. *American Journal of Physiology-Endocrinology and Metabolism* **324**, E437-E448 (2023).

50. Barrès R, *et al.* Acute exercise remodels promoter methylation in human skeletal muscle. *Cell Metab* **15**, 405-411 (2012).

51. Nitert MD, *et al.* Impact of an Exercise Intervention on DNA Methylation in Skeletal Muscle From First-Degree Relatives of Patients With Type 2 Diabetes. *Diabetes* **61**, 3322-3332 (2012).

1708 52. Geiger C, *et al.* DNA methylation of exercise-responsive genes differs between trained and
1709 untrained men. *BMC Biology* **22**, 147 (2024).
1710

1711 53. Furrer R, *et al.* Molecular control of endurance training adaptation in male mouse skeletal
1712 muscle. *Nature Metabolism* **5**, 2020-2035 (2023).
1713

1714 54. Lindholm ME, *et al.* An integrative analysis reveals coordinated reprogramming of the
1715 epigenome and the transcriptome in human skeletal muscle after training. *Epigenetics* **9**,
1716 1557-1569 (2014).
1717

1718 55. Jacques M, *et al.* Molecular landscape of sex- and modality-specific exercise adaptation in
1719 human skeletal muscle through large-scale multi-omics integration. *Cell Rep* **44**, 115750
1720 (2025).
1721

1722 56. Egan B, Sharples AP. Molecular responses to acute exercise and their relevance for
1723 adaptations in skeletal muscle to exercise training. *Physiol Rev* **103**, 2057-2170 (2023).
1724

1725 57. Voisin S, *et al.* Exercise is associated with younger methylome and transcriptome profiles in
1726 human skeletal muscle. *Aging cell*, e13859 (2023).
1727

1728 58. Sailani MR, *et al.* Lifelong physical activity is associated with promoter hypomethylation of
1729 genes involved in metabolism, myogenesis, contractile properties and oxidative stress
1730 resistance in aged human skeletal muscle. *Sci Rep* **9**, 3272 (2019).
1731

1732 59. Turner DC, *et al.* DNA methylation across the genome in aged human skeletal muscle tissue
1733 and muscle-derived cells: The role of HOX genes and physical activity. *Scientific reports* **10**,
1734 15360 (2020).
1735

1736 60. Voisin S, *et al.* Meta-analysis of genome-wide DNA methylation and integrative omics of age
1737 in human skeletal muscle. *J Cachexia Sarcopenia Muscle* **12**, 1064-1078 (2021).
1738

1739 61. Zykovich A, *et al.* Genome-wide DNA methylation changes with age in disease-free human
1740 skeletal muscle. *Aging cell* **13**, 360-366 (2014).
1741

1742 62. Barrès R, *et al.* Non-CpG methylation of the PGC-1alpha promoter through DNMT3B controls
1743 mitochondrial density. *Cell Metab* **10**, 189-198 (2009).
1744

1745 63. Ruple BA, *et al.* Resistance training rejuvenates the mitochondrial methylome in aged human
1746 skeletal muscle. *Faseb j* **35**, e21864 (2021).
1747

1748 64. Alibegovic AC, *et al.* Insulin resistance induced by physical inactivity is associated with
1749 multiple transcriptional changes in skeletal muscle in young men. *American Journal of
1750 Physiology-Endocrinology and Metabolism* **299**, E752-E763 (2010).
1751

1752 65. Tomiga Y, *et al.* One week, but not 12 hours, of cast immobilization alters promotor DNA
1753 methylation patterns in the nNOS gene in mouse skeletal muscle. *The Journal of Physiology*
1754 **597**, 5145-5159 (2019).
1755

1756 66. Fisher AG, *et al.* Transcriptomic and epigenetic regulation of disuse atrophy and the return to
1757 activity in skeletal muscle. *Faseb j* **31**, 5268-5282 (2017).
1758

1759 67. Falick Michaeli T, *et al.* Muscle injury causes long-term changes in stem-cell DNA
1760 methylation. *Proc Natl Acad Sci U S A* **119**, e2212306119 (2022).
1761

1762 68. Lee PH, Macfarlane DJ, Lam TH, Stewart SM. Validity of the International Physical Activity
1763 Questionnaire Short Form (IPAQ-SF): a systematic review. *Int J Behav Nutr Phys Act* **8**, 115
1764 (2011).
1765

1766 69. Li Y, *et al.* Active Ankle Movements Prevent Formation of Lower-Extremity Deep Venous
1767 Thrombosis After Orthopedic Surgery. *Med Sci Monit* **22**, 3169-3176 (2016).

1768
1769 70. Sarto F, *et al.* Effects of short-term unloading and active recovery on human motor unit
1770 properties, neuromuscular junction transmission and transcriptomic profile. *J Physiol* **600**,
1771 4731-4751 (2022).
1772
1773 71. Skattebo Ø, Peci D, Clauss M, Johansen EI, Jensen J. Increased Mass-Specific Maximal Fat
1774 Oxidation Rate with Small versus Large Muscle Mass Exercise. *Med Sci Sports Exerc* **54**,
1775 974-983 (2022).
1776
1777 72. Seaborne RA, *et al.* UBR5 is a novel E3 ubiquitin ligase involved in skeletal muscle
1778 hypertrophy and recovery from atrophy. *J Physiol* **597**, 3727-3749 (2019).
1779
1780 73. Elhassan YS, *et al.* OR03-06 NAD+ Availability Modulates 11β-HSD1-Mediated
1781 Glucocorticoid Regeneration in Mouse Skeletal Muscle. *J Endocr Soc* **4**, (2020).
1782
1783 74. Alsharidah M, Lazarus NR, George TE, Agley CC, Velloso CP, Harridge SD. Primary human
1784 muscle precursor cells obtained from young and old donors produce similar proliferative,
1785 differentiation and senescent profiles in culture. *Aging cell* **12**, 333-344 (2013).
1786
1787 75. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat*
1788 *Protoc* **3**, 1101-1108 (2008).
1789
1790 76. Dobin A, *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
1791
1792 77. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
1793 seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
1794
1795 78. Tang D, *et al.* SRplot: A free online platform for data visualization and graphing. *PLoS One*
1796 **18**, e0294236 (2023).
1797
1798 79. Hulsen T. DeepVenn - a web application for the creation of area-proportional Venn diagrams
1799 using the deep learning framework Tensorflow.js. *ArXiv* **abs/2210.04597**, (2022).
1800
1801 80. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists
1802 using DAVID bioinformatics resources. *Nature Protocols* **4**, 44-57 (2009).
1803
1804 81. Smith T, Heger A, Sudbery I. UMI-tools: modeling sequencing errors in Unique Molecular
1805 Identifiers to improve quantification accuracy. *Genome Res* **27**, 491-499 (2017).
1806
1807 82. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
1808 applications. *Bioinformatics* **27**, 1571-1572 (2011).
1809
1810 83. Akalin A, *et al.* methylKit: a comprehensive R package for the analysis of genome-wide DNA
1811 methylation profiles. *Genome Biol* **13**, R87 (2012).
1812
1813 84. Rooney JP, *et al.* PCR based determination of mitochondrial DNA copy number in multiple
1814 species. *Methods Mol Biol* **1241**, 23-38 (2015).
1815
1816 85. Lapatto HAK, *et al.* Nicotinamide riboside improves muscle mitochondrial biogenesis, satellite
1817 cell differentiation, and gut microbiota in a twin study. *Sci Adv* **9**, eadd5163 (2023).
1818
1819 86. Kemper MF, Stirone C, Krause DN, Duckles SP, Procaccio V. Genomic and non-genomic
1820 regulation of PGC1 isoforms by estrogen to increase cerebral vascular mitochondrial
1821 biogenesis and reactive oxygen species protection. *Eur J Pharmacol* **723**, 322-329 (2014).
1822
1823 87. Arduini A, *et al.* Mitochondrial biogenesis fails in secondary biliary cirrhosis in rats leading to
1824 mitochondrial DNA depletion and deletions. *Am J Physiol Gastrointest Liver Physiol* **301**,
1825 G119-127 (2011).
1826

88. Stevanović-Silva J, Beleza J, Coxito P, Oliveira PJ, Ascensão A, Magalhães J. Gestational Exercise Antagonises the Impact of Maternal High-Fat High-Sucrose Diet on Liver Mitochondrial Alterations and Quality Control Signalling in Male Offspring. *Int J Environ Res Public Health* **20**, (2023).

89. Dall M, *et al.* Hepatic NAD+ levels and NAMPT abundance are unaffected during prolonged high-fat diet consumption in C57BL/6JBomTac mice. *Molecular and Cellular Endocrinology* **473**, 245-256 (2018).

90. Van der Stede T, *et al.* Deconvoluting fiber type proportions from human skeletal muscle transcriptomics and proteomics data using FibeRtypeR. *bioRxiv*, 2025.2009.2017.676778 (2025).

91. Damgaard MV, Treebak JT. What is really known about the effects of nicotinamide riboside supplementation in humans. *Sci Adv* **9**, eadi4862 (2023).

92. Bodine SC, Sinha I, Sweeney HL. Mechanisms of Skeletal Muscle Atrophy and Molecular Circuitry of Stem Cell Fate in Skeletal Muscle Regeneration and Aging. *The Journals of Gerontology: Series A* **78**, 14-18 (2023).

93. Zuccarelli L, *et al.* Mitochondrial Sensitivity to Submaximal [ADP] Following Bed Rest: A Novel Two-Phase Approach Associated With Fibre Types. *J Cachexia Sarcopenia Muscle* **16**, e13775 (2025).

94. Sarto F, *et al.* Twenty-one days of bed rest alter motor unit properties and neuromuscular junction transmission in young adults. *J Appl Physiol (1985)* **138**, 1411-1424 (2025).

95. Damgaard MV, *et al.* Intravenous nicotinamide riboside elevates mouse skeletal muscle NAD(+) without impacting respiratory capacity or insulin sensitivity. *iScience* **25**, 103863 (2022).

96. Stearns-Reider KM, *et al.* Aging of the skeletal muscle extracellular matrix drives a stem cell fibrogenic conversion. *Aging cell* **16**, 518-528 (2017).

97. Pavan P, *et al.* Alterations of Extracellular Matrix Mechanical Properties Contribute to Age-Related Functional Impairment of Human Skeletal Muscles. *International Journal of Molecular Sciences* **21**, 3992 (2020).

98. Pillon NJ, *et al.* Transcriptomic profiling of skeletal muscle adaptations to exercise and inactivity. *Nature Communications* **11**, 470 (2020).

99. White JR, Confides AL, Moore-Reed S, Hoch JM, Dupont-Versteegden EE. Regrowth after skeletal muscle atrophy is impaired in aged rats, despite similar responses in signaling pathways. *Exp Gerontol* **64**, 17-32 (2015).

100. Pearen MA, *et al.* The nuclear receptor, Nor-1, markedly increases type II oxidative muscle fibers and resistance to fatigue. *Mol Endocrinol* **26**, 372-384 (2012).

101. Goode JM, *et al.* The Nuclear Receptor, Nor-1, Induces the Physiological Responses Associated With Exercise. *Mol Endocrinol* **30**, 660-676 (2016).

102. Smith JAB, *et al.* Inactivity-induced NR4A3 downregulation in human skeletal muscle affects glucose metabolism and translation: Insights from in vitro analysis. *Mol Metab* **99**, 102200 (2025).

103. Brocca L, Rossi M, Canepari M, Bottinelli R, Pellegrino MA. Exercise Preconditioning Blunts Early Atrogenes Expression and Atrophy in Gastrocnemius Muscle of Hindlimb Unloaded Mice. *International Journal of Molecular Sciences* **23**, 148 (2022).

1886   104.   Michel JM*, et al.* Effects of leg immobilization and recovery resistance training on skeletal
1887   muscle-molecular markers in previously resistance-trained versus untrained adults. *Journal of*
1888   *Applied Physiology* **138**, 450-467 (2025).
1889
1890