Title: Calcium: Modulator of Post-transcriptional and post-translational process in mESCs

Shikha Sharma, Dasaradhi Palakodeti, Tina Mukherjee

Short title: Calcium and mESCs pluripotency

Author List

Dr Shikha Sharma: First author, lead contact and corresponding author

Affiliation: Institute for Stem Cell Science and Regenerative Medicine (inStem), Bangalore,
India, 560065

E-mail: shikha.ss19254@gmail.com

Orcid ID: https://orcid.org/ 0000-0002-0364-3162

Dr Dasaradhi Palakodeti

Affiliation: Institute for Stem Cell Science and Regenerative Medicine (inStem), Bangalore,
India, 560065

E-mail: dasaradhip@instem.res.in

Dr Tina Mukherjee

Affiliation: Institute for Stem Cell Science and Regenerative Medicine (inStem), Bangalore,
India, 560065

E-mail: tinam@instem.res.in

Abstract

Calcium ion (Ca^{2+}) is a ubiquitous intracellular and extracellular messenger that regulates several cellular activities. Previous findings have reported the presence of active Ca^{2+} receptors in mouse embryonic stem cells (mESCs) but the fundamental requirement of Ca^{2+} signalling remains unclear. We have noted the presence of high Ca^{2+} in undifferentiated mESCs and showed that its depletion exerts G2/M cell cycle arrest, spontaneous differentiation of mESCs towards mesoderm lineage and mitochondrial biogenesis. Further, our data demonstrates that Ca^{2+} regulates the homeostasis and stability of Oct4 and Nanog at the post-translational level through pCamkIIα dependent mechanism independent of polyubiquitin system, JAK-STAT3 pathway, transcriptional and translational control. Our data also signifies the role of Ca^{2+} at the post-transcriptional level in regulating p-bodies and stress granule markers (Dcp1a, XRN1, Tudor and EDC4), splicing-dependent and 3'UTR-dependent NMD activity. Together, this study identifies the broad role of Ca^{2+} in modulating key processes in mESCs.

Keywords: Calcium; p-bodies; cell-cycle arrest; pluripotency; mitochondrial biogenesis

Highlights

- Ca^{2+} is present at elevated levels in mESCs and is important for pluripotency

- Ca^{2+} depletion accelerates G2/M cell cycle arrest and mesoderm differentiation

- Ca^{2+} regulates pluripotency and p-bodies markers at the post-translational level

- Ca^{2+} regulates Oct4, Nanog, dcp1a through pCaMKIIα dependent mechanism

Introduction

Embryonic stem cells (ESCs) are derived from the inner cell mass (ICM) of embryos defined by eminent features such as pluripotency, self-renewal, and limitless proliferation (Ye et al., 2024). Several studies have shown that the ESCs' core pluripotency network is regulated by diverse molecular mechanisms, extracellular signalling pathways, post-transcriptional regulation, epigenetic mechanisms, and chromatin remodelling (Yeo and Ng, 2013, Kobayashi and Kikyo, 2015). The ability of ESCs to differentiate into all three lineages of an organism makes them an ideal candidate for basic research on pluripotency, genomic engineering, cell identity transition, disease modelling, cell therapy, and regenerative medicine (Kobayashi and Kikyo, 2015, Czechanski et al., 2014). Further understanding the regulation of ESCs' pluripotency network will provide more insight into the nature of pluripotent stem cells, cell fate decision, and lineage specification.

Ca^{2+} is a fundamental regulator of several processes such as cell cycle, cell proliferation, neuronal plasticity, gene regulation, and apoptosis (Clapham, 2007, Berridge et al., 2003). Despite the broad role of Ca^{2+} signalling in different cell types, there are only a few studies that have reported the presence of Ca^{2+} signalling in pluripotent stem cells. In 2004, Yanagida et al. documented the activation of various Ca^{2+} receptors upon stimulation in mESCs, indicating the presence of calcium-regulating pathways (Yanagida et al., 2004). Todorova et al. (2009) observed the activation of Ca^{2+} signalling by lysophosphatidic acid (LPA) in mESCs. In 2010, Schwirtlich et al. reported that GABA-mediated Ca^{2+} signalling is important for the proliferation and differentiation of mESCs. Recently, it was demonstrated that high intracellular calcium is paramount for regulating naïve pluripotent state and self-renewal of mESCs, suggesting the importance of calcium in controlling the pluripotency of

mESCs (Zhu and Zhang, 2019, MacDougall et al., 2019). However, the role of Ca^{2+}

signalling in the regulation of pluripotency remains unclear. In this study, we have reported

the role of Ca^{2+} signalling in regulating proliferation, cell cycle, p-bodies dynamic and core

pluripotency markers of mESCs. Our data indicates that high Ca^{2+} is important for the

maintenance of pluripotency in mESCs at the post-transcriptional and post-translational

levels.

Results

Pluripotent stem cells maintain an elevated level of cytosolic calcium

Using fluo4 staining, we observed a high level of Ca^{2+} in undifferentiated mESCs compared

to spontaneously differentiated mESCs cultured in the absence of LIF (Fig. 1A and 1D). It is

not yet clear why undifferentiated mESCs maintain a high level of Ca^{2+} within their

cytoplasm. Therefore, we became interested in understanding the role of Ca^{2+} in mESCs. We

manipulated the level of Ca^{2+} in the presence of LIF using BAPTA-AM (Bapta) and

thapsigargin. Bapta is a highly selective cell-permeable Ca^{2+} chelator (Collatz et al., 1997),

and thapsigargin is a SERCA pump inhibitor known to increase intracellular Ca^{2+} in various

cell types including ESCs (Lytton et al., 1991, Huang et al., 2017). We observed using Fluo-4

staining that supplementation of Bapta 2uM (B2uM) for 72hours was able to reduce the level

of Ca^{2+} in mESCs, comparable to without LIF condition (C-L), while thapsigargin 1nM

(T1nM) treatment increased the calcium level (Fig. 1A and 1D). We also noted that Ca^{2+} was

downregulated in the C-L condition, and supplementation of T1nM in C-L condition was able

to increase the Ca^{2+} level slightly but was unable to bring it back to the control condition

(C+L) (Fig. 1A and 1D). We also detected heterogeneity in the level of Ca^{2+} in cells within

mESCs colonies (Fig. 1A). In addition, we noted that B1uM also reduced the Ca^{2+} level in

the different organelles, such as endoplasmic reticulum (ER) (Fig. 1B and 1E) and lysosomes

(Fig. 1C) of mESCs. Conversely, T1nM increased the overall Ca^{2+} level in the lysosomes

(Fig. 1C) but not in the ER (Fig. 1B and 1E).

Further, we have observed a decrease in the proliferation of mESCs with the increase in the

concentration of bapta (B1uM, B2uM, and B5uM) in 72 hrs (Fig. 1F). We noted that mESCs

colonies exhibited fibroblast morphology upon differentiation and observed increased

differentiation with increasing concentrations of bapta (B1uM and B2uM) (Supplementary

file S1 (S1)_Supplementary Fig. S2C and S2D). Indeed, we also observed that differentiation

was less in bapta-treated samples compared to C-L, but they exhibited fibroblast morphology

similar to C-L condition (S1_ Supplementary Fig. S2C and S2D). We further replaced B2uM

media after 72 hours with control ESC media (B2R) for 48 hours and observed an increase in

the proliferation of mESCs, indicating bapta treatment is reversible (Fig. 1G). However, we

noted that mESCs that underwent complete morphological change (fibroblast appearance)

were unable to proliferate further and had retained their phenotype. Several studies have

revealed that cMyc is important for maintaining the unlimited proliferation of mESCs, and its

depletion exerts dormancy and proliferation arrest (Scognamiglio et al., 2016; Fagnocchi et

al., 2016; Ruan et al., 2017). We next checked the level of cMyc within these mESCs and

observed that cMyc expression goes down upon B2uM treatment in 72 hours, independent of

transcriptional control (Fig. 1H and 1K), which indicated that Ca^{2+} might be regulating

proliferation in mESCs by controlling cMyc expression. We further observed an increase in

the expression of cMyc in B2R compared to B2uM in 5 days (Fig. 1I and 1J). Besides this,

we analysed the cell cycle profile of C+L, B2uM, and B2R mESCs in 72 hrs (Fig. 1L) and 5

days (Fig. M), and observed that B2uM treatment resulted in the cell cycle arrest at G2/M,

which was reversible with B2R treatment (Fig. M). Earlier reports in mesenchymal stem cells

(MSCs) and epidermal stem cells suggested that high calcium is crucial for stem cell

proliferation and cell cycle progression; and G2 phase of cell cycle is important for the

determination of Ca^{2+} signalling pattern (Moore et al., 2024, Vallet et al., 2025) . Further, we

also noted that B2uM and C-L resulted in the fluctuation in the expression of calcium

receptors of the plasma membrane, ER and mitochondria including SERCA PMCA, IP3,

STIM, ORAI, Ryanodine, and MCU (S1_Supplementary Fig. S1C, D, E, F, G, H), which

signifies that bapta treatment might be affecting the level of Ca^{2+} within different organelles.

Calcium homeostasis is critical for pluripotency

To explore the role of Ca^{2+} in the maintenance of pluripotency in mESCs, we checked the

expression of pluripotency markers upon bapta and thapsigargin treatment. Western blot and

immunofluorescence (IF) data confirmed the increase/decrease in the protein level of core

pluripotency markers including Oct4 and Nanog upon low/high Ca2+ condition using bapta

(B1uM and B2uM) and thapsigargin 250 pM, 500pM and 1nM (T250pM, T500pM, T1nM)

in the presence of LIF, suggesting that Ca^{2+} has role in the regulation of pluripotency markers

(Fig. 2 A, B, E, F and M). Moreover, we noted the heterogeneity in the expression of Oct4

and Nanog in cells within mESCs colonies (Fig. 2A and 2B), which also corroborates with

the heterogeneous level of Ca^{2+} in cells within mESCs (Fig. 1A). Next, we have not observed

any significant difference in the expression of Oct4, Nanog, and cMyc in the absence of LIF

and the presence of thapsigargin (Fig. 2 A, B, G, H, and M). CHIR 99021 (CH) and

PD0325901 (PD) are inhibitors of glycogen synthase kinase 3 beta (GSK3β) and

MAPK/ERK signalling pathways, also known as 2i inhibitors implicated for the maintenance

of the ground state of mESCs (Sim et al., 2017). Furthermore, we subjected mESCs to 2i

inhibitor treatment in the presence of LIF and B1uM and observed a decrease in the

expression of Oct4 and Nanog (S1_Supplementary Fig. S2A), which signifies that Ca^{2+}

regulates the expression of pluripotency markers under naive and ground state of mESCs. We

also discerned that some of the mESCs colonies exhibited fibroblast morphology upon B1uM

treatment in the presence of a 2i inhibitor (S1_Supplementary Fig. 2B).

Phosphorylated CaMKIIα (pCaMKIIα) is an eminent downstream target of Ca^{2+} signalling

(Clapham 2007). When intracellular calcium level increases, calcium interact with

calmodulin and activates CaMKIIα by phosphorylating it at T286, which induces

autonomous activation of CaMKIIα making it sustain pCaMKIIα level for some time, when

calcium level drops (Clapham 2007, Chi et al., 2016). We checked the expression of

pCaMKIIα using the pCaMKIIα (Thr 286) monoclonal antibody. Our data showed Ca^{2+}-

dependent simultaneous upregulation/downregulation of pCaMKIIα and Oct4 in mESCs

upon B2uM and T1nM treatment for 72 hours (Fig. 2B, 2C, and 2M). We further noted that

pCaMKIIα is present mainly in the nucleus of mESCs and observed the co-localization of

pCaMKIIα with Oct4 (Fig. 2E, yellow), indicating that Ca^{2+} might have a role in the

regulation of phosphorylation of pluripotency markers through a pCaMKIIα-dependent

mechanism. Previous reports suggest that LIF plays a major role in maintaining the

pluripotency of mESCs mainly through the JAK-STAT3 pathway (Tang and Tian, 2013,

Ohtsuka et al., 2015). We asked if Ca^{2+} regulates the LIF signalling by controlling the JAK-

STAT pathway in mESCs. We observed that the protein level of pSTAT3 was unchanged

upon B1uM, B2uM, T250pM, T500pM, and T1nM treatment in the presence and absence of

LIF, suggesting that Ca^{2+} might not be playing role through JAK-STAT3 pathway for the

maintenance of pluripotency of mESCs (Fig. 2 E, F, and H). Further, we checked the

transcript level of Oct4, Nanog, Sox2 and Klf4, and early differentiation markers of

mesoderm (Mixl1, Brachyury), endoderm (Sox17, and MeoX1), and ectoderm (Sox1, and

Pax6) upon Ca^{2+} manipulation using B2uM and T1nM and observed no change in the

transcript level of these markers in 48hrs (S1_Sipplementary Fig. 1A and 1B) and 4 days

(Fig. 2I and 2J). Therefore, our data indicate that Ca^{2+} might be playing a role in the

regulation of pluripotency and differentiation markers at the post-transcriptional,

translational, and post-translational levels.

Besides this, we observed an increase in the cytoplasm-to-nucleus ratio of mESCs by

decreasing the level of Ca^{2+} using B2uM and C-L conditions, and a decrease in the

cytoplasm-to-nucleus ratio upon T1nM treatment by phalloidin staining, shown by the arrow

(Fig. 2K and 2L). Park et al. have also demonstrated that high cytosolic Ca^{2+} results in a

decreased cytoplasm/nuclear ratio in drosophila C4da neurons (Park et al., 2020), which also

corroborates with our findings and indicates the possible role of Ca^{2+} in cytoskeleton

remodelling for cell shape transition. Cadherin-adhesion complex is important for bridging

cell-to-cell contact and drives various morphogenetic changes (Mege and Ishiyama, 2017).

Therefore, we checked the expression of calcium-dependent adhesion protein E-cadherin and

found its expression downregulated in mESCs upon B2uM treatment and increased upon

T1nM condition, indicating the involvement of Ca^{2+}-dependent cadherin family proteins for

driving morphological changes in ESCs upon Ca^{2+} manipulation (Fig. 2N).

Calcium is important for the regulation of ribonucleoprotein complexes and NMD

activity

To understand whether Ca^{2+} has any role at the post-transcriptional level. We examined the

expression of p-bodies and stress granule markers upon Ca^{2+} manipulation in mESCs. Our

data showed that dcp1a, xrn1, Tudor and b-actin are present abundantly in both cytoplasm

and nucleus of undifferentiated mESCs (Fig. 3A and 3B, S1_Supplementary Fig. S3A, B, C,

and D) and observed a decrease/increase in the expression of these mRNP granules upon

B2uM and T1nM treatment (Fig. 3A, B, D, S1_Supplementary Fig S3 A, B, C, D, E). We

also noted the variation in the size of these granules, as some were large in size and others

small (Fig. 3B). The variations in size could be due to several reasons, such as a) these

markers may be a part of several complexes having different compositions, either present as a

monomer or dimer b) they may be at different stages of maturation for post-transcriptional

processes to carry out their functions. Further, we have observed the co-localization of dcp1a

with b-actin, xrn1, tudor, and pCaMKIIα, indicating that Ca^{2+} might be regulating p-body

markers through a pCaMKIIα-dependent mechanism (Fig. 3B). However, we were unable to

check the co-localization of other markers with pCaMKIIα as both the antibodies were raised

in the same species. Our data also shows that Ca^{2+} regulates p-bodies and stress granule

markers independent of transcriptional control (Fig. 3C). Next, we have quantified splicing-

dependent nonsense-mediated mRNA decay (NMD) activity and long 3' UTR-dependent

NMD activity in B2uM and C-L conditions using two fluorescent reporter proteins, green

TagGFP2 and far-red Katushka, as described by Pereverzev et al. 2015. We have observed

increased splicing-dependent NMD activity (Fig. 3E and 3F) and decreased 3' UTR NMD

activity (Fig. 3G and 3H) in mESCs treated with B2uM and C-L conditions, suggesting that

Ca^{2+} exerts the opposite effect on splicing-dependent and 3'UTR-dependent NMD pathways

in mESCs. As we have not observed any change in the transcript level of pluripotency and

differentiation markers. This also raises a possibility that p-bodies, stress granule markers,

and NMD in mESCs might be playing a major role in the regulation of markers other than

pluripotency and differentiation markers, and further studies are needed to identify those

markers.

Calcium doesn't control the expression of pluripotency and differentiation markers at

the post-transcriptional and translational level

We checked the global translation rate through polysome profiling and observed no change in

the global translational rate upon Ca^{2+} manipulation with B2uM in 72 hours (Fig. 4A).

However, we observed that the expression of early translation initiation marker elf1a was

downregulated in B2uM-treated mESCs in 72 hours (Fig. 4B). which indicates a possibility

of differential expression of translation markers upon Ca^{2+} manipulation. We have observed

no change in the transcription of pluripotency and differentiation markers in control and

B2uM-treated mESCs (S1_Sipplementary Fig. 1A and 1B, Fig. 2I and 2J). We sought to find

whether Ca^{2+} suppresses the translation of pluripotency/differentiation markers by holding

them in p-bodies/stress granules. To address this, we performed polysome profiling of

mESCs treated with B2uM for 72 hours and collected cytoplasmic fraction (composed of

mRNP granules), 80s ribosomal subunit (80s), and polysome fraction, and checked the

expression of Oct4, Nanog, Sox2, and Klf4 and early differentiation markers of mesoderm,

and ectoderm, such as Mixl1, Brachyury, Pax6 exhibiting high CT value. We were unable to

detect any significant difference in the transcript level of both pluripotency and

differentiation markers in the cytoplasmic fraction of mRNP granules, the 80S and polysome

fraction of control and B2uM-treated samples (Fig. 4D and 4E), indicating that pluripotency

and differentiation markers are not held by p-bodies and might also not be regulated at the

translational level. As we observed no change in the transcript level of core pluripotency and

early differentiation markers, To affirm this further, we blocked the transcription and

translation of control, B1uM, T1nM and C-L treated mESCs after 72hrs with combined

actinomycin D (ACD) cycloheximide (CHX) treatment for 2hr and checked the expression of

pluripotency and mesodermal differentiation markers and observed no significant change in

their expression, which illustrate that Ca^{2+} doesn't regulate core pluripotency and

differentiation markers at transcriptional and post-transcriptional level (Fig. 4F and 4G).

It is known that undifferentiated ESCs either do not express or express low levels of

differentiation markers (Ghimire et al., 2018). We noted that the transcript level of early

differentiation markers such as MixL1 and Brachyury was expressed at CT values of 28±1

and 28±1, respectively, while MeoX1 expressed a CT value of 28±1, and SOX17 was mostly

undetermined or expressed at a CT value of 34±1, Pax6 and SOX1 were present at CT values

of 26±1 and 26±1 respectively in undifferentiated mESCs. These CT values signify that

transcripts of differentiation markers are present at a low level in undifferentiated mESCs. In

addition, we detected transcript of differentiation markers (MixL1, Brachyury, and Pax6) in a

polysome fraction of control and B2uM treated mESCs (Fig. 4D and 4E), which indicates

that differentiation markers are undergoing translation and might be expressed in

undifferentiated mESCs or getting degraded. Since these mESCs showed a tendency to

spontaneous differentiation towards mesoderm lineage (fibroblast morphology). We have

checked the expression of only brachyury protein in control, B2uM and C-L condition (Fig.

4H). We have observed some background expression of brachyury in undifferentiated mESCs

and its increased expression in B2uM and C-L conditions (Fig. 4H), which indicates the role

of calcium in the degradation of brachyury protein in C+L condition. Next, we also

performed mitotracker-red staining and noted the increase in the size and number of

mitochondria upon calcium reduction with B2uM and C-L condition in 48hrs (Fig. 4I, and

4J), we observed globular mitochondria in control mESCs and a network of branched

mitochondria in B2uM and C-L in 48hrs (Fig. 4I), indicating mitochondria undergo a

morphological change from globular to tubular form upon Ca^{2+} reduction.

Calcium is important for the stability of pluripotency markers, p-body and stress

granule markers, and degradation of differentiation markers

Further, we asked whether Ca^{2+} plays a role in the degradation of proliferation,

differentiation, core pluripotency markers, polyubiquitin marker (pUBQ), p-body, and stress

granule markers at post-translational level. Our data demonstrate an increase in the

expression of cMyc, dcp1a, tudor, Oct4, and pUBQ in both control and B2uM-treated mESCs

upon MG132 treatment (Fig. 5 A, B, C, D, E, F, G), which suggests that Ca^{2+} level might be

important for the maintenance of homeostasis and stability of these proteins. Further, we also

observed increased expression of brachyury in the control condition after MG2132 treatment,

which also corroborates with the increased expression of brachyury upon B2uM treatment but

MG132 was unable to increase the brachyury protein expression in B2uM treated mESCs but

rather it led to its decreased expression (Fig. 5A and 5D). Earlier findings have suggested that

proteasomal inhibition can lead to the degradation of proteins through the activation of other

degradation pathways such as autophagy (Wang et al., 2019). Furthermore, we noted that

MG132 treatment was able to rescue only partial protein levels of dcp1a and tudor in B2uM-

treated mESCs compared to control mESCs (Fig. 5B and 5F), which also implies the role of

other signalling pathways in the regulation of p-body and stress granule markers upon Ca^{2+}

reduction. Interestingly, we also noted the decrease in the expression of pUBQ upon B2uM

treatment and its upregulation with MG132 treatment (Fig. 5C and 5D), which suggests that

Ca^{2+} might be regulating the stability of pluripotency markers independent of PUBQ

proteasomal system upon Ca^{2+} reduction. Next, we observed a very low expression of LC3 in

the undifferentiated mESCs, which further decreased upon B2uM and C-L condition,

suggesting autophagy might not be playing a role in the degradation of pluripotency and

differentiation markers upon Ca^{2+} reduction (Fig. 5H). Our study also indicates that the

pUBQ proteasomal system might be important for the degradation of differentiation markers

in undifferentiated mESCs, and its low expression upon Ca^{2+} reduction is important for the

stability of brachyury protein and activation of other signalling pathways apart from

autophagy to determine lineage specificity. We have further checked the stability of Oct4 and

Nanog using a CHX chase assay as described by Kao et al., 2015. We chased the expression

of Oct4 and Nanog for 1hr 20 min with a 20 min time interval and observed that their

degradation was faster in B2uM-treated mESCs, indicating that Ca^{2+} is important for the

maintenance of the stability of core pluripotency markers (Fig. 5I). Furthermore, we observed

an increase in the level of Oct4 and Nanog proteins in B2uM-treated mESCs with MG132

treatment (Fig. 5J), suggesting that Ca^{2+} maintains the stability of pluripotency markers at the

post-translational level.

304   Calcium maintains the stability of OCT4, NANOG, and DCP1a by a pCaMKII alpha-

305   dependent mechanism

In order to affirm the role of calcium signalling in the regulation of pluripotency and p-body

markers, we performed the knockdown the pCaMKIIα, the main downstream target of

calcium signalling to estimate the decreased calcium level effect and serca-R2 knockdown to

observe increase calcium effect on calcium signalling. We used the electroporation method to

deliver the CaMKIIα shRNA to silence the pCaMKIIα protein expression. The CaMKIIα

shRNA targets and silence the CaMKIIα mRNA and prevents the production of

unphosphorylated and active pCaMKIIα protein (Moore et al., 2010, Hirst et al., 2022). We

noted that 80-90% of mESCs expressed CaMKIIα shRNA as shown by their GFP expression

(Fig. 6A). Moreover, we observed that CaMKIIα shRNA treated mESCs exhibited fibroblast-

like morphology (Fig. 6B and 6C), which was similar to the phenotype shown by B2uM

treated mESCs (Fig. 2K). However, we were unable to detect any significant decrease in the

level of CaMKIIα mRNA but observed a decrease in the level of pCaMKIIα

protein (pCaMKIIα KD) upon CaMKIIα shRNA treatment (Fig. 6D and 6I), which suggests

the possibility that CaMKIIα shRNA might have exerted an indirect mechanism in regulating

pCaMKIIα protein expression rather than canonical knockdown efficiency, and this might

be through stress granules, translational control, or protein instability, or a decrease in RNA

pool was not detected by qPCR. Furthermore, we detected a significant decrease in the

expression of Oct4 and Nanog proteins upon pCaMKIIα KD by western blotting (Fig. 6F);

however, we did not observe any change in the transcript level of both pluripotency and

differentiation markers (Fig. 6M and 6N), similar to B2uM-treated mESCs (Figure 2I and 2J).

In addition, we observed that the pSTAT3 protein level was unaffected in pCaMKIIα KD

mESCs, which again signifies that Ca^{2+} might not be playing role through JAK-STAT

pathway (Fig. 6F). We further observed a decrease in the expression of p-body marker dcp1a

upon pCaMKIIα KD (Fig. 6I), which indicates that Ca^{2+} is important for the stability of

dcp1a through a pCaMKIIα-dependent mechanism. Furthermore, we also performed SERCA-

R2 receptor knockdown (SER-R2 KD) by using shRNA against its highly expressed

receptor2, which we have noted in our QPCR data. We observed a decrease in the mRNA

pool of the SERCA-R2 receptor upon SER-R2 KD (Fig. 6D). We noted that SER-R2 KD

mESCs exhibited compact colonies as compared to control mESCs (Fig. 6B and 6C), which

was similar to T1nM-treated mESCs (Fig. 2K). Furthermore, we observed an increase in the

expression of Oct4 and Nanog proteins upon SER-R2 KD (Fig. 6G). We observed no change

in the transcript level of pluripotency and differentiation marker (Fig. 6M and 6N) as well as

in the protein level of pSTAT3 in SER-R2 KD mESCs (Fig. 6G), again signifying that

calcium might not be acting through JAK-STAT pathway. Furthermore, we also observed an

increase in the expression of dcp1a and pCaMKIIα upon SER-R2 KD (Fig. 6H), which

indicates that cytosolic Ca^{2+} plays a role in the biogenesis of dcp1a through pCaMKIIα.

Identification of phosphoproteins altered upon calcium manipulation using a

phosphoproteomics approach

We explored the pooled phosphoproteomics profile of control and B2uM-treated mESCs. We

identified 53 phosphoproteins (PP) that were significantly upregulated and 35 PP

downregulated in B2uM samples (n=2 each set pooled phosphoproteomic) shown by the heat

map (Fig.7A and Supplementary file S2 (S2)_Supplementary Table 1, 2, 3, 5). Using

Proteome Discoverer Software (Thermo Fisher Scientific), we detected the phosphosites in

the identified phosphopeptides from mass spectrometry (MS) data and observed majorly

serine-threonine phosphosites along with a few tyrosine sites, which (S2_Supplementary

Table 1 and 4), which indicate that Ca^{2+} manipulation through bapta majorly influence broad

category of serine-threonine PP. Further using ShinyGO 0.80 software, we carried out the

KEGG pathway, tree plot, GO cellular component (GOC), and string function enrichment

analysis for the differentially expressed phospho-proteins (DEPP) represented in the heat map

(Fig. 7B, C, D, E, F). We observed that spliceosome and mRNA surveillance PP were highly

enriched in our dataset and carried out further KEGG pathway enrichment analysis for them

(Fig. 7G and 7H) and observed that our PP are part of these complexes labelled as red. We

further observed that DEPP exhibits 1385 protein-protein interactions (PPI) with an expected

689 PPI (p=0) using string PPI network analysis (Fig. 7I). Next, we identified DEPP related

to different pathways using string bio process enrichment analysis (SBPEA). We conducted

GO biological function (GOB), GO molecular function (GOF) and GOC analysis for DEPP

related to metabolism listed in the legend of Supplementary Fig. 4 A, and observed that they

are the positive and negative regulator of metabolism, part of mRNP complex and displayed

nuclear and organelle localization which suggests a widespread functional role of these

proteins (S1_Supplementary Fig. S4A, 4B, 4C). Further, we carried out the GOC analysis of

a few DEPP, including Hk2, Hspa8, Atp5a1, Cox4i1, Pkm, and Gapdh, identified for ATP

metabolism, and observed they have mitochondrial localization (S1_Supplementary Fig.

S4J), which signifies their role in mitochondrial biogenesis and oxidative phosphorylation.

We also noted that several PP related to translational machinery including Rps18, Rp16,

Eif3c, Abcf1, Eef1a1, Eef2, Eif5a, Rpl18a, Eif2s1, Rpl5, Rplp2, Rpl34, Eef1d, Rpl27, Eif3b,

Rpl23a, Rps8, and Larp1 were highly differentially regulated (Fig. 7A), despite global

translation rate was unaffected upon Ca^{2+} reduction and conducted GOC, GOB network and

GOF analysis for them (S1_Supplementary Fig. S4G, 4H and 4I). We observed that these

proteins are also part of the mRNP complex and have a role in macromolecule biosynthetic

pathway, which exemplify their other role apart from controlling translation rate. Next, we

further performed the GOB and GOF analysis for DEPP implicated in nucleosome and

chromatin assembly, including Smarcd1, Nasp, Hat1, Atrx, and Nap1l1 (S1_Supplementary

Fig. S4D and 4E), and observed they have chromatin and histone binding properties and has a

role in DNA replication-independent chromatin assembly. Further, we conducted GOB

network analysis of DEPP, including CSe1, smarcd1, Hk2, Pkm, Tfpt, Tcof1, champ1, nasp,

Macf1, Hat1, Bcam, Atrx, Nmd3, Sde2, Ubr4, Hsph1, Trap1, nucks, Cdk1, Mki67, Tpx2,

champ1 (S1_Supplementary Fig. 4F), and observed that they exhibit cross-talk in the

kinetochore process, chromatin condensation, nuclear division, mitotic cell cycle, and

chromatin remodelling. We have also detected some DEPP, including Cdk1, Mki67, Nucks1,

and Tpx2, crucial for mitotic cell division and mitochondrial ATP synthesis through GOB

network analysis, implicating their role in G2/M cell cycle arrest and oxidative

phosphorylation (S1_Supplementary Fig. S5A and 5B). However, further studies are needed

to confirm the role of Ca^{2+} in the regulation of above processes in mESCs.

Discussion

In recent years, pluripotent stem cells have received extensive importance in the field of

regenerative medicine to study cell development and differentiation. However, the role of

Ca^{2+} signalling in the maintenance of pluripotency is not yet explored. Previous work in

drosophila, mesenchymal stem cells and mESCs has demonstrated that stem cells undergo

differentiation upon Ca^{2+} reduction (Shim et al., 2013, Vallet et al., 2025, MacDougall et al.,

2025, Zhu and Zhang 2019, Apáti et al., 2016). Our study showed that low calcium leads to

the spontaneous differentiation of mESCs, reflecting the broadly conserved phenomenon in

stem/progenitor cells across the species. In stem cell biology, there is a persistent effort to

understand the mechanism underlying ESC pluripotency. Nevertheless, the comprehensive

repertoire of molecules involved in controlling the ESC state is not yet completely

understood. Several studies have reported the heterogeneous expression of Nanog in mESCs

but the mechanism underlying the same is still unclear (Torres-Padilla and Chambers 2014).

Previous finding in hESCs has also indicated the heterogenous expression of Ca^{2+} signalling

in hESCs (Huang et al., 2017). Our data indicates a correlation between the heterogenous

expression of Oct4 and Nanog with the heterogeneous level of Ca^{2+} and pCaMKIIα in

mESCs, which suggest that calcium might be related with the heterogenous expression of

pluripotency markers. Earlier reports have suggested that activation of calcium receptor

RyR2, calcium-sensing receptor (CaSR) and increasing calcium level promotes the

differentiation of human and mouse ESCs towards cardiac differentiation (Meng et al., 2025,

Sun et al., 2013). Interestingly, we noted that upon Ca^{2+} reduction, mESCs showed the

mESCs tendency to differentiate towards mesoderm lineage, suggesting that Ca^{2+} signalling

might be important in undifferentiated mESCs to mask the expression of mesoderm markers.

Furthermore, our data signifies the role of high Ca^{2+} in mediating the degradation of

brachyury protein, and low calcium and low pUBQ proteasomal system for its stability,

which also suggests the activation of other signalling pathways upon calcium reduction to

determine lineage specificity. Moreover, our data demonstrate that calcium might not be

controlling LIF-JAK-STAT3 pathway for the regulation of core pluripotency and

differentiation markers and has no role at the transcriptional level but controls the stability of

these fundamental regulators at the post-translational level. Our pCaMKIIα KD data indicate

that pCaMKIIα is important for the stability of Oct4 and Nanog proteins in mESCs. We have

also observed the co-localization of Oct4 and pCaMKIIα, but have not checked whether and

how pCaMKIIα phosphorylates core pluripotency markers Oct4 and Nanog and how it results

in their downregulation. It is well known that pCaMKIIα is a serine-threonine kinase and is

involved in the phosphorylation of a wide array of proteins (Clapham 2007), thus suggesting

that it might have a role in regulating the phosphorylation status of these core pluripotency

markers. Earlier studies have demonstrated that phosphorylation of Oct4 and Nanog proteins

at serine residues decreases their stability and dispenses them towards the pUBQ-mediated

proteasomal degradation pathway in mESCs (Bae et al., 2017, Kim et al., 2014). Our data

demonstrate that Ca^{2+} might be regulating the stability of Oct4 and Nanog proteins

independent of the pUBQ system. Several reports suggest that protein degradation can

happen independent of polyubiquitin tagging or without the 19S proteasome subunit by 20S

or 26S mediated proteolysis through structural disorder (Shang and Taylor 2011, Ben-Nissan

and Sharon 2014). Previous findings have advocated that Ca^{2+} and Ca2+-binding proteins

promote conformational change upon binding to target proteins and modulate their activity

and stability (Clapham, 2007, . Strynadka, and James, 1991, Hołubowicz et al., 2017)

Therefore, further structural, biochemical, and molecular studies are needed to understand the

in-depth mechanism of how Ca^{2+} regulate the activity and stability of pluripotency markers

either through phosphorylation at specific residues or conformational change and structural

disorder.

Interestingly, our data also signifies that Ca^{2+} has a role in controlling the expression of p-

bodies and stress granule markers, but has no role in controlling the expression of

pluripotency and differentiation markers at a post-transcriptional level, which indicates that

Ca^{2+} might be regulating other signalling pathways at the post-transcriptional level important

for the maintenance of pluripotency of mESCs. Our data demonstrate that 3'UTR-mediated

NMD is important for the maintenance of pluripotency, while the splicing-dependent NMD

mechanism is more active during the differentiation process, which signifies that Ca^{2+} plays a

role in the modulation of different post-transcriptional processes. Our data has identified

several RNA-binding and ribosomal PP including edc4, dcp1a, xrn1, tudor, Pa2g4, Nmd3,

Ddx21, Srsf5, Srsf6, Srsf1, EDC4, Fkbp4, Cpsf3, Fip1l1, Hnrnpab, Hnrnpa, Hnrnpa3,

Rbm39, Dhx15, Acin1, Snw1, Larp, Srrm1, Clasrp, Rps18, Rars, Eif3c, Eef1a1, Eif5a,

Rpl18a, Eif2s1, Rpl5, Rps8, Rplp2, Rpl34, Eef1d, Rpl27, Eif3b, Rpl23a implicated in mRNP

complex formation, alternative splicing, alternative polyadenylation, mRNA stability, 5'TOP

mRNAs stability, miRNA biogenesis, mRNA decapping and NMD through GOB analysis

and literature review (Martinez et al., 2013, Hochstoeger et al., 2024, Chiang et al., 2013,

Decker and Parker, 2012), which suggests a widespread role of Ca^{2+} at a post-transcriptional

level rather than global translational control. Our data have revealed decreased expression of

dcp1a, tudor, and xrn1 and increased expression of edc4 upon Ca^{2+} reduction, which signifies

differential expression of p-bodies markers upon Ca^{2+} manipulation. Moreover, our data

indicate that pCaMKIIα is also a part of p-bodies component machinery, which suggests that

Ca^{2+} might be regulating the dynamic, composition, interaction, and activity of p-bodies

markers through regulating their phosphorylation status. Chiang et al. have revealed that edc4

binding to dcp1a is independent of its phosphorylation status, while the phosphorylation

status of dcp1a is important for its strong binding to dcp2 (decapping co-factor) (Chiang et

al., 2013). Moreover, studies on yeast demonstrated that Ca^{2+} is required for p-bodies

formation and disclosed variations in the size of p-body granules (Kilchert et al., 2010, Park

et al., 2016). Park et al. also observed co-localization of calcineurin (calcium-calmodulin-

activated serine/threonine-specific phosphatase) with p-bodies and stress granules markers in

yeast (Park et al., 2016). Taken together, our data and these studies indicate that Ca^{2+} might

have a role in mediating the interaction of dcp1a with other p-bodies and stress granule

markers and the formation of mRNP granule composition through regulating phosphorylation

and dephosphorylation. Therefore, further detailed studies are needed to elucidate the role of

Ca^{2+} in the regulation of mRNP granule composition in undifferentiated and differentiated

mESCs.

In our phosphoproteomic data, we have observed the upregulation of Atp5f1a, Cox4i1, Hk2,

and the downregulation of Pkm upon Ca^{2+} reduction, which are important for oxidative

phosphorylation and mitochondrial fusion (Lis et al., 2016, Li et al., 2019, Čunátová et al.,

2021). It has already been reported that undifferentiated ESCs have globular immature

mitochondria and prefer glycolysis over oxidative phosphorylation, whereas during

differentiation, mitochondria undergo conformation change and differentiated cells favour

oxidative phosphorylation for energy production (Fu et al., 2019). Our data also suggest

globular mitochondria in undifferentiated mESCs and long tubular mitochondria, as well as

an increase in the number of mitochondria upon Ca^{2+} reduction in mESCs, which illustrates

that low calcium might be needed for mitochondrial morphological remodelling and further

studies are needed to examine the role of Ca^{2+} in mitochondrial biogenesis and metabolic

switching in mESCs.

Furthermore, we have not observed any apoptosis in mESCs upon bapta treatment, nor did

mESCs show any stress-related phenotype. Indeed, our phosphoproteomic data detected

apoptosis marker BCL-2-associated transcription factor 1, but we have not observed any

significant change in its protein profile (S2_Supplementary Table 1), which indirectly

supports that apoptosis might not be dysregulated. In addition, we have also not observed any

ER stress-related key phosphoproteins such as PERK, eIF2α, and IRE1α (Chen et al., 2023)

in our phosphoprotein dataset (S2_Supplementary Table 1). Moreover, ER stress triggers heat

shock protein response (Liu and Chang 2008). We have detected several heat shock

phosphoproteins, such as HSP-90 alpha, HSP 90 beta, Hspa8, Hspa4, Hspb1, Ahsa1, Hsph1,

and Carhsp1 in our phosphoproteomic dataset (S2_Supplementary Table 1). Among these,

Hspa8 and Hsph1 were only significantly upregulated and downregulated, which indicates

that ER stress might not be majorly altered.

Taken together, our data indicate that calcium might be having a broad role in modulating

various biological processes such as cell cycle arrest, protein stability, RNA splicing, mRNA

stability, ribosome biogenesis, mRNP granule formation, ubiquitination, metabolism,

telomere maintenance, oxidative phosphorylation, mitochondrial biogenesis, chromatin

regulation, RNA polyadenylation for the maintenance of mESCs identity.

Materials and Methods

Culturing of mESCs and calcium modulation

mESCs E14TG2a and E14 cell lines were maintained on 0.1% gelatin at 37 $^{\circ}$C in a 5% CO2

incubator in ES medium containing high glucose DMEM (GibcoTM), ESGRO recombinant

mouse LIF, 15% ES cell FBS (GibcoTM), β-Mercaptoethanol (GibcoTM), 1x NEAA

(GibcoTM), and gentamycin (GibcoTM). To induce spontaneous differentiation, mESCs

were trypsinized with 0.05% Trypsin-EDTA (GibcoTM) and cultured in ES media in the

absence of LIF. For Bapta (InvitrogenTM) and thapsigargin treatment, mESCs were cultured

in the presence of LIF with different concentrations of bapta (1μM and 2μM) and

thapsigargin (250pm, 500pm, and 1nM) for 72 hours. Bapta and thapsigargin stock solutions

were prepared using DMSO. An equal amount of DMSO was added to the control media as

compared to bapta and thapsigargin. The medium was changed every day. All experiments

were done at 72 hour time interval except few such as mitochondrial staining was done at 48

hours where mitochondrial morphological change was detected early and transcript profile

was studied at 48 hours and 4 days to ascertain transcript are unchanged upon calcium

manipulation at shorter and longer time interval and cell cycle, proliferation and cMyc

staining was studied for 5 days with 72 hour bapta treatment and 48 hour rescue condition.

Cell proliferation assay

mESCs proliferation was calculated using the trypan blue method, and cell cycle analysis was

done by PI staining. Briefly, mESCs were fixed in 70% ethanol and stored at -20 °C. Fixed

cells were centrifuged at 1000 rpm for 3-5 min and washed with PBS 2-3 times and PI

staining solution (80 µg/ml PI, 100 µg/ml RNase A and 0.1 % Triton X-100) for 7 hrs at 4 °C.

After incubation, mESCs were washed with PBS 2-3 times and cell cycle analysis was done

by using flow cytometry and BD software.

Mitotracker-red   staining,   Fluo4   staining,   lysotracker   Staining,   and

Immunofluorescence staining

Cells were grown on coverslips in ES media containing bapta (2µM) and thapsigargin (1nM).

After 72 hours, cells were incubated with media containing Fluo4 (1µM), lysotracker

(100nM), and mitotracker red (100nM) for 45 minutes, followed by 2-3 washes with media

and immediately taken for live cell imaging. In the case of Fluo 4 staining, cells were

incubated at 37 °C in a 5% CO_2 incubator for 15 minutes before live cell imaging. The

imaging was done using a confocal microscope at 40x magnification. For immunostaining,

cells were grown on coverslips and treated with bapta (2 µM) and thapsigargin (1nM) for 72

hours, followed by fixation in 4% paraformaldehyde (PFA) overnight at 4 °C. After

incubation, PFA was removed and cells were washed with PBS twice. The fixed cells were

permeabilised using 0.4% Triton X-100 for 2 hours at room temperature (RT). Then the cells

were blocked using the blocking buffer (5% BSA and 0.1% Tween-20 in PBS) for 2 hrs at

RT. After blocking, cells were incubated with primary antibodies at 4 °C overnight or 3 hours

at RT, followed by twice washing with PBS for 5 minutes each. The primary antibodies used

were OCT3/4 (Invitrogen), Nanog (Invitrogen), elf41a (Abcam), DCP1a (sigma), XRN1,

beta-actin (mRNP) (Santacruz), pCAMKIIα (Cell Signalling Technology), pSTAT3

(Abcam), cmyc (Abcam), and Brachyury. Further, secondary antibodies (Jackson

Immunoresearch Laboratories (West Grove, PA) were added and incubated for 1-2 hr at RT.

After incubation, cells were washed with PBS twice for 5 minutes and mounted on a glass

slide using mounting media containing DAPI. For IF experiments, confocal images were used

for calculating mean intensity using ImageJ software for the gene expression analysis.

Gene expression analysis

RNA was extracted using the Trizol (Invitrogen, Carlsbad, CA, USA) method. Briefly, cells

were lysed and homogenised using Trizol, and 1/5 volume of chloroform was added to the

Trizol to extract RNA by centrifuging at 12000 rpm at 4 °C for 20 mins. After centrifugation,

an aqueous layer was collected, and an equal volume of isopropanol was added, and samples

were incubated at -80 oC overnight. After incubation, samples were centrifuged to precipitate

RNA. After centrifugation, precipitated RNA was washed with 70% ethanol, dried, and

dissolved in nuclease-free water and stored at -80°C. Further, 1 μg of RNA was reverse

transcribed with Superscript II or Superscript III reverse transcriptase (Invitrogen) according

to the manufacturer's protocol. Real-time polymerase chain reaction (PCR) was performed

using AB Invitrogen with Power Sybr Green (Thermo Fisher Scientific) according to the

manufacturer's instructions. The gene expression analysis of pluripotency and differentiation

markers was calculated using the formula $2=\Delta\Delta ct$ and normalised to glyceraldehyde 3

phosphate (GAPDH).

Protein extraction and western blotting

For protein extraction, mESCs were lysed using radioimmunoprecipitation buffer (RIPA

buffer). The samples were then centrifuged at 12000 rpm for 10 minutes at 4 °C. The

supernatant was collected, and protein estimation was quantified by the Bradford method

using BSA as a standard. An equal amount of proteins (5-10 μg) was separated on a 12%

SDS-PAGE gel. Proteins were electrophoretically transferred into polyvinylidine difluoride

membranes (Bio-Rad) for western blot analysis. The blots were blocked with blocking buffer

containing 5% BSA in TBST (20mM Tris pH 7.5, 150mM NaCl, and 0.1% Tween-20)

followed by incubation with primary antibody at 4 °C overnight or 3-4 hours at RT. After

incubation, blots were washed twice with TBST, and membranes were incubated with

horseradish peroxidase (HRP)-conjugated secondary antibody for 2 hrs. Signals were

detected with ECL substrate (Amersham GE) using ImageQuant. The primary antibodies

used were OCT3/4 (Invitrogen), Nanog (Invitrogen), GAPDH (Invitrogen), cMyc (Abcam),

and pSTAT3 (Abcam), and secondary antibodies were anti-rabbit HRP and anti-goat HRP.

GAPDH was used as a loading control.

Cycloheximide chase assay, MG132 treatment for analysing protein stability, and

actinomycin D and CHX assay for posttranscriptional activity detection

mESCs were cultured in the absence and presence of B 2μM for 72hrs, followed by treatment

with 50ug/ml CHX for 1hr, and protein levels of Oct4 and Nanog were detected every 20 min

using western blot. For the MG132 experiment, mESCs were cultured in the absence and

presence of bapta (2uM) for 72 hours, followed by 10uM MG132 treatment for 1 1hr and

detected using western blot and IF. For posttranscriptional analysis of pluripotency markers,

mESCs were treated with 5ug/ml of actinomycin D (ACD) and CHX (50ug/ml) for 2 hr after

culturing the mESCs in C+L, B2uM, T1nM, C-L, C-LT1nM, and samples were extracted in

Trizol for RNA purification.

Electroporation

Electroporation was performed using an Invitrogen Neon Transfection System (100ul)

according to the manufacturer's instructions. Briefly, mESCs were trypsinized and washed

with PBS and centrifuged at 1000rpm for 3-5 min. The mESCs pellet was then re-suspended

in the resuspension buffer (Buffer R) containing the required plasmid (pCAMKIIa shRNA,

SERCA shRNA, Design 1 Endoplasmic reticulum (D1ER) calcium sensor for measuring

endoplasmic reticulum calcium as described by Waldeck-Weiermair et al., (2015), and NMD

reporter plasmids [NMD reporter plasmids were kind from Konstantin A. Lukyanov from

Sergey A. Lukyanov lab, Russian Academy of Sciences, Russia]). Further, electroporation

was performed using a parameter of 900 volts, 3 pulses, 23 width. After electroporation,

mESCs were immediately transferred into the ES media without antibiotics and incubated at

37 $^{\circ}$C in a 5% CO2 incubator. Fresh media containing antibiotics was added after 24 hours,

and GFP expression was analysed between 48 hours and 72 hours. After 48 hours, mESCs

were collected in RIPA buffer for western blot, TRIzol for RNA extraction and fixed in 4%

for immunofluorescence staining.

Polysome profiling

mESCs were cultured in the presence of B2μM for 72hour and pellet was resuspended in

polysome lysis buffer (20mM Tris pH-7.4, 200mM KCL, 5mM Mgcl2, 0.7% triton X-100, 1x

protease inhibitor, RnaseoOUT, 100ug/ml CHX, 1mM DTT). A 15-45% sucrose gradient

was prepared in gradient buffer (20mM Tris, pH 7.4, 5mM MgCl2, 100mM KCL, 1mM

DTT, 100 μg/ml CHX) using a gradient mixer. An equal amount of RNA/260OD was loaded

into the gradient and centrifuged at 39000 rpm at 4 ^{o}C for 2.5 hr. Polysome peak was

analysed using polysome profiler, and cytoplasmic mRNP granule, 80s ribosome, and

polysome fractions were collected in absolute ethanol. Samples were stored at -80 ^{o}C

overnight, and RNA isolation was performed by collecting pellets from ethanol fractions in

Trizol or by using an RNA extraction kit (Qiagen). cDNA synthesis was carried out using

Superscript II or Superscript III according to the manufacturer's instructions.

630   Phosphoproteomics

Phosphoproteomics experiment was conducted n=2 times. mESCs were collected from one

60mm dish and one 35mm dish, each dish cultured on different days and pooled to form one

set. mESCs were cultured for 72 hours in the presence of B 2μM, and proteins were extracted

using RIPA buffer containing protease and phosphatase inhibitors. Further, Proteins were

precipitated using 15% TCA followed by a 90% acetone wash. The protein pellet was

dissolved in 8% urea buffer (8% urea in 100mM ammonium bicarbonate buffer) containing 5

mM DTT, 1x phosphor stop inhibitor and incubated at RT for 35 min for complete

denaturation of proteins. Protein levels were estimated using the Bradford method. Equal

amounts of proteins (500ug and 200ug) were taken for control and bapta-treated samples and

incubated at 56 ^{o}C for 45 min with 5mM DTT concentration. Further samples were treated

with 10mM iodoacetamide and incubated at 56 ^{o}C for 45 min. After incubation, Samples

were diluted 10 times with ammonium bicarbonate buffer and incubated at 37 °C overnight

with trypsin gold (mass spec grade, Promega) in a 1:100 ratio. After incubation, peptides

were collected using 1% FA. Alternatively, during standardization of protocol, the remaining

protein pellet after first digestion was again dissolved twice in ammonium bicarbonate buffer

to re-activate trypsin gold and supplemented with 500ng trypsin gold and incubated at 37 °C

for 8-9 hrs, and overnight respectively, and peptides were collected using 1% formic acid.

Peptides were purified using the Oasis HLB column according to manufacturer instructions

(Waters). The eluted peptides were dried using a speed vacuum. Dried peptides were

resuspended in 50% acetonitrile in 0.2% formic acid, and phosphopeptide enrichment was

done using Fe-NTA resin. Peptides were incubated in Fe-NTA for 3-4 hrs and washed twice

with 50% acetonitrile containing 0.2% formic acid. Finally, phosphopeptides were eluted

using a 2.5% ammonium hydroxide buffer. An equal amount of phosphopeptides was used in

LC/MS/MS analysis using the Thermo Orbitrap Fusion system. Fe-NTA resin was prepared

by chelating Ni-NTA resin according to the method described by Sanford & Smolka, 2021

with slight modification. After mass spec analysis, raw data of phosphopeptides were

analysed by Proteome Discoverer software from Thermo Fischer Scientific with standardised

consensus and processing workflow parameters with slight modifications, including s/n ratio

5, threshold 50, unique peptides, static carbamidomethylation modification (+57.021 Da), and

phosphosites detection was done by selecting dynamic phosphorylation modification for

serine-threonine and tyrosine. Further log2fold change was calculated from abundance values

detected by the software, and p-value was calculated using the one-way Chi-square test. The

heat map was plotted for significantly differentially regulated genes using an SR plot web

server as described by Tang et al., 2023.

Statistical analysis: p-value calculated by the two-sample t-test method from the mean and

standard deviation as described by Xu et al. (2017) using MedCalc online statistical software

calculator (MedCalc Software, 2024).

Author contributions

671 TM initiated the project. TM and SS have designed the experiments and performed the

672 analysis. DP has given suggestions on p-body markers. SS performed all the experiments and

673 wrote the manuscript with input from all authors.

Acknowledgements:

676 The work was supported by the inStem core funds.

Resource availability: Further information and requests for reagents may be directed to and

679 will be fulfilled by the Shikha Sharma (shikha.ss19254@gmail.com)

Materials availability

682 This study did not generate new unique reagents.

Funding declaration

685 This work has not received any specific funding.

Declaration of interests

The authors declare no competing interests

References

1. Apáti Á, Berecz T, Sarkadi B (2016) Calcium signaling in human pluripotent stem
cells. Cell Calcium 59(2-3):117-23.

2. Bae KB, Yu DH, Lee KY, Yao K, Ryu J, Lim DY, Zykova TA, Kim MO, Bode AM,
Dong Z (2017) Serine 347 Phosphorylation by JNKs Negatively Regulates OCT4
Protein Stability in Mouse Embryonic Stem Cells. Stem Cell Reports 9:2050-2064.

3. Ben-Nissan G, Sharon M (2014) Regulating the 20S proteasome ubiquitin-
independent degradation pathway. Biomolecules 4:862-84.

4. Berridge MJ, Bootman MD, Roderick HL (2003) Calcium signalling: dynamics,
homeostasis and remodelling. Nat Rev Mol Cell Biol. 4:517-529.

5. Chen X, Shi C, He M, Xiong S, Xia X (2023) Endoplasmic reticulum stress:
molecular mechanism and therapeutic targets. Signal Transduct Target Ther 8(1):352.

6. Chi M, Evans H, Gilchrist J, Mayhew J, Hoffman A, Pearsall EA, Jankowski H,
Brzozowski JS, Skelding KA (2016) Phosphorylation of calcium/calmodulin-
stimulated protein kinase II at T286 enhances invasion and migration of human breast
cancer cells. Sci Rep 6:33132.

7. Chiang PY, Shen YF, Su YL, Kao CH, Lin NY, Hsu PH, Tsai MD, Wang SC, Chang
GD, Lee SC, Chang CJ (2013). Phosphorylation of mRNA decapping protein Dcp1a
by the ERK signaling pathway during early differentiation of 3T3-L1 preadipocytes.
PLoS One. 8, e61697.

8. Clapham DE (2007) Calcium signalling. Cell 131:1047-1058.

9. Collatz MB, Rüdel R, Brinkmeier H (1997) Intracellular calcium chelator BAPTA

protects cells against toxic calcium overload but also alters physiological calcium

responses. Cell Calcium 21:453-9.

10. Čunátová K, Reguera DP, Vrbacký M, Fernández-Vizarra E, Ding S, Fearnley IM,

Zeviani M, Houštěk J, Mráček T, Pecina P (2021) Loss of COX4I1 Leads to

Combined Respiratory Chain Deficiency and Impaired Mitochondrial Protein

Synthesis. Cells 10(2): 369.

11. Czechanski A, Byers C, Greenstein I, Schrode N, Donahue LR, Hadjantonakis AK,

Reinholdt LG (2014) Derivation and characterization of mouse embryonic stem cells

from permissive and nonpermissive strains. Nat Protoc. 9:559-74.

12. Decker CJ, Parker R (2012) P-bodies and stress granules: possible roles in the control

of translation and mRNA degradation. Cold Spring Harb Perspect Biol. 4(9):

a012286.

13. Fagnocchi L, Cherubini A, Hatsuda H, Fasciani A, Mazzoleni S, Poli V, Berno V,

Rossi RL, Reinbold R, Endele M, Schroeder T, Rocchigiani M, Szkarłat Ż, Oliviero

S, Dalton S, Zippo A (2016). A Myc-driven self-reinforcing regulatory network

maintains mouse embryonic stem cell identity. Nat Commun. 7:11903.

14. Fu W, Liu Y, Yin H (2019) Mitochondrial Dynamics: Biogenesis, Fission, Fusion,

and Mitophagy in the Regulation of Stem Cell Behaviors. Stem Cells Int

2019:9757201.

15. Ghimire S, Van der Jeught M, Neupane J, Roost MS, Anckaert J, Popovic M, Van

Nieuwerburgh F, Mestdagh P, Vandesompele J, Deforce D, Menten B, Chuva de

Sousa Lopes S, De Sutter P, Heindryckx B (2018) Comparative analysis of naive,

primed and ground state pluripotency in mouse embryonic stem cells originating from

the same genetic background. Sci Rep 8:5884.

16. Hirst NL, Lawton SP, Walker AJ (2022) CaMKII regulates neuromuscular activity
and survival of the human blood fluke Schistosoma mansoni. Sci Rep 12(1):19831.

17. Hochstoeger T, Papasaikas P, Piskadlo E, Chao JA (2024) Distinct roles of LARP1
and 4EBP1/2 in regulating translation and stability of 5'TOP mRNAs. Sci Adv
10(7):eadi7830.

18. Hołubowicz R, Wojtas M, Taube M, Kozak M, Ożyhar A, Dobryszycki P (2017)
Effect of calcium ions on structure and stability of the C1q-like domain of otolin-1
from human and zebrafish. FEBS J 284:4278-4297.

19. Huang Jj, Wang Yj, Zhang M, Zhang P, Liang H, Bai HJ, Yu XJ, Yang HT
(2017) Functional expression of the Ca^{2+} signaling machinery in human embryonic
stem cells. Acta Pharmacol Sin 38:1663–1672.

20. Kao SH, Wang WL, Chen CY, Chang YL, Wu YY, Wang YT, Wang SP, Nesvizhskii
AI, Chen YJ, Hong TM, Yang PC (2015) Analysis of Protein Stability by the
Cycloheximide Chase Assay. Bio Protoc. 5:e1374.

21. Kilchert C, Weidner J, Prescianotto-Baschong C, Spang A (2010) Defects in the
secretory pathway and high Ca2+ induce multiple P-bodies. Mol Biol Cell 21(15):
2624-38.

22. Kim SH, Kim MO, Cho YY, Yao K, Kim DJ, Jeong CH, Yu DH, Bae KB, Cho EJ,
Jung SK, Lee MH, Chen H, Kim JY, Bode AM, Dong Z (2014) ERK1 phosphorylates
Nanog to regulate protein stability and stem cell self-renewal. Stem Cell Res 13:1-11.

23. Kobayashi H, Kikyo N (2015) Epigenetic regulation of open chromatin in pluripotent
stem cells. Transl Res 165:18-27.

24. Li T, Han J, Jia L, Hu X, Chen L, Wang Y (2019) PKM2 coordinates glycolysis with
mitochondrial fusion and oxidative phosphorylation. Protein Cell 10:583-594.

25. Lis P, Dyląg M, Niedźwiecka K, Ko YH, Pedersen PL, Goffeau A, Ułaszewski S

(2016) The HK2 Dependent "Warburg Effect" and Mitochondrial Oxidative

Phosphorylation in Cancer: Targets for Effective Therapy with 3-Bromopyruvate.

Molecules. 21(12): 1730.

26. Liu Y, Chang A (2008) Heat shock response relieves ER stress. EMBO J 27(7):1049-

59.

27. Lytton J, Westlin M, Hanley MR (1991) Thapsigargin inhibits the sarcoplasmic or

endoplasmic reticulum Ca-ATPase family of calcium pumps. J Biol Chem

266:17067-71.

28. MacDougall MS, Clarke R, Merrill BJ (2019) Intracellular Ca^{2+} Homeostasis and

Nuclear Export Mediate Exit from Naive Pluripotency. Cell Stem Cell 25(2):210-224.

29. Martinez NJ, Chang HM, Borrajo Jde R, Gregory RI (2013) The co-chaperones

Fkbp4/5 control Argonaute2 expression and facilitate RISC assembly. RNA

19(11):1583-93.

30. MedCalc Software Ltd. Comparison of means calculator.

https://www.medcalc.org/calc/comparison_of_means.php (Version 23.0.1; accessed

August 22, 2024)

31. Mège RM, Ishiyama N (2017) Integration of Cadherin Adhesion and Cytoskeleton

at Adherens Junctions. Cold Spring Harb Perspect Biol 9(5):a028738.

32. Meng Y, Deng C, Xiao X, Wei S, Song C, Wang J, Lei CL, Liu W, Chen G (2025)

The modulation of calcium and chloride channels induces cardiomyocytes from

human pluripotent stem cells. Int J Biol Sci 21(1):95-108.

33. Moore CB, Guthrie EH, Huang MT, Taxman DJ (2010) Short hairpin RNA (shRNA):

design, delivery, and assessment of gene knockdown. Methods Mol Biol 629:141-58.

34. Moore JL, Bhaskar D, Gao F, Matte-Martone C, Du S, Lathrop E, Ganesan S, Shao L,

Norris R, Sanz NC, Annusver K, Kasper M, Cox A, Hendry C, Rieck B,

Krishnaswamy S, Greco V (2024) Correction: Cell cycle controls long-range calcium

signaling in the regenerating epidermis. J Cell Biol 223(4):e20230209503052024c.

Erratum for: J Cell Biol. 2023 Jul 3;222(7):e202302095.

35. Ohtsuka S, Nakai-Futatsugi Y, Niwa H (2015) LIF signal in mouse embryonic stem

cells. JAKSTAT 4(2):e1086520.

36. Park HS, Chow EW, Fu C, Soderblom EJ, Moseley MA, Heitman J, Cardenas ME

(2016) Calcineurin Targets Involved in Stress Survival and Fungal Virulence. PLoS

Pathog 12(9):1005873.

37. Park JH, Chung CG, Park SS, Lee D, Kim KM, Jeong Y, Kim ES, Cho JH, Jeon YM,

Shen CJ, Kim HJ, Hwang D, Lee SB (2020) Cytosolic calcium regulates cytoplasmic

accumulation of TDP-43 through Calpain-A and Importin α3. Elife 9:e60132.

38. Pereverzev A, Gurskaya N, Ermakova GV, Kudryavtseva EI, Markina NM, Kotlobay

AA, Lukyanov SA, Zaraisky AG, Lukyanov KA (2015) Method for quantitative

analysis of nonsense-mediated mRNA decay at the single cell level. Sci Rep 5:7729.

39. Ruan Y, He J, Wu W, He P, Tian Y, Xiao L, Liu G, Wang J, Cheng Y, Zhang S, Yang

Y, Xiong J, Zhao K, Wan Y, Huang H, Zhang J, Jian R (2017) Nac1 promotes self-

renewal of embryonic stem cells through direct transcriptional regulation of c-Myc.

Oncotarget 8(29):47607-47618.

40. Sanford EJ, Smolka MB (2021) Fe-NTA Microcolumn Purification of

Phosphopeptides from Immunoprecipitation (IP) Eluates for Mass Spectrometry

Analysis. Bio Protoc. 11(15):e4113.

41. Schwirtlich M, Emri Z, Antal K, Mate Z, Katarova Z, Szabo G (2010) GABA(A) and

GABA(B) receptors of distinct properties affect oppositely the proliferation of mouse

embryonic stem cells through synergistic elevation of intracellular Ca(2+). FASEB J

24:1218–1228.

42. Scognamiglio R, Cabezas-Wallscheid N, Their MC, Altamura S, Reyes A,

Prendergast ÁM, Baumgärtner D, Carnevalli LS, Atzberger A, Haas S, von Paleske L,

Boroviak T, Wörsdörfer P, Essers MA, Kloz U, Eisenman RN, Edenhofer F, Bertone

P, Huber W, van der Hoeven F, Smith A, Trumpp A (2016) Myc Depletion Induces a

Pluripotent Dormant State Mimicking Diapause. Cell 164(4):668-80.

43. Shang F, Taylor A (2011) Ubiquitin-proteasome pathway and cellular responses to

oxidative stress. Free Radic Biol Med 51:5-16.

44. Shim J, Mukherjee T, Mondal BC, Liu T, Young GC, Wijewarnasuriya DP, Banerjee

U (2013) Olfactory control of blood progenitor maintenance. Cell 155:1141-53.

45. Sim YJ, Kim MS, Nayfeh A, Yun YJ, Kim SJ, Park KT, Kim CH, Kim KS (2017) 2i

Maintains a Naive Ground State in ESCs through Two Distinct Epigenetic

Mechanisms. Stem Cell Reports. 8:1312-1328.

46. Strynadka NCJ, and James MNJ (1991) Towards an understanding of the effects of

calcium on protein structure and function. Current Opinion in Structural Biology 1:

905-914.

47. Sun J, He W, Bai SZ, Peng X, Zhang N, Li HX, Zhang WH, Wang LN, Shao XQ, He

YQ, Yang GD, Wu LY, Wang R, Xu CQ (2013) The expression of calcium-sensing

receptor in mouse embryonic stem cells (mESCs) and its influence on differentiation

of mESC into cardiomyocytes. Differentiation. 5(1-2):32-40.

48. Tang Y, Tian XC (2013) JAK-STAT3 and somatic cell reprogramming. JAKSTAT

2:e24935.

49. Tang D, Chen M, Huang X, Zhang G, Zeng L, Zhang G, Wu S, Wang Y

(2023) SRplot: A free online platform for data visualization and graphing. PLoS One

18:e0294236.

50. Todorova MG, Fuentes E, Soria B, Nadal A, Quesada I (2009) Lysophosphatidic acid

induces Ca2+ mobilization and c-Myc expression in mouse embryonic stem cells via

the phospholipase C pathway. Cell Signal 21:523–528.

51. Torres-Padilla ME, Chambers I (2014) Transcription factor heterogeneity in

pluripotent stem cells: a stochastic advantage. Development 141(11):2173-81.

52. Vallet L, Sánchez-Petidier M, Fernandes R, Naumova N, Merla C, Consales C,

Innamorati G, André FM, Mir LM (2025) Calcium oscillations in mesenchymal stem

cells, a control on cell cycle progression to influence cell fate towards proliferation or

differentiation?. Stem Cell Res Ther 16:446.

53. Waldeck-Weiermair M, Bischof H, Blass S, Deak AT, Klec C, Graier T, Roller C,

Rost R, Eroglu E, Gottschalk B, Hofmann NA, Graier WF, Malli R (2015) Generation

of Red-Shifted Cameleons for Imaging Ca²$\square$ Dynamics of the Endoplasmic

Reticulum. Sensors (Basel) 15(6):13052-68.

54. Wang D, Xu Q, Yuan Q, Jia M, Niu H, Liu X, Zhang J, Young CY, Yuan H (2019)

Proteasome inhibition boosts autophagic degradation of ubiquitinated-AGR2 and

enhances the antitumor efficiency of bevacizumab. Oncogene 38:3458–3474.

55. Xu M, Fralick D, Zheng JZ, Wang B, Tu XM, Feng C (2017) The Differences and

Similarities Between Two-Sample T-Test and Paired T-Test. Shanghai Arch

Psychiatry 29(3):184-188..

56. Yanagida E, Shoji S, Hirayama Y, Yoshikawa F, Otsu K, Uematsu H, Hiraoka M,

Furuichi T, Kawano S (2004) Functional expression of Ca2+ signalingpathwaysin

mouse embryonic stem cells. Cell Calcium 36:135–146.

57. Ye Y, Xie W, Ma Z, Wang X, Wen Y, Li X, Qi H, Wu H, An J, Jiang Y, Lu X, Chen

858   G, Hu S, Blaber EA, Chen X, Chang L, Zhang W (2024) Conserved mechanisms of

859   self-renewal and pluripotency in mouse and human ESCs regulated by simulated

860   microgravity using a 3D clinostat. Cell Death Discov. 10, 68.

861   58. Yeo JC and Ng HH (2013) The transcriptional regulation of pluripotency. Cell Res.

862   23, 20-32.

863   59. Zhu Y, Zhang J Calcium: (2019). A New Guardian of Naive Pluripotency. Cell Stem

864   Cell. 25(2), 169-170.

Figure Legends

Figure 1: Calcium level in mESCs: A) Fluo4 staining, B) DIER reporter expression, C)

Fluo4 and lysotracker staining of mESCs under different conditions (C+L, B2uM, T1nM, C-

L, C-L+T1nM) (n=3-5).

D and E) Bar plot represents the mean intensity of fluo4 staining and D1ER reporter in

mESCs calculated from the confocal images of the above experiments.

F) Bar graph showing the percentage of the proliferation of mESCs treated with B1uM,

B2uM, and B5uM for 72 hrs compared to control using the trypan blue method (n=4).

G) Bar graph showing percentage of the proliferation of mESCs treated with B2uM and

B5uM for 5 days and B2R condition compared to control D2 and D5 respectively (n=4)

H) Western blot detection of cMyc in the presence and absence of B2uM for 72hrs (n=2)

I) Immunofluorescence (IF) detection of cMyc expression in mESCs in the presence of

B2uM for 5 days and B2R condition (n=3)

J) The bar plot represents the mean intensity of cMyc expression calculated from the

confocal images of the above experiments.

K) QPCR of cMyc expression in mESCs treated with B2uM for 72hrs (n=3).

L) Cell cycle analysis of control (D2) and B2uM treated mESCs for 72 hrs (n=4). Significant p-value (0.0085) was detected for B2/D2 (G2/M)

M) Cell cycle analysis of D2, B2uM treated mESCs for 5 days and B2R condition (n=5). Significant p-value (0.0003 and 0.01) was detected for B2/D2 and B2R/B2 (G2/M)

C+L= Control media + Lif; B1uM= C+L + Bapta 1uM; B2uM = C+ L + Bapta 2uM; T1nM= C+L + thapsigargin 1nM; C-L= control media without Lif; C-L+T1nM = C-L + thapsigargin 1nM; D2= C+L+ dmso equivalent to B2uM condition; B5uM = C+L+ Bapta 5uM; D5= C+L+ dmso equivalent to B5uM condition, B2R= B2 media for 72 hour + D2 media for 48 hour

The mean intensity was calculated from the confocal images for all experiments using ImageJ software. p-value representation with asterisk <0.05 (*), <0.01 (**), <0.001 (***), <0.0001 (****).

Figure 2: Role of calcium in pluripotency: IF detection of **A)** Nanog (red), **B)** Oct4 (green), **C)** pCamkIIα (red), **D)** Oct4, pCamkIIα and Dapi co-staining in mESCs cultured under different conditions (C+L, B2uM, T1nM, C-L, C-L+T1nM) (n=3)

E) Western blot shows the expression of Oct4, Nanog, pSTAT3 and Gapdh in mESCs cultured under different conditions (C+L, B1uM, B2uM, C-L) (n=3).

F) Western blot shows the expression of Oct4, Nanog, pSTAT3 and Gapdh in mESCs cultured under different condition (C+L+T250pM, C+L+T500pM, C+L+T1nM) (n=3)

G) Western blot showed expression of Nanog in mESCs cultured under different conditions (C-L, C-L+T250pM, C-L+T500pM, C-L+T1nM)

H) Western blot showed expression of pSTAT3 and cMyC in mESCs cultured under different conditions (C-L, C-L+T1nM)

I, J) QPCR of pluripotency markers and early differentiation markers under different conditions (C+L, B2uM, T1nM, C-L, C-L+T1nM) (n=5)

K) Brightfield images of the mESCs cultured under different conditions (C+L, B2uM, T1nM, C-L, C-L+T1nM) (n=5)

L) IF detection of Phalloidin in mESCs cultured under different conditions as above (n=3)

M) The bar graph represents the mean intensity of Oct4, Nanog and pCamkIIα expression calculated from the confocal images of the above experiments (A, B, C).

N) IF detection of E-cadherin staining in mESCs cultured under different conditions (C+L, B2uM, T1nM, C-L)

Note: C+L= Control media + Lif; B1uM= C+L + Bapta 1uM; B2uM = C+ L + Bapta 2uM; T1nM= C+L + thapsigargin 1nM; C-L= control media without Lif; C-L+T1nM = C-L + thapsigargin 1nM; T250 pM, T500pM, and T nM= C+L + thapsigargin (250 pM, 500pM, and 1nM)

The mean intensity was calculated from the confocal images for all experiments using ImageJ software. p-value representation with asterisk <0.05 (*), <0.01 (**), <0.001 (***), <0.0001 (****).

Figure 3: Role of calcium at post-transcriptional level: A) IF detection of dcp1a, β-actin, xrn1a, pCamkIIα, tudor in mESCs cultured under different conditions (C+L, B2uM, T1nM, C-L) (n=3-5) using confocal microscopy at 60x magnification.

B) Co-localization expression profile of dcp1a and β-actin; dcp1a and xrn1; dcp1a and pCamkIIα; dcp1a and tudor in mESCs cultured under different condition (C+L, B2uM, T1nM, C-L, C-L+T1nM). To observe granule size and co-localization of p-bodies and stress

granule markers, the images obtained using a confocal microscope at 60x magnification were

cropped to one nucleus size and normalized using a 5uM marker at the bottom.

C) QPCR of p-bodies and stress granule markers (dcp1a, xrn1, **t**udor and pCamkIIα) in the

mESCs cultured in the presence of B2uM for 72hrs.

D) The bar graph represents the mean intensity of dcp1a, xrn1, pCamkIIα and **t**udor

expression calculated from confocal images from the above experiments (A).

E) IF detection of splicing dependent NMD by using NMD+ GFP, NMD+RFP, NMD-GFP,

NMD-RFP reporter in mESCs cultured under different conditions (C+L, B2uM, C-L) (n=3).

F) Splicing-dependent NMD activity was calculated from the confocal images of the above

experiments described by Pereverzev et al. 2015.

G) IF detection of long 3' UTR dependent NMD by using SMG+ GFP, SMG+RFP, pTRAM-

GFP, pTRAM-RFP reporter in mESCs cultured under different conditions (C+L, B2uM, C-

L) (n=3).

H) 3' UTR-dependent NMD activity was calculated from the confocal images of the above

experiments as described by Pereverzev et al. 2015.

Note: C+L= Control media + Lif; B2uM = C+ L + Bapta 2uM; T1nM= C+L + thapsigargin

1nM; C-L= control media without Lif; C-L+T1nM = C-L + thapsigargin 1nM

The mean intensity was calculated from the confocal images for all experiments using

ImageJ software. p-value representation with asterisk <0.05 (*), <0.01 (**), <0.001 (***),

<0.0001 (****).

Figure 4: **Role of calcium at translational level: A)** Polysome profile of C+L and B2uM

treated mESCs. (n=5)

B) IF detection of Elf1a expression in mESCs cultured under different conditions (C+L, B2uM, C-L) using confocal microscopy at 60x magnification. (n=3)

C) The bar graph represents the mean intensity of Elf1a expression calculated from the confocal images of the above experiments.

D, E) QPCR of core proliferation and early differentiation markers in the cytoplasmic, 80s, polysome fraction of polysome profiling of control (D2) and B2uM treated mESCs. (n=4)

F, G) QPCR of core proliferation markers and early differentiation markers of mESCS cultured under different conditions (C+L, B2uM, C-**L, C-L+Tti)** for 72hrs followed by combined CHX and ACD treatment (TTi) for 2hr. (n=5)

H) IF detection of brachyury and Nanog expression in mESCs cultured under different conditions (C+L, B2uM, C-L). (n=3)

I) Confocal detection of mitochondria using mitotracker-red staining in mESCs cultured under different conditions (C+L, B2uM, C-L) for 48 hours using confocal microscopy at 40x magnification. To observe the mitochondria shape, images were cropped to one nucleus and normalized using a 5uM marker at the bottom. (n=4)

J) Confocal detection of mitochondria and calcium level using mitotracker-red and fluo-4 co-staining in mESCs cultured under condition (C+L and B2uM) (n=3).

Note: C+L= Control media + Lif; B2uM = C+ L + Bapta 2uM; C-L= control media without Lif; C-L+T1nM = C-L + thapsigargin 1nM

The mean intensity was calculated from the confocal images for all experiments using ImageJ software. p-value representation with asterisk <0.05 (*), <0.01 (**), <0.001 (***), <0.0001 (****).

Figure 5: Role of calcium at post-translational level: IF detection of **A)** Brachyury, and cMyc, **B)** dcp1a, and Tudor, **C)** Oct4 and pUBQ expression mESCs cultured under different

conditions (D2, B2uM and C-L) for 72hrs followed by MG132 (10uM) treatment for 1 hr (n=3).

The bar plot represents the mean intensity of the expression profile of **D)** Brachyury, **E)** cMyc, **F)** dcp1a, and Tudor, **G)** Oct4 and pUBQ from the confocal images of the above experiments.

H) Western blot detection of autophagy marker LC3 in mESCs cultured under different conditions (C+L, B2uM, C-L). (n=3)

I) Cycloheximide chase assay was performed for oct4 and Nanog protein in mESCs cultured under D2, B2uM conditions for 72hrs followed by addition of CHX (50 ug/ml) and chased for every 20min for 1 hr 20 min using western blotting. (n=3)

J) Oct4 and Nanog expression was detected by western blot in mESCs cultured under D2, B2uM conditions for 72hrs followed by the addition of MG132 (10uM) for 1hr. (n=3)

Note: C+L= Control media + Lif; C-L= control media without Lif; D2= C+L+ dmso equivalent to B2uM condition; D2_MG = D2+MG132; B2uM= C+L+ Bapta 2uM; B2_MG = B2uM+MG132; C-L_MG= C-L+MG132

The mean intensity was calculated from the confocal images for all experiments using ImageJ software. p-value representation with asterisk <0.05 (*), <0.01 (**), <0.001 (***), <0.0001 (****).

Figure 6: knockdown of pCamkIIα in mESCsA) Detection of GFP expression of pCamkIIα shRNA and SERCA shRNA transfected by electroporation method in mESCs using fluorescent microscopy at 10X.

Brightfield images showing the morphology of the **B)** control mESCs, **C)** pCamkIIα shRNA (pCamkIIα KD) and SERCA shRNA (SER_R2_KD) transfected mESCs at 48hrs.

D) QPCR of pCamkIIα and serca R2 receptor expression in the mESCs at 48hrs upon pCamkIIα KD and SER_R2_ KD respectively

E) QPCR of pCamkIIα expression in the SER_R2 KD mESCs and serca_R2 receptor expression in the pCamkIIα KD mESCs at 48hrs.

F, G) Western blot detection of Oct4, Nanog, pSTAT3, and Gapdh expression in the pCamkIIα KD and SER_R2_KD mESCs at 48hrs (n=3)

H, I) IF detection of DCP1a and pCamkIIα co-expression in pCamkIIα and SER_R2 KD mESCs at 48hrs. (n=3)

J, K) Bar plot represents the mean intensity of pCamkIIα and dcp1a expression in mESCs calculated from the confocal images of the above experiments

L) QPCR of p-bodies and stress granule markers in the mESCs at 48hrs upon pCamkIIα KD and ser_R2 KD. (n=4)

M, N) QPCR of pluripotency and differentiation markers in the mESCs at 48hrs upon pCamkIIα KD and SER_R2 KD. (n=3)

Note: The mean intensity was calculated from the confocal images for all experiments using ImageJ software. p-value representation with asterisk <0.05 (*), <0.01 (**), <0.001 (***), <0.0001 (****).

Figure 7: Calcium and phosphoproteomics in mESCs: A) Heat map shows log2fold change of DEPP in B2uM treated mESCs compared to D2 (control) mESCs detected by mass spectrometry (n=2)

B, C, E) KEGG pathway and GOC enrichment analysis was done using ShinyGO 0.80 software for the DEPP represented in the heat map (A).

D) The hierarchical clustering tree shows the correlation among the most significant pathways found in KEGG pathway enrichment analysis for all DEPP.

F) The bar plot shows the string enrichment function analysis of all the DEPP demonstrated in the heat map.

G, H) KEGG pathway enrichment analysis was done for spliceosome and mRNA surveillance pathways all the DEPP represented in heat map (A).

I) String PPI network analysis was performed for all DEPP represented in a heat map (A).

Note: Arrow represent the upregulated and downregulated genes

Note: C+L= Control media + Lif; B2uM = C+ L + Bapta 2uM; D2= C+L+ dmso equivalent to B2uM condition

p-value representation with asterisk <0.05 (*), <0.01 (**), <0.001 (***), <0.0001 (****).