A long non-coding RNA regulates in vitro and in vivo triazole antifungal susceptibility in *Aspergillus fumigatus*

Nava Raj Poudyal[1,2], Ryan T. Mehlem[2,3], Priyanka Doneparthi[1,2*], Tabitha L. Cady[1,2^],
Garrett Kaufman[2,4], Sven D. Willger[5#], Maximiliano Ortiz[6], Rooksana E. Noorai[6], Jason
E. Stajich[7] and Sourabh Dhingra[1,2,8]

[1]Department of Biological Sciences, Clemson University, Clemson, SC 29634,
USA
[2]Eukaryotic Pathogen Innovation Center, Clemson University, Clemson, SC
29634, USA
[3]Department of Genetics and Biochemistry, Clemson University, Clemson, SC
29634, USA
[4]USC School of Medicine-Greenville, Greenville, SC 29605, USA
[5]Department of Immunology and Infectious Disease, Montana State University,
Bozeman, MT 59717, USA
[6]Clemson University Genomics & Bioinformatics Facility, Clemson University,
Clemson, SC 29634, USA
[7]Department of Microbiology and Plant Pathology, University of California-
Riverside, Riverside, CA 92521, USA

* Present address: Department of Oncology, Georgetown University, Washington, DC,
USA, pd771@georgetown.edu

^ Present address: Biology Department, Morrissey College of Arts and Sciences, Boston
College, Chestnut Hill, MA 02467, USA, tabithacady@gmail.com

Present address: Independent Researcher, Düsseldorf, Germany,
sven.willger@gmail.com

[8]Address correspondence to: sdhingr@clemson.edu

Abstract:

Azole-resistant *Aspergillus* infections are a source of increasing concern with limited alternative therapeutic options. However, as most infections are still caused by azole-susceptible *Aspergillus* strains, there is a need to better understand fungal responses to azole antifungals. To this end, we discover that a long non-coding RNA, *afu-182,* is a major regulator of *cyp51*-independent sub-MIC azole response. We observe that loss of *afu-182* leads to increased surface attached growth and poor treated disease outcomes in a murine model of invasive pulmonary aspergillosis upon azole treatment. In contrast, overexpression of *afu-182* significantly reduces fungal burden in animals treated with the azole drug, posaconazole. Importantly, *afu-182* levels decrease upon azole exposure and in an azole adaptation experiment, continuous exposure to low dose azole led to MIC increase in an *afu-182* dependent manner. Whole transcriptome analyses revealed that azole drug treatment leads to an increase in transcripts of genes encoding 7-transmembrane domain proteins of the RTA1 family, and these proteins are negatively regulated by *afu-182*. Two RTA1 family genes have individual and combined effects and are sufficient to increase fungal susceptibility to azole drugs in the WT strain. Taken together, our data show a role of the long non-coding RNA *afu-182* in regulating *Aspergillus fumigatus* response to azole drugs both *in vitro* and *in vivo.*

Importance:

Drug resistance in *Aspergillus* is a major challenge that is often associated with the agricultural use of azoles in the environment. How drug resistance arises *in vivo* is still an active area of research. Here, we show that azole exposure results in fungal adaptation by lowering the RNA levels of lncRNA, which upon low dose azole exposure leads to an increase in azole drug MIC.

Introduction:

Aspergillus fumigatus causes a spectrum of infections termed as Aspergillosis, mainly in patients with compromised immune systems (1). The patient population most susceptible to infections is those who have acute neutropenia due to underlying conditions or immunotherapies, or patients on long-term steroid use (2, 3). Recent studies have shown that *A. fumigatus* is a significant cause of secondary infections, leading to increased morbidity and mortality in patients with viral infections, including COVID-19 and influenza A virus (4, 5). Importantly, *A. fumigatus* is increasingly a source of infection in tertiary care centers in patients with mild to no immune suppression, highlighting the prevalence of the infection and increase in the patient population that is susceptible to aspergillosis (6). The World Health Organization (WHO) classified drug resistant *A. fumigatus* as a critical pathogen in 2023 (7).

Azoles, polyenes, and echinocandins are the current classes of antifungal drugs available for aspergillosis treatment; azoles are the most widely used (8). Clinically, voriconazole is the frontline therapy; however, posaconazole, itraconazole and isavuconazole are also used (8). Concerningly, azole drug resistance is on the rise (9). Azoles inhibit the sterol 14-α-demethylase gene, *cyp51* (*erg11* in yeast), which leads to accumulation of toxic sterols (10). Molecularly, resistance to azoles is primarily attributed to mutational hotspots in the *cyp51* gene sequence that affect drug-enzyme binding (11); with or without the replication of a 34 base pair or 46 base pair tandem repeat (TR) in the promoter region of the *cyp51a* gene (commonly referred to as TR_{34} and TR_{46}, respectively) (12–14). TRs in promoter regions increase the binding site of transcription factor SrbA, which positively regulates *cyp51a* transcription (15). SrbA also regulates ergosterol biosynthesis by transcriptional regulation of *erg3*, *erg25* and *erg24* genes (16, 17). In addition, transcription factor *atrR* cooperates with SrbA for azole resistance (18).

SrbA is a class of ER resident sterol regulator binding element protein (SREBP) that, in association with SREBP cleavage activating protein (Scap) and Insulin induced gene (INSIG), regulates cholesterol levels in mammalian cells (19, 20). In *A. fumigatus*, SrbA plays a significant role in the hypoxia response, low iron response, virulence, and

azole susceptibility (19); however, a Scap homolog has not been identified in any

Eurotiomycete fungus to date, and the Insig homolog *insA* does not regulate SREBPs or

subsequent hypoxia response in these fungi (21). In the mammalian system, INSIG

senses the accumulation of sterol pathway intermediates and negatively regulates

HMG-CoA reductase, thus blocking the rate-limiting step in sterol biosynthesis (22). In

A. fumigatus, mutations in HMG-CoA reductase (HMG-CoA) have been recently

identified in clinical isolates (23–26), and these mutations, especially in the sterol-

sensing domain (SSD), are predicted to inhibit sterol intermediate sensing by changing

HMG-CoA-Insig stochiometric balance (27). Thus, *insA* provides a feedback mechanism

that contributes to azole resistance (27). Interestingly, overexpression (OE) of *insA*

reverses the azole resistance in OE HMG-CoA mutants but not in SSD mutants,

highlighting the importance of sterol sensing by HMG-CoA (27).

In a laboratory setting, drug resistance is defined as a change in minimum

inhibitory concentration (MIC) above the clinical cut-off value measured in standardized

MIC testing assays (28). Multiple methods are used to determine the MICs of the

antifungal drugs in the laboratory setting, including broth microdilution, disk diffusion

assay and E-Test strips (28, 29), and strains are then classified as susceptible,

resistant, tolerant, persistent or exhibiting hetero-resistance (30) based on the ability of

all cells or a sub-population of cells to grow at or above MIC (30, 31).

Here, we show that *insA* adjacent long non-coding RNA *afu-182* negatively

regulates and aids in fungal adaptation to the azole drugs at sub-MIC concentrations. In

an animal model of IPA, the Δ*afu-182* strain causes azole recalcitrant infections,

whereas OE-182 strains show increased clearance from the lungs when treated with the

azole drug, posaconazole. Upon low dose exposure to azoles, Δ*afu-182* strain adapts

and has an increase in the MIC for voriconazole, whereas the WT strain does not show

a change in MIC.

Collectively, we observed that *afu-182* regulates fungal azole response

independent of *cyp51* regulation and plays a significant role in azole treatment

outcomes, especially in strains classified as azole-susceptible. Thus, our study has

uncovered a novel role of long ncRNA *afu-182* in azole susceptibility both *in vitro* and *in*

vivo.

Results:

afu-182 is an intergenic long non-coding RNA:

afu-182 was previously characterized and annotated as an anti-sense ncRNA to the
gene *insA* (32, 33). To determine if *afu-182* is an anti-sense or intergenic ncRNA, we
performed 3' rapid amplification of cDNA ends (RACE) (34) for *insA* and showed that
insA transcript ends 191bp after the translational stop codon (Figure 1). A non-species
specific computational analysis assigned *afu-182* sequence a Fickett score of 0.40586
with a protein coding probability of 0.016 (35), and is classified as an intergenic long
non-coding RNA.

afu-182 regulates sub-MIC response to azole class of anti-fungal drugs

afu-182 neighboring INSIG homolog gene *insA,* was recently shown to play a role in
azole drug susceptibility (27) and Supplementary Figure 1A. However, when we
complemented *insA* ectopically, we did not see phenotypic reversion (Supplementary
Figure 1A, red arrows); even when the *insA* RNA levels were rescued (Supplementary
Figure 1B). The ∆*insA* strain does have inherent *afu-182* RNA (Supplementary Figure
1C); however, to determine the cis-acting roles of ncRNA, we also complemented the
∆*insA* strain with DNA fragment containing both *insA* fragment and *afu-182* fragment
and saw partial reversal of phenotype (Supplementary Figure 1A, orange arrows);
however, no change in voriconazole MIC was observed (Supplementary Figure 1D).

We compared the RNA levels of *afu-182* in ∆*insA* and comp-*insA* strains. Even
though, statistically insignificant, we did see 2-fold lower levels of *afu-182* RNA levels in
comp-insA strain that did not revert the phenotype (Supplementary Figure 1C). Thus, to
determine if *afu-182* levels regulate fungal response to sub-MIC azole concentrations,
we made a ∆*afu-182* and over-expression 182 (OE-182) strains. We also
complemented *afu-182* back at the original locus (Comp-182). We confirmed the levels
of *afu-182* in the complement and OE strain and the absence of RNA in the ∆*afu-182*

strain (Supplementary Figure 2A). Our qPCR data show the absence of *afu-182*

transcript in the Δ*afu-182* strain and WT level of expression in the Comp-182 strain

(p=0.5469, One-Way ANOVA, Supplementary Figure 2A). We observed a 4-fold

overexpression in the OE-182 strain (p = 0.0025, One-Way ANOVA, Supplementary

Figure 2A). To assess the role of *afu-182* in fungal azole response, we serially

inoculated the strains (10^5-10^2 conidial density) in the absence and presence of

0.25µg/ml voriconazole, itraconazole (0.075µg/ml), posaconazole (0.03µg/ml), and

isavuconazole (0.25µg/ml). The *afu-182* null mutant strain showed increased diametric

growth against all azoles, confirming that *afu-182* lncRNA regulates sub-MIC fungal

azole response to azole class of drugs (Figures 2A – 2E), and importantly this effect

was not due to a change in MIC (Supplementary Figure 2B). OE-182 showed

decreased diametric fungal growth to azoles, suggesting a causal link between *afu-182*

levels and fungal azole response (Figures 2A-2E).

It is possible that *afu-182* inhibits fungal germination and not the azole stress

response. Thus, we assessed the role of *afu-182* in fungal azole response post-

germination in surface attached cultures. We allowed all strains to germinate and

adhere to the abiotic surface (i.e., cell culture plates) for 12 hours. We then washed the

fungal biofilms with PBS and grew the biofilms in the presence of voriconazole

(0.2µg/ml) for 8 additional hours. The final biofilms were washed, stained with crystal

violet and de-stained with ethanol as described previously (36). The binding of crystal

violet was measured as an indirect measure of fungal biomass. We observed increased

biofilm formation in the Δ*afu-182* upon azole treatment (41% increase compared to WT,

p=0.0049, One-Way ANOVA), whereas the OE-182 mutant showed severe and

significant impairment in biofilm development after drug treatment (51% decrease

compared to WT, p=0.0011, One-Way ANOVA) (Figure 2F), indicating *afu-182* mediated

fungal azole response is independent of spore germination. No difference across strains

was observed in initial biofilm formation after 12 hours (Supplementary Figure 2C).

Additionally, to understand the role of growth condition-specific role of *afu-182*,

we grew AF as liquid submerged cultures in the presence and absence of voriconazole

for 48 hours and weighed dry fungal biomass. The Δ*afu-182* strain showed significantly

increased biomass (116% increase compared to WT, p=0.0017, One-Way ANOVA),

whereas the OE-182 strain showed significant impairment (76% decrease compared to

WT, p=0.0219, One-Way ANOVA, Figure 2G). Thus, our data show that *afu-182*

influences fungal azole response in both submerged and surface-attached cultures.

RNA levels of *afu-182* are negatively regulated by azole drugs, and correlate with

azole MIC:

To understand the role of *afu-182* in fungal azole response, we tested the RNA levels of

afu-182 in the presence and absence of azoles. A significant reduction in levels of *afu-*

182 was observed in strains grown in the presence of either voriconazole or

posaconazole compared to GMM (Figure 3A). Multiple strains of *A. fumigatus,* including

A1163, CEA10 and Af293, are used as laboratory WT strains (37). Additionally, multiple

reports have shown significant strain virulence heterogeneity, thus making it hard to

establish a true WT strain (38); however, a comprehensive study of azole response in

ASAF strains is lacking. To further determine the role of *afu-182* in regulating fungal

azole response, we quantified the relative abundance (to CEA10) of *afu-182* RNA levels

in strain Af293, environmental isolates 47-57, 47-7, 08-19-02-46 (referred as - 46) and

08-19-02-10 (referred as -10). Interestingly, *afu-182* RNA levels differ in AF strains and

negatively correlate with azole response i.e. we observed lower MIC for strains trending

higher for *afu-182* levels compared to CEA10 (47-7 and 47-57) and increased MIC for

strains trending lower for *afu-182* RNA levels compared to CEA10 (Figures 3B and 3C).

To confirm if *afu-182* is sufficient to regulate the azole response in multiple isolates, we

made a Δ*afu-182* mutant in the Af293.1 background, and the resulting strain showed

increased fungal growth at sub-MIC voriconazole concentration of 0.4μg/ml (Figures 3D

and 3E). Environmental isolate 47-57 has increased RNA levels of *afu-182* compared to

CEA10 and is more susceptible to voriconazole compared to CEA10 (Figure 3C).

Deletion of *afu-182* in the 47-57 background also resulted in increased fungal growth of

Δ*afu-182* strain compared to the host strain at sub-MIC voriconazole concentration of

0.1μg/ml (Figure 3F). These two isolates with azole sensitivity on either side of CEA10

showed increased fungal growth upon *afu-182* loss, highlighting the role of *afu-182*

irrespective of the strain's background. To further characterize the role of *afu-182* in

fungal azole response, we identified the *afu-182* homolog in the related species *A.*

nidulans. The deletion of *an-182* (Δ*an-182*) in two distinct AN strains (A1145 and GR5)

also leads to increased fungal growth (Supplementary Figure 3, orange arrows),

indicating roles of lncRNA *afu*-182 homologs in multiple Aspergilli species.

afu-182 does not regulate transcription of azole target gene cyp51 or transcription

factors srbA and atrR

To further test the role *of afu-182* in regulating fugal sub-MIC azole response, we

quantified the mRNA levels of azole target genes *cyp51a* and *cyp51b,* both in the

presence and absence of azoles and TFs *srbA* and *atrR* in GMM. Importantly, we did

not see a change in the transcript levels of *cyp51* genes in the absence of azoles

(Supplementary Figures 4A and 4B). While in the presence of azoles, we clearly see an

upregulation of *cyp51 genes* based on drug response, no upregulation was seen due to

genotype (Two-Way ANOVA, interaction p-value >0.5, Supplementary Figures 4A and

4B). We also did not see a change in mRNA levels of transcription factors, *srbA* and

atrR, that are mainly associated with *cyp51a/b* expression. (Supplementary Figure 4C

and 4D).

Loss of afu-182 results in an azole recalcitrant infection in a murine model of

invasive pulmonary aspergillosis.

To test the role of *afu-182* during infection, we used a corticosteroid model of invasive

pulmonary aspergillosis (39). We challenged mice with 2×10^6 spores of the respective

strains and monitored mice for 14 days for survival (Figure 4A). We did not see a

difference in survival rates across strains (Figure 4B, log-rank test, p-value 0.1166

between WT andΔ*afu-182* strains). Also, we did not see a qualitative difference in fungal

growth or histopathology as determined by GMS and H&E staining, respectively

(Supplementary Figure 5). We next assessed the impact of *afu-182* loss on azole

treatment outcome. Posaconazole was selected due to its higher bioavailability in the

absence of grapefruit juice compared to voriconazole in mice (40, 41). Mice were

challenged as previously described (39) and were treated with 5mg/kg of posaconazole

at 24, 48 and 72 hours of infection (Figure 4C) . Mice challenged with Δ*afu-182* strains

had significantly more adverse outcomes compared to mice infected with the WT strain

(p = 0.0103, log-rank test) (Figure 4D). Thus, the Δ*afu-182* strain that is as virulent as

WT in the absence of drug is recalcitrant to treatment in the presence of the azole drug.

Additionally, we challenged and treated mice and harvested lungs after 72 hours (Figure

4E, schematic). We measured the fungal DNA as previously described (40). Our data

indicate a significantly higher fungal burden in mice challenged with the Δ*afu-182* strain

(p=0.0159, Mann-Whitney U-test) and a significantly reduced fungal burden in mice

challenged with OE-182 strain (p=0.0159, Mann-Whitney U-test) upon azole treatment

(Figure 4F); however, survival of animals infected with OE-182 strain and treated with

posaconazole was not different compared to WT (p=0.4065, log-rank test, Figure 4D).

Sub-MIC azole exposure results in increased azole Minimum inhibitory

concentration in an *afu-182* dependent manner:

Lower drug concentrations at the infectious lesions compared to serum drug level is a

challenging problem in treating fungal infections and fungus might be exposed to sub-

MIC drug concentrations during infections. To determine the role of *afu-182* mediated

sub-MIC response in the development of drug resistance i.e., change in MIC, we

exposed WT and Δ*afu-182* strains to low dose azole at 0.1 μg/ml voriconazole every 48

hours for 12 generations in ten independent lineages (Supplementary Figure 6). After 12

generations, we see an increase in voriconazole MIC only in Δ*afu-182* strain in 6

lineages out of 10, but never in the WT strain, indicating that sub-MIC azole adaptation

leads to an increase in MIC as measured by broth microdilution or E-Test strip in an *afu-*

182 dependent manner (Figures 5A and 5B). To further see the growth difference on

solid medium we grew the adapted strains at varying spore densities (10^5-10^2) in

presence of 0.5μg/ml voriconazole and see robust growth in Δ*afu-182* adapted strains

but not in WT adapted strain (Figure 5C, red arrows). We did observe increase in sub-

MIC (0.25μg/ml) growth of WT adapted strain (Figure 5C, orange arrow) but no

discernable growth was observed at 0.5μg/ml voriconazole. We further tested the levels

of azole target genes *cyp51a* and *cyp51b* and did not see a difference in RNA levels,

indicating that the change in resistance in voriconazole adapted strain is not associated
with transcriptional regulation of azole target genes (Supplementary Figure 7).
278
279 ***afu-182* regulates the RTA family of proteins in the presence of azoles:**
280
To determine the effects of azole on global transcription in WT on solid medium, we
compared the transcriptomic profile of WT strain in the presence and absence of azoles
with a stringent threshold change of 4-fold or more. We identified 4 genes
(*afub_000050*, *afub_000060*, *afub_049700* and *afub_046760*) that are upregulated 4
fold or more in presence of azole in WT (Supplementary Figure 8) that contain a domain
with predicted functions in stress response and belong to the RTA1family of proteins
(32). In *Saccharomyces cerevisiae*, members of this family are involved in resistance to
7-aminocholestrol (RTA), resistance to molasses (RTM), and lipid and heme transport
(42). Previous report has suggested that a member of the RTA1 family is the most
increased transcript in response to itraconazole in clinical samples (43). In *A. fumigatus*,
we identified 23 genes in the genome encoding RTA1 family proteins. Interestingly, most
of these genes are expressed at very low RNA levels (FPKM <5 for 17 genes in the
glucose minimal medium conditions examined) and thus did not make the cut-off for
global transcriptomic analysis. We selected 12 genes that showed significant change (2-
fold or more) in transcript level and have an FPKM value >1 in GMM samples and
performed qPCR. Comparative analysis revealed that all 12 gene transcripts were
increased in the presence of azoles in WT (Figure 6).
To determine the role of *afu-182* in regulating fungal azole response, we
compared the transcriptome of WT and Δ*afu-182* strain and saw 15 genes to be
upregulated and 10 genes to be downregulated in Δ*afu-182* strain by 4-fold strain
compared to WT in the presence of azoles out of about 10,000 variables (Figure 7A).
Interestingly, two genes of the RTA1 family were the most upregulated genes in the
Δ*afu-182* strain compared to WT indicating a negative regulation (Figure 7A, orange
arrows) were upregulated in presence of azoles in WT. In the absence of azoles, their
expression was not different (Figure 7B).

Of the 23 genes in the RTA1 class, *afub_00050* and *afub_00060* are annotated

as RTM1 proteins, a homolog of the *S. cerevisiae* gene providing resistance to

molasses (44). We termed them as *rtmA* and *rtmB,* respectively.

rtmA and rtmB regulate sub-MIC fungal azole response:

To ascertain the role of *rtmA* and *rtmB* in fungal azole response, we deleted them in the

CEA10 background. Additionally, we made a double null mutant Δ*rtmA*Δ*rtmB* strain. We

also ectopically complemented *rtmA* and *rtmB* single mutants. Both the Δ*rtmA* (31%

decrease in growth ratio, p=0.0012 One-Way ANOVA) and Δ*rtmB* (42% decrease in

growth ratio, p<0.0001, One-Way ANOVA) strain showed reduced fungal growth in

response to voriconazole (Figures 8A-B, and 8C-D), whereas the Δ*rtmA*Δ*rtmB* strain

also showed significant growth attenuation in the presence of voriconazole (Figures 8E

and 8F, 46% decrease in growth ratio p<0.001, One-Way ANOVA), indicating a

combined role of these genes in regulating fungal azole response, where *rtmB* is the

major driver of the azole response phenotype. The complement strains reverted to the

WT levels, indicating a causal role of these genes in fungal azole response downstream

of *afu-182* (Figures 8A-D, inset – macro lens images). Further studies are needed to

determine the functional redundancy of other genes in this family; however, recently, an

RTA1 family protein, *rtaA,* was shown to play no role in azole response (45). The

functions of these genes appear to be diverse.

Discussion:

While increasing azole resistance in *A. fumigatus* is a major concern, azole-

susceptible strains (ASAF) still cause most infections (46–48). Agricultural use of azoles

is a major source of *cyp51* mediated environmental azole resistance (49). This hints at a

broader response to AF drugs, especially in ASAF, during infections. Here, we show that

the long non-coding RNA *afu-182* negatively regulates fungal response to sub-MIC

azole drug levels in ASAF (Figure 2), and deletion of *afu-182* is not associated with a

change in MIC (Supplementary Figure 2D). *afu-182* was previously identified as an anti-

sense ncRNA to *insA* (32) that was recently shown to play a role in azole drug

resistance (27); however, the role of *afu-182* has been obscure. (32, 33). Here, we show

that *afu-182* lncRNA regulate fungal response to sub-MIC azole drugs.

Fungal response to drugs has been characterized based on changes in genetic

sequence (i.e., resistance) or non-genetic mechanisms (tolerance, persistence, and

hetero-resistance) (28, 30, 31). We see robust growth of Δ*afu-182* strain at azole

concentrations approaching MIC but all strains fail to germinate at 2x MIC concentration

(Supplementary Figure 9). Thus, the Δ*afu-182*-mediated phenotype at sub-MIC azole

concentration is likely a stress adaptation mechanism to tolerate azole drugs.

Furthermore, the Δ*afu-182*-mediated sub-MIC growth phenotype is independent of the

strain's background and is conserved in related species *A. nidulans* (Supplementary

Figure 3).

We also showed the increased fungal growth upon azole treatment in fungal

surface attached cultures, thus indicating effects are independent of spore germination

(Figure 2F). Drug resistance is a classical feature of the biofilms (50), and even though

we saw increased attached fungal biomass for the Δ*afu-182* strain upon azole treatment

(Figure 2F), it is possible that different mechanisms are playing roles in this distinct

growth conditions. For example, metabolic adaptation, drug binding to the ECM, and

oxygen gradients have all been shown to play a role in biofilm mediated drug resistance

(51, 52). Additionally, biofilms can lead to persister cells (53). Further research will

uncover if distinct mechanisms confer difference in azole response in surface attached

cultures.

Different AF isolates show differential *afu-182* levels, implying that *afu-182* might

be involved in fungal adaptation to various stresses, though in our murine model, it was

not required for virulence (Figure 4B and Supplementary Figure 5). Thus, the

evolutionary significance of *afu-182* and its role in adaptation against other stresses in

AF is currently under investigation. Importantly, the Δ*afu-182* strain shows increased

fungal growth only in response to azole stress and does not play a role in response to

the polyene and echinocandin classes of drugs (Supplementary Figure 10). Over 90%

of infections are still caused by ASAF, with treatment failure rates approaching 50-90%

(54, 55). Thus, it is imperative to better understand the fungal response to azole drugs.

To understand the role of *afu-182* in fungal azole response *in vivo*, we used a
steroid model of IPA. In the absence of azole drugs, the Δ*afu-182* strain is pathogenic
and equally virulent as WT (Figure 4B); however, in the presence of 5mg/kg of
posaconazole, our data show increased virulence in the Δ*afu-182* strain as measured
by survival analysis and fungal burden analysis (Figure 4D and 4F). The overexpression
strain did not show a change in survival rate compared to WT when treated with azole
drugs (Figure 4D); however, it showed a 90% reduction in fungal burden, highlighting
the role of *afu-182* in azole treatment outcomes (Figure 4F). In a steroid model of
invasive aspergillosis, deaths due to exacerbated immune response have been reported
(55), and it is plausible that this is the case in OE-182-infected azole-treated animals. In
vitro, we see that *afu-182* regulates sub-MIC response to all azoles, including
posaconazole (Figures 2A and 2D). Like *in vitro* response, it is possible that low levels
of azole exposure in fungal lesions lower the *afu-182* RNA levels that allow the AF to
grow better in the presence of the drug, thus increasing fungal burden in Δ*afu-182* strain
and lower burden in OE-182 strain. As drug resistant isolates are recovered from
patients during azole treatment if IA or IPA (56), this sub-MIC azole adaptation may
eventually lead to drug resistance. To test this *in vitro*, we adapted the WT and the Δ*afu-*
182 strain in presence of 0.1µg/ml of voriconazole (Supplementary Figure 6) and
observed a 2-fold change in MIC of voriconazole only in Δ*afu-182* strain but not in WT
strain (Figures 5A and 5B). Thus, it is possible that during the long-term infection low
dose interactions leads to *afu-182* mediated fungal adaptation, and this adaptation
eventually leads to increase in MIC above clinical breakpoints (Figure 9).

Drug concentration at the site of infection is a major indicator of successful
therapy (57). Crucially, the inherent rise of drug resistance in 43% of isolates could not
be attributed to changes in the *cyp51A* sequence (56), indicating non-*cyp51*
mechanisms at play. It is known that continuous exposure to low level of drug leads to
resistance whereas transient exposure to high concentration of drugs lead to tolerance
(58, 59). How fungus adapts to anti-fungal stress before genetic changes associated
with resistance occur is still unknown. Many studies have shown differences in serum
level vs tissue level concentrations of anti-fungal drugs (57). Posaconazole has been
shown to accumulate in lipid rich membrane of alveolar cells in high concentration in

healthy subjects (60); however, the lesion concentration might still be low. Also, during

infection tissue necrosis may limit interaction of fungal cells and posaconazole rich cell

membrane thus limiting lesion concentrations (61). Thus, drug treatment challenges

may lead to low dose exposure in the fungal lesions in patients with IPA even when

serum concentrations are high, leading to lower *afu-182* RNA levels, stress adaptation

and rise in inherent drug resistance in AF during infection (Figure 9).

Different isolates of *A. fumigatus* have been previously shown to differ in

virulence (62). Some of the factors attributed to this difference are nutrient plasticity and

response to light and oxygen (62–64); however, how different azole-susceptible isolates

respond to azole drugs and their virulence potential in relation to azole treatment

requires further research. *afu-182* regulates fungal azole response in ASAF irrespective

of the strain's background, as indicated by deletion of *afu-182* in the laboratory isolates

CEA10 and Af293 backgrounds (Figures 2 and 3D-E) (37). Additionally, deletion of *afu-*

182 in the environmental isolate 47-57 also leads to differential fungal response to sub-

MIC concentrations of azoles (Figures 3D and 3F).

The emergence of new strains provides a unique challenge in studying

Aspergillus pathogenicity, but *afu-182* appears to play a role irrespective of the strain's

background. One possible reason for this might be the mechanisms by which ncRNA

functions. LncRNA provide stress adaptation in the absence of evolution that is

associated with genomic changes (65). A rapid adaptation will increase the likelihood of

fungal survival during stress, leading to DNA changes or the accumulation of beneficial

mutations. A recent study identified 1089 novel long non-coding RNAs; eleven of which

negatively regulate fungal response to azole drugs; however, *afu-182* fell below the size

threshold identified in this study (66). How *afu-182* regulates fungal response is not yet

clear but it is independent of *cyp51* regulation at the transcriptional level

(Supplementary Figures 4A and 4B). Also, mRNA levels of *srbA* and *atrR*, the TFs that

regulate *cyp51* did not change in Δ*afu-182* strain (Supplementary Figures 4C and 4D)

indicating either the effect in post-transcriptional, downstream of these TFs or

completely independent of these known mechanisms.

As long non-coding RNAs function in diverse ways, we used a transcriptomic

approach to identify the genes directly or indirectly regulated by *afu-182* at the RNA

levels. In the presence of azoles, we see the RTA1 family of proteins upregulated in WT

(Figure 6). This is a 7-transmembrane domain family of proteins and has been

associated with lipid and heme export and resistance to 7-amonicholesterol in *S.*

cerevisiae and *Cryptococcus neoformans* (42, 67) and consists of 23 members in *A.*

fumigatus. Recently, *rtaA* (Afu3g12830, CEA10 homolog - *afub_036340*) was shown to

play a role in response to the polyene class of drug, amphotericin B and not azoles (45).

Here, we identified *rtmA* and *rtmB,* two divergent genes downstream of *afu-182*,

situated next to each other to regulate fungal azole response (Figure 8). Both *rtmA* and

rtmB are negatively regulated by *afu-182* (Figures 7); however, *rtmB* mutations might be

the primary contributor to the azole response phenotype, as the double deletion strains

show nearly the same inhibition as Δ*rtmB* (Figures 8D and 8F). As both these genes are

next to each other, it is not clear whether *afu-182*, by itself or in coordination with other

complex members, if any, regulates the promoter region of these genes. It is also

possible that these genes evolved due to a duplication event to maintain tight regulation

for stress adaptation. If *afu-182* regulates fungal azole response via direct regulation of

RTM/RTA1 domain-containing proteins is an active and ongoing area of research.

Importantly, *rtmA* and *rtmB* belong to the accessory genome of AF and were detected in

only 7 of the 300 AF genomes previously analyzed (68). Thus it is possible that the

phenotypes observed in Δ*rtmA* and Δ*rtmB* are correlative with Δ*afu-182* phenotypes as

these genes are absent in Af293 strains of AF (68) but we still see the Δ*afu-182*

mediated sub-MIC phenotype in Af293 background (Figures 3D and 3E). It is also

possible that in absence of *rtmA* and *rtmB* genes, other RTA1 family proteins play a

similar role in Af293. More research will uncover the function of this family of proteins in

AF in the future.

Long non-coding RNAs mainly function through their secondary structure and are

difficult to predict based on the nucleotide sequence alone (69). Here, using synteny, we

identified *afu-182* homolog in related fungal species *A. nidulans*. Deletion of *afu-182*

homolog, *an-182,* also lead to differential sub-MIC azole response in *A. nidulans,*

showing its role in multiple Aspergilli (Supplementary Figure 3). Whether *afu-182*

homologs play a role in the phytopathogens *A. flavus* and *A. parasiticus* is unknown and

needs more research; complex approaches are needed to identify the homologs of *afu-*

182 in other fungal pathogens. However, our data clearly show that long non-coding

RNA, *afu-182,* regulates fungal adaptation in presence of sub-MIC azole concentration

in AF and AN irrespective of the strain's background.

In conclusion, the lncRNA *afu-182* regulates fungal response to azole drugs in both

ASAF and the related fungal species AN. This adaptation is beneficial for AF to gain

azole resistance. This effect, led to identification of *rtmA* and *rtmB* genes in RTA1 family

of protein that regulate fungal azole response and whose transcription is negatively

regulated by *afu-182* in CEA10 strain. As MIC is a poor predictor of azole treatment

outcomes in ASAF strains, further characterization of *afu-182* levels will help us

determine the fungal response to azole drugs *in vivo* that might contribute to inherent

rise in azole drug resistance during azole treatment. Given advancements in RNA

biology, *afu-182* also presents a target to augment the efficacy of current azole drugs.

Materials and Methods

Fungal strains

Aspergillus fumigatus (AF) strain CEA10 or CEA17 was used as the host strain to

generate subsequent strains. For *A. nidulans*, A1145 (FGSC) and GR5 (FGSC) were

used as the host strains to generate subsequent strains. For *A. nidulans*, the strains

were supplemented with Uridine (1.26g/L), Uracil (0.56g/L), pyridoxine (0.05μg/ml), p-

amino-benzoic acid (PABA) (5μM) and riboflavin (0.1μg/ml). All experiments with

stationary fungal growth were performed at 37°C in the presence of 5% CO_2. Liquid

biomass and germination assay were performed in the absence of supplied CO_2. All

strains were stored as 30% glycerol stocks in -70°C for long-term storage. The list of

strains used and generated is listed in Supplementary Table 1. All primers and crRNAs

used in this study are listed in supplementary tables 2 and 3 respectively.

Δ*insA* strain construction: The *insA* deletion strain was created using homologous

recombination. Fusion PCR was used to create the DNA deletion cassette. Briefly, the 5'

and 3' fragments of *insA* were amplified with primers RAC 815 and RAC 816, and

RAC819 and RAC820, respectively. A gene encoding for *pyrG* was amplified from

pJW24 (70) using primers RAC817 and RAC818. The three fragments were fused

together using primers RAC 821 and RAC 822. The resulting DNA fragment was

transformed into the CEA17 strain via PPEG-mediated transformation of protoplast (71).

Strains were checked with PCR, Southern blot for single integration and qPCR for lack

of detectable RNA. This strain does express *afu-182* at the native locus.

Complementation plasmid construction: A plasmid encoding for *pyrithiamine*

resistance, BS309, was created as follows. Two oligo fragments (SD37 and SD38)

encoding for 5' SacI overhang-AscI-BglII-MluI-Pac1-SacII-3' overhang were ordered

from IDT and were allowed to anneal in the thermocycler. The resulting dsDNA fragment

with SacI-SacII overhangs was ligated into pBS KS (+) plasmid previously digested with

SacI and SacII, resulting in plasmid BS309 intermediate. A DNA fragment encoding for

pyrithiamine resistance was amplified from pSD51 with primers RAC3280 and

RAC3154 expressing an XhoI restriction site (72). The DNA fragment was digested with

XhoI and ligated into BS309 intermediate previously digested with SalI, resulting in the

plasmid BS309.

Plasmid construction for Comp-*insA* with cis acting *afu-182*: A DNA fragment

consisting of ~1kb 5' region of *insA* along with *insA* and 485bp after the transcription

end site (as determined by RACE) expressing AscI and PacI restriction sites at the 5'

and 3' ends, respectively, was amplified with primers RAC3589 and RAC3591 using

Takara Prime Star HS polymerase per manufacturer's recommendation. The resulting

DNA fragment was digested with AscI-Pac1 and was ligated into BS309, resulting in

comp-cis *insA* plasmid.

Plasmid Construction for Comp-*insA* without cis-acting *afu-182*: A DNA fragment

consisting of 5' region of *insA* along with *insA* and 74bp after the transcription end site

(without ncRNA *afu-182*) expressing AscI and PacI restriction sites at 5' and 3' ends

respectively was amplified with primers RAC3589 and RAC3590 using Takara Prime

Star HS polymerase per manufacturer's recommendation. The resulting DNA fragment

was digested with AscI-Pac1 and was ligated into BS309, resulting in comp-without cis-
182 *insA* plasmid.

The resulting plasmids were transformed into the Δ*insA* strain using PEG-mediated
transformation of protoplast as previously described (71).

Δ*afu-182* strain construction: *afu-182* was deleted from the strain CEA17 (*pyrG*-)
using CRISPR-Cas9 as previously described (73). A homology-directed repair cassette
was amplified from plasmid pSD38.1 using primers SD149_182 homology F and
SD150_182 homology. Briefly, 5μl of 100mM crRNAs (on either side of *afu-182*) were
combined with 5μl of 100mM tracrRNA and 5 μl of nuclease free duplex buffer (IDT),
heated to 95°C for 5 min and allowed to cool to room temperature, yielding 33mM gRNA
complex. Then, 1.5 μl of each guide RNA complex on either side of *afu-182* were
combined with 1.5μl of Cas9 (1mg/ml) and 22μl of Cas9 working buffer as previously
described (73). The samples were allowed to combine at RT for 5 min to allow the
assembly of the RNP complex. RNP complex and repair cassette were transformed via
PEG-mediated transformation of the CEA17 strain as previously described (71, 73). The
same gRNAs and HDR were also used to delete *afu-182* in the Af293.1 strain and 47-7
strain (62). The sequence of crRNAs and primers used is provided in Tables S2 and S3.

Comp-182 strain construction: *afu-182* was put back at its genomic location using
CRISPR-Cas9. Briefly, the *afu-182* fragment was amplified with primers SD76 and 164
using CEA10 gDNA as template. A gene encoding for pyrithiamine resistance was
amplified from plasmid pSD51.1 (72) using primers SD23 and SD165. Both fragments
were fused using primers SD183 and SD166, generating the HDR cassette. A single
gRNA (gRNA 13 SD 1) targeting the linker sequence in the *pyrG* gene was used for
double-strand break, and HDR was used to insert *afu-182* back at the locus.

OE *afu-182*pyrG plasmid construction: The *A. nidulans gpdA* promoter was amplified
from strain FGSC A4 (FGSC) using primers SD134 and SD103. The *afu-182* fragment
was amplified from CEA10 gDNA using primers SD104 and SD105. Subsequently,

these fragments were fused using primers SD135_EcoRI and SD139_NotI and digested

with EcoRI and NotI. The resulting fragment ligated into plasmid pSD21 previously

digested with EcoR1 and Not1, yielding plasmid OE*afu-182*pyrG. The *afu-182* RNA

expression starts at the +1 transcription start site (TSS).

CEA10 OE *afu-182* strain construction: CRISPR-Cas9 was used to target the OE *afu-*

182 pyrG HDR cassette to the safe-haven *atf4* locus (74) as previously described (64).

The HDR was amplified from plasmid OE *afu-182pyrG* using primers

SD162_OE_homology-F and SD163_OE_homology-R. gRNA was assembled as

described above for the Δ*afu-182* strain. A single guide (atf4_gRNA7) was used to

initiate a double-stranded break targeting the *atf4* locus (74).

Polyethylene glycol-mediated fungal transformation was done as described above. All

resulting transformants were checked with PCR and confirmed with Southern blot and

qPCR for gene expression.

Aspergillus nidulans Δ*182* (Δ*an-182*) **construction:** A DNA sequence homologous to

afu-182 was identified 240bp after the translational stop site of the *insA* homolog gene

AN4465 (32). The resulting 200bp sequence was deleted from the *A. nidulans* genome

using CRISPR-Cas9 in the A1145 strain using the *pyrG* marker or the GR5 strain using

the *pyroA* marker as previously described (73). The HDR cassette for A1145 was

amplified using primers SD181 and SD182 that amplified the sequence of orotidine 5'-

phosphate decarboxylase (75) gene from *A. parasiticus* using plasmid pSD38.1 (76) as

template. The HDR cassette for GR5 was amplified with SD 352 and SD 353, which

amplified the pyridoxine biosynthesis gene (77) from *A. fumigatus* using CEA10 as the

DNA template.

Δ***rtmA*, Δ*rtmB* and Δ*rtmA*Δ*rtmB* strains construction:** Δ*rtmA* and Δ*rtmB* strains were

constructed using the CRISPR-Cas9 protocol as previously described for the Δ*afu-182*

strain (64). The HDR cassette primers and crRNA sequences are provided in Table S1.

Δ*rtmA* and Δ*rtmB* strains were constructed in the CEA10 background using the

pyrithiamine resistance marker, *ptrA* from pSD51.1. For Δ*rtmA*, crRNA 50L and 50R

were used, for Δ*rtmB*, crRNA 60L and 60R were used. For double deletion strains,

crRNAs 60R and 50R were used. HDRs were amplified using primers SD151 and

SD152 for Δ*rtmA* deletion; SD153 and SD154 for Δ*rtmB* deletion and primers SD152

and SD153 for double deletion.

Complement *rtmA* and complement *rtmB* strain construction: Hygromycin

resistance gene expressing plasmid: Hyg resistance gene was amplified using the

primers RAC2939 and RAC2940 from the plasmid pBC Hygro (FGSC) and was ligated

into pJET1.2 (Thermo Fisher). The fragment was digested using XhoI-XbaI and was

ligated into the plasmid BS309 (see above), previously digested with XhoI and XbaI,

resulting in plasmid BS311.

Plasmid construction for Comp *rtmA* strain: The *rtmA* fragment, along with the

promoter and terminator region, was amplified (CEA10 gDNA template) using the

primers SD738 and SD739 with Asc1 and Pac1 restriction sites, respectively, using NEB

Q5 hot start polymerase per manufacturer's recommendation. The resulting DNA

fragment was digested with AscI and PacI and was ligated into the plasmid BS311

previously digested with AscI and PacI, resulting in the Comp-*rtmA* plasmid.

Plasmid construction for comp-*rtmB* strain: The *rtmB* fragment, along with the

promoter and terminator regions, was amplified (CEA10 gDNA template) using the

primers SD740 and SD741 with Asc1 and Pac1 sites, respectively, using NEB Q5 hot

start polymerase per manufacturer's recommendation. The resulting DNA fragment was

digested with AscI and PacI and was ligated into BS311 previously digested with AscI

and PacI, resulting in the Comp-*rtmB* plasmid.

Comp-*rtmA* and comp-*rtmB* plasmids were transformed into Δ*rtmA* and Δ*rtmB* strains,

respectively, as described above.

Spot assays: Plates containing solid GMM were spot inoculated with serially diluted AF

spores (10^5-10^2 spores) and grown at 37°C as indicated. Anti-fungal drugs (azoles,

amphotericinB and Caspofungin) were used at the indicated concentrations. The colony

diameter was measured and reported as the growth ratio, dividing the colony diameter

in the presence of azole by the colony radial diameter in GMM. All experiments were

done at least in triplicates and were repeated three times. Statistical analyses (t-test or

one-way ANOVA) were done to determine the significant differences using GraphPad

Prism.

Biomass assay: Spores were grown in 50 ml liquid GMM (1 x 10^6 spores/ml) at 37°C

for 24 hours in the presence or absence of 0.2 µg/ml voriconazole with continuous

shaking at 250rpm. Mycelium collected after 24 hours was lyophilized, and the biomass

was weighed. Statistical difference between means was assessed with One-way

ANOVA followed by Tukey's post hoc analysis to compare individual groups using

GraphPad Prism.

Biofilm formation assay: AF spore suspension (1.5 x 10^5 spores/ml) in liquid GMM

was inoculated in three biological replicates in a 12-well plate. The plates were

centrifuged at 250g for 10 min to settle the conidia at the bottom of the wells and

incubated at 37°C for 12 hours. Then, the wells were washed twice with PBS to remove

planktonic cells. The biofilms at the bottom of the wells were further allowed to grow in

the presence of 0.2 µg/ml Voriconazole for 8 hours before the crystal violet assay was

performed as previously described (36). Absorbance was measured using a plate

reader (Multiscan Sky, Thermo scientific) at 600 nm. All experiments were done in

triplicate, and three independent experiments were performed. Statistical significance

was determined with One-way ANOVA using GraphPad Prism.

E-test assay: Five milliliters of GMM top-agar (0.75%) containing AF spores (1x 10^5/5

ml) were spread on GMM plates and allowed to solidify. A voriconazole E-test strip

(Biomerieux) was carefully laid in the center of the plate per the manufacturer's

recommendation. The plates were incubated at 37°C for 48 hours, and the minimum

inhibitory concentration (MIC) was recorded. The experiment was done in triplicate for

each strain and repeated three times.

Broth Microdilution assay: AF spores collected from a 48-hour-old colony and (5 x

10^4) were inoculated in liquid GMM in a 96-well plate containing voriconazole (16µg/ml

– 0.015µg/ml) and incubated at 37°C for 48 hours. The visible growth of AF strains was

observed in each of the wells, and the lowest concentration of azoles showing no

growth was recorded as MIC. The experiment was repeated three times.

Murine model of pulmonary aspergillosis, lung histopathology, and fungal

burden: Outbred female CD-1 mice (Charles River Laboratories) with body weights of

18-21 g were housed (5 per cage) in micro-isolator cages. The mice were kept in a soft-

lit, well-ventilated cubicle with a 12h:12h light/dark cycle. Mice were immunosuppressed

with a single subcutaneous injection of 40 mg/kg body weight of Triamcinolone

acetonide (Kenalog-10, Bristol-Myers Squibb) on day -1. Mice were briefly anesthetized

with isoflurane and infected by intranasal instillation of 2 x 10^6 spores in 40µL PBS. Ten

mice were infected per AF strain for survival analysis, whereas five mice per strain were

used for fungal burden analysis. The mice were observed for 14 days for their survival.

For azole treatment, 5 mg/kg Posaconazole (Noxafil, oral suspension, Merck) diluted in

PBS was used.

For fungal burden analyses, mice were treated with 5 mg/kg Posaconazole on day +1

and day +2, and their lungs were harvested on day +3. The mouse lungs were

lyophilized overnight, homogenized with 2.3 zirconia beads, and lysed with LETS buffer

(0.1M LiCl, 10mM EDTA pH 8.0, 10mM Tris-Cl pH 8.0, 0.5% SDS). Total DNA was

extracted in an aqueous phase using phenol: chloroform: isoamyl alcohol and

precipitated with isopropanol. Fungal burden was determined by calculating the levels of

18s rRNA encoding gene against a standard curve in 500ng total DNA using iTaq

universal probe mix and hydrolysis probe as previously described (40). The statistical

significance of survival analysis was determined using log-rank analysis, and fungal

burden analysis was done using a non-parametric Mann-Whitney U Test.

For histological analysis, lungs were harvested from untreated mice on day +3 and were

fixed in 10% formalin for 48 hours before microdissection at the Clemson Light Imaging

Facility (CLIF) at Clemson University. Adjacent slides obtained from microdissection

were stained with Grocott's Methenamine Silver and Hematoxylin and Eosin stains as

previously described (41). The stained slides were pictured with the Leica Laser

Microdissection Microscopy System at CLIF.

Voriconazole adaptation: Azole adaptation experiment was carried out for CEA10 and

Δafu-182 strains by growing the strains in 0.1 µg/ml voriconazole for 12 generations at

37°C. Briefly, CEA10 and Δafu-182 strains were spot inoculated out of glycerol stock

(0G) in glucose minimal media (GMM) plate containing 0.1 µg/ml voriconazole at 37°C

for 48 hours. Spores ($1G^V$) were collected in 0.01% Tween-80, and again spot

inoculated onto fresh GMM plate containing 0.1 µg/ml voriconazole for 48 hours. The

remaining $1G^V$ spores were stored as glycerol stock for further experiments in -70°C.

This process was repeated for 12 generations.

RNA extraction: For RNA sequencing, total RNA was extracted from AF strains CEA10,

Δafu-182, comp-182, and OE-182 strains. Briefly, 1×10^4 spores were spot inoculated

and incubated at 37°C for 48 hours on solid GMM plates in the absence or presence of

voriconazole (0.15 µg/ml). The whole colonies from samples were excised with a clean

scalpel and flash frozen in liquid N_2. The frozen samples were homogenized with 2.3

mm zirconia beads in acid guanidinium thiocyanate phenol, and RNA was extracted

using the acid guanidinium thiocyanate phenol chloroform (AGPC) method as described

with some modifications (78). Briefly, the aqueous phase was mixed with ethanol to a

final concentration of 35%. The resulting mix was passed through RNA-binding columns

(Epoch Life Sciences), washed twice with 70% Ethanol and eluted in nuclease-free

water. The RNA was treated with Turbo DNase (Thermo Fisher) per the manufacturer's

recommendation and purified using magnetic beads (Beckman Coulter). RNA quantity

and quality were estimated using the AccuBlue Broad Range RNA Quantitation Kit

(Biotium) and the Quantum Fluorometer (Promega), and via gel electrophoresis,

respectively, before submission of the samples to LC Sciences.

Sample quality was checked by Agilent Technologies 2100 Bioanalyzer at LC Sciences.

The poly(A) RNA sequencing library was prepared following Illumina's TruSeq-stranded-

mRNA sample preparation protocol. Poly(A) tail-containing mRNAs were purified using

oligo-(dT) magnetic beads with two rounds of purification. After purification, poly(A) RNA

was fragmented using a divalent cation buffer at elevated temperature. Quality control

analysis and quantification of the sequencing library were performed using Agilent

Technologies 2100 Bioanalyzer High Sensitivity DNA Chip. Paired-end sequencing was

performed on Illumina's NovaSeq 6000 sequencing system.

RNA-seq analyses: Transcript expression was determined by first aligning RNA-Seq

reads to the *A. fumigatus* A1163 genome obtained from FungiDB v68 (32) with the

splice-aware aligner STAR v2.7.11a (79). The resulting SAM file for each experimental

replicate was processed with samtools v1.19.2 (80) and featureCount v2.0.6 (81) to

produce a table of read counts per annotated gene locus. The read counts were

analyzed with DESeq v1.46.0 (82) and plotted with EnhancedVolcano v1.24.0 (83) in

the R v4.4.2 statistical environment(84). The pipeline for analyzing these is archived in a

GitHub repository (https://github.com/stajichlab/DhingraLab_ncRNA_Afumigatus) and

Zenodo (85)

Assessment of gene expression by RT-qPCR: For qPCR, 48-hour-old AF colonies

grown at 37°C in solid GMM plates with or without voriconazole (0.15µg/ml) were

extracted as described above. RNA samples were then treated with Turbo DNase per

the manufacturer's recommendations. Equal amounts of DNase-treated RNA were used

for first-strand cDNA synthesis using MMLV reverse transcriptase (Promega) per the

manufacturer's recommendation. Quantitative PCR (CFX real-time PCR system - Bio-

Rad) was done using the sso Advanced universal SYBR green supermix (Bio-Rad).

Histone H4 was used as the reference gene for normalization. All primers used in the

study are listed in Supplementary Table 2.

Data Availability: The RNA-seq sequencing data and per-gene read counts are

available from Sequence Read Archive associated to BioProject PRJNA1334547 and in

Gene Expression Omnibus under accession GSE309441.

Statistical analyses: GraphPad Prism 10 for macOS was used to perform all statistical

analyses.

References:

1. Morrissey CO, Kim HY, Duong T-MN, Moran E, Alastruey-Izquierdo A, Denning DW,

Perfect JR, Nucci M, Chakrabarti A, Rickerts V, Chiller TM, Wahyuningsih R,

Hamers RL, Cassini A, Gigante V, Sati H, Alffenaar J-W, Beardsley J. 2024.

Aspergillus fumigatus—a systematic review to inform the World Health

Organization priority list of fungal pathogens. Med Mycol 62:myad129.

2. Baddley JW. 2011. Clinical risk factors for invasive aspergillosis. Med Mycol 49

Suppl 1:S7–S12.

3. Latgé J-P, Chamilos G. 2019. Aspergillus fumigatus and Aspergillosis in 2019. Clin

Microbiol Rev 33:e00140-18.

4. Liu K-W, Grau MS, Jones JT, Wang X, Vesely EM, James MR, Gutierrez-Perez C,

Cramer RA, Obar JJ. 2022. Postinfluenza Environment Reduces Aspergillus

fumigatus Conidium Clearance and Facilitates Invasive Aspergillosis In Vivo. mBio

13:e02854-22.

5. Sarden N, Sinha S, Potts KG, Pernet E, Hiroki CH, Hassanabad MF, Nguyen AP,

Lou Y, Farias R, Winston BW, Bromley A, Snarr BD, Zucoloto AZ, Andonegui G,

Muruve DA, McDonald B, Sheppard DC, Mahoney DJ, Divangahi M, Rosin N,

Biernaskie J, Yipp BG. 2022. A B1a–natural IgG–neutrophil axis is impaired in viral-

and steroid-associated aspergillosis. Science Translational Medicine 14:eabq6682.

6. Tio SY, Chen SC-A, Hamilton K, Heath CH, Pradhan A, Morris AJ, Korman TM,

Morrissey O, Halliday CL, Kidd S, Spelman T, Brell N, McMullan B, Clark JE,

Mitsakos K, Hardiman RP, Williams P, Campbell AJ, Beardsley J, Van Hal S, Yong

MK, Worth LJ, Slavin MA. 2023. Invasive aspergillosis in adult patients in Australia

and New Zealand: 2017–2020. The Lancet Regional Health - Western Pacific

40:100888.

7. WHO fungal priority pathogens list to guide research, development and public

health action. https://www.who.int/publications/i/item/9789240060241. Retrieved 19

September 2025.

8. Patterson TF, Thompson GR, Denning DW, Fishman JA, Hadley S, Herbrecht R,

Kontoyiannis DP, Marr KA, Morrison VA, Nguyen MH, Segal BH, Steinbach WJ,

Stevens DA, Walsh TJ, Wingard JR, Young J-AH, Bennett JE. 2016. Practice

Guidelines for the Diagnosis and Management of Aspergillosis: 2016 Update by the

Infectious Diseases Society of America. Clin Infect Dis 63:e1–e60.

9. Verweij PE, Lucas JA, Arendrup MC, Bowyer P, Brinkmann AJF, Denning DW, Dyer

PS, Fisher MC, Geenen PL, Gisi U, Hermann D, Hoogendijk A, Kiers E, Lagrou K,

Melchers WJG, Rhodes J, Rietveld AG, Schoustra SE, Stenzel K, Zwaan BJ,

Fraaije BA. 2020. The one health problem of azole resistance in *Aspergillus*

fumigatus: current insights and future research agenda. Fungal Biology Reviews

34:202–214.

10. Lee Y, Robbins N, Cowen LE. 2023. Molecular mechanisms governing antifungal

drug resistance. npj Antimicrob Resist 1:5.

11. Warrilow AGS, Parker JE, Price CL, Nes WD, Kelly SL, Kelly DE. 2015. In Vitro

Biochemical Study of CYP51-Mediated Azole Resistance in Aspergillus fumigatus.

Antimicrobial Agents and Chemotherapy 59:7771–7778.

12. Verweij PE, Lucas JA, Arendrup MC, Bowyer P, Brinkmann AJF, Denning DW, Dyer

PS, Fisher MC, Geenen PL, Gisi U, Hermann D, Hoogendijk A, Kiers E, Lagrou K,

Melchers WJG, Rhodes J, Rietveld AG, Schoustra SE, Stenzel K, Zwaan BJ,

Fraaije BA. 2020. The one health problem of azole resistance in Aspergillus

fumigatus: current insights and future research agenda. Fungal Biology Reviews

34:202–214.

13. Mellado E, Garcia-Effron G, Alcázar-Fuoli L, Melchers WJG, Verweij PE, Cuenca-

Estrella M, Rodríguez-Tudela JL. 2007. A new Aspergillus fumigatus resistance

mechanism conferring in vitro cross-resistance to azole antifungals involves a

combination of cyp51A alterations. Antimicrob Agents Chemother 51:1897–1904.

14. Chowdhary A, Sharma C, Meis JF. 2017. Azole-Resistant Aspergillosis:

Epidemiology, Molecular Mechanisms, and Treatment. J Infect Dis 216:S436–

S444.

15. Gsaller F, Hortschansky P, Furukawa T, Carr PD, Rash B, Capilla J, Müller C,

Bracher F, Bowyer P, Haas H, Brakhage AA, Bromley MJ. 2016. Sterol

Biosynthesis and Azole Tolerance Is Governed by the Opposing Actions of SrbA

and the CCAAT Binding Complex. PLOS Pathogens 12:e1005775.

16. Blosser SJ, Cramer RA. 2012. SREBP-Dependent Triazole Susceptibility in

Aspergillus fumigatus Is Mediated through Direct Transcriptional Regulation of

erg11A (cyp51A). Antimicrob Agents Chemother 56:248–257.

17. Blatzer M, Barker BM, Willger SD, Beckmann N, Blosser SJ, Cornish EJ, Mazurie

A, Grahl N, Haas H, Cramer RA. 2011. SREBP coordinates iron and ergosterol

homeostasis to mediate triazole drug and hypoxia responses in the human fungal

pathogen Aspergillus fumigatus. PLoS Genet 7:e1002374.

18. Paul S, Stamnes M, Thomas GH, Liu H, Hagiwara D, Gomi K, Filler SG, Moye-

Rowley WS. 2019. AtrR Is an Essential Determinant of Azole Resistance in

Aspergillus fumigatus. mBio 10:10.1128/mbio.02563-18.

19. Willger SD, Puttikamonkul S, Kim K-H, Burritt JB, Grahl N, Metzler LJ, Barbuch R,

Bard M, Lawrence CB, Jr RAC. 2008. A Sterol-Regulatory Element Binding Protein

Is Required for Cell Polarity, Hypoxia Adaptation, Azole Drug Resistance, and

Virulence in Aspergillus fumigatus. PLOS Pathogens 4:e1000200.

20. Yan R, Cao P, Song W, Li Y, Wang T, Qian H, Yan C, Yan N. 2021. Structural basis

for sterol sensing by Scap and Insig. Cell Rep 35:109299.

21. Butler G. 2013. Hypoxia and Gene Expression in Eukaryotic Microbes. Annu Rev

Microbiol 67:291–312.

22. Sever N, Yang T, Brown MS, Goldstein JL, DeBose-Boyd RA. 2003. Accelerated

degradation of HMG CoA reductase mediated by binding of insig-1 to its sterol-

sensing domain. Mol Cell 11:25–33.

23. Rybak JM, Ge W, Wiederhold NP, Parker JE, Kelly SL, Rogers PD, Fortwendel JR.

2019. Mutations in hmg1, Challenging the Paradigm of Clinical Triazole Resistance

in Aspergillus fumigatus. mBio 10:10.1128/mbio.00437-19.

24. Hagiwara D, Arai T, Takahashi H, Kusuya Y, Watanabe A, Kamei K. 2018. Non-

cyp51A Azole-Resistant Aspergillus fumigatus Isolates with Mutation in HMG-CoA

Reductase. Emerg Infect Dis 24:1889–1897.

25. Gonzalez-Jimenez I, Lucio J, Roldan A, Alcazar-Fuoli L, Mellado E. 2021. Are Point

Mutations in HMG-CoA Reductases (Hmg1 and Hmg2) a Step towards Azole

Resistance in Aspergillus fumigatus? Molecules 26:5975.

26. Souza ACO, Ge W, Wiederhold NP, Rybak JM, Fortwendel JR, Rogers PD. 2023.

hapE and hmg1 Mutations Are Drivers of cyp51A-Independent Pan-Triazole

Resistance in an Aspergillus fumigatus Clinical Isolate. Microbiology Spectrum

11:e05188-22.

27. Rybak JM, Xie J, Martin-Vicente A, Guruceaga X, Thorn HI, Nywening AV, Ge W,

Souza ACO, Shetty AC, McCracken C, Bruno VM, Parker JE, Kelly SL, Snell HM,

Cuomo CA, Rogers PD, Fortwendel JR. 2024. A secondary mechanism of action

for triazole antifungals in Aspergillus fumigatus mediated by hmg1. Nat Commun

15:3642.

28. Berman J, Krysan DJ. 2020. Drug resistance and tolerance in fungi. Nat Rev

Microbiol 18:319–331.

29. Scott J, Valero C, Mato-López Á, Donaldson IJ, Roldán A, Chown H, Rhijn NV,

Lobo-Vega R, Gago S, Furukawa T, Morogovsky A, Ami RB, Bowyer P, Osherov N,

Fontaine T, Goldman GH, Mellado E, Bromley M, Amich J. 2022. Aspergillus

fumigatus can display persistence to the fungicidal drug voriconazole. bioRxiv

https://doi.org/10.1101/2022.05.16.491816.

30. Scott J, Valero C, Mato-López Á, Donaldson IJ, Roldán A, Chown H, Rhijn NV,

Lobo-Vega R, Gago S, Furukawa T, Morogovsky A, Ami RB, Bowyer P, Osherov N,

Fontaine T, Goldman GH, Mellado E, Bromley M, Amich J. 2023. Aspergillus

fumigatus Can Display Persistence to the Fungicidal Drug Voriconazole.

Microbiology Spectrum https://doi.org/10.1128/spectrum.04770-22.

31. Amich J, Bromley M, Goldman GH, Valero C. 2025. Toward the consensus of

definitions for the phenomena of antifungal tolerance and persistence in

filamentous fungi. mBio 16:e03475-24.

32. Basenko EY, Shanmugasundram A, Böhme U, Starns D, Wilkinson PA, Davison

HR, Crouch K, Maslen G, Harb OS, Amos B, McDowell MA, Kissinger JC, Roos

DS, Jones A. 2024. What is new in FungiDB: a web-based bioinformatics platform

for omics-scale data analysis for fungal and oomycete species. Genetics

227:iyae035.

33. Jöchl C, Rederstorff M, Hertel J, Stadler PF, Hofacker IL, Schrettl M, Haas H,

Hüttenhofer A. 2008. Small ncRNA transcriptome analysis from Aspergillus

fumigatus suggests a novel mechanism for regulation of protein synthesis. Nucleic

Acids Res 36:2677–2689.

34. Scotto–Lavino E, Du G, Frohman MA. 2006. 3′ End cDNA amplification using
classic RACE. Nat Protoc 1:2742–2745.

35. Kang Y-J, Yang D-C, Kong L, Hou M, Meng Y-Q, Wei L, Gao G. 2017. CPC2: a fast
and accurate coding potential calculator based on sequence intrinsic features.
Nucleic Acids Res 45:W12–W16.

36. Snarr BD, Baker P, Bamford NC, Sato Y, Liu H, Lehoux M, Gravelat FN, Ostapska
H, Baistrocchi SR, Cerone RP, Filler EE, Parsek MR, Filler SG, Howell PL,
Sheppard DC. 2017. Microbial glycoside hydrolases as antibiofilm agents with
cross-kingdom activity. Proc Natl Acad Sci U S A 114:7124–7129.

37. Bertuzzi M, van Rhijn N, Krappmann S, Bowyer P, Bromley MJ, Bignell EM. 2020.
On the lineage of Aspergillus fumigatus isolates in common laboratory use. Med
Mycol 59:7–13.

38. Keller NP. 2017. Heterogeneity Confounds Establishment of "a" Model Microbial
Strain. mBio 8:10.1128/mbio.00135-17.

39. Dhingra S, Kowalski CH, Thammahong A, Beattie SR, Bultman KM, Cramer RA.
2016. RbdB, a Rhomboid Protease Critical for SREBP Activation and Virulence in
Aspergillus fumigatus. mSphere 1:e00035-16.

40. Dhingra S, Buckey JC, Cramer RA. 2018. Hyperbaric Oxygen Reduces Aspergillus
fumigatus Proliferation In Vitro and Influences In Vivo Disease Outcomes.
Antimicrob Agents Chemother 62:e01953-17.

41. Graybill JR, Najvar LK, Gonzalez GM, Hernandez S, Bocanegra R. 2003.

Improving the mouse model for studying the efficacy of voriconazole. J Antimicrob

Chemother 51:1373–1376.

42. Manente M, Ghislain M. 2009. The lipid-translocating exporter family and

membrane phospholipid homeostasis in yeast. FEMS Yeast Res 9:673–687.

43. Hokken MWJ, Zoll J, Coolen JPM, Zwaan BJ, Verweij PE, Melchers WJG. 2019.

Phenotypic plasticity and the evolution of azole resistance in Aspergillus fumigatus;

an expression profile of clinical isolates upon exposure to itraconazole. BMC

Genomics 20:28.

44. Ness F, Aigle M. 1995. Rtm1: A Member of a New Family of Telomeric Repeated

Genes in Yeast. Genetics 140:945–956.

45. Abou-Kandil A, Tröger-Görler S, Pschibul A, Krüger T, Rosin M, Schmidt F,

Akbarimoghaddam P, Sarkar A, Cseresnyés Z, Shadkchan Y, Heinekamp T, Gräler

MH, Barber AE, Walther G, Figge MT, Brakhage AA, Osherov N, Kniemeyer O.

2025. The proteomic response of Aspergillus fumigatus to amphotericin B (AmB)

reveals the involvement of the RTA-like protein RtaA in AmB resistance. microLife

6:uqae024.

46. Verweij PE, Chowdhary A, Melchers WJG, Meis JF. 2016. Azole Resistance in

Aspergillus fumigatus: Can We Retain the Clinical Use of Mold-Active Antifungal

Azoles? Clin Infect Dis 62:362–368.

47. Cuypers L, Aerts R, Van de gaer O, Vinken L, Merckx R, Gerils V, Vande Velde G,

Reséndiz-Sharpe A, Maertens J, Lagrou K. 2025. Doubling of triazole resistance

rates in invasive aspergillosis over a 10-year period, Belgium, 1 April 2022 to 31

March 2023. Euro Surveill 30:2400559.

48. Kang Y, Ma W, Li Q, Wang P, Jia W. 2025. Epidemiology, antifungal susceptibility

and biological characteristics of clinical Aspergillus fumigatus in a tertiary hospital.

Sci Rep 15:16906.

49. Burks C, Darby A, Londoño LG, Momany M, Brewer MT. 2021. Azole-resistant

Aspergillus fumigatus in the environment: Identifying key reservoirs and hotspots of

antifungal resistance. PLOS Pathogens 17:e1009711.

50. Liu S, Le Mauff F, Sheppard DC, Zhang S. 2022. Filamentous fungal biofilms:

Conserved and unique aspects of extracellular matrix composition, mechanisms of

drug resistance and regulatory networks in Aspergillus fumigatus. npj Biofilms

Microbiomes 8:83.

51. Kowalski CH, Morelli KA, Schultz D, Nadell CD, Cramer RA. 2020. Fungal biofilm

architecture produces hypoxic microenvironments that drive antifungal resistance.

Proceedings of the National Academy of Sciences 117:22473–22483.

52. Kowalski CH, Morelli KA, Stajich JE, Nadell CD, Cramer RA. 2021. A

Heterogeneously Expressed Gene Family Modulates the Biofilm Architecture and

Hypoxic Growth of Aspergillus fumigatus. mBio 12:e03579-20.

53.   Wuyts J, Van Dijck P, Holtappels M. 2018. Fungal persister cells: The basis for
recalcitrant infections? PLoS Pathog 14:e1007301.

54.   De Francesco MA. 2023. Drug-Resistant Aspergillus spp.: A Literature Review of Its
Resistance Mechanisms and Its Prevalence in Europe. Pathogens 12:1305.

55.   Dagenais TRT, Keller NP. 2009. Pathogenesis of Aspergillus fumigatus in Invasive
Aspergillosis. Clin Microbiol Rev 22:447–465.

56.   Bueid A, Howard SJ, Moore CB, Richardson MD, Harrison E, Bowyer P, Denning
DW. 2010. Azole antifungal resistance in Aspergillus fumigatus: 2008 and 2009. J
Antimicrob Chemother 65:2116–2118.

57.   Zhao Y, Prideaux B, Baistrocchi S, Sheppard DC, Perlin DS. 2019. Beyond tissue
concentrations: antifungal penetration at the site of infection. Med Mycol 57:S161–
S167.

58.   Moreillon P, Tomasz A. 1988. Penicillin resistance and defective lysis in clinical
isolates of pneumococci: evidence for two kinds of antibiotic pressure operating in
the clinical environment. J Infect Dis 157:1150–1157.

59.   Windels EM, Michiels JE, Van den Bergh B, Fauvart M, Michiels J. 2019.
Antibiotics: Combatting Tolerance To Stop Resistance. mBio 10:e02095-19.

60.   Conte JE, Golden JA, Krishna G, McIver M, Little E, Zurlinden E. 2009.
Intrapulmonary Pharmacokinetics and Pharmacodynamics of Posaconazole at
Steady State in Healthy Subjects. Antimicrob Agents Chemother 53:703–707.

61. Campoli P, Al Abdallah Q, Robitaille R, Solis NV, Fielhaber JA, Kristof AS,

Laverdiere M, Filler SG, Sheppard DC. 2011. Concentration of Antifungal Agents

within Host Cell Membranes: a New Paradigm Governing the Efficacy of

Prophylaxis ▯. Antimicrob Agents Chemother 55:5732–5739.

62. Kowalski CH, Beattie SR, Fuller KK, McGurk EA, Tang Y-W, Hohl TM, Obar JJ,

Cramer RA. 2016. Heterogeneity among Isolates Reveals that Fitness in Low

Oxygen Correlates with Aspergillus fumigatus Virulence. mBio

7:10.1128/mbio.01515-16.

63. Caffrey-Carr AK, Kowalski CH, Beattie SR, Blaseg NA, Upshaw CR, Thammahong

A, Lust HE, Tang Y-W, Hohl TM, Cramer RA, Obar JJ. 2017. Interleukin 1α Is

Critical for Resistance against Highly Virulent Aspergillus fumigatus Isolates. Infect

Immun 85:e00661-17.

64. Fuller KK, Cramer RA, Zegans ME, Dunlap JC, Loros JJ. 2016. Aspergillus

fumigatus Photobiology Illuminates the Marked Heterogeneity between Isolates.

mBio 7:e01517-16.

65. Traubenik S, Charon C, Blein T. 2024. From environmental responses to

adaptation: the roles of plant lncRNAs. Plant Physiol 195:232–244.

66. Bowyer P, Weaver D, Qi T, Chown H, Fraczek M, Lebedinec R, Dineen L, Valero C,

Rhijn N van, Furukawa T, Bromley M, Delneri D. 2025. Genome-wide discovery

and phenotyping of non-coding transcripts in A. fumigatus reveals lncRNAs with a

role in antifungal drug sensitivity. Research Square https://doi.org/10.21203/rs.3.rs-

6048166/v1.

67. Smith-Peavler ES, Patel R, Onumajuru AM, Bowring BG, Miller JL, Brunel JM,

Djordjevic JT, Prabu MM, McClelland EE. 2022. RTA1 Is Involved in Resistance to

7-Aminocholesterol and Secretion of Fungal Proteins in Cryptococcus neoformans.

Pathogens 11:1239.

68. Barber AE, Sae-Ong T, Kang K, Seelbinder B, Li J, Walther G, Panagiotou G,

Kurzai O. 2021. Aspergillus fumigatus pan-genome analysis identifies genetic

variants associated with human infection. Nat Microbiol 6:1526–1536.

69. Mercer TR, Dinger ME, Mattick JS. 2009. Long non-coding RNAs: insights into

functions. Nat Rev Genet 10:155–159.

70. Thammahong A, Caffrey-Card AK, Dhingra S, Obar JJ, Cramer RA. 2017.

Aspergillus fumigatus Trehalose-Regulatory Subunit Homolog Moonlights To

Mediate Cell Wall Homeostasis through Modulation of Chitin Synthase Activity.

mBio 8:e00056-17.

71. Szewczyk E, Nayak T, Oakley CE, Edgerton H, Xiong Y, Taheri-Talesh N, Osmani

SA, Oakley BR. 2006. Fusion PCR and gene targeting in Aspergillus nidulans. Nat

Protoc 1:3111–3120.

72. Thammahong A, Dhingra S, Bultman KM, Kerkaert JD, Cramer RA. 2019. An Ssd1

Homolog Impacts Trehalose and Chitin Biosynthesis and Contributes to Virulence

in Aspergillus fumigatus. mSphere 4:e00244-19.

989 73. Al Abdallah Q, Ge W, Fortwendel JR. 2017. A Simple and Universal System for

990   Gene Manipulation in Aspergillus fumigatus: In Vitro-Assembled Cas9-Guide RNA

991   Ribonucleoproteins Coupled with Microhomology Repair Templates. mSphere

992   2:e00446-17.

993 74. Furukawa T, van Rhijn N, Chown H, Rhodes J, Alfuraji N, Fortune-Grant R, Bignell

994   E, Fisher MC, Bromley M. 2022. Exploring a novel genomic safe-haven site in the

995   human pathogenic mould Aspergillus fumigatus. Fungal Genet Biol 161:103702.

996 75. Oakley BR, Rinehart JE, Mitchell BL, Oakley CE, Carmona C, Gray GL, May GS.

997   1987. Cloning, mapping and molecular analysis of the pyrG (orotidine-5'-phosphate

998   decarboxylase) gene of Aspergillus nidulans. Gene 61:385–399.

999 76. Dhingra S, Lind AL, Lin H-C, Tang Y, Rokas A, Calvo AM. 2013. The fumagillin

1000   gene cluster, an example of hundreds of genes under veA control in Aspergillus

1001   fumigatus. PLoS One 8:e77147.

1002 77. Osmani AH, May GS, Osmani SA. 1999. The extremely conserved pyroA gene of

1003   Aspergillus nidulans is required for pyridoxine synthesis and is required indirectly

1004   for resistance to photosensitizers. J Biol Chem 274:23565–23569.

1005 78. Zepeda B, Verdonk JC. 2022. RNA Extraction from Plant Tissue with Homemade

1006   Acid Guanidinium Thiocyanate Phenol Chloroform (AGPC). Current Protocols

1007   2:e351.

79.   STAR: ultrafast universal RNA-seq aligner | Bioinformatics | Oxford Academic.

https://academic.oup.com/bioinformatics/article/29/1/15/272537. Retrieved 9

September 2025.

80. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,

Keane T, McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and

BCFtools. Gigascience 10:giab008.

81. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. Bioinformatics 30:923–

930.

82. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

83. Blighe K, Rana S, Lewis M. EnhancedVolcano: Publication-ready volcano plots with

enhanced colouring and labeling

https://doi.org/10.18129/B9.bioc.EnhancedVolcano.

84.   R Core Team (2024). R: A Language and Environment for Statistical Computing

(version 3.5.1). R Foundation for Statistical Computing. https://www.R-project.org/.

Retrieved 9 September 2025.

85. Stajich J. 2025. stajichlab/DhingraLab_ncRNA_Afumigatus: Archive of Analysis and

Data archive for publication (v0.1). Zenodo.

1027

Acknowledgements:

We would like to acknowledge members of Dhingra laboratory and Eukaryotic Pathogen innovation center for discussions surrounding the experiments in the manuscript. The research in Dhingra lab is supported by start-up funds from the Department of Biological Science at Clemson University and National Institute of General Medical Sciences (NIGMS) award (P20GM146584) (PI - James Morris). We thank Dr. Robert Cramer at the Dartmouth Geisel School of Medicine (NIAID/NIH grants R01AI181215 and 2R0AI130128) for providing resources and critical reading of the manuscript. Analyses were performed on the UC Riverside High Performance Computing Cluster supported by grants from the National Science Foundation (DBI-1429826 & DBI-2215705) and the National Institutes of Health (NIH) (S10-OD016290). JES is a CIFAR fellow in the program Fungal Kingdom: Threats and Opportunities and supported by NIH grant R01AI130128.0. The Clemson University Genomics and Bioinformatics Facility is supported by the College of Science and Grants P20GM146584 and P20GM139769 Institutional Development Awards (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health. This work was supported, in part, by equipment housed in the Clemson Light Imaging Facility. We utilized a Leica Laser Microdissection system for brightfield imaging, and a Leica Pearl Automatic Tissue Processor and a Leica Histocore Biocut Rotary Microtome for histology preparation. Thank you to Mr. Avery Herren for assistance with histology. The Clemson Light Imaging Facility is supported by the Division of Research and EPIC COBRE P20GM146584. We thank Dr. Jarrod Fortwendel for the Fdel262 strain, Anne Strong for technical help and Dr. James Morris for the critical reading of the manuscript. All animal experiments were approved by Institutional Animal Care and Use Committee (IACUC approval # AUP2022-0236). Garret Kaufman was supported by MENTOR: Medical Enrichment Through Opportunities in Research Grant (1T35AI134643-01A1, PI – Dr. Kerry Smith). We acknowledge the support of veterinary and technical staff at the Godley-Snell Research Center for help with the animal studies. The funding sources had no role in the study design, data collection and interpretation, preparation of this manuscript or the decision to submit the manuscript.

Figure1:

Figure 1. *insA* and *afu-182* are independent transcripts. Schematic showing the genomic location of *afu-182* with respect to *insA* gene. The 3' end of *insA* was determined via 3' RACE. The inset sequence shows the sequencing data matching the end of *insA* transcript.

Figure 2:

Figure 2. *afu-182* regulates sub-MIC fungal response to azoles. (A) AF strains were serially inoculated ($10^5 - 10^2$) on GMM in the absence and presence of azole drugs as indicated. The Δ*afu-182* strain showed increased fungal growth in the presence of the sub-MIC concentration of voriconazole, whereas the OE-182 strain is more inhibited compared to the WT strain. (B-E) Growth ratio. Colony diameters were measured for colonies as grown in A (for 10^4 spore density) and represented as diametric growth ratio (Drug/no drug). (F) *afu-182* plays a role in fungal biofilm response to azole drugs. AF strains were allowed to adhere to cell culture plates, and biofilms were treated with voriconazole and stained with crystal violet. The OD$_{600}$ values are represented. Δ*afu-182* formed azole recalcitrant

biofilms (p=0.0049, One-Way ANOVA), whereas OE-182 showed a significant decrease in biofilm formation (p=0.0011, One-Way ANOVA) when treated with voriconazole. (G) AF strains were grown in liquid submerged culture with continuous shaking in the absence or presence of 0.2µg/ml voriconazole. After 48 hours, fungal biomass was collected and lyophilized. Dry weight ratio (Drug/no drug) is represented. One-way ANOVA was used to assess the differences in mean; p-values are listed for all samples compared to WT (One-Way ANOVA followed by Tukey post-hoc analysis). a.u. – arbitrary units.

Figure 3:

Figure 3. *afu-182* RNA levels are negatively regulated by azoles and correlate with MIC.
(A) Quantitative reverse transcriptase PCR was used to determine the *afu-182* RNA levels in the WT strain grown in the absence and presence of azoles. Significant downregulation of *afu-182* was observed in the presence of voriconazole (p=0.0019, One-Way ANOVA) and posaconazole (p=0.0088, One-Way ANOVA). (B) Quantitative reverse transcriptase PCR was used to determine the *afu-182* RNA levels in Af293, 47-57, 47-7, -46 and -10 isolates compared to CEA10. (C) Minimum inhibitory concentration. Voriconazole MIC as measured by broth microdilution method for strains in B. (D) Growth ratio. *afu-182* was deleted from laboratory reference strain Af293.1 and environmental isolate 47-57. WT and corresponding deletion strains were serially inoculated on GMM in the absence and presence of voriconazole. Growth ratio represented as the ratio of colony diameter in the

presence of drug over no drug is represented. Significantly more growth was observed in *Δafu-182* strains for Af293 and 47-57 strains (two-tailed t-test, p =0.0038 and 0.0019, respectively). AF strains were serially inoculated ($10^5 – 10^2$) on GMM in the absence and presence of voriconazole as indicated. The *Δafu-182* strain showed increased fungal growth in the presence of the sub-MIC concentration of voriconazole.

Figure 4:

Figure 4. *afu-182* plays a role in azole treatment outcomes but is indispensable for virulence in a corticosteroid murine model of IPA. (A) Schematic showing experimental details for animal infection. (B) Kaplan-Meier plot of animal survival probability for animals infected as in A. No significant difference was observed (p=0.1166 WT vs. Δ*afu-182*, log-rank test). (C) Schematic showing experimental details for animal infection and azole treatment. (D) Kaplan-Meier plot of animal survival probability for animals infected as in C. A significantly higher mortality (90% vs 55%) was observed in animals infected with the Δ*afu-182* strain compared to animals infected with the WT strain (p=0.0103, log-rank test) upon azole treatment. No difference was observed in the mice infected with the OE-182 strain compared to WT (40% mortality vs 50% mortality, p=0.4065). (E) Schematic showing experimental details for infection and treatment. (F) Mice were infected and treated as in F.

Fungal DNA was quantified as described in Methods. Significantly more fungal DNA was recovered from animals infected with the Δ*afu-182* strains (p = 0.0159, Mann-Whitney U-test), and significantly less DNA was recovered from animals infected with the OE-182 strain (p = 0.0159, Mann-Whitney U-test) after posaconazole treatment. No difference was observed in fungal DNA isolated from animals infected with WT or comp *afu-182* strains (p=0.6905).

Figure 5:

Figure 5. Continuous exposure to low level azoles leads to azole drug resistance in an afu-182 dependent manner. (A) Broth microdilution was used to determine the voriconazole MIC for the un-adapted and adapted strains (G12^V – 12 defines the generation and V defines adaptation in presence of voriconazole). Only the Δ*afu-182* strain exposed to low level voriconazole show increase in MIC. (B) E-test strip was used to assess the MIC of voriconazole of the adapted strains. Like broth microdilution, E-Test showed increase in MIC in Δ*afu-182* adapted strains. MIC values represented at the bottom as µg/ml. (C) Serially diluted spores (10^5-10^2) were spot inoculated on GMM or GMM + voriconazole (0.25µg/ml) or (0.5µg/ml) and grown for 48 hours at 37°C. The WT adapted strain showed increased fungal growth in the presence of the sub-MIC concentration of voriconazole (orange arrow), whereas the Δ*afu-182* adapted strain showed increased fungal growth at 0.5µg/ml (red arrows).

Figure 6:

Figure 6 . Genes encoding RTA1 domain proteins are upregulated in response to voriconazole. Quantitative reverse transcription PCR was used to determine the mRNA levels of RTA1 genes in the presence and absence of azoles. Out of 23, we selected 12 genes that have an FPKM value >1. Eleven of the 12 genes showed significant upregulation in the presence of azoles (p<0.05, two-tailed Student's t-test). Importantly, *rtaA,* which plays a role in polyene response but not azole response, showed the lowest upregulation (1.6-fold, p=0.0204, two-tailed student's t-test). Numbers in orange represent the FPKM values from the RNAseq data set.

Figure 7:

Figure 7. *afu-182* negatively regulates genes encoding for RTM1 proteins. Volcano plots showing the 2-log$_2$ fold change (4-fold upregulated or downregulated). (A) Comparison between CEA10 azole and Δ*afu-182* azole. Global transcriptomic analysis revealed genes encoding RTM1 domain proteins *rtmA* (*afub_000050*) and *rtmB* (*afub_000060*) are

significantly upregulated (orange arrows). (B) Comparison between CEA10 and Δ*afu-182* in GMM. No significant change in *rtmA* and *rtmB* level was observed. Quantitative reverse transcriptase PCR was used to assess the mRNA levels of (C) *rtmA* and (D) *rtmB*. Significant upregulation of *rtmA* (3-fold compared to WT, p=0.0005, Two-Way ANOVA followed by Tukey's post-hoc analysis) and *rtmB* (4.6-fold compared to WT, p=0.0015, Two-Way ANOVA followed by Tukey's post-hoc analysis) was observed in Δ*afu-182* strains compared to WT in the presence of voriconazole. No significant difference in *rtmA* and *rtmB* expression was observed in the absence of azoles.

Figure 8:

Figure 8. *rtmA* and *rtmB* play a role in fungal azole response. AF strains (A) Δ*rtmA* and comp-*rtmA*, (C) Δ*rtmB* and comp-*rtmB* and (E) Δ*rtmA* Δ*rtmB* were point inoculated in the absence or presence of 0.25 µg/ml of voriconazole. The plates were incubated at 37°C for 48 hours and photographed. The inset shows images taken with a macro lens. After 48 hours, colony diameter was measured and growth ratio of drug over no drug for (B) Δ*rtmA* and comp-*rtmA*, (E) Δ*rtmB* and comp-*rtmB* and (H) Δ*rtmA*Δ*rtmB* (three independent

deletion strains) were plotted. One-way ANOVA was used to assess the difference in the mean. Numerical p-values are represented based on Tukey's post-hoc analysis to compare the means of all groups.

Figure 9:

Figure 9. AF responds to sub-MIC azole drugs by regulating *afu-182* levels resulting in azole drugs resistance. (A) Sub-MIC azole exposure lowers the *afu-182* levels. (B) This adaptation affects the clinical outcomes in an animal model of IPA treated with azole drugs. (C) Continuous exposure to low level drugs that is possible during infection or environmental use leads to increase in MIC. (D) The mechanistic pathway controlling *afu-182* mediated resistance are unknown and under investigation; however, appear to be independent of Cyp51 transcriptional regulation. (E) Acute exposure of supra-MIC drugs is linked to tolerance and persistence, and it is currently unknown if that plays a role in drug resistance or (F) Spontaneous mutations can lead to azole drug resistance.