Lactic Acid Bacteria Dominate Urban Bokashi: A culture-independent study of Microbial Diversity and Functional Potential in Household-Scale Food Waste Fermentation

Katharina Kujala[1,2], Veera Kinnunen[3,4]

[1] Natural Resources Institute Finland, Oulu, Finland

[2] Water, Energy and Environmental Engineering Research Unit, University of Oulu, Oulu, Finland

[3] Faculty of Education and Psychology, University of Oulu, Oulu, Finland

[4] Faculty of Social Sciences, University of Lapland, Rovaniemi, Finland

Corresponding author:

Katharina Kujala

Natural Resources Institute Finland (Luke)

Paavo Havaksen tie 3

FI-90570 Oulu, Finland

Katharina.kujala@luke.fi

Phone: +358 29 5322772

Keywords: Bokashi compost, lactic acid bacteria, urban biodiversity, citizen science, One Health

Abstract

In recent years, concerns over declining (microbial) biodiversity in urban spaces have increased. Urban Bokashi composting (i.e., the microaerobic or anaerobic fermentation of food waste indoors on the household scale) has been suggested as a possibility to promote microbial diversity in the domestic environment. However, studies on microbial communities in household-scale Bokashi and their potential impacts on health and environment are lacking. Thus, the present study investigated microbial communities in different stages of the Bokashi composting process in collaboration with Bokashi practitioners by looking into physicochemical characteristics as well as microbial community composition (16S amplicon sequencing) and functional potential (shotgun metagenome sequencing). The collective results indicate that i) microbial communities in Bokashi compost differed between stages, but also between households, ii) microbial communities were dominated by lactic acid bacteria like *Lentilactobacillus* or *Lacticaseibacillus*, iii) metabolic pathways for the production of diverse organic acids were detected, iv) application of Bokashi ferment or leachate to soil can supply nutrients and organic acids to promote plant growth but does not substantially affect soil microbial community composition, and v) potentially pathogenic organisms were detected in extremely low abundances. Thus, urban Bokashi is likely not associated with increased health risks and positive impacts are feasible.

Introduction

In recent years, there has been growing concern over the declining microbial biodiversity of urban spaces and the subsequent disconnect between humans and natural environments (Flandroy *et al.* 2018). In the course of urbanization, the original green spaces have largely been converted into "grey spaces" (United Nations 2004). In many places, natural soil has been replaced by synthetic, impervious materials such as concrete, asphalt or rubber, which has effectively sealed surfaces and impacted thermal and hydrological characteristics of urban spaces. As there is a continuous exchange in microbiota between humans and their environment, low-biodiversity environments can negatively impact the biodiversity of human microbiomes (Hanski *et al.* 2012). Lack of biodiversity in the environment and the human microbiome can in turn have negative effects on the human immune system and lead to a higher prevalence of allergies and autoimmune disorders (Hanski *et al.* 2012; Haahtela 2019). To address this issue, attempts have been made to enhance urban biodiversity by creating green

spaces in urban areas such as green walls or roofs and urban gardens. A more recent, promising and increasingly explored option to promote urban microbial biodiversity is the incorporation of soil, rocks, humus and organic matter in public spaces such as playgrounds or parks (Sun *et al.* 2023; Manninen *et al.* 2025). Even enrichment of materials like sand with microbial communities from organic soils has been considered to promote contact of humans with microbial biodiversity (Hui *et al.* 2019).

In addition to larger-scale measures, citizen-driven initiatives can diversify microbial communities and increase exposure to microorganisms in their living spaces such as their homes or their garden. One such practice that has become more popular in recent years in urban settings is Bokashi. Bokashi originates from an East Asian tradition of treating organic waste such as food waste by fermenting it and using the ferment as a soil amendment (Park and DuPonte 2008; Christel 2017). It can be used on the household scale (e.g., (Lew *et al.* 2021)) or in larger settings (Formowitz *et al.* 2007). In contrast to conventional composting, household-scale Bokashi composting is generally practiced indoors in urban settings and can thus lead to a higher exposure of the Bokashi practitioner to microbial communities during the composting process. On the household scale, food waste is collected over the course of several weeks and subsequently fermented microaerobically or anaerobically in closed containers (Kinnunen 2021). To facilitate the fermentation process, the material is typically inoculated with activated Bokashi starter bran which has been prepared with enriched fermenting microorganisms or directly with a solution of fermenting microorganisms. One source of enriched fermenters are the commercially available effective microorganisms (EM®). EM® is a microbial inoculum consisting of about 80 microbial species that was developed in Japan by Teruo Higa (Higa 1991), and is a trademark licensed by EMRO. EM® is used for the preparation of many of the available commercial starter brans. While the detailed composition of EM is kept secret, some of the main species include lactic acid bacteria like *Lactiplantibacillus plantarum, Lacticaseibacillus casei* and *Streptococcus lactis*, photosynthetic purple non-sulfur bacteria like *Rhodopseudomonas palustris* and *Rhodobacter sphaeroides* as well as yeasts like *Saccharomyces cerevisiae* (Xu 2001). Activated Bokashi bran is prepared by mixing enriched fermenting microbial consortia such as EM with wheat or rice bran, molasses and water. This activated bran is subsequently mixed with food waste for fermentation. Alternatively, ready-made Bokashi bran (i.e., prefermented and dried) can be purchased from different suppliers. After a minimum of two weeks of fermentation, the fermented food waste is mixed with soil to complete the process. In addition, leachate is

collected from the fermenting Bokashi containers and used for different purposes such as fertilizer (often strongly diluted) or cleaning agent (e.g., to unclog pipes; (Kinnunen 2021)).

Bokashi composting has been surrounded by bold claims for its environmental benefits, such as improving climate smartness and microbial diversity (Fan *et al.* 2018). Although Bokashi practitioners are reporting positive experiences, independent, scientific research demonstrating the environmental impact of the method is still scarce and partly contradictory (see e.g., (Thorslund 2020; Vicente *et al.* 2020; Lew *et al.* 2021; Safwat and Matta 2021)). Despite these inconsistent results, there is a shared experiential knowledge among the Bokashi practicing community that using Bokashi as a soil amendment results in a healthy soil by improving soil porosity and water-holding capacity as well as by addition of nutrients, which in turn results in healthier, bigger crops (e.g., (Murillo-Amador *et al.* 2015; Christel 2017; Quiroz and Céspedes 2019)). In addition, bokashi composting has proven to have positive effect especially in odor control and humification (Fan *et al.* 2018). Moreover, vernacular waste fermentation practices, such as bokashi, are also incorporated into urban everyday life in order to remediate toxins and to enrich domestic microbiota and thus enhance wellbeing of ecosystems and inhabitants (Zhang 2021). These holistic health claims are in line with the popularized One Health paradigm which seeks to envision health of people, animals, and environments through the same lens (Evans and Leighton 2014).

While larger-scale nature-based solutions and their impact on health, environment and microbial communities have been studied to some extent, studies on microbial communities in household-scale Bokashi and their potential impacts on health and environment are lacking. Thus, this study set out to characterize microbial communities in Bokashi from several urban practitioners along with collecting information about their experiences with Bokashi composting. Bokashi practitioners were invited to participate in study design and sample collection. The main goals of the study were to i) identify points-of-interest of the Bokashi practitioners and adjust the conducted analyses accordingly, ii) characterize the microbial community composition in different stages of the Bokashi process, iii) investigate the process potential of actively fermenting Bokashi, iv) identify similarities and differences in Bokashi communities from different households, v) visualize the microbial communities in Bokashi in an easily graspable way for their practitioners, and vi) outline potential health benefits and risks associated with household-scale Bokashi. To address these goals, Bokashi practitioners provided samples from different stages of their Bokashi process, which were analyzed for microbial community composition and other characteristics. Moreover, participants provided

background information on their Bokashi and were able to shape the project focus and data visualization in pre- and post-analysis workshops (Figure 1).

Figure 1: Overview of the project workflow. Samples were collected by six Bokashi practitioners, leachate characteristics and microbial community composition were analyzed and visualized for sharing with the practitioners. The results were discussed with the practitioners to give them insights on the microbial communities in their Bokashi compost and answer questions they brought forward. Icons used to construct this workflow overview were modified from bioicons.com (Agilent_HPLC icon by Sasha Sundstrom, ph-meter and nanopore_sequencing icons by DBCLS, cuvette-pinkdark and microtube-closed-translucent icons by Servier, bubble icon by dbs-sticky, column-stacked-100, heatmap and tree-map icons by dbs-sticky, DNA icon by Kumar) and a graphic designed by Freepick ("People business in office").

Material and methods

Recruitment of Bokashi practitioners

The participants for the study were recruited by circulating an open invitation to a face-to-face Bokashi workshop organized at the University of Oulu in May 2023. The invitation was written both in English and in Finnish and it was circulated in two largest social media groups dedicated to Bokashi making ("Bokashi Suomi" and "Bokashi kompostointi (riippumaton)"), on mailing

lists of Oulu-based environmental associations as well as mailing lists for University of Oulu researchers. The invitation was addressed to households in the Oulu region which were already practicing Bokashi-composting. The pre-experiment workshop was open for all interested Bokashi practitioners, from which volunteer households were to be selected to participate in sample collection from their Bokashi. Despite a number of people expressing interest in the study, only five Bokashi practitioners attended the face-to-face workshop, so all of them were selected to provide samples for the study. In the beginning of the workshop, the participants were informed about the aims of the study and their rights as research participants, emphasizing that the Bokashi samples will be handled anonymously without any personal identifiers. After the first workshop, two participants withdrew from the study, so two additional Bokashi households were recruited for sample collection through direct contacts.

Questions brought forward by and interaction with Bokashi practitioners

In the pre-experiment workshop, potential participation in the project was discussed with interested Bokashi practitioners. After giving some initial information about the project, participants were organized into two small groups to discuss what questions intrigued them about Bokashi composting and what issues about Bokashi they would like to have addressed in the experimental phase. The researchers facilitated the discussion and took notes. Practical questions of participants were related to the fermentation process ("What do the microbes produce?", "What is the pH of the Bokashi/leachate?", "Is it possible to ferment green waste with EM-liquid without molasses?", "Do additions like biochar or egg shells affect the process?"), to the use of ferment or leachate ("Does leachate need to be diluted before being used as fertilizer?", "What type of soil can be used?", "Is acidity a problem later on?") and to health implications ("What will stay on my hands if I mix Bokashi/soil without gloves?", "Are there any proven health benefits?").

It transpired that Bokashi practitioners wished a rather complete picture of the whole process (from starter to soil) but were especially interested in the analysis of the activated Bokashi starter bran, which is often a commercial product. Based on the participant input and the results of a preliminary study (see next section), four different stages of the Bokashi composting process were selected to be sampled by the Bokashi practitioners (activated Bokashi starter bran, actively fermenting Bokashi, leachate, soil factory).

Citizen sampling of Bokashi

Two rounds of sampling were conducted. In a preliminary study in 2022, one household (= household 1) sampled actively fermenting Bokashi in different stages (start, after 1 week, after 3 weeks) as well as Bokashi leachate (after 1 week). Based on the outcome of the preliminary study, which showed no major differences between the different stages of actively fermenting Bokashi, and on the input from Bokashi practitioners during the workshop, the protocol was amended to include the following fractions: 1) activated Bokashi starter bran that was used as a Bokashi starter, 2) actively fermenting Bokashi (after 2-5 weeks), 3) Bokashi leachate sampled at the same time, 4) soil collected from the soil factory (i.e., fermented Bokashi mixed with soil; after 2-7 weeks). Samples were taken by the Bokashi practitioners into clean containers (such as glasses or plastic jars) and frozen in a home freezer. Some practitioners provided several samples for stage 2) (5 households) and stage 4) (2 households), either from different fermentation batches or as replicate from the same batch. Samples were then collected by the project team or delivered to the laboratory by the Bokashi practitioners and stored at -20°C until further processing. In addition, Bokashi practitioners were asked to provide information about their Bokashi system (e.g., type of food waste that was composted, composting bin type, soil used for soil factory, how long which stage was run).

Analytical methods

Bokashi leachates were filtered (0.2 µm). The leachate pH was analyzed used a handheld pH meter (WTW Sentix). Organic acids were quantified on an Agilent 1200 HPLC system equipped with an Aminex HPX-87H column (300 x 7.8 mm; Biorad) and a UV-visible light diode array detector (detection at 210 nm) using a mobile phase of 4 mM H_3PO_4, a column temperature of 60°C and a flow rate of 0.6 mL/min. Organic acid standards with known concentrations of formate, acetate, lactate, propionate and butyrate were used for calibration. Ammonium and phosphate concentrations were determined using a YSI9300 photometer (YSI Incorporated, USA) and the kit/protocols Phot 62 (ammonium) and Phot 28 (phosphate). Nitrate concentrations were determined colorimetrically (Miranda, Espey and Wink 2001).

DNA extraction, Nanopore 16S amplicon and metagenomic sequencing

DNA was extracted from all samples and a negative extraction control using the ZymoBiomics DNA Miniprep Kit according to the manufacturer's instructions. Solid materials were used directly, while leachates were centrifuged for 10 minutes at 14,000 *g* to pellet solid materials (2 mL) and the supernatant was discarded. 16S rRNA genes were amplified using the primer set 515F (5'-GTGYCAGCMGCCGCGGTAA; (Parada, Needham and Fuhrman 2016)) - 926R

(5'-CCGYCAATTYMTTTRAGTTT; (Quince *et al.* 2011)). The primers were preceded by adapters (5'-TTTCTGTTGGTGCTGATATTGC-515F, 5'-ACTTGCCTGTCGCTCTATCTTC-926R) to allow for the attachment of barcodes in a second round of PCR using the Oxford Nanopore Technologies (ONT) 96 PCR barcoding extension (EXP-PBC096). The first round of PCR consisted of initial denaturation (5 mins at 95°C), followed by 35 cycles (45 sec at 95°C, 45 sec at 50°C, 90 sec at 68°C) and a final elongation (5 mins at 68°C), while the second round of PCR consisted of initial denaturation (5 mins at 95°C), followed by 15 cycles (45 sec at 95°C, 45 sec at 62°C, 90 sec at 68°C) and a final elongation (5 mins at 68°C). After the second PCR, PCR products were pooled and purified using 1x AmpureXP magnetic beads (Beckman Coulter) according to the manufacturer's instructions. Purified PCR products were end-prepped and sequencing adapters were ligated according to the ONT protocol. Amplicons were sequenced on a MinION using a Flongle R10.4.1 flow cell as given in the ONT protocol.

Shotgun metagenomes were sequenced only from DNA extracts of actively fermenting Bokashi (11 samples). Due to low sample amount, the rapid PCR barcoding kit (SQK-RPB114.24) was chosen. Input DNA was tagmented which generated DNA fragments of on average 3.5kb (Fig. S1) in size flanked by adapters on both ends. Tagmented DNA was then amplified for 14 cycles using the kit's barcoded primers and processed as described in the ONT protocol. Metagenomes were sequenced on a MinION using a R10.4.1 flow cell.

Bioinformatics analysis and representation of the results

Basecalling and demultiplexing of raw POD5 data was conducted with Dorado ([GitHub - nanoporetech/dorado: Oxford Nanopore's Basecaller](#)) using the super accurate basecalling model ([dna_r10.4.1_e8.2_400bps_sup@v5.2.0](#)) for both amplicon and metagenome sequencing reads. Demultiplexed amplicon sequence reads were filtered to remove reads with an average quality score < 7 and imported into qiime2 (Bolyen *et al.* 2019). For each barcode, forward and reverse primers were initially processed separately due to the mixed orientation of Nanopore reads. Forward/reverse primer sequences were identified and trimmed at the 5'-end of the reads and the reverse complements of the reverse/forward primer sequences were identified and trimmed at the 3'-end of the reads using the q2-cutadapt plugin (Martin 2011). Reads in which primers were lacking at one or both ends were discarded. The sequence orientation of the reverse reads was adjusted using the q2-rescript plugin (Robeson *et al.* 2021),

after which forward and reverse reads and OTU tables were merged. Sequences were classified against the greengenes2 database (McDonald *et al.* 2024) using the consensus blast q2-feature-classifier (Camacho *et al.* 2009). Unassigned reads and reads with taxonomy assigned only at domain-level were removed from the OTU table. The OTU table was collapsed at species-level and rarefied to an even sampling depth of 7000 sequences. Alpha (Shannon diversity (Shannon 1948) and Pielou Evenness (Pielou 1966)) parameters were calculated based on the rarefied OTU table using the q2-diversity plugin.

Read quantities and qualities of metagenomic reads were visualized as histograms and density plots using Nanoplot (De Coster and Rademakers 2023) and the "ggplot2" (Wickham 2016) and "ggExtra" (Attali and Baker 2025) packages in R version 4.4.2 (R Core Team 2024). The highest density of reads was observed at ~3.5 kb and an average quality score of 20 in all samples (Fig. S1), while the reads in the negative control where mainly short (~200 bp) and of low quality (< 10). Metagenomic reads where filtered using SeqKit2 (Shen, Sipos and Zhao 2024), removing any reads with an average quality score < 10. *In silico* translated reads were aligned against the NCBI non-redundant protein database using DIAMOND (Buchfink, Reuter and Drost 2021) for taxonomic and functional annotation and visualized in MEGAN6 (Huson *et al.* 2007, 2016). Reads were assembled into contigs using myloasm, a tool optimized for the assembly of long-read sequences (Shaw, Marin and Li 2025), which yielded 13,336 contigs with an average length of 11,376 bp (1026 – 2,708,123 bp). Reads were mapped onto the contigs using bowtie2 (Langmead and Salzberg 2012) with mapping rates ranging from 81 to 95% as part of the anvi'o pipeline v8 (Eren *et al.* 2020) and binned using CONCOCT v1.1.0 (Alneberg *et al.* 2014), Maxbin2 v2.2.7 (Wu, Simmons and Singer 2016) and MetaBAT 2 v2.15 (Kang *et al.* 2019). DAS Tool v1.1.7 (Sieber *et al.* 2018) was applied to integrate the results of the three binning methods to produce a set of non-redundant bins, and bins with >50% completeness and <10% redundancy were retained as MAGs. MAGs were imported into Kbase (Arkin *et al.* 2018) for taxonomic classification with GTDB-Tk v2.3.2 (Chaumeil *et al.* 2022; Chivian *et al.* 2023) and functional annotation with DRAM v0.1.2 (Shaffer *et al.* 2020). Phylogenetic trees from GTDB-Tk were modified in iTOL v6 (Letunic and Bork 2024). DRAM output was screened for functions and pathways involved in degradation of organic substrates, including degradation of polymers and monomers as well as fermentations. MAGs were moreover inspected for the presence of potential antibiotic resistance genes using the Resistance Gene Identifier (RGI) web tool of the Comprehensive Antibiotic Resistance Database (CARD) (Alcock *et al.* 2023). Strict and loose hits were filtered to only retain hits

with identities >30% to known antibiotic resistance genes and grouped based on the drug classes to which they confer resistance. Notably, some genes confer resistance to multiple drug classes and where thus included multiple times. MAGs were classified as high-quality or medium-quality drafts according to the MIMAG criteria (Bowers *et al.* 2018).

To illustrate microbial community diversity and the dominant genera in each stage of the process to the participating Bokashi practitioners, genus-level relative abundances were visualized as Voronoi plots using the "Weighted Treemap" (Jahn *et al.* 2024) package in R. Only genera with a relative abundance greater than 0.01% were included, and the error tolerance for differences between the target and actual cell sizes was set to 5%. The 15 genera with the highest relative abundance were labelled for each plot. Concentrations of organic acids were visualized using the R package "packcircles" (Bedward, Eppstein and Menzel 2024).

Results

Bokashi setup comparison

Activated Bokashi starter brans from three different companies readily available in Finland were used by the Bokashi practitioners. According to the information provided on the suppliers' website and/or the product label, all starters were prepared with wheat bran, sugar molasses, water, and EM. Two of the products mentioned additional ingredients such as spelt husk, sunflower oil, and lime (Table 1). No information was provided about the bran activation and drying process (e.g., conditions, duration). Two of the products mentioned some of the microorganisms contained, which included *Lactobacillus plantarum* (= *Lactiplantibacillus plantarum*), *Lactobacillus casei* (= *Lactocaseibacillus casei*), *Rhodopseudomonas palustris* and *Saccharomyces cerevisiae* in concentrations of $10^4 - 10^5$ cfu g^{-1}. While it was not explicitly stated, it seemed like the suppliers listed the species and concentrations that were applied for the preparation of the activated Bokashi starter bran rather than referring to the final product.

All study participants set up their Bokashi in closed buckets equipped with a stopcock for withdrawal of leachate. In all households, the food waste collected included fruits and vegetables/vegetable peels. In addition, some households reported that they had also added bread (household 2), berries (households 2 and 4), fish (household 3), dairy products (household 3), coffee grounds and tea leaves (household 5), porridge (household 6) and meal leftovers (household 6). However, no guarantee can be given for completeness of information,

as the collection of food waste was conducted over a period of several weeks, and study participants did not provide an exhaustive list of the contents of their Bokashi containers. For the soil factory, study participants used garden soil, sandy soil and leftover soil from previous soil factories.

Table 1: Information on activated Bokashi starter brans provided by the supplier.

Starter bran	Consistency	Recommended use	Ingredients	EM microorganisms mentioned on label
1	dried grits	add 1 tablespoons to surface for 1L food waste	wheat bran, spelt husk, EM, water, sunflower oil, sugar cane molasses, coralline algae lime	*Lactobacillus plantarum, Lactobacillus casei, Rhodopseudomonas palustis, Saccharomyes cerevisiae*
2	dried grits	add 2 tablespoons to surface after each addition of food waste	wheat bran, water, sugar molasses, EM-1®	*Lactobacillus plantarum, Lactobacillus casei, Saccharomyces cerevisiae, Rhodopseudomonas palustris, Rhodospirillum rubrum*
3	dried grits	mix 1-2 tablespoon with 1L food waste	wheat bran, spelt husk, EM, sugar cane molasses, EM ceramic powder, seaweed lime, sunflower oil	not specified

Physicochemical characteristics of Bokashi leachate

All leachates were acidic with pH ranging from 3.7 to 4.9 and contained large quantities of organic acids (Table 2). In most households, lactate was the dominant organic acid produced with concentrations ranging from 14 to 19 mM (Table 2). Only leachate from household 5 had considerably lower lactate concentrations (< 5 mM). Acetate and propionate were present in

similar concentrations in all leachates (9-14 mM and 1.5-2.5 mM, respectively), while butyrate had only accumulated to >5 mM in four of the leachates (households 1, 3, 5, 6). Formate concentrations were rather low in all leachates (< 0.25 mM). Ammonium concentrations in leachate ranged from <10 to 167 µM, while nitrate concentrations ranged from 29 to 860 µM. Phosphate concentrations ranged from 3 to 70 µM.

Table 2: Characteristics of Bokashi
leachate

Household	pH	EC (ms/cm)	Ammonium (µM)	Nitrate (µM)	Phosphate (µM)	Formate (mM)	Acetate (mM)	Propionate (mM)	Butyrate (mM)	Lactate (mM)
1	4.661	13.5	<10	189.5	20.7	0.20	11.8	2.3	5.0	14.3
2	3.778	10.9	11.1	857.6	2.7	n.d.[a]	12.7	1.5	0.37	13.0
3	3.852	15.2	111.1	44.2	69.7	0.05	14.6	2.3	6.3	17.2
4	3.706	13.73	111.1	595.3	6.7	n.d.	13.0	1.8	n.d.	19.4
5	4.909	10.63	166.7	23.3	38.7	0.15	8.9	2.5	7.1	4.6
6	4.335	14.89	138.9	29.1	41.9	0.21	10.3	1.9	5.8	18.5

[a] not detected

Microbial communities during different stages of the Bokashi composting process

16S rRNA gene amplicon sequencing revealed that microbial communities in actively fermenting Bokashi and its leachate were dominated by *Firmicutes* (50-99% relative abundance; Figure 2 A). *Proteobacteria* were the second most abundant group, accounting for up to 30% of the microbial communities in this stage. However, *Proteobacteria* were only abundant in Bokashi and leachate from households 2, 5 and 6. The dominant genera in actively fermenting Bokashi and its leachate differed between the households: While *Lentilactobacillus* was rather abundant (20-60% relative abundance) in samples from most households (apart from household 5), the abundance of other genera was more individual (Figure 2 B). In household 1, *Companilactobacillus* was prominent, in household 3, *Lactobacillus*, in households 2 and 4 *Lactiplantibacillus* and *Levilactobacillus* were prominent and households 2 and 5 had many genera below the threshold abundance of 5% in at least one sample. Microbial community composition in Bokashi starter bran was often similar to the composition in actively fermenting Bokashi of the same household, and *Lentilactobacillus* was the dominant genus in starter brans from supplier 1 and 3. Notably, photosynthetic genera like *Rhodopseudomonas, Rhodobacter*

and *Rhodospirillum* (which are advertised as a beneficial part of EM) were not detected in any of the analyzed activated Bokashi starter brans (at a sequencing depth of >7000 sequences), indicating that they were not present in high numbers in the used starters .

Figure 2: Phylum- (A) and genus-level (B) microbial community composition in different stages of the Bokashi process based on 16S rRNA gene amplicon sequencing. Phyla or genera with a relative abundance < 5% in all individual samples were grouped ("other"). Average relative abundances are presented for cases in which duplicate samples were analyzed (marked with *).

Microbial diversity in soil factories was considerably higher than in the activated Bokashi starter brans, actively fermenting Bokashi or leachate, with a median number of observed species of 1100 (vs. 200-500) and a median Shannon diversity of 8 (vs. 4-5; Figure 3). Microbial communities in soil factories were dominated by *Proteobacteria* (37-60%; Figure 2 A). In households 2 and 3, *Firmicutes* were prominent (33-39% relative abundance), while in the other households they were below 10%. Unlike the actively fermenting Bokashi samples, most soil factories were not strongly dominated by one or two genera but had a more even distribution (Figure 2 B), indicating that soil factory microbial communities were likely not determined by the Bokashi ferment they received but rather by the residing soil microbes.

Figure 3: Species-level microbial diversity in different stages of the Bokashi process based on 16S rRNA gene amplicon sequencing. No. of observed species, Shannon diversity and Pielou Evenness diversity indices were calculated based on the rarefied species table using the q2-diversity plugin after rarefication to an even sampling depth of 7000 sequences.

Metagenomic analysis of actively fermenting Bokashi

Long-read metagenomes were created from eleven samples of actively fermenting Bokashi. Analysis of unassembled reads mostly replicated the prokaryotic relative abundance patterns obtained from 16S rRNA gene amplicon sequencing (Fig. S2). A small percentage of metagenomic reads (0.6 to 1%) was affiliated to *Eukaryota* and viruses (Table S1). Some species like *Saccharomyces cerevisiae* or *Aureobasidium melanogenum* were only detected in one or two samples, while others like *Trichuris trichiura* or *Mortierella* sp. were detected in all samples. The detected viruses were mainly viruses of *Lactobacillus/Lactobacillaceae*. SEED

functional annotation revealed the functional potential for organic matter degradation and fermentation processes (Fig. S3).

Assembly of long-read sequencing data resulted in the recovery of 19 non-redundant medium- to high-quality metagenome-assembled genomes (MAGs; Table S2). Most MAGs were classified as *Lactobacillaceae* (18/19), representing the genera *Lentilactobacillus* (7 MAGs), *Lacticaseibacillus* (2 MAGs), *Lactobacillus* (2 MAGs), *Pediococcus* (2 MAGs), *Levilactobacillus* (1 MAG), *Loigolactobacillus* (1 MAG), *Lactiplantibacillus* (1 MAG), *Companilactobacillus* (1 MAG) and *Schleiferilactobacillus* (1 MAG), while one MAG was classified as *Bifidobacterium* (Fig. 4).

Figure 4: Phylogenomic tree of the 19 metagenome-assembled genomes (MAGs) recovered from actively fermenting Bokashi. MAGs were placed in GTDB species trees for lactic acid bacteria and Bifidobacteria using GTDB-tk Classify in KBase. The tree was imported into iTOL v7 for midpoint rooting and editing. MAGs are indicated by (*) on the branches and bold, blue font.

In each household, actively fermenting Bokashi was typically dominated by one to three MAGs (Fig. 5). The functional potential of MAGs was inspected focusing on pathways and functions important for organic matter degradation as well as for antimicrobial resistance. MAGs encoded pathways for the degradation of glucose as well as enzymes needed for the degradation of complex organic compounds, including beta-glucosidase, beta-galactosidase and arabinosidase. Moreover, fermentative pathways that produce short chain organic acids or

alcohols were also encoded, with potential for lactate, acetate and alcohol production being present in nearly all MAGs, while the potential for butyrate, propionate or formate production was detected in only a few MAGs. Hits to potential antibiotic resistance genes were found in most MAGs, with the highest number of hits for genes conveying resistance to fluoroquinolones, macrolides and tetracyclines (Fig. S4). However, the majority of the hits were classified as "loose" by the CARD search tool with only 24 "strict" hits, which mainly were related to resistance to the glycopeptide antibiotic vancomycin (15), disinfecting agents like benzalkonium chloride (5) and fluoroquinolones like norfloxazin and ciprofloxazin. Loose hits, while showing good sequence similarity to known antimicrobial resistance genes, might not actually encode functional proteins involved in antimicrobial resistance.

Figure 5: Taxonomic identification (A), relative abundance (B) and metabolic potential (C-E) of metagenome-assembled genomes (MAGs) recovered from actively fermenting Bokashi. In (B) averages of duplicates for households 1-3 and 5-6 are shown. Pathways and functions were identified based on DRAM annotations. Multiple variants of organic acid production pathways provided by DRAM were combined in (E). The tree and heatmaps were generated using iTOL v7.

Communicating the results to study participants

For study participants, information cards were generated to illustrate the microbial community composition in different stages and the characteristics of the leachate (Figure 6). These results were shared at a post-experiment workshop, in which the questions brought forward by participants in the previous meeting or on the day were also addressed. Participants were concerned when names of know pathogenic genera appeared in their chart, e.g., *Pseudomonas, Klebsiella* or *Streptococcus*. Therefore, the workshop was used as an opportunity to discuss the diversity of the genera in question, which are metabolically versatile and contain also many non-pathogenic species as well as microbes common in soil and water. Species-level classifications of sequences were not shared with the participants due to potential inaccuracies caused by incomplete coverage of the 16S rRNA genes and potential sequencing errors. The results of the metagenomic analysis were not yet available at the time of the workshop and were thus not discussed. Overall, participants showed great interest in the study results, and some of the questions from the pre-experiment workshop could be answered (e.g., regarding pH or what the Bokashi microbes produce). However, many open questions remained that could not directly be answered with the experimental results and further questions were brought forward on the day during discussions. E.g., Bokashi practitioners questioned whether the addition of activated starter bran was actually needed, or whether inoculation with leachate or ferment from a previous batch would be enough, and brought forward many practical concerns e.g., related to regulations for composting by local authorities. In hindsight, a second post-experiment workshop might have been needed to better connect with the Bokashi practitioners. Many questions (e.g., related to health impacts of Bokashi) could not be addressed directly due to the study design, but answers might in part be found in the scientific literature or might be addressed in follow-up studies.

Figure 6: Example (household 6) of visualization provided for the Bokashi practitioners in the final workshop. (A) Microbial community composition in each stage, illustrated as Voronoi plots in which the polygon area correlates with the relative abundance of the genera. Top 10 genera are labelled in each plot. (B) Leachate characteristics. Organic acid concentrations are visualized in a bubble plot, in which the bubble size corresponds to the measured concentration in µM. Visualizations for all households are given in the supplementary material. Note that the original version of the visualization used during the workshop were prepared in Finnish language.

Discussion

Plant growth-promoting benefits of Bokashi compost application on farm- and household-scale

Bokashi composting for the recycling of household food waste is increasing in popularity and comes with many promises, for many of which there is only anecdotal evidence reported by Bokashi practitioners. Open questions surrounding the practice of Bokashi include its useability as fertilizer, potential health risks through pathogens or potential health benefits through enrichment of beneficial microbes in the domestic ecosystem. While this study is among the first that has studied Bokashi microbial communities on the household scale, more research has been done looking at the use of Bokashi for treatment of various waste materials on a larger scale as well as the use of the final product as an organic soil amendment in agriculture. Bokashi compost has been used as an organic fertilizer e.g., in South America for a variety of crops including bananas, coffee, bell peppers, cucumber, tomatoes, maize and strawberries (Formowitz *et al.* 2007; Mosquera *et al.* 2016; Sarmiento, Amézquita and Mena 2019; Pandit *et al.* 2020; Abo-Sido *et al.* 2021; Mondaca *et al.* 2025; García-Hernández *et al.*). The addition of Bokashi compost increases crop production and soil fertility (e.g., (Gómez-Velasco *et al.* 2014; Jaramillo-López, Ramírez and Pérez-Salicrup 2015; Ferreira *et al.* 2017; Pandit *et al.* 2020)), and addition of biochar during or after the composting has an additional beneficial effect (Pandit *et al.* 2020; Pagliaccia *et al.* 2024; Dhakal *et al.* 2025). Observed benefits of Bokashi addition include i) increased nutrient availability and organic matter content (Gashua *et al.* 2023); ii) production of plant growth-promoting substances such as 3-phenyllactic acid (Maki *et al.* 2021), iii) increased microbial abundance (Gómez-Velasco *et al.* 2014; Mondaca *et al.* 2025), diversity (Luo *et al.* 2022), functional potential (Park and Kremer 2010) and activity (Gómez-Velasco *et al.* 2014) in soil, which might in turn accelerate nutrient turnover, iv) mitigation of the impact of heavy metal contamination (Barajas-Aceves and Rodríguez-Vázquez 2013), and v) increased suppressiveness of soils to plant pathogens such as fungi or root-knot nematodes in some cases (Ferreira *et al.* 2017; Shin *et al.* 2017; Van Der Sloot *et al.* 2024). These reported benefits appeal also to Bokashi practitioners on the household-scale, who often use their Bokashi compost as a soil amendment for gardening. However, waste material for Bokashi composting for agricultural use is more diverse than on the household-scale and often includes waste products such as chicken, pig or cow manure, fishery waste or even sludge from municipal wastewater treatment (e.g., (Epelde *et al.* 2018; Urra *et al.* 2019; Abo-Sido *et al.* 2021; De Oliveira *et al.* 2025)). These waste products typically have a higher nitrogen content than plant-based waste products, and the effect of plant-based

Bokashi on plant growth might thus not be 1:1 comparable, providing lower nitrogen amendment. On the other hand, application of organic amendments of animal or anthropogenic origin can pose a higher risk of introduction of inorganic or organic contaminants, pathogens and AMR (especially for manure- or sewage-based products) (Epelde *et al.* 2018; Urra *et al.* 2019), which is likely less problematic in household-scale fermentations using plant-based waste products.

Impact of Bokashi application on soil microbial communities

Microbial community composition in actively fermenting Bokashi has been only rarely assessed even on the agricultural scale, with slightly more studies looking at the microbial community composition in soils amended with Bokashi compost. Some studies have looked at the microbial community composition in Bokashi produced from animal manure, and the observed community composition is rather different from that observed in the present study: The community was more diverse and the most abundant taxa included *Actinobacteriota*, *Proteobacteria* and *Chloroflexota* (Abo-Sido *et al.* 2021; De Oliveira *et al.* 2025), and lactic acid bacteria were not a prominent part of the microbial communities. In comparison, relative abundances of *Proteobacteria* exceeded 10% only in half of the actively fermenting Bokashi samples in the present study, and *Actinobacteria* accounted for <2% of the microbial community in all actively fermenting Bokashi samples (Fig. 2). The impact of soil amendment with Bokashi compost is inconclusive with some studies reporting changes in microbial community composition after addition of Bokashi fertilizer (Luo *et al.* 2022) and increases in bacterial CFU counts (Gashua *et al.* 2023), while others suggest no significant differences in bacterial community composition between Bokashi-amended and unamended soil, suggesting that Bokashi-microbes could not outcompete the native soil microbes (Shin *et al.* 2017). Indeed, microbial communities in soil factory samples in the present study strongly differed from starter, fermenting Bokashi and leachate samples (Fig. 2), and only the most abundant genus, *Lentilactobacillus*, was detected in three of the soil factories in notable abundances (Fig. 2B). The communities were rather similar to microbial communities reported from urban and suburban soils (e.g., (Baruch *et al.* 2021; Gómez-Brandón *et al.* 2022; Baldi *et al.* 2023; Probst *et al.* 2023; Tessler *et al.* 2023)). The most abundant genera detected in actively fermenting Bokashi, *Lentilactobacillus, Lactiplantibacillus, Companilactobacillus,* and *Levilactobacillus*, are homo- and heterofermentative lactic acid bacteria that are frequently found in fermented foods as well as food spoilage (Zheng *et al.* 2020). They can produce a variety of organic acids as well as ethanol. In the present study, lactate and acetate were the dominant organic acids

produced, and fermentation pathways detected in metagenomic reads and MAGs confirmed the potential for diverse fermentations in actively fermenting Bokashi. Lactate, acetate and ethanol are also among the main fermentation products detected in Bokashi compost produced from plant-based waste on a larger scale (Pian *et al.* 2023). In addition, other studies have also reported the formation of butyrate (Alattar *et al.* 2012), which was likewise detected in the present study, albeit in low concentrations. Thus, while application of Bokashi compost to soils might not have a strong impact on the soil's microbial community composition, it nonetheless adds organics acids, alcohols and other nutrients produced by lactic acid bacteria. Indeed, studies suggest that the benefits of Bokashi amendment are more likely due to the input of nutrients than the addition of living "effective" microorganisms to the soil (Schenck Zu Schweinsberg-Mickan and Müller 2009; Mayer *et al.* 2010; Kok *et al.* 2022). Bokashi addition does lead to a priming effect but not to long-term enhancement of SOC sequestration (Kok *et al.* 2022).

Potential health benefits and risks of household-scale Bokashi composting

Use of Bokashi composting at the household scale features several routes of potential exposure to microorganisms from the Bokashi process: 1) inhalation or ingestion (e.g., after deposition on foods) of aerosolized microbes, 2) skin exposure during handling of Bokashi compost or soil factory, or 3) ingestion e.g., when using Bokashi as fertilizer for plants including fruits and vegetables. In general, lactic acid bacteria are known for health-promoting properties (Gilliland 1990), and many probiotics contain strains of lactic acid bacteria (Zielińska and Kołożyn-Krajewska 2018). Aerosolized lactic acid bacteria have been suggested as therapeutics to help combat lung diseases like infections with *P. aeruginosa* (Glieca *et al.* 2024), pulmonary neutrophilic inflammation (Nicola *et al.* 2024) or virus infections (Spacova *et al.* 2023). Bacteria and fungi can be aerosolized from household compost bins, but most of the aerosols are detected rather close to the compost bin (Naegele *et al.* 2016). Any negative effects (like occurrence of storage mites and molds) have been observed only close to the waste bin, not in the whole kitchen or home (Naegele *et al.* 2016), and it can be assumed that any potential positive impacts by aerosolized lactic acid bacteria would likewise be limited to the close proximity of the Bokashi bin or places in which leachate has been applied (e.g., flower pots). Microbes from compost including lactic acid bacteria have been detected on the hands of compost facility workers (Madsen, Rasmussen and Frederiksen 2024), which might result in health risks if no proper hand hygiene is practiced. Potential pathogens were only rarely detected in amplicons and metagenomes from Bokashi (0.005-0.5% of all reads), and most of

them were opportunistic pathogens commonly found in soils or in the human microbiome. However, risks for immuno-compromised people cannot be excluded. Among the non-bacterial species detected in the metagenomic reads was the parasitic whipworm *Trichuris trichiura*, which was detected in all samples (0.13-0.37% relative read abundance). Whipworm infections are not common in Europe (Behniafar *et al.* 2024), so it is highly unlikely that there has been contamination through a diseased person in the household. However, fruit and vegetable produce imported from places where *T. trichiura* is endemic such as South America, Africa or East Asia could have had traces of said organism on the peel due to contamination (Alemu, Nega and Alemu 2020; Eslahi *et al.* 2022). *T. trichiura* has been detected on fruits and vegetables (e.g., banana, mango, cucumber, tomato) with a prevalence of 5-10% depending on the country of origin (Eslahi *et al.* 2022). Detection of DNA, however, does not give any indication whether there are actually living and/or infectious parasites present, as it might also persist in dead cells or as extracellular DNA.

Another concern is the introduction of antimicrobial resistance (AMR) into the home environment. Indeed, lactic acid bacteria have been discussed as potential donors of AMR genes, and prospective probiotic food supplements have to undergo testing to prove that AMR transfer from these strains is not possible. Vancomycin resistance is intrinsic to many lactic acid bacteria like *Lactobacillus* and *Pediococcus* (Tynkkynen 1998), and genes encoding vancomycin resistance were detected in nearly all MAGs in the present study. However, as vancomycin resistance in lactic acid bacteria is chromosomally encoded and not closely related to enterococcal vancomycin resistance, it is believed not to be easily transferable (Tynkkynen 1998). Resistance to other antimicrobials like tetracycline, ciprofloxacin or chloramphenicol is fairly widespread in Lactobacilli with some carrying resistance to multiple antibiotics (Campedelli *et al.* 2019; Rivera-Rodriguez *et al.* 2025). Transfer of genes conveying resistance to streptomycin or erythromycin from Lactobacilli to other gram-positive and -negative species has been described (Gevers, Huys and Swings 2003; Ouoba, Lei and Jensen 2008; Tian *et al.* 2025). The MAGs recovered in the present study did encode a wide range of potential AMR genes and there might thus be a risk of AMR genes from Bokashi microbes to other species. However, it has to be kept in mind that after fermentation the Bokashi ferment was mixed with garden soil for the soil factory. Soils including garden soils are a huge reservoir of diverse AMR genes, and the interaction of humans with soils of any kind is thus associated with a certain risk (Popowska *et al.* 2012; Knapp *et al.* 2017; Lee *et al.* 2018; Zhao *et al.* 2025). It thus can be

argued that handling the Bokashi ferment and using the soil from the soil factory does not carry a heightened risk compared to e.g., gardening.

Conclusions, limitations and future perspectives

Bokashi as a household-scale composting method suitable for urban lifestyle provides a promising low-technology, citizen-lead means to enhance microbial diversity and health of urban ecosystems. Bokashi practitioners, who have developed skills to make Bokashi with a sensory feeling for the process, achieve good fermentation results and productive soil without having a detailed microbiological understanding of the process. However, more information on the microbiological promises and risks of Bokashi are needed e.g., regarding waste recycling, health or environmental authorities to make recommendations or regulations concerning Bokashi composting. Bokashi composting on the household scale is surrounded by optimistic expectations to improve ecosystems and even human health, but the scientific basis for these claims is not well studied. In collaboration with Bokashi practitioners, this study assessed the microbial community composition in different stages of the Bokashi composting process. Lactic acid bacteria dominated starters and actively fermenting Bokashi, and metabolic pathways for the production of diverse organic acids were detected. While potentially pathogenic organisms and AMR genes were detected, their abundance was extremely low. Thus, handling Bokashi and keeping it in the house is likely not associated with increased health risks. Microbes from the Bokashi might be enriched in the household, e.g., due to aerosols near the compost bin. However, this was not assessed in the present study, and further studies are needed to assess the potential spread of microorganisms from the Bokashi compost in the household. Moreover, only a limited number of samples was analyzed in this study. This was in part due to limitations in the amount of funding, but also in part in the difficulty of recruiting suitable participants. The results gained here can be used to inform the design of future studies, e.g., concerning the sampling procedure, replication efforts or sampling locations of interest. The study thus forms an important steppingstone for more detailed and more targeted studies on household-scale Bokashi composting.

Acknowledgements

The authors thank the Bokashi practitioners for active participation in the workshops and for providing samples as well as Sara Rinne for help with laboratory work.

Author contributions

The article is based on interdisciplinary collaboration between Veera Kinnunen, who provided understanding of the lay Bokashi practice based on her ethnographic study, and Katharina Kujala, who contributed with microbiological expertise.

Katharina Kujala: Conceptualization (decision on sampling strategy, planning of physicochemical and molecular analyses), Formal analysis (bioinformatics), Investigation (laboratory work), Resources, Writing – Original draft, Writing – Review and editing, Visualization, Project administration, Funding acquisition

Veera Kinnunen: Conceptualization (General understanding of the lay Bokashi practice, planning and organization of participatory citizen workshops, participant recruitment), Investigation (leading discussions during workshops and summarizing input from participants), Resources, Writing – Review and editing, Project administration, Funding acquisition

Funding

This work was supported by CitizenANTS funding provided by the Biodiverse Anthropocenes research program (supported by the University of Oulu and the Research Council of Finland PROFI6 funding (2021-2026), project Urbiolabs awarded to VK), the Research Council of Finland (project 322753 awarded to KK, and project number 362001 supporting VK) as well as the Jane and Aatos Erkko foundation (project ICE-FISHING awarded to KK).

Data availability

The sequence data generated in this study has been deposited in the European Nucleotide Archive ENA under study accession number PRJEB100587.

References

Abo-Sido N, Goss JW, Griffith A *et al.* Microbial transformation of traditional fermented fertilizer bokashi alters chemical composition and improves plant growth. 2021, DOI: 10.1101/2021.08.01.454634.

Alattar MA, Green TR, Henry J *et al.* Effect of microaerobic fermentation in preprocessing fibrous lignocellulosic materials. *Appl Biochem Biotechnol* 2012;**167**:909–17.

Alcock BP, Huynh W, Chalil R *et al.* CARD 2023: expanded curation, support for machine learning, and resistome prediction at the comprehensive antibiotic resistance database. *Nucleic Acids Res* 2023;**51**:D690–9.

Alemu G, Nega M, Alemu M. Parasitic contamination of fruits and vegetables collected from local markets of Bahir Dar City, Northwest Ethiopia. *Res Rep Trop Med* 2020; **11**:17–25.

Alneberg J, Bjarnason BS, De Bruijn I *et al.* Binning metagenomic contigs by coverage and composition. *Nat Methods* 2014;**11**:1144–6.

Arkin AP, Cottingham RW, Henry CS *et al.* KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* 2018;**36**:566–9.

Attali D, Baker C. *ggExtra: Add Marginal Histograms to "Ggplot2", and More "ggplot2" Enhancements. R Package Version 0.11.0.9000.* https://github.com/daattali/ggextra

Baldi DS, Humphrey CE, Kyndt JA *et al.* Native plant gardens support more microbial diversity and higher relative abundance of potentially beneficial taxa compared to adjacent turf grass lawns. *Urban Ecosyst* 2023;**26**:807–20.

Barajas-Aceves M, Rodríguez-Vázquez R. Effects of organic amendments on the mobility of Pb and Zn from mine tailings added to semi-arid soils. *J Environ Sci Health Part B* 2013;**48**:226–36.

Baruch Z, Liddicoat C, Cando-Dumancela C *et al.* Increased plant species richness associates with greater soil bacterial diversity in urban green spaces. *Environ Res* 2021;**196**:110425.

Bedward M, Eppstein D, Menzel P. *Packcircles: Circle Packing. R Package Version 0.3.7.* https://CRAN.R-project.org/package=packcircles

Behniafar H, Sepidarkish M, Tadi MJ *et al.* The global prevalence of *Trichuris trichiura* infection in humans (2010-2023): A systematic review and meta-analysis. *J Infect Public Health* 2024;**17**:800–9.

Bolyen E, Rideout JR, Dillon MR *et al.* Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 2019;**37**:852–7.

Bowers RM, Kyrpides NC, Stepanauskas R *et al.* Correction: Corrigendum: Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol* 2018;**36**:660–660.

Buchfink B, Reuter K, Drost H-G. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* 2021;**18**:366–8.

Camacho C, Coulouris G, Avagyan V *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* 2009;**10**:421.

Campedelli I, Mathur H, Salvetti E *et al.* Genus-Wide Assessment of Antibiotic Resistance in *Lactobacillus* spp. Björkroth J (ed.). *Appl Environ Microbiol* 2019;**85**:e01738-18.

Chaumeil P-A, Mussig AJ, Hugenholtz P *et al.* GTDB-Tk v2: memory friendly classification with the genome taxonomy database. Borgwardt K (ed.). *Bioinformatics* 2022;**38**:5315–6.

Chivian D, Jungbluth SP, Dehal PS *et al.* Metagenome-assembled genome extraction and analysis from microbiomes using KBase. *Nat Protoc* 2023;**18**:208–38.

Christel DM. The use of bokashi as a soil fertility amendment in organic spinach cultivation. 2017.

De Coster W, Rademakers R. NanoPack2: population-scale evaluation of long-read sequencing data. Alkan C (ed.). *Bioinformatics* 2023;**39**:btad311.

De Oliveira AL, Kruker G, Guidi ES *et al.* Rhizobacteria and bokashi-type biofertilizer interaction in common bean crop. *Pesqui Agropecuária Trop* 2025;**55**:e81195.

Dhakal G, Fujino T, Magar ST *et al.* Co-application of bokashi and biochar alleviates water stress, improves soil fertility and enhances wheat production under water-deficit conditions. *Soil Syst* 2025;**9**:33.

Epelde L, Jauregi L, Urra J *et al.* Characterization of composted organic amendments for agricultural use. *Front Sustain Food Syst* 2018;**2**:44.

Eren AM, Kiefl E, Shaiber A *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat Microbiol* 2020;**6**:3–6.

Eslahi AV, Olfatifar M, Karim MR *et al.* Global incidence of helminthic contamination of vegetables, cucurbits and fruits: A systematic review and meta-analysis. *Food Control* 2022;**133**:108582.

Evans BR, Leighton FA. A history of One Health: -EN- A history of One Health -FR- Histoire du concept « Une seule santé » -ES- Historia de «Una sola salud». *Rev Sci Tech OIE* 2014;**33**:413–20.

Fan YV, Lee CT, Klemeš JJ *et al.* Evaluation of Effective Microorganisms on home scale organic waste composting. *J Environ Manage* 2018;**216**:41–8.

Ferreira JC, Hernandes I, Brito OD *et al.* Dosages of bokashi in the control of *Meloidogyne javanica* in lettuce, in greenhouse. *Hortic Bras* 2017;**35**:224–9.

Flandroy L, Poutahidis T, Berg G *et al.* The impact of human activities and lifestyles on the interlinked microbiota and health of humans and of ecosystems. *Sci Total Environ* 2018;**627**:1018–38.

Formowitz B, Elango F, Okumoto S *et al.* The role of "effective microorganisms" in the composting of banana (*Musa* ssp.) residues. *J Plant Nutr Soil Sci* 2007;**170**:649–56.

García-Hernández SA, Marín-Sánchez J, Romero-Méndez M *et al.* Productive and quality response of six varieties of bell pepper (*Capsicum annuum* L.) to organic fertilization in Gualdalzacar, S.L.P. *Rev Bio Cienc* **7**.

Gashua AG, Sulaiman Z, Mohammad Yusoff M *et al.* Short-term effects of bokashi fertilizer with reduced NPK fertilization on soil fertility, growth, and yield of rubber trees. *Pertanika J Trop Agric Sci* 2023;**46**:839–59.

Gevers D, Huys G, Swings J. In vitro conjugal transfer of tetracycline resistance from *Lactobacillus* isolates to other Gram-positive bacteria. *FEMS Microbiol Lett* 2003;**225**:125–30.

Gilliland SE. Health and nutritional benefits from lactic acid bacteria. *FEMS Microbiol Lett* 1990;**87**:175–88.

Glieca S, Quarta E, Bottari B *et al.* Development of inhalation powders containing lactic acid bacteria with antimicrobial activity against *Pseudomonas aeruginosa*. *Int J Antimicrob Agents* 2024;**63**:107001.

Gómez-Brandón M, Herbón C, Probst M *et al.* Influence of land use on the microbiological properties of urban soils. *Appl Soil Ecol* 2022;**175**:104452.

Gómez-Velasco DA, Álvarez-Solís JD, Ruiz-Valdiviezo VM *et al.* Enzymatic activities in soil cultivated with coffee (*Coffea arabica* L. *cv.* 'Bourbon') and amended with organic material. *Commun Soil Sci Plant Anal* 2014;**45**:2529–38.

Haahtela T. A biodiversity hypothesis. *Allergy* 2019;**74**:1445–56.

Hanski I, Von Hertzen L, Fyhrquist N *et al.* Environmental biodiversity, human microbiota, and allergy are interrelated. *Proc Natl Acad Sci* 2012;**109**:8334–9.

Higa T. Effective microorganisms: A biotechnology for mankind. *Proceedings of the 1st International Conference on Kyusei Nature Farming*. Washington, DC: USDA, 1991, 8–14.

Hui N, Grönroos M, Roslund MI *et al.* Diverse environmental microbiota as a tool to augment biodiversity in urban landscaping materials. *Front Microbiol* 2019;**10**:536.

Huson DH, Auch AF, Qi J *et al.* MEGAN analysis of metagenomic data. *Genome Res* 2007;**17**:377–86.

Huson DH, Beier S, Flade I *et al.* MEGAN community edition - interactive exploration and analysis of large-scale microbiome sequencing data. Poisot T (ed.). *PLOS Comput Biol* 2016;**12**:e1004957.

Jahn M, Leslie D, Dicko A *et al. WeightedTreemaps: Generate and Plot Voronoi or Sunburst Treemaps from Hierarchical Data. R Package Version 0.1.4.* https://CRAN.R-project.org/package=WeightedTreemaps

Jaramillo-López PF, Ramírez MI, Pérez-Salicrup DR. Impacts of Bokashi on survival and growth rates of *Pinus pseudostrobus* in community reforestation projects. *J Environ Manage* 2015;**150**:48–56.

Kang DD, Li F, Kirton E *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* 2019;**7**:e7359.

Kinnunen V. Knowing, living, and being with bokashi. In: Brives C, Rest M, Sariola S (eds.), *With Microbes*. Mattering Press, 2021, 64–83.

Knapp CW, Callan AC, Aitken B *et al.* Relationship between antibiotic resistance genes and metals in residential soil samples from Western Australia. *Environ Sci Pollut Res* 2017;**24**:2484–94.

Kok DD, Scherer L, De Vries W *et al.* Relationships of priming effects with organic amendment composition and soil microbial properties. *Geoderma* 2022;**422**:115951.

Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012;**9**:357–9.

Lee JH, Park KS, Jeon JH *et al.* Antibiotic resistance in soil. *Lancet Infect Dis* 2018;**18**:1306–7.

Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. *Nucleic Acids Res* 2024;**52**:W78–82.

Lew PS, Nik Ibrahim NNL, Kamarudin S *et al.* Optimization of bokashi-composting process using Effective Microorganisms-1 in smart composting bin. *Sensors* 2021;**21**:2847.

Luo Y, Gonzalez Lopez JB, Van Veelen HPJ *et al.* Bacterial and fungal co-occurrence patterns in agricultural soils amended with compost and bokashi. *Soil Biol Biochem* 2022;**174**:108831.

Madsen AM, Rasmussen PU, Frederiksen MW. Fungal and bacterial species on biowaste workers' hands and inhalation zone, and potential airway deposition. *Waste Manag* 2024;**183**:290–301.

Maki Y, Soejima H, Kitamura T *et al.* 3-Phenyllactic acid, a root-promoting substance isolated from Bokashi fertilizer, exhibits synergistic effects with tryptophan. *Plant Biotechnol* 2021;**38**:9–16.

Manninen J, Saarenpää M, Roslund M *et al.* Microbial communities on dry natural rocks are richer and less stressed than those on man-made playgrounds. Gralnick JA (ed.). *Microbiol Spectr* 2025;**13**:e01930-24.

Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 2011;**17**:3.

Mayer J, Scheid S, Widmer F *et al.* How effective are 'Effective microorganisms® (EM)'? Results from a field study in temperate climate. *Appl Soil Ecol* 2010;**46**:230–9.

McDonald D, Jiang Y, Balaban M *et al.* Greengenes2 unifies microbial data in a single reference tree. *Nat Biotechnol* 2024;**42**:715–8.

Miranda KM, Espey MG, Wink DA. A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 2001;**5**:62–71.

Mondaca P, Vergara A, Velásquez A *et al.* Comparative effects of compost, bokashi, and *Trichoderma* on soil and tomato quality in a mediterranean area. *J Soil Sci Plant Nutr* 2025;**25**:351–62.

Mosquera AT, Melo M, Quiroga C *et al.* Evaluación de fertilización orgánica en cafeto (*Coffea arabica*) con pequeños productores de Santander, Colombia. *Temas Agrar* 2016;**21**:90–101.

Murillo-Amador B, Morales-Prado LE, Troyo-Diéguez E *et al.* Changing environmental conditions and applying organic fertilizers in *Origanum vulgare* L. *Front Plant Sci* 2015;**6**, DOI: 10.3389/fpls.2015.00549.

Naegele A, Reboux G, Vacheyrou M *et al.* Microbiological consequences of indoor composting. *Indoor Air* 2016;**26**:605–13.

Nicola T, Wenger N, Xu X *et al.* A lactobacilli-based inhaled live biotherapeutic product attenuates pulmonary neutrophilic inflammation. *Nat Commun* 2024;**15**:7113.

Ouoba LII, Lei V, Jensen LB. Resistance of potential probiotic lactic acid bacteria and bifidobacteria of African and European origin to antimicrobials: Determination and transferability of the resistance genes to other bacteria. *Int J Food Microbiol* 2008;**121**:217–24.

Pagliaccia D, Ortiz M, Rodriguez MV *et al.* Enhancing soil health and nutrient availability for *Carrizo citrange* (X *Citroncirus* sp.) through bokashi and biochar amendments: An exploration into indoor sustainable soil ecosystem management. *Sci Hortic* 2024;**326**:112661.

Pandit NR, Schmidt HP, Mulder J *et al.* Nutrient effect of various composting methods with and without biochar on soil fertility and maize growth. *Arch Agron Soil Sci* 2020;**66**:250–65.

Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ Microbiol* 2016;**18**:1403–14.

Park H, DuPonte MW. How to cultivate indigenous microorganisms. 2008.

Park K-C, Kremer R. Establishment of an inoculated bacterial strain on plant root surfaces in soils with different microbial community. *J Korean Soc Appl Biol Chem* 2010;**53**:379–83.

Pian LB, Guerra JGM, Berbara RLL *et al.* Characterization, nitrogen availability, and agronomic efficiency of fermented composts in organic vegetable production. *Org Agric* 2023;**13**:461–81.

Pielou EC. The measurement of diversity in different types of biological collections. *J Theor Biol* 1966;**13**:131–44.

Popowska M, Rzeczycka M, Miernik A *et al.* Influence of soil use on prevalence of tetracycline, streptomycin, and erythromycin resistance and associated resistance genes. *Antimicrob Agents Chemother* 2012;**56**:1434–43.

Probst M, Gómez-Brandón M, Herbón C *et al.* Fungal-bacterial associations in urban allotment garden soils. *Appl Soil Ecol* 2023;**188**:104896.

Quince C, Lanzen A, Davenport RJ *et al.* Removing noise from pyrosequenced amplicons. *BMC Bioinformatics* 2011;**12**:38.

Quiroz M, Céspedes C. Bokashi as an amendment and source of nitrogen in sustainable agricultural systems: a review. *J Soil Sci Plant Nutr* 2019;**19**:237–48.

R Core Team. *R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.* https://www.R-project.org

Rivera-Rodriguez DE, Busby C, Cervantes-Barragan L *et al.* Widespread heteroresistance to antibiotics in *Lactobacillus* species. 2025, DOI: 10.1101/2025.03.24.644958.

Robeson MS, O'Rourke DR, Kaehler BD *et al.* RESCRIPt: Reproducible sequence taxonomy reference database management. Pertea M (ed.). *PLOS Comput Biol* 2021;**17**:e1009581.

Safwat SM, Matta ME. Environmental applications of Effective Microorganisms: a review of current knowledge and recommendations for future directions. *J Eng Appl Sci* 2021;**68**:48.

Sarmiento G, Amézquita M, Mena L. Use of bocashi and effective microorganisms as an ecological alternative in strawberry crops in arid zones. *Sci Agropecu* 2019;**10**:55–61.

Schenck Zu Schweinsberg-Mickan M, Müller T. Impact of effective microorganisms and other biofertilizers on soil microbial characteristics, organic-matter decomposition, and plant growth. *J Plant Nutr Soil Sci* 2009;**172**:704–12.

Shaffer M, Borton MA, McGivern BB *et al.* DRAM for distilling microbial metabolism to automate the curation of microbiome function. *Nucleic Acids Res* 2020;**48**:8883–900.

Shannon CE. A Mathematical Theory of Communication. *Bell Syst Tech J* 1948;**27**:379–423.

Shaw J, Marin MG, Li H. High-resolution metagenome assembly for modern long reads with myloasm. 2025, DOI: 10.1101/2025.09.05.674543.

Shen W, Sipos B, Zhao L. SeqKit2: A Swiss army knife for sequence and alignment processing. *iMeta* 2024;**3**:e191.

Shin K, Van Diepen G, Blok W *et al.* Variability of Effective Micro-organisms (EM) in bokashi and soil and effects on soil-borne plant pathogens. *Crop Prot* 2017;**99**:168–76.

Sieber CMK, Probst AJ, Sharrar A *et al.* Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat Microbiol* 2018;**3**:836–43.

Spacova I, De Boeck I, Cauwenberghs E *et al.* Development of a live biotherapeutic throat spray with lactobacilli targeting respiratory viral infections. *Microb Biotechnol* 2023;**16**:99–115.

Sun X, Liddicoat C, Tiunov A *et al.* Harnessing soil biodiversity to promote human health in cities. *Npj Urban Sustain* 2023;**3**:5.

Tessler M, David FJ, Cunningham SW *et al.* Rewilding in miniature: suburban meadows can improve soil microbial biodiversity and soil health. *Microb Ecol* 2023;**85**:1077–86.

Thorslund R. Bokashis respiration i jorden – Betydelse för dess klimatpåverkan. 2020.

Tian Q, Ye H, Zhou X *et al.* Evaluating the health risk of probiotic supplements from the perspective of antimicrobial resistance. Parveen S (ed.). *Microbiol Spectr* 2025;**13**:e00019-24.

Tynkkynen S. Vancomycin resistance factor of Lactobacillus rhamnosus GG in relation to enterococcal vancomycin resistance (van) genes. *Int J Food Microbiol* 1998;**41**:195–204.

Urra J, Alkorta I, Lanzén A *et al.* The application of fresh and composted horse and chicken manure affects soil quality, microbial composition and antibiotic resistance. *Appl Soil Ecol* 2019;**135**:73–84.

Van Der Sloot M, Maerowitz-Mcmahan S, Postma J *et al.* Soil-borne disease suppressiveness after short and long term application of fermented, composted or fresh organic amendment treatments in arable soils. *Appl Soil Ecol* 2024;**195**:105268.

Vicente NFDP, Marafeli ÉAM, Oliveira JADC *et al.* Uma revisão bibliográfica sobre bokashi dos últimos 20 anos. *Res Soc Dev* 2020;**9**:e279108339.

Wickham H. *Ggplot2: Elegant Graphics for Data Analysis.* https://ggplot2.tidyverse.org

Wu Y-W, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* 2016;**32**:605–7.

Xu H-L. Effects of a microbial inoculant and organic fertilizers on the growth, photosynthesis and yield of sweet corn. *J Crop Prod* 2001;**3**:183–214.

Zhang A. From immunity to collaboration: microbes, waste, and antitoxic politics. *Curr Anthropol* 2021;**62**:S298–310.

Zhao Y, Li L, Huang Y *et al.* Global soil antibiotic resistance genes are associated with increasing risk and connectivity to human resistome. *Nat Commun* 2025;**16**:7141.

Zheng J, Wittouck S, Salvetti E *et al.* A taxonomic note on the genus *Lactobacillus*: Description of 23 novel genera, emended description of the genus *Lactobacillus* Beijerinck 1901, and union of Lactobacillaceae and Leuconostocaceae. *Int J Syst Evol Microbiol* 2020;**70**:2782–858.

Zielińska D, Kołożyn-Krajewska D. Food-origin lactic acid bacteria may exhibit probiotic properties: review. *BioMed Res Int* 2018;**2018**:1–15.