Neural network-based cross-species chromatin annotation goes beyond sequence conservation

Noémien Maillard[1], Julie Demars[1*], Raphaël Mourad[2,3*]

[1] GenPhySE, Université de Toulouse, INRAE, ENVT, F-31326, Castanet Tolosan, France

[2] MIAT, INRAE, F-31326 Castanet-Tolosan, France

[3] University of Toulouse, UPS, 31062 Toulouse, France

*equal corresponding authors

noemien.maillard@inrae.fr, https://orcid.org/0009-0002-8840-6939

julie.demars@inrae.fr, https://orcid.org/0000-0003-2451-0059

raphael.mourad@univ-tlse3.fr, https://orcid.org/0000-0001-6700-5728

Abstract

Analogous to the Encyclopedia of DNA Elements (ENCODE) project, the Functional Annotation of ANimal Genomes (FAANG) consortium has produced chromatin annotations for domesticated animals, albeit in smaller amounts. Classical methods based on sequence conservation can be used to infer missing annotations, but are inappropriate for non-conserved sequences. Here, we demonstrate the ability of neural networks trained with human data to infer the missing chromatin annotations in livestock species. For this purpose, we comprehensively assessed predictions of transcription factors, chromatin accessibility, and histone marks in several species. Our results showed good predictions for various annotations in mammalian genomes, and surprisingly, also for bird genomes, despite the large phylogenetic distance from the human genome. Moreover, predictions were accurate even for non-conserved sequences, unlike conservation-based methods. Our results advocate the widespread use of neural networks in cross-species genome annotation, a key step in understanding the genetic architecture of complex traits.

Introduction

The chromatin, the complex between the DNA double helices and proteins, regulates the organization of DNA in three dimensions within the nucleus over time. Interactions between transcription factors (TF), insulators, histones, and their modifications, and DNA modulate chromatin condensation, leading to a continuum of DNA accessibility degrees that participate in gene expression regulation[1]. Therefore, the characterization of chromatin regulation is required to improve the annotation of genomes for a better understanding of noncoding genetic variation in complex traits[2].

For this purpose, the Encyclopedia of DNA Elements (ENCODE) consortium was created in the

2000s in order to generate data and help to elucidate how this regulation is modulated in model

organisms, including human[3] and mouse[4]. To date, almost 8000 human and 1400 murine ChIP-

seq of both TF and histone marks, DNase-seq, and ATAC-seq experiments are referenced in the

ENCODE dataportal[5,6] (Supplementary Data 1). Later, in the 2010s, the Functional Annotation

of ANimal Genomes (FAANG) initiative was organised to collect and analyse similar chromatin

regulation data for species of agronomic interest, such as chicken, cattle, and pigs[7]. Initial efforts

were made to standardize both protocols and computational analyses across different species,

rather than conducting a large number of experiments. Although the FAANG dataportal is still in

expansion through additional international projects[8], much less data is available for livestock

species (https://data.faang.org/). To date, 2,288 pig ChIP-seq and ATAC-seq experiments have

been referenced, including biological and technical replicates for a limited number of

experimental conditions (e.g., experiment type, tissue, developmental stage) (Supplementary

Data 1).

As well as continuing to conduct costly and time-consuming wet lab experiments across species,

computational methods based on sequence conservation and pairwise or multiple whole-genome

alignment have been used over the last two decades to transfer annotations from well-annotated

to poorly-annotated genomes[9,10]. However, multiple works have shown that non-coding

sequences, including regulatory elements, often show weak conservation signals, thereby

limiting the use of conservation-based tools[11,12].

As an alternative to conservation-based methods, deep learning methods have been trained with

ENCODE data to predict annotations of chromatin regulation, and a growing number of models

are now available[13]. For example, transposable elements[14] and regulatory activity predictions[15,16]

can be detected from sequence information. On the one hand, models are becoming increasingly

sophisticated in order to improve performance through precision and recall metrics, while on the

other hand, they are often used for a similar purpose in the study of model organisms[17]. Although

chromatin regulation is conserved across large evolutionary distances[12,18–20], few studies have

evaluated the use of such novel tools for cross-species functional annotation by focusing on

enhancer predictions[21–24].

Here, we comprehensively assessed the ability of three artificial neural networks (DeepBind[25],

DeepSEA[26], and Enformer[27]), trained using human and mouse chromatin regulation data

(ENCODE project, https://www.encodeproject.org/), to predict functional annotations from the

reference genomes of livestock species, including pig, cattle, chicken, and seabass. We first

compared the cross-species abilities of the different neural networks to predict a wide range of

experiments. We then evaluated the performance of neural networks trained on human and

mouse data to predict annotations across the phylogenetic tree in vertebrates and their

limitations. Finally, we investigated the influence of sequence conservation and genomic features

on the performance of chromatin prediction annotations.

Results

From both human and mouse-trained neural networks to cross-species predictions

To investigate the relevance of human and mouse-trained neural networks for translational

studies, especially to identify genome-to-phenome links in livestock, we used three artificial

neural networks presenting different properties (Supplementary Data 2) and trained on various

biological experiments related to chromatin regulation (DeepBind[25], DeepSEA[26], and

Enformer[27]) (Fig. 1a and Supplementary Data 2). DeepBind[25] and DeepSEA[26] are based on

Convolution layers, while Enformer also uses Transformer layers[27]. In addition, the size of the

input sequence ranges from 200 bp for DeepBind[25], 2 kb for DeepSEA[26], and up to 196 kb for

Enformer[27] (Fig. 1a and Supplementary Data 2). Cross-species predictions of functional

chromatin annotations, including transcription factor binding sites and histone marks obtained

from ChIP-seq data, as well as chromatin accessibility obtained from ATAC-seq and DNase-seq

data, were performed using the reference genomes of the pig (Sscrr11.1), cattle (ARS-UCS1.2),

chicken (bGalGal1) and European seabass (dlabrax2021) (Fig. 1b). The performance of the

predictions was evaluated using standard metrics (area under the receiver operating characteristic

curve [AUC-ROC] and area under the precision-recall curve [AUC-PR]) by comparing the

predicted data with the observed data from the FAANG data portal (https://data.faang.org/) (Fig.

1c). AUC-ROC is classically used for benchmarks, but does not account for class imbalance (i.e.

most bins do not overlap a peak). Hence, AUC-PR has also been used to account for class

imbalance.

Prediction of chromatin annotations in mouse using human-trained models

To evaluate the ability of neural networks trained with human data to predict diverse annotations

on the genome of another mammal, we first predicted annotations from the mouse reference

genome (mm39) using only DeepBind[25] and DeepSEA[26]. Since Enformer[27] has been trained

using both human and mouse data, predicting experiments from the mouse genome would likely

lead to data leakage. We split the mouse genome into bins and then predicted the presence or

absence of a ChIP-seq or DNase-seq peak in the bin. The comparisons included 9 transcription

factors (TFs), 4 histone mark ChIP-seq, and DNase experiments (Fig. 2 and Supplementary Data

3). For human predictions, AUC-ROC and AUC-PR values were extracted from DeepBind[25] and

DeepSEA[26] publications. For mouse data, 2 cell lines have been used as proxies according to the

human training data (WEHI, close to GM12801, used to train DeepBind CTCF model, and

mESC equivalent to hESC used to train other DeepBind and DeepSEA models).

Predictions on the mouse genome showed high accuracy for CTCF peaks with AUC-ROC values

of 0.934 and 0.989 using DeepBind and DeepSEA respectively, which were close to AUC-ROC

values for the human genome (0.997 for DeepBind and 0.980 for DeepSEA) (Fig. 2a). For

DeepSEA AUC-PR, which accounts for class imbalance, we obtained a value of 0.500 for mouse

predictions, whereas it was 0.664 for human predictions (Fig. 2a). For all evaluated TFs,

DeepSEA showed better AUC-PR values than DeepBind (Fig. 2b-c and Supplementary Data 3).

For DNase and histone marks predicted only with DeepSEA, AUC-ROC values for mouse

predictions were similar to those for human predictions, but there was a more significant gap

between AUC-PR values between mouse and human predictions (Fig. 2d-e). In addition, we also

observed that ubiquitous TFs, including CTCF (CCCTC-Binding Factor) and RBBP5 (RB

Binding Protein 5, Histone Lysine Methyltransferase Complex Subunit), showed higher AUC-

PR values than TFs involved in specific biological processes or developmental stages, such as

NANOG (Nanog Homeobox) (Fig. 2a-c and Supplementary Data 3).

Overall, the correlations between human and mouse AUC-PR values appeared to be stronger for

TF binding sites, regardless of the AUC-PR values, than for histone and DNase experiments,

despite the greater amount of peak sequence in the training data (Fig. 3a). Indeed, when training

a machine learning model, it is expected that the amount of training data (here the amount of

peak sequences) will strongly influence prediction performances. In agreement, we observed a

strong correlation between the number of peaks used to train the models and the AUC-PR

calculated with DeepSEA predictions on both human (Fig. 3b) and mouse (Fig. 3c) genomes. We

then focused on the *Sox2* and *Igf2* loci, which are known to carry binding sites for NANOG and

CTCF, respectively (Fig. 3d). For NANOG predictions, 1 true positive out of 4 was predicted in

the *Sox2* region. However, a high error rate has also been measured with 3 false negatives and 1

false positive (Fig. 3d). Conversely, CTCF experimental peaks are well predicted in the *Igf2*

region, albeit at a high false positive rate (Fig. 3d). Like CTCF, DNase and H3K4me3 observed

peaks were correctly predicted despite false positive peak predictions (Fig. 3d).

Taken together, our results demonstrated that the amount of data for training (i.e., the number of

peaks) is a critical factor for accurate cross-species predictions, which makes it more difficult to

predict TFs that are specific to a particular biological process.

Prediction of chromatin annotations in livestock using human and mouse-trained models

To assess the predictive ability of the neural networks across the phylogenetic tree, we then

predicted CTCF, four histone marks, and chromatin accessibility experiments in five tissues

(adipose, liver, lung, muscle, and spleen) with DeepSEA[26] and Enformer[27] from reference

genomes of livestock species. We chose two mammals (pig, Sscr11.1, and cattle, ARS-UCS1.2),

one bird (chicken, bGalGal1), and one fish (European seabass, dlabrax2021). We used the

DeepSEA model to compare the metrics between the human, the mouse, and the livestock

species genomes. We also used the Enformer model, which is expected to outperform other

neural networks given its additional use of Transformer layers (Fig. 4 and Supplementary Data

4).

First, the AUC-PR metrics obtained using DeepSEA for the comparison between the mouse and

the four species studied varied according to the experiments (Fig. 2a and 2d-e and

Supplementary Data 4a). The binding of CTCF (0.500) and DNase (0.287) appeared to be better

predicted in mice than in pigs (0.353 ± 0.035 se and 0.220 ± 0.012 se), cattle (0.309 ± 0.035 se

and 0.230 ± 0.017 se), and chickens (0.284 ± 0.010 se and 0.230 ± 0.021 se). Similarly,

H3K27me3 was better predicted in mice. These results can be the consequence of overall low

predictive performance and the absence of replicates for the evaluation of predictions from the

mouse genome. Conversely, H3K4me1, H3K4me3, and H3K27ac in pigs, cattle, and chickens

seem to be predicted more accurately than in mouse (Fig. 2d-e and Supplementary Data 4a).

Second, Enformer showed better predictive ability than DeepSEA, since all AUC-PR values

obtained with Enformer (Fig. 4a-b) are higher than those measured with DeepSEA

(Supplementary Data 4a-b) for all four species and tissues.

Overall, we observed more variability in the prediction metrics across different experiments than

between different tissues (Fig. 4a-b and Supplementary Data 4a-b). However, AUC-PR values

are lower in cattle adipose, liver, and muscle tissues than in pig and chicken tissues, reflecting

metabolic specificities of those tissues in ruminants (Fig. 4b). Indeed, monogastric and ruminant

energetic metabolism evolved differently due to their respective energetic substrate (glucose and

short fatty acids), as reviewed in[28,29]. Thus, our results seem to support this by showing particular

chromatin regulation related to energetic metabolism in cattle[30], which might be the consequence

of no ruminant data in the training dataset. For non-mammalian genomes, although we found

slightly less good metrics for the chicken genome, much lower metrics for the European seabass

genome were found, suggesting a shift in the predictive ability of neural networks across

phylogeny (Fig. 4a-b).

To illustrate predictions, we focused on two genomic regions: the *IGF2* locus, which undergoes

genomic imprinting in pigs (Fig. 4c) but not in chickens (Fig. 4d)[31], and the *CEBPA* region,

which is a master transcription factor gene for adipogenesis[32]. Overall, as previously observed

from genome-wide AUC-PR values, the Enformer neural network showed better predictive

ability than the DeepSEA model (Fig. 4c-d and Supplementary Data 4c-d). For CTCF

predictions in the IGF2 region, all peaks have been correctly predicted in pigs and half of the

peaks in chickens (Fig. 4c-d). Conversely, more false positives are observed for H3K4me3 peaks

in pigs than in chickens in both regions, while the opposite is shown for DNase (Fig. 4c-d).

Our results revealed that cross-species chromatin predictions based on neural networks trained

on human data go beyond mammals, especially for CTCF, H3K4me3, and H3K27ac (Fig. 4 and

Supplementary Data 4).

Prediction performances in regards to the phylogeny and sequence conservation

In order to better understand the relationship between predictive annotation metrics across

species, we first evaluated the correlation between the AUC-PR of H3K4me3 and the

phylogenetic distance calculated as substitutions per site (Fig.5a-c). The differences of DeepSEA

AUC-PR values measured among the species seem explained by the phylogenetic distances (Fig.

5a), since strong correlations are obtained between the four livestock species, mouse and human

(Fig. 5b). A similar trend is observed for the Enformer neural network (Fig. 5c). This suggests a

breakpoint in prediction performance at a certain sequence evolution distance beyond 1.166

substitutions per site.

In addition, we used a metric for sequence conservation - the Genomic Evolutionary Rate

Profiling (GERP) score - to investigate the impact of conserved and unconserved sequences on

prediction performances (Fig. 5d-f). For H3K4me3, we kept 1% of bins along the pig genome,

and the maximum and minimum GERP scores were used per bin. We evaluated the distribution

of these GERP scores in function of the prediction scores of each bin (Fig. 5d-e). We showed

that a high proportion of nucleotides displaying negative GERP scores, reflecting low sequence

conservation across species, presented high prediction scores as well as highly conserved

nucleotides with positive GERP scores (Fig. 5d-e). To focus on the prediction scores, we

extracted bins with a minimum GERP score lower than the mean of the minimum GERP scores

(the least conserved). We did the same for bins with a maximum GERP score greater than the

mean of maximum GERP scores (the most conserved). We then split the bins into two classes,

depending on whether or not they overlapped with an observed peak (Fig. 5f). For H3K4me3, we

show that a statistical difference exists between conserved and unconserved sequences, whether

bins overlap an observed peak or not (p-value=0.0463 and 2.160×10^{-9}, respectively, on a two-

sided unpaired Wilcoxon test). However, the differences measured between classes were

negligible (0.070 and 0.036 for bins overlapping a peak or not, respectively). CTCF bins

overlapping a peak showed no statistical difference between conserved and unconserved groups,

but this may be due to smaller sample sizes (Fig. 5g). In contrast, all other comparisons showed

the same tendency as H3K4me3 (Fig.5f-h).

Overall, our results showed that cross-species neural network prediction accuracy goes beyond

sequence conservation.

Prediction performances in pigs with respect to genomic organization

We then assessed whether prediction performances depend on genomic organization. For this

purpose, we predicted 6 experiments in 5 tissues on the pig reference genome and evaluated

AUC-ROC and AUC-PR metrics depending on different genomic features (Fig. 6a).

We found that CpG islands (CGIs) and 5'UTR regions are particularly well predicted with AUC-

PR values above 0.564 for all experiments, except H3K4me1, which has a mean AUC-PR of

0.422 ± 0.172 SE and 0.474 ± 0.141 SE for CpG and 5'UTR, respectively (Fig. 6a). In contrast,

repeated sequences and 3'UTR regions showed AUC-PR under 0.521 for all experiments, except

CTCF and H3K4me3 in 3'UTR, which cap respectively at 0.640 ± 0.221 SE and 0.614 ± 0.091

SE (Fig. 6a).

In addition, we measured the difference in prediction performance between human and pig

genomes in muscle tissue (Fig. 6b). The results showed that the strongly conserved regulatory

features (CpG, 5'UTR) seem to have the least AUC-PR loss. Moreover, the regulation pattern of

promoter activity seemed well predicted by the neural networks, as shown by poor AUC-PR loss

of H3K4me3 and DNase-seq. In contrast, H3K27ac and H3K4me1, the enhancer-associated

histone marks, showed AUC-PR loss ranging from 23.4% to 34.3%, and from 26.0% to 39.5%,

respectively. Those results are in concordance with DeepGCF[33] functional conservation scores

and suggest that neural networks improve the overall predictions.

Altogether, our results showed that prediction performances vary across genomic features, with

CpG islands and 5'UTR regions being the most accurately predicted.

Discussion

Here, we demonstrated the accuracy of neural networks trained with human and mouse data to

predict annotations on genomes of livestock species. Extensive benchmarking revealed great

prediction performances on genomes from the same phylogenetic clade as the genome used for

the training of neural networks (Mammals), but more strikingly, also for a more distant clade

(Aves). However, the results suggested a breakpoint at a phylogenetic distance beyond 1.166

substitutions per site, since for the seabass genome, prediction metrics were very low.

A few studies have highlighted conservation of regulatory elements in mammals[19,34],

vertebrates[18,20], and other animals[12]. These fundamental results demonstrate that the function of

gene regulatory elements, such as enhancers, is maintained over the course of evolution

regardless of their sequence conservation[35,36]. In parallel, during the last two decades,

computational methods based on sequence conservation have been used for cross-species

prediction, but are inherently not appropriate for unconserved sequences[10,37,38]. For example,

GERP++ relies on the alignment of the genome sequences of many divergent species and

searches for sequences whose similarity has been maintained during evolution. In contrast, more

recently, machine learning methods, which do not rely on sequence conservation but instead on

sequence features such as DNA motifs, have shown that enhancers share functional properties

across species, supporting biological data, but this was often limited to a few species of

mammals[22,23,33,39]. For instance, the two-step neural network approach, DeepGCF, first predicts

epigenomic functional effects and then uses these effects as input to predict the functional

conservation between humans and pigs[33]. Here, our findings are in concordance with the

literature, supporting that we can predict chromatin regulation beyond mammals, regardless of

sequence conservation.

Although we were able to evaluate the ability of neural networks to predict across several species

and point out strengths as well as some limits, our analysis was limited by the data available in

FAANG for the evaluated livestock species. It would be of great interest to evaluate such

performances for an even broader range of experiments. For example, the evaluation of other

histone modifications would have been interesting, such as H3K9me3, which has been shown to

be enriched for silencing LTRs, LINEs, and SVAs repeated sequences, and with tissue

specificities[40]. In addition, it would be valuable to study intermediate species across the tree of

life, such as reptiles and amphibians, to accurately define the shift in chromatin prediction

annotation abilities. However, accessing valuable chromatin-omics data for experimental

purposes may present an obstacle to performing AUC-PR.

As pointed out here and by the literature, machine and deep learning models are not perfect, and

several options exist in order to improve the cross-species predictions. Over Enformer and its use

of Transformer layers, it has been shown that model architecture is a key element in highlighting

different features and interactions between them[22,41]. Moreover, Kelley showed that using multi-

species data during the deep learning model training phase improves predictions[42], and while

annotation data from not commonly studied species are often lacking, alternatives like pseudo-

labeling have been proposed[43]. Finally, the improvement of molecular techniques used to study

chromatin regulation and annotate genomes might have some biases (sequence affinity of

enzymes, for example). Novel innovative methods, such as CUT&Tag, for efficient epigenomic

profiling, that have not been used in the training set nor in the evaluation set, may likely improve

predictions of chromatin accessibility[44]. Here, we used three artificial neural networks relying on

different parameters (DeepBind[25], DeepSEA[26], and Enformer[27]) as a first step to evaluate their

abilities for cross-species predictions of chromatin annotation, mainly in livestock species. This

area of research is in constant evolution, and novel models are regularly released. As an example,

ChromBPNet seems to show prediction performances close to Enformer performances but with a

lighter architecture based on convolution layers[45]. While it has not been used for cross-species

predictions, this suggests that other models might be used in order to improve cross-species

predictions.

In summary, our research opens the field of predicting experiments for species of agronomical

interest with artificial neural networks trained with human and mouse data. While neural

networks are generally developed to predict variant impact in human pathologies, we show here

that we can also use them for cross-species predictions and consider their use in the study of

complex traits for species of agronomical interest.

Methods

DeepBind model. DeepBind[25] is a convolutional neural network taking 200 bp sequences as

input (http://tools.genes.toronto.edu/deepbind/). The output is a pseudo-probability of a given

protein, ranging from 0 to 1. The network has been used to predict the binding of CTCF, RBBP5,

NANOG, MAX, MYC, POU5F1, SIN3A, SUZ12, YY1 with the corresponding models,

respectively D00328.018, D00797.001, D00786.001, D00504.005, D00785.001, D00795.001,

D00804.002, D00812.001, D00710.007. DeepBind predictions have been computed on the

Genobioinfo cluster of the Genotoul-Bioinfo platform, parallelized on 64 CPU cores

(approximately 11 hours for the predictions with all 9 models on the whole mouse genome).

DeepSEA model. DeepSEA[26] is a convolutional-based model taking a 2000 bp (context and

sequence) as input, and which outputs a pseudo-probability to observe a peak in a 200 bp

centered sequence. The model has been downloaded from Kipoi. DeepSEA was used to predict

2002 experiments (DNase, ChIP-seq of transcription factors, and histone marks) for each

sequence. Transcription factors (CTCF, RBBP5, NANOG, MAX, MYC, POU5F1, SIN3A,

SUZ12, YY1), DNase, and histone ChIP-seq experiments (H3K4me1, H3K4me3, H3K27ac,

H3K27me3) from tissues of interest (liver (HepG2), lung (A549), muscle (HSMM)) have been

selected. DeepSEA predictions have been computed with an NVIDIA RTX4090 GPU

(prediction times ranging from 50 minutes for the seabass genome to 2h30 for the pig and cattle

genomes).

Enformer model. Enformer[27] is a deep learning model composed of convolutional layers followed by transformer layers. It takes a 196,608 bp length sequence as input and outputs 896 bins of 128 bp length. Enformer predicts 5313 human genomic tracks (DNase-seq, ChIP-seq of transcription factors and histone marks, ATAC-seq, and CAGE) for each sequence. We extracted ChIP-seq experiments of CTCF, H3K4me1, H3K4me3, H3K27ac, H3K27me3, DNase-seq, and ATAC-seq related to adipose, liver, lung, muscle, and spleen tissues. Enformer predicts a normalized read count. Enformer predictions have been computed with an NVIDIA RTX4090 GPU (prediction times ranging from 30 minutes for the seabass genome to 2h50 for the human genome).

Mouse genome. Mouse reference genome (GRCm39, mm39) from Ensembl release 112 has been used to predict peaks with DeepBind and DeepSEA. For this purpose, the genome has been binned into 200 bp and 2000 bp bins for DeepBind and DeepSEA, respectively, and then DNA sequences have been extracted according to bins. For each sequence, the models were used to predict the pseudo-probability of binding for the different experiments.

Mouse ChIP-seq and DNase-seq data. All mouse experimental data have been obtained in the embryonic stem cell line (ESC). ChIP-seq bed files of transcription factors (CTCF (GSE60917 and GSE29218), RBBP5 (GSE22934), NANOG (GSE44286), MAX (GSE48175), MYC (GSE90893), POU5F1 (GSE44286), SIN3A (GSE81081), SUZ12 (GSE127804), YY1 (GSE68195)) have been extracted from ReMap2022[46] and bed files of DNase-seq and histone mark experiments have been extracted from ENCODE (DNase-seq (ENCFF178MGP), ChIP-seq

H3K4me1 (ENCFF426IIV), ChIP-seq H3K4me3 (ENCFF247GVM), ChIP-seq H3K27ac

(ENCFF360VIS), ChIP-seq H3K27me3 (ENCFF008XKX)) (Supplementary Data 5). Reads have

been aligned on the mm10 genome, then peaks called have been lifted over to the mm39 genome

version with the liftOver tool from UCSC (Supplementary Data 6).

AUC-ROC and AUC-PR values extraction from publications. For human (GRCh37, hg19 for

both DeepBind and DeepSEA) ChIP-seq peak predictions, we extracted available AUC-ROC

and AUC-PR values from publications.

AUC-ROC and AUC-PR computation. AUC-ROC and AUC-PR have been computed using the

PRROC R package (version 1.3.1). For class labeling, a bin has been considered positive (class

1) if at least half of the bin overlaps with an observed peak. For this, the findOverlaps function

from the GenomicRanges package was used (version 1.56.2) with default parameters (maxgap=-

1L, ignore.strand=TRUE) except minoverlap set at half of the bin width (100L for DeepBind and

DeepSEA and 64L for Enformer).

Livestock species genomes. Pig reference genome (Sscrofa11.1), chicken reference genome

(GRCg7b, bGalGal1), and European seabass reference genome (dlabrax2021) have been

downloaded from Ensembl release 112. The cattle reference genome (ARS-UCD1.2, bosTau9)

has been downloaded from Ensembl release 99. Each livestock species' genome has been used to

predict peaks with DeepSEA and Enformer. Each genome has been binned into 2000 bp and

196,608 bp bins for DeepSEA and Enformer predictions, respectively. For each sequence,

DeepSEA was used to predict the pseudo-probability to observe a peak, and Enformer predicted

a normalized read count. For DeepSEA and Enformer predictions, AUC-ROC and AUC-PR have

been calculated with the same method as for mouse experiments.

Livestock species ChIP-seq and chromatin accessibility data. For each experiment, pig,

chicken, and cattle data have been performed in adipose, liver, lung, muscle, and spleen tissues.

ChIP-seq (CTCF, H3K4me1, H3K4me3, H3K27ac, H3K27me3), DNase-seq (chicken), and

ATAC-seq (pig and cattle) bed files have been downloaded from the FAANG consortium

(GSE158430). European seabass bigBed of the same experiments in lung and spleen have been

extracted from the AQUA-FAANG consortium (ensembl.org database), then bed files have been

created from bigBed using bigBedToBed from UCSC (Supplementary Data 5). The pig reads

have been aligned on the Sscrofa11.1 pig genome assembly. The chicken ChIP-seq and DNase-

seq reads have been aligned on the galGal6 genome assembly, and peaks have been lifted over to

the bGalGal1 genome assembly using the liftOver tool from UCSC. The cattle reads have been

aligned on the ARS-UCD1.2 genome assembly. The seabass reads have been aligned on the

dlabrax2021 genome assembly.

Pig genome annotations. The GTF file of genomic feature annotations (Gene, Transcript, CDS,

Exon, 5'UTR, 3'UTR) has been extracted from Ensembl database release 112. Repeats have

been extracted from the UCSC database. CpG islands have been extracted from the UCSC Table

Browser. For conserved regions analysis, GERP scores have been extracted from Ensembl

release 113 (Supplementary Data 6).

Human genome. To calculate AUC-ROC and AUC-PR differences from human to pig with

Enformer predictions, the Human reference genome (GRCh38, hg38) from Ensembl release 112

has been used. For this purpose, the genome has been binned into 196,608 bp bins for Enformer,

and then DNA sequences have been extracted according to bins. For each sequence, the models

were used to predict the normalized read count for the different experiments. AUC-ROC and

AUC-PR have been calculated with the same method as for mouse experiments.

Human genome annotations. The GTF file of genomic feature annotations (Gene, Transcript,

CDS, Exon, 5'UTR, 3'UTR) has been extracted from Ensembl database release 112. Repeats

have been extracted from the UCSC database. CpG islands have been extracted from the UCSC

Table Browser (Supplementary Data 5).

NGS read alignments. For Integrative Genome Viewer[47] (IGV) panels, mouse, pig, and chicken

raw reads have been aligned on mm39, Sscr11.1, and bGalGal1 reference genomes, respectively.

Mouse raw reads have been extracted with the SRA run selector for transcription factors and

from the ENCODE consortium for H3K4me3 and DNaseI-Hypersensitivity experiments. Pig and

chicken raw reads have been extracted from data referenced in the FAANG consortium

corresponding to GSE158430 (PRJEB14330, PRJNA665197, PRJNA665199, PRJNA665209,

PRJNA665214, PRJNA665216, PRJNA665216). For ChIP-seq, inputs have been merged for

each tissue-sample couple (Supplementary Data 7). For the alignment of raw reads, the nf-core

ChIP-seq (for CTCF and H3K4me3 experiments) and ATAC-seq (for ATAC-seq and DNase-seq

experiments) pipelines have been used with default parameters. The macs_fdr parameter has

been set to 0.01 for ChIP-seq experiments and 0.05 for ATAC-seq/DNase-seq experiments.

Integrative Genome Viewer. Integrative Genome Viewer[47] (IGV) has been used to show how predictions fit observations for genomic regions. Predicted peak heights have been set with the autoscale parameter.

Statistical information

Wilcoxon tests have been computed using a two-tailed *wilcox_test* function from the coin R library. The r effect size has been computed by dividing the Z statistic from the Wilcoxon tests by the square root of the sample sizes for each test.

Code and data availability

All the code is available at: https://forge.inrae.fr/genesis/EAGLE.

All the bigwig prediction files are available at: https://entrepot.recherche.data.gouv.fr/ under the accession numbers https://doi.org/10.57745/TN3R6F, https://doi.org/10.57745/92QIQS, https://doi.org/10.57745/EEQDSP.

Acknowledgements

We would like to thank Sarah Djebali, Laurent Bréhélin, Bertrand Servin, and Juliette Riquet for their help and advice. We would also like to thank Guillaume Devailly for his help in the R analysis.

Funding

456 This study is supported by the PEPR Agro-écologie et Numérique PDOC2 – ANR-22-PEAE-

457 0015 PEAE 2022 and by the DIGIT-BIO Metaprogram Research from INRAE.

Contributions

460 N.M. performed analyses, wrote the manuscript, and drew figures. J.D. and R.M. conceived the

461 study, secured funding, supervised N.M., and read and edited the manuscript.

Competing interests

The authors declare no competing interests.

References

1. Klemm, S. L., Shipony, Z. & Greenleaf, W. J. Chromatin accessibility and the regulatory epigenome.

Nat. Rev. Genet. **20**, 207–220 (2019).

2. Ward, L. D. & Kellis, M. Interpreting noncoding genetic variation in complex traits and human

disease. *Nat. Biotechnol.* **30**, 1095–1106 (2012).

3. The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science* **306**, 636–640 (2004).

4. Yue, F. *et al.* A comparative encyclopedia of DNA elements in the mouse genome. *Nature* **515**, 355–

364 (2014).

5. Dunham, I. *et al.* An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**,

57–74 (2012).

6. Luo, Y. *et al.* New developments on the Encyclopedia of DNA Elements (ENCODE) data portal.

Nucleic Acids Res. **48**, D882–D889 (2020).

7. Andersson, L. *et al.* Coordinated international action to accelerate genome-to-phenome with

FAANG, the Functional Annotation of Animal Genomes project. *Genome Biol.* **16**, 57 (2015).

8.   Clark, E. L. *et al.* From FAANG to fork: application of highly annotated genomes to improve farmed

animal production. *Genome Biol.* **21**, 285 (2020).

9.   Hinrichs, A. S. *et al.* The UCSC Genome Browser Database: update 2006. *Nucleic Acids Res.* **34**,

D590–D598 (2006).

10. Zhao, H. *et al.* CrossMap: a versatile tool for coordinate conversion between genome assemblies.

Bioinformatics **30**, 1006–1007 (2014).

11. Nelson, A. C. & Wardle, F. C. Conserved non-coding elements and cis regulation: actions speak

louder than words. *Development* **140**, 1385–1395 (2013).

12. Wong, E. S. *et al.* Deep conservation of the enhancer regulatory code in animals. *Science* **370**,

eaax8137 (2020).

13. Chen, Z., Ain, N. ul, Zhao, Q. & Zhang, X. From tradition to innovation: conventional and deep

learning frameworks in genome annotation. *Brief. Bioinform.* **25**, bbae138 (2024).

14. Yan, H., Bombarely, A. & Li, S. DeepTE: a computational method for de novo classification of

transposons with convolutional neural network. *Bioinformatics* **36**, 4269–4275 (2020).

15. Ji, Y., Zhou, Z., Liu, H. & Davuluri, R. V. DNABERT: pre-trained Bidirectional Encoder Representations

from Transformers model for DNA-language in genome. *Bioinformatics* **37**, 2112–2120 (2021).

16. Kelley, D. R. *et al.* Sequential regulatory activity prediction across chromosomes with convolutional

neural networks. *Genome Res.* **28**, 739–750 (2018).

17. Eraslan, G., Avsec, Ž., Gagneur, J. & Theis, F. J. Deep learning: new computational modelling

techniques for genomics. *Nat. Rev. Genet.* **20**, 389–403 (2019).

18. Pérez-Rico, Y. A. *et al.* Comparative analyses of super-enhancers reveal conserved elements in

vertebrate genomes. *Genome Res.* **27**, 259–268 (2017).

19. Woo, Y. H. & Li, W.-H. Evolutionary Conservation of Histone Modifications in Mammals. *Mol. Biol.*

Evol. **29**, 1757–1767 (2012).

20. Phan, M. H. Q. *et al.* Conservation of regulatory elements with highly diverged sequences across

large evolutionary distances. *Nat. Genet.* **57**, 1524–1534 (2025).

21. Minnoye, L. *et al.* Cross-species analysis of enhancer logic using deep learning. *Genome Res.* **30**,

1815–1834 (2020).

22. Chen, L., Fish, A. E. & Capra, J. A. Prediction of gene regulatory enhancers across species reveals

evolutionarily conserved sequence properties. *PLOS Comput. Biol.* **14**, e1006484 (2018).

23. MacPhillamy, C., Alinejad-Rokny, H., Pitchford, W. S. & Low, W. Y. Cross-species enhancer prediction

using machine learning. *Genomics* **114**, 110454 (2022).

24. Hong, J., Gao, R. & Yang, Y. CrepHAN: cross-species prediction of enhancers by using hierarchical

attention networks. *Bioinformatics* **37**, 3436–3443 (2021).

25. Alipanahi, B., Delong, A., Weirauch, M. T. & Frey, B. J. Predicting the sequence specificities of DNA-

and RNA-binding proteins by deep learning. *Nat. Biotechnol.* **33**, 831–838 (2015).

26. Zhou, J. *et al.* Deep learning sequence-based ab initio prediction of variant effects on expression and

disease risk. *Nat. Genet.* **50**, 1171–1179 (2018).

27. Avsec, Ž. *et al.* Effective gene expression prediction from sequence by integrating long-range

interactions. *Nat. Methods* **18**, 1196–1203 (2021).

28. Hocquette, J. F. & Bauchart, D. Intestinal absorption, blood transport and hepatic and muscle

metabolism of fatty acids in preruminant and ruminant animals. *Reprod. Nutr. Dev.* **39**, 27–48

(1999).

29. Bauchart, D., Gruffat, D. & Durand, D. Lipid absorption and hepatic metabolism in ruminants. *Proc.*

Nutr. Soc. **55**, 39–47 (1996).

30. Chen, S. *et al.* Comparative epigenomics reveals the impact of ruminant-specific regulatory

elements on complex traits. *BMC Biol.* **20**, 273 (2022).

31. Frésard, L. *et al.* Transcriptome-wide investigation of genomic imprinting in chicken. *Nucleic Acids*

Res. **42**, 3768–3782 (2014).

32. Lefterova, M. I. *et al.* PPARγ and C/EBP factors orchestrate adipocyte biology via adjacent binding on

a genome-wide scale. *Genes Dev.* **22**, 2941–2952 (2008).

33. Li, J. *et al.* Learning functional conservation between human and pig to decipher evolutionary

mechanisms underlying gene expression and complex traits. *Cell Genomics* **3**, 100390 (2023).

34. Villar, D. *et al.* Enhancer Evolution across 20 Mammalian Species. *Cell* **160**, 554–566 (2015).

35. Snetkova, V. *et al.* Ultraconserved enhancer function does not require perfect sequence

conservation. *Nat. Genet.* **53**, 521–528 (2021).

36. Yang, S. *et al.* Functionally conserved enhancers with divergent sequences in distant vertebrates.

BMC Genomics **16**, 882 (2015).

37. Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral substitution rates

on mammalian phylogenies. *Genome Res.* **20**, 110–121 (2010).

38. Davydov, E. V. *et al.* Identifying a High Fraction of the Human Genome to be under Selective

Constraint Using GERP++. *PLoS Comput. Biol.* **6**, e1001025 (2010).

39. Kaplow, I. M. *et al.* Relating enhancer genetic variation across mammals to complex phenotypes

using machine learning. *Science* **380**, eabm7993 (2023).

40. Trizzino, M., Kapusta, A. & Brown, C. D. Transposable elements generate regulatory novelty in a

tissue-specific fashion. *BMC Genomics* **19**, 468 (2018).

41. Cochran, K. *et al.* Domain-adaptive neural networks improve cross-species prediction of

transcription factor binding. *Genome Res.* **32**, 512–523 (2022).

42. Kelley, D. R. Cross-species regulatory sequence activity prediction. *PLOS Comput. Biol.* **16**, e1008050

(2020).

43. Phan, H., Brouard, C. & Mourad, R. Semi-supervised learning with pseudo-labeling compares

favorably with large language models for regulatory sequence prediction. *Brief. Bioinform.* **25**,

bbae560 (2024).

44. Kaya-Okur, H. S. *et al.* CUT&Tag for efficient epigenomic profiling of small samples and single cells.

Nat. Commun. **10**, 1930 (2019).

45. Pampari, A. *et al.* ChromBPNet: bias factorized, base-resolution deep learning models of chromatin

accessibility reveal cis-regulatory sequence syntax, transcription factor footprints and regulatory

variants. 2024.12.25.630221 Preprint at https://doi.org/10.1101/2024.12.25.630221 (2025).

46. Hammal, F., de Langen, P., Bergon, A., Lopez, F. & Ballester, B. ReMap 2022: a database of Human,

Mouse, Drosophila and Arabidopsis regulatory regions from an integrative analysis of DNA-binding

sequencing experiments. *Nucleic Acids Res.* **50**, D316–D325 (2022).

47. Robinson, J. T., Thorvaldsdottir, H., Turner, D. & Mesirov, J. P. igv.js: an embeddable JavaScript

implementation of the Integrative Genomics Viewer (IGV). *Bioinforma. Oxf. Engl.* **39**, btac830

(2023).

Figure

Fig. 1: Approach to perform cross-species functional chromatin annotation predictions in

livestock using human and mouse-trained neural networks. a. Neural networks trained on

human and mouse data. Three neural networks are used, including DeepBind[25], DeepSEA[26], and

Enformer[27]. b. Cross-species predictions. References genomes of mouse (mm39), pig (Sscr11.1),

cattle (ARS-UCS1.2), chicken (bGalGal1), and European seabass (dlabrax2021) have been used.

c. Evaluation of prediction performances. Standard metrics (AUC-ROC and AUC-PR) were

calculated by comparing the predicted data with the observed data from the ENCODE

(Encyclopedia   of   DNA   Elements,   https://www.encodeproject.org/),   ReMap2022

(https://remap2022.univ-amu.fr/), and FAANG (Functional Annotation of Animal Genomes,

https://data.faang.org/) databases. For a finer analysis, Genomic Evolutionary Rate Profiling

(GERP) conservation scores have been used to show whether only sequences with conserved

nucleotides can be well predicted.

Fig. 2: Predictive performances of two human-trained neural networks on the mouse genome. AUC-ROC (left) and AUC-PR (right) for CTCF (CCCTC-binding factor) (a), RBBP5 (RB Binding Protein 5, Histone Lysine Methyltransferase Complex Subunit) (b), NANOG (Nanog Homeobox) (c), DNase (d), and histone marks (e). The DeepBind prediction metric values are in grey and black for the human (hg19) and mouse (mm39) genomes, respectively. The DeepSEA prediction metric values are in orange and yellow for the human (hg19) and mouse (mm39) genomes, respectively. The human AUC values have been extracted from the original articles for DeepBind[25] and DeepSEA[26]. Except for the DeepBind-CTCF prediction (model trained with GM12801 cell line), which was compared to CTCF conducted in WEHI cell line, the observations used for comparisons with predictions were conducted on mouse embryonic stem cells (ESCs) (n=1 for each comparison).

Fig. 3: Comparison of the predictive performances of DeepSEA across different types of experiments. a. Correlation between human (hg19, x axis) and mouse (mm39, y axis) AUC-PR values (blue dotted line and statistics). The size of the dots indicates the number of peaks counted in the experiments used to train DeepSEA. b and c. Correlations between peak count of experiments used for DeepSEA training and AUC-PR of human (b) or mouse (c). The colors of the dots represent the type of experiments with transcription factor binding sites (red), DNase (orange), and histone marks (yellow). The correlations take only TF (red dotted lines and statistics) into account or all experiments (blue dotted lines and statistics). d. Visualization of the *Sox2* and *Igf2* regions through the Integrative Genome Viewer. The mouse annotated genes are shown on the first track with *Sox2* and *Igf2* on the left and right panels, respectively. For CTCF (red), NANOG (blue), DNase (green), and H3K4me3 (purple), both observed experiments in

610 mouse embryonic stem cells (ESC) (top) and predicted annotations (bottom) are shown for

611 comparison. Annotations of the mouse genome are from Ensembl release 112.

Fig. 4: Predictive performances of the Enformer human and mouse-trained neural network on different livestock genomes. a and b. AUC–ROC (left) and AUC–PR (right) for the same predictions from pigs, cattle, chicken, and European seabass reference genomes with Enformer[27] grouped by experiments (a, purple) or by tissue (b, green). Data shown are means ± SE. Grouped by experiments, n=36, 30, 44, 52, 34, and 44 for CTCF, Chromatin accessibility, H3K4me1, H3K4me3, H3K27ac, and H3K27me3, respectively. Grouped by tissues, n=50, 62, 34, 56, and 38 for adipose, liver, lung, muscle, and spleen, respectively. c and d. Visualization of the *IGF2* (left) and *CEBPA* (right) regions in pig (c) and chicken (d) through the Integrative Genome Viewer. The annotated genes are shown on the first track. For CTCF (red), DNase (green), and H3K4me3 (purple), both observed experiments in pig and chicken liver tissue (top), and predicted experiments (bottom) are shown for comparison. Annotations of pig and chicken genomes are from Ensembl release 112.

Fig. 5: Impact of the sequence conservation on the performance of predictions in pigs. a. Phylogenetic distance (substitutions per site) of commonly studied species and species of interest (bold). gadMor1 is used as a proxy for European seabass. b and c. Relationship between AUC-PR of H3K4me3 predicted by DeepSEA[26] (b) and Enformer[27] (c) and phylogenetic distances, relative to the human genome, of species of interest. Pearson (R) and Spearman (Rs) correlations have been calculated excluding human AUC-PR. d and e. Distribution of minimum GERP scores

(d) and maximum GERP scores (e). 1% of bins have been sampled all along the pig reference

genome (n=1,897,280). For each bin, the maximum and minimum GERP scores have been

extracted. Vertical line is the mean of minimum (d) and maximum (e) GERP used to differentiate

unconserved (pink, n=372,291) or conserved (green, n=372,267) bins. f to h. Distribution of

Enformer prediction scores of H3K4me3 (f), CTCF (g), and H3K4me1 (h) in adipose tissue

according to their conservation status (pink vs green) and the overlap (dark pink or green) or not

(light pink or green) with an observed peak. For each experiment, a two-sided unpaired

Wilcoxon test has been realized between unconserved not overlapping a peak bins (light pink,

n=13781, 13960, 13565 for H3K4me3, CTCF and H3K4me1 respectively) and conserved not

overlapping a peak bins (light green, n=13744, 13988, 13657 for H3K4me3, CTCF and

H3K4me1 respectively), and between unconserved overlapping a peak bins (dark pink, n=394,

215, 610 for H3K4me3, CTCF and H3K4me1 respectively) and conserved overlapping a peak

bins (dark green, n=407, 163, 494 for H3K4me3, CTCF and H3K4me1 respectively).

Additionally, r effect sizes have been calculated for each group: r=-0.036, -0.070, 0.154, -0.021,

-0,022 and -0.111 for H3K4me3 no peak overlap, H3K4me3 peak overlap, CTCF no peak

overlap, CTCF peak overlap, H3K4me1 no peak overlap, and H3K4me1 peak overlap groups,

respectively. A negative r effect size indicates a prediction score higher in the unconserved

group, and a positive r effect size indicates a prediction score higher in the conserved group.

Fig. 6: Impact of pig annotated genomic features on the performance of Enformer predictions. a. AUC-ROC (left) and AUC-PR (right) calculated for 8 genomic features between observations from the FAANG project and predictions from the pig reference genome with Enformer[27], grouped by experiments (yellow) or tissues (pink). Data shown are means ± SE. Grouped by experiments, for each genomic feature, n=36, 30, 44, 52, 34, and 44 for CTCF, Chromatin accessibility, H3K4me1, H3K4me3, H3K27ac, and H3K27me3, respectively. Grouped by tissues, for each genomic feature, n=50, 62, 34, 56, and 38 for adipose, liver, lung, muscle, and spleen, respectively. b. Difference in percentage of AUC-ROC (left) and AUC-PR (right) from human (GRCh38/hg38) to pig (Sscr11.1) muscle. For each genomic feature, 1 human AUC has been subtracted from pig AUCs (n=2, 6, 12, 14, 10, and 12 for CTCF, Chromatin accessibility, H3K4me1, H3K4me3, H3K27ac, and H3K27me3, respectively). Data shown are means ± SE.