Disentangling the reproductive and metabolic transcriptional responses to diet in *Drosophila melanogaster*

M. Florencia Camus[1], Avishikta Chakraborty[1,2] and Max Reuter[1,2]

[1] *Research Department of Genetics, Evolution and Environment, University College London, Gower Street, London, WC1E 6BT, United Kingdom*

[2] *Centre for Life's Origins and Evolution, University College London, Gower Street, London, WC1E 6BT, United Kingdom*

Keywords: fitness, nutrition, reproduction, nutritional plasticity, life history, gene regulation

15 **Abstract**

16
Nutrition significantly influences various life-history traits in organisms, impacting decisions
about growth, reproduction, and longevity. Previous studies in *Drosophila* have demonstrated
that diet affects gene transcription, with many genes exhibiting altered expression between
protein- and carbohydrate-rich diets. Some responses, such as the upregulation of oogenesis-
related genes in protein-rich conditions, are physiologically intuitive, yet it remains
challenging to distinguish between metabolic adaptations to different diets and regulation
pertaining to reproductive investment in response to nutrient availability. In this study, we
explore the transcriptomic responses of virgin and mated flies to changes in nutritional
environments. We confirm previous phenotypic findings by observing responses to dietary
preferences and nutritional needs. Our results indicate that both nutritional conditions and
mating status lead to significant changes in gene expression in females. By comparing
responses between virgin and mated flies, we differentiate between basal dietary responses
and reproductive adaptations, with the latter involving eight times as many genes as the
former. Notably, we identify GATA family transcription factors and the heat-shock factor
(*Hsf*) as crucial regulators of diet-dependent reproductive genes. These findings enhance our
understanding of the complex interactions between nutrition and reproductive strategies in
Drosophila.

Introduction

Nutrition plays a central role in life-histories across the animal kingdom, shaping traits such as growth, development, the onset of reproductive maturity, investment in gamete production, and survival. It does so by providing not only building blocks and energetic fuel for organisms, but also environmental cues that are used to make active decisions about how to modulate life histories to maximise expected lifetime fitness. One of the best-studied traits that is heavily dependent on nutrition is reproduction, where the quality and quantity of resources affect investment in energetically costly offspring. In sexual species, females in particular allocate significant resources to the production of a small number of large-sized gametes and accordingly nutrition has a large impact on female fecundity [1, 2]. For example, *Drosophila* females housed in higher protein environments produce up to four times as many eggs as females in carbohydrate-rich (and more protein-poor) environments [3].

Recent work in fruit flies assessing the transcriptional responses to diet suggests that diet-dependent variation in female fecundity is at least in part due to active regulation of reproductive investment rather than nutrient limitation [4]. Thus, gene expression changes observed between an optimal protein-rich and a suboptimal carbohydrate-rich diet mirror those observed in response to dietary restriction; similar to limiting dietary input, providing a diet with a sub-optimal macronutrient balance triggers an evolutionarily conserved life history response, where reproduction is actively down-regulated. This life history response relies heavily on the insulin-signalling and especially the TOR pathway [4-6], via members of the GATA transcription factor family [4, 7]. The presence of these responses makes it difficult to interpret diet-dependent physiological and transcriptional changes, and discriminate which transcriptional responses are to the diet itself, and which are a consequence of the altered reproductive investment.

One way to disentangle these effects on the female transcriptome, is to combine diet treatments with manipulation of a female's mating status. In flies, like in most other organisms, a female's first mating triggers the transition from a non-reproductive to a reproductive state [8-10]. This also brings about permanent physiological changes to be able to meet reproductive demands [11, 12]. For example, studies have documented a drastic shift in nutritional preference before and after mating; while virgin females prefer to feed on

carbohydrate-rich foods similar to males, mated females not only consume more food, but

switch to preferring diets higher in protein content in order to be able to meet the nutritional

demands of egg production [13, 14]. Interestingly, this shift occurs even in genetically

sterilized females that do not possess a germline and are unable to produce eggs [15]. In large

parts, these changes are reliant on male-derived sex peptide, which is delivered as part of

accessory gland fluids during mating [16-18]. Several transcriptomic studies have

demonstrated that, as expected, the phenotypic changes after mating are associated with

widespread responses at the gene expression level [15, 19-21]. But again, it is not clear to

which degree these changes are a general, hard-wired response to mating and the onset of

reproduction or diet-dependent, and if the latter, how expression changes triggered by mating

are aligned with responses to diet.

Here, we report experiments that manipulate both diet and mating status in *Drosophila*

melanogaster to dissect the complex relationship between nutritional and reproductive

responses. Examining gene expression of virgin and mated flies fed either protein- or

carbohydrate-rich synthetic diets, we identify and characterise expression changes that

represent physiological adaptations to diet (shared by all flies), those that are generic

responses to mating (invariant across diets) and those that modulated by both mating status

and diet composition. We further compare these classes functionally and in terms of their

regulation.

We find several hundred genes respond to mating and nutrition. A large proportion of genes

show diet-dependent expression changes that are similar between virgin and mated flies.

These genes include a set of core metabolic genes involved in physiological diet responses

that are independent of mating status. But we also find evidence for a larger set of genes

where expression differences tend to be more pronounced in the mated cohort. Concordant

with previous work, we show that the diet response is mainly regulated by GATA

transcription factors. However, we also identify *Hsf* (heat shock factor) as an important

transcription factor modulating the upregulation of diet-specific reproductive response. In

line with a role of *Hsf*, additional experiments reveal effects of mating status and diet on

resistance to heat shock, adding support to an emerging role of heat shock factors in

reproduction.

Materials and Methods

<u>Fly Stocks and Maintenance</u>

We used *D. melanogaster* from the laboratory population LH_M. This population has been maintained as a large outbred population for over 400 non-overlapping generations [22, 23]. It is kept on a strict 14-day regime, with larval densities kept constant (~150–200 eggs per vial containing ca. 7ml of media) and a fixed adult population size (56 vials of 16 male and 16 females) at each generation. Flies were reared at $25^{\circ}C$, under a 12h:12h light:dark regime, on a cornmeal-molasses-yeast-agar food medium (see **Supplementary Methods**). For all experiments involving capillary feeding, we used high relative humidity (>80%) to minimise evaporation.

<u>Synthetic Diet</u>

For experiments, we used a modified liquid version of the holidic diet described in Piper et al. [24]. This food is prepared entirely from synthetic components to enable precise control over nutritional value (see **Supplementary Methods**). Two artificial liquid diets were made which differed in the ratio of protein (incorporated as individual amino acids) and carbohydrate (sucrose); with all other nutritional components provided in fixed concentrations. The nutrient ratios [P:C] used were 2:1 and 1:4, with the final concentration of each diet being 32.5g/L. These ratios were identified in previous work [4] as maximising the reproductive fitness of female and male LH_M flies, respectively. We note that P:C ratios may not be directly comparable to standard fly media, as nutrients in synthetic diets appear to be more readily accessible [24].

<u>Phenotypic data collection</u>

Diet preference: Flies from each sex were collected as virgins using CO_2 anaesthesia. Triplets of virgins were placed in individual vials containing molasses-yeast-agar culture medium (see Supplementary Methods for full recipe) with no added live yeast. We used triplets of flies as a means to balance between, on the one hand, replication across vials and, on the other hand, between-vial variance and workload [25]. Twenty vials of triplets were collected for each sex and mating status. All flies were aged for 2 days for them to become sexually mature. Subsequently, females in the "mated" treatment were provided with three virgin males (collected at the same time as females) as mating partners. These hextets of flies were left for 5 hours to mate, before being re-split by sex and transferred to new vials containing an 0.8% agar-water mixture. Virgin flies were not perturbed during this period and

placed on the 0.8% agar medium at the same time as mated flies. The agar-water vials

provide water for the flies but have no nutritional value.

In accordance with previous literature using this methodology [25, 26], flies were kept in

agar-water vials overnight, then supplied with two 5µl microcapillary tubes (ringcaps©,

Hirschmann) containing amino acid and carbohydrate solutions, respectively. Capillary tubes

were replaced daily, and food consumption for each fly trio was recorded for a period of three

days. As a control, the rate of evaporation for all diet treatments was measured in six vials

that contained the two solution-bearing capillary tubes but no flies, placed alongside the

experimental vials in the controlled temperature room. Their average evaporation per day was

subtracted from the values of volume consumed in experimental vials to correct diet

consumption for evaporation.

Nutritional requirement: The setup for nutritional requirement is almost identical to the diet

choice experiment, except at the point of synthetic diet allocation. Instead of two capillary

tubes (as per diet choice), flies were supplied with one microcapillary tube (ringcaps©,

Hirschmann) containing one of the two allocated diets. These diets varied in their protein-to-

carbohydrate ratios and captured the following nutritional rails (P:C): 2:1, 1:4. We had 20

vials with fly triplets per allocated diet and mating status. Capillary tubes were replaced daily,

and food consumption for each fly trio was recorded for a total period of three days. Flies

were exposed to diet treatments in a controlled temperature room (25°C), 12L:12D light

cycle and high relative humidity >80%. The rate of evaporation for all diet treatments was

measured by using five vials per diet that contained no flies, placed randomly in the constant

temperature chamber. The average evaporation per day was used to correct diet consumption

for evaporation.

Female fitness: Female adult fitness was measured as the number of eggs produced over a

fixed period of time (18h). This performance proxy is expected to correlate closely with other

fitness measures, such as the total number of offspring [27, 28]. Following the feeding

period, trios of virgin or mated females were placed in new agar vials and presented with

three males of the same age from the LH_M stock population. We note that this mating event is

the second mating opportunity for the "mated fly" group, and this was done so that we could

compare fecundity of recently inseminated females across both groups. Flies were left to

mate/oviposit for 18 hours in vials containing *ad libitum* food corresponding to their diet

treatment provided via capillary tubes. All flies were removed after this 18-hour mating window. Following removal of the flies, the total number of eggs laid were determined by taking pictures of the agar surface and counting eggs using the software *QuantiFly* [29].

Male fitness: Adult male fitness was quantified by counting the number of offspring produced in competitive mating trials, a metric previously validated in our lab as a reliable indicator of reproductive success, given that total adult progeny production in our population primarily depends on mating success [30]. Our method closely followed the experimental design outlined in [31], in which focal experimental males competed against standard competitor males for mating opportunities. After the previously described feeding period, a group consisting of three focal males (from both virgin and mated treatments), three virgin competitor males, and six virgin females was transferred to a fresh vial containing molasses-yeast-agar medium lacking live yeast, the primary food source for both sexes [32, 33]. Competitor males and females carried an LH_M genetic background and were homozygous for the recessive *bw−* eye-colour mutation. These competitor flies were raised under identical conditions and matched the age of the focal males. Following a 24-hour egg-laying period, adults were removed from the vials. The resulting eggs developed for 12 days, after which adult offspring were counted and classified based on eye colour to determine paternity: red-eyed offspring indicated paternity by the focal experimental males, whereas brown-eyed offspring were fathered by the competitor males.

Thermal tolerance: Fly thermal tolerance was measured only on female flies following established heat knockdown assays [34]. Following the feeding period in triplets, individual females were placed into 4ml water-tight glass vials, clipped onto a two-sided rack and immersed in a water bath with temperature set to 39°C. Thermal tolerance was measured as the time (minutes) it took an individual fly to stop locomotory activity (enter a coma-like state). The experiment was performed over two runs, with each run measuring heat knockdown time of 90 individual flies. Thus, a total of 180 flies were measured, corresponding to 15 fly triplets for each combination of diet and mating status.

Statistical analysis: To determine if dietary choices differed, we used a multivariate analysis of variance (MANOVA). The main model had protein and carbohydrate consumption as response variables, with mating status, sex, and their interaction as fixed effect. We performed subsequent univariate analysis of variance (ANOVA) to determine which

nutrient(s) contributed to the overall multivariate effect. All analyses were performed using
the *manova* function.

To examine whether the flies varied in the quantity they consumed of each diet, we used a
linear model to investigate differences in dietary consumption. We modelled total food
consumption as a response variable with diet treatment, sex, and their interaction as fixed
effects. Female fitness was measured as total number of eggs produced within an 18-hour
timeframe following a mating event. Male fitness was measured as the proportion of
offspring sired from the focal male. For this we modelled the response as a binomial vector
comprising the number of offspring sired by the focal male and the number sired by the
competitor male and diet composition as a categorical fixed effect. Given data followed a
normal distribution, we used a Gaussian linear model to analyse the data. Number of eggs
was the response variable, with mating status, and diet plus their interaction as fixed factors.

Thermal tolerance data was analysed with a Gaussian mixed model, fitted with the *lmer*
function in the *lme4* package [35]. The response variable was time (minutes) for flies to
become unconscious, with mating status and diet (plus their interactions) as fixed effects. We
included the nested term of "side of tank" within "run" of experiment as a random effect.

<u>Transcriptomic analysis</u>
Experimental Setup: Flies from each sex were collected as virgins using CO_2 anaesthesia.
For the virgin treatment, groups of six virgin flies were collected and placed into vials
containing culture medium (molasses-yeast-agar) with no added live yeast. For the mated
treatment, three virgin females and three virgin males were placed in individual vials. For
both treatments there was a total of six flies in the vials. Flies were left in these vials for a
period of 36-hours in which the mated treatment had the chance to mate. Following this
period, flies were split into triplets, and placed on 0.8% agar-water mixture. Agar-water vials
provide water for the flies but have no nutritional value. Flies were kept in these vials
overnight before being supplied with a microcapillary tube (ringcaps©, Hirschmann)
containing one of the two allocated diets.

Sample collection and RNA extraction: For each sex, we generated 3 biological replicates for
each diet treatment (virgins on protein diet, virgins on carbohydrate diet, mated on protein
diet, mated on carbohydrate diet), a total of twelve samples. For each replicate sample, we

pooled 4 triplets (a total of 12 flies) to ensure we collected sufficient RNA. Total RNA was

extracted using the *Qiagen RNeasy Minikit* (Qiagen BV, Venlo, The Netherlands) according

to manufacturer's instructions. Quantity and quality of RNA was first inspected using a

Nanodrop 2000 spectrophotometer (Wilmington, USA), and later verified using an Agilent

Tapestation 2200 (Agilent, USA) at the UCL Genomics facility.

Sequencing and read mapping: Library construction and sequencing were performed at the

UCL Institute of Child Health Genomics facility. Barcoded cDNA libraries were constructed

using the *KAPA Hyper mRNA Library prep kit* (Roche, USA) and mixed at equal

concentrations. This multiplexed sample was sequenced (43bp paired-end reads) on four

flowcell lanes on an Illumina Nextgen 500 instrument to an average of 18M reads per

sample.

Having verified that there was no bias towards particular libraries across the sequencing lanes

using the Illumina Basespace online server, we merged reads from all four lanes. Adaptors

and low-quality base pairs (below quality value of 3) were trimmed using trimmomatic v0.36

[36]. Trimmed reads from each sample were independently mapped to the *D. melanogaster*

genome release 6.19 using HISAT2 [37]. Mapped reads were manipulated using *samtools*

[38].

Statistical analyses: Exon-level read counts for each annotated gene were obtained using

htseq-count [39], using release 6.19 annotations obtained from the ENSEMBL Biomart, and

then summed across all exons within a single gene. Total read counts for each gene for the

twelve samples were then used for differential gene expression analysis using the

Bioconductor package *edgeR* [40], analysing each sex separately.

We first removed lowly expressed genes (read count < 2 across all samples). Read count data

were then normalised across libraries and expression dispersion parameters calculated using

the entire dataset. We tested for differential gene expression between our experimental

groups using the negative binomial models implemented in *edgeR*. We fitted a full model

where expression of each transcript was a function of mating status, diet and their interaction.

The significance of each model term was tested using a specific contrast matrix. To obtain

estimates of expression fold changes between the two diets for each mating status, we further

fitted separate models for each mating status with diet as the sole term.

Gene ontogeny enrichment was performed using the Bioconductor package *clusterProfiler* [41]. We further compared our list of genes that responded to mating to previous work that has examined transcriptomic responses of changes in mating status [20]. For this, we used the R package *GeneOverlap* [42] uses contingency table tests to identify greater than expected overlap between gene lists. In order to assess whether genes that showed similar diet responses were regulated by common transcription factors we used the Bioconductor package *RcisTarget* [43], which tests for enrichment of cis-regulatory motifs in 5kb windows upstream of genes in a given sets. In all analyses, we used a statistical significance threshold of 5% False Discovery Rate (FDR) [44].

All analyses were performed in R version 3.3.2 [45].

Results

<u>Phenotypic responses</u>

Phenotypic data collected alongside our expression data recapitulated previously described patterns of diet preference and diet-dependent choice and fitness. Briefly, we observe sexual dimorphism for diet preference, diet consumption and diet-dependent fitness, with females both preferring and performing better on high-protein diets and males on high-carbohydrate diets (**Supplementary Results 1**). Preference assays showed that females, but not males, show a large shift in nutritional preference following mating. While in a choice assay virgin females consumed separately offered protein and carbohydrate diets in relatively small quantities and in proportions that resulted in a protein-to-carbohydrate ratio of 1:2.5 (not dissimilar to the composition chosen by males), mated females consumed a much larger total quantity of food and almost tripled their protein intake, resulting in a significantly altered protein-to-carbohydrate ratio of $1:0.92$ (**Figure SR1.1A, Table SR1.1**). In no-choice assays, we found that mating has a significant effect on female total food consumption (**Figure SR.1B**, **Table SR1.2**), with mated females consuming more than virgin females. We also found that mated females consumed similar amounts when supplied with high-protein or high-carbohydrate carbohydrate diets, whereas virgin females consumed more on the high-protein diet (**Figure SR1.1B**, **Table SR1.2**). Reproductive fitness measurements showed sex-specific effects, whereby virgin females (freshly mated in our assay) were able to lay more eggs than mated females (who had already been previously reproductively in our assay) across both diets, but this fitness difference was not diet-dependant. Unlike in females,

mating did not influence male fitness. Regardless of mating status however, females performed best on protein and males on carbohydrate (**Figure SR1.1, Table SR1.3**).

Differential gene expression

After QC, our dataset included expression data for 12,880 genes for males and 9,862 genes for females. For males, we found 139 genes responding significantly in expression to changes in diet; no genes were differentially expressed in relation to mating or its interaction with diet (**Table 1**). Genes that were differentially expressed in relation to diet included a number of that are associated with male ejaculates, specifically accessory gland proteins (24A4, 53C14a, 53C14c, 53Ea, 62F, 63F, 76A), seminal fluid proteins (24Bb, 33A3, 53C14a, 60F, 87B) and serpins (28F, 38F, 77Bb), as well as *Seminal metalloprotease-1*, *Ejaculatory bulb protein* and *Ejaculatory bulb protein II* (see online archived data for a full list of genes [46]). All of these are down-regulated on protein-rich compared to carbohydrate-rich food, in line with the diet-specific reproductive investment observed in our phenotypic assays.

Transcriptomic responses in females were considerably more complex (**Table 1**). We identified 651 genes that showed differential expression between the diet treatments ('Diet' set), 345 genes that showed differential expression between virgin and mated flies ('Mating' set), 110 genes that showed additive expression changes in response to both diet and mating ('Diet+Mating' set) and 14 genes that showed a significant diet-by-mating interaction ('Diet×Mating' set), some in combination with additive diet and/or mating effects.

Genes in the Diet set represent a core metabolic response to nutritional composition. Expression changes occur in both directions (up- and down-regulation when going from a carbohydrate to protein environment, **Figure 1A**) and carbohydrate-to-protein fold changes are positively correlated between virgin and mated flies (r = 0.826, p < 0.001). In line with these genes being part of a basal metabolic response to diet, female Diet genes significantly overlap with the genes that show a significant diet effect in males (overlap = 41 genes, p < 0.001). Furthermore, genes in the female Diet set overlap with genes that we previously found to show concordant diet responses between mated males and females (S category in [4]: overlap = 305 genes, p < 0.001, **Table 2**). But in addition, the Diet set also includes smaller numbers of genes that show female-specific responses to diet, and were previously shown to respond differently to diet between mated females and males (DxS category in [4]: overlap = 15 genes, p < 0.001; D+(D×S): overlap = 70 genes, p < 0.001, **Table 2**). GO

analysis showed that Diet genes are significantly enriched for terms that relate to neuronal signalling and response to stimulus (**Figure 2**).

Genes in the Mating set represent female responses to fertilisation. Fold-changes from virgin to mated state are mostly positive (indicating increased expression after mating) and—as expected—are highly correlated across diets (**Figure 1B**). The set of mating-dependent genes identified here overlapped significantly with those previously reported as showing differential expression between mated and virgin females in the abdomen ([20], overlap = 19 genes, p< 0.001), but not in head-thorax (overlap = 102 genes, p = 0.5). They are also significantly enriched for previously identified reproductive genes with female-limited expression ([4], overlap = 45 genes, $p < 0.001$, **Table 2**). Gene Ontology (GO) enrichment analysis revealed that mating-dependent genes enriched for terms related to cuticle development, chitin catabolic process and cell adhesion (**Figure 2**).

For the small Diet+Mating category we find significant overlap with genes previously shown to have concordant diet-responses across the sexes (D category in [4]: overlap = 47 genes, p< 0.001, **Table 2**), as well as sex-specific diet-dependent regulation (DxS category in [4]: overlap = 5 genes, $p < 0.001$; D+(D×S): overlap = 31 genes, $p < 0.001$, **Table 2**). The genes show concordant responses to mating between diets (r = 0.884, $p < 0.001$, **Figure 1C**) and concordant responses to diets between mating treatments (r = 0.811, $p < 0.001$, **Figure 1D**, possibly with some undetected carbohydrate-specific expression changes following mating). In terms of function, genes in the Diet+Mating category were significantly enriched for carbohydrate metabolic processes (**Figure 2**).

Finally, for the Diet×Mating interaction category, we find patterns of fold change that indicate genes that are up-regulated in the carbohydrate environment in mated flies only, while being unaffected by diet in virgins (**Figures 1E and 1F**). Despite its small size, we found significant overlap between the gene set and those previously shown to have concordant diet responses across the sexes (D category in [4]: overlap = 7 genes, $p < 0.001$, **Table 2**) and those showing opposing diet-dependent regulation between males and females (D×S category [4]: overlap = 5 genes, $p < 0.001$, **Table 2**). Functionally, Diet×Mating genes are significantly enriched for biological processes of amino acid transport (**Figure 2**).

Expression regulation

We used analyses that detect the enrichment of binding motifs in the regulatory regions of sets of genes to infer the transcription factors that are the best candidates for driving the observed patterns of differential expression (**Table 3**, see data sets 6 and 7 in online repository [46] for full results).

For males, the genes responding in expression to diet are enriched for transcription factors of the GATA family, such as *grain* (*grn*), *GATAd*, *GATAe*, *pointer* (*pnr*) and *serpent* (*srp*). In the more complex female dataset, we found both the Diet and the Diet+Mating sets to be enriched for regulation by transcription factors of the GATA family, similar to regulation in males. Genes in the Mating set were enriched in binding motifs for *Heat shock factor* (*Hsf*), as well as *Signal-transducer and activator of transcription protein at 92E* (*Stat92E*), *Synaptobrevin* (*Syb*), *ftz transcription factor 1* (*ftz-f1*), and *Blimp-1*. Finally, genes in the Diet×Mating category had upstream regions enriched for binding motifs for *Hepatocyte nuclear factor 4* (*Hnf4*), *Ets96B*, *Hormone receptor 4* (*Hr4*), CG7786, and *bigmax*.

Thermal tolerance

With gene expression in response to mating enriched for regulation by *Hsf*, we assessed thermal tolerance in virgin and mated females on the two diets. The data showed that overall, mated females were significantly more heat-tolerant than virgin females (**Table SR1.4**, **Figure 3**). Additionally, we found that females fed on high-protein diet were also significantly more heat-tolerant than flies fed on high-carbohydrate diet (**Table SR1.4**, **Figure 3**). Both effects were of similar size (coefficients $\pm$ SE: mating = -2.86 $\pm$ 1.19, diet = -2.16 $\pm$ 1.17) and additive (mating status-by-diet interaction not significant, **Table SR1.4**). Accordingly, out of the four treatment groups, mated females fed with the protein-rich diet had the highest heat tolerance levels, with mated females on the carbohydrate-rich diet having a similar thermal tolerance to virgin flies on protein-rich diet and virgins fed the protein-rich diet having the lowest heat tolerance (**Figure 3**).

Discussion

In this study, we aimed to clarify the relationship between nutrition and reproduction in *Drosophila*. We placed virgin and mated flies into two nutritional environments that differed in their protein-to-carbohydrate ratios and had previously been shown to represent the female- and male-optimal diets [4]. We performed phenotypic measurements to confirm that

our diet manipulations replicated previously described patterns and then investigated the
transcriptome across these treatments to separate genes involved in nutritional responses from
those that are regulated in line with reproductive investment.

Our phenotypic experiments replicated previous findings, thus validating our approach. In
no-choice assays, we found the expected patterns of sex-specific fitness on the two types of
diet across virgin and mated flies [5, 9, 16], where females are more fecund on protein-rich
than carbohydrate-rich food while males gain greater mating success on the carbohydrate-rich
food. Choice assays provided evidence for a limited response to mating of male diet
preference [13, 24, 25], compared to a large shift of diet preference in females [13, 18].

Transcriptome analysis revealed sex differences in diet- and mating-dependent gene
expression that match those observed in feeding behaviour. Males show a very restricted
change in expression between the two types of diet, but this response is independent of
mating status, and mating in itself does not result in any transcriptomic changes. Females, in
contrast, not only share the core dietary response that is seen in males but also exhibit
profound changes in gene expression patterns that are triggered by mating, as well as some
mating-dependent changes in their responses to diet.

The fact that the male diet response includes many ejaculate-associated genes whose
expression is regulated in parallel to reproductive output suggests that male reproductive
investment is modulated in line with the nutritional environment. This contrasts with the
absence of any plasticity in response to mating status, implying that adult males are in a
reproductively active state whether mating takes place or not. This fact is consistent with an
onset of spermatogenesis already in the pupal state [47]. It should be noted, however, that the
absence of a transcriptional mating response in males reported here is at least somewhat at
odds with the only two previous studies that have compared gene expression in virgin and
mated males. Ellis and Carney [48] measured gene expression in heads using microarrays and
found a modest response, with 47 genes differentially expressed between males that had
mated vs. controls that had not been presented with a female. Fowler et al. [20] used RNA-
seq to assay gene expression in two tissues, head-thorax and abdomen. While they detected
no expression change in head-thorax, they found a much larger response to mating in the
abdomen with 2068 differentially expressed genes [20]. The fact that these effects of mating
were not detected in our study could be due to signals being obscured in the whole flies

(rather than specific tissues) analysed here. Furthermore, differences could arise due to the
timing of samples collection. The two other studies were conducted by taking samples soon
after an observed mating (2 hours for Ellis and Carney, 3h for Fowler et al.). It is possible
that expression changes induced by mating are transient and could then have been missed in
our design, where males were sampled from mating groups independently of the occurrence
of recent mating events. It is worth noting, however, that the differential expression of
accessory and seminal fluid proteins between mated and virgin males that was detected by
Fowler et al. ([20], table S2d), presumably due to replenishment after mating, is also apparent
in our dataset, but as correlate to differences in overall reproductive investment between the
diet treatments.

In contrast to results in males, the large transcriptional response to mating that we observe in
females is in line with previous results. For example, Fowler et al. [20] found 125 genes to be
differentially expressed in response to mating in the female abdomen and 2040 genes in
head-thorax. Similarly, Delbare et al. [49] compared the mating response across lines of
different geographic origins and found a shared set of 272 genes that were consistently
differentially expressed between mated and virgin females (in addition to a further 77 genes
that showed differential expression in specific combinations of female and male origin).
Beyond these quantitative similarities, we also find correspondence between the specific
genes that we and previous work identify as being regulated in response to mating. Thus,
genes in our Mating category overlap significantly with those that Fowler et al. [20] detected
as differentially expressed between the abdomens of mated and virgin females. We note that
a similar result was not obtained when comparing our gene set with Fowler et al.'s
differentially expressed genes from head-thorax. However, it is worth noting that their gene
set is very large (more than 2000 genes) and contains a smaller proportion of genes that are
also detected as mating-responsive in other studies than the thorax set (29% vs. 65% of genes
also detected elsewhere [20]).

While expression changes associated with mating have been well studied, we were also able
to assess the effect of diet alongside and in combination with the mating response. Doing so,
we were able to generate several new insights. First, our analysis allowed us to extend the
inference of a core metabolic response to macronutrient composition from mated females and
males (4) to virgin females and males. The existence of such a fundamental physiological
response to diet is not surprising, but the fact that it also occurs in reproductively inactive

individuals corroborates this finding. More interestingly, our analysis of transcription factor
binding sites revealed that the core response to diet composition, shared between virgin and
mated females as well as males, is heavily reliant on transcription factors of the GATA
family. This could seem surprising, as members of this transcription factor family have been
associated with a number of reproductive and ageing phenotypes. Thus, GATA transcription
factors are involved in the regulation of reproductive genes in fruit flies [50] and mosquitoes
[51], as well as in the ageing response to dietary restriction in *Caenorhabditis elegans* [52]
and regulation associated with dietary restriction and rapamycin treatment in flies [7]. While
these functions should not apply to virgins, our results are more in line with previous studies
that implicated GATA factors in the regulation of physiology and feeding, especially in
response to sugar [53]. Further work will be needed to investigate whether the regulation of
life history decisions and more general physiology are distinct functions of GATA factors, or
whether regulators of physiology are deployed in different contexts.

Beyond the core response to diet composition, the analyses presented here suggest that diet
composition and mating responses act largely independently and additively, with the number
of genes for which we detect a significant diet-by-mating status interaction being very small.
This is not what we had expected, as we had predicted that diet and mating would act
synergistically. Specifically, we had expected that a significant number of genes would show
differential responses to diet in virgin and mated females, with a larger increase in expression
under the more favourable conditions in mated females than in virgins, matching feeding
rates and reproductive investment. The fact that we observe an additive effect of mating
would imply that the mating response is largely qualitative (mating induces the production of
gene products) rather than quantitative (the amount of gene product produced scales with
reproductive output).

When interpreting the above results, however, it is important to keep in mind that the number
of genes with a significant interaction effect between diet and mating status is likely to be
limited by the statistical power to pick up these more subtle effects. It is interesting in this
respect to contrast the findings of our factorial analysis with results obtained by separate
univariate analyses performed in each mating regime (see **Supplementary Results 2**). These
analyses detect a large excess of differentially expressed genes in the mated compared to the
virgin mating regime (887 vs. 102 genes). Of course, this result is also limited by statistical
power, which here reduces the ability to detect shared diet-specific regulation when analysing

two datasets of half the size compared to the joint analysis in the main text. So taken together,
these results suggest that there is a sizeable set of genes that show quantitative differences
between mated and virgin flies in their level of diet-specific regulation, where fold changes
are larger in the former and smaller in the latter. These differences in differential expression
are small enough to not result in significant interaction effects in the joint analysis, but large
enough to lead to contrasting significance in the separate analyses.

Another interesting result from this work is the identification of *Heat shock factor* (*Hsf*) as
enriched in the regulation of genes related to reproduction and diet. As their name suggests,
heat-shock factors were first described as drivers for responses to thermal stress [54].
However, there have been several studies suggesting roles in important biological functions
in the absence of stress. Most notably, *Hsf* is required for oogenesis in *Drosophila* in a way
that does not depend on the expression of heat-shock proteins [55] and shows its highest level
of tissue-specific expression in ovaries [56]. A similar role for heat-shock factors in fecundity
has been described in mice [57-59], where they also have been shown to play a role in sperm
production (50). In line with the role of heat-shock factors in reproduction, a large number of
Hsf binding sites in *Drosophila* are associated with genes that are not involved in the heat-
shock response and *Hsf*-regulated genes are enriched for GO categories associated with
gamete formation and oogenesis [60]. In light of these previous findings, it seems that *Hsf* is
involved in the diet-dependent regulation of female reproduction, binding to genes that are
differentially regulated in response to the macronutrient composition of the food consumed
by females.

Not much is known about the mechanistic links between *Hsf* and diet. Work in rodent models
has also shown that the amino acid glutamine is involved in the activation of *Hsf* [61], while
experiments with the nematode *C. elegans* have demonstrated links between *Hsf* and the
insulin-signalling network, with inputs from both being required for development, stress
response, survival and reproduction [62]. In *Drosophila*, larvae raised in high-protein
environments are better able to tolerate thermal stress than flies raised in carbohydrate-rich
environments [63, 64]. A recent study further found that males fed a control food take longer
to be knocked down by heat than males fed a restricted diet rich in complex carbohydrate–
cellulose [65]. However, no study to date has examined how mating status interacts with the
dietary environment to shape the heat-shock response. Our experiments confirmed a role of
diet in modulating heat resistance, showing that flies fed a protein-rich diet were more heat

tolerant than flies fed a carbohydrate-rich food, possibly because a protein-rich diet favours

the production of protective proteins such as heat-shock proteins, antioxidants, and enzymes

involved in repair. Moreover, we found that independently of the diet treatment, mated flies

exhibited better tolerance to heat compared to virgins, suggesting that the metabolic

conditioning induced post-mating extends beyond egg or offspring production, generating

additional protective mechanisms without the trade-offs previously assumed. Future studies

should dissect this further by editing one or more known heat-shock factors to determine

which are specifically involved in oogenesis versus those contributing to stress tolerance. It is

unclear whether and to which degree these two roles are related. Previous work has proposed

that the role of *Hsf* in oogenesis is independent of the heat-shock response [55], yet our

observations suggest some possible links.

In conclusion, our aim was to investigate the metabolic pressures placed upon an organism as

a consequence of reproduction. We find that in female *Drosophila*, there is a large metabolic

rewiring that occurs after mating, and this is mainly to fuel the process of oogenesis. We

were able to dissect the nutritional response into basal and reproduction-related categories

and identified several transcription factors as possible drivers of these responses, including

GATA family transcription factors and *Hsf*. Our work provides further understanding on how

organisms are able to cope with changes in the internal and external physiology.

561
562

Data Availability Statement

RNA-seq raw data will be deposited on the Gene Expression Omnibus before publication.

Files with the data for phenotypic assays (VM-datafile1-fitness.csv, VM-datafile2-

preference.csv, VM-datafile3-heat.csv), lists of differentially expressed genes (VM-datafile4-

genelist-female.xlsx, VM-datafile5-genelist-male.xlsx) and enriched transcription factor

binding sites (VM-datafile6-genelist-female-TF.xlsx, VM-datafile7-genelist-male-TF.xlsx)

have been uploaded to the GSA figshare.

571
572

573   **Acknowledgements**

575 We would like to thank Nazif Alic and Adam Dobson for their help with planning the
576 experiments, Matt Piper for his input into the interpretation and Rebecca Finlay for help with
577 laboratory work.
578
579

580 **Study funding**

582 This study was supported by a Marie Skłodowska-Curie Research Fellowship (#708362), a
583 UKRI Natural Environment Research Council Independent Research Fellowship
584 (NE/V014307/1) and a Leverhulme Trust Research Project Grant (RPG-2023-198) to MFC
585 and Leverhulme Trust Research Project Grant (RPG-2021-414) to MR.
586

References

1.   Partridge, L., A. Green, and K. Fowler, *Effects of Egg-Production and of Exposure to Males on Female Survival in Drosophila-Melanogaster.* Journal of Insect Physiology, 1987. **33**(10): p. 745-749.

2.   Chapman, T. and L. Partridge, *Female fitness in Drosophila melanogaster: An interaction between the effect of nutrition and of encounter rate with males.* Proceedings of the Royal Society B-Biological Sciences, 1996. **263**(1371): p. 755-759.

3.   Lee, K.P., et al., *Lifespan and reproduction in Drosophila: New insights from nutritional geometry.* Proceedings of the National Academy of Sciences of the United States of America, 2008. **105**(7): p. 2498-2503.

4.   Camus, M.F., M.D. Piper, and M. Reuter, *Sex-specific transcriptomic responses to changes in the nutritional environment.* Elife, 2019. **8**.

5.   Alic, N. and L. Partridge, *Death and dessert: nutrient signalling pathways and ageing.* Current Opinion in Cell Biology, 2011. **23**(6): p. 738-743.

6.   Hoedjes, K.M., M.A. Rodrigues, and T. Flatt, *Amino acid modulation of lifespan and reproduction in Drosophila.* Current Opinion in Insect Science, 2017. **23**: p. 118-122.

7.   Dobson, A.J., et al., *Tissue-specific transcriptome profiling of Drosophila reveals roles for GATA transcription factors in longevity by dietary restriction.* Aging and Mechanisms of Disease, 2018. **4**: p. 5.

8.   Chapman, T., et al., *Cost of mating in Drosophila melanogaster females is mediated by male accessory gland products.* Nature, 1995. **373**(6511): p. 241-244.

9.   Sgrò, C.M., T. Chapman, and L. Partridge, *Sex-specific selection on time to remate in Drosophila melanogaster.* Animal Behaviour, 1998. **56**(5): p. 1267-1278.

10.   Liu, H.F. and E. Kubli, *Sex-peptide is the molecular basis of the sperm effect in Drosophila melanogaster.* Proceedings of the National Academy of Sciences of the United States of America, 2003. **100**(17): p. 9929-9933.

11.   Flatt, T., *Survival costs of reproduction in Drosophila.* Exp Gerontol, 2011. **46**(5): p. 369-75.

12.   Mirth, C.K., A. Nogueira Alves, and M.D.W. Piper, *Turning food into eggs: insights from nutritional biology and developmental physiology of Drosophila.* Current Opinion in Insect Science, 2019. **31**: p. 49-57.

13.   Camus, M.F., et al., *Dietary choices are influenced by genotype, mating status, and sex in Drosophila melanogaster.* Ecology and Evolution, 2018. **8**(11): p. 5385-5393.

14.   Corrales-Carvajal, V.M., A.A. Faisal, and C. Ribeiro, *Internal states drive nutrient homeostasis by modulating exploration-exploitation trade-off.* eLife, 2016. **5**: p. e19920.

15. Newell, N.R., et al., *The Drosophila Post-mating Response: Gene Expression and*
Behavioral Changes Reveal Perdurance and Variation in Cross-Tissue Interactions.
G3 (Bethesda), 2020. **10**(3): p. 967-983.

16. Chapman, T., et al., *The sex peptide of Drosophila melanogaster: Female post-mating*
responses analyzed by using RNA interference. Proceedings of the National Academy
of Sciences of the United States of America, 2003. **100**(17): p. 9923-9928.

17. Wigby, S. and T. Chapman, *Sex peptide causes mating costs in female Drosophila*
melanogaster. Current Biology, 2005. **15**(4): p. 316-321.

18. Ribeiro, C. and B.J. Dickson, *Sex peptide receptor and neuronal TOR/S6K signaling*
modulate nutrient balancing in Drosophila. Current Biology, 2010. **20**(11): p. 1000-
1005.

19. Innocenti, P. and E.H. Morrow, *Immunogenic males: a genome-wide analysis of*
reproduction and the cost of mating in Drosophila melanogaster females. J Evol Biol,
2009. **22**(5): p. 964-73.

20. Fowler, E.K., et al., *Divergence in Transcriptional and Regulatory Responses to*
Mating in Male and Female Fruitflies. Sci Rep, 2019. **9**(1): p. 16100.

21. McGraw, L.A., A.G. Clark, and M.F. Wolfner, *Post-mating gene expression profiles*
of female Drosophila melanogaster in response to time and to four male accessory
gland proteins. Genetics, 2008. **179**(3): p. 1395-408.

22. Chippindale, A.K., J.R. Gibson, and W.R. Rice, *Negative genetic correlation for adult*
fitness between sexes reveals ontogenetic conflict in Drosophila. Proceedings of the
National Academy of Sciences of the United States of America, 2001. **98**(4): p. 1671-
1675.

23. Rice, W.R., *Sexually antagonistic male adaptation triggered by experimental arrest*
of female evolution. Nature, 1996. **381**(6579): p. 232-4.

24. Piper, M.D.W., et al., *A holidic medium for Drosophila melanogaster.* Nature
Methods, 2014. **11**(1): p. 100-105.

25. Camus, M.F., et al., *Sex- and genotype-effects on nutrient-dependent fitness*
landscapes in Drosophila melanogaster. Proceedings of the Royal Society B-
Biological Sciences, 2017. **284**(1869): p. 20172237.

26. Reddiex, A.J., et al., *Sex-specific fitness consequences of nutrient intake and the*
evolvability of diet preferences. American Naturalist, 2013. **182**(1): p. 91-102.

27. Tanaka, T. and T. Yamazaki, *Fitness and its components in Drosophila melanogaster.*
Japanese Journal of Genetics, 1990. **65**(6): p. 417-26.

28. Hoffmann, A.A. and L.G. Harshman, *Male effects on fecundity in Drosophila*
melanogaster. Evolution, 1985. **39**(3): p. 638-644.

29. Waithe, D., et al., *QuantiFly: Robust trainable software for automated Drosophila*
egg counting. PLoS ONE, 2015. **10**(5): p. e0127659.

30.   Pischedda, A. and W.R. Rice, *Partitioning sexual selection into its mating success*
and fertilization success components. Proceedings of the National Academy of
Sciences of the United States of America, 2012. **109**(6): p. 2049-2053.

31.   Collet, J.M., et al., *Rapid evolution of the intersexual genetic correlation for fitness in*
Drosophila melanogaster. Evolution, 2016. **70**(4): p. 781-795.

32.   Sang, J.H., *The nutritional requirements of Drosophila melanogaster*, in *The Genetics*
and Biology of Drosophila. 1978, Academic Press, London. p. 159-192.

33.   Colinet, H. and D. Renault, *Dietary live yeast alters metabolic profiles, protein*
biosynthesis and thermal stress tolerance of Drosophila melanogaster. Comparative
Biochemistry and Physiology Part A Molecular & Integrative Physiology, 2014. **170**:
p. 6-14.

34.   Hoffmann, A.A., A. Anderson, and R. Hallas, *Opposing clines for high and low*
temperature resistance in Drosophila melanogaster. Ecology Letters, 2002. **5**(5): p.
614-618.

35.   Bates, D., M. Maechler, and B. Bolker, *lme4: Linear mixed-effects models using S4*
classes. 2012.

36.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina*
sequence data. Bioinformatics, 2014. **30**(15): p. 2114-20.

37.   Kim, D., B. Langmead, and S.L. Salzberg, *HISAT: a fast spliced aligner with low*
memory requirements. Nature Methods, 2015. **12**(4): p. 357-60.

38.   Li, H., et al., *The sequence alignment/map format and SAMtools.* Bioinformatics,
2009. **25**(16): p. 2078-9.

39.   Anders, S., P.T. Pyl, and W. Huber, *HTSeq - a Python framework to work with high-*
throughput sequencing data. Bioinformatics, 2015. **31**(2): p. 166-9.

40.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package*
for differential expression analysis of digital gene expression data. Bioinformatics,
2010. **26**(1): p. 139-40.

41.   Yu, G., et al., *clusterProfiler: an R package for comparing biological themes among*
gene clusters. OMICS: A Journal of Integrative Biology, 2012. **16**(5): p. 284-287.

42.   Shen, L. and M. Sinai, *GeneOverlap: Test and visualize gene overlaps.* R package
version 1.18.0, 2018.

43.   Aibar, S., et al., *SCENIC: single-cell regulatory network inference and clustering.*
Nature Methods, 2017. **14**(11): p. 1083-1086.

44.   Benjamini, Y. and Y. Hochberg, *Controlling the false discovery rate: A practical and*
powerful approach to multiple testing. Journal of the Royal Statistical Society. Series
B (Methodological), 1995. **57**(1): p. 289-300.

45.   Team, R.C., *R: A Language and Environment for Statistical Computing.* 2016.

46.   DRYAD, <Enter DRYAD/other repository data here>.

47.   Gärtner, S.M., et al., *Ex vivo culture of Drosophila pupal testis and single male germ-*
701   *line cysts: dissection, imaging, and pharmacological treatment.* J Vis Exp, 2014(91):
702   p. 51868.

703   48.   Ellis, L.L. and G.E. Carney, *Mating alters gene expression patterns in Drosophila*
704   *melanogaster male heads.* BMC Genomics, 2010. **11**: p. 558.

705   49.   Delbare, S.Y.N., et al., *Roles of Female and Male Genotype in Post-Mating*
706   *Responses in Drosophila melanogaster.* Journal of Heredity, 2017. **108**(7): p. 740-
707   753.

708   50.   Lossky, M. and P.C. Wensink, *Regulation of Drosophila yolk protein genes by an*
709   *ovary-specific GATA factor.* Molecular Cell Biology, 1995. **15**(12): p. 6943-52.

710   51.   Attardo, G.M., et al., *RNA interference-mediated knockdown of a GATA factor*
711   *reveals a link to anautogeny in the mosquito <i>Aedes aegypti</i>.* Proceedings of
712   the National Academy of Sciences, 2003. **100**(23): p. 13374-13379.

713   52.   Budovskaya, Y.V., et al., *An elt-3/elt-5/elt-6 GATA Transcription Circuit Guides*
714   *Aging in C. elegans.* Cell, 2008. **134**(2): p. 291-303.

715   53.   Erickson, P.A., et al., *Unique genetic signatures of local adaptation over space and*
716   *time for diapause, an ecologically relevant complex trait, in Drosophila*
717   *melanogaster.* PLoS Genet, 2020. **16**(11): p. e1009110.

718   54.   Ritossa, F., *A new puffing pattern induced by temperature shock and DNP in*
719   *drosophila.* Experientia, 1962. **18**(12): p. 571-573.

720   55.   Jedlicka, P., M.A. Mortin, and C. Wu, *Multiple functions of Drosophila heat shock*
721   *transcription factor in vivo.* The EMBO Journal, 1997. **16**(9): p. 2452-2462.

722   56.   Leader, D.P., et al., *FlyAtlas 2: a new version of the Drosophila melanogaster*
723   *expression atlas with RNA-Seq, miRNA-Seq and sex-specific data.* Nucleic Acids Res,
724   2018. **46**(D1): p. D809-D815.

725   57.   Christians, E., et al., *Maternal effect of Hsf1 on reproductive success.* Nature, 2000.
726   **407**(6805): p. 693-4.

727   58.   Akerfelt, M., R.I. Morimoto, and L. Sistonen, *Heat shock factors: integrators of cell*
728   *stress, development and lifespan.* Nat Rev Mol Cell Biol, 2010. **11**(8): p. 545-55.

729   59.   Xiao, X., et al., *HSF1 is required for extra-embryonic development, postnatal growth*
730   *and protection during inflammatory responses in mice.* The EMBO Journal, 1999.
731   **18**(21): p. 5943-5952.

732   60.   Gonsalves, S.E., et al., *Whole-genome analysis reveals that active heat shock factor*
733   *binding sites are mostly associated with non-heat shock genes in Drosophila*
734   *melanogaster.* PLoS One, 2011. **6**(1): p. e15934.

61.   Brasse-Lagnel, C., A. Lavoinne, and A. Husson, *Control of mammalian gene*
expression by amino acids, especially glutamine. FEBS J, 2009. **276**(7): p. 1826-44.

62.   Barna, J., et al., *Heat shock factor-1 intertwines insulin/IGF-1, TGF-beta and cGMP*
signaling to control development and aging. BMC Dev Biol, 2012. **12**: p. 32.

63.   Andersen, L.H., et al., *Protein and carbohydrate composition of larval food affects*
tolerance to thermal stress and desiccation in adult Drosophila melanogaster. J
Insect Physiol, 2010. **56**(4): p. 336-40.

64.   Sisodia, S. and B.N. Singh, *Experimental evidence for nutrition regulated stress*
resistance in Drosophila ananassae. PLoS One, 2012. **7**(10): p. e46131.

65.   Bak, N.K., et al., *The Role of Genetic Variation in Shaping Phenotypic Responses to*
Diet in Aging Drosophila melanogaster. bioRxiv, 2025: p.
2025.01.09.632132.
747
748

Figures and Tables

Figure 1: Correlations in log2-fold changes across treatments for genes classified as showing: only a diet effect (Diet) or only a mating effect (Mating) (first row of panels), an additive effect of both (Diet + Mating, second row) or a diet-by-mating interaction (Diet × Mating, third row). Fold changes in the left column are from the carbohydrate-rich to protein-rich diet and shown between virgins (x-axis) and mated females (y-axis), fold changes in the right column are from virgins to mated females and shown between the protein-rich diet (x-axis) and the carbohydrate-rich diet (y-axis). Orange dashed lines indicate zero-fold change for the two dimensions, blue and red diagonal lines designate equal fold change in the two conditions (slope = 1).

761
762

763
764 **Figure 2:** GO enrichment for the transcriptomic response to changes in diet and mating
765 status. Enrichment analyses for 'biological process' were performed for all gene categories
766 indicated on the x-axis, and p-values were adjusted for FDR < 0.05 ('p.adjust').
767

768
769 **Figure 3:** Boxplots of heat knockdown response (time to knock down, y-axis) of mated (red)
770 and virgin (blue) females fed a protein-rich or a carbohydrate-rich diet (x-axis). Bold lines
771 indicate medians, boxes the interquartile range, whiskers the most extreme values within 1.5
772 interquartile ranges. Dots indicate data points beyond this range.
773
774
775

776 **Table 1:** Numbers of genes that do or do not show significant differential expression in
777 response to Mating, Diet, and their interaction in males and females. Based on the
778 significance of the three predictor terms, genes were classified into four functional categories
779 (last column).
780

significance (FDR<0.05)

mating	diet	mating × diet	n. genes male	n. genes female	category
-	-	-	9723	8263	
Y	-	-	0	345	mating
-	Y	-	139	651	diet
Y	Y	-	0	110	diet + mating
-	-	Y	0	3	diet × mating
-	Y	Y	0	9	diet × mating
Y	Y	Y	0	2	diet × mating

781
782
783
784

785 **Table 2:** Gene overlap analysis between expression categories for female-expressed genes
786 obtained from our current analysis (see Tab. 1) and categories defined in previous work
787 investigating sex-specific expression responses to diet (Camus et al. 2019, ref. [4]).
788

		Current Female-specific Dataset			
		Diet 651	Mating 345	D + M 110	D × M 14
Camus et al. (2019)	Diet 639	**305**	19	**47**	**7**
	Diet × Sex 51	**15**	5	**5**	0
	D + (D × S) 116	**70**	2	**31**	**5**
	f. reproductive 165	8	**45**	0	0

789
790
791
792

793 **Table 3:** Enriched transcription factor binding motifs for each of the four expression
794 categories defined in Table 1.
795

	Category	Transcription factor prediction
Male	Diet	GATA family (GATAd, GATAe, grn, pnr, srp)
Female	Diet	GATA family (GATAd, GATAe, grn, pnr, srp)
	Mating	Hsf, Stat92E, Syb, ftz-f1, Blimp-1,
	Diet + mating	GATA family (GATAd, GATAe, grn, pnr, srp)
	Diet × mating	Hnf4, Est96B, Hr4, CG7786, bigmax