Dietary sulfur amino acid restriction improves glucose homeostasis through hepatic *de novo* serine synthesis

Andres F. Ortega[1], Cha Mee Vang[1], Ferrol I. Rome[1], Kaitlyn M. Andreoni[2], Aiden M. Phoebe[2],

Alisa B. Nelson[1], Peter A. Crawford[1,3], James J. Galligan[2], Stanley Ching-Cheng Huang[4,5],

Curtis C. Hughey[1]*

[1]Division of Molecular Medicine, Department of Medicine, University of Minnesota, Minneapolis, MN

[2]Department of Pharmacology and Toxicology, University of Arizona, Tucson, AZ

[3]Department of Biochemistry, Molecular Biology, and Biophysics, University of Minnesota, Minneapolis, MN

[4]Pelotonia Institute for Immuno-oncology, Comprehensive Cancer Center – James Cancer Hospital and Solove Research Institute, The Ohio State University, Columbus, OH

[5]Department of Microbial Infection and Immunity, College of Medicine, The Ohio State University, Columbus, OH

Running Title: PSAT1 is needed for glucose responses to SAAR

*To whom the correspondence should be addressed: Curtis C. Hughey from the Department of Medicine, Division of Molecular Medicine, University of Minnesota, Dwan 316, 425 East River Parkway, Washington Ave SE, Minneapolis, MN, 55455. Tel.: 612-301-2677, Email: chughey@umn.edu

ABSTRACT

Dietary sulfur amino acid restriction (SAAR) improves whole-body glucose homeostasis, elevates liver insulin action, and lowers liver triglycerides. These adaptations are associated with an increased expression of hepatic *de novo* serine synthesis enzymes, phosphoglycerate dehydrogenase (PHGDH) and phosphoserine aminotransferase 1 (PSAT1). This study tested the hypothesis that enhanced hepatic serine synthesis is necessary for glucose and lipid adaptations to SAAR. Hepatocyte-specific PSAT1 knockout (KO) mice and wild type (WT) littermates were fed a high-fat control or SAAR diet. In WT mice, SAAR increased liver PSAT1 protein (~70-fold), serine concentration (~2-fold), and ^{13}C-serine (~20-fold) following an intravenous infusion of [U-^{13}C]glucose. The elevated liver serine and partitioning of circulating glucose to liver serine by SAAR were attenuated in KO mice. This was accompanied by a blunted improvement in glucose tolerance in KO mice fed a SAAR diet. Interestingly, SAAR decreased liver lysine lactoylation, a SAA-supported post-translational modification known to inhibit PHGDH enzymatic activity. This suggests dietary SAAR may increase serine synthesis, in part, by lowering lysine lactoylation. Beyond glucose metabolism, dietary SAAR reduced body weight, adiposity, and liver triglycerides similarly in WT and KO mice. Collectively, these results demonstrate that hepatic PSAT1 is necessary for glucose, but not lipid, adaptations to SAAR.

ARTICLE HIGHLIGHTS

- Dietary SAAR increases glucose partitioning to liver *de novo* serine synthesis

- Deletion of liver PSAT1 abrogates improvements in glucose tolerance by SAAR.

- SAAR lowers lysine lactoylation, a SAA-supported post-translational modification known to inhibit serine synthesis enzymes.

- Hepatic PSAT1 is not necessary for SAAR to lower adiposity and liver triglycerides.

INTRODUCTION

Dietary sulfur amino acid restriction (SAAR) provokes metabolic adaptations that are advantageous in preventing and treating obesity and its comorbidities (1). In rodents, SAAR enhances glucose and insulin tolerance (2-7). The use of hyperinsulinemic-euglycemic clamps combined with isotope tracers revealed that the increased insulin action is due to a greater suppression of endogenous glucose production and heightened glucose disposal by skeletal muscle, heart, and adipose tissue (8-11). Concurrently, SAAR decreases adiposity and liver triglycerides (1–4). It is important to note that these metabolic responses to SAAR are not confined to rodents. Short-term dietary SAAR in humans elicits decreases in body weight, body mass index (BMI), and liver fat (12-14). The SAAR-mediated adaptations are partly due to a liver-derived increase in fibroblast growth factor 21 (FGF21) (1). The transcriptional activation of hepatic FGF21 increases whole-body energy expenditure, which contributes to reduced adiposity by SAAR (9). FGF21 is necessary for the enhanced insulin-stimulated cardiac glucose uptake (10, 11). In contrast, SAAR lowers liver triglycerides similarly in wild type (WT) and FGF21-deficient mice (9). Prior work suggests the enhanced liver insulin signaling by SAAR is also independent of FGF21 (1; 11). Together, these findings indicate that the mechanisms underlying the liver glucose and lipid adaptations to SAAR, and the extent to which these responses are hepatocyte-dependent, remain to be clearly defined.

Elevated expression of liver *de novo* serine synthesis pathway enzymes, phosphoglycerate dehydrogenase (PHGDH) and phosphoserine aminotransferase 1 (PSAT1), is among the most reproducible responses to dietary SAAR, making their increase a marker of a successful dietary intervention (2; 9; 10; 15-23). *De novo* serine synthesis is sourced from 3-phosphoglycerate, an intermediate of both glycolysis and gluconeogenesis (24). Thus, a SAAR-mediated increase in

hepatic serine synthesis could promote greater glucose disposal and/or divert gluconeogenic flux

away from glucose production. Liver-specific PHGDH knockout (KO) mice fed a chow diet show

higher adiposity and reduced glucose tolerance (25). In agreement with PHGDH KO mice, PSAT1

knockdown impairs glucose and insulin tolerance (26). Conversely, PSAT1 overexpression

enhances glucose and insulin tolerance (26). Beyond glucose homeostasis, the overexpression of

PHGDH has been shown to lower liver lipids (27). These prior results support that PHGDH and

PSAT1 action impacts whole-body and liver-specific glucose and lipid homeostasis.

The experiments presented herein tested the hypothesis that hepatic PSAT1 is required for

the liver glucose and lipid adaptations to SAAR to be fully realized. Liver-specific PSAT1 KO

mice and WT littermates were fed a high-fat control or SAAR diet. Comprehensive phenotyping

approaches were completed. These included glucose and insulin tolerance tests to define whole-

body glucose homeostasis, ^{13}C-isotope tracing in conscious unrestrained mice to determine the

tissue-specific fate of glucose and serine *in vivo*, and metabolomics to quantify metabolite

concentrations. Immunoblotting and proteomics were used to identify molecular drivers of liver

metabolism. The results of this study show that hepatic PSAT1 is necessary for whole-body and

liver-specific glucose, but not lipid, adaptations to dietary SAAR.

MATERIALS AND METHODS

Mouse models and husbandry

All procedures on mice were approved by the University of Minnesota Institutional Animal

Care and Use Committee. Hepatic-specific PSAT1 KO were generated by breeding mice

expressing Cre recombinase under control of the albumin promoter (B6.Cg-Tg(Alb-cre)21Mgn/J)

with PSAT1 floxed mice (28) that were backcrossed onto a C57BL/6J background for 5

generations. Floxed PSAT1 littermates lacking expression of Cre recombinase were used as WT

controls. C57BL/6J mice were used for select studies. Mice were maintained on a chow diet

(Teklad Global 18% Protein Rodent Diet 2918, Madison, WI) from weaning at three weeks of age

until six weeks of age. At six weeks of age mice received one of three diets (unless otherwise

stated): low-fat SAA control (0.85% kcal from methionine; 10% kcal from fat; LF-CTRL;

A11051302B, Research Diets Inc., New Brunswick, NJ), high-fat SAA control (0.85% kcal from

methionine; 60% kcal from fat; HF-CTRL; A11051306, Research Diets Inc., New Brunswick, NJ),

and HF-SAAR (0.125% kcal from methionine; 60% kcal from fat; A11051305, Research Diets

Inc., New Brunswick, NJ). Food and water were provided ad libitum. Mice were housed with

cellulose bedding (Cellu-nest™, Shepherd Specialty Papers, Watertown, TN) in temperature- and

humidity-controlled conditions maintained on a 14:10 hour light/dark cycle. At 12 weeks of age

(6 weeks on diet), mice underwent metabolic testing and were euthanized via cervical dislocation.

Tissues were rapidly excised, freeze-clamped in liquid nitrogen, and stored at -80°C. One cohort

of mice was euthanized at 26 weeks of age (20 weeks on diet) using the same procedures described

above. All mice used in this study were males.

Body Composition

Body composition was measured in mice fed ad libitum at 10-11 weeks of age using an

EchoMRI-100™ Body Composition Analyzer (EchoMRI LLC, Houston, TX).

Glucose and insulin tolerance tests

For glucose tolerance tests (GTT), mice were fasted for five hours and injected

intraperitoneally (i.p.) with dextrose (Hospira, Inc., Lake Forest, IL) at a dose of 2 $g \cdot kg^{-1}$ lean

weight. For insulin tolerance tests (ITT), mice were fasted for five hours and received insulin

(Humulin R human, Eli Lilly and Company, Indianapolis, IN) i.p. at a dose of 0.75 U·kg^{-1} lean

weight. During both procedures, tail-blood glucose was measured at 0, 15, 30, 45, 60, 90, and 120

minutes with a Contour blood glucose meter (Ascensia Diabetes Care, Parsippany, NJ). A cohort

of mice had blood samples obtained at 0 and 15 minutes after the insulin administration for plasma

insulin concentrations and were euthanized immediately after for assessing indices of insulin

signaling in tissues.

Insulin pellet implantation.

At 7 weeks of age, C57BL/6J mice had one sustained-release insulin pellet implanted

subcutaneously (~0.1 U/day per implant; LinBit; LinShin Canada, Toronto, ON) according to

manufacturer guidelines. Briefly, under isoflurane anesthesia, a 1 cm diameter area of hair was

clipped and disinfected. A small incision in the skin was made with a 16-gauge needle, and a sterile

12-gauge trocar was used to deliver the pellet subcutaneously. Procedures in control mice were

identical, except the skin incisions were not performed. Mice were group housed following the

procedure.

Vascular catheter surgeries

At 11 weeks of age, catheters were surgically implanted in the jugular vein and carotid

artery for isotope infusion and sampling protocols as previously described (21; 29). The

exteriorized ends of the implanted catheters were flushed with 200 U·mL^{-1} heparinized saline and

sealed with stainless-steel plugs. Following surgery, mice were housed individually and provided

5-7 days of post-operative recovery prior to stable isotope infusion studies. All mice were within

10% of pre-surgical weight prior to stable isotope infusions.

Stable-isotope tracing: [U-^{13}C]glucose intravenous infusion in the conscious, unrestrained mouse

During the first hour of the housing light cycle, both food and water were withdrawn for

the duration of the experiment (5.5 hours). Four hours into the fast, the exteriorized vascular

catheters were connected to infusion syringes. Following a one-hour acclimation, a 60 mg·kg^{-1}·min^{-1}

continuous intravenous infusion of [U-^{13}C]glucose was performed for 30 minutes.

C57BL/6J mice had blood glucose measured prior to and every 10 minutes during the [U-

^{13}C]glucose infusion to ensure the infusion rate was sufficient to achieve glucose concentrations

comparable to those observed during a GTT. The experiments in PSAT1 WT and KO mice did not

measure blood glucose. Mice were euthanized and tissue collected immediately following the [U-

^{13}C]glucose infusion.

Stable-isotope tracing: In-vivo [U-^{13}C]serine intraperitoneal injection

Food and water were withdrawn for 5 hours starting within the first hour of the light cycle.

[U-^{13}C]serine was administered i.p. at a dose of 100 mg·kg^{-1}. Fifteen minutes following the i.p.

injection, mice were euthanized and tissues harvested.

Hormone and metabolite analysis

Plasma insulin was determined using the Mercodia Ultrasensitive Mouse Insulin ELISA

(Mercodia AB, Winston Salem, NC). Liver triglycerides were measured with the Triglycerides -

Liquid Reagent Set (Pointe Scientific, Inc., Lincoln Park, MI). Liver glycogen was quantified as

previously described (21; 30). All other metabolites were measured using untargeted metabolomics

as detailed below. Tissue metabolite levels were quantified from mice that were fasted 7 hours.

Untargeted metabolomics and ^{13}C-isotope tracing untargeted metabolomics (^{13}C-ITUM) pipeline

Extraction and analytical methods were completed as previously described with minor

modifications (31-33). Briefly, metabolites were separated using a Thermo Fisher Scientific

Vanquish UHPLC system and an Atlantis Premier UPLC BEH Z-HILIC column (186009979,

Milford, MA). Mobile Phase A was 90% water with 15 mM ammonium bicarbonate and Mobile

Phase B was 90% ACN with 15 mM ammonium bicarbonate. The total run time was 10 minutes,

flow rate was 0.5 mL·min^{-1}, and column chamber temperature was 30°C. Separated metabolites

were detected using a Thermo Fisher Scientific QExactive Plus hybrid quadrupole-orbitrap mass

spectrometer, fitted with a heated electrospray ionization source operated in negative and positive

polarity mode. Two sample types were run to aid in chemical feature identification: 1) a standard

mix consisting of authentic standards for all expected analytes and 2) a pooled sample containing

only the naturally occurring samples, analyzed via data-dependent analysis (DDA)–tandem mass

spectrometry (MS/MS), using IE omics script and R-Studio (34). Data processing and initial

analysis were performed using Thermo Compound Discoverer 3.3.

Targeted analysis of glyoxalase cycle intermediates

Reduced glutathione (GSH), lactoylglutathione (LGSH), and hemithioacetal (HTA) were

quantified in liver as previously described (35). Briefly, samples were prepared in a sulfosalicylic

acid solution containing the internal standard, GSH-(Gly-^{13}C$_2$,^{15}N). The protein precipitate was

removed via centrifugation and supernatants chromatographed using a Shimadzu LC system

equipped with an Atlantis C$_{18}$ column (186001291; Waters, Milford, MA). Multiple reaction

monitoring-mass spectrometry (MRM-MS) was performed in positive ion mode using an AB

SCIEX 6500+ QTRAP.

Quantitative analysis of arginine and lysine modifications

Lysine lactoylation (lactoyllys), lysine acetylation (acetyllys), methylglyoxal-derived

hydroimidazolone 1 (MG-H1), and carboxyethylarginine (CEA) were determined as previously

described (35; 36). Liver protein was precipitated and resuspended in ammonium bicarbonate

containing internal standards. Samples were then digested to single amino acids, acidified with

heptofluorobutyric acid, chromatographed using a Shimadzu LC system equipped with an Eclipse

XDB-C8 column (930990-906; Agilent, Santa Clara, CA), and analyzed by MRM-MS. The post-

translational modifications (PTMs) were normalized to Leu to control for digestion (36; 37).

Immunoblotting

Tissue homogenates were prepared as previously detailed (21; 29). Tissue proteins (15 µg)

were separated by electrophoresis on a NuPAGE 4-12% Bis-Tris gel (Invitrogen, Carlsbad, CA,

USA) and transferred to a PVDF membrane. Immunoblotting was performed with the primary

antibodies and secondary antibodies listed in Supplementary Table 1. The PVDF membranes were

treated with a chemiluminescent substrate (Thermo Fisher Scientific, Waltham, MA, USA) and

imaged using a ChemiDoc Imaging system and Image Lab software (Bio-Rad, Hercules, CA,

USA). Total protein was assessed via BLOT-FastStain (G-Bioscience, St. Louis, MO, USA) and

used as the loading control. Densitometry was evaluated using ImageJ software.

Statistical Analysis

GraphPad Prism software (GraphPad Software LLC., San Diego, CA) was used to perform

Student's t-test, two-way ANOVAs followed by Tukey's post-hoc tests, and repeated measures

ANOVAs followed by Šidák's post-hoc tests as appropriate to detect statistical differences

($p < 0.05$). All data are reported as mean ± SEM.

RESULTS

Elevated glucose partitioning to liver serine by SAAR is linked to improved glucose tolerance

In C57BL/6J mice, 6 weeks of dietary intervention showed a greater gain in body weight,

higher adiposity, and similar lean mass in the HF-CTRL compared to LF-CTRL group (Fig. 1A

and B). Mice fed a HF-SAAR diet had a lower body weight, decreased fat mass, and increased

lean mass percentage relative to both the LF-CTRL and HF-CTRL mice (Fig. 1B). A HF-CTRL

diet elevated liver triglycerides compared to mice fed the LF-CTRL diet, while the HF-SAAR diet

prevented the increase in liver triglycerides and lowered liver weight (Fig. 1C). Dietary SAAR

also lowered liver glycogen and fasting blood glucose (Fig. 1D and E). HF-CTRL mice showed

impaired glucose tolerance, but not insulin tolerance, compared to LF-CTRL mice (Fig. 1F and

G). Conversely, glucose and insulin tolerance were improved in mice fed the HF-SAAR diet

relative to mice fed the LF-CTRL and HF-CTRL diets (Fig. 1F and G). Plasma insulin was

significantly reduced in HF-SAAR mice (Fig 1H). This is notable because chronically elevated

insulin has been proposed to promoted gain in adiposity and dysregulated glucose control (38-40).

To test if the lowered circulating insulin levels in HF-SAAR mice was contributing to the

metabolic adaptations to SAAR, a subcutaneous, sustained-release insulin pellet was implanted in

HF-SAAR mice to raise insulin. The persistent delivery of exogenous insulin from the

subcutaneous pellet over 5 weeks did not impact the glucose and lipid adaptations to SAAR (Suppl.

Fig. 1). This suggests that the mechanisms underlying SAAR are independent of circulating insulin

concentrations.

Untargeted metabolomics revealed that liver cysteine was higher in HF-CTRL compared

to LF-CTRL mice (Fig. 2A and 2B). As expected, liver cysteine and taurine were decreased ~5-

fold in mice fed the HF-SAAR diet (Fig. 2B). Liver PSAT1 protein (Fig. 2C) and serine

concentrations (Fig. 2E) were lower in HF-CTRL versus LF-CTRL. In contrast, the HF-SAAR

diet led to a markedly increased in liver PHGDH (~5-fold) and PSAT1 (~70-fold) protein

expression (Fig. 2D). The elevated expression of serine synthesis enzymes was associated with

higher liver serine and glycine levels in HF-SAAR relative to both CTRL diet groups (Fig. 2E).

To test if the increased liver serine and glycine were being sourced from circulating glucose, [U-

^{13}C]glucose was intravenously infused in conscious, unrestrained mice fed a LF-CTRL or a LF-

SAAR diet for 6 weeks (Fig. 2F). ^{13}C-labeling of liver serine and glycine was significantly higher

in LF-SAAR mice compared to LF-CTRL, with no alterations observed in labeled glucose (Fig.

2H). Collectively, our data show that dietary SAAR improves glucose and insulin tolerance, which

is associated with increased partitioning of circulating glucose to liver serine synthesis.

Liver PSAT1 is required for SAAR-induced improvements in systemic glucose homeostasis

To assess whether hepatic serine synthesis is essential for the metabolic benefits of dietary

SAAR on glucose control, we studied liver-specific PSAT1 KO mice and WT littermates fed a

HF-CTRL or -SAAR diet (Fig. 3A). The HF-SAAR diet prevented a gain in body weight and

adiposity in both genotypes (Fig. 3B and C). When fed a HF-CTRL diet, WT and liver PSAT1 KO

mice showed similar glucose tolerance (Fig. 3D). This was expected given that high-fat feeding

lowers liver PSAT1 in WT mice (Fig. 2D), a phenotype closely resembling liver PSAT1 KO mice.

However, on a HF-SAAR diet, liver-specific PSAT1 KO mice showed a greater excursion of

glucose during the glucose tolerance test compared to WT mice (Fig. 3D). Loss of hepatic PSAT1

in mice fed HF-CTRL diet delayed the normalization of circulating glucose during an insulin

tolerance test relative to WT littermates (Fig. 3E). Six weeks of HF-SAAR feeding improved

insulin tolerance similarly in WT and KO mice compared to HF-CTRL groups (Fig. 3E).

Concordant with the C57BL/6J findings (Fig. 1), HF-SAAR feeding lowered plasma insulin levels

relative to mice fed a HF-CTRL diet in both genotypes (Fig. 3F). Fifteen minutes after an insulin

injection, the liver pAkt-to-Akt ratio was higher in both genotypes fed the HF-SAAR diet (Fig.

3G). This suggests liver insulin signaling was enhanced similarly in WT and KO mice following

SAAR. Dietary SAAR lowered liver glycogen in WT and liver PSAT1 KO mice (Fig. 3H).

However, this was independent of changes key enzymes regulating glycogen synthesis and

breakdown. Specifically, total protein and the phospho-to-total protein ratio for glycogen synthase

(GS) and glycogen phosphorylase (PYGL) were similar between groups (Fig. 3I). Together, these

data indicate that liver PSAT1 is necessary for dietary SAAR to improve whole-body glucose

tolerance.

Increased glucose partitioning to serine following SAAR is abolished by loss of hepatic PSAT1

Next, we aimed to further connect the attenuated improvement in glucose tolerance

observed in liver PSAT1 KO mice fed a HF-SAAR diet to inability of hepatocytes to synthesize

serine from glucose (Fig. 4A). The HF-SAAR diet elevated serine and glycine synthesis enzymes,

PHGDH, PSAT1 (WT mice only), and serine hydroxymethyltransferase 2 (SHMT2), in the livers

of mice (Fig. 4B). Under post-absorptive conditions, liver glucose, serine, and glycine were

comparable between WT and KO mice fed a HF-CTRL diet (Fig. 4C). Dietary SAAR lowered

liver glucose in WT, but not liver PSAT1 KO mice (Fig. 4C). Liver serine and glycine were higher

in WT mice fed a HF-SAAR diet compared to WT mice on a HF-CTRL diet (Fig. 4C); however,

the increase in liver serine and glycine by SAAR was blunted in liver PSAT1 KO mice (Fig. 4C).

Consistent with these metabolite levels, a HF-SAAR diet prompted a rise in the ^{13}C-labeling of

liver serine and glycine in WT mice intravenously infused with [U-^{13}C]glucose (Fig. 4D). Loss of

liver PSAT1 attenuated the increased in serine and glycine ^{13}C-labeling by SAAR (Fig. 4D). These

data imply that the improved glucose tolerance by dietary SAAR is associated with enhanced

partitioning of circulating glucose to liver serine synthesis.

Serine sourced from glucose following SAAR is preferentially fated to glutathione

Serine is a precursor for cysteine synthesis via transsulfuration and donates a one-carbon

units to support methionine regeneration within the methionine cycle (41). Given this, one could

speculate that the elevated diversion of glucose to serine is an adaptive response to prevent

limitations in SAA availability that may arise from dietary SAAR. To further test this hypothesis,

liver proteins, metabolite concentrations, and metabolite labeling from [U-^{13}C]glucose and [U-

^{13}C]serine administration were determined (Fig. 5). Regardless of diet, loss of hepatic PSAT1 did

not impact proteins involved in transsulfuration or cysteine catabolism (Fig. 5A and 5B). Dietary

SAAR increased cystathionine γ-lyase (CSE; involved in cysteine production), decreased cysteine

dioxygenase 1 (CDO1; involved in cysteine catabolism to taurine), and elevated GSH synthesis

enzymes, glutamate-cysteine ligase catalytic subunit (GCLC) and glutathione synthetase (GSS),

in both genotypes (Fig. 5A and 5B). SAAs and related metabolites were comparable between WT

and liver PSAT1 KO mice fed a HF-CTRL diet (Fig. 5C). The HF-SAAR diet decreased liver

cysteine, taurine, GSH, and oxidized glutathione (GSSG) in both genotypes compared to their

counterparts fed the HF-CTRL diet (Fig. 5C). Consistent with metabolite levels, ^{13}C-labeling of

liver SAA and related metabolites following a 30-minute [U-^{13}C]glucose infusion was similar

between WT and Liver PSAT1 KO mice on a HF-CTRL diet (Fig. 5D). HF-SAAR feeding

increased ^{13}C-labeling from glucose in liver methionine in both WT and liver PSAT1 KO mice

(Fig. 5D). Fractional enrichment in liver cysteine was not detectable due to the low cysteine pool

in mice fed a HF-SAAR diet; however, downstream fates of cysteine were measured. The [U-

^{13}C]glucose infusion showed lower ^{13}C-taurine, and increased ^{13}C-GSH and ^{13}C-GSSG in the

livers of both WT and KO mice fed the HF-SAAR diet relative to the HF-CTRL mice (Fig. 5D),

with the SAAR-mediated increase in GSH and GSSG ^{13}C-labeling from [U-^{13}C]glucose being

blunted in liver PSAT1 KO mice compared to their WT littermates (Fig. 5D). To more directly

assess the impact of dietary SAAR on serine fate, [U-^{13}C]serine was administered to WT mice fed

a HF-CTRL or a HF-SAAR diet for 6 weeks (Fig. 5E). ^{13}C-labeling was increased in liver

methionine, taurine, GSH, and GSSG in the HF-SAAR mice compared to the HF-CTRL (Fig. 5E).

The rise in ^{13}C-labeling was particularly pronounced for GSH (~12-fold) and GSSG (~37-fold)

(Fig. 5E). Together, these results suggest that serine synthesized *de novo* from glucose is important

for supporting GSH provision during SAAR.

SAAR reduces glyoxalase cycle intermediates and associated protein PTMs

To further explore the potential connectivity between serine synthesis and GSH

availability, we assessed markers of glyoxalase cycle metabolism (Fig. 6A). GSH can be used to

generate LGSH in the glyoxalase cycle and recent work identified LGSH-derived lysine

lactoylation inhibits PHGDH (42). Liver glyoxalase cycle enzymes, glyoxalase 1 (GLO1) and

hydroxyacylglutathione hydrolase (GLO2), were comparable between genotypes on both diets

(Fig. 6B). However, SAAR decreased liver GLO1 protein (Fig. 6B). Consistent with data from our

untargeted metabolomic approaches (Fig. 5C), targeted quantification showed lower liver GSH in

WT and KO mice fed a HF-SAAR diet compared to the HF-CTRL diet (Fig. 6C). Dietary SAAR

also decreased liver LGSH and HTA in both genotypes (Fig 6C). Liver lactate was lower in liver

PSAT1 KO mice relative to WT mice fed HF-CTRL (Fig. 6C). SAAR decreased liver lactate in

WT, but not liver PSAT1 KO mice (Fig. 6C). PTMs derived from glyoxalase cycle metabolites

were broadly suppressed by SAAR. Specifically, lysine lactoylation was decreased in HF-SAAR

WT mice compared to WT HF-CTRL (Fig. 6D). In contrast, lysine lactoylation was comparable

between liver PSAT1 KO mice fed the HF-CTRL and -SAAR diets (Fig. 6D). Methylglyoxal

(MGO)-derived modifications, MG-H1 and CEA, were significantly lower in HF-SAAR- versus

HF-CTRL-fed mice regardless of genotype (Fig. 6D). Global lysine acetylation was not changed

by diet or genotype (Fig. 6D). Overall, SAAR reduces glyoxalase-cycle intermediates and the

protein modifications they support.

SAAR diverts glucose toward asparagine synthesis

Given that loss of hepatic PSAT1 blunted, but did not completely abolish the improvement

in glucose tolerance by SAAR, we further examined how SAAR could be influencing glucose

metabolism in the liver. On a HF-CTRL diet, liver TCA cycle, anaplerotic, and cataplerotic

metabolites were mostly comparable between WT and KO mice (Fig. 7A and B). SAAR increased

pyruvate, decreased malate (WT mice only), and elevated aspartate (Fig. 7B). [U-^{13}C]glucose

tracing showed lower ^{13}C-pyruvate, increased ^{13}C-aspartate, and significantly elevated ^{13}C-

asparagine (~370-fold) labeling in WT and KO mice on the HF-SAAR diet compared to the HF-

CTRL diet (Fig. 7C). Interestingly, SAAR lowered the ^{13}C-labeling of liver succinate in WT mice

(Fig. 7C). This decreased ^{13}C-labeling of liver TCA cycle intermediates was more pronounced in

PSAT1 KO mice on a HF-SAAR diet as indicated by lower ^{13}C-citrate, ^{13}C-succinate, and ^{13}C-

malate (Fig. 7C). Next, we assessed three isotopologue ratios to provide indices of nutrient flux

(43-48). Livers of WT mice fed the HF-SAAR diet had a higher citrate M+3-to-pyruvate M+3

ratio relative to WT mice fed a HF-CTRL and KO mice fed a HF-SAAR diet (Fig. 7D). This

suggests that TCA cycle anaplerosis via pyruvate carboxylase (PC) was stimulated by SAAR. The

aspartate M+3-to-succinate M+3 and aspartate M+3-to-citrate M+3 ratios were elevated in HF-

SAAR mice compared to their HF-CTRL counterparts (Fig. 7D). The increase in these two ratios

suggest a preferential partitioning of pyruvate to aspartate compared to flux through the TCA

cycle. Consistent with the metabolite and ^{13}C-isotope tracing data, liver asparagine synthetase

(ASNS) protein was ~25-fold higher in mice fed a HF-SAAR diet compared to those receiving the

HF-CTRL diet (Fig. 7E). Most TCA cycle proteins tested were not altered by diet or genotype

(Fig. 7E). The cataplerotic enzyme, mitochondrial glutamic-oxaloacetic transaminase (GOT2),

was increased in PSAT1 KO mice fed a HF-SAAR diet compared to both their WT littermates and

KO mice fed HF-CTRL (Fig. 7E). These results indicate that SAAR shifts the fate of glucose, via

the TCA cycle anaplerosis, towards liver asparagine synthesis.

SAAR prevents liver lipid accumulation independently of hepatic PSAT1

In addition to glucose control, we investigated the role of hepatic PSAT1 action in liver

lipid adaptations to dietary SAAR. After 6 weeks of diet, liver triglycerides were lower in WT

mice fed a HF-SAAR diet compared to WT mice on the HF-CTRL diet (Fig. 8A). This dietary

effect was not observed in liver PSAT1 KO mice (Fig. 8A). The expression of liver *de novo*

lipogenesis enzymes, ATP-citrate lyase (ACLY) and stearoyl-CoA desaturase 1 (SCD1), were

suppressed with HF-SAAR feeding in both genotypes (Fig. 8B and D). SAAR decreased FASN

protein in WT mice only (Fig. 8B and D). As we previously observed, SAAR increased liver

mitochondrial respiratory proteins (Fig. 8C and D). Specifically, complex I (NADH

dehydrogenase (ubiquinone) iron-sulfur protein 4; NDUFS4) and complex V (ATP synthase F1

subunit alpha; ATP5A) were higher in WT mice fed HF-SAAR compared to WT mice fed the HF-

CTRL diet. (Fig. 8C and D). Complex IV (mitochondrial encoded cytochrome c oxidase subunit

II; MTCO2) protein was elevated in PSAT1 KO mice on the HF-SAAR diet versus KO mice fed

the HF-CTRL diet. An additional mechanism through which SAAR may lower liver triglycerides

is the formation of acylglycines. Protein expression of liver glycine N-acyltransferase (GLYAT),

which catalyzes the conjugation of acyl-CoA to glycine, was similar between groups (Fig. 8F).

However, HF-SAAR raised the levels of short-chain acylglycines (acetylglycine,

isobutyrylglycine, and isovalerylglycine) in the liver (Fig. 8G).

To further test if hepatic PSAT1 action was required for SAAR to lower liver triglycerides,

mice were fed the HF-SAAR diet for 20 weeks. Both WT and KO mice fed the HF-SAAR diet

had lower body weight, fat mass, and liver triglycerides compared to mice on the HF-CTRL diet

(Fig. 8H-J). This suggest that hepatic PSAT1 is not needed, particularly with increasing duration,

for SAAR to promotes a decline in liver triglycerides.

DISCUSSION

Dietary SAAR prevents obesity, dysregulated glucose control, and liver steatosis (1).

Analysis of changes at the level of the transcriptome has identified that the liver is particularly

responsive to SAAR (1). These changes include an increased expression of liver *de novo* serine

synthesis pathway enzyme, PSAT1, which is among the most reproducible responses to dietary

SAAR (2; 9; 10; 15-23). Given the interconnectedness of *de novo* serine synthesis with glycolysis

and gluconeogenesis pathways, this study tested the hypothesis that hepatic PSAT1 is needed for

the liver glucose and lipid adaptations to SAAR. The pertinent findings of this study are: 1)

enhanced hepatic *de novo* serine synthesis is required for SAAR to improve glucose tolerance, 2)

concurrent with transcriptional activation, a serine synthesis-GSH-lysine lactoylation feedback

loop may underlie the increased hepatic serine synthesis in response to SAAR, and 3) hepatic

serine synthesis is not necessary for the SAAR-mediated decline in adiposity and liver

triglycerides.

Role of hepatic PSAT1 in glucose adaptations to SAAR

Consistent with previous studies using rodents (2-7), we found that dietary SAAR

enhances glucose tolerance. Untargeted metabolomics showed this whole-body glucose phenotype

was associated with increased liver serine. Exogenous serine supplementation can (49), but does

not always (50), improve glucose tolerance. Thus, the SAAR-mediated increase in serine may

actively promote glucose tolerance via one or more downstream functions (51). Our [13]C-isotope

tracing experiments showed elevated partitioning of circulating glucose to hepatic serine following

SAAR. Moreover, loss of hepatic PSAT1 impaired glucose tolerance in mice fed the SAAR diet

and abolished the increase in liver serine sourced from circulating glucose without completely

preventing the rise in liver serine levels. These results suggest that a higher rate of hepatic glucose

disposal and the diversion of glucose to the synthesis of serine is, in part, responsible for the

SAAR-mediated adaptations in whole-body glucose control.

PSAT1 overexpression in mice enhances both glucose and insulin tolerance (26). In the

experiments presented here, we found that the heightened insulin tolerance following SAAR was

not impacted by the loss of hepatic PSAT1. Prior work has reported that the enhanced insulin

action in response to PSAT1 overexpression is mediated by decreased expression of tribbles

homolog 3 (TRB3), which is a pseudokinase that inhibits insulin signaling by binding to Akt and

impairing its activation by phosphorylation (26). Our studies showed the phosphorylation of liver

Akt was elevated similarly in both WT and liver PSAT1 KO mice. Together, these data suggest

the elevated liver PSAT1 increases glucose tolerance independent of insulin action. However,

future work should assess the role of liver PSAT1 on liver insulin action during hyperinsulinemic-

euglycemic clamps combined with isotope tracers. These conditions would allow for assessing

liver-specific insulin sensitivity under euglycemic conditions so that the impact of SAAR on

extrahepatic glucose flux and concentrations do not mask alterations at the liver.

Potential mechanisms promoting increased hepatic de novo serine synthesis from glucose

Work by independent groups indicate that SAAs negatively regulate serine synthesis.

Cysteine suppresses the expression of PHGDH by decreasing mRNA stability (52). In addition to

post-transcriptional regulation, mechanisms at the level of transcription may also contribute to the

enhanced serine synthesis by SAAR. Deletion of the transcription factor, nuclear factor erythroid

2-related factor 2 (NRF2), in the livers of mice blunted but did not abolish the increase in PSAT1

mRNA (16). These mechanisms of action are consistent with the elevated liver PHGDH and

PSAT1 mRNA and protein in response to SAAR. Serine is a precursor for cysteine, which is

generated via transsulfuration wherein the sulfur from methionine is transferred to serine (41). It

may be that increased cysteine impedes the synthesis of its precursor, serine, to control SAA levels.

Conversely, when cysteine is low, serine synthesis is stimulated to promote SAA availability.

Ultimately, these findings suggest that SAAR-mediated increase in glucose tolerance is the result

of liver responses to maintain SAA homeostasis via serine synthesis.

A prominent fate of methionine and cysteine, particularly under conditions of low SAA

levels, is the synthesis of GSH (53-56). Aligning with prior studies (23), we observed liver GSH

was decreased by SAAR. However, our [U-^{13}C]glucose and [U-^{13}C]serine isotope tracing

experiments showed that ^{13}C-labeling of liver GSH and GSSG was elevated in mice fed the SAAR

diet. While the [U-^{13}C]serine tracing showed SAAR increased labeling of other fates of cysteine

such as taurine, loss of hepatic PSAT1 uniquely attenuated the increased liver ^{13}C-GSH and ^{13}C-

GSSG from [U-^{13}C]glucose. These results indicate that serine synthesized *de novo* from glucose

is important to support GSH provision (via cysteine) during SAAR.

Beyond its antioxidant properties, GSH is a metabolite within the glyoxalase cycle. This

highly conserved metabolic cycle comprised of two enzymes, GLO1 and GLO2, uses GSH to

catabolize MGO to lactate through the intermediate, LGSH (57). LGSH is an acyl-donor in a non-

enzymatic reaction with free lysine residues, yielding the PTM, lysine lactoylation (35; 58). This

is notable because PHGDH and PSAT1 are heavily lactoylated and LGSH inhibits PHGDH

activity (35; 42). Our results showed that SAAR lowered liver LGSH and lysine lactoylation.

Together, our data and the work of others suggest that SAAR may increase serine synthesis by

preventing the inhibition of PHGDH and/or PSAT1 by LGSH-derived lysine lactoylation.

Consistent with GSH regulating serine synthesis via post-translational mechanisms, the

pharmacologic inhibition of GSH synthesis using buthionine sulfoximine increased liver serine

without altering NRF2 or PHGDH expression (22). While it remains to be fully tested, SAAR may

promote glucose tolerance through a hepatic serine synthesis-GSH-lysine lactoylation feedback

loop in which low SAA decreases liver GSH, lowers lysine lactoylation, and thereby prevents

inhibition of *de novo* serine synthesis from glucose.

SAAR shifts the fate of glucose towards liver asparagine

In addition to *de novo* serine synthesis, this study demonstrates that SAAR stimulates

glucose partitioning towards asparagine synthesis. Liver asparagine concentration was similar

between mice fed the HF-CTRL and SAAR diets. However, ^{13}C-labeling of liver asparagine from

[U-^{13}C]glucose increased markedly by ~370-fold in SAAR-fed mice. We also assessed indices of

nutrient flux (43-48). Isotopologue ratios from our [U-^{13}C]glucose tracing experiments are

consistent with SAAR increasing TCA cycle anaplerosis through PC (i.e., the generation of OAA

from pyruvate) and the resulting OAA being preferentially directed to asparagine rather than flux

through the TCA cycle. The SAAR-induced shift in glucose and TCA cycle intermediates to

asparagine was associated with a ~25-fold increase in liver ASNS, which generates asparagine

from aspartate. SAAR did not alter the expression of other anaplerotic and cataplerotic enzymes

(i.e., PC, MDH2, PCCA, and GOT2) linked to asparagine synthesis. Interestingly, these

aforementioned enzymes have been reported to be modified by lysine lactoylation (35). Thus,

further research is warranted to determine if the SAAR-driven reduction in lactoylation prevents

the inhibition of these enzymes and favors the diversion of glucose to asparagine. Future studies

should also test whether the stimulation glucose partitioning to asparagine is an additional means

through which SAAR increases glucose tolerance.

The shift in glucose fate to asparagine during SAAR could be an additional means of

enhancing serine availability. Intracellular asparagine exchanges with extracellular serine to

promote serine uptake (59). Liver PSAT1 KO mice receiving a HF-SAAR diet showed higher liver

serine levels compared to KO mice fed a HF-CTRL diet. This was accompanied by a trend towards

greater [13]C-labeling of liver serine [U-[13]C]glucose in KO mice fed a HF-SAAR diet compared to

KO mice fed a HF-CTRL diet. To further explore if the liver was acquiring this serine from

extrahepatic sources, we administered [U-[13]C]serine to mice. Liver [13]C-serine labeling was similar

between mice fed a HF-CTRL and HF-SAAR diet. Given that the liver serine pool size is larger

in mice fed a HF-SAAR diet, a greater amount of exogenous [13]C-serine would need to be taken up

by the liver to achieve a similar proportion of [13]C-serine labeling in livers of mice fed a HF-SAAR

diet. Thus, our data aligns with the hypothesis that SAAR increases asparagine synthesis from

glucose to promote liver serine uptake.

Role of hepatic PSAT1 in lipid adaptations to SAAR

Our findings agree with prior studies reporting that SAAR lowers adiposity and liver

triglycerides (2; 5; 9; 18; 21). In the current study, the decline in liver triglycerides following

SAAR was associated with decreased *de novo* lipogenesis enzymes (ACLY, FASN, and SCD-1).

There was a modest increase in mitochondrial respiratory chain proteins (complexes I, IV, and V)

in response to SAAR. These results are consistent with our prior work showing dietary SAAR

increases the expression of select mitochondrial proteins without enhancing mitochondrial

respiration (21). While terminal oxidation may not be enhanced by SAAR (21), other means of

lipid disposal could contribute to lower triglycerides. Studies using [14]C-palmitate in liver

homogenates have determined that SAAR increases liver β-oxidation (60). We identified short-

chain acylglycines (acetylglycine, isobutyrylglycine, and isovalerylglycine) were increased. These

acylglycines are generated in the mitochondria by conjugating acyl-CoAs to glycine in a reaction

mediated by GLYAT (61). Thus, the SAAR-mediated rise in glycine may facilitate acylglycine

synthesis and, consequently, promote the disposal of fatty acids during SAAR. Importantly, SAAR

elicited the molecular responses in lipid metabolism and lowered liver triglycerides similarly in

WT and PSAT1 KO mice. Overall, the prevailing signature under SAAR is a coordinated decrease

in *de novo* lipogenesis and stimulation of acyl-CoA disposal that is independent of PSAT1 action.

In summary, hepatic PSAT1 is necessary for adaptations to SAAR involving serine

synthesis that allow improvements in glucose tolerance to be fully realized. In contrast, PSAT1 is

dispensable for the adaptations in lipid accretion and disposal pathways that contribute to the

SAAR-mediated decline in liver lipids. Thus, the serine synthesis pathway is critical to the glucose,

but not lipid, adaptations to SAAR. These studies provide a mechanistic foundation for how SAAR

impacts the liver to prevent dysregulated glucose control.

ACKNOWLEDGEMENTS

The authors acknowledge the technical assistance provided by Ruth Pfeiffer. Figure schematics

were created using BioRender.com.

FUNDING

The National Institutes of Health Grants DK136772 (C.C.H.), DK091538 (P.A.C.), AG069781

(P.A.C.), DK133196 (J.J.G.) supported this research. A.F.O. was supported by the National

Institutes of Health Ruth L. Kirschstein National Research Service Award T32 DK007293.

DUALITY OF INTEREST

No potential conflicts of interest relevant to this article were reported.

AUTHOR CONTRIBUTIONS

A.F.O and C.C.H. conceived and designed the research; A.F.O., C.V., F.I.R., K.M.A., A.M.P., A.B.N., and C.C.H. performed experiments; A.F.O., C.C.H., J.J.G. analyzed data; A.F.O. and C.C.H. interpreted results of experiments; A.F.O. and C.C.H. prepared figures; A.F.O. drafted the manuscript; A.F.O., C.V., F.I.R., K.M.A., A.M.P., A.B.N., P.A.C., S.C.H., J.J.G., and C.C.H edited and revised manuscript; A.F.O., C.V., F.I.R., K.M.A., A.M.P., A.B.N., P.A.C., S.C.H., J.J.G., and C.C.H approved final version of the manuscript. C.C.H. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

REFERENCES

1. Fang H, Stone KP, Wanders D, Forney LA, Gettys TW: The Origins, Evolution, and Future of Dietary Methionine Restriction. Annu Rev Nutr 2022;42:201-226

2. Forney LA, Wanders D, Stone KP, Pierse A, Gettys TW: Concentration-dependent linkage of dietary methionine restriction to the components of its metabolic phenotype. Obesity (Silver Spring) 2017;25:730-738

3. Malloy VL, Krajcik RA, Bailey SJ, Hristopoulos G, Plummer JD, Orentreich N: Methionine restriction decreases visceral fat mass and preserves insulin action in aging male Fischer 344 rats independent of energy restriction. Aging Cell 2006;5:305-314

4. Thyne KM, Salmon AB: Metabolic benefits of methionine restriction in adult mice do not require functional methionine sulfoxide reductase A (MsrA). Sci Rep 2022;12:5073

5. Lees EK, Banks R, Cook C, Hill S, Morrice N, Grant L, Mody N, Delibegovic M: Direct comparison of methionine restriction with leucine restriction on the metabolic health of C57BL/6J mice. Sci Rep 2017;7:9977

6. Lees EK, Krol E, Grant L, Shearer K, Wyse C, Moncur E, Bykowska AS, Mody N, Gettys TW, Delibegovic M: Methionine restriction restores a younger metabolic phenotype in adult mice with alterations in fibroblast growth factor 21. Aging Cell 2014;13:817-827

7. Ables GP, Perrone CE, Orentreich D, Orentreich N: Methionine-restricted C57BL/6J mice are resistant to diet-induced obesity and insulin resistance but have low bone density. PLoS One 2012;7:e51357

8. Wanders D, Burk DH, Cortez CC, Van NT, Stone KP, Baker M, Mendoza T, Mynatt RL, Gettys TW: UCP1 is an essential mediator of the effects of methionine restriction on energy balance but not insulin sensitivity. FASEB J 2015;29:2603-2615

9. Wanders D, Forney LA, Stone KP, Burk DH, Pierse A, Gettys TW: FGF21 Mediates the Thermogenic and Insulin-Sensitizing Effects of Dietary Methionine Restriction but Not Its Effects on Hepatic Lipid Metabolism. Diabetes 2017;66:858-867

10. Wanders D, Stone KP, Forney LA, Cortez CC, Dille KN, Simon J, Xu M, Hotard EC, Nikonorova IA, Pettit AP, Anthony TG, Gettys TW: Role of GCN2-Independent Signaling Through a Noncanonical PERK/NRF2 Pathway in the Physiological Responses to Dietary Methionine Restriction. Diabetes 2016;65:1499-1510

11. Stone KP, Wanders D, Orgeron M, Cortez CC, Gettys TW: Mechanisms of increased in vivo insulin sensitivity by dietary methionine restriction in mice. Diabetes 2014;63:3721-3733

12. Plaisance EP, Greenway FL, Boudreau A, Hill KL, Johnson WD, Krajcik RA, Perrone CE, Orentreich N, Cefalu WT, Gettys TW: Dietary methionine restriction increases fat oxidation in obese adults with metabolic syndrome. J Clin Endocrinol Metab 2011;96:E836-840

13. Olsen T, Stolt E, Ovrebo B, Elshorbagy A, Tore EC, Lee-Odegard S, Troensegaard H, Johannessen H, Doeland B, Vo AAD, Dahl AF, Svendsen K, Thoresen M, Refsum H, Rising R, Barvikova K, van Greevenbroek M, Kozich V, Retterstol K, Vinknes KJ: Dietary sulfur amino acid restriction in humans with overweight and obesity: a translational randomized controlled trial. J Transl Med 2024;22:40

14. Richie JP, Jr., Sinha R, Dong Z, Nichenametla SN, Ables GP, Ciccarella A, Sinha I, Calcagnotto AM, Chinchilli VM, Reinhart L, Orentreich D: Dietary Methionine and Total Sulfur Amino Acid Restriction in Healthy Adults. J Nutr Health Aging 2023;27:111-123

15. Fang H, Stone KP, Ghosh S, Forney LA, Gettys TW: The Role of Reduced Methionine in Mediating the Metabolic Responses to Protein Restriction Using Different Sources of Protein. Nutrients 2021;13

16. Fang H, Stone KP, Ghosh S, Forney LA, Sims LC, Vincik L, Gettys TW: Hepatic Nfe2l2 Is
Not an Essential Mediator of the Metabolic Phenotype Produced by Dietary Methionine
Restriction. Nutrients 2021;13

17. Fang H, Stone KP, Forney LA, Sims LC, Gutierrez GC, Ghosh S, Gettys TW:
Implementation of dietary methionine restriction using casein after selective, oxidative deletion
of methionine. iScience 2021;24:102470

18. Stone KP, Ghosh S, Kovalik JP, Orgeron M, Wanders D, Sims LC, Gettys TW: The acute
transcriptional responses to dietary methionine restriction are triggered by inhibition of ternary
complex formation and linked to Erk1/2, mTOR, and ATF4. Sci Rep 2021;11:3765

19. Forney LA, Fang H, Sims LC, Stone KP, Vincik LY, Vick AM, Gibson AN, Burk DH, Gettys
TW: Dietary Methionine Restriction Signals to the Brain Through Fibroblast Growth Factor 21
to Regulate Energy Balance and Remodeling of Adipose Tissue. Obesity (Silver Spring)
2020;28:1912-1921

20. Ghosh S, Forney LA, Wanders D, Stone KP, Gettys TW: An integrative analysis of tissue-
specific transcriptomic and metabolomic responses to short-term dietary methionine restriction in
mice. PLoS One 2017;12:e0177513

21. Rome FI, Hughey CC: Disrupted liver oxidative metabolism in glycine N-methyltransferase-
deficient mice is mitigated by dietary methionine restriction. Mol Metab 2022;58:101452

22. Ommi NB, Mattocks DAL, Kalecky K, Bottiglieri T, Nichenametla SN: Pharmacological
recapitulation of the lean phenotype induced by the lifespan-extending sulfur amino acid-
restricted diet. Aging (Albany NY) 2025;17:960-981

23. Nichenametla SN, Mattocks DAL, Cooke D, Midya V, Malloy VL, Mansilla W, Ovrebo B,
Turner C, Bastani NE, Sokolova J, Pavlikova M, Richie JP, Jr., Shoveller AK, Refsum H, Olsen
T, Vinknes KJ, Kozich V, Ables GP: Cysteine restriction-specific effects of sulfur amino acid
restriction on lipid metabolism. Aging Cell 2022;21:e13739

24. Kalhan SC, Hanson RW: Resurgence of serine: an often neglected but indispensable amino
Acid. J Biol Chem 2012;287:19786-19791

25. Hamano M, Esaki K, Moriyasu K, Yasuda T, Mohri S, Tashiro K, Hirabayashi Y, Furuya S:
Hepatocyte-Specific Phgdh-Deficient Mice Culminate in Mild Obesity, Insulin Resistance, and
Enhanced Vulnerability to Protein Starvation. Nutrients 2021;13

26. Yu J, Xiao F, Guo Y, Deng J, Liu B, Zhang Q, Li K, Wang C, Chen S, Guo F: Hepatic
Phosphoserine Aminotransferase 1 Regulates Insulin Sensitivity in Mice via Tribbles Homolog 3.
Diabetes 2015;64:1591-1602

27. Sim WC, Lee W, Sim H, Lee KY, Jung SH, Choi YJ, Kim HY, Kang KW, Lee JY, Choi YJ,
Kim SK, Jun DW, Kim W, Lee BH: Downregulation of PHGDH expression and hepatic serine
level contribute to the development of fatty liver disease. Metabolism 2020;102:154000

28. Raines LN, Zhao H, Wang Y, Chen H-Y, Gallart-Ayala H, Hsueh P-C, Cao W, Koh Y,
Alamonte-Loya A, Liu P-S: PERK is a critical metabolic hub for immunosuppressive function in
macrophages. Nature immunology 2022;23:431-445

29. Rome FI, Shobert GL, Voigt WC, Stagg DB, Puchalska P, Burgess SC, Crawford PA, Hughey
CC: Loss of hepatic phosphoenolpyruvate carboxykinase 1 dysregulates metabolic responses to
acute exercise but enhances adaptations to exercise training in mice. Am J Physiol Endocrinol
Metab 2023;324:E9-E23

30. Chan TM, Exton JH: A rapid method for the determination of glycogen content and
radioactivity in small quantities of tissue or isolated hepatocytes. Analytical biochemistry
1976;71:96-105

31. Puchalska P, Crawford PA: Application of stable isotope labels for metabolomics in studies in
fatty liver disease. In *Metabolomics: Methods and Protocols*, Springer, 2019, p. 259-272
32. Nelson AB, Reese LE, Rono E, Queathem ED, Qiu Y, McCluskey BM, Crampton A, Conniff
E, Cummins K, Boytim E: Deciphering Colorectal Cancer–Hepatocyte Interactions: A
Multiomics Platform for Interrogation of Metabolic Crosstalk in the Liver–Tumor
Microenvironment. International journal of molecular sciences 2025;26:1976
33. Queathem ED, Moazzami Z, Stagg DB, Nelson AB, Fulghum K, Hayir A, Seay A,
Gillingham JR, d'Avignon DA, Han X: Ketogenesis supports hepatic polyunsaturated fatty acid
homeostasis via fatty acid elongation. Science Advances 2025;11:eads0535
34. Koelmel JP, Kroeger NM, Gill EL, Ulmer CZ, Bowden JA, Patterson RE, Yost RA, Garrett
TJ: Expanding lipidome coverage using LC-MS/MS data-dependent acquisition with automated
exclusion list generation. Journal of the American Society for Mass Spectrometry 2017;28:908-
917
35. Gaffney DO, Jennings EQ, Anderson CC, Marentette JO, Shi T, Schou Oxvig AM, Streeter
MD, Johannsen M, Spiegel DA, Chapman E, Roede JR, Galligan JJ: Non-enzymatic Lysine
Lactoylation of Glycolytic Enzymes. Cell Chem Biol 2020;27:206-213 e206
36. Galligan JJ, Wepy JA, Streeter MD, Kingsley PJ, Mitchener MM, Wauchope OR, Beavers
WN, Rose KL, Wang T, Spiegel DA, Marnett LJ: Methylglyoxal-derived posttranslational
arginine modifications are abundant histone marks. Proc Natl Acad Sci U S A 2018;115:9228-
9233
37. Galligan JJ, Kingsley PJ, Wauchope OR, Mitchener MM, Camarillo JM, Wepy JA, Harris
PS, Fritz KS, Marnett LJ: Quantitative Analysis and Discovery of Lysine and Arginine
Modifications. Anal Chem 2017;89:1299-1306
38. Mehran AE, Templeman NM, Brigidi GS, Lim GE, Chu KY, Hu X, Botezelli JD, Asadi A,
Hoffman BG, Kieffer TJ, Bamji SX, Clee SM, Johnson JD: Hyperinsulinemia drives diet-
induced obesity independently of brain insulin production. Cell Metab 2012;16:723-737
39. Thomas DD, Corkey BE, Istfan NW, Apovian CM: Hyperinsulinemia: An Early Indicator of
Metabolic Dysfunction. J Endocr Soc 2019;3:1727-1747
40. Wiebe N, Ye F, Crumley ET, Bello A, Stenvinkel P, Tonelli M: Temporal Associations Among
Body Mass Index, Fasting Insulin, and Systemic Inflammation: A Systematic Review and Meta-
analysis. JAMA Network Open 2021;4:e211263-e211263
41. Werge MP, McCann A, Galsgaard ED, Holst D, Bugge A, Albrechtsen NJW, Gluud LL: The
Role of the Transsulfuration Pathway in Non-Alcoholic Fatty Liver Disease. J Clin Med 2021;10
42. Trujillo MN, Jennings EQ, Farrera DO, Kitamura N, Anderson CC, Gehrke S, Reisz JA,
Johannsen M, Roede JR, D'Alessandro A, Galligan JJ: Glyoxalase 2 Coordinates de Novo Serine
Metabolism. Chembiochem 2025;26:e202401086
43. Buescher JM, Antoniewicz MR, Boros LG, Burgess SC, Brunengraber H, Clish CB,
DeBerardinis RJ, Feron O, Frezza C, Ghesquiere B: A roadmap for interpreting 13C metabolite
labeling patterns from cells. Current opinion in biotechnology 2015;34:189-201
44. Christen S, Lorendeau D, Schmieder R, Broekaert D, Metzger K, Veys K, Elia I, Buescher
JM, Orth MF, Davidson SM: Breast cancer-derived lung metastases show increased pyruvate
carboxylase-dependent anaplerosis. Cell reports 2016;17:837-848
45. Bruntz RC, Lane AN, Higashi RM, Fan TW-M: Exploring cancer metabolism using stable
isotope-resolved metabolomics (SIRM). Journal of Biological Chemistry 2017;292:11601-11609

46. Johnston K, Pachnis P, Tasdogan A, Faubert B, Zacharias LG, Vu HS, Rodgers-Augustyniak L, Johnson A, Huang F, Ricciardo S: Isotope tracing reveals glycolysis and oxidative metabolism in childhood tumors of multiple histologies. Med 2021;2:395-410. e394

47. Lau AN, Li Z, Danai LV, Westermark AM, Darnell AM, Ferreira R, Gocheva V, Sivanand S, Lien EC, Sapp KM: Dissecting cell-type-specific metabolism in pancreatic ductal adenocarcinoma. Elife 2020;9:e56782

48. Pachnis P, Wu Z, Faubert B, Tasdogan A, Gu W, Shelton S, Solmonson A, Rao AD, Kaushik AK, Rogers TJ: In vivo isotope tracing reveals a requirement for the electron transport chain in glucose and glutamine metabolism by tumors. Science Advances 2022;8:eabn9550

49. Holm LJ, Haupt-Jorgensen M, Larsen J, Giacobini JD, Bilgin M, Buschard K: L-serine supplementation lowers diabetes incidence and improves blood glucose homeostasis in NOD mice. PLoS One 2018;13:e0194414

50. Lopez-Gonzales E, Lehmann L, Ruiz-Ojeda FJ, Hernandez-Bautista R, Altun I, Onogi Y, Khalil AE, Liu X, Israel A, Ussar S: L-Serine Supplementation Blunts Fasting-Induced Weight Regain by Increasing Brown Fat Thermogenesis. Nutrients 2022;14

51. Handzlik MK, Metallo CM: Sources and Sinks of Serine in Nutrition, Health, and Disease. Annu Rev Nutr 2023;43:123-151

52. Achouri Y, Robbi M, Van Schaftingen E: Role of cysteine in the dietary control of the expression of 3-phosphoglycerate dehydrogenase in rat liver. Biochem J 1999;344 Pt 1:15-21

53. Garcia RA, Stipanuk MH: The splanchnic organs, liver and kidney have unique roles in the metabolism of sulfur amino acids and their metabolites in rats. J Nutr 1992;122:1693-1701

54. Stipanuk MH, Coloso RM, Garcia RA, Banks MF: Cysteine concentration regulates cysteine metabolism to glutathione, sulfate and taurine in rat hepatocytes. J Nutr 1992;122:420-427

55. Coloso RM, Hirschberger LL, Stipanuk MH: Uptake and metabolism of L-2-oxo-[35S]thiazolidine-4-carboxylate by rat cells is slower than that of L-[35S]cysteine or L-[35S]methionine. J Nutr 1991;121:1341-1348

56. Bagley PJ, Stipanuk MH: Rats fed a low protein diet supplemented with sulfur amino acids have increased cysteine dioxygenase activity and increased taurine production in hepatocytes. J Nutr 1995;125:933-940

57. Farrera DO, Galligan JJ: The Human Glyoxalase Gene Family in Health and Disease. Chem Res Toxicol 2022;

58. Trujillo MN, Jennings EQ, Hoffman EA, Zhang H, Phoebe AM, Mastin GE, Kitamura N, Reisz JA, Megill E, Kantner D, Marcinkiewicz MM, Twardy SM, Lebario F, Chapman E, McCullough RL, D'Alessandro A, Snyder NW, Cusanovich DA, Galligan JJ: Lactoylglutathione promotes inflammatory signaling in macrophages through histone lactoylation. Mol Metab 2024:101888

59. Krall AS, Xu S, Graeber TG, Braas D, Christofk HR: Asparagine promotes cancer cell proliferation through use as an amino acid exchange factor. Nat Commun 2016;7:11457

60. Hasek BE, Boudreau A, Shin J, Feng D, Hulver M, Van NT, Laque A, Stewart LK, Stone KP, Wanders D, Ghosh S, Pessin JE, Gettys TW: Remodeling the integration of lipid metabolism between liver and adipose tissue by dietary methionine restriction in rats. Diabetes 2013;62:3362-3372

61. Badenhorst CPS, van der Sluis R, Erasmus E, Van Dijk AA: Glycine conjugation: importance in metabolism, the role of glycine N-acyltransferase, and factors that influence interindividual variation. Expert opinion on drug metabolism & toxicology 2013;9:1139-1153

 FIGURE LEGENDS

Figure 1. Sulfur amino acid restriction (SAAR) prevents obesity and improves systemic
glucose control in C57BL/6J mice. A. A schematic representation of the study design. C57BL/6J
male mice were fed a low-fat control (LF-CTRL), high-fat control (HF-CTRL), or HF-SAAR diet
for 6 weeks starting at 6 weeks of age. **B.** Body weight time course (g), body weight change over
the 6 weeks on diet (g), fat mass (% of body weight), and lean mass (% of body weight). **C.** Liver
triglycerides (mg·g^{-1}) and liver weight (g). **D.** Liver glycogen (mg·g^{-1}). **E.** Blood glucose (mg·dL^{-1}
^{1}). **F.** Glucose excursion and area under the curve (AUC) during a glucose tolerance
test. **G.** Glucose excursion and AUC during an insulin tolerance test. **H.** Plasma insulin (pmol·L^{-1}
^{1}). n=3-17. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.
769

Figure 2. Sulfur amino acid restriction (SAAR) increases liver glucose partitioning to *de novo*
***serine synthesis in C57BL/6J mice. A.** A schematic representation of the select reactions involved
in *de novo* serine synthesis and SAA synthesis and catabolism. **B.** Liver methionine, cysteine, and
taurine (arbitrary units) in C57BL/6J male mice fed a low-fat control (LF-CTRL), high-fat control
(HF-CTRL), or HF-SAAR diet for 6 weeks starting at 6 weeks of age. **C.** Liver phosphoserine
aminotransferase 1 (PSAT1) as determined by immunoblotting and representative immunoblots in
mice fed a LF-CTRL and HF-CTRL diet only. **D.** Liver phosphoglycerate dehydrogenase
(PHGDH) and PSAT1 as determined by immunoblotting and representative immunoblots in mice
fed a LF-CTRL, HF-CTRL, and HF-SAAR diet. **E.** Liver serine and glycine (arbitrary units). **F.** A
schematic representation of the study design. C57BL/6J mice fed LF-CTRL (0.88% kcal from
methionine, 510072, Dyets Inc., Bethlehem, PA) and low-fat SAAR (LF-SAAR; 0.18% kcal from
methionine, 510071, Dyets Inc., Bethlehem, PA) for 6 weeks starting at 6 weeks of age. A 30 -
minute intravenous infusion of [U-^{13}C]glucose in the conscious, unrestrained mouse was
completed at 12 weeks of age. **G.** Arterial blood glucose (mg·dL^{-1}). **H.** Fractional enrichment (%
total pool) of liver glucose, serine and glycine from [U-^{13}C]glucose. n=6-10. [#]p<0.05 by *t*-test,
*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by ANOVA.
786

Figure 3. Liver PSAT1 is required for systemic glucose adaptations to sulfur amino acid
restriction (SAAR). A. A schematic representation of the study design. Liver-specific PSAT1
knockout (KO) mice and wild type (WT) littermates were fed a high-fat control (HF-CTRL) or
HF-SAAR diet for 6 weeks. **B.** Body weight time course (g) and body weight change over 6 weeks
on diet (g). **C.** Fat mass and lean mass (% of body weight). **D.** Glucose excursion curves and area
under the curve (AUC) for glucose tolerance tests. **E.** Glucose excursion curves and AUC for
insulin tolerance tests. **F.** Plasma insulin (pmol·L^{-1}) before (0 min) and 15 minutes after an insulin
injection. **G.** Liver phosphoAkt (pAkt)-to-Akt ratio as determined by immunoblotting and
representative immunoblots. **H.** Liver glycogen (mg·g^{-1}). **I.** Liver phospho-glycogen synthase
(pGS)-to-GS ratio and phospho-glycogen phosphorylase (pPYGL)-to-PYGL ratio as determined
by immunoblotting and representative immunoblots. n=5-18. *p<0.05, **p<0.01, ***p<0.001,
****p<0.0001.
799

800 **Figure 4. Sulfur amino acid restriction (SAAR) increases liver *de novo* serine synthesis from**
801 **glucose. A.** A schematic representation of select metabolites and enzymes involved in *de novo*
802 serine synthesis from glucose. **B.** Liver phosphoglycerate dehydrogenase (PHGDH),
803 phosphoserine aminotransferase 1 (PSAT1), and serine hydroxymethyltransferase 2 (SHMT2) as

determined by immunoblotting and representative immunoblots in wild type (WT) and liver-specific PSAT1 knockout (KO) mice fed a high-fat control (HF-CTRL) or HF-SAAR diet for 6 weeks. **C.** Liver glucose, serine, and glycine (arbitrary units) in 7-hour fasted mice. **D.** Fractional enrichment (% of total pool) of liver glucose, serine, and glycine following a 30-minute intravenous infusion of [U-^{13}C]glucose. n=7-9. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 5. Sulfur amino acid restriction (SAAR) increases partitioning of glucose-sourced serine to liver glutathione. A. A schematic representation of the select reactions involved in *de novo* serine synthesis and SAA synthesis and catabolism. **B.** Liver cystathionine β-synthase (CBS), cystathionine γ-lyase (CSE), cysteine dioxygenase 1 (CDO1), glutamate-cysteine ligase catalytic subunit (GCLC), and glutathione synthetase (GSS) as determined by immunoblotting and representative immunoblots in wild type (WT) and liver-specific PSAT1 knockout (KO) mice fed a high-fat control (HF-CTRL) or HF-SAAR diet for 6 weeks. **C.** Liver methionine, cysteine, taurine, reduced glutathione (GSH), and oxidized glutathione (GSSG; arbitrary units) in 7-hour fasted mice. **D.** Fractional enrichment (% of total pool) of liver methionine, taurine, GSH, and GSSG following a 30-minute intravenous infusion of [U-^{13}C]glucose. **E.** Fractional enrichment (% of total pool) of liver serine, methionine, taurine, GSH, and GSSG at 15 minutes following an i.p. injection of [U-^{13}C]serine. n=6-9. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 6. Sulfur amino acid restriction (SAAR) lowers glyoxalase cycle metabolites and associated post-translational modifications (PTMs). A. A schematic of select reactions, metabolites, and PTMs related to the glyoxalase cycle. **B.** Liver glyoxalase 1 (GLO1) and hydroxyacylglutathione hydrolase (GLO2) as determined by immunoblotting and representative immunoblots in wild type (WT) and liver-specific PSAT1 knockout (KO) mice fed a high-fat control (HF-CTRL) or HF-SAAR diet for 6 weeks. **C.** Liver glutathione (GSH; arbitrary units), hemithioacetal (HTA; arbitrary units), lactoylglutathione (arbitrary units), and lactate (arbitrary units). **D.** Global PTMs (pmol·nmol Leu^{-1}): lactoyllysine, methylglyoxal-derived hydroimidazolone 1 (MG-H1), carboxyethylarginine (CEA), and acetyllysine. n=6-9. *p<0.05, **p<0.01, ****p<0.0001.

Figure 7. Sulfur amino acid restriction (SAAR) increases glucose partitioning toward asparagine synthesis. A. A schematic representation of select reactions and metabolites linking tricarboxylic acid (TCA)-cycle intermediates to asparagine synthesis. **B.** Liver TCA cycle, anaplerotic and cataplerotic metabolites (arbitrary units) in wild type (WT) and liver-specific PSAT1 knockout (KO) mice fed a high-fat control (HF-CTRL) or HF-SAAR diet for 6 weeks. **C.** Fractional enrichment (% of total pool) of liver TCA cycle, anaplerotic and cataplerotic metabolites following a 30-minute intravenous infusion of [U-^{13}C]glucose. **D.** Ratios of isotopologues used as indices of anaplerotic and cataplerotic flux. **E.** Liver asparagine synthase (ASNS), pyruvate carboxylase (PC), propionyl-CoA carboxylase subunit A (PCCA), malate dehydrogenase 2 (MDH2), and mitochondrial glutamic-oxaloacetic transaminase (GOT2) as determined by immunoblotting and representative immunoblots. n=6-9. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 8. Sulfur amino acid restriction (SAAR) reduces lipid accumulation independently of hepatic PSAT1. Panels A-G are measurements in in wild type (WT) and liver-specific PSAT1 knockout (KO) mice fed a high-fat control (HF-CTRL) or HF-SAAR diet for 6 weeks. Panels H-

J are measurements in WT and KO mice fed the HF-CTRL and -SAAR diets for 20 weeks. **A.** Liver triglycerides (mg·g^{-1}). **B.** Liver *de novo* lipogenesis enzymes: ATP-citrate lyase (ACLY), acetyl-CoA carboxylase (ACC), fatty acid synthase (FASN), and stearoyl-CoA desaturase 1 (SCD1) as determined by immunoblotting (arbitrary units). **C.** Liver mitochondrial respiratory complexes I-V (CI-V) as determined by immunoblotting. **D.** Representative immunoblots for panels B, C, and F. **E.** A schematic representation of select metabolites and enzymes related to the synthesis of acylglycines. **F.** Liver glycine N-acyltransferase (GLYAT) as determined by immunoblotting. **G.** Liver acylglycines (arbitrary units): acetylglycine, isobutyrylglycine, isovalerylglycine, hexanoylglycine, and capryloylglycine. **H.** Body weight time course (g) and change in body weight (g) over 20 weeks on diet. **I.** Fat mass and lean mass (% body weight). **J.** Liver triglycerides (mg·g^{-1}). n=5-9 *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Supplementary Figure 1. Sulfur Amino Acid Restriction (SAAR) impacts systemic glucose control and hepatic glucose partitioning in C57BL/6J mice independent of insulin levels. A. A schematic representation of the study design. C57BL/6J male mice were fed a low-fat control (LF-CTRL), high-fat control (HF-CTRL), or HF-SAAR diet for 6 weeks starting at 6 weeks of age. A subset of mice fed the HF-SAAR diet had a subcutaneous insulin pellet (HF-SAAR + Insulin) implanted at 7 weeks of age for sustained release of insulin until 12 weeks of age. **B.** Body weight time course (g), body weight change over the 6 weeks on diet (g), fat mass (% of body weight), and lean mass (% of body weight). **C.** Liver triglycerides (mg·g^{-1}) and liver weight (g). **D.** Liver glycogen (mg·g^{-1}). **E.** Blood glucose (mg·dL^{-1}). **F.** Glucose excursion and area under the curve (AUC) during a glucose tolerance test. **G.** Glucose excursion and AUC during an insulin tolerance test. **H.** Plasma insulin (pmol·L^{-1}). **I.** Liver methionine, homocysteine, cysteine, and taurine (arbitrary units). **J.** Liver phosphoglycerate dehydrogenase (PHGDH) and phosphoserine aminotransferase 1 (PSAT1) as determined by immunoblotting and representative immunoblots. **K.** Liver serine and glycine (arbitrary units). n=3-17. [#]p<0.05 by *t*-test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by ANOVA.

Figure 1. SAAR prevents obesity and improves systemic glucose control in C57BL/6J mice

Figure 2. SAAR increased glucose partitioning to de novo serine synthesis in C57BL/6J mice

Figure 3. Liver PSAT1 is required for systemic glucose adaptations to SAAR

Figure 4. SAAR increases liver de novo serine synthesis from glucose

Figure 5. SAAR increases partitioning of glucose-sourced serine to glutathione

Figure 6. SAAR lowers liver glyoxalase cycle metabolites and associated post-translational modifications

Figure 7. SAAR increased glucose partitioning toward liver asparagine synthesis

Figure 8. SAAR reduces lipid accumulation independently of hepatic PSAT1

Graphical abstract