Sexual dimorphism in sensorimotor transformation of optic flow

Sarah Nicholas[1], Katja Sporar Klinge[1#], Luke Turnbull[1], Annabel Moran[1], Aika Young[1], Yuri Ogawa[1], Karin Nordström[1,2*]*

[1]*Flinders Health and Medical Research Institute, Flinders University, GPO Box 2100, Adelaide, SA, 5001, Australia;* [2]*Department of Medical Cell Biology, Uppsala University, 75123 Uppsala, Sweden.*

[#]Current address: Institute for Developmental Biology, RWTH Aachen University, 52074 Aachen, Germany

*Shared authorship; email: yuri.ogawa@flinders.edu.au; karin.nordstrom@flinders.edu.au

ORCIDs

Sarah Nicholas: 0000-0002-5555-9421

Katja Sporar Klinge: 0009-0004-7224-6568

Aika Young: 0009-0000-5028-6164

Yuri Ogawa: 0000-0002-9708-7063

Karin Nordström: 0000-0002-6020-6348

CRediT author statement

Sarah Nicholas: conceptualization, methodology, validation, formal analysis, investigation, data curation, writing – original draft, visualization, project administration; **Katja Sporar**

Klinge: methodology, validation, formal analysis, investigation, data curation, writing –

review and editing, visualization; **Luke Turnbull:** methodology, validation, investigation,

writing – review and editing; **Annabel Moran:** investigation, writing – review and editing;

Aika Young: investigation, writing – review and editing; **Yuri Ogawa:** methodology,

validation, formal analysis, investigation, data curation, writing – original draft, visualization,

supervision, project administration; **Karin Nordström:** conceptualization, validation, formal

analysis, resources, writing – original draft, visualization, supervision, project administration,

funding acquisition.

Abstract

Motion vision underpins a wide range of adaptive behaviours essential for individual and species survival. In hoverflies, some visual behaviours are sexually dimorphic, including for example male high-speed pursuit of conspecifics, matched by improved optics, and faster photoreceptors. Other visual behaviours are sexually monomorphic, with for example similar foraging flight speeds in male and female hoverflies. However, whether the descending neurons responsible for sensorimotor transformation of optic flow are sexually dimorphic is unknown. To address this, we combined morphological analysis with electrophysiology of optic flow sensitive descending neurons and compared neural responses to the behavioural output in tethered hoverflies. We found that while optomotor flight behaviour is largely sexually monomorphic, the underlying neural responses are sexually dimorphic, especially at higher optic flow velocities. Additionally, behavioural responses were noticeably slower than neural responses. Together, our findings uncover a nuanced, sex- and stimulus- dependant sensorimotor transformation, shaped by both neural architecture and behavioural demands.

Keywords

Sexual dimorphism, hoverfly, descending neuron, optic flow, optomotor response

Introduction

Motion vision is a fundamental sensory modality across the animal kingdom, enabling animals to navigate, to maintain a straight trajectory, to avoid collisions, and to identify prey, predators or mates. Among the most potent cues for self-motion is widefield optic flow, the coherent motion of the entire visual field, generated by an animal's own movement through the world. In insects, the neural mechanisms underlying optic flow processing have been studied extensively (for a review, see e.g. Ref [1]), offering a powerful model for understanding how compact nervous systems extract behaviourally relevant information from dynamic visual scenes.

To generate appropriate responses to widefield optic flow, the visual input needs to be integrated across space. In flies, this spatial pooling occurs in 45-60 lobula plate tangential cells (LPTCs)[2,3], each matched to a particular type of self-generated optic flow[4], where the horizontal system (HS) and vertical system (VS) cells are the most well described. HS cells respond optimally to rotations around the yaw axis, whereas VS cells are tuned to pitch and roll rotations[5]. LPTCs project to the inferior posterior slope[6], where they synapse with descending neurons[7,8]. In *Drosophila* at least 35 descending neuron types have their inputs in the posterior surface of the brain (named DNp1-35)[9]. Furthermore, in *Drosophila* and blowflies, three of these descending neurons have been shown to respond robustly to widefield optic flow. Their axons project to the dorsal part of the thoracic ganglion, where motor neurons controlling the neck, wings and halteres are located[7,10-13].

64 DNp15, also called DNHS1, receives input from HS cells[9-11] and is physiologically similar to

65 the optic flow sensitive descending neuron type 1 (OFS DN1) in the hoverfly *Eristalis*

66 *tenax*[14], although direct LPTC coupling has not yet been demonstrated in the hoverfly.

67 DNHS1 projects to the neck and haltere motor neuropils[9] and has been implicated in the

68 control of head yaw movements, abdominal ruddering and flight stabilization via the haltere

69 motor system[10]. DNp20, or DNOVS1, receiving input from the ocelli and VS cells[7,9,10,15], is

70 likely involved in rapid head movements[9], possibly facilitating gaze stabilization during

71 flight[10]. DNp22, or DNOVS2, also receives input from the ocelli but a different subset of VS

72 cells[7,9,10,16]. DNOVS2 is physiologically similar to the *Eristalis tenax* optic flow sensitive

73 descending neuron type 2 (OFS DN2)[14], but synaptic coupling with VS cells has not been

74 shown in hoverflies. DNOVS2 projects to the neck, wing and haltere motor neuropils[9] and

75 has been implied in the initiation of the fast body saccades that support rapid re-orientation[10]

76 during flight in response to dynamic visual cues.

77 Hoverflies are interesting in the context of motion vision, which they use both to maintain a

78 hovering stance, and to fly at high speed[17]. Indeed, hoverflies show striking sexually

79 dimorphic flight behaviour, where males establish territories which they guard rigorously

80 from a hovering stance[18], followed by high-speed flight to chase away any intruding insects

81 and/or pursue conspecific females for courtship and mating[17,19]. Male hoverflies are also

82 smaller than females[20,21], which introduces an aerodynamic component by reducing inertia

83 and enabling finer control over rapid flight adjustments[22]. Accompanying this sexually

84 dimorphic pursuit behaviour, male *E. tenax* have larger lenses than females in a dorso-frontal

85 bright zone[23], with faster motion detection and increased signal-to-noise ratio[24]. In many fly

species, the photoreceptors in this part of the eye are also faster in males[25]. In hoverflies, even

if the LPTCs are typically implied in optomotor responses[1], males have a smaller HSN

receptive field[26] and velocity tuning shifted to higher velocities[23,27]. These adaptations are

likely useful in regulating optomotor responses during high-speed target pursuit[28,29]. Indeed,

as target selective descending neurons (TSDNs) are suppressed when target and background

move in the same direction[28,30], the optic flow sensitive descending neurons may serve a

complementary role in stabilizing flight under these conditions.

The differences in optics, photoreceptor dynamics, and LPTC receptive field size and

velocity tuning, have been interpreted as required by males in the fast flight used during

sexually dimorphic territorial behaviours. Interestingly, there is no sexual dimorphism in

flight speed during behaviours likely to be governed primarily by the LPTCs, such as

foraging between flowers[31] and when flying within the confines of an indoor arena[32]. To

investigate the discrepancy between sexually dimorphic visual processing and sexually

monomorphic flight speed, we compared the electrophysiological response characteristics

and morphology of optic flow sensitive descending neurons in male and female hoverflies

and linked these findings to the behaviour of tethered hoverflies viewing similar stimuli. We

used the wing beat amplitude (WBA) as a measure of the optomotor response, and found no

sexual dimorphism at speeds up to 2 m/s for translation and 200°/s for rotation. While the

head movements were largely sexually monomorphic, the extension of the fore- and hind legs

exhibited clearer sexual dimorphism. Furthermore, while neural morphology, receptive fields

and direction sensitivity of the descending neurons showed minimal sex differences, there

was a significant and noticeable difference in the velocity response functions between males

and females, especially at higher speeds. Critically, neural differences were not only velocity

dependent but also varied between stimuli (occurring only for sideslip, lift and thrust, but not

roll) and neuron type. These neuron-, stimulus-, and sex-specific differences uncovers a

previously unrecognized complexity in the neural encoding of visual motion, revealing for

the first time, a sex-dependent transformation from sensory input to motor output.

Results

Two distinct types of optic flow sensitive descending neurons can be identified by their receptive field location and preferred direction

Optic flow sensitive descending neurons can be readily identified by mapping their receptive field using small sinusoidal gratings[14,23]. Based on the receptive fields of 100 reference neurons recorded from 90 male hoverflies, we found that two key parameters, the azimuthal position of the receptive field centre and its preferred direction of motion, are sufficient to reliably ascertain neuron type (Fig. 1, Supplementary Fig. 1). OFS DN1 has a preferred direction up and away from the midline, either leftward (range from 137° to 171°) or rightward (range from 16° to 40°) for neurons on the left- and right-hand side of the visual field, respectively (Fig. 1a, b and c; green, Fig. 1g, h and i). OFS DN2 neurons respond preferentially to downward motion (range from 228° to 293°; Fig. 1d, e and f; yellow and orange, Fig. 1g, h and i) with the azimuthal position of the receptive field centre separating left-hand side (LHS) from right-hand side neurons (RHS, Fig. 1g, h and i).

We used the maximum local motion sensitivity (LMS, Fig. 1c, f), the extent of the receptive

fields (number of positions with LMS over 50%, arrows in Fig. 1c, f) and the local preferred

direction (LPD, Fig. 1c, f) variance from these 100 reference neurons (grey data,

Supplementary Fig. 2a, b and c) to set strict exclusion criteria (dashed red, Supplementary

Fig. 2a, b and c) of the neurons used in the rest of the paper. This resulted in the exclusion of

two neurons from males and seven from females (grey, Supplementary Fig. 2d, e) due to

either low LMS (less than 20 spikes/s, Supplementary Fig. 2a), a small number of locations

where LMS exceeded 50% of the maximum (four positions or less, Supplementary Fig. 2b)

or high LPD variance (above 30°, Supplementary Fig. 2c).

We found no sexual dimorphism in either neuron type when comparing receptive field width

and height of the remaining 33 male and 29 female neurons (Supplementary Fig. 2f, unpaired

t-test, $p = 0.52$ and 0.09 for width and height of OFS DN1, and $p = 0.19$ and 0.13 for width

and height of OFS DN2).

Directional tuning of optic flow sensitive descending neurons exhibits limited sexual

dimorphism

Some LPTCs, which are upstream of optic flow sensitive descending neurons, show distinct

sexual dimorphism, whilst others do not[23,26]. The receptive field data used for classifying

OFS DN1 and DN2 showed that they are strongly directional (Fig. 1g, i). To investigate if

this direction tuning is sexually dimorphic, we used a separate experiment quantifying

responses to full-screen sinusoidal grating stimuli (wavelength 7°, 5 Hz, Supplementary Fig.

3). We found that the resulting preferred direction of both OFS DN1 and OFS DN2 matched

their receptive field preferred directions (compare polar plots in Supplementary Fig. 2d, e

with Supplementary Fig. 3g, h), i.e. up and away from the visual midline for OFS DN1

(range from 359° to 52°) or downwards for OFS DN2 (range from 273° to 297°, Fig. 2a).

Whilst OFS DN1 showed no difference in preferred direction between the sexes (Watson-

Williams two-sample test, $p = 0.38$), male OFS DN2 had a slightly more lateral preferred

direction compared to females (Fig. 2a; median = 285.4° compared to 281.5°, Watson-

Williams two-sample test, $p = 0.046$).

We quantified the spontaneous rate and found that neither this nor the responses of OFS DN1

to a stationary starfield pattern differed between the sexes (circles, Supplementary Fig. 4a,

two-way ANOVA, $p = 0.29$). Conversely, OFS DN2 exhibited significant sexual

dimorphism, with females displaying a higher spontaneous rate and response to stationary

stimuli than males (circles, Supplementary Fig. 4b, two-way ANOVA, $p < 0.001$). The

response to a stationary stimulus was larger than spontaneous rate in males ($p = 0.015$), but

not in females (red circles, Supplementary Fig. 4b, $p = 0.35$, two-way ANOVA).

We next looked at the responses to moving full-screen 3-dimensional starfield stimuli

simulating the type of optic flow that would be generated by self-motion through space[33] (at

0.5 m/s for translations and 50°/s for rotations), after subtracting the response to the

stationary stimulus (filled circles, Supplementary Fig. 4a, b). As predicted from the receptive

fields (Fig. 1) OFS DN1 was excited by stimuli moving either up or away from the midline,

such as rightward sideslip and yaw, and upwards lift and pitch (Fig. 2b), whilst OFS DN2

showed the strongest responses to downwards lift and pitch (Fig. 2c). Both neuron types

respond strongly to clockwise roll (Fig. 2b, c), as predicted by their receptive fields (Fig. 1a,

b) and shown previously for males[14]. However, neither neuron type showed any sexual

dimorphism (Fig. 2b, c, two-way ANOVA, $p = 0.81$ and 0.92, OFS DN1 and OFS DN2,

respectively).

Sexual dimorphism in optic flow descending neurons is velocity dependent

As the velocity tuning of some LPTCs is sexually dimorphic[23,27], we tested the responses of

optic flow sensitive descending neurons using a continuous velocity step stimulus (Fig. 3,

Supplementary movie 1). Six velocities for each direction of the stimuli (e.g. anticlockwise

roll -10 to -200°/s and clockwise roll +10 to +200°/s) and a stationary control were presented

three times each in random order for 2 s each (see example, Fig. 3a).

We confirmed that the spontaneous rate and response to stationary stimuli was significantly

higher in females compared to males for OFS DN2 but not for OFS DN1 (triangles,

Supplementary Fig. 4a, b, two-way ANOVA, $p = 0.18$ and $p < 0.001$, OFS DN1 and OFS

DN2, respectively). As above, we subtracted the average response to the stationary stimulus

from the responses to moving optic flow. The resulting data show that both neuron types

exhibit sexual dimorphism to certain types of optic flow (Fig. 3e, f and Table 1). For

example, the OFS DN1 response to thrust show a significant interaction between sex and

velocity, with female neurons responding stronger to positive thrust compared to males (Fig.

3e and Table 1). The OFS DN2 response to sideslip was significantly different between males

and females, and there was a significant interaction between sex and velocity in the responses

to sideslip, lift, and thrust, with males responding stronger to sideslip, lift and thrust (Fig. 3f

and Table 1).

Morphological reconstruction suggests optic flow sensitive descending neurons could

control the wings

By recording from the descending neurons intracellularly we could iontophoretically fill them

with 3% neurobiotin following identification based on receptive field (as in Fig. 1). We found

that OFS DN1 and OFS DN2 of either sex receive their input in the part of the brain where

LPTCs have their output (Fig. 4a, b, d, e and f). Both neuron types project along the length of

the thoracic ganglion, with several fine branches along the way (Fig. 4a, b, c, g, h and i), like

Drosophila DNHS1 and DNOVS2[34]. In *Drosophila*, DNOVS2 projects more medially than

DNHS1, which also has a more distal projection, close to the haltere nerve (HN)[34]. However,

we saw no such differences between the projections of OFS DN1 and DN2 (Fig. 4a, b, c, g, h

and i). In addition, both hoverfly neuron types have prominent outputs where the prothoracic

and pterothoracic nerves likely get their inputs (T1 LN and PtN, Fig. 4a, b, c, g, h and i),

which is more similar to the outputs of *Drosophila* DNOVS1[34]. These outputs suggest that

OFS DN1 and DN2 could contribute to controlling the neck, wings and/or the forelegs.

Whilst there was no indication of extensive sexual dimorphism in the structural morphology

of these neurons, OFS DN2 appears to be slightly wider along the length of the cervical

connective in females (Supplementary Fig. 5a, b). In part, this may be due to female

hoverflies being bigger[20,21] and therefore having a wider cervical connective (Supplementary

Fig. 5c).

Wing beat amplitude changes are velocity dependent but not sexually dimorphic

Given that the morphological data suggest that both neuron types could control the wings (Fig. 4), we conducted behavioural experiments in an open-loop tethered flight arena using the same continuous velocity stimulus as in electrophysiology (see Fig. 3a). We tracked the wing beat amplitude (WBA) for each wing (Fig. 5a, b, Supplementary movie 2) using DeepLabCut models[35,36] while the tethered animal was viewing different starfield velocities (Fig. 5c, d, e and f). Changes in the sum of the left and right WBA (WBAS, Fig. 5b) reflect variations in flight force generation, potentially contributing to either lift or thrust, while the difference between left and right WBA (WBAD) indicates yaw turning behaviour[37-39] or adjustments to body orientation[40].

The male WBAS is higher when viewing stationary stimuli compared with pre-stimulation ("pre stim" in Fig. 5c, d; blue, Supplementary Fig. 4c, two-way ANOVA followed by Uncorrected Fisher's LSD test, $p < 0.01$). However, there was no sexual dimorphism in the WBAS during pre-stimulation, nor when viewing stationary stimuli (Supplementary Fig. 4c, two-way ANOVA, $p = 0.35$). When comparing responses to each type of optic flow, after subtracting the response to stationary stimuli, both the WBAD and WBAS depended strongly on velocity, however, there was no sexual dimorphism (Fig. 5g, h and Table 2).

The lack of sexual dimorphism observed in behaviour (Fig. 5g, h and Table 2) contrasts with the neural responses (Fig. 3e, f and Table 1). Additionally, it is difficult to align the WBA responses to different types of optic flow with the neuronal input. For example, in response to higher clockwise roll velocities of (100 - 200°/s), hoverflies exhibit a decreased WBAD

(positive roll, Fig. 5g) and a slightly increased WBAS (positive roll, Fig. 5h) indicating either

a leftward turn or an anticlockwise body correction. Similarly, both RHS neuron types give

strong responses to clockwise roll (positive roll, Fig. 3e, f). However, sideslip moving

leftwards also elicits a decreased WBAD (negative sideslip, Fig. 5g) and an increased WBAS

(negative sideslip, Fig. 5h) but neither neuron type responds strongly to this (negative

sideslip, Fig. 3e, f).

We found that neither lift nor thrust evoke large turning responses (Fig. 5g), however, large

and significant increases in the WBAS were observed in hoverflies experiencing the

sensation of falling (positive lift, Fig. 5h and Table 2) or being pushed backwards (negative

thrust, Fig. 5h and Table 2). Interestingly, each neuron type responds to lift and thrust in a

different manner, with OFS DN1 being excited by upwards lift and approaching thrust (Fig.

3e), whereas OFS DN2 is excited by downwards lift and receding thrust (Fig. 3f).

In electrophysiology we used a horizontally oriented visual monitor (Supplementary Fig. 6a),

whereas in behaviour the monitor was rotated 90° (Supplementary Fig. 6b and Ref [36]). To

ensure that this did not substantially affect the shape of the velocity-response functions, we

reduced the stimuli to the central square in both behaviour and electrophysiology. This

reduction caused a slight but significant change in neuronal (central square; Supplementary

Fig. 6c, two-way ANOVA, $p = 0.04$) and WBAS responses (central square; Supplementary

Fig. 6d, two-way ANOVA, $p = 0.04$). However, it did not alter the shape of the velocity

response function nor produce a significant interaction between velocity and screen size

(two-way ANOVA, $p = 0.36$ and 0.49, OFS DN2 and WBAS, respectively).

Response onset depends on stimulus type

It has been previously shown that while neuronal responses are fast[33] behavioural responses occur on a much slower time scale and differ depending on stimulus type[41,42]. We here found that there is no significant effect of sex on either neuronal or WBAS response onset (Fig. 6, two-way ANOVA, $p = 0.21$, 0.79 and 0.31, OFS DN1, OFS DN2 and WBAS, respectively). However, the onset for roll compared to lift is significantly longer for OFS DN2 and WBAS but not for OFS DN1 (Fig. 6, two-way ANOVA, $p < 0.0001$ for both OFS DN2 and WBAS compared to $p = 0.08$ for OFS DN1). Note that we recorded the WBA from above (Fig. 5a), and as such did not quantify wing rotations and adjustments outside of the horizontal plane. The WBA response onset to roll optic flow (Fig. 6c) should thus be viewed as highly conservative.

Sideslip and lift optic flow generate strong head, fore- and hind leg movements

The lack of sexual dimorphism in WBA suggests that behavioural responses to optic flow are the same between the sexes. To investigate this further, we next extracted the head angle (Fig. 7a, b), and the extension of the fore- (Fig. 7a, c) and hind legs (Fig. 7a, d). We found that the head turned significantly when viewing sideslip (Supplementary movie 2 and 3, Fig. 7e and Table 2, two-way ANOVA, $p < 0.001$ for velocity), but there was no response to the other three types of optic flow. We found no sexual dimorphism (Fig. 7e and Table 2).

We found that while the forelegs could not be seen in the dorsal view for most of the flight (see e.g. Fig. 5a), they extended anteriorly in response to high velocity sideslip, lift and

thrust, but not to roll (Fig. 7f , Table 2 and Supplementary movie 2-9). The foreleg extension

to sideslip was sexually dimorphic, with significant interaction between sex and velocity for

sideslip and lift motions (Fig. 7f and Table 2, two-way ANOVA, p = 0.04, 0.01, 0.03,

respectively). Post hoc pairwise comparisons revealed females extended their forelegs further

than males to upwards lift at 1.5 m/s (Fig. 7f, Šídák's multiple comparisons test, p = 0.01).

The hind legs were generally extended during flight (Fig. 5a, Fig. 7a) with males extending

their legs further than females (Supplementary Fig. 4d, two-way ANOVA, p = 0.032). After

subtracting the hind leg extension when viewing the stationary starfield (filled symbols,

Supplementary Fig. 4d), we found similar patterns to foreleg extension (Fig. 7f), with a

significant interaction between sex and velocity only in response to sideslip (Fig. 7g and

Table 2, two-way ANOVA, p = 0.02). We quantified the difference between the right and the

left hind leg and found that there was a significant interaction between sex and velocity in

response to roll and thrust (Fig. 7h and Table 2, two-way ANOVA, p = 0.02 for roll and

p < 0.0001 for thrust).

Looking at the different body parts together we noted that in response to upwards lift

(generating a falling sensation) the WBAS increases substantially (Fig. 5h), and the fore-

(Fig. 7f) and hind legs (Fig. 7g) all extend, potentially to brace for impact (Supplementary

movie 4 and 5). We also noted that in response to sideslip, the WBAS increases (Fig. 5h), the

WBAD indicates a turn in the direction of the optic flow (Fig. 5g), the head rotates (Fig. 7e),

and both the forelegs and hind legs extend (Fig. 7f, g). The hind legs appear to be ruddering,

with the right leg extending more when the animal is turning right (Fig. 5g, 7h,

Supplementary movie 2 and 3). There was no significant difference in the response onset of

these body parts, nor was there any sexual dimorphism (Supplementary Fig. 7).

Discussion

In the hoverfly, the optics, photoreceptors, and HS cells exhibit clear sexual

dimorphism[23,24,26,27]. In line with this, our findings reveal that optic flow sensitive descending

neurons also show sexually dimorphic velocity response functions (Fig. 3). Yet, despite the

multi-level anatomical and neural sexual dimorphism, the wing beat amplitude (Fig. 5) and

head movements (Fig. 7) remain mostly monomorphic, while leg movements exhibit clear

sex-specific differences. Moreover, differences between neural and behavioural velocity

response functions (Fig. 3, 5, 7) and response latency (Fig. 6) highlight the likely

involvement of downstream processing mechanisms that could reconcile sex-specific sensory

encoding with conserved flight control.

Some hoverfly flight behaviours are strongly sexually dimorphic, such as those related to

courtship and mating. For example, male hoverflies defend territories from a hovering stance

from which they pursue and capture female conspecifics at high speeds[19]. In contrast, other

behaviours, including cruising speeds in field[31] and indoor settings[32] do not differ between

sexes, and are much slower (median around 0.3 m/s[31,32]) than the 10 m/s that can be reached

during outdoor pursuit[19]. In comparison, our results show limited sex-related differences in

WBA in response to optic flow speeds up to 2 m/s (Fig. 5), yet we detected pronounced

sexual dimorphism in neural responses at velocities as low as 0.5 m/s (Fig. 3, Table 1). This

suggests that sensory processing differences emerge at lower velocities than the velocities

where motor output diverges. Moreover, the WBAS response latencies were much longer

than the corresponding neural activity (Fig. 6), implying that temporal integration or

additional circuit-level modulation may delay and refine motor execution. Together, these

findings suggest a complex transformation between sex-specific sensory encoding and

conserved motor behaviour, potentially mediated by downstream integration or additional

alternate neural pathways that bypass the descending neurons studied here.

In *Drosophila* there are around 1000 descending neurons, of which at least 35 types receive

their input in the posterior surface of the brain, which includes the inferior posterior slope,

where the LPTC outputs are found[6,7,9]. However, only a subset of these constitute the 29

descending neuron types that project to the wing motor neuropil[9]. It is thus likely that the

WBA that we recorded here (Fig. 5) reflects contributions from descending neurons beyond

the two that we have characterised (Fig. 1, 3) in this study. For example, many looming

sensitive neurons also respond to widefield optic flow[14,43] and could therefore play a role in

the WBA changes that we detected (Fig. 5). While the morphology of both neurons (Fig. 4)

suggested involvement in wing motor control, there is currently no direct evidence

implicating them in the control of wingbeat amplitude. Interestingly, DNOVS2

in *Drosophila*, a physiologically similar neuron to OFS DN2[14], has been indirectly linked to

rapid turning behaviour[10]. Furthermore, silencing HS and VS cells, the presumed presynaptic

LPTCs to these descending neurons, results in reductions in WBA only at higher stimulus

speeds[44]. These findings suggest that HS and VS cells, along with their downstream targets,

may be specialized for driving fast optomotor responses under high-speed visual motion

(>180°/s), rather than broadly regulating WBA across all velocities, including those

examined in this study.

Beyond their potential involvement in wingbeat amplitude modulation, OFS DNs may play a

role in coordinating head and body positioning during flight, especially in contexts requiring

precise visual alignment and rapid manoeuvring. Indeed, the *Drosophila* physiological

homologs of OFS DN1 and OFS DN2[14], DNHS1 and DNOVS2, project to the neck motor

neuropil[9], and have been implicated in controlling head movements, abdominal ruddering,

and engagement with the haltere motor system for flight stabilization[10]. While we did not

quantify abdominal or haltere movements, we found that the head turned when viewing

sideslip optic flow (Fig. 7e), and that the velocity dependence was similar to neural (Fig. 3e)

and WBAD (Fig. 5g) responses. While we did not detect any head movements to roll, lift or

thrust, this could be a technical limitation of recording from above (Fig. 5a). Indeed, many of

the 29 descending neuron types that project to the wing motor neuropil, also project to the

neck motor neuropil[9], suggesting that synchronized responses are not unexpected. Neither are

the similar responses of the fore- and hind legs surprising (Fig. 7, f and g), considering that in

Drosophila many of the ~30 descending neuron types that project to the forelegs, also project

to the mid- and hind legs[9].

Our WBA recordings would also be affected by filming from above and thus lacking other

nuanced changes in wing angles. Thus, the sexual dimorphism in neural responses (Fig. 3)

could reflect an evolutionary tuning of visuomotor pathways in males, optimized for fast,

directional adjustments rather than gross changes in WBA in the horizontal plane. For

instance, whilst changes in WBAS are similar when generating either lift or thrust (Fig. 5h),

body pitch dynamically adjusts their ratio, driving behavioural variation[45]. Thus, OFS DNs

may serve as integral components of a broader flight control architecture, interfacing optic

flow detection with dynamic body and head positioning systems to support complex, sex-

specific behavioural outcomes, particularly in males engaging in high-speed pursuits.

Indeed, flight speed in insects results from a complex integration of multiple kinematic

parameters, not solely from changes in wingbeat amplitude. Many species refine their

aerodynamic output by modulating wingbeat frequency, angle of attack, wing tip trajectory,

deviations from the mean stroke plane and through precise adjustments to the timing and

duration of the up- and downstrokes[46]. These control strategies support agile manoeuvring,

particularly during visually guided behaviours such as the high-speed pursuits undertaken by

male hoverflies. Furthermore, the smaller body size of male hoverflies compared to

females[20,21] may confer biomechanical advantages, including reduced mass, facilitating faster

acceleration, heightened responsiveness, and lower metabolic costs for executing flight

manoeuvres[22,47], which do not appear in a tethered flight set-up. Such enhanced agility is

likely crucial for rapid direction changes required during courtship and may enable male

hoverflies to outperform female flies without relying on increased wingbeat amplitude.

Taken together, our findings reveal significant differences between sexually dimorphic

sensory encoding and conserved motor output in hoverfly flight at cruising speeds. Although

OFS DNs and their upstream visual circuits display clear sex-specific tuning, wingbeat

amplitude and head angle changes in response to optic flow stimuli remain relatively similar

between the sexes, suggesting additional mechanisms downstream of sensory input. This

likely reflects a complex interplay of biomechanical properties, multisensory integration and

circuit-level modulation, each shaping and refining behavioural outcomes to meet distinct

demands, preserving the consistency of low-speed manoeuvres whilst enabling sex-specific

tuning during high-speed pursuits.

Methods

Animals

For all experiments, male and female *Eristalis tenax* were reared and housed as described

previously[21]. Briefly, eggs were collected from females captured under permit in Wittunga

Botanic Garden, Adelaide, South Australia. Upon hatching, larvae were reared in a rabbit

dung slurry until third instar larvae emerged to pupate. Eclosion occurred 1-2 weeks post-

pupation. Adult hoverflies were used for behavioural experiments at 17-87 days post-

eclosion, for intracellular electrophysiology and subsequent morphological reconstruction at

38-54 days, and for extracellular electrophysiology at 8-204 days.

Electrophysiology

Before recording the animal was immobilised, mounted dorsal side down (Supplementary

Fig. 6a) and secured using a mixture of beeswax and resin. A small region of cuticle was

removed at the anterior end of the thorax to expose the cervical connective. If required, any

excessive gut or tracheal tissue obstructing the recording site was removed and a small

volume of PBS was added to prevent drying within the ventral cavity. A fine wire hook was

positioned under the cervical connective for mechanical support, and a silver wire was

inserted into the cavity to serve as a reference electrode and grounding wire[14].

For extracellular recordings, a sharp tungsten microelectrode (2 MΩ, polyimide-insulated;

Microprobes) was inserted into the cervical connective[14]. Signals were amplified 1000 times

and band-pass filtered between 10–3000 Hz using a DAM50 differential amplifier (World

Precision Instruments), followed by noise reduction with a HumBug (Quest Scientific). Data

were digitized via a PowerLab 4/30 interface (ADInstruments) and acquired at 40 kHz. Spike

sorting was performed in LabChart 7 Pro (ADInstruments) based on the amplitude and width

of individual action potentials.

For intracellular recordings, aluminosilicate electrodes were pulled using a Sutter P-1000

micropipette puller, achieving a resistance of approximately 40-70 MΩ. Electrode tips were

filled with 3% neurobiotin (Vector Laboratories), then backfilled with 1 M KCl using a

syringe, leaving a small air bubble between the two solutions. Electrodes were inserted into

the cervical connective for recording, and the resulting signal was amplified using an

Axoclamp-2B amplifier (Axon Instruments), followed by 50 Hz noise reduction with a

HumBug (Quest Scientific). Data acquisition and digitization were performed at 10 kHz

using an NI USB-6210 16-bit data acquisition card (National Instruments) and the MATLAB

Data Acquisition Toolbox (Mathworks), using in house software

(https://github.com/HoverflyLab/SampSamp).

Morphological reconstructions

Following intracellular recordings, neurons were stained iontophoretically with neurobiotin using currents in the 1 nA range for 3–12 minutes. The nervous system was then carefully dissected and fixed in 4% paraformaldehyde overnight. Tissue was incubated with a Cy3-streptavidin conjugate (1:100; Jackson ImmunoResearch) for 2 hours, then dehydrated through an ethanol series (50–100%) for 15-20 minutes per step. After washing in PBT, the tissue was cleared in RapiClear (SUNJIN Lab) and mounted with spacers. Imaging was performed using a Zeiss LSM 880 Fast Airyscan confocal microscope at the institutional microscopy facility. Neuron morphology and cervical connective width were quantified using ImageJ[48].

Tethered flight

Prior to flight recordings, hoverflies were tethered at a 32° angle (Supplementary Fig. 6b) using a beeswax–resin mixture to a small pin inserted into a hypodermic needle (BD Microlance, 23G × 1¼"). Flight was initiated by manually providing airflow for 1–10 minutes until consistent flight behaviour was observed. Once positioned facing the centre of the visual monitor (Supplementary Fig. 6b), hoverflies were filmed from above at 100 Hz using a Sony PlayStation 3 Move Eye Camera (SLEH- 00448, Sony) with the IR filter removed, and equipped with an infrared pass filter (R72 INFRARED, 49 mm, HOYA; for details see Ref[36]). Illumination was provided by infrared LEDs inserted into USB lights (JANSJÖ LED USB lamp, IKEA) and a Musou Black (Shin Kokushoku Musou black,

KOYO Orient Japan) surface was placed beneath the hoverfly to enhance contrast and

minimize optical interference.

We used DeepLabCut (DLC) version 2.3.3[35] to train a model to track the thorax, the peak

downstroke angle, referred to as the wing beat amplitude (WBA), of the left and right wing

(WBA$_L$ and WBA$_R$; Fig. 5a, b), as described previously[36]. In addition, we tracked the dorso-

medial head (yellow, green, Fig. 7a, b), the proximal and distal points of the forelegs (blue,

Fig. 7a, c), the hind leg knee and the lateral, mid-abdomen (magenta, green, Fig. 7a, d). For

this purpose, we first manually labelled the anatomical landmarks (Fig. 5a and Fig. 7a),

across 16 extracted video frames per individual from four hoverflies (2 males, 2 females). In

addition to examples where the hoverfly was not flying, these frames included responses to

yaw rotation and forward translation. The DLC model was trained for 300,000 iterations,

yielding train and test errors of 1.2 and 1.16 pixels, respectively.

To identify potential tracking errors from DLC, we smoothed the WBA time series using

MATLAB's *smooth* function with both *loess* and *rloess* methods. Because *rloess* is more

resistant to outliers, we used it to detect abnormal data points. For each wing, if the absolute

difference between the *loess*- and *rloess*-smoothed signals exceeded 5% for more than 1 s,

the data were excluded due to suspected tracking artifacts. The smoothing was used only for

error detection; all subsequent analyses were performed on the unsmoothed data. In addition,

we excluded data if the WBA of either wing dropped below 40° for at least 0.5 s, as this

indicated a cessation of flight. Finally, entire trials were excluded if the hoverfly was not

flying for more than 50% of the trial duration. Head and leg kinematic data were excluded

whenever the corresponding WBA data were excluded, ensuring consistent trial inclusion

across behavioural metrics.

Visual stimuli

Visual stimuli were generated using custom software

(https://github.com/HoverflyLab/FlyFly/releases/tag/v4.2.3) written in MATLAB,

incorporating the Psychophysics Toolbox[49,50]. All screens had a refresh rate of 165 Hz and a

linearized contrast with a mean illuminance of 200 Lux. For intracellular recordings, the

hoverfly was placed 13 cm away from a ViewSonic screen with a resolution of 2560 × 1440

pixels, corresponding to 143° × 107° of the visual field. For extracellular recordings, the

hoverfly was placed 6.5 cm away from a 2560 × 1440 pixel Asus screen, yielding a projected

visual field of 155° × 138° (Supplementary Fig. 6a). For behavioural recordings, the hoverfly

was placed 10 cm from a vertically orientated Asus screen (1440 × 2560 pixel,

Supplementary Fig. 6b) producing a projected size of 118° × 142°. To evaluate the impact of

different screen orientations in electrophysiology and behaviour, visual stimuli were

presented either full-screen or using the central 1440 × 1440 pixel square (Supplementary

Fig. 6c, d).

Receptive field mapping

To map each neuron's receptive field, we presented local sinusoidal gratings (average

38 × 38°) drifting in 8 directions for 0.36 s each, across 48 overlapping locations, as

described in detail previously[14]. Stimuli were full contrast, with an average spatial frequency

of 0.14 cycles/° and a temporal frequency of 5 Hz. At each location, we calculated the local

maximum spike frequency (red, inset, Fig. 1b, e). After subtracting the spontaneous rate,

calculated for 0.8 s preceding stimulus onset (dotted line, inset, Fig. 1b, e), we interpolated

the resulting local maximum responses to a ten-fold higher spatial resolution (colour coding,

Fig. 1a, d). We then quantified the 50% response maximum using Matlab's *contour* function

(black, Fig. 1b, e). We created a polygon-shape of this 50% contour line using Matlab's

polyshape function to calculate the receptive field size, defined as its width and height

(Supplementary Fig. 2f), and used the *centroid* function to identify the centre of this

polyshape (red circle, Fig. 1b, e). As recordings were performed with the animal ventral side

up (Supplementary Fig. 3a, 6a), receptive fields were rotated to display them with the dorsal

visual field up.

Next, at each location, we fit a cosine function to the responses to the different directions of

motion, to extract the local preferred direction (LPD) and local motion sensitivity (LMS;

inset, Fig. 1c, f). These were visualized as vectors where angle indicates LPD and length

indicates LMS (arrows, Fig. 1a, d). We defined the receptive field preferred direction as the

median of LPDs at locations where LMS exceeded 50% of the maximum (black and red

arrows; Fig. 1c, f), using *circ_median* from the CircStat toolbox for MATLAB[51]. We

calculated the LPD variance using the *circ_var* functions from CircStat toolbox for

MATLAB[51], of LPDs at locations where LMS exceeded 50% of the maximum (black and red

arrows; Fig. 1c, f).

We extracted unpublished receptive field data from 100 reference neurons (other data from

five of these optic flow sensitive descending neurons has been published previously[28]) to set

exclusion criteria (red shading, Supplementary Fig. 2a, b and c) and to classify the OFS DNs

used in the rest of this study. OFS DN1 LHS neurons were defined by receptive field

preferred directions between 120° and 200° (light green, Fig. 1g, i). OFS DN1 RHS neurons

were defined by receptive field preferred directions between 340° and 60° (dark green, Fig.

1g, i). OFS DN2 neurons were classified by receptive field preferred directions between 220°

and 320° (yellow and orange, Fig. 1g, i), with the azimuthal location of the receptive field

centre determining whether they were LHS or RHS (yellow and orange, Fig. 1h).

A polar plot where the distance from origo indicates azimuthal position, and the location

along the radius the preferred direction, confirms that the LHS and RHS of OFS1 and OFS2

cluster distinctly (Supplementary Fig. 1a). To test whether incorporating additional receptive

field parameters would reveal further neuronal subtypes, we performed k-means clustering in

Matlab using z-score normalised data. The additional parameters included receptive field

centre elevation, receptive field height and width (see Supplementary Fig. 2f), maximum

LMS (see Supplementary Fig. 2a), the number of positions with LMS values exceeding 50%

of the maximum (see Supplementary Fig. 2b), and LPD variance (see Supplementary Fig.

2c). However, the Callinski-Harabasz[52] values indicate that adding additional receptive field

parameters actually reduced clustering performance (Supplementary Fig. 1b). In contrast,

clustering based solely on preferred direction and azimuthal location yielded four well-

defined groups (Supplementary Fig. 1c), and produced the highest Callinski-Harabasz value

(Supplementary Fig. 1b).

Directional sensitivity

Full-screen sinusoidal gratings were presented at full contrast, using the same spatial (0.14

cycles/°) and temporal (5 Hz) frequencies as in receptive field mapping. For each stimulus

direction, mean spike frequency was calculated over the 1 s stimulus duration, excluding the

first 100 ms to avoid onset transients[53].We fit a cosine function to the responses to the eight

different directions of motion, to extract the preferred direction and response amplitude

(Supplementary Fig. 3b), and then plotted these values for OFS DN1 (Supplementary Fig. 3c)

and DN2 neurons (Supplementary Fig. 3d). To account for the ventral-side-up recording

position (Supplementary Fig. 3a) directional responses were rotated 180° (Supplementary

Fig. 3f, g and h). In addition, to display all neurons as RHS, and assuming mirror-symmetry,

we flipped LHS neurons across the midline (Supplementary Fig 2i, j, k and l).

Responses to optic flow

We used a starfield stimulus to generate the type of perspective-corrected optic flow that

would have been seen by the hoverfly if it was moving through a space of 2 cm diameter

spheres (for details, see Ref[14,33]). These simulated translations at 0.5 m/s (sideslip, lift, and

thrust) or rotations (pitch, yaw, and roll), at 50°/s. To quantify neural responses, the mean

spike frequency was calculated over the 0.97 s stimulus duration, excluding the first 0.1 s to

avoid onset-related transients[53]. The spontaneous firing rate, averaged across 0.48 s

immediately preceding stimulus onset (open circles, Supplementary Fig 4a, b), was

subtracted from the response.

Velocity response functions

We used four types of optic flow: three translations (sideslip, lift and thrust, Supplementary

movie 2-7) and one rotation (roll, Supplementary movie 1, 8, 9). Translations were presented

at velocities of −2, −1.5, −1, −0.4, −0.2, −0.1, 0, 0.1, 0.2, 0.4, 1, 1.5, and 2 m/s, while

rotations were presented at 200, −150, −100, −40, −20, −10, 0, 10, 20, 40, 100, 150, and

200°/s. The sign of the velocity indicates the direction of motion as seen by the fly when

corrected for its position, with positive values corresponding to counterclockwise roll,

leftward sideslip, downward lift, and thrust moving away. Conversely, negative values

indicate clockwise roll, rightward sideslip, upward lift, and thrust moving toward the

hoverfly. The upper limit was defined by the movement of the individual dots within the

starfield stimulus between frames, i.e. was limited by the refresh rate of the screen.

Each trial consisted of 39 stimuli (13 unique velocities × 3 repetitions), presented in a random

order, with each stimulus lasting 2 s, immediately followed by the next velocity. Trials began

with a 1 s blank screen, which served as the pre-stimulation baseline (Fig. 3a and Fig. 5c).

Each optic flow condition was repeated multiple times, resulting in at least nine repetitions

for each velocity and neuron, and five repetitions for each velocity and animal in behaviour.

For neuronal recordings, velocity response function trials were interleaved with the optic

flow stimuli described above. In behavioural experiments, a flight refresher sequence

interspersed each velocity tuning trial. This consisted of sinusoidal gratings (200°

wavelength, 5 Hz) drifting rightward, leftward, and rightward again for 4 s each.

Neural responses were quantified as the mean spike frequency during the final second of

stimulation (grey shaded areas, Fig. 3a–c). We then calculated the median across trials for

each velocity (Fig. 3d). Pre-stimulation activity (spontaneous rate) was measured over a 2 s

window immediately preceding stimulus onset. Neural responses are displayed after

subtracting the response when viewing a stationary stimulus (filled symbols, Supplementary

Fig. 4a, b).

In behavioural experiments, we extracted the mean wing beat amplitude of the left and right

wings ($\overline{WBA_L}$, $\overline{WBA_R}$, Fig. 5b) during the final second of stimulation (grey shaded areas, in

Fig. 5 c, d and e). For each velocity, we calculated the WBA difference (WBAD, defined as

$\overline{WBA_L} - \overline{WBA_R}$) and the WBA sum (WBAS, defined as $\overline{WBA_L} + \overline{WBA_R}$, Fig. 5b). We then

calculated the median across repetitions for each individual (Fig. 5f). Pre-stimulation

responses were averaged over a 0.5 s time window immediately preceding stimulus onset.

WBA responses are displayed after subtracting the response when viewing a stationary

stimulus (filled symbols, Supplementary Fig. 4c).

We defined the head angle as the angle between a straight line joining the 2 tracked points on

the head (using *polyfit* in matlab, Fig. 7a, b) and the longitudinal axis of the thorax (Fig. 7a,

b). As we filmed in one plane only, this measurement can be caused by a combination of

head rotations, but seem to be dominated by roll rotations (Supplementary movie 2-9). We

measured the distance between the proximal and distal points of the forelegs (Fig. 7a, c).

Note that the forelegs were mostly hidden under the animal from our dorsal view, and could

only be seen when extended anteriorly (e.g. Fig. 5a, see also Supplementary movie 2-9). Data

are displayed as the mean extension of the left and right foreleg. We measured the distance

between the hind leg knee and a lateral point on the mid-abdomen (Fig. 7a, d). Hind leg data

are displayed as the mean or the difference of the left and right hind leg. All body parts data

are displayed after subtracting the response when viewing a stationary stimulus (filled

symbols show hindleg data, Supplementary Fig. 4d).

Response onset

For neuronal recordings, we compared onset times for clockwise roll (+50°/s) and either

upwards lift (+0.5 m/s) for OFS DN 1 or downwards lift (-0.5 m/s) for OFS DN2, stimuli

which generated strong responses in these neurons (Fig. 3 e, f). Mean spike frequency was

calculated over the 0.97 s stimulus duration, excluding the first 0.1 s to avoid onset

transients[53]. Onset was defined as the first time point after the first 0.1 s, where spike rate

exceeded 80% of the mean.

For behaviour, we used WBA responses to roll (-200°/s) and lift (+2 m/s). Mean WBAS was

defined as the average response during the final second of stimulation, and onset was defined

as the first time point where WBAS exceeded 80% of this mean. We also quantified WBAS,

head, fore-and hind leg onsets to sideslip (+2 m/s). For each body part, we quantified the

mean response during the final second of stimulation, and onset was defined as the first time

point where the response exceeded 80% of this mean.

Data availability and statistics

All data and analysis scripts have been submitted to DataDryad (private link for peer review: doi:10.5061/dryad.tb2rbp0fd).

Throughout the text n refers to individual repetitions, whereas N refers to individual neurons (electrophysiology) or animals (behaviour). All data are presented as median and interquartile range, unless otherwise indicated.

Statistical analyses were performed in Prism 10.4.0 (GraphPad Software), except for circular statistics, which were conducted using the CircStat toolbox for MATLAB[51]. The results of the statistical tests are given in the text, figure legends and in Table 1 and Table 2. For behavioural data the p-values were Bonferroni adjusted[54] for multiple comparisons (Table 2).

Acknowledgements

We thank Biomedical Engineering at SAHLN and the Botanic Gardens of Adelaide for their ongoing support. We thank the editor and the two Reviewers for their feedback which greatly improved our paper. This research was funded by the US Air Force Office of Scientific Research (AFOSR, FA9550-19-1-0294 and FA9550-23-1-0473) and the Australian Research Council (ARC, DP210100740, DP230100006 and DP250104770).

Declaration of interests

The authors declare no competing interests.

References

1   Mauss, A. S. & Borst, A. Optic flow-based course control in insects. *Curr Opin*
Neurobiol **60**, 21-27, doi:10.1016/j.conb.2019.10.007 (2020).

2   Pierantoni, R. A look into the cock-pit of the fly. The architecture of the lobular plate.
Cell Tissue Res **171**, 101-122 (1976).

3   Zhao, A. *et al.* A comprehensive neuroanatomical survey of the
Drosophila Lobula Plate Tangential Neurons with predictions for their
optic flow sensitivity. *bioRxiv*, 2023.2010.2016.562634,
doi:10.1101/2023.10.16.562634 (2023).

4   Franz, M. O. & Krapp, H. G. Wide-field, motion-sensitive neurons and matched
filters for optic flow fields. *Biol Cybern* **83**, 185-197 (2000).

5   Krapp, H. G., Hengstenberg, B. & Hengstenberg, R. Dendritic structure and
receptive-field organization of optic flow processing interneurons in the fly. *J*
Neurophysiol **79**, 1902-1917. (1998).

6   Wei, H., Kyung, H. Y., Kim, P. J. & Desplan, C. The diversity of lobula plate
tangential cells (LPTCs) in the Drosophila motion vision system. *Journal of*
Comparative Physiology A **206**, 139-148, doi:10.1007/s00359-019-01380-y (2020).

7   Strausfeld, N. J. & Bassemir, U. K. Lobula plate and ocellar interneurons converge
onto a cluster of descending neurons leading to neck and leg neuropil in *Calliphora*
erythrocephala. Cell Tissue Res **240**, 617-640 (1985).

8   Strausfeld, N. J. & Seyan, H. S. Convergence of visual, haltere, and prosternai inputs
at neck motor neurons of *Calliphora erythrocephala. Cell and Tissue Research* **240**,
601-615 (1985).

9   Namiki, S., Dickinson, M. H., Wong, A. M., Korff, W. & Card, G. M. The functional
organization of descending sensory-motor pathways in *Drosophila. eLife* **7**, e34272,
doi:10.7554/eLife.34272 (2018).

10 Suver, M. P., Huda, A., Iwasaki, N., Safarik, S. & Dickinson, M. H. An Array of
Descending Visual Interneurons Encoding Self-Motion in Drosophila. *J Neurosci* **36**,
11768-11780, doi:10.1523/JNEUROSCI.2277-16.2016 (2016).

11 Erginkaya, M. *et al.* A competitive disinhibitory network for robust optic flow
processing in Drosophila. *bioRxiv*, 2023.2008.2006.552150,
doi:10.1101/2023.08.06.552150 (2023).

12 Haikala, V., Joesch, M., Borst, A. & Mauss, A. S. Optogenetic control of fly
optomotor responses. *J Neurosci* **33**, 13927-13934, doi:10.1523/jneurosci.0340-
13.2013 (2013).

13 Pokusaeva, V. O., Satapathy, R., Symonova, O. & Joesch, M. Bilateral interactions of
optic-flow sensitive neurons coordinate course control in flies. *Nat Commun* **15**, 8830,
doi:10.1038/s41467-024-53173-w (2024).

14 Nicholas, S., Leibbrandt, R. & Nordström, K. Visual motion sensitivity in descending
neurons in the hoverfly. *J Comp Physiol A* **206**, 149-163, doi:10.1007/s00359-020-
01402-0 (2020).

15 Haag, J., Wertz, A. & Borst, A. Integration of lobula plate output signals by
DNOVS1, an identified premotor descending neuron. *J Neurosci* **27**, 1992-2000,
doi:10.1523/jneurosci.4393-06.2007 (2007).

16 Wertz, A., Borst, A. & Haag, J. Nonlinear integration of binocular optic flow by
DNOVS2, a descending neuron of the fly. *J Neurosci* **28**, 3131-3140,
doi:10.1523/jneurosci.5460-07.2008 (2008).

17 Collett, T. S. & Land, M. F. Visual control of flight behaviour in the hoverfly, *Syritta*
pipiens L. *J Comp Physiol A* **99**, 1-66 (1975).

18 Wellington, W. & Fitzpatrick, S. Territoriality in the drone fly, *Eristalis tenax*
(Diptera, Syrphidae). *Can Entomol* **113**, 695-704 (1981).

19 Collett, T. S. & Land, M. F. How hoverflies compute interception courses. *Journal of*
comparative physiology **125**, 191-204, doi:10.1007/BF00656597 (1978).

20 Dankova, K., Nicholas, S. & Nordstrom, K. Temperature during pupal development
affects hoverfly developmental time, adult life span, and wing length. *Ecol Evol* **13**,
e10516, doi:10.1002/ece3.10516 (2023).

21 Nicholas, S., Thyselius, M., Holden, M. & Nordström, K. Rearing and long-term
maintenance of *Eristalis tenax* hoverflies for research studies. *JoVE*, e57711,
doi:doi:10.3791/57711 (2018).

22 Dudley, R. Mechanisms and implications of animal flight maneuverability. *Integr*
Comp Biol **42**, 135-140, doi:10.1093/icb/42.1.135 (2002).

23 Straw, A. D., Warrant, E. J. & O'Carroll, D. C. A 'bright zone' in male hoverfly
(*Eristalis tenax*) eyes and associated faster motion detection and increased contrast
sensitivity. *J Exp Biol* **209**, 4339-4354, doi:10.1242/jeb.02517 (2006).

24 van Hateren, J. H., Hardie, R. C., Rudolph, A., Laughlin, S. B. & Stavenga, D. G. The
bright zone, a specialized dorsal eye region in the male blowfly *Chrysomyia*
megacephala. J Comp Physiol A **164**, 297-308 (1989).

25 Hornstein, E. P., O'Carroll, D. C., Anderson, J. C. & Laughlin, S. B. Sexual
dimorphism matches photoreceptor performance to behavioural requirements. *Proc R*
Soc Lond B Biol Sci **267**, 2111-2117., doi:10.1098/rspb.200.1257 (2000).

26 Nordström, K., Barnett, P. D., Moyer de Miguel, I. M., Brinkworth, R. S. A. &
O'Carroll, D. C. Sexual dimorphism in the hoverfly motion vision pathway. *Curr Biol*
18, 661-667, doi:10.1016/j.cub.2008.03.061 (2008).

27 Barnett, P. D., Nordström, K. & O'Carroll, D. C. Motion adaptation and the velocity
coding of natural scenes. *Curr Biol* **20**, 994-999, doi:10.1016/j.cub.2010.03.072
(2010).

28 Nicholas, S. & Nordström, K. Facilitation of neural responses to targets moving
against optic flow. *Proc Natl Acad Sci U S A* **118**, doi:10.1073/pnas.2024966118
(2021).

29 Ghosh, A., Nicholas, S., Nordström, K., Nowotny, T. & Knight, J. Understanding the
mechanism of facilitation in hoverfly TSDNs. *PLOS Computational Biology* **21**,
e1012986, doi:10.1371/journal.pcbi.1012986 (2025).

30 Nicholas, S., Supple, J., Leibbrandt, R., Gonzalez-Bellido, P. T. & Nordström, K.
Integration of small- and wide-field visual features in Target-Selective Descending
Neurons of both predatory and non-predatory dipterans. *J Neurosci* **38**, 10725-10733,
doi:10.1523/jneurosci.1695-18.2018 (2018).

31 Thyselius, M., Gonzalez-Bellido, P. T., Wardill, T. J. & Nordstrom, K. Visual
approach computation in feeding hoverflies. *J Exp Biol* **221**, doi:10.1242/jeb.177162
(2018).

32 Thyselius, M. *et al.* Hoverfly (Eristalis tenax) pursuit of artificial targets. *J Exp Biol*
226, doi:10.1242/jeb.244895 (2023).

33 Leibbrandt, R., Nicholas, S. & Nordström, K. The impulse response of optic flow-
sensitive descending neurons to roll m-sequences. *J Exp Biol* **224**,
doi:10.1242/jeb.242833 (2021).

34 Suver, M. P., Huda, A., Iwasaki, N., Safarik, S. & Dickinson, M. H. An array of
descending visual interneurons encoding self-motion in *Drosophila*. *J Neurosci* **36**,
11768 (2016).

35 Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts
with deep learning. *Nat Neurosci* **21**, 1281-1289, doi:10.1038/s41593-018-0209-y
(2018).

36 Ogawa, Y. *et al.* Combining Unity with machine vision to create low latency, flexible
and simple virtual realities. *Methods in Ecology and Evolution* **16**, 126-144,
doi:https://doi.org/10.1111/2041-210X.14449 (2025).

37 Götz, K. G. Flight control in *Drosophila* by visual perception of motion. *Kybernetik*
4, 199-208 (1968).

38 Maimon, G., Straw, A. D. & Dickinson, M. H. Active flight increases the gain of
visual motion processing in *Drosophila*. *Nat Neurosci* **13**, 393-399, doi:nn.2492 [pii]
10.1038/nn.2492 [doi] (2010).

39 Tammero, L. F., Frye, M. A. & Dickinson, M. H. Spatial organization of visuomotor
reflexes in Drosophila. *J Exp Biol* **207**, 113-122, doi:10.1242/jeb.00724 (2004).

40 Ristroph, L. *et al.* Discovering the flight autostabilizer of fruit flies by inducing aerial
stumbles. *Proc Natl Acad Sci U S A* **107**, 4820-4824, doi:10.1073/pnas.1000615107
(2010).
41 Schnell, B., Weir, P. T., Roth, E., Fairhall, A. L. & Dickinson, M. H. Cellular
mechanisms for integral feedback in visually guided behavior. *Proc Natl Acad Sci U*
S A **111**, 5700-5705, doi:1400698111 [pii]
10.1073/pnas.1400698111 [doi] (2014).
42 Theobald, J. C., Ringach, D. L. & Frye, M. A. Dynamics of optomotor responses in
Drosophila to perturbations in optic flow. *J Exp Biol* **213**, 1366-1375, doi:213/8/1366
[pii]
10.1242/jeb.037945 [doi] (2010).
43 Nicholas, S., Ogawa, Y. & Nordström, K. Dual receptive fields underlying target and
wide-field motion sensitivity in looming-sensitive descending neurons. *eNeuro* **10**,
ENEURO.0188-0123.2023, doi:10.1523/eneuro.0188-23.2023 (2023).
44 Kim, A. J., Fenk, L. M., Lyu, C. & Maimon, G. Quantitative Predictions Orchestrate
Visual Signaling in Drosophila. *Cell* **168**, 280-294 e212,
doi:10.1016/j.cell.2016.12.005 (2017).
45 Zanker, J. M. The Wing Beat of Drosophila-Melanogaster .3. Control. *Philos T R Soc*
B **327**, 45-64, doi:DOI 10.1098/rstb.1990.0042 (1990).
46 Sane, S. P. The aerodynamics of insect flight. *Journal of Experimental Biology* **206**,
4191-4208, doi:10.1242/jeb.00663 (2003).
47 Niven, J. E. & Scharlemann, J. P. W. Do insect metabolic rates at rest and during
flight scale with body mass? *Biol Letters* **1**, 346-349, doi:10.1098/rsbl.2005.0311
(2005).
48 Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* **9**, 671-675, doi:10.1038/nmeth.2089 (2012).
49 Brainard, D. H. The Psychophysics Toolbox. *Spat Vis* **10**, 433-436 (1997).
50 Pelli, D. G. The VideoToolbox software for visual psychophysics: transforming
numbers into movies. *Spat Vis* **10**, 437-442 (1997).

51   Berens, P. CircStat: A MATLAB toolbox for circular statistics. *Journal of Statistical*
Software **31**, 1 - 21, doi:10.18637/jss.v031.i10 (2009).
52   Caliński, T. & Harabasz, J. A dendrite method for cluster analysis. *Communications*
in Statistics **3**, 1-27, doi:10.1080/03610927408827101 (1974).
53   Nicholas, S. & Nordstrom, K. Persistent Firing and Adaptation in Optic-Flow-
Sensitive Descending Neurons. *Curr Biol* **30**, 2739-2748 e2732,
doi:10.1016/j.cub.2020.05.019 (2020).
54   Abdi, H. in *Encyclopedia of measurement and statistics*   (ed N. J. Salkind) (Sage,
2007).
55   Kane, G. A., Lopes, G., Saunders, J. L., Mathis, A. & Mathis, M. W. Real-time, low-
latency closed-loop feedback using markerless posture tracking. *eLife* **9**,
doi:10.7554/eLife.61909 (2020).
768
769

 Figure legends

 Fig. 1. Classification of optic flow sensitive (OFS) descending neurons (DN) in *Eristalis*

tenax. **a** Receptive field of a representative OFS DN1 recorded from a male hoverfly. Colour

coding indicates the local maximum spike frequency (red, inset, panel b), the direction of the

arrows shows the local preferred direction (LPD, red arrowhead, panel c) and their length the

local motion sensitivity (LMS, red, panel c). As the stimuli were not perspective corrected,

the receptive field map reflects the visual monitor. **b** Contour line representing the 50%

receptive field boundary based on local maximum spike frequency (colour coding, panel a).

Inset: example response at one central location to eight directions of motion (black circles),

showing local maximum spike frequency (local max, red line) above spontaneous rate (spont,

black dotted line). **c** Local preferred direction (LPD) map of the same example neuron at

locations where local motion sensitivity (LMS) exceeds 50% of the maximum. Inset:

example response at one location to eight directions of motion with a sinusoidal fit (black

line), illustrating LMS (red line) and LPD (red arrowhead). The data in the inset corresponds

to the red arrow. **d** Receptive field of a representative male OFS DN2. **e** 50% receptive field

contour and receptive field centre of the neuron shown in panel d. **f** Preferred direction map

of the same OFS DN2. **g** Distribution of receptive field preferred directions across 100 male

reference neurons, colour-coded by neuron classification. Dashed lines in corresponding

colours indicate the thresholds used for neuron type classification. **h** Receptive field centres

of the same 100 reference neurons, using the same colour coding. **i** Relationship between

preferred direction and receptive field centre for the 100 neurons. Note that the scale is the

same for the azimuth and elevation, and the scale bar in panels a, d, h and i correspond to

both the x- and y-axis.

Fig. 2. Sex-based comparison of direction sensitivity in OFS DNs. a Polar plot showing

the preferred direction of male (blue) and female (red) OFS DNs in response to a full-screen,

full-contrast sinusoidal grating (spatial wavelength 7º, temporal frequency 5 Hz) moving in

eight different directions (see also Supplementary Fig. 3). Individual data points represent the

response amplitude and preferred direction of each OFS DN (red, Supplementary Fig. 3j).

Larger, salient circles indicate the population median, with error bars showing the

interquartile range. The dashed lines indicate the directional thresholds used to classify

neuron type, as OFS DN1 (male: N = 9; female: N = 12) or OFS DN2 (male: N = 20; female:

N = 14). Asterisk indicates a statistically significant difference ($p < 0.05$, Watson-Williams

two-sample test). **b** Comparison of male and female OFS DN1 responses to translational

optic flow at 0.5 m/s: sideslip, lift and thrust; and rotational optic flow at 50 º/s: pitch, yaw

and roll (male: N = 9; female: N = 12). **c** Comparison of male and female OFS DN2

responses to the same optic flow stimuli (male: N = 20; female: N = 14). Data presented as

median and interquartile range.

Fig. 3. Velocity response functions in male and female OFS DNs. a Example stimulus

profile over time, with roll velocity on the y-axis. **b** Representative spike histogram from a

single trial from a male OFS DN2, smoothed using a 100 ms square-wave filter with 0.025

ms resolution, and time-aligned to the stimulus shown in panel a. Grey shading in panels a

and b highlight the analysis windows used to calculate response. **c** Example extracellular raw

data traces extracted from the analysis windows in panel b, illustrating neuronal responses to

roll velocities of –150, 10, and 200 º/s. **d** Average spike frequency (grey circles) was

calculated for each repetition from each neuron (N = 1 neuron, n = 18 repetitions), and the

median of these (black) was used for further analysis. **e** Velocity response functions of OFS

DN1 in male (blue) and female (red) hoverflies in response to roll (N = 4 males, 5 females),

sideslip (N = 5, 6), lift (N = 4, 4), and thrust (N = 3, 5). **f** Velocity response functions of OFS

DN2 to roll (N = 10 males, 8 females), sideslip (N = 6, 7), lift (N = 6, 7), and thrust (N = 6,

7). Data in panels e and f are presented as median and interquartile range. Asterisks indicate

statistically significant differences, two-way ANOVA with Šídák's multiple comparisons test

(** $p < 0.01$ and **** $p < 0.0001$), see also Table 1.

Fig. 4. Morphological reconstruction of OFS DNs. a Schematic diagram of the hoverfly

central brain and thoracic ganglia showing the projections of OFS DN1 (green) and OFS

DN2 (orange). Key anatomical landmarks are labelled: CB, central brain; OL, optic lobe; CC,

cervical connective; T1 LN, prothoracic leg nerve; T2 LN, mesothoracic leg nerve; T3 LN,

metathoracic leg nerve; TAG, thoracic-abdominal ganglion; FN, frontal nerve; PtN,

pterothoracic nerve; HN, haltere nerve; AbN, abdominal nerve. **b** Confocal image of a

reconstructed male OFS DN2. White boxes indicate regions magnified in panels d - i. **c** A

magnification of the thoracic ganglion in panel b. **d** Input dendrites of the same OFS DN2

around the sub-oesophageal ganglion. **e** Input dendrites of a female OFS DN2. **f** Input

dendrites of a male OFS DN1. **g** Output projections of the same OFS DN2 neuron as in

panels b, c and d. **f** Output projections within the thoracic ganglia of the neuron in panel e. **h**

Output projections of the neuron in panel f.

Fig. 5. WBA velocity response functions in male and female hoverflies. a A single frame

from a video labelled with trained DeepLabCut models[35,36,55]. Coloured dots correspond to

those in panel b, used to extract the wing beat amplitude (WBA). **b** Pictogram illustrating a

higher wing beat amplitude on the left wing (WBA_L) compared to the right wing (WBA_R),

suggesting a turn to the right. Equations used to calculate wing beat amplitude difference

(WBAD) and wing beat amplitude sum (WBAS). **c** Example stimulus with sideslip velocity

on the y-axis. **d** Representative WBA for the left (pale colour) and right (salient colour)

wings from a single trial, time-aligned with the stimulus shown in panel c. Grey shading in

panels c and d indicate the analysis windows used to calculate WBAD and WBAS. **e**

Magnified view of example WBA, showing behavioural responses to sideslip velocities of -2,

0.2, and 2 m/s. **f** Average WBAS (grey circles) calculated across repetitions (N = 1 animal,

n = 19-36 repetitions). Black circles represent the median WBAS per stimulus velocity, used

for further analysis. **g** WBAD in male (blue) and female (red) hoverflies in response to

different optic flow velocities: roll (N = 5 males, 7 females), sideslip (N = 6, 6), lift (N = 7,

5), and thrust (N = 6, 5). **h** WBAS to the same stimuli in the same animals. Data in panels g

and h are presented as median and interquartile range.

Fig. 6. Response onset to roll and lift stimuli in male and female hoverflies. a Time to

response onset in OFS DN1 to roll (+50 °/s) or lift (+0.5 m/s) stimuli in males (blue, N = 5)

and females (red, N = 8). **b** Response onset in OFS DN2 to roll (+50 °/s) or lift (-0.5 m/s)

measured in males (N = 20) and females (N = 14). **c** Onset of WBAS responses to roll

(-200 °/s) and lift (+2 m/s) in males (N = 5 for roll, 9 for lift) and females (N = 7 for roll, 5

for lift). Data are presented from individual neurons (a, b) or animals (c) with lines indicating

median. Asterisks indicate statistically significant differences, two-way ANOVA with

uncorrected Fisher's LSD test (** $p < 0.01$ and *** $p < 0.001$).

Fig. 7. Velocity response functions of the head, fore- and hind legs in male and female

hoverflies. a A single frame highlighting the points used to track the head (yellow and dark

green), thorax (orange), forelegs (pale and dark blue), and hind legs (magenta and pale

green). **b** The head angle (HA) was defined as the angle between the medio-posterior head

and the longitudinal thorax axis. **c** The foreleg extension was defined as the distance between

the proximal and distal point, after calculating the average for the left and the right leg. **d** The

hind leg extension was defined as the distance between the knee and a lateral point on the

mid-abdomen, after calculating the average for the left and the right leg. We also calculated

the difference between the right and the left hind leg. **e** The head angle in male (blue) and

female (red) hoverflies in response to different optic flow velocities: roll (N = 5 males, 7

females), sideslip (N = 6, 6), lift (N = 7, 5), and thrust (N = 6, 5). **f** The foreleg extension to

the same stimuli in the same animals. **g** The hind leg extension in the same animals. **h** The

difference between the left and right hind leg extension in the same animals. The data in

panels e, f, g and h are presented as median and interquartile range. Asterisks indicate

statistically significant differences, two-way ANOVA with Šídák's multiple comparisons test

(* $p < 0.05$, ** $p < 0.01$), see also Table 2.

Table legends

Table 1. Statistical summary of neural velocity response functions. Results from two-way

ANOVA analyses evaluating the effects of stimulus velocity and sex on neural responses

(OFS DN1 and DN2). Asterisks denote levels of statistical significance:

*** $p < 0.001$, **** $p < 0.0001$.

Table 2. Statistical summary of behavioural velocity response functions. Results from

two-way ANOVA analyses evaluating the effects of stimulus velocity and sex on behavioural

responses (WBAS, WBAD, head angle, foreleg extension, hind leg extension and hind leg

difference). Asterisks denote levels of statistical significance: ** $p < 0.01$,

*** $p < 0.001$, **** $p < 0.0001$, after performing a Bonferroni correction[54] for 6 comparisons

of data from the same videos.

Figure 1

Figure 2

a

90°

400
300
200

Response [spikes/s]

OFS DN1

0°

OFS DN2

270°

- ● Male
- ● Female

b

OFS DN1 [spikes/s]

-25

c

OFS DN2 [spikes/s]

-50

Left	Right	Down	Up	Away	Towards	Down	Up	Left	Right	CCW	CW
Sideslip		Lift		Thrust		Pitch		Yaw		Roll	

Figure 3

Figure 4

a
OFS DN1 OFS DN2
CB
OL
CC
FN T1 LN
PtN
TAG T2 LN
HN
AbN T3 LN

b
CB
TAG
200 µm

c
OFS DN2
Male
100 µm

d
OFS DN2
Male
100 µm

e
OFS DN2
Female
100 µm

f
OFS DN1
Male
100 µm

g
50 µm

h
50 µm

i
50 µm

Figure 5

Figure 6

Figure 7

Table 1

Stimulus		Velocity	Sex	Interaction		Velocity	Sex	Interaction
Roll	**OFS DN1**	****	0.35	0.14	**OFS DN2**	****	0.31	>0.99
Sideslip		****	0.90	0.12		****	**0.01**	****
Lift		**0.002**	0.86	>0.99		****	0.18	***
Thrust		****	0.11	****		****	0.10	**0.02**

		Velocity	Sex	Interaction		Velocity	Sex	Interaction
	OFS DN1	****	0.35	0.14	**OFS DN2**	****	0.31	>0.99
		****	0.90	0.12		****	**0.01**	****
		0.002	0.86	>0.99		****	0.18	***
		****	0.11	****		****	0.10	**0.02**

Table 2

Stimulus		Velocity	Sex	Interaction		Velocity	Sex	Interaction
Roll	WBAD	****	0.66	>0.99	WBAS	****	0.78	>0.99
Sideslip		****	>0.99	>0.99		****	>0.99	>0.99
Lift		0.18	>0.99	>0.99		****	0.54	0.12
Thrust		>0.99	>0.99	>0.99		**	>0.99	>0.99
Roll	Head	0.94	>0.99	>0.99	Foreleg	0.89	>0.99	0.26
Sideslip		**	>0.99	>0.99		**	**0.04**	**0.01**
Lift		>0.99	>0.99	>0.99		****	0.38	**0.03**
Thrust		>0.99	>0.99	0.08		**	>0.99	>0.99
Roll	Hind leg	0.21	>0.99	0.17	Δ Hind leg	**	>0.99	**0.02**
Sideslip		**0.04**	>0.99	**0.02**		****	>0.99	>0.99
Lift		0.39	>0.99	>0.99		0.41	>0.99	>0.99
Thrust		0.07	>0.99	>0.99		**	0.06	****

        Supplementary information

      Sexual dimorphism in sensorimotor transformation of optic flow

Sarah Nicholas[1], Katja Sporar Klinge[1#], Luke Turnbull[1], Annabel Moran[1], Aika Young[1], Yuri

Ogawa[1], Karin Nordström[1,2*]*

Supplementary Figure legends

Supplementary Fig 1. a The four clusters as identified by us (see Fig. 1g, h and i) displayed

on a polar plot where the distance from origo illustrates the azimuthal position and the

position along the circumference the preferred direction, colour coded as in the main paper

(pale green: OFS DN1 LHS; dark green: OFS DN1 RHS; yellow: OFS DN2 LHS; orange:

OFS DN2 RHS). **b** The Calinski-Harabasz value[1] as a function of cluster number when using

only the azimuthal location of the receptive field centre and the preferred direction (red), or

when adding one other receptive field criterion (grey, receptive field centre elevation,

receptive field height or width, maximum LMS, the number of positions with LMS values

exceeding 50% of the maximum, or LPD variance, see Methods), or all receptive field

criteria (blue). **c** The resulting z-score for azimuthal location and preferred direction, and the

four clusters using k-means clustering in Matlab.

Supplementary Fig 2. Exclusion criteria and receptive field comparisons in male and

female OFS DNs. a Distribution of maximum local motion sensitivity (LMS) in 100 male

reference neurons (grey histogram, N = 100) compared to the neurons used for quantification

in the rest of the paper (N = 35 males, 36 females). Six neurons with a maximum LMS below

20 spikes/s (red cross-hatched region or dashed line) were excluded from further analysis. **b**

Number of stimulus positions with LMS greater than 50% of the neuron's maximum LMS

(black and red arrows in Fig. 1c and f). Neurons with fewer than 4 positions were excluded.

c Distribution of LPD variance. Neurons with directional variance exceeding 30° were

excluded. **d** Classification of male neurons based on receptive field centre and preferred

direction. The top panel shows the preferred directions across neurons, with dashed lines

indicating the directional thresholds used for neuron type classification. The middle panel

shows receptive field centre locations, and the bottom panel illustrates the relationship

between receptive field centre and the receptive field preferred direction for all 35 male

neurons, with 2 neurons excluded from further analysis (grey). Note that the scale is the same

for the azimuth and elevation, and the scale bar corresponds to both the x- and y-axis. **e** Same

classification criteria applied to 36 female neurons, with 7 neurons excluded (grey). **f** The

receptive field width and height at the 50% contour line for the remaining OFS DN 1 (N = 10

males, 12 females) and OFS DN2 (N = 23 males, 17 females). Individual data points from

independent neurons, with black lines representing the median.

Supplementary Fig 3. Schematic illustration showing the steps used to standardize data

across recordings. a. Top panel, ventral-side-up position of hoverfly during

electrophysiology recordings. **b** Response of an example OFS DN2 neuron to a full-screen,

full-contrast sinusoidal grating (spatial wavelength 7°; temporal frequency 5 Hz) moving in

eight different directions. The red arrowhead shows the neuron's preferred direction, and the

red line indicates response amplitude. **c** Preferred direction of male hoverfly OFS DNs

(N = 29). Neurons are colour coded by classification. **d** Preferred direction of female hoverfly

OFS DNs (N = 26). Neurons are colour coded by classification. **e** We rotated the data 180° to

display the dorsal visual field up. **f** Direction sensitivity of the same neuron as in panel b after

rotating the data 180°. **g** Preferred direction of male hoverfly OFS DNs after rotating the data

180°. Neurons are colour coded by classification, with dashed lines indicating the receptive

field thresholds. **h** Preferred direction of female hoverfly OFS DNs after rotating the data

180°. **i** To display all neural data as RHS neurons, we flipped LHS neurons along the midline

(yellow to black). **i** Resulting direction sensitivity after flipping along the midline of the same

neuron as in panels b and f. **k** Preferred direction of male hoverfly OFS DNs when displayed

as RHS only. **l** Preferred direction of female hoverfly OFS DNs when displayed as RHS only.

Supplementary Fig. 4. Comparison of spontaneous activity and responses to stationary

stimuli. a Spontaneous activity (open shapes) and responses of a male OFS DN1 to the

stationary starfield (filled shapes), from data extracting directional sensitivity (as in Fig. 2;

circles) or velocity response functions (as in Fig. 3; triangles). **b** Spontaneous activity and

responses of OFS DN2, displayed as in panel a. **c** WBAS before stimulus presentation (open

squares) or when viewing the stationary starfield (filled squares), quantified during velocity

response function experiments (as in Fig. 5). **d** Hind leg extension before stimulus

presentation (open diamonds) or when viewing the stationary starfield (filled diamonds),

quantified during velocity response function experiments (shown in Fig. 7). The data points

show data from individual neurons (a and b) or animals (c and d), with lines representing the

median. Asterisks denote statistical significance, two-way ANOVA with uncorrected Fisher's

LSD test: $* p < 0.05$, $** p < 0.01$ and $*** p < 0.001$.

Supplementary Fig 5. Morphological quantification of OFS DN2. a Schematic diagram of

the hoverfly central brain and thoracic ganglia showing locations where the cervical

connective (CC) and the OFS DN2 (orange) widths were measured. **b** Width measurements

of OFS DN2 axons along the cervical connective at four anatomical landmarks: (1)

immediately posterior to the central brain, (2) upper third of the CC, (3) lower third of the

CC, and (4) immediately anterior to the thoracic ganglion. Data are shown for 3 male (blue)

and 3 female (red) hoverflies. **c** Width of the anterior end of the cervical connective in the

same individuals. All data are presented as individual animals with the horizontal lines

indicating median.

Supplementary Fig 6. Impact of stimulus size on neural and behavioural velocity

response functions. a Electrophysiological recordings were performed with the hoverfly

orientated ventral side up, facing the centre of a horizontally oriented visual display. **b** In

behavioural experiments, the hoverfly was tethered at 32º angle, parallel to the ground, facing

the centre of a vertically aligned visual display. **c** Velocity response functions of male OFS

DN2 neurons when the stimulus covered either the full screen (blue) or the central square

(black, N = 7). **d** Velocity response functions of WBAS in male hoverflies when the stimulus

covered either the full screen (blue) or the central square (black, N = 5). All data are

presented as median and interquartile range.

Supplementary Fig 7. Response onset to sideslip. Onset of head angle, WBAS, fore- and

hind legs to sideslip (+2 m/s) in males (N = 6) and females (N = 6). There was no significant

difference between body parts, nor males and females (two-way ANOVA, $p = 0.08$, $p = 0.67$

and $p = 0.84$, for body parts, sex and interaction respectively). All data are presented as

individual animals with the horizontal lines indicating median.

Supplementary Movies

Supplementary Movie 1. Neural response to roll optic flow. The video shows the raw

extracellular response of an OFS DN2 in red when the hoverfly was viewing an example roll

stimulus, as in Fig. 3 of the main paper.

Supplementary Movie 2. Female behavioural response to sideslip optic flow. The video

on the left shows a female hoverfly filmed from above, with coloured dots indicating the

body parts tracked by DeepLabCut to extract wing beat amplitude (WBA), head angle, and

fore- and hind leg kinematics. The stimulus viewed by the hoverfly is shown on the right,

with stimulus velocity indicated above the hoverfly video. The movie relates to Fig. 5 of the

main paper.

Supplementary Movie 3. Male behavioural response to sideslip optic flow. The video

shows a male hoverfly filmed from above, viewing sideslip at different velocities.

Supplementary Movie 4. Female behavioural response to lift. A female hoverfly filmed

from above, viewing lift optic flow at different velocities.

Supplementary Movie 5. Male behavioural response to lift. A male hoverfly filmed from

above, viewing lift optic flow at different velocities.

Supplementary Movie 6. Female behavioural response to thrust. A female hoverfly

filmed from above, viewing thrust optic flow at different velocities.

Supplementary Movie 7. Male behavioural response to thrust. A male hoverfly filmed

from above, viewing thrust optic flow at different velocities.

Supplementary Movie 8. Female behavioural response to roll. A female hoverfly filmed

from above, viewing roll optic flow at different velocities.

Supplementary Movie 9. Male behavioural response to roll. A male hoverfly filmed from

above, viewing roll optic flow at different velocities

111   Reference cited in Supplementary Information

112   1   Caliński, T. & Harabasz, J. A dendrite method for cluster analysis. *Communications*
113   *in Statistics* **3**, 1-27, doi:10.1080/03610927408827101 (1974).
114

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7