Single Domain Antibody Inhibitors Target the Coiled Coil Arms of the _Bacillus subtilis_ SMC complex

Ophélie Gosselin[1], Michael Taschner[1], Lea M. Huber-Hürlimann[2], Markus A. Seeger[2] and Stephan Gruber[1]*

[1]Department of Fundamental Microbiology (DMF), Faculty of Biology and Medicine (FBM), University of Lausanne (UNIL), 1015 Lausanne, Switzerland

[2]Institute of Medical Microbiology, University of Zurich, Zürich, Switzerland

*Corresponding Author: stephan.gruber@unil.ch

Key words: Single domain antibodies, Sybodies, nanobodies, SMC complex, bacterial condensin, DNA loop extrusion, chromosome segregation

Abstract

Synthetic nanobodies—also called sybodies—have proven valuable for stabilizing conformations of

purified proteins, advancing structural and functional studies for example of transmembrane protein

complexes. However, their utility in modulating protein function in living cells has remained less well

explored. Structural Maintenance of Chromosomes (SMC) complexes facilitate chromosome

organization by DNA loop extrusion, a fundamental process in all domains of life. In this study, we target

the bacterial SMC complex, Smc-ScpAB, in *Bacillus subtilis* with synthetic nanobodies, aiming to identify

key functional regions of the protein complex in a largely unbiased manner. We first isolate sybodies

that specifically bind purified Smc-ScpAB and then express them in *B. subtilis* to select binders capable

of disrupting Smc-ScpAB function, leading to chromosome segregation defects and cell death. Mapping

and biochemical characterization show that the fourteen disruptive sybodies belong to one of three

library designs, target the Smc subunit near the same coiled coil arm interface and modulate its ATPase

activity in two principal ways, highlighting the mid-region of the Smc coiled coil as critical feature of

the DNA folding process. These findings underscore the potential of sybodies—and, by extension,

designed binders—as versatile tools for probing dynamic protein function in living cells, with potentially

broad applications in cell and synthetic biology.

Introduction

Rapid and specific interference with protein activity and dynamics in living cells is essential for studying and understanding biological mechanisms. However, current approaches—primarily based on small-molecule inhibitors—remain laborious, time-consuming, and costly. The advent of conformation-specific synthetic nanobody (sybody) selection [1], and more recently, computational design of protein binders [2], offer unique opportunities for protein-based inhibition and modulation of cellular targets. Here, we generate sybodies against the SMC complex in *B. subtilis* to efficiently inhibit its function within cells.

Structural Maintenance of Chromosomes (SMC) complexes are multi-subunit, ring-shaped ATP-hydrolyzing DNA motors that structure chromosomal DNA by loop extrusion. They are essential for chromosome organization and segregation, gene expression, DNA repair, and defense against non-self DNA across all domains of life [3]. Each SMC protein (Smc in *B. subtilis, bsu*Smc) is a long polypeptide with its N- and C-termini folding together to form a globular ATP binding cassette (ABC) "head" domain. The head is connected to a "hinge" dimerization domain via a ~50 nm antiparallel coiled coil "arm", creating an elongated dimer (Figure 1Aii and 1Aiii). ATP binding promotes head engagement, bringing the heads together to form a functional ATPase. Non-SMC subunits, a kleisin (ScpA in *B. subtilis*) and a dimer of KITE (ScpB in *B. subtilis*) or two HAWK proteins, bridge the heads, together forming a ring around DNA (Figure 1Aiii). In *B. subtilis*, Smc-ScpAB is recruited to the origin of replication (*oriC*) region on the chromosome by ParB, a DNA-binding protein that recognizes centromere-like *parS* sequences (Figure 2Ai). Once loaded, Smc-ScpAB translocates at a rate of ~1 kb/s onto flanking DNA, aligning the two chromosome arms and individualizing nascent sister chromosomes[4-6], likely bypassing obstacles on the chromosome through the SMC hinge gate [7]. Null mutants of *smc*, *scpA*, or *scpB* fail to segregate chromosomes properly and lose viability under conditions promoting rapid growth[5,6,8].

Several models have been proposed for the mechanism of DNA loop extrusion by SMC complexes, including the "segment capture" model, in which DNA segments are transiently trapped between the

SMC arms in partially open states and fused into larger loops through iterative ATP-driven cycles (Figure

1Aiii and S1B) [9-11]. However, detailed structural understanding and experimental testing *in vivo* are

lacking.

Antibodies can block biochemical reactions by stabilizing reaction intermediates. Synthetic single-

domain antibodies called sybodies are small, robust antigen-binding proteins that were engineered

based on three camelid nanobody structures [1]. They are selected from synthetic libraries that encode

a wide diversity of binding surfaces and epitope shapes based on the length and geometry of their

CDR3 loop, allowing strong binding of diverse antigen surfaces (Figure 1Ai). Their small size (~15 kDa),

stability, and ability to bind transient epitopes make them ideal tools for targeting specific states of

biomolecules [12]. Selection is carried out using purified and immobilized Smc-ScpAB by ribosome display

and phage display, typically followed by an ELISA-based screening [1].

Here, we demonstrate that sybodies can be used to interfere with the function of *B. subtilis* Smc-ScpAB

in vivo. We first isolate binders that specifically target purified Smc-ScpAB *in vitro* and then select those

that eliminate Smc function when expressed in *B. subtilis*. Fourteen sybodies were found to disrupt

chromosome segregation, mimicking *smc* deletion phenotypes. Biochemical assays confirmed that

selected sybodies alter the ATP hydrolysis rate of Smc-ScpAB and its stimulation by DNA addition, likely

stabilizing intermediates of the conformational cycle. Mapping experiments using chimeric Smc

constructs revealed that most binders, unexpectedly, target the coiled coil, specifically in a defined

region near the Smc 4N arm-to-arm contact [13], highlighting the potential of sybodies and designed

binders, as genetic tools to identify essential functional domains and underscoring the importance of

coiled coil dynamics for Smc function.

Results

Generation of *bsuS*mc-ScpAB-specific sybodies

To isolate sybodies that impede the function of *B. subtilis* Smc-ScpAB complex, we first performed *in vitro* selection with purified Smc-ScpAB in the presence of 40 bp duplex DNA, and an ATPase-deficient mutant (E1118Q) of *bsuS*mc. These conditions favour ATP-engaged complexes alongside the typically predominant ATP-disengaged rod-shaped state [13]. *bsuS*mc was also biotinylated at the hinge at residue R643C for immobilization (Figure 1Aii-iii and S1A) [1,14-16]. Starting from three synthetic sybody libraries encoding distinct epitope-binding geometries (denoted as concave, loop, convex, respectively, Figure 1Ai), we performed one round of ribosome display starting from a large sybody library followed by two rounds of phage display, thereby enriching sybodies that bind to Smc-ScpAB (Table 1). From each library, 95 sybody-expressing *E. coli* clones were randomly chosen for *in vivo* characterization. ELISA data revealed that nearly all clones bind purified Smc-ScpAB (Table 1). However, the ELISA signals of only few Sybodies showed clear dependence on the presence or absence of ATP and DNA (Table S1).

Identification of *bsuS*mc-ScpAB-blocking sybodies through phenotypic screening in *B. subtilis*

We therefore decided to screen for inhibiting sybodies directly in *B. subtilis* cells. Despite containing a conserved disulfide bond, sybodies have been successfully expressed inside cells and were shown to bind their targets in the reducing environment of the cytoplasm, likely owing to their robust folding [17]. To assess whether the selected sybodies interfere with Smc function *in vivo*, each of the 285 sybody sequences was cloned under a xylose-inducible promoter (P_{xyl}) into an *E. coli–B. subtilis* shuttle vector and integrated at the *amyE* locus of the wild-type 1A700 strain (Figure 1B) [18]. Fourteen sybodies from the loop library failed to yield transformants in the presence (but not absence) of xylose, indicating cell lethality due to sybody inhibition of Smc-ScpAB (Figure 1C, S2A-B). These Smc-disruptive sybodies (denoted as Sb002, 003, 006, 010, 015, 020, 021, 031, 156, 160, 162, 169, 188, and 194, respectively) harbor fourteen distinct sequences(Table S2). Intriguingly, no such disruptive sybodies were isolated

from the concave and convex libraries (Figure S2B), suggesting that the CDR3 geometry as present in

the loop library is particularly effective at inhibiting *bsuS*mc-ScpAB. Likely, sybodies of the other

libraries bound to other epitopes of *bsuS*mc-ScpAB that are not sensitive to conformational trapping,

and it is further possible that they were not stably expressed in *B. subtilis*. However, at least for sybodies

from the loop library, we did not notice any obvious correlation between expression levels and

phenotypes. Selected sybodies showed similar expression and were appreciably expressed even

without inducer (Figure S4, S5). Notably, tendencies of preferential isolation of binders from one of the

three libraries have previously been observed, although at milder levels [1,17,19].

Selected sybodies target *bsuS*mc-ScpAB function *in vivo*

To determine whether the fourteen sybodies indeed impair chromosome segregation, we used

fluorescence microscopy to monitor *oriC* positioning in a *B. subtilis* strain expressing a ParB-GFP fusion

protein together with a P_{xyl}-inducible sybody gene. In these strains, ParB-GFP marks the replication

origin region by binding to *parS* sites near *oriC* (Figure 2Ai). As expected, the *Δsmc* strain displayed

fewer ParB-GFP foci and elongated cells (Figure 2Ai) (Table S3). These elongated cells are known to

harbour expanded nucleoids, consistent with delayed *oriC* separation rather than delayed DNA

replication [6,20]. A time course experiment using sybody Sb006 revealed chromosome organization

defects as early as 30 minutes post induction, with a significant reduction in ParB-GFP foci per μm cell

length at 40 minutes (p = 0.0145) (Figure 2Aii). Subsequent imaging was performed ~35 minutes after

induction of sybody expression.

All disruptive sybodies reduced ParB-GFP foci density compared to the control (1.03 foci/μm), with

values ranging from 0.56 foci/μm (Sb020) to 0.22 foci/μm (Sb003). These defects were all robustly

detected, while being less severe than the *Δsmc* mutant (~0.05 foci/μm), conceivably due to the short

induction time or incomplete inhibition. Milder defects were also observed without induction, likely

attributed to leaky expression from the P_{xyl} promoter (Figure S3A). Sybody expression also caused cell

elongation and growth delays, both hallmarks of impaired Smc activity, consistent with chromosome

segregation defects that delay cell division (Figure S3B–C). Altogether, these results show that all

selected sybodies induce *Δsmc*-like phenotypes likely by interfering with *bsu*Smc-ScpAB activity *in vivo*.

Selected sybodies specifically target *B. subtilis* Smc-ScpAB

To test whether sybodies specifically target *B. subtilis* Smc-ScpAB, we assessed their effects in a strain

expressing *Streptococcus pneumoniae* (*Spn*) Smc-ScpAB and ParB proteins in place of the endogenous

B. subtilis proteins. The *Spn* sequences can functionally replace the corresponding *Bsu* sequences

despite significant sequence divergence (~38% sequence identity) [21]. Strikingly, none of the sybodies

impaired growth in this background, even though a cognate ParAB pair is absent in this strain (*S.*

pneumoniae lacks *parA*), sensitizing cells to chromosome segregation defects (Figure 2C). These results

confirm that the sybodies are specific to *bsu*Smc-ScpAB and that off-target toxicity is not noticeable.

***bsu*Smc-ScpAB-disrupting sybodies target two distinct coiled coil regions adjacent to the *bsu*Smc 4N**

arm-to-arm contact

To map the sybody-binding site on the *bsu*Smc-ScpAB complex, we utilized five chimeric Smc constructs

(Smc Chimera 1-5), in which the hinge and progressively longer segments of the adjacent coiled coils

were replaced with the corresponding sequences from the *S. pneumoniae* Smc protein (Figure 3A).

Chimeric junctions were designed to preserve coiled coil integrity based on available crystal structures,

coiled coil predictions by DeepCoil, and AlphaFold2 structural models [1,22,23]. All chimeric Smc strains

retained the native *B. subtilis scpA*, *scpB*, and *parB* genes and were viable under conditions promoting

fast growth, demonstrating proper protein folding and functioning of chimeric Smc-ScpAB complexes

(Figure S6A).

Chimeras 1 and 2, which carry the central hinge and coiled coil sequences of *Spn* origin (residues 363-

812 and 339-836, respectively), showed the same sybody-induced growth defects as the wild-type

strain, indicating that the sybodies bind further away from the hinge, within the remaining *B. subtilis*

portion comprising of the Smc head and the first ~230 coiled coil residues, and ScpAB (Figure 3A). By

contrast, Chimera 5 (residues 248–927 of *Spn* origin), supported robust growth even in the presence of

the *bsu*Smc-disrupting sybodies, indicating that the region comprising the heads and first ~50 residues

of the coiled coil (including the joint) does not comprise the binding site. The binding region for the

sybodies is thus located on the coiled coil between residues 248 and 339 (or 836 and 927),

demonstrating that all fourteen sybodies bind to a central region of the coiled coil.

Chimera 3 (*spn*Smc residues 318-857) and chimera 4 (*spn*Smc residues 276-899) revealed two distinct

sybody responses. Sb003, 031, 156, 160, 162, and 169 impaired growth of both strains, mapping their

binding sites to *B. subtilis* residues 248-276 (and 899-927), present in chimeras 1–4 but absent in

chimera 5. In contrast, Sb002, 006, 010, 015, 020, 021, 188, and 194 had no impact on the viability of

chimera 3 or 4, suggesting they target residues 318-339/836-857 (Figure 3A, 3D).

In sum, these experiments mapped the binding sites of all fourteen disruptive sybodies to one of two

~twenty amino acid segments in the central region of the *bsuS*mc coiled coil (Figure 3D). These regions

harbor relatively poorly conserved sequences but flank the 4N arm-arm contact (~at residue 295),

previously proposed to serve as an essential structural switch between closed and open

conformations[13]. These results reveal that all isolated sybodies target the same functional element of

the Smc arms, and that interfering with this element disrupts Smc-dependent chromosome

organization *in vivo*.

Smc-disrupting sybodies affect the ATPase activity in one of two ways

To investigate how sybodies may influence *bsuS*mc-ScpAB, we next measured ATP hydrolysis rates using

purified proteins in the presence and absence of DNA. SMC ATPase activity serves as an indirect

readout for state transitions, including arm opening and head engagement. In absence of DNA, *bsuS*mc-

ScpAB hydrolyzes ATP at ~4 ATP/min/Smc [13]. DNA binding stimulates ATP hydrolysis to ~9 ATP/min/Smc,

presumably by DNA binding promoting arm opening and head engagement, creating a more open

conformation.

All disruptive sybodies measurably affected the ATP hydrolysis rate of *bsuS*mc-ScpAB. Ten maintained

a near-basal ATPase rate even in presence of DNA (Sb002, Sb003, Sb010, Sb031, Sb156, Sb160, Sb162,

Sb169, Sb188, and Sb194) (Figure 3B-C and S6B). This suggests that they stabilize the complex in a

closed or partially closed conformation, thus hindering DNA-dependent ATPase stimulation. Notably,

in this group, Sb002 and Sb194 showed slightly elevated ATPase rates with DNA, hinting at a mildly

more flexible conformation.

In contrast, the other sybodies (Sb006, Sb015, Sb020, and Sb021) stimulated ATP hydrolysis in the

absence of DNA, with Sb006 reaching levels comparable to DNA-induced activation (~10 ATP/min). This

implies that these sybodies stabilize Smc-ScpAB in a more open state promoting head engagement.

Curiously, addition of DNA in the presence of Sb006 and Sb021 reduced—rather than stimulated—

ATPase activity (Figure 3B-C), suggesting that concurrent DNA and sybody binding may trap the complex

in a non-productive conformation, perhaps by favoring the open state excessively. These findings are

consistent with the idea that 4N contact region plays a critical role in arm opening and ATPase

stimulation.

Interestingly, the observed ATPase profiles correlate with distinct binding regions along the Smc arms.

All ATPase-stimulating sybodies (Sb006, Sb015, Sb020, Sb021) bind above the 4N contact

("Opening^{UP}"), while sybodies that hinder DNA stimulation fall into two subgroups: those binding above

the 4N contact (Sb002, Sb010, Sb188, Sb194—"Closing^{UP}") and those binding below it (Sb003, Sb031,

Sb156, Sb160, Sb162, Sb169—"Closing^{DOWN}") (Figure 3D). These results indicate that the sybodies

perturb *bsuS*mc-ScpAB in two principal ways: either by favoring the opening of the arms or by

constraining the arms in a more closed state. The results highlight arm opening and closure as central

regulatory features of bacterial Smc-ScpAB [9,11,24,25] and confirm the key role of the 4N arm contact in

balancing the opening and closing of SMC arms [13].

Discussion

Over the past three decades, nanobodies and their synthetic counterparts have transformed biomedical research and structural biology by enabling the targeting and stabilization of transient protein conformations. Despite these advances, their application to probing protein function in living cells has remained limited [17,26]. Here, we expand the utility of sybody selection by employing it as a tool to screen for genetic probes targeting the function of selected molecular machines *in vivo*. This strategy combines the advantages of controlled biochemical reconstitution with the ability to study the function of proteins in their native context. Importantly, sybody selection requires no prior knowledge of vulnerable regions, providing an unbiased means to interrogate protein function that is difficult to achieve through rational design. Our study based on the *B. subtilis* Smc-ScpAB DNA motor as a model uncovers a central regulatory region in the coiled coil arms of Smc and highlight how distinct arm conformations shape SMC function in loop extrusion and chromosome segregation.

Revealing Smc arm dynamics through synthetic binders

All fourteen Smc-disrupting sybodies share the loop library design. Moreover, their Smc binding sites map to the two short coiled coil segments flanking the conserved 4N arm-to-arm contact, a region that has previously been implicated in key conformational transitions during arm opening [13]. While the fourteen sybodies harbour different epitope binding sequences, we suspect that they may all bind to the Smc dimer in an analogous manner. With the long CDR3 loops known to bind narrow structural cavities[12,27], we speculate that these sybodies may intercalate between the two coiled coil arms (Figure 3D). Structural studies are needed to test this hypothesis directly, however, our attempts at X-ray crystallography and electron microscopy of Smc-sybody complexes have failed so far.

Notably, while the coiled coil dimer appears similarly structured from the elbow to the hinge [9] (Figure 3D), no Smc-disrupting sybodies mapped to regions further away from the 4N contact (Figure 3D). This indicates that opening and closure of the Smc arms might be dispensable at these places, although

formal testing would require isolating binders efficiently targeting these regions, potentially facilitated

by binder design, and expressing them in *B. subtilis*. This parallels with recent findings with the bacterial

SMC defense system Wadjet, where hinge-proximal arm opening is dispensable for loop extrusion, but

essential for obstacle bypass [7]. Alternatively, these arm-to-arm contacts might be more stable and thus

less likely to be interfered with by sybody binding. Similarly, sybodies targeting below the 4N contact

all reduce the ATPase rate (or rather its DNA stimulation), indicating that artificial opening at this

position may not disrupt Smc function (Figure S7). Again, these hypotheses need to be tested more

directly. We envision that the design of protein binders may allow us to confirm these observations and

further dissect the underlying mechanisms.

This work establishes sybodies as precision tools for blocking ATP-driven machines inside bacterial cells.

By targeting allosteric control points in Smc, they enable mechanistic dissection of loop extrusion and

open new avenues for studying dynamic complexes *in vivo*. More broadly, the study demonstrates how

synthetic binders can trap or block conformations of active chromatin-associated machines, providing

a powerful means to probe their mechanisms in living cells. Looking ahead, the rational design of

protein binders with tailored geometry and allosteric potential could allow researchers to manipulate

and visualize specific conformational states with even greater control and refinement, across both

prokaryotic and eukaryotic organisms, while the generation and *in vivo* screening of sybodies remains

an attractive approach to gain unexpected biological insights in an unbiased manner.

Material and Methods

***Bacillus subtilis* strains construction and growth**

B. subtilis strains used in this study were derived from the parental strain 1A700. Allelic replacement

was achieved via double-crossover recombination at the endogenous *smc* locus using the natural

competence method described by [18,28]. Chromosome-integration of sybody genes were performed by

transforming a vector containing an *amyE h*omologous regions on each side of the recombinant gene

of interest, as described in [18]. Transformants were selected on ONA solid medium containing the

appropriate antibiotics. Following transformation, strains were purified by single colony isolation and

verified through a combination of marker testing, phenotype assessment, PCR, and Sanger sequencing

of the *smc* locus, when appropriate. For spot assays, cells were cultured to stationary phase in liquid

LB medium, and 9^{-2} and 9^{-5} dilutions were then spotted onto solid ONA medium with chloramphenicol

(5 ug/mL) and xylose (0.5% w/v) when induction of the P_{xyl} promoter was required. The strains,

plasmids, and oligonucleotides used in this study are listed in Table S4, Table S5 and Table S6,

respectively.

For viability assessment, 200 LB medium was inoculated with a 1:1000 dilution of a dense *B. subtilis*

culture and grown to stationary phase. The culture was then serially diluted in LB medium. From each

dilution (9^{-1} to 9^{-8}), 5 µL was spotted onto LB agar plates containing the appropriate antibiotic selection.

Plates were incubated, and viability was assessed by imaging the spots at 16- and 24-hours post-

incubation.

Sybody selection

Sybody selection was performed following the protocol by [16], with modifications to include binders

specific for the ATP/DNA-stabilized "open" form of the *bsuS*mc-ScpAB complex.

Ribosome Display

The standard WTB buffer (50 mM Tris/acetate pH7.4, 150 mM NaCl, 50 mM MgAc2) and its derivatives

were adjusted to 50 mM NaCl to facilitate *bsuS*mc-ScpAB binding to dsDNA. The target complex was

assembled by mixing 500 nM *bsuS*mc(C119S, C437S, C826S, E1118Q, R643C)-ScpAB, 2 mM ATP, and 5

μM 40 bp dsDNA in WTB-D-BSA buffer (WTB including 0.5% (w/v) BSA and 0.1% Tween-20), incubated

for 15 min at room temperature (RT). The ribosome display panning solution was supplemented with

2 mM ATP and 5 μM dsDNA to maintain binding conditions. Solution panning and biotinylated target

capture were conducted at RT to optimize selection for the "open" complex. Washing steps

incorporated 2 mM ATP and 5 μm 40 bp dsDNA, while elution, reverse transcription, and cDNA

amplification followed the original protocol. Reverse transcribed RNA molecules were quantified by

qPCR.

First round of phage display

Phagemid libraries were cloned and electroporated into *E. coli* SS320 as previously described[16]. Phage

production was performed using M13KO7 helper phage. The phage display buffer (TBSM) consisted of

20 mM Tris-HCl (pH 7.4), 50 mM NaCl, and 2 mM MgCl2. Target preparation involved incubating 500

nM biotinylated *bsuS*mc(C119S, C437S, C826S, E1118Q, R643C)ScpAB with 2 mM ATP and 5 μM dsDNA

in TBSM-BSA-D (TBSM including 0.5% (w/v) BSA and 0.05% Tween-20). Phages (10^{12}/ml) were incubated

with the target at 50 nM for 20 min at RT before immobilization on neutravidin-coated plates. Washing

(with added 2 mM ATP and 5 μm 40 bp dsDNA) and elution otherwise followed the original protocol,

with enrichment assessed by qPCR. Amplified phages were used for a second selection round.

Second round of phage display

Purified phages from the first round were used at 5×10^{13} phages/ml, and the target was prepared as

in the first round. Phage-target binding occurred at 50 nM in the presence of ATP and dsDNA, followed

by capture on magnetic beads. A competition step using non-biotinylated *bsuS*mc-ScpAB (1 μM) was

included before washes with TBSM-D. Phages were eluted and enrichment was determined by qPCR.

Phagemids were purified, sub-cloned into the pNb_init vector, and transformed into *E. coli* MC1061.

Growth curve analyses

Growth curves were generated from *B. subtilis* strains grown overnight to exponential phase in LB

medium supplemented with 0.5% glucose, incubated at 30°C with shaking. The following day, fresh 5

mL LB cultures were inoculated with the overnight culture to an OD of 0.005 and incubated at 37°C

with shaking until reaching an optical density (OD) of 0.05. Subsequently, each culture was subjected

to a twofold dilution in a 96-well plate (Costar #3596), achieving a final volume of 200 μL per well.

Where necessary, cultures were induced with 0.5% xylose. Plates were incubated at 37°C with

continuous shaking in a Thermo Scientific Multiskan FC plate reader. Growth curves were determined

by light scattering at 620 nm. The BactEXTRACT app was used to perform Analysis and visualization of

the data [29].

Protein purification

Purification of *bsuS*mc and biotinylated *bsuS*mc(C119S, C437S, C826S, E1118Q, R643C)

*bsuS*mc proteins were purified according to [14]. pET-22 or pET-28 plasmids encoding the *Smc*

recombinant sequences were transformed into *E. coli* BL21-Gold (DE3) cells. Protein expression was

carried out in ZYM-5052 autoinduction medium for 23 hours at 24°C. The cells were harvested and

resuspended in lysis buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM DTT, 10% (w/v)

sucrose) supplemented with protease inhibitor cocktail. Cell lysis was achieved by sonication, and the

lysate was clarified by centrifugation. The supernatant was filtered through a 0.45 μm membrane and

loaded onto two HiTrap Blue HP 5 mL columns connected in series. Elution was performed using lysis

buffer with 1 M NaCl. The main peak fractions were pooled and diluted with salt-free buffer (50 mM

Tris-HCl, pH 7.5; 1 mM EDTA; 1 mM DTT) to a final conductivity equivalent to 50 mM NaCl (~8 mS/cm).

This diluted sample, supplemented with protease inhibitor cocktail, was applied to a HiTrap Heparin

HP 5 mL column, and the proteins were eluted using a linear gradient up to 2 M NaCl. The main peak

fractions (~ 5 mL) were collected and subjected to further purification by gel filtration using an XK 16/70

Superose 6 PG column equilibrated with 50 mM Tris-HCl, pH 7.5, 200 mM NaCl, 1 mM EDTA, and 1 mM

TCEP. The peak fractions were collected, concentrated using a Vivaspin 15 10K MWCO filter, flash frozen

in liquid nitrogen, and stored at -80°C. Protein concentration was determined by absorbance, utilizing

theoretical molar absorption and molecular weight values.

For the biotinylated *bsu*Smc(C119S, C437S, C826S, E1118Q, R643C) protein used in the sybody

selection, the protein was first purified as described above, except that the reducing agent (TCEP) was

omitted from the final gel filtration buffer. Labelling was performed by incubating 600 µL of

*bsuS*mc(C119S, C437S, C826S, E1118Q, R643C) (83 µM in 50 mM Tris-HCl pH 7.5, 200 mM NaCl) with 1

mM PEG2-biotin maleimide for 10 minutes at 4 °C. The reaction was quenched by addition of 0.5 mM

2-mercaptoethanol. Excess label was removed using Zeba spin desalting columns (Thermo Fisher) in

multiple parallel runs due to volume limitations. The final yield was 600 µL at 72.35 µM (subunit

concentration). Labelled *bsuS*mc(C119S, C437S, C826S, E1118Q, R643C) was loaded onto a Superose 6

10/300 Increase column pre-equilibrated in 50 mM Tris-HCl pH 7.5, 200 mM NaCl. Peak fractions were

pooled and mixed with ScpA and ScpB at a 1:1:2 molar ratio relative to the *bsuS*mc dimer. The final

concentration of the full complex was adjusted to 5.7 µM. Complexes were aliquoted in 100 µL

portions, flash-frozen in liquid nitrogen, and stored at −80 °C.

Purification of *bsu*ScpA

ScpA was purified using the method described by[25]. *E. coli* BL21-Gold (DE3) cells, transformed with a

pET-28 derived plasmid encoding the ScpA protein, were used for expression. Cultivation was

performed in ZYM-5052 autoinduction medium at 16°C for 28 hours. Cells were then harvested and

resuspended in lysis buffer (50 mM Tris-HCl, pH 7.5; 200 mM NaCl; 5% glycerol) supplemented with

protease inhibitor cocktail. The cells were lysed by sonication, and the lysate was clarified by

centrifugation. The supernatant was applied to a 5 mL HiTrap Q ion exchange column and eluted with

a gradient up to 2 M NaCl. Peak fractions were pooled and adjusted to a final concentration of 3 M

NaCl by mixing with 4 M NaCl buffer. This mixture was loaded onto a HiTrap Butyl HP column and eluted

with a reverse gradient to 50 mM NaCl. Eluted peak fractions were concentrated to 5 mL using Vivaspin

15 10K MWCO filters and further purified by size exclusion chromatography (SEC) on a HiLoad 16/600

Superdex 75 pg column equilibrated with 20 mM Tris-HCl, pH 7.5, and 200 mM NaCl. The purified

protein was concentrated, flash frozen, and stored at -80°C.

Purification of *bsu*ScpB

ScpB was purified following the protocol outlined by [25]. The coding sequence of ScpB, cloned into a

pET-22 derived plasmid, was transformed into chemically competent BL21-Gold (DE3) *E. coli* cells.

These cells were cultivated in ZYM-5052 autoinduction medium at 24°C for 23 hours. Subsequently, the

cells were harvested and resuspended in lysis buffer (50 mM Tris-HCl, pH 7.5; 150 mM NaCl; 1 mM

EDTA; 1 mM DTT) supplemented with protease inhibitor cocktail. Cell lysis was performed by

sonication, followed by centrifugation to remove cell debris. The resulting supernatant was diluted to

a final NaCl concentration of 50 mM and loaded onto a 5 mL HiTrap Q HP column. Elution was achieved

using a gradient up to 2 M NaCl. The eluate was then diluted with lysis buffer containing 4 M NaCl to

achieve a final concentration of 3 M NaCl. This sample was applied to two 5 mL HiTrap Butyl columns

connected in series, and the protein was eluted with a reverse gradient to 50 mM NaCl. The peak

fractions from this column were concentrated and subjected to size exclusion chromatography (SEC)

using a HiLoad 16/600 Superdex 200 pg column equilibrated with 50 mM Tris-HCl, pH 7.5, 100 mM

NaCl, and 1 mM DTT. Fractions containing ScpB were concentrated, flash frozen, and stored at -80°C.

Medium scale sybody purification

Individual sybody plasmids were transformed into chemically competent *E. coli* MC1061 cells via heat

shock at 42 °C for 45 s, followed by recovery in LB medium at 37 °C. Transformed cells were cultured

overnight in TB medium supplemented with 25 µg/ml chloramphenicol. Precultures were used to

inoculate 50 mL TB cultures, which were grown at 37 °C before shifting to 22 °C. Expression was induced

with 0.02% (wt/vol) L-(+)-arabinose and continued overnight at 22 °C with shaking. Cells were

harvested by centrifugation and resuspended in periplasmic extraction buffer. Following incubation at

4 °C, cells were pelleted, and the supernatant was supplemented with imidazole to 15 mM. The extract

was incubated with His MultiTrap HP resin, followed by centrifugation and washing with TBS containing

30 mM imidazole. Elution was performed using TBS with 300 mM imidazole. Purified sybodies were

analyzed on a Sepax SRT-10C SEC100 column at 1 ml/min flow rate. Monomeric sybodies eluted

between 11–12.5 ml, while retention volumes <11 ml indicated oligomerization, and >14 ml suggested

strong column interaction. Non-expressed, oligomeric, or highly interacting sybodies were discarded.

Typical yields ranged from 200 µg to 1 mg [16].

Fluorescence imaging of ParB-GFP strains

Image acquisition

B. subtilis cells were first cultured in LB supplemented with 5 µg/mL chloramphenicol and 0.5% glucose

at 30°C, from which a day culture was inoculated to 0.005 in LB with 5 µg/mL chloramphenicol. When

needed, cultures were induced at OD_{600} ~ 0.02 with 0.5% xylose and grown an extra 30 minutes until

imaging at an OD_{600} of 0.04. For microscopy analysis, 0.5 µL of the cell suspension was spotted onto

agarose-coated microscopy slides. Images were acquired using a Leica DMi8 microscope equipped with

an sCMOS DFC9000 (Leica) camera, a SOLA light engine (Lumencor), and a 100×/1.40 oil-immersion

objective. Exposure time for image acquisition was set to 600 ms. Acquired images were processed

using LAS X Office software (v.1.4.7.28982, Leica Microsystems).

Image analysis

Microscopy images were analyzed using a fully automated image processing pipeline. Cell

segmentation was performed with the pretrained Omnipose model[30] implemented in Cellpose[31], which

is optimized for bacterial morphologies. Prior to segmentation, a normalization step was applied to

standardize image contrast. Following segmentation, intracellular foci were detected using Spotiflow[32],

a dedicated model trained for spot identification. Each detected spot was assigned to an individual cell

based on spatial location, enabling quantification of the number of foci per cell. Cell morphology

analysis was also performed to estimate cell length.

Quantitative results were compiled into a structured dataset, including the number of cells per image,

percentage of cells lacking foci, number of foci per cell, cell length, and number of foci per micron. Data

visualization and statistical analysis was performed using GraphPad Prism 10.4.1 (627).

ELISA

E. coli MC1061 colonies were screened for sybody expression, and periplasmic extracts were obtained

as per the original protocol [16]. ELISA plates were coated with Protein A and anti-c-Myc antibody before

incubation with sybody-containing extracts. Three wells per sybody were incubated with biotinylated

maltose binding protein (MBP, negative control) or with 50 nM biotinylated bsSmc-ScpAB(E1118Q) with

ATP/dsDNA or without. Washes and ELISA development were performed using TBSM-D. Absorbance

was measured at 650 nm.

ATPase measurements

ATPase activity was assessed using the coupled pyruvate kinase/lactate dehydrogenase reaction as

described by [14]. ADP production was monitored over a period of 1 hour by measuring the absorbance

changes of NADH at 340 nm. Data collection was performed using a Synergy Neo Hybrid Multi-Mode

Microplate Reader. The reaction mixture comprised 1 mM NADH, 3 mM phosphoenolpyruvic acid, 100

U pyruvate kinase, 20 U lactate dehydrogenase, and varying concentrations of ATP. For assays requiring

double-stranded oligonucleotides, a 40 bp oligonucleotide (5'-TTAGTTGTTC GTAGTGCTCG TCTGGCTCTG

GATTACCCGC-3') was added to a final concentration of 3 mM. The final protein concentration in the

assay was 0.15 μM *bsuS*mc dimers in ATPase assay buffer (50 mM HEPES-KOH, pH 7.5; 50 mM NaCl; 2

mM MgCl2). All measurements were conducted at 25°C.

Acknowledgments

We are grateful to Roberto Vazquez Nunez for the biotinylated *bsuS*mc(C119S, C437S, C826S, E1118Q,

R643C) protein preparation, Arianna Ravera from the DCSR computing facility for help in image analysis

and data extraction of the fluorescence microscopy images. Members of the Gruber Lab for critical

feedback of the manuscript and for stimulating discussions.

Funding

This study was supported by internal funding and by the European Research Council (Horizon 2020 ERC

CoG 724482).

Author contributions

O.G. performed all *in vivo* experiments from sybody selection, fluorescence microscopy imaging,

chimeric phenotypic screen. M.T. and L.H-H. performed the *in vitro* sybody screening (ribosome display,

phage display, and first sybody purification). O.G. performed protein purifications, cross-linking and

ATPase measurements, as well as data analysis, figures and writing. O.G. wrote the first manuscript

draft. All authors reviewed and edited the manuscript. Funding acquisition and supervision, S.G.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All raw data will be made available at Mendeley Data DOI: 10.17632/k6k62p7z2s.1

(https://data.mendeley.com/datasets/k6k62p7z2s/1).

Figure Legends

Figure 1: Selection of sybodies targeting *Smc*-ScpAB in *B. subtilis*.

(A) (i) Structural design and randomization scheme of the three synthetic sybody libraries: concave, loop, and convex. Complementarity-determining regions (CDRs) 1, 2, and 3 are shown in yellow, orange, and red, respectively; randomized residues are shown as sticks. Adapted from [1]. **(ii)** SMC domain organization. The N- and C-terminal regions encode half an ATPase head, while the middle domain constitutes the hinge dimerization domain. The elongated sequence in-between encodes for long helical sequences that will form an antiparallel intramolecular coiled coil. Hd: head, Hng: hinge, 4N: 4N arm-to-arm contact, J: joint. **(iii)** SMC complexes harbor juxtaposed arms in their resting state. Upon ATP binding at the heads, the heads dimerize (head engagement), leading to a conformational change of the entire complex in which the arms open. This state is thought to fit a loop of DNA (not shown) in the S compartment formed by the SMC arms, that will be pushed towards the K compartment formed by the heads and kleisin, after ADP release. We hypothesized that SMC-blocking sybodies can stabilize distinct conformation of the *bsuS*mc-ScpAB complex promoted by ATP hydrolysis and DNA binding, potentially inducing a *Δsmc*-like phenotype *in vivo* and enabling biochemical characterization of trapped conformations *in vitro*. Blue: SMC dimer, Green: kleisin ScpA. The ScpB KITE subunits are not represented for the sake of simplicity. HngDBS: Hinge DNA-binding site, HdDBS: Head DNA-binding site.

(B) Framework for sybody selection. *In vitro* selection starts with ~10^{12} sybody variants per library subjected to ribosome display for pre-enrichment, followed by two rounds of phage display. For *in vivo* selection, 95 randomly selected sybody genes were integrated into the *B. subtilis* chromosome under the control of a xylose-inducible promoter by allelic replacement at the *amyE* locus. Growth defects on rich medium were tested on ONA agar plates with or without 0.5% xylose. Example shown: Sb164 (loop library) did not affect growth, whereas Nb31 impaired growth upon induction, suggesting interference with *bsuS*mc function.

(C) Growth assay results for the 95 *B. subtilis* strains expressing individual xylose-inducible sybodies of

the loop library. Bars show colony counts "without" (green) on top of "with" (pink) xylose. Strains are

ordered by their original position in the 96-well plate. Fourteen sybodies consistently impaired colony

formation under inducing conditions (marked by dotted lines). Sybody numbers indicated above the

plots correspond to selected candidates used in subsequent experiments, numbering according to

order of first use. Notably, the strain corresponding to the E09 sybody (Sb018), showed an absence of

colonies upon sybody induction. However, this sybody candidate gave intermediate phenotypes in later

experiments, which is why it was excluded from detailed analysis.

Figure 2: Sybody-induced chromosome segregation defects visualized by ParB-GFP imaging.

(A) i, *Top*: Schematic illustrating ParB-GFP binding near *oriC*, enabling visualization of origin positioning.

Bottom: Representative images showing ParB-GFP foci in wild-type and *Δsmc B. subtilis*. *Smc* WT cells

typically display 2–4 foci per cell, whereas *Δsmc* cells exhibit reduced foci numbers. ii, Number of ParB-

GFP foci per μm of cell length in a strain carrying inducible sybody Sb006. A significant decrease in foci

is detected from 40 minutes post-induction (p_{t0-10} = 0.9674; p_{t0-20} = 0.9033; p_{t0-30} = 0.3744; p_{t0-40} =

0.0145; p_{t0-50} = 0.0004; p_{t0-60} < 0.001). Based on this, a standardized induction time of ~35 minutes

was used in subsequent experiments.

(B) ParB-GFP foci density (foci/μm) in WT, *Δsmc,* and sybody-expressing strains after 35 minutes of

xylose induction. Violin plots show distribution per condition; solid lines denote the mean, and dotted

lines indicate quartiles. Several hundreds of cells were analyzed (between 351 and 1735), except for

Δsmc and Sb010 were fewer cells were available (134 and 73, respectively).

(C) Spot assay to assess colony formation of *B. subtilis* strains *bsu* or *spn* variants of *S*mc-ScpAB and

ParB. *Top*: Schematic shows gene origins (blue: *B. subtilis*; grey: *S. pneumoniae*). The leftmost column

corresponds to the parental *B. subtilis* strain 1A700 carrying the *S. pneumoniae* genes without

chloramphenicol resistance (non-growing). The next spots represent the same strains carrying no

sybody gene but the Cm resistance (EV for empty vector). Remaining spots are sybody-expressing

strains; sybody numbers are indicated above. Cells were grown for 16 h at 37 °C on ONA supplemented

with 0.5% xylose and chloramphenicol. Hd: head, Hng: hinge, 4N: 4N arm-to-arm contact, J: joint.

Figure 3: Mapping sybody binding sites on Smc-ScpAB.

(A) Colony formation of *B. subtilis* strains expressing chimeric Smc-ScpAB complexes comprising *S.*

pneumoniae and *B. subtilis* sequences. The schematic above the spot assay depicts the species origin

of the *smc* gene, the *scpAB* operon and *parB* gene (blue: *B. subtilis*; grey: *S. pneumoniae*). "–" indicates

no insertion at the *amyE* locus; "EV" refers to the empty vector control containing only the

chloramphenicol resistance cassette that was inserted in *amyE*; numbered labels correspond to

sybodies. Cells were spotted on rich medium (ONA) supplemented with chloramphenicol and xylose

and incubated for 16 h at 37 °C. Hd: head, Hng: hinge, 4N: 4N arm-to-arm contact, J: joint.

(B) ATP hydrolysis rates of *bsu*Smc-ScpAB in the presence of sybodies but absence of DNA. Significant

effects by one-way ANOVA are indicated by p values.

(C) ATPase rates in the presence of 40 bp dsDNA. All sybodies significantly reduced DNA-stimulated ATP

hydrolysis. Reported p-values: Sb020 ($p = 0.340$), Sb194 ($p = 0.0049$), Sb002 ($p = 0.0007$), Sb015 ($p =$

0.0003); all others, $p < 0.0001$.

(D) Schematic summary of sybody binding sites mapped onto the Smc dimer, categorized by their effect

on ATPase activity. Sybodies are grouped based on functional impact and mapped to corresponding

structural regions: pink/yellow boxes indicate residues 318–339 and 836–857; green boxes mark

residues 248–276 and 899–927; and the red box highlights the 4N contact region (approx. residues

290–320).

Figure 4: Proposed models for sybody interactions with *bsuS*mc-ScpAB during the ATPase cycle.

Model for sybodies of the Opening^{UP}, Closing^{UP} and Closing^{DOWN} group. Opening^{UP} sybodies, likely

prevent complete arm closure. In the presence of DNA, these sybodies may stabilize a suboptimal

conformation for head disengagement, resulting in reduced ATPase activity (not shown here). Closing^{UP}

and Closing^{DOWN} sybodies stabilize a closed-arm conformation. These likely hinder DNA segments from

entering the inter-arm space and accessing the hinge-proximal DNA binding site.

Figure S1: Preparation of *bsuS*mc(C119S, C437S, C826S, E1118Q, R643C)-ScpAB complex and loop

extrusion model

(A) Size-exclusion chromatography profile of biotinylated *bsuS*mc(C119S, C437S, C826S, E1118Q,

R643C)-ScpAB complex. Peak fractions (elution at ~13 mL, pooled fractions indicated by dotted lines

for a final volume of 1.8 mL) were collected for downstream use; the final concentration was 5.9 µM

(dimer). * : A secondary peak appeared at ~14 mL; its identity was uncertain and was excluded from

the peak. SDS-PAGE confirms successful biotinylation of the *bsu*Smc(C119S, C437S, C826S, E1118Q,

R643C)-ScpAB complex.

(B) Segment-capture model for *bsuS*mc-ScpAB, in which ATP binding and hydrolysis drive transitions

between open and closed conformations to mediate loop extrusion.

Figure S2: Representative results from *in vivo* sybody selection based on colony formation.

(A) Representative examples of *in vivo* transformation assays for nine sybody-expressing *B. subtilis*

strains. Sybody genes were integrated into the *amyE* locus under xylose-inducible control.

Transformants were grown on oxoid nutrient agar (ONA) plates supplemented with chloramphenicol

and 0.5% xylose to assess growth defects. Shown are candidates from the concave library: Sb051 (well

B08) and Sb078 (D12). Loop library: Sb007 (E05), Sb020 (E12), Sb031 (F05), Sb164 (D12) and NbH07

(Sb194). Convex library: Sb239 (D12) and Sb248 (E10).

(B) Growth assay results for the 285 *B. subtilis* strains expressing individual xylose-inducible sybodies.

Bars show each colony counts "without" (green) on top of "with" (pink) xylose. Strains are ordered by

their original 96-well plate. Only the loop library yielded sybodies that consistently impaired growth

under inducing conditions (dotted line). Sybody numbers indicated above the plots correspond to

positive candidates used in subsequent experiments, numbered according to order of use. Notably, the

strain corresponding to the E09 M sybody (Sb018), also showed an absence of growth upon sybody

induction. However, this sybody candidate gave intermediate phenotypes in later experiments, which

is why it was excluded from this study. Results for the loop library are also shown in Figure 1.

Figure S3: Functional impact of sybody expression on chromosome organization, cell length, and

growth in *B. subtilis*.

(A) Average number of ParB-GFP foci per µm of cell length in strains expressing individual sybodies,

with and without xylose induction. Most uninduced strains differed significantly from the Smc WT

control, likely due to leaky expression from the P_{xyl} ($p < 0.0001$), except Sb015 ($p > 0.9999$), Sb156

($p = 0.2345$), and Sb016 (negative control, $p > 0.9999$).

(B) Average cell length across the same conditions, measured in ParB-GFP strains with or without Smc,

and with or without each sybody. Bars indicate standard deviation. Tested sybodies caused cell

elongation, a characteristic phenotype of impaired Smc activity, as chromosome segregation defects

delay cell division, with mean lengths from 4.70 ± 2.16 µm (Sb020) to 8.68 ± 2.99 µm (Sb010), compared

to 4.51 ± 1.95 µm in wild-type and 9.96 ± 8.92 µm in Δsmc. **(C)** Growth curves of *B. subtilis* strains

expressing individual sybodies compared to the WT strain, with and without xylose induction. Each

curve represents the mean of two biological replicates. Six sybodies were randomly picked and showed

in this figure. Fluorophore-free strains showed wild type-like growth without xylose but strong delays

post-induction, followed by partial growth recovery after 12 hours, likely due to xylose depletion or

suppressor emergence. In sybody-expressing strains, a clear drop in cell density was observed ~2.5

hours post-induction.

Figure S4: Sybody-GFP expression at different inducer concentration

Selected Sybody-GFP proteins were tested by GFP imaging at different inducer concentrations (0, 0.005,

0.05 and 0.5 % xylose). Representative images are shown. Notably, sybody Sb007 (E5, loop library)

generates a mild growth phenotype (smaller colonies but normal colony numbers, Figure 1A and 1B)

but shows good expression levels and focal localization (top row).

Figure S5: Imaging of various Sybody-GFP proteins without inducer.

Selected Sybody-GFP constructs were grown in the absence of inducer and tested for gfp expression.

Representative images are shown.

Figure S6: ATPase hydrolysis rate of the *bsuS*mc-ScpAB complex with each selected sybody and

viability of *B. subtilis* strains expressing chimeric Smc-ScpAB complexes in the absence of sybody

induction.

(A) Spotting assay assessing the growth of *B. subtilis* strains carrying chimeric Smc-ScpAB operons

composed of *S. pneumoniae* and *B. subtilis* components. Strains harbor individual sybody constructs

integrated at the *amyE* locus under control of a xylose-inducible promoter but were grown without

inducer. "–" indicates no integration at *amyE*; "EV" corresponds to an empty vector control containing

only the chloramphenicol resistance cassette. Numbered labels indicate strains carrying specific sybody

constructs. Cells were spotted on rich medium (ONA) containing chloramphenicol and incubated for

16 h at 37 °C.

(B) Overall ATPase hydrolysis rate of the *bsuS*mc-ScpAB complex in presence of each sybody and +/-

DNA. Sb06, 021 and 194 triggered a significant difference in ATP hydrolysis between +/- dsDNA

conditions. The rest is non-significant.

Figure S7: Hypothetical model for an Opening^{DOWN} class of sybodies, not recovered in this study.

This may indicate that stabilizing an open conformation near the ATPase heads does not interfere with

Smc function.

Tables

Table 1: Sybody enrichments at different steps of the selection procedure.

Ribosome display output was quantified by qPCR, while phage display results include final phage titers

and enrichment values from rounds one and two, also measured by qPCR.

Table S1: ELISA sybody binding assay

Table S2: Sybody sequences

Table S3: Percentage of cells lacking ParB-GFP foci, indicating the absence of chromosome due to

segregation defect. In presence and absence of xylose for sybody induction.

Table S4: Strains

Table S5: Plasmids

Table S6: Oligonucleotides

References

1.   Zimmermann, I., Egloff, P., Hutter, C.A., Arnold, F.M., Stohler, P., Bocquet, N., Hug, M.N., Huber,
S., Siegrist, M., Hetemann, L., et al. (2018). Synthetic single domain antibodies for the
conformational trapping of membrane proteins. Elife *7*. 10.7554/eLife.34317.
2.   Pacesa, M., Nickel, L., Schellhaas, C., Schmidt, J., Pyatova, E., Kissling, L., Barendse, P.,
Choudhury, J., Kapoor, S., Alcaraz-Serna, A., et al. (2025). One-shot design of functional protein
binders with BindCraft. Nature *646*, 483-492. 10.1038/s41586-025-09429-6.
3.   Yatskevich, S., Rhodes, J., and Nasmyth, K. (2019). Organization of Chromosomal DNA by SMC
Complexes. Annu Rev Genet *53*, 445-482. 10.1146/annurev-genet-112618-043633.
4.   Burmann, F., and Gruber, S. (2015). SMC condensin: promoting cohesion of replicon arms. Nat
Struct Mol Biol *22*, 653-655. 10.1038/nsmb.3082.
5.   Gruber, S., and Errington, J. (2009). Recruitment of condensin to replication origin regions by
ParB/SpoOJ promotes chromosome segregation in B. subtilis. Cell *137*, 685-696.
10.1016/j.cell.2009.02.035.
6.   Wang, X., Tang, O.W., Riley, E.P., and Rudner, D.Z. (2014). The SMC condensin complex is
required for origin segregation in Bacillus subtilis. Curr Biol *24*, 287-292.
10.1016/j.cub.2013.11.050.
7.   Liu, H.W., Roisne-Hamelin, F., Taschner, M., Collier, J., Srinivasan, M., and Gruber, S. (2025). The
SMC Hinge is a Selective Gate for Obstacle Bypass. Nat Commun *16*, 10457. 10.1038/s41467-
025-65408-5.
8.   Sullivan, N.L., Marquis, K.A., and Rudner, D.Z. (2009). Recruitment of SMC by ParB-parS
organizes the origin region and promotes efficient chromosome segregation. Cell *137*, 697-707.
10.1016/j.cell.2009.04.044.
9.   Diebold-Durand, M.L., Lee, H., Ruiz Avila, L.B., Noh, H., Shin, H.C., Im, H., Bock, F.P., Burmann,
F., Durand, A., Basfeld, A., et al. (2017). Structure of Full-Length SMC and Rearrangements
Required for Chromosome Organization. Mol Cell *67*, 334-347 e335.
10.1016/j.molcel.2017.06.010.
10.   Marko, J.F., De Los Rios, P., Barducci, A., and Gruber, S. (2019). DNA-segment-capture model
for loop extrusion by structural maintenance of chromosome (SMC) protein complexes. Nucleic
Acids Res *47*, 6956-6972. 10.1093/nar/gkz497.
11.   Minnen, A., Burmann, F., Wilhelm, L., Anchimiuk, A., Diebold-Durand, M.L., and Gruber, S.
(2016). Control of Smc Coiled Coil Architecture by the ATPase Heads Facilitates Targeting to
Chromosomal ParB/parS and Release onto Flanking DNA. Cell Rep *14*, 2003-2016.
10.1016/j.celrep.2016.01.066.
12.   Rasmussen, S.G., Choi, H.J., Fung, J.J., Pardon, E., Casarosa, P., Chae, P.S., Devree, B.T.,
Rosenbaum, D.M., Thian, F.S., Kobilka, T.S., et al. (2011). Structure of a nanobody-stabilized
active state of the beta(2) adrenoceptor. Nature *469*, 175-180. 10.1038/nature09648.
13.   Vazquez Nunez, R., Polyhach, Y., Soh, Y.M., Jeschke, G., and Gruber, S. (2021). Gradual opening
of Smc arms in prokaryotic condensin. Cell Rep *35*, 109051. 10.1016/j.celrep.2021.109051.
14.   Burmann, F., Basfeld, A., Vazquez Nunez, R., Diebold-Durand, M.L., Wilhelm, L., and Gruber, S.
(2017). Tuned SMC Arms Drive Chromosomal Loading of Prokaryotic Condensin. Mol Cell *65*,
861-872 e869. 10.1016/j.molcel.2017.01.026.
15.   Hirano, M., and Hirano, T. (2004). Positive and negative regulation of SMC-DNA interactions by
ATP and accessory proteins. EMBO J *23*, 2664-2673. 10.1038/sj.emboj.7600264.
16.   Zimmermann, I., Egloff, P., Hutter, C.A.J., Kuhn, B.T., Brauer, P., Newstead, S., Dawson, R.J.P.,
Geertsma, E.R., and Seeger, M.A. (2020). Generation of synthetic nanobodies against delicate
proteins. Nat Protoc *15*, 1707-1741. 10.1038/s41596-020-0304-x.
17.   Deneka, D., Rutz, S., Hutter, C.A.J., Seeger, M.A., Sawicka, M., and Dutzler, R. (2021). Allosteric
modulation of LRRC8 channels by targeting their cytoplasmic domains. Nat Commun *12*, 5435.
10.1038/s41467-021-25742-w.

18. Diebold-Durand, M.L., Burmann, F., and Gruber, S. (2019). High-Throughput Allelic
Replacement Screening in Bacillus subtilis. Methods Mol Biol *2004*, 49-61. 10.1007/978-1-
4939-9520-2_5.
19. Hutter, C.A.J., Timachi, M.H., Hurlimann, L.M., Zimmermann, I., Egloff, P., Goddeke, H., Kucher,
S., Stefanic, S., Karttunen, M., Schafer, L.V., et al. (2019). The extracellular gate shapes the
energy profile of an ABC exporter. Nat Commun *10*, 2260. 10.1038/s41467-019-09892-6.
20. Gruber, S., Veening, J.W., Bach, J., Blettinger, M., Bramkamp, M., and Errington, J. (2014).
Interlinked sister chromosomes arise in the absence of condensin during fast replication in B.
subtilis. Curr Biol *24*, 293-298. 10.1016/j.cub.2013.12.049.
21. Bock, F.P., Liu, H.W., Anchimiuk, A., Diebold-Durand, M.L., and Gruber, S. (2022). A joint-ParB
interface promotes Smc DNA recruitment. Cell Rep *40*, 111273. 10.1016/j.celrep.2022.111273.
22. Gabler, F., Nam, S.Z., Till, S., Mirdita, M., Steinegger, M., Soding, J., Lupas, A.N., and Alva, V.
(2020). Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. Curr Protoc
Bioinformatics *72*, e108. 10.1002/cpbi.108.
23. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction
with AlphaFold. Nature *596*, 583-589. 10.1038/s41586-021-03819-2.
24. Soh, Y.M., Burmann, F., Shin, H.C., Oda, T., Jin, K.S., Toseland, C.P., Kim, C., Lee, H., Kim, S.J.,
Kong, M.S., et al. (2015). Molecular basis for SMC rod formation and its dissolution upon DNA
binding. Mol Cell *57*, 290-303. 10.1016/j.molcel.2014.11.023.
25. Vazquez Nunez, R., Ruiz Avila, L.B., and Gruber, S. (2019). Transient DNA Occupancy of the SMC
Interarm Space in Prokaryotic Condensin. Mol Cell *75*, 209-223 e206.
10.1016/j.molcel.2019.05.001.
26. Seeger, M.A., Mittal, A., Velamakanni, S., Hohl, M., Schauer, S., Salaa, I., Grutter, M.G., and van
Veen, H.W. (2012). Tuning the drug efflux activity of an ABC transporter in vivo by in vitro
selected DARPin binders. PLoS One *7*, e37845. 10.1371/journal.pone.0037845.
27. Kruse, A.C., Ring, A.M., Manglik, A., Hu, J., Hu, K., Eitel, K., Hubner, H., Pardon, E., Valant, C.,
Sexton, P.M., et al. (2013). Activation and allosteric modulation of a muscarinic acetylcholine
receptor. Nature *504*, 101-106. 10.1038/nature12735.
28. Burmann, F., Shin, H.C., Basquin, J., Soh, Y.M., Gimenez-Oya, V., Kim, Y.G., Oh, B.H., and Gruber,
S. (2013). An asymmetric SMC-kleisin bridge in prokaryotic condensin. Nat Struct Mol Biol *20*,
371-379. 10.1038/nsmb.2488.
29. Denereaz, J., and Veening, J.W. (2024). BactEXTRACT: an R Shiny app to quickly extract, plot and
analyse bacterial growth and gene expression data. Access Microbiol *6*.
10.1099/acmi.0.000742.v3.
30. Cutler, K.J., Stringer, C., Lo, T.W., Rappez, L., Stroustrup, N., Brook Peterson, S., Wiggins, P.A.,
and Mougous, J.D. (2022). Omnipose: a high-precision morphology-independent solution for
bacterial cell segmentation. Nat Methods *19*, 1438-1448. 10.1038/s41592-022-01639-4.
31. Stringer, C., Wang, T., Michaelos, M., and Pachitariu, M. (2021). Cellpose: a generalist algorithm
for cellular segmentation. Nat Methods *18*, 100-106. 10.1038/s41592-020-01018-x.
32. Dominguez Mantes, A., Herrera, A., Khven, I., Schlaeppi, A., Kyriacou, E., Tsissios, G., Skoufa, E.,
Santangeli, L., Buglakova, E., Durmus, E.B., et al. (2025). Spotiflow: accurate and efficient spot
detection for fluorescence microscopy with deep stereographic flow regression. Nat Methods
22, 1495-1504. 10.1038/s41592-025-02662-x.

	Ribosome Display (qPCR)		Phage Display (qPCR)	
Library	Total #of RNAs	Titer (PFU/mL)	Enrichment 1	Enrichment 2
Convex (S)	1.36×10^8	8.96×10^{13}	3.1×	1474.7×
Loop (M)	6.95×10^7	5.46×10^{13}	1.4×	1209.9×
Concave (L)	1.25×10^8	8.32×10^{13}	1.8×	1808.0×

Table 1

	Ribosome Display (qPCR)		Phage Display (qPCR)	
Library	Total #of RNAs	Titer (PFU/mL)	Enrichment 1	Enrichment 2
	1.36×10^8	8.96×10^{13}	3.1×	1474.7×
	5.46×10^{13}		1.4×	1209.9×
	1.25×10^8		1.8×	1808.0×

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1

Figure S3

Figure S4: ΔamyE::Sybody-mGFP::CamR::amyE (xylose titration)

No xylose	0.5 % xylose	0.05 % xylose	0.005 % xylose
20210825 BSG1001 Sb007 No Xyl 1 001	20210825 BSG1001 Sb007 Xyl 2 002	20210825 BSG1001 Sb007 Xyl 3 002	20210825 BSG1001 Sb007 Xyl 4 003
20210825 BSG1001 Sb015 No Xyl 1 002	20210825 BSG1001 Sb015 Xyl 2 002	20210825 BSG1001 Sb015 Xyl 3 001	20210825 BSG1001 Sb015 Xyl 4 005

Figure S5: ΔamyE::Sybody-mGFP::CamR::amyE (No induction)

20210826 BSG1001 Sb002-GFP No Xyl002	20210826 BSG1001 Sb007-GFP No Xyl002	20210826 BSG1001 Sb010-GFP No Xyl001	20210826 BSG1001 Sb015-GFP No Xyl001
20210826 BSG1001 Sb016-GFP No Xyl003	20210826 BSG1001 Sb018-GFP No Xyl003	20210826 BSG1001 Sb021-GFP No Xyl001	20210826 BSG1067 Smc-GFP003

A

Sybody: - EV 002 003 006 010 015 020 021 031 156 160 162 169 188 194

bsuSmc-ScpAB

spnSmc-ScpAB

Smc chimera

ONA + CM
9⁻⁵ dilution
16hrs growth

B

** < 0.0001
* = 0.02
Rest = NS

Figure S6

Opening^DOWN

?

Figure S7