Noncanonical amino acid incorporation enables minimally disruptive labeling of stress granule and TDP-43 proteinopathy

Hao Chen[1, 2], Haocheng Wang[1], Yuning Lu[1, 2], Peng Chen[1, 2], Zhongfan Zheng[1, 2], Tao Zhang[1, 2], Jiou Wang[1, 2*]

[1] Department of Biochemistry and Molecular Biology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, 21205, USA.

[2] Department of Neuroscience, School of Medicine, Johns Hopkins University, Baltimore, MD, 21205, USA.

*To whom correspondence should be addressed:

Jiou Wang, E-mail: jiouw@jhmi.edu

Key words: Genetic code expansion, Anap labeling, Stress granule, TDP-43

Abstract

We report a minimally disruptive labeling strategy for stress granule protein G3BP1 and ALS-

linked protein TDP-43 using the fluorescent noncanonical amino acid Anap. By integrating

genetic code expansion with rational site selection, we achieved precise incorporation of Anap

that preserves protein structure and function. In live cells and neurons, Anap labeling faithfully

recapitulated localization, stress-induced dynamics, and recovery behavior, outperforming

conventional fluorescent tags and enabling physiologically relevant visualization of protein

pathobiology.

Introduction

Fluorescent protein labeling remains a cornerstone of live-cell biology, yet conventional techniques rely heavily on large fusion tags, such as auto-fluorescent tags (AFPs) or small-molecule-binding motifs[1], at limited positions (typically N-or C-terminal). These tags might potentially affect the structure, function, and even localization pattern of proteins, limiting their use for studying proteins with complex dynamics[2]. Alternatively, genetic code expansion (GCE) has emerged as a versatile labeling strategy to label proteins site-specifically in a minimally disruptive manner[3-8].

GCE employs an engineered orthogonal aminoacyl tRNA synthetase/tRNA pair to incorporate non-canonical amino acids (ncAAs) at desired positions of proteins according to reassigned codons, most commonly the amber stop codon (TAG), thereby introducing a single-residue substitution within the protein of interest. Among these ncAAs, L-Anap (3-(6-acetylnaphthalen-2-ylamino)-2-aminopropanoic acid) is especially attractive for live imaging. Anap is intrinsically fluorescent, exhibits polarity-sensitive emission spectra, and requires no post-incorporation modification[2, 5]. Despite these advantages, GCE-based Anap labeling has rarely been systematically applied to track disease-relevant protein dynamics in live mammalian cells. In this study, we developed an Anap-based labeling platform optimized for minimally disruptive labeling of two important proteins, G3BP1 and TDP-43, involved in membraneless organelles and neurodegenerative diseases, such as amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD).

G3BP1 is a core protein in stress granules, dynamic membraneless organelles related to stress response[9]. Chronically aberrant dynamics of stress granules with observed cytotoxicity have also been implicated as a common feature of ALS/FTD[10, 11]. TDP-43 cytoplasmic inclusion is a hallmark of ALS/FTD pathology[12] and is closely associated with dysregulation of RNA metabolism, ultimately leading to cellular defects[13]. Both proteins undergo dynamic and stress-

49 dependent redistribution[9], making them ideal candidates for minimally invasive, site-specific

50 fluorescent labeling.

Results and Discussion

To implement the site-specific Anap incorporation system, we selected and generated two

amber mutants, G3BP1^{F337TAG} and TDP-43^{V100TAG}, using a combination of structural and

functional criteria: exclusion from functional domains or localization signals, absence of disease-

associated mutations, lack of post-translational modification, and low predicted structural impact

by AlphaFold models. Incorporation of Anap was achieved via co-expression of an orthogonal

tRNA/synthetase pair in cells.

We first tested the feasibility of the Anap labeling system for G3BP1. In HeLa cells, G3BP1-

Anap localized diffusely in the cytoplasm under basal conditions, closely matching antibody

staining. Interestingly, a nuclear signal was detected with Anap but not with antibody, indicating

the presence of nuclear pools of G3BP1 inaccessible to antibody detection. Upon sodium

arsenite treatment, both the Anap and antibody signals colocalized within stress granules (Fig.

1A), validating the ability of Anap labeling to visualize the dynamics of G3BP1-driven stress

granule formation. To assess labeling fidelity, we compared the performance of G3BP1-Anap

with G3BP1-GFP using fluorescence recovery after photobleaching (FRAP). Following stress,

both labeled proteins localized to granules, but G3BP1-Anap exhibited significantly higher

fluorescence recovery (~53%) than G3BP1-GFP (~33%) (Fig. 1C, 1F). These results suggested

that Anap labeling better preserved the native mobility and biophysical properties of G3BP1

than the conventional GFP tag.

We next applied Anap labeling to TDP-43. Under basal conditions, the signal of TDP-43-Anap

overlapped with that of anti-TDP-43 antibody staining, predominantly within the nucleus (Fig.

1B). Following sodium arsenite treatment, TDP-43-Anap mislocalized to cytoplasmic inclusions,

validated by antibody staining. By contrast, TDP-43-YFP failed to recapitulate this cytoplasmic

mislocalization, instead forming prominent nuclear puncta under stress conditions (Fig. 1E),

suggesting that large C-terminal tags may distort native localization of the protein. We then

performed FRAP analysis to test the mobility of TDP-43-Anap in these cytoplasmic inclusions.

Fluorescence recovery of TDP-43-Anap reached ~45% within 20 seconds post-bleaching (Fig.

1D, 1G), consistent with liquid-like dynamics. This differs from previous reports of immobile

aggregates in nuclear localization signal (NLS)-deleted TDP-43[14], illustrating the value of Anap

labeling in preserving native protein properties, allowing real-time assessment of pathological

dynamics without requiring disruptive mutations.

To determine whether TDP-43-Anap retains biological function, we expressed it in a TDP-43

knockout HeLa cell line (Fig. 1H) and tested cell viability under oxidative stress. Following 12.5

µM sodium arsenite treatment for 24 hours, the expression of TDP-43-Anap significantly

rescued cell survival, reaching levels comparable to wild-type TDP-43 overexpression (45% vs

46%; Fig. 1I). We further validated this finding in a mouse embryonic stem (ES) cell model with

an inducible TDP-43 knockout (iTDPKO). For mouse ES cells, TDP-43 KO alone was sufficient

to induce cell death. Here, either wild-type TDP-43 or TDP-43-Anap was expressed in iTDPKO

mouse ES cells, where TDP-43 was deleted following induction with 4-hydroxytamoxifen (4-HT).

Expression of TDP-43-Anap restored ES cell viability nearly to wild-type TDP-43 (47% vs 51%;

Fig. 1J). Together, these results confirmed that Anap incorporation preserves the biological

function of TDP-43.

Furthermore, to extend Anap labeling to neuronal systems, we applied this approach to label

both proteins in primary mouse cortical neurons. The neurons were co-stained with human-

specific anti-TDP-43 or human-specific anti-G3BP1 antibodies. Under basal conditions, the

signal of G3BP1-Anap colocalized with antibody staining in the cytoplasm and relocalized to

stress granule upon sodium arsenite treatment (Fig. 2A, 2B). Notably, nuclear Anap signal was

again observed in neurons, suggesting additional pools of G3BP1 not captured by antibody

staining.

TDP-43-Anap localized to the neuronal nucleus under basal conditions and showed strong

colocalization with anti-TDP-43 antibody. Interestingly, antibody staining appeared more

diffusely cytoplasmic than Anap, suggesting improved signal specificity of Anap labeling with

direct genetic incorporation. Under oxidative stress, both Anap and antibody signals colocalized

within cytoplasmic inclusions (Fig. 2C, 2D). These findings demonstrated that the Anap system

for G3BP1 and TDP-43 performs robustly in neuronal environments, a key setting for ALS/FTD

research.

In summary, our results demonstrated that Anap-based genetic code expansion provides a

minimally disruptive strategy for tracking the dynamic behavior of G3BP1 and TDP-43 in live

cells (Fig. 3). In G3BP1, Anap faithfully reported stress granule assembly and preserved native

mobility, unlike GFP fusions that impaired dynamics. In TDP-43, Anap labeling maintained

nuclear localization under basal conditions and revealed liquid-like behavior of cytoplasmic

inclusions during stress, in contrast to the aberrant nuclear puncta produced by YFP-tagged

TDP-43. Critically, Anap-labeled TDP-43 retained biological activity, rescuing cell survival in

TDP-43–deficient HeLa and stem cells. Moreover, we have generated stable TDP-43–Anap cell

lines that exhibited consistent expression, protein localization, and stress-induced aggregation,

providing stable cell models for TDP-43 research.

By enabling high-fidelity visualization of both stress granule dynamics and TDP-43 aggregation

in live cells and primary neurons, Anap labeling bridges a critical gap between structural

preservation and functional readout. The ability to monitor native protein behavior without

perturbation provides a unique opportunity to study early events in ALS/FTD progression, such

as stress granule maturation, protein cytoplasmic mislocalization, and aggregate fluidity, at a

resolution inaccessible with conventional tagging approaches.

Acknowledgements

This work was supported by grants from NIH (NS110098). We thank Dr. Peter Schultz from

Scripps Research for providing the pAnap plasmid, Dr. Shawn Ferguson from Yale University for

the TDP-43 knockout HeLa cell line, and Dr. Philip Wong from Johns Hopkins University for the

iTDPKO mouse ES cell line. We also thank the members of Wang's lab for discussion and

suggestions.

Author contributions

H.C. performed and analyzed most of the experiments. H.C.W. helped with plasmid preparation,

western blot, and imaging. Y.N.L. provides cells and helped with data preparation. P.C. provides

mouse primary cortical neurons. Z.F.Z helped with imaging. T.Z. helped culture iTDPKO mouse

ES cells. H.C. and J.W. designed the studies and wrote the paper with inputs from other

authors.

Methods and materials

General information

Oligonucleotide synthesis was performed by IDT, and Sanger sequencing of DNA plasmids and

PCR products was performed by Quintara. L-ANAP (trifluoroacetate salt) (15436) used in this

study was purchased from Cayman. For immunostaining, the following primary and secondary

antibodies were used: mouse anti-human G3BP (BD Biosciences, 611126), rabbit anti-TDP-43

(Proteintech, 10782-2-AP), mouse anti-TDP-43 (human-specific, monoclonal; Proteintech,

60019-2-Ig), chicken anti-βIII-tubulin (Tuj1; GeneTex, GTX85469), donkey anti-mouse Alexa

Fluor 555 (Thermo Fisher, A-31570), donkey anti-rabbit Alexa Fluor 488 (Thermo Fisher, A-

21202), donkey anti-chicken Alexa Fluor 647 (Thermo Fisher, A78952), and donkey anti-rabbit

Alexa Fluor Plus 800 (Thermo Fisher, A32808).

Plasmids

pAnap plasmid was a gift from Peter Schultz's lab[2]. pCMV-G3BP1 was purchased from Sino

Biologics. pRK5-TDP-43, pEGFP-G3BP1, and pCMV/TO-TDP-43-YFP were conducted by our

lab. Site-directed mutagenesis was conducted using the NEB Q5 site-directed mutagenesis kit

(NEB, E0554). TAG substitutions were introduced by PCR amplification with Q5 Hot Start High-

Fidelity DNA Polymerase using primers designed with NEBaseChanger. Then the PCR products

were incubated with an enzyme mix consisting of a kinase, a ligase, and DpnI to rapidly

circularize the PCR products and remove the template DNA. And then the mix will be

transformed into DH5α cells.

Cell culture

Mammalian cell lines were cultured at 37°C and 5% CO_2 in humidified incubators. HeLa cells

(ATCC) were cultured in DMEM/F12 (Corning, 10-013-CV) supplemented with 10% FBS (Gibco,

A5256801). The TDP-43 knockout HeLa cell line was a gift from Dr. Shawn M. Ferguson[15].

Primary cortical neurons were prepared as previously described [16]. Briefly, the cortex of the

mouse embryo was separated into HBSS on ice, then digested into single-cell suspension with

0.25% trypsin and 0.1mg/ml DNase I at 37°C for 20 min. Dissociated cells were washed twice

and resuspended in plating media (DMEM supplemented with 10% FBS) before seeding onto

poly-D-lysine (Gibco, A3890401)-coated plates. Cells were cultured for 3-4 hours in plating

medium to settle down, and then the medium was replaced with maintenance medium

(neurobasal medium (Gibco, 21103049)) supplemented with 2% B-27 supplement

(Gibco,17504044), 1% GlutaMax (Gibco, 35050061), and 1% penicillin/streptomycin (Gibco,

15140163). The iTDPKO inducible mouse embryonic stem ES cell line was a gift from Wong's

lab[17]. Cells were maintained on attachment factor (Gibco, S-006-100) coated plates in 2i media

containing half of DMEM/F12 and half of Neurobasal media complemented with L-Glutamine

(Gibco, 25030081), B-27 supplement, N2 supplement (Gibco, 17502048), BSA (Gibco,

15260037), 1% penicillin/streptomycin (Gibco, 15140163), PD0325901 (MedChemExpress,

391210-10-9), CHIR99021(MedChemExpress, 252917-06-9), monothioglycerol(Millipore

Sigmal, M6145), and mlif(Millipore Sigma, ESG1107).

Transfection

HeLa cells were transfected using Lipofectamine 2000 reagent (ThermoFisher, 11668030). Cells

were seeded 24 h before transfection, and plasmid DNA was mixed with Lipofectamine 2000 at

a ratio of 1 μg DNA:2 μL reagent in Opti-MEM medium (Thermo Fisher, 31985070). After 15 min

incubation at room temperature, the complexes were added to the cells. Six hours post-

transfection, the medium was replaced with fresh growth medium supplemented with or without

20 μM L-Anap, and cells were incubated overnight before further experiments.

Mouse primary cortical neurons and inducible TDP-43 knockout (iTDPKO) mouse ES cells were

transfected using Lipofectamine 3000 (Thermo Fisher, L3000015) according to the

manufacturer's instructions. For neurons, first, the total DNA was diluted and mixed with p3000

reagent in Opti-MEM media, and the Lipofectamine 3000 was diluted in Opti-MEM media

separately. Second, both diluted DNA and Lipofectamine 3000 were mixed and incubated at

room temperature for 15 minutes. The final mix would then be added to cells seeded on poly-D-

lysine-coated plates at DIV5. After incubation overnight, the media were half-replaced with fresh

neurobasal media with 10μM Anap, and the neurons were incubated for two additional days (the

media were half-replaced each day to keep the Anap supply). Differently, for mouse ES cells,

the DNA-lipo3000 mix was added directly into the cell suspension, and the suspension was

added to attachment factor-coated plate. After overnight incubation, the media for ES cells was

replaced with fresh 2i media supplemented with 10μM Anap and any further treatments.

Anap labeling and immunofluorescence in cell fixation

6h post-transfection, the cells were cultured in medium supplemented with L-Anap for another

20-24 h. Cells were then washed three times with DPBS to remove excess Anap and incubated

in fresh medium for 1 h before treatment with 250 μM sodium arsenite for anther 1h. After the

treatment, the cells were washed with DPBS three times and then fixed with 4%

paraformaldehyde (PFA; Millipore Sigma, 158127) for 15-20 mins at room temperature. Fixed

cells were permeabilized and blocked in immunofluorescence blocking buffer (Cell Signaling

Technology, 12411) containing 0.1% Triton X-100 (Millipore Sigma, X100) for 45 min.

Immunostaining was performed using anti-TDP-43 and anti-G3BP1 antibodies in both HeLa

cells and primary cortical neurons, with anti-Tuj1 antibody included as a neuronal marker.

Images were acquired using a Leica SP8 confocal microscope.

Anap labeling and GFP/YFP tagging in live-cell imaging

For Anap labeling, HeLa cells were seeded on 35-mm glass-bottom dishes (Cellvis, D35-20-

1.5H), transfected, and incubated with L-Anap following the same protocol used for fixed-cell

imaging. After removal of excess Anap, cells were cultured in fresh medium for 2-3 hours before

live cell imaging. For GFP/YFP tagging, cells expressing the tagged proteins were directly ready

for live cell imaging. Before FRAP, cells expressing Anap-labeled or GFP/YFP-tagged TDP-43

or G3BP1 were treated with 250 µM sodium arsenite for 1 hour. Next, regions of interest (ROIs)

corresponding to protein signals were selected for FRAP analysis using a Leica SP8 confocal

microscope.

Cell survival tests

Cell viability was assessed using Calcein AM staining (Invitrogen, C1430). HeLa cells were

treated with 12.5 µM sodium arsenite for 24 h, followed by incubation with 3 µM Calcein AM.

Fluorescence from viable cells was measured using a Synergy H1 Hybrid Multi-Mode Plate

Reader (BioTek) at excitation/emission wavelengths of 485/535 nm.

For mouse ES cells, the conditional knockout of TDP-43 was induced by 4-hydroxytamoxifen (4-

HT, Millipore Sigma, H7904-5MG) treatment (300ng/ml) for 5 days. Cell viability was assessed

using the Cell Counting-Lite 2.0 Luminescent Cell Viability Assay Kit (Vazyme, DD1101-02).

Immunoblotting

Cells were washed three times with PBS and lysed in ice-cold RIPA buffer [50 mM Tris-HCl (PH

7.5), 150 mM NaCl, 1% NP40, 0.1% SDS, 100 mM NaF, 0.5% sodium deoxycholate, 17.5 mM

beta-glycerophosphate, 1 mM PMSF, and protease inhibitor cocktail (1:200, Millipore Sigma,

P8340)]. Lysates were sonicated and clarified by centrifugation (12,000 rpm, 20 min, 4 °C), and

supernatants were collected. Protein concentrations were determined using the BCA assay

(Thermo Fisher, Cat. No. 23225). Equal amounts of protein were resolved by SDS-PAGE and

transferred to membranes, which were blocked with 5% BSA and incubated with primary

antibodies overnight at 4 °C. After three washes with TBST, membranes were incubated with

fluorescence-conjugated secondary antibody dilution at room temperature for 2h. Blots were

scanned and imaged by LI-COR Odyssey M scanner.

Image analysis and statistical analysis

All images were processed and analyzed by Fuji/ImageJ software, and the colocalization of

signals was analyzed by the colocalization threshold analysis plugin in ImageJ. Experiments

requiring quantification were repeated at least three times independently. The FRAP results

were analyzed by ImageJ to quantify the fluorescence intensity at each time point. The average

relative intensities (normalized to pre-bleaching intensity) of each time point were analyzed in

Prism 10 software. For survival tests, the average relative rates (normalized to the control

group) for each group were analyzed in Prism 10 software, and the one-way analysis of

variance (ANOVA) was used to compare the significance between each group. All data were

presented as means ± SEM.

References:

1. Crivat, G. & Taraska, J.W. Imaging proteins inside cells with fluorescent tags. *Trends in biotechnology* **30**, 8–16 (2012).
2. Chatterjee, A., Guo, J., Lee, H.S. & Schultz, P.G. A Genetically Encoded Fluorescent Probe in Mammalian Cells. *Journal of the American Chemical Society* **135**, 12540–12543 (2013).
3. Hao, M. et al. Tracking endogenous proteins based on RNA editing-mediated genetic code expansion. *Nat Chem Biol* **20**, 721–731 (2024).
4. Nikić, I., Kang, J.H., Girona, G.E., Aramburu, I.V. & Lemke, E.A. Labeling proteins on live mammalian cells using click chemistry. *Nature Protocols* **10**, 780–791 (2015).
5. Nygaard, A., Zachariassen, L.G., Larsen, K.S., Kristensen, A.S. & Loland, C.J. Fluorescent non-canonical amino acid provides insight into the human serotonin transporter. *Nature Communications* **15**, 9267 (2024).
6. Bessa-Neto, D. et al. Bioorthogonal labeling of transmembrane proteins with non-canonical amino acids unveils masked epitopes in live neurons. *Nature Communications* **12**, 6715 (2021).
7. Eddins, A.J. et al. Quantitative Protein Labeling in Live Cells by Controlling the Redox State of Encoded Tetrazines. *Journal of the American Chemical Society* **147**, 23625–23634 (2025).
8. Arsić, A., Hagemann, C., Stajković, N., Schubert, T. & Nikić-Spiegel, I. Minimal genetically encoded tags for fluorescent protein labeling in living neurons. *Nature Communications* **13**, 314 (2022).
9. Yang, P. et al. G3BP1 is a tunable switch that triggers phase separation to assemble stress granules. *Cell* **181**, 325–345 (2020).
10. Kassouf, T. et al. Targeting the NEDP1 enzyme to ameliorate ALS phenotypes through stress granule disassembly. *Science Advances* **9**, eabq7585 (2023).
11. Van Nerom, M. et al. C9orf72-linked arginine-rich dipeptide repeats aggravate pathological phase separation of G3BP1. *Proceedings of the National Academy of Sciences* **121**, e2402847121 (2024).
12. Neumann, M. et al. Ubiquitinated TDP-43 in Frontotemporal Lobar Degeneration and Amyotrophic Lateral Sclerosis. *Science* **314**, 130–133 (2006).
13. Suk, T.R. & Rousseaux, M.W.C. The role of TDP-43 mislocalization in amyotrophic lateral sclerosis. *Molecular Neurodegeneration* **15**, 45 (2020).
14. Yan, X. et al. Intra-condensate demixing of TDP-43 inside stress granules generates pathological aggregates. *Cell* **188**, 4123–4140.e4118 (2025).
15. Roczniak-Ferguson, A. & Ferguson, S.M. Pleiotropic requirements for human TDP-43 in the regulation of cell and organelle homeostasis. *Life science alliance* **2** (2019).
16. Chen, P. et al. Spine impairment in mice high-expressing neuregulin 1 due to LIMK1 activation. *Cell Death & Disease* **12**, 403 (2021).
17. Chiang, P.-M. et al. Deletion of TDP-43 down-regulates Tbc1d1, a gene linked to obesity, and alters body fat metabolism. *Proceedings of the National Academy of Sciences* **107**, 16320–16324 (2010).

Figure legend

Fig. 1: Anap labeling of TDP-43 and G3BP1 in HeLa cells.

A, B. HeLa cells expressing G3BP1-Anap and TDP-43-Anap under basal conditions or 250 μM

sodium arsenite treatment. Anap and antibody signals are shown in blue and green,

respectively; for merged panels, Anap was pseudo-colored red. Scale bars: 10 μm (overview), 3

μm (zoom). C, D. FRAP of G3BP1-Anap, G3BP1-GFP, and TDP-43-Anap, following 250 μM

sodium arsenite treatment. ROI signal intensities are displayed in rainbow RGB (red–high, blue–

low). Scale bars: 5 μm (cells), 1 μm (ROI). E. Comparison of HeLa cells expressing TDP-43-

Anap and TDP-43-YFP under basal conditions or 250 μM sodium arsenite treatment. Here,

Anap labeling and YFP labeling yield a blue signal and a yellow to green signal, respectively. F,

G. Relative fluorescence recovery of each time point after photobleaching for G3BP1-Anap,

G3BP1-GFP, and TDP-43-Anap. H. Immunoblotting of wild-type TDP-43 and TDP-43-Anap. The

Rabbit anti-TDP-43 antibody was used. KO: TDP-43 knockout; V100Stop/Anap: the expression

of TDP-43^{V100TAG} via Anap labeling without or with the addition of Anap. I. Survival of TDP-43

knockout HeLa cells expressing TDP-43-Anap after being treated with 12.5 μM sodium arsenite

for 24h. Here, the calcien EM staining was used to detect the survival rate of cells, and the

relative survival rate= sodium arsenite treatment/ nontreatment for each group. OE:

overexpression of TDP-43. J. Survival of iTDPKO inducible mouse ES cells expressing TDP-43-

Anap. Here, cell counting-Lite 2.0 Luminescent cell viability assay kit was used to detect the

survival rate of ES cells. TDP-43 knockout was induced by 4-HT (300ng/ml) for 5 days. The

relative survival rate= 4-HT induction/DMSO for each group. All quantitative data (F, G, I, J) are

shown as mean ± SEM. ***P = 0.0001; ****P < 0.0001; n.s., not significant.

Fig. 2: Anap labeling of TDP-43 and G3BP1 in neurons.

A, C. Primary mouse cortical neurons expressing G3BP1-Anap and TDP-43-Anap under basal conditions or 250 μM sodium arsenite treatment. Cells were stained with anti-G3BP1 (human-specific) or anti-TDP-43 (human-specific) antibodies with chicken anti-Tuj1 as a neuron marker. Signals: Anap (blue, pseudo-colored red in merged images), antibody (green), Tuj1 (gray). Scale bar, 10 μm. B, D. The colocalization level of each region of interest for G3BP1 and TDP-43. Colocalization threshold analysis in Fuji ImageJ was used to analyze the R value of each region (a higher R value means a higher colocalization level). Scale bar, 1 μm.

Fig. 3: Schematic of the Anap labeling system for G3BP1 and TDP-43 using genetic code expansion.

Briefly, two plasmids were required to express the protein with site-specific Anap incorporation, one for Anap incorporation and one for the mutated protein of interest (TAG introduction). As the plasmids were transfected into cells, the orthogonal Anap-tRNA synthetase would charge the Anap to its cognate tRNA, and the tRNA would incorporate the Anap site-specifically into the protein of interest in response to the TAG stop codon. Cells expressing Anap-labeled TDP-43 and G3BP1 were subsequently imaged by confocal microscopy, either after fixation or in live-cell conditions.

Fig. 1

Fig. 2

Fig. 3