Gain and loss of plasmid-borne antibiotic resistance genes are associated with their chromosomal counterparts and the broader plasmid gene repertoire in Enterobacteriaceae

Yang Liu[1, *], Yue Liu[1,2, *]

[1] Department of Microbiology, Zhejiang University School of Medicine, Zhejiang, China

[2] State Key Laboratory for Diagnosis and Treatment of Infectious Diseases of the First Affiliated Hospital, Zhejiang University School of Medicine, Zhejiang, China

[*]Correspondence: yang.liu29@zju.edu.cn (Yang L.); yuel@zju.edu.cn (Yue L.)

ABSTRACT

Plasmids are central vehicles for the dissemination of antibiotic resistance genes (ARGs). They are among the most mobile and evolvable genetic elements, with broad host ranges and high rates of gene turnover, making them especially effective in spreading antibiotic resistance across bacterial lineages. Using the phylogeny-aware gene gain and loss model applied to 6,895 Enterobacteriaceae genomes, we quantified four evolutionary processes—gene gain, loss, expansion, and reduction—for plasmid-borne genes. We found that, overall, plasmid-borne ARGs (pARGs) exhibit similar gain and loss rates compared with other plasmid genes, but significantly higher expansion and reduction rates. All four processes were strongly species dependent, with only a minor influence of antibiotic class. Further, pARGs were co-transferred with a broad range of plasmid functional genes, including defense, metabolic, and mobilome modules throughout evolutionary history. Moreover, bacterial clades harboring chromosomal ARGs (cARGs) showed significantly higher acquisition of plasmid-borne resistance than did their sister clades lacking corresponding cARGs. Our results demonstrate that pARGs in Enterobacteriaceae are co-transferred with the broader plasmid gene repertoire on an evolutionary timescale, whereas copy-number dynamics are distinct from plasmids. Moreover, cARGs can serve as markers of lineages predisposed to accumulate plasmid-mediated resistance.

KEYWORDS antibiotic resistance genes; Enterobacterales; gene gain and loss, plasmid; mobile genetic elements.

Importance

Plasmids play a central role in the spread of antibiotic resistance genes (ARGs), and the long-term evolutionary behavior of plasmid-borne ARGs (pARGs) could provide insights into the emergence of novel multidrug resistance. We studied nearly 7,000 Enterobacteriaceae genomes and show that pARGs evolve through the same gain and loss processes as other plasmid genes but exhibit markedly higher and species-dependent copy number changes. Moreover, bacterial lineages carrying chromosomal ARGs are significantly more likely to acquire additional plasmid-mediated resistance. This association suggests that plasmid and chromosomal ARGs could act synergistically to accelerate resistance accumulation within specific lineages. Identifying such evolutionarily high-risk lineages may provide a basis for predicting and monitoring the emergence of multidrug resistance.

INTRODUCTION

Antibiotic resistance is a major, ongoing challenge in both medicine and evolutionary biology (Davies and Davies 2010). In bacterial genome evolution, gene gain through horizontal gene transfer (HGT) and gene loss (via selection) are the primary forces shaping genomic repertoires (Koonin and Wolf 2008, Koonin and Wolf 2012). Mobile genetic elements (MGEs), particularly conjugative plasmids, serve as principal vehicles of HGT and are central to the spread of antibiotic resistance genes (ARGs) (Bennett 2008, Partridge, Kwong et al. 2018, Redondo-Salvo, Fernández-López et al. 2020).

Comparative genomic analyses show that plasmid taxonomic units carrying ARGs exhibit significantly higher mobility, broader host ranges, and elevated rates of

homologous recombination and gene turnover—indicating that resistance

dissemination is facilitated by intrinsically plastic and mobile plasmid lineages (Coluzzi

and Rocha 2025). Their plasticity is enhanced by high densities of transposable

elements and recombination systems, which accelerate gene repertoire turnover

(Amos, Ploumakis et al. 2018, Che, Yang et al. 2021). Moreover, evolutionary

innovations such as replication-control mutations can increase plasmid copy number,

thereby amplifying resistance levels and enhancing transfer efficiency (Dimitriu,

Matthews et al. 2021). Thus, plasmids are highly dynamic replicons characterized by

frequent horizontal transfer, rapid turnover, and extensive genetic plasticity (San Millan

and MacLean 2017, San Millan 2018).

Given their evolutionary plasticity, understanding plasmid-mediated ARG turnover

is particularly critical in clinically important bacterial families such as

Enterobacteriaceae (San Millan 2018). The Enterobacteriaceae family harbors

extensive plasmid diversity and constitutes one of the largest reservoirs of clinically

relevant ARGs (Carattoli 2013, Mathers, Peirano et al. 2015, Rozwandowicz, Brouwer

et al. 2018). Members of this family, including *Escherichia*, *Klebsiella*, *Enterobacter*,

and *Salmonella*, are major contributors to multidrug-resistant infections in hospitals

and communities worldwide (van Duin and Paterson 2016). Although numerous clinical

studies have documented outbreaks of antibiotic resistance within the

Enterobacteriaceae family (Grundmann, Livermore et al. 2010, Mathers, Peirano et al.

2015, van Duin and Paterson 2016), most of these reports have focused on the

dissemination of particular resistance plasmids or the epidemiology of individual ARGs

(Carattoli 2013, Mathers, Peirano et al. 2015, Rozwandowicz, Brouwer et al. 2018).

However, key questions remain as to how plasmid-borne ARGs in Enterobacteriaceae

are maintained and amplified across lineages, and whether their evolutionary

dynamics differ from those of other plasmid-encoded functions.

Here, we systematically investigate ARG turnover dynamics in

Enterobacteriaceae plasmids by applying a phylogeny-aware birth-death model to

6,895 genomes. Specifically, we (i) quantify the balance of ARG gain, loss, expansion,

and reduction across diverse species, (ii) identify non-ARG gene modules that

consistently co-transfer with ARGs, and (iii) assess the association between

chromosomal and plasmid-borne resistance repertoires.

RESULTS

Plasmid-borne antibiotic resistance genes show higher expansion and

reduction rates but similar gain and loss dynamics compared with other

plasmid genes

To investigate the evolutionary dynamics driving the current distribution of antibiotic

resistance genes (ARGs) in Enterobacteriaceae plasmids, we applied a phylogenetic

birth-death model using Count (Csuros 2010) to quantify gene turnover rates across

6,895 genomes of which 6,130 contained at least one identified plasmid sequence

(Supplementary Table S1). The approach estimates four key evolutionary processes:

gain (appearance of new gene families), loss (disappearance of entire gene families),

expansion (increase in gene copy number within existing families), and reduction

(decrease in gene copy number within existing families). Since we focused exclusively

on plasmid-borne genes, expansion and reduction can be interpreted as copy number

increases or decreases of plasmid gene families within individual genomes. We

compared overall rates between plasmid borne ARGs (pARGs) and all other plasmid

genes (non-ARGs) across 98 species. Our analysis revealed that pARGs exhibit

substantially different dynamics for gene gain and loss versus copy number variation

(Figure 1, Supplementary Table S2). Both gain and loss rates showed no significant

differences between pARGs and non-ARGs (n = 98, gain: adjusted p = 0.173, loss:

adjusted p = 0.080; Figure 1A-B), with data points forming relatively compact

distributions across species. This suggests that the fundamental processes governing

the acquisition and elimination of entire gene families operate similarly for resistance

and non-resistance genes. In contrast, pARGs exhibited significantly elevated rates for

both expansion (n = 97, adjusted p = 3.21×10^{-5}, median $\log_2$(ARG/ALL) = 1.09; Figure

1C) and reduction (adjusted p = 4.17×10^{-4}, median $\log_2$(ARG/ALL) = 0.39; Figure 1D)

events. The high scatter in these data points indicated substantial species-specific

variation in the extent to which expansion and reduction dynamics differ between

pARGs and other plasmid genes, suggesting that genes copy number fluctuations on

the genomes are strongly influenced by which bacterial species carries the plasmid.

The analysis of copy-number changes revealed that pARGs showed balanced

dynamics between expansion and reduction rates (median $\log_2$ ratio = -1.38, adjusted

p = 0.47, Supplementary Figure S1). In contrast, the overall gain/loss ratios displayed

gain bias for both pARGs and non-ARGs (ARGs: median $\log_2$(gain/loss) = 3.87,

adjusted p = 4.27×10^{-8}; non-ARGs: median = 4.69, adjusted p = 6.35×10^{-15}; Figure

1E-F).

In summary, across Enterobacteriaceae, gain and loss rates show no pARG-

specific signal, indicating that acquisition and elimination of pARG families follow

common plasmid evolutionary pathways. In contrast, pARGs exhibit significantly

elevated expansion and reduction rates with pronounced species-level variation,

revealing that copy-number dynamics are shaped by lineage-specific factors.

Turnover dynamics of antibiotic resistance genes are primarily species

dependent

To test whether the species-level patterns in Figure 1 are consistent across resistance

mechanisms or vary among drug classes, which are defined by their mechanisms of

action, we assigned orthologous gene families to CARD-defined drug classes and

estimated gain, loss, expansion, and reduction rates for each class across species.

Using the CARD database (Alcock, Huynh et al. 2023), we identified and analyzed the

turnover dynamics of specific drug classes, which are groups of antibiotics with similar

chemical structures and mechanisms of action. Our analysis revealed that the

evolutionary rates for all four event types were strongly species-dependent (Kruskal–

Wallis test; gain: $H = 735.9$, $p = 7.6 \times 10^{-99}$, $\eta^2 = 0.84$; loss: $H = 660.9$, $p = 9.1 \times 10^{-85}$, η^2

$= 0.74$; expansion: $H = 511.5$, $p = 5.9 \times 10^{-58}$, $\eta^2 = 0.56$; reduction: $H = 583.2$, $p =$

7.5×10^{-71}, $\eta^2 = 0.66$), with substantial variation across species for gain, loss, expansion,

and reduction (Supplementary Figure S2A–B; Figure 2A–B). In contrast, drug class

effects were weak: only marginal signals were detected for gain ($p = 0.021$, $\eta^2 = 0.02$)

and expansion ($p = 0.019$, $\eta^2 = 0.02$), while loss ($p = 0.97$, $\eta^2 \approx 0$) and reduction (p

$= 0.35$, $\eta^2 < 0.01$) showed no significant drug class dependence. Full results are

provided in Supplementary Table S3.

Furthermore, species that exhibited high expansion rates typically also showed

elevated reduction rates for the same drug classes (n=96, Spearman's rank correlation

with 5,000 permutations: ρ=0.57, permutation p≈0.0002). By contrast, species with

high gain rates did not necessarily display the corresponding high loss rates (n=98,

ρ=−0.15, p≈0.145), indicating that plasmid-borne gene copy number increases and

decreases within existing gene families are more tightly coupled than complete gene

family acquisition and loss.

In summary, antibiotic resistance gene evolution shows distinct signatures across

event types: gain and loss are strongly species dependent with negligible drug class

influence, expansion is likewise dominated by species with only minor drug class

effects, and reduction is also primarily species dependent. Moreover, expansion and

reduction are coupled, whereas gain and loss are largely independent.

Plasmid-borne antibiotic resistance genes co-transfer with diverse plasmid

functional modules and often entire plasmids

To identify genes that are potentially co-transferred with antibiotic resistance genes on

plasmids, we quantified the relationship between per-branch gain and loss rates of

pARGs with those of individual COG-annotated gene families within each species. For

each species tree, branch-level rates were pooled to compute overall correlations, and

significance was evaluated using Spearman correlation followed by phylogenetic

Brownian motion simulations to control for phylogenetic effects.

Focusing on 85 species with plasmid borne pARGs exhibiting non-zero gain or

loss rates, we found correlated evolutionary dynamics between pARGs and genes

from diverse functional categories within plasmids (Figure 3, Supplementary Table S4).

Although our study focuses on plasmid-encoded genes, many correlated COG

categories were not mobilomes (COG category X), but rather belonged to categories

such as defense system (V), various metabolic functions, and cellular processes, often

with high numbers of unique COG families involved (Figure 3A; line thickness reflects

this number). Considering both the number of unique COG families involved (Figure

3A) and the number of species showing significant correlations (Figure 3B), gain and

loss correlations occurred at similar frequencies, indicating that co-dynamics were not

restricted to a single event type (full results in Supplementary Table S4). At the species

level, genes that tended to co-gain with pARGs also tended to co-loss with them over

evolutionary timescales (Figure 3B).

Additionally, 29 species showed significant correlations when all COG categories

were analyzed together, 18 of which involved both gain and loss correlations

(Supplementary Table S4), suggesting that in these species, pARGs may be

transferred together with entire plasmids rather than as isolated genes. A comparison

of plasmid length between the 29 species with significant correlations across all COG

categories and the 56 species without such correlations revealed a statistically

significant difference (Mann–Whitney U test, $p < 0.001$); however, the effect size was

negligible (Cliff's delta = –0.034, Cohen's d = –0.081), indicating that plasmid size does

not account for the observed correlations.

These findings provide quantitative evidence that co-transfer patterns extend

beyond a few functional categories and encompass a broad spectrum of plasmid-

encoded genes, with different COG categories contributing to varying degrees. The

consistent coupling of high gain rates with high loss rates across gene categories

indicates that plasmid-borne genes experience bidirectional turnover, with frequent

acquisition accompanied by frequent loss. Importantly, in nearly one-third of the

species analyzed, pARGs co-transferred together with the broader plasmid gene

repertoire, indicating that entire plasmids often represent the fundamental units of

resistance dissemination.

Chromosomal antibiotic resistance genes (cARGs) are associated with elevated

pARG acquisition rates

Our previous analyses revealed that pARGs exhibit similar gain and loss rates

compared to other plasmid genes. We further investigated whether additional factors

might be associated with the evolutionary dynamics of pARGs, specifically examining

whether the presence of antibiotic resistance genes in bacterial chromosomes is

associated with altered evolutionary rates of pARGs. We employed a comparative

phylogenetic approach using sister clade analysis (Figure 4A), identifying 436 pairs of

phylogenetic clades where one clade (lineage) contained ≥ 80% of genomes

harboring a specific ARG family in their chromosome (ARG-containing clade) while

<20% of genomes in the sister clade (lineage) contained that homologous ARG family.

For each clade pair, we calculated gain and loss rates for plasmid genes by summing

evolutionary events across all branches within each clade and normalizing by total

branch length. In total, 78 species and 79 RGI-defined ARG families were included in

the analysis (Supplementary Table S5 and S6). Linear mixed-effects models (LMM)

were used as statistical test for the analysis, including fixed effects for clade type, gene

class, and their interaction, with random effects for species, ARG family, and sister-

clade comparisons to capture lineage-specific variation (see Methods).

Our analysis revealed that bacterial clades with cARGs exhibit significantly

different plasmid gene gain rates compared with their sister clades (Figure 4B-C).

ARG-containing clades showed approximately 10-fold higher gain rates for pARGs

(estimate = 2.30 on the log scale, p = 1.9×10⁻⁷; Figure 4B, Supplementary Table S7)

and modestly but significantly lower loss rates (p = 0.034). In contrast, other plasmid

genes showed significantly lower gain rates in ARG-containing clades compared with

sister clades (about threefold lower; estimate = −1.12, p = 0.0069), while loss rates

showed no significant differences between clade types (p = 0.30). The interaction term

indicated that the correlation between cARG presence and evolutionary rates differs

significantly between gene types for gain events (p = 1.7 × 10⁻⁸) but not for loss events

(p = 0.418, Figure 4C, Supplementary Table S7).

In addition to the overall association, we performed per-family LMM analyses for

those with at least three available clades in Enterobacteriaceae (Supplementary Table

S8), the overall gene gain rate remained higher in cARG clades (Supplementary Figure

S3; p < 0.05). For each ARG family, we found significantly higher gene-gain rates on

the ARG clade relative to its sister clade for ANT(3"), chloramphenicol

acetyltransferase (CAT), fluoroquinolone-resistant *gyrB*, sulfonamide-resistant *sul*, and

trimethoprim-resistant dihydrofolate reductase (*dfr*) (Supplementary Figure S4). We

further summarized the species distribution and lineage-level genome counts for these

five ARG families (Figure 5), suggesting that in these species, ARGs tend to be more

frequently acquired on plasmids over evolutionary timescales. In contrast, several

families showed lower gene-loss rates in the ARG-containing clade, including APH(3'),

OXA-10-like β-lactamase, and SHV β-lactamase (Supplementary Figure S4). These

results suggest that elevated gene gain rates are not consistent across all ARG families.

These results indicate that bacterial lineages harboring cARGs exhibit enhanced

acquisition of plasmid-borne resistance, with preferential gain of pARGs but not other

plasmid genes in cARG-containing clades.

DISCUSSION

Across 6,895 Enterobacteriaceae genomes, we show that plasmid-borne antibiotic

resistance genes (pARGs) and other plasmid genes exhibit similar species-level gain

and loss rates (Figure 1A–B). This parallels our co-transfer analyses, in which pARG

gain and loss dynamics were significantly correlated with those of diverse plasmid

functions, including defense, metabolic, and mobilome modules (Figure 3A-B).

Moreover, we observed 29 species with significant correlations between pARGs and

overall plasmid gene dynamics (Supplementary Table S4). Together, these findings

suggest that pARGs are integrated into broader plasmid evolutionary processes, with

plasmids acting as units of turnover and selection rather than ARGs following distinct

evolutionary rules. This observation supports the view that plasmid genomes evolve

as integrated genetic modules rather than collections of independently evolving genes

(Puigbò, Lobkovsky et al. 2014), and suggesting monitoring and controlling resistance

may require targeting plasmid populations as whole evolutionary units.

Importantly, our observed gain bias in pARGs (Figure 1E-F) contrasts with the loss

bias dynamics that characterize bacterial genomes and other mobile genetic elements

more generally (Puigbò, Lobkovsky et al. 2014, Puigbo, Makarova et al. 2017). One

possible explanation is that lineage-specific features of Enterobacteriaceae favor

acquisition, consistent with recent evidence that mobile element dynamics are strongly

species dependent (Liu, Botelho et al. 2025). Moreover, plasmid turnover may not

conform to the general loss-biased evolutionary rule governing bacterial genome

dynamics.

In contrast to gene family gain and loss, pARGs exhibited significantly elevated

expansion and reduction rates compared to other plasmid genes (Figure 1C-D). These

copy-number dynamics were highly variable across species, and statistical tests

showed that species identity, rather than resistance class, explained most of the

observed variance (Figure 2A-B, Supplementary Table S3). Weak drug class signals

were detectable for expansion and gain, but effect sizes were negligible

(Supplementary Table S3). Elevated copy-number turnover may reflect the activity of

gene-level mobile elements such as IS26 translocatable units (He, Hickman et al. 2015,

Harmer and Hall 2016) and integron cassette rearrangements (Partridge, Tsafnat et al.

2009, Souque, Escudero et al. 2021), as well as selective amplification events that

duplicate or individual genes without affecting entire plasmids (Nicoloff, Hjort et al.

2024). Besides, recent study revealed that 77% of studied ARG-carrying plasmids

contain insertion sequences, facilitating inter-plasmid resistance gene transfer(Coluzzi

and Rocha 2025). And the replication-control mutations (e.g., *copA*) may also elevate

plasmid copy number and increase ARG transfer rates (Dimitriu, Matthews et al. 2021)

Beyond species-specific variation in copy-number turnover, which likely reflects

differing balances between antibiotic-driven selection and fitness costs(Vogwill and

MacLean 2015, Pal and Andersson 2024), we observed that bacterial lineages

harboring cARGs acquired pARGs at ~10-fold higher rates than their sister clades

(Figure 4B, Supplementary Table S7). This pattern mirrors clinical observations of high-

risk clones that accumulate resistance genes across both chromosomal and plasmid

compartments (Mathers, Peirano et al. 2015, Rozwandowicz, Brouwer et al. 2018).

Rather than implying causation, the association suggests that ARG accumulation is a

lineage-level property: clades predisposed to high plasmid turnover are more likely to

fix ARGs in their chromosomes, functioning as 'super-accumulators' of resistance

(Woodford, Turton et al. 2011, Wang and Dagan 2024, Zongo, Cabanel et al. 2024).

From a clinical perspective, cARGs may thus serve as genomic markers for identifying

bacterial lineages with elevated capacity for plasmid-mediated resistance acquisition,

underscoring the need for enhanced surveillance and infection control targeting these

high-risk lineages.

We note that the cARG families showing elevated acquisition of pARGs in our

analyses (e.g., ANT(3"), CAT, *sul*, and *dfr*) are consistently highlighted across lineages

(Figure 5). These families correspond to antibiotic classes that remain widely applied

or frequently detected in clinical and agricultural settings (Manyi-Loh, Mamphweli et al.

2018, Bacanli and Basaran 2019, Dominguez, Miranda et al. 2019). For example,

Providencia rettgeri and *Citrobacter spp*. have been reported in agricultural and

zoonotic contexts to carry resistance to aminoglycosides and sulfonamides (Liu, Zhang

et al. 2021, Ahmed, Islam et al. 2023, Wang, Li et al. 2023, Valiatti, Santos et al. 2024).

Such lineages are likely exposed to overlapping selective pressures from multiple

antibiotic classes, which may accelerate pARG turnover, promote their integration into

chromosomes, and in turn facilitate the acquisition of additional pARGs. Thus, the

combined action of chromosomal and plasmid-borne ARGs may drive the rapid

accumulation of multidrug resistance within these taxa.

Limitations

This study focused exclusively on plasmid-associated genes, excluding other

mobile elements such as Integrative and conjugative elements (ICEs) and integrative

mobilizable elements (IMEs), which also mediate chromosomal integration and

excision of ARGs (Johnson and Grossman 2015, Cury, Touchon et al. 2017, Botelho

2023). Bacteriophages further can contribute to ARG dissemination through

transduction and phage–plasmid hybrids (Touchon, Moura de Sousa et al. 2017).

Consequently, our observations represent only a subset of the overall resistance

mobilome. Moreover, we lacked ecological information about the habitats and antibiotic

exposure histories of the analyzed organisms. Expanding future analyses to include

additional mobile genetic element and ecological datasets would provide a more

comprehensive understanding of ARG dissemination in natural environments.

Conclusion

Our findings highlight a duality in plasmid-mediated resistance evolution in

Enterobacteriaceae. At the level of whole gene families, pARGs follow the same

turnover dynamics as do other plasmid genes, indicating that resistance dissemination

largely reflects plasmid-level processes rather than ARG-specific rules. At the level of

copy-number changes, however, pARGs show elevated expansion and reduction with

strong lineage dependence and negligible drug-class effects. Such species-driven

volatility suggests recurrent expansion–reduction cycles as a flexible resistance

strategy. Moreover, chromosomal resistance content predicts a lineage's propensity

for further plasmid-mediated resistance acquisition, positioning cARGs as potential

markers of high-risk lineages.

METHODS

Genome Collection and Gene Annotation

We parsed the Genome Taxonomy Database (GTDB; https://gtdb.ecogenomic.org)

release 220 (Parks, Chuvochina et al. 2020) to identify genomes belonging to

Enterobacteriaceae. For species with more than 300 genomes, we randomly selected

300 representatives with random seed 19940421. We applied MIMAG criteria (Bowers,

Kyrpides et al. 2017) with completeness >99%, contamination <1%, mean contig

length >5,000 bp, and contig count <500 to filter high-quality genomes. To reduce

computational cost, species with more than 100 genomes were subsampled to retain

at most 100 genomes (random seed 19940421). Only species with more than 10

genomes were considered for further analysis. The genomes that passed these filters

were downloaded from the NCBI FTP site (https://ftp.ncbi.nlm.nih.gov).

Bacterial genes were predicted using pyrodigal-gv v0.3.2 integrated in geNomad.

Plasmids and their encoded genes were identified using geNomad v1.11.0 with

database v1.9 and default parameters (Camargo, Roux et al. 2024). Antibiotic

resistance genes were identified with Resistance Gene Identifier (RGI) v6.0.5 using

protein sequences with "include_nudge" option and default parameters, CARD version

4.0.1 (Alcock, Huynh et al. 2023). Pan-genomes of each species were constructed

using the orthology-based clustering method OrthoFinder v2.5.5 (Emms and Kelly

2019) following established guidelines (Manzano-Morales, Liu et al. 2023).

For COG functional annotation, we constructed HMM profiles from the COG 2024

database accessed on 2025-02-20 (Galperin, Vera Alvarez et al. 2025). Protein

sequences from each COG family were grouped and aligned using MAFFT v7.526

(Katoh and Standley 2013) with --auto for algorithm selection. Multiple sequence

alignments were trimmed using ClipKIT v2.6.1 (Steenwyk, Buida et al. 2020) with

smart-gap mode to remove poorly aligned regions while preserving phylogenetic signal.

HMM profiles were built for each COG family using HMMER v3.4 (http://hmmer.org)

hmmbuild with default parameters. Predicted proteins from each genome were

searched against the COG HMM database using hmmsearch with an E-value cutoff of

1e-3.

After applying these criteria, 6,895 complete or nearly complete genomes

belonging to 159 Enterobacteriaceae species (sensu GTDB) were included in the

analysis (Supplementary Table S1), of which 98 contained antibiotic resistant genes

on plasmids (Supplementary Data).

Applying Phylogenetic Birth-Death Model for Gene Gain and Loss Inference

From the plasmid gene annotations obtained through geNomad and the orthologous

gene families and species trees generated by OrthoFinder for each species, we

extracted plasmid orthologous gene families. An orthologous gene family was

considered plasmid-associated if the proportion of plasmid genes within that

orthologous gene family exceeded 70% of all genes in the family. Species trees were

pruned to retain only genomes contributing to the filtered orthologous gene families.

Gene gain and loss dynamics were quantified using the phylogenetic birth-death

model implemented in Count v10.04 (Csuros 2010). This probabilistic framework

models the evolution of gene family sizes along phylogenetic trees as a continuous-

time Markov process. The model accounts for three fundamental processes: gene gain

(κ), gene loss (μ), and gene duplication (λ). A gene family of size n changes according

to the following rates: the family decreases at rate $n\mu$ due to individual gene losses,

increases through duplication at rate $n\lambda$, and gains new genes at rate κ through

horizontal transfer or other acquisition mechanisms.

For each species dataset, we performed iterative maximum likelihood optimization

to estimate birth-death model parameters (Puigbò, Lobkovsky et al. 2014). We

implemented a systematic search strategy starting with initial parameter values of

κ=μ=λ=1 and incrementally exploring the parameter space up to κ, μ, λ ≤ 4. This

resulted in a total of 10 optimization rounds. The iterative approach followed the pattern:

if all parameters are equal, increase loss rate (μ); if duplication and gain rates are equal,

increase duplication rate (λ); otherwise increase gain rate (κ). Each round uses the

previous round's results as starting values. The optimization utilized maximum

likelihood estimation with maximum paralogs of 200, optimization tolerance of 0.1, and

minimum lineages of 2.

After parameter optimization, we calculated posterior probabilities for ancestral

gene family states and evolutionary events using the Posteriors module of Count. This

reconstructs the evolutionary history of each gene family, providing probabilistic

estimates for gain, loss, expansion, and reduction events along each branch. Count

analysis was performed separately for each species dataset.

Gene Gain and Loss Data Integration and Rate Calculation

Antibiotic resistance gene families were identified using a guilt-by-association

approach. If any gene within a plasmid-associated orthologous gene family matched

CARD reference sequence through RGI analysis, the entire orthologous gene family

was classified as ARG gene families. Branch lengths were extracted from the original

species trees and combined with corresponding Count posterior probabilities. For rate

calculations, we summed gain, loss, expansion, and reduction events across all

branches within each orthologous gene family and divided by the corresponding total

branch length. Only branches with branch lengths > 1e-12 were included to avoid

computational artifacts from extremely short branches. Evolutionary rates were

calculated per species per orthologous gene family as:

$$rate_i = \frac{\sum(events_{\{i,j\}})}{\sum(branch\_length_j)}$$

where $events_{\{i,j\}}$ is the number of events for orthologous gene family i on branch j,

and $branch\_length_{\{j\}}$ is the length of branch j in substitutions per site.

Comparative Analysis of Evolutionary Rates

For comparative analysis between ARG and non-ARG plasmid gene families, data

were aggregated by species, event type, and ARG status. Median rates were

calculated within each group, and the data were restructured into a paired format

where each species served as a matched pair with corresponding median rates for

ARG and non-ARG families. Paired Wilcoxon signed-rank tests compared ARG versus

non-ARG families within species to control for species-specific evolutionary factors.

Additionally, gain/loss and expansion/reduction ratios were calculated per species to

assess the relative balance of these complementary processes. Multiple testing

correction was performed using the Benjamini-Hochberg procedure. Effect sizes were

quantified as median log2-fold changes, and statistical significance was assessed at p

< 0.05 and q < 0.05 (FDR-adjusted p).

To examine rate variation across antibiotic resistance classes, drug class

annotations were extracted from RGI output. Each ARG-containing orthologous gene

family was annotated to its corresponding drug classes. If an orthologous gene family

matched multiple antibiotic classes, we count it in each matching class. Evolutionary

rates were then visualized across different species and drug class within

Enterobacteriaceae. To assess concordance between event types across species, we

compared median per-species rates using Spearman's rank correlation with

permutation tests (5,000 iterations, random seed = 19940421) to obtain empirical p-

values.

Identification of Genes Co-transferred with Antibiotic Resistance Genes

To identify genes potentially co-transferred with ARGs through horizontal gene transfer,

we assessed whether ARG and non-ARG plasmid genes exhibited similar evolutionary

patterns across phylogenetic branches within each species.

For data preparation, branch-specific evolutionary events from Count analysis

were merged with ARG annotations from RGI and COG functional annotations. In

particular, gene families were classified as ARG-containing based on RGI matches,

and all plasmid gene families were annotated with their consensus COG assignments

(requiring ≥75% consistency within each orthologous gene family). Only COG families

with at least 5 genes per species were retained for analysis. Event rates were

calculated by summing gain and loss events for each gene family at each phylogenetic

branch (both tips and internal nodes). ARG event rates were aggregated across all

resistance genes, while rates of non-ARG plasmid genes were calculated both globally

and per individual COG family.

We then compared the branch-specific event rates between ARGs and individual

COG-annotated gene families using phylogenetic Brownian motion (BM) simulations

to test for correlations while accounting for phylogenetic structure. The analysis used

Spearman correlation on both tip and internal node data, with BM simulations providing

null distributions for significance testing. Initial screening was performed with a

minimum of 5 data points per comparison and 100 simulations. Gene families showing

significant correlations (adjusted p < 0.05 after Benjamini-Hochberg correction) were

subjected to progressively stringent re-analyses: first with a minimum of 8 data points

and 200 simulations, then with 10 data points and 500 simulations. Only gene families

maintaining significance through all three rounds were considered as potential co-

transfer candidates. Species trees were pre-processed using midpoint rooting, with

zero-length branches adjusted by adding small random values (1e-13 to 5e-13) to

avoid artifacts. Phylogenetic Brownian-motion traits were simulated using fastBM, and

ancestral states were reconstructed using fastAnc, both in the phytools package v2.4.4

(Revell 2024) for R v4.3.0.

The identified co-transfer candidate genes were then categorized according to

COG functional categories. Individual COG categories were mapped to four major

process groups: information storage and processing, cellular processes and signaling,

metabolism, and poorly characterized functions.

Comparison of Plasmid Gene Dynamics Between ARG-containing and Sister

Clades

To assess whether the presence of antibiotic resistance genes in bacterial

chromosomes influences the evolutionary dynamics of plasmid-borne genes, we

identified pairs of phylogenetic clades that differed in the chromosomal presence of an

ARG family. For each combination of species tree and ARG family, we scanned for

internal nodes where ≥80% of descendant genomes harbored the specific ARG family

in their chromosome (ARG-containing clade), while <20% of genomes in the

corresponding sister clade contained that ARG family. Only clade pairs with at least

two genomes in each clade were retained for analysis.

For each identified clade pair, we calculated the gain and loss rates of plasmid

genes, distinguishing between plasmid-borne ARGs and other plasmid genes. For

each orthologous gene family, rates were computed as:

492
$$Rate_{OG, clade} = \frac{\sum_{b \in clade} Events_{OG, b}}{\sum_{b \in clade} Branch\ Length_b}$$

where events were summed across all branches within each clade, normalized by the

total branch length of that entire clade. Plasmid ARGs that matched the chromosomal

ARG family used to define the clade were excluded from the analysis. The median rate

across all orthologous gene families within each gene class (ARGs or non-ARGs) was

then used as the representative value for each clade.

Event rates were analyzed using linear mixed-effects models on the log scale, with

gain and loss modeled separately; the response was $\log(rate + \varepsilon)$, where $\varepsilon =$

$0.1 \times \min(\text{non-zero rate})$ within the corresponding event type. Fixed effects were clade

type (ARG-containing vs sister clade), gene class (plasmid ARGs vs other plasmid

genes), and their interaction; random effects were species, resistance gene family, and

the clade pair linking each ARG-containing clade with its sister (implemented as a

random intercept on Clade ID). Models were fitted using restricted maximum likelihood

and restricted to paired observations with both clade types available within a given

gene class. Post hoc contrasts from estimated marginal means assessed (i) ARG-containing vs sister differences within each gene class and (ii) the clade-type × gene-class interaction. In addition to the overall model, we performed family-specific analyses. For each resistance gene family with ≥ 4 available paired clades, we re-fit the model within that family using the same fixed-effects structure and random intercepts for species and clade pair, but omitting the family random effect (not identifiable within a single family). Estimates and 95% CIs were back-transformed to the original rate scale, with two-sided tests and Benjamini–Hochberg adjustment for multiple contrasts. Analyses were performed in R v4.4.0 using lme4 v1.1.35.5 (Bates, Mächler et al. 2015), lmerTest v3.1.3 (Kuznetsova, Brockhoff et al. 2017), and emmeans v 1.11.2.8.

ACKNOWLEDGEMENTS

This work was funded by the ZJU100 Young Professor Start-up Fund from Zhejiang University of China (to Yue Liu).

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES IN THE WRITING PROCESS

During the preparation of this work, the author(s) used ChatGPT to polish the language. After using this tool, the authors reviewed and edited the content as needed and took full responsibility for the content of the publication.

DECLARATION OF COMPETING INTERESTS

The authors declare no competing financial interests.

ETHICS APPROVAL

There are no ethical concerns in this study.

DATA AVAILABILITY

Supplementary material is available for this article. Also, the supplement data, and core

script are available on Figshare under the DOI: 10.6084/m9.figshare.30186307.

AUTHOR CONTRIBUTIONS

Yang Liu: conceptualization, bioinformatics analysis, writing – original draft, writing –

review & editing. **Yue Liu:** writing – review & editing, funding acquisition.

References

Ahmed, T., M. S. Islam, N. Haider, L. Elton, B. Hasan, M. Nuruzzaman, M. T. Rahman,
S. M. L. Kabir and M. S. R. Khan (2023). "Phenotypic and Genotypic Characteristics
of Antimicrobial Resistance in Citrobacter freundii Isolated from Domestic Ducks (Anas
platyrhynchos domesticus) in Bangladesh." Antibiotics (Basel) **12**(4).

Alcock, B. P., W. Huynh, R. Chalil, K. W. Smith, A. R. Raphenya, M. A. Wlodarski, A.
Edalatmand, A. Petkau, S. A. Syed, K. K. Tsang, S. J. C. Baker, M. Dave, M. C.
McCarthy, K. M. Mukiri, J. A. Nasir, B. Golbon, H. Imtiaz, X. J. Jiang, K. Kaur, M. Kwong,
Z. C. Liang, K. C. Niu, P. Shan, J. Y. J. Yang, K. L. Gray, G. R. Hoad, B. F. Jia, T.
Bhando, L. A. Carfrae, M. A. Farha, S. French, R. Gordzevich, K. Rachwalski, M. M.
Tu, E. Bordeleau, D. Dooley, E. Griffiths, H. L. Zubyk, E. D. Brown, F. Maguire, R. G.
Beiko, W. W. L. Hsiao, F. S. L. Brinkman, G. Van Domselaar and A. G. McArthur (2023).
"CARD 2023: expanded curation, support for machine learning, and resistome
prediction at the Comprehensive Antibiotic Resistance Database." Nucleic Acids
Research **51**(D1): D690-D699.

Amos, G. C. A., S. Ploumakis, L. Zhang, P. M. Hawkey, W. H. Gaze and E. M. H.

Wellington (2018). "The widespread dissemination of integrons throughout bacterial communities in a riverine system." ISME J **12**(3): 681-691.

Bacanli, M. and N. Basaran (2019). "Importance of antibiotic residues in animal food." Food Chem Toxicol **125**: 462-466.

Bates, D., M. Mächler, B. M. Bolker and S. C. Walker (2015). "Fitting Linear Mixed-Effects Models Using lme4." Journal of Statistical Software **67**(1): 1-48.

Bennett, P. M. (2008). "Plasmid encoded antibiotic resistance: acquisition and transfer of antibiotic resistance genes in bacteria." Br J Pharmacol **153 Suppl 1**(Suppl 1): S347-357.

Botelho, J. (2023). "Defense systems are pervasive across chromosomally integrated mobile genetic elements and are inversely correlated to virulence and antimicrobial resistance." Nucleic Acids Res **51**(9): 4385-4397.

Bowers, R. M., N. C. Kyrpides, R. Stepanauskas, M. Harmon-Smith, D. Doud, T. B. K. Reddy, F. Schulz, J. Jarett, A. R. Rivers, E. A. Eloe-Fadrosh, S. G. Tringe, N. N. Ivanova, A. Copeland, A. Clum, E. D. Becraft, R. R. Malmstrom, B. Birren, M. Podar, P. Bork, G. M. Weinstock, G. M. Garrity, J. A. Dodsworth, S. Yooseph, G. Sutton, F. O. Glöckner, J. A. Gilbert, W. C. Nelson, S. J. Hallam, S. P. Jungbluth, T. J. G. Ettema, S. Tighe, K. T. Konstantinidis, W. T. Liu, B. J. Baker, T. Rattei, J. A. Eisen, B. Hedlund, K. D. McMahon, N. Fierer, R. Knight, R. Finn, G. Cochrane, I. Karsch-Mizrachi, G. W. Tyson, C. Rinke, A. Lapidus, F. Meyer, P. Yilmaz, D. H. Parks, A. M. Eren, L. Schriml, J. F. Banfield, P. Hugenholtz, T. Woyke and G. S. Consortium (2017). "Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea." Nature Biotechnology **35**(8): 725-731.

Camargo, A. P., S. Roux, F. Schulz, M. Babinski, Y. Xu, B. Hu, P. S. G. Chain, S. Nayfach and N. C. Kyrpides (2024). "Identification of mobile genetic elements with geNomad." Nat Biotechnol **42**(8): 1303-1312.

Carattoli, A. (2013). "Plasmids and the spread of resistance." Int J Med Microbiol **303**(6-7): 298-304.

Che, Y., Y. Yang, X. Xu, K. Brinda, M. F. Polz, W. P. Hanage and T. Zhang (2021). "Conjugative plasmids interact with insertion sequences to shape the horizontal transfer of antimicrobial resistance genes." Proc Natl Acad Sci U S A **118**(6).

Coluzzi, C. and E. P. C. Rocha (2025). "The Spread of Antibiotic Resistance Is Driven by Plasmids Among the Fastest Evolving and of Broadest Host Range." Molecular Biology and Evolution **42**(3).

Csuros, M. (2010). "Count: evolutionary analysis of phylogenetic profiles with
parsimony and likelihood." <u>Bioinformatics</u> **26**(15): 1910-1912.

Cury, J., M. Touchon and E. P. C. Rocha (2017). "Integrative and conjugative elements
and their hosts: composition, distribution and organization." <u>Nucleic Acids Res</u> **45**(15):
8943-8956.

Davies, J. and D. Davies (2010). "Origins and evolution of antibiotic resistance."
<u>Microbiol Mol Biol Rev</u> **74**(3): 417-433.

Dimitriu, T., A. C. Matthews and A. Buckling (2021). "Increased copy number couples
the evolution of plasmid horizontal transmission and plasmid-encoded antibiotic
resistance." <u>Proc Natl Acad Sci U S A</u> **118**(31).

Dominguez, M., C. D. Miranda, O. Fuentes, M. de la Fuente, F. A. Godoy, H. Bello-
Toledo and G. Gonzalez-Rocha (2019). "Occurrence of Transferable Integrons and sul
and dfr Genes Among Sulfonamide-and/or Trimethoprim-Resistant Bacteria Isolated
From Chilean Salmonid Farms." <u>Front Microbiol</u> **10**: 748.

Emms, D. M. and S. Kelly (2019). "OrthoFinder: phylogenetic orthology inference for
comparative genomics." <u>Genome Biol</u> **20**(1): 238.

Galperin, M. Y., R. Vera Alvarez, S. Karamycheva, K. S. Makarova, Y. I. Wolf, D.
Landsman and E. V. Koonin (2025). "COG database update 2024." <u>Nucleic Acids Res</u>
53(D1): D356-D363.

Grundmann, H., D. M. Livermore, C. G. Giske, R. Canton, G. M. Rossolini, J. Campos,
A. Vatopoulos, M. Gniadkowski, A. Toth, Y. Pfeifer, V. Jarlier, Y. Carmeli and C. W.
Group (2010). "Carbapenem-non-susceptible Enterobacteriaceae in Europe:
conclusions from a meeting of national experts." <u>Euro Surveill</u> **15**(46).

Harmer, C. J. and R. M. Hall (2016). "IS26-Mediated Formation of Transposons
Carrying Antibiotic Resistance Genes." <u>mSphere</u> **1**(2).

He, S., A. B. Hickman, A. M. Varani, P. Siguier, M. Chandler, J. P. Dekker and F. Dyda
(2015). "Insertion Sequence IS26 Reorganizes Plasmids in Clinically Isolated
Multidrug-Resistant Bacteria by Replicative Transposition." <u>mBio</u> **6**(3): e00762.

Johnson, C. M. and A. D. Grossman (2015). "Integrative and Conjugative Elements
(ICEs): What They Do and How They Work." <u>Annu Rev Genet</u> **49**: 577-601.

Katoh, K. and D. M. Standley (2013). "MAFFT multiple sequence alignment software
version 7: improvements in performance and usability." <u>Mol Biol Evol</u> **30**(4): 772-780.

Koonin, E. V. and Y. I. Wolf (2008). "Genomics of bacteria and archaea: the emerging
dynamic view of the prokaryotic world." <u>Nucleic Acids Res</u> **36**(21): 6688-6719.

Koonin, E. V. and Y. I. Wolf (2012). "Evolution of microbes and viruses: a paradigm
shift in evolutionary biology?" <u>Front Cell Infect Microbiol</u> **2**: 119.

Kuznetsova, A., P. B. Brockhoff and R. H. B. Christensen (2017). "lmerTest Package:
Tests in Linear Mixed Effects Models." <u>Journal of Statistical Software</u> **82**(13): 1-26.

Liu, L., L. Zhang, H. Zhou, M. Yuan, D. Hu, Y. Wang, H. Sun, J. Xu and R. Lan (2021).
"Antimicrobial Resistance and Molecular Characterization of Citrobacter spp. Causing
Extraintestinal Infections." <u>Front Cell Infect Microbiol</u> **11**: 737636.

Liu, Y., J. Botelho and J. Iranzo (2025). "Timescale and genetic linkage explain the
variable impact of defense systems on horizontal gene transfer." <u>Genome Res</u> **35**(2):
268-278.

Manyi-Loh, C., S. Mamphweli, E. Meyer and A. Okoh (2018). "Antibiotic Use in
Agriculture and Its Consequential Resistance in Environmental Sources: Potential
Public Health Implications." <u>Molecules</u> **23**(4).

Manzano-Morales, S., Y. Liu, S. Gonzalez-Bodi, J. Huerta-Cepas and J. Iranzo (2023).
"Comparison of gene clustering criteria reveals intrinsic uncertainty in pangenome
analyses." <u>Genome Biol</u> **24**(1): 250.

Mathers, A. J., G. Peirano and J. D. Pitout (2015). "The role of epidemic resistance
plasmids and international high-risk clones in the spread of multidrug-resistant
Enterobacteriaceae." <u>Clin Microbiol Rev</u> **28**(3): 565-591.

Nicoloff, H., K. Hjort, D. I. Andersson and H. Wang (2024). "Three concurrent
mechanisms generate gene copy number variation and transient antibiotic
heteroresistance." <u>Nat Commun</u> **15**(1): 3981.

Pal, A. and D. I. Andersson (2024). "Bacteria can compensate the fitness costs of
amplified resistance genes via a bypass mechanism." <u>Nat Commun</u> **15**(1): 2333.

Parks, D. H., M. Chuvochina, P. A. Chaumeil, C. Rinke, A. J. Mussig and P. Hugenholtz
(2020). "A complete domain-to-species taxonomy for Bacteria and Archaea." <u>Nat</u>
<u>Biotechnol</u> **38**(9): 1079-1086.

Partridge, S. R., S. M. Kwong, N. Firth and S. O. Jensen (2018). "Mobile Genetic
Elements Associated with Antimicrobial Resistance." <u>Clin Microbiol Rev</u> **31**(4).

Partridge, S. R., G. Tsafnat, E. Coiera and J. R. Iredell (2009). "Gene cassettes and
cassette arrays in mobile resistance integrons." <u>FEMS Microbiol Rev</u> **33**(4): 757-784.

Puigbò, P., A. E. Lobkovsky, D. M. Kristensen, Y. I. Wolf and E. V. Koonin (2014).
"Genomes in turmoil: quantification of genome dynamics in prokaryote
supergenomes." <u>Bmc Biology</u> **12**.

Puigbo, P., K. S. Makarova, D. M. Kristensen, Y. I. Wolf and E. V. Koonin (2017).
"Reconstruction of the evolution of microbial defense systems." BMC Evol Biol **17**(1):
94.

Redondo-Salvo, S., R. Fernández-López, R. Ruiz, L. Vielva, M. de Toro, E. P. C. Rocha,
M. P. Garcillán-Barcia and F. de la Cruz (2020). "Pathways for horizontal gene transfer
in bacteria revealed by a global map of their plasmids." Nature Communications **11**(1).

Revell, L. J. (2024). "phytools 2.0: an updated R ecosystem for phylogenetic
comparative methods (and other things)." Peerj **12**.

Rozwandowicz, M., M. S. M. Brouwer, J. Fischer, J. A. Wagenaar, B. Gonzalez-Zorn,
B. Guerra, D. J. Mevius and J. Hordijk (2018). "Plasmids carrying antimicrobial
resistance genes in Enterobacteriaceae." J Antimicrob Chemother **73**(5): 1121-1137.

San Millan, A. (2018). "Evolution of Plasmid-Mediated Antibiotic Resistance in the
Clinical Context." Trends Microbiol **26**(12): 978-985.

San Millan, A. and R. C. MacLean (2017). "Fitness Costs of Plasmids: a Limit to
Plasmid Transmission." Microbiol Spectr **5**(5).

Souque, C., J. A. Escudero and R. C. MacLean (2021). "Integron activity accelerates
the evolution of antibiotic resistance." Elife **10**.

Steenwyk, J. L., T. J. Buida, 3rd, Y. Li, X. X. Shen and A. Rokas (2020). "ClipKIT: A
multiple sequence alignment trimming software for accurate phylogenomic inference."
PLoS Biol **18**(12): e3001007.

Touchon, M., J. A. Moura de Sousa and E. P. Rocha (2017). "Embracing the enemy:
the diversification of microbial gene repertoires by phage-mediated horizontal gene
transfer." Curr Opin Microbiol **38**: 66-73.

Valiatti, T. B., F. F. Santos, F. O. Bessa-Neto, R. Veiga, S. Simionatto, G. H. de Almeida
Souza, M. S. M. Vaz, A. C. C. Pignatari, R. Cayo, A. C. Gales and N. Guarani (2024).
"Emergence of multidrug-resistant Providencia rettgeri clone in food-producing
animals: A public health threat." One Health **19**: 100887.

van Duin, D. and D. L. Paterson (2016). "Multidrug-Resistant Bacteria in the
Community: Trends and Lessons Learned." Infect Dis Clin North Am **30**(2): 377-390.

Vogwill, T. and R. C. MacLean (2015). "The genetic basis of the fitness costs of
antimicrobial resistance: a meta-analysis approach." Evol Appl **8**(3): 284-295.

Wang, P., C. Li, Z. Yin, X. Jiang, X. Li, X. Mu, N. Wu, F. Chen and D. Zhou (2023).
"Genomic epidemiology and heterogeneity of Providencia and their bla(NDM-1)-
carrying plasmids." Emerg Microbes Infect **12**(2): 2275596.

Wang, Y. and T. Dagan (2024). "The evolution of antibiotic resistance islands occurs
within the framework of plasmid lineages." <u>Nat Commun</u> **15**(1): 4555.

Woodford, N., J. F. Turton and D. M. Livermore (2011). "Multiresistant Gram-negative
bacteria: the role of high-risk clones in the dissemination of antibiotic resistance."
<u>FEMS Microbiol Rev</u> **35**(5): 736-755.

Zongo, P. D., N. Cabanel, G. Royer, F. Depardieu, A. Hartmann, T. Naas, P. Glaser and
I. Rosinski-Chupin (2024). "An antiplasmid system drives antibiotic resistance gene
integration in carbapenemase-producing Escherichia coli lineages." <u>Nat Commun</u>
15(1): 4093.

Figure legends

Figure 1. Evolutionary dynamics of plasmid-borne antibiotic resistance genes

compared to other plasmid genes in Enterobacteriaceae.

(A) Gain rates (appearance of new gene families per substitution per site) for ARGs

versus other plasmid genes. (B) Loss rates (disappearance of gene families per

substitution per site) for ARGs versus other plasmid genes. (C) Expansion rates

(addition of gene copies within existing families per substitution per site) for ARGs

versus other plasmid genes. (D) Reduction rates (loss of gene copies within existing

families per substitution per site) for ARGs versus other plasmid genes. (E) Log_2 ratio

of gain to loss rates for ARGs. (F) Log_2 ratio of gain to loss rates for other plasmid

genes. Each point represents one Enterobacteriaceae species. Box plots show median

and quartiles. Gene families were defined as orthologous groups (OGs).

Figure 2. Expansion and reduction rate patterns across antibiotic resistance

classes and Enterobacteriaceae species.

Heatmaps showing evolutionary rates for different antibiotic resistance classes across

bacterial species. (A) Expansion rates. (B) Reduction rates. Rows represent

Enterobacteriaceae species, and columns represent CARD-defined antibiotic

resistance classes. Color intensity indicates log10 scaled rate magnitude, with darker

colors representing higher rates. White cells indicate absence of the ARG class in the

species. Color scales are truncated for visualization clarity: gain rates (20th-80th

percentile), loss/expansion/reduction rates (5th-95th percentile); all statistical analyses

were performed on complete untruncated datasets.

Figure 3. Correlation between ARGs and other plasmid functional genes on

evolutionary timescales.

Sankey diagram showing the size of significant COG gene families that correlated with

ARGs across phylogenetic tree. Flow width represents the number of unique COG

families within each functional category that show significant correlation with antibiotic

resistance genes. (B) Bar plot showing the number of species have at least a significant

correlation for each COG functional category. X-axis represents the count of species

showing significant case between ARGs and genes from each COG category. The

triangle represents the median branches number involved in the statistical test for each

category. The border color of the bar indicates whether both gain and loss are

significant or only one of the tests is significant. Abbreviations of functional categories,

A: RNA processing and modification; B: Chromatin structure and dynamics; C: energy

production and conversion; D: cell cycle control, cell division, chromosome partitioning;

E: amino acid transport and metabolism; F: nucleotide transport and metabolism; G:

carbohydrate transport and metabolism; H: coenzyme transport and metabolism; I:

lipid transport and metabolism; J: translation, ribosomal structure and biogenesis; K:

transcription; L: replication, recombination and repair; M: cell wall/membrane/envelope

biogenesis; N: cell motility; O: posttranslational modification, protein turnover,

chaperones; P: inorganic ion transport and metabolism; Q: secondary metabolites

biosynthesis, transport and catabolism; R: general function prediction only; S: function

unknown; T: signal transduction mechanisms; U: intracellular trafficking, secretion, and

vesicular transport; V: defense mechanisms; W: extracellular structures; X: mobilome:

prophages, transposons; Y: nuclear structure; Z: cytoskeleton.

Figure 4. Association between presence of cARGs in bacterial lineages and plasmid borne ARG acquisition.

(A) Schematic showing sister clade comparison approach. Phylogenetic clades

containing high frequencies (≥80%) of specific cARG families are compared to sister

clades with low frequencies (<20%). Colored circles represent different bacterial clades.

(B) Fold-change comparison of evolutionary rates between ARG-containing clades and

sister clades. Y-axis shows the fold-change ratio (ARG clade / Sister clade) for gain

(purple) and loss (orange) rates of plasmid ARGs versus other plasmid genes. Dashed

line at y=1 indicates no difference between clades. (C) Interaction effect showing the

differential impact of cARG presence on gain versus loss dynamics. Y-axis represents

the interaction effect (fold-change, log scale) between clade type and evolutionary

process. Error bars show 95% confidence intervals. Statistical significance indicators:

***$p < 0.001$, *$p < 0.05$, ns = not significant.

Figure 5. Species distribution and lineage-level genome counts for ARG families

showing significantly higher plasmid gene-gain rates.

The y-axis indicates the antibiotic classes conferring resistance by the corresponding

gene families: aminoglycosides (ANT(3")), phenicols (CAT), fluoroquinolones (gyrB),

sulfonamides (sul), and diaminopyrimidines (dfr). The x-axis represents species as

defined by the Genome Taxonomy Database (GTDB).

Gain and loss of plasmid-borne antibiotic resistance genes are associated with their chromosomal counterparts and the broader plasmid gene repertoire in Enterobacteriaceae

Yang Liu[1, *], Yue Liu[1,2, *]

[1] Department of Microbiology, Zhejiang University School of Medicine, Zhejiang, China

[2] State Key Laboratory for Diagnosis and Treatment of Infectious Diseases of the First Affiliated Hospital, Zhejiang University School of Medicine, Zhejiang, China

[*]Correspondence: yang.liu29@zju.edu.cn (Yang L.); yuel@zju.edu.cn (Yue L.)

Supplementary Figure Legends

Figure S1. Ratio of gene copy number across species. (E) Log$_2$ ratio of expansion

to reduction rates for ARGs. (F) Log$_2$ ratio of expansion to reduction rates for other

plasmid genes. Each point represents one Enterobacteriaceae species. Box plots

show median and quartiles. Gene families were defined as orthologous groups (OGs).

Figure S2. Gain and Loss rate patterns across antibiotic resistance classes and

Enterobacteriaceae species. Heatmaps showing evolutionary rates for different

antibiotic resistance classes across bacterial species. (A) Gain rates. (B) Loss rates.

Rows represent Enterobacteriaceae species, and columns represent CARD-defined

antibiotic resistance classes. Color intensity indicates log10 scaled rate magnitude,

with darker colors representing higher rates. White cells indicate absence of the ARG

class in the species. Color scales are truncated for visualization clarity: gain rates

(20th-80th percentile), loss/expansion/reduction rates (5th-95th percentile); all

statistical analyses were performed on complete untruncated datasets.

Figure S3. Comparison of evolutionary rates between ARG-containing clades

and their sister clades in per-family analyses. Y-axis shows the fold-change ratio

(ARG clade / Sister clade) of plasmid ARGs versus other plasmid genes. Dashed line

at y=1 indicates no difference between clades. Violin plots show data distribution, and

box plots indicate the median and interquartile range.

Figure S4. Per-family comparison of evolutionary rates between ARG-containing clades and their sister clades. (A) Gain rates and (B) loss rates are shown for individual antibiotic resistance gene (ARG) families. A dashed line at y = 1 indicates no difference between clades. Asterisks denote statistical significance (* p < 0.05; ** p < 0.01; *** p < 0.001).

55